Multi-model inference of non-random mating from an information theoretic approach

Running title: Multi-model inference of non-random mating

1   Keywords: Mate choice, intra-sexual competition, sexual selection, assortative mating,

2   information criteria, multi-model inference

Abstract

Non-random mating has a significant impact on the evolution of organisms. Here, I

developed a modelling framework for discrete traits (with any number of phenotypes) to

explore different models connecting the non-random mating causes (intra sexual

competition and/or mate choice) and their consequences (sexual selection and/or assortative

mating).

I derived the formulas for the maximum likelihood estimates of each model and used

information criteria for performing multimodel inference. Simulation results showed a good

performance of both model selection and parameter estimation. The methodology was

applied to data from Galician *Littorina saxatilis* ecotypes, to show that the mating pattern is

better described by models with two parameters that involve both mate choice and

intrasexual competition, generating positive assortative mating plus female sexual

selection.

As far as I know, this is the first standardized methodology for model selection and

multimodel inference of mating parameters for discrete traits. The advantages of this

framework include the ability of setting up models from which the parameters connect

causes, as intrasexual competition and mate choice, with their outcome in the form of data

patterns of sexual selection and assortative mating. For some models, the parameters may

have a double effect i.e. they cause both kind of patterns, while for others models there are

separated parameters for one kind of pattern or another.

The full methodology was implemented in a software called InfoMating (available at

http://acraaj.webs6.uvigo.es/InfoMating/Infomating.htm).

1. Introduction

In a given population, mating is random when an individual of one sex has an equal chance

of mating with any other individual of the opposite sex. Thus, random mating implies that

the overall frequency of mating between female and male types is equal to the product of

frequencies of these types in the population (Gavrilets 2004). Deviation from random

mating may occur as a consequence of variation in mating preferences, and/or due to

mechanical or physiological constraints. Detecting and understanding the behavioral

mechanisms causing deviations from random mating, in relation to traits used by the

organism for mating discrimination has been of great interest since such deviation may

have important evolutionary consequences (Rolan-Alvarez et al. 2015b).

Mate choice (or intersexual choice) is a process driven by different preference between

different mating types. This preferences may be mediated by phenotypic (sensorial or

behavioural) properties that affect the propensity of individuals to mate with certain

phenotypes (Jennions and Petrie 1997). Mate choice may generate a non-random mating

pattern.

In addition, non-random mating patterns can also be obtained due to intra-sexual

competition i.e. differential mating propensity, or vigour, among the types from one sex for

mating with whatever individual from the other sex (Spieth and Ringo 1983; Casares et al.

1998; Rolán-Alvarez and Caballero 2000).

From a population genetics point of view, a pattern of non-random mating can be divided

into sexual selection and/or assortative mating patterns. The decomposition of non-random

mating into sexual selection and assortative mating effects has been made originally from

multiple-choice experiments in *Drosophila* (Merrell 1950; Spieth and Ringo 1983;

Knoppien 1985), see also (Rolán-Alvarez and Caballero 2000).

Assortative mating is defined as the a posteriori deviation from random mating within

mated individuals (Rolán-Alvarez and Caballero 2000). Mate choice may produce an

assortative mating pattern. When similar phenotypes mate more often than expected, the

mating process leads to positive assortative mating, while negative assortative pattern is

obtained when dissimilar phenotypes mate more often (Merrell 1950; Lewontin et al. 1968;

Spieth and Ringo 1983).

Sexual selection is defined as the a posteriori observed change in gene or phenotype

frequencies in mated individuals with respect to population frequencies (Hartl and Clark

1997). Intrasexual competition may produce a sexual selection pattern. When some

phenotypes from one sex mate more frequently than other phenotypes from the same sex,

the mating process leads to a sexual selection pattern. However, a frequency-dependent

sexual selection pattern can also be produced by mate choice.

In a previous work (Carvajal-Rodríguez 2018b), the processes of intrasexual competition

and mate choice were modelled for discrete traits by means of the parameters m_{ij}, that

represent the mutual mating propensity between a female of type i and a male j. Therefore,

if A-type females prefer A-type males, this mate choice is modelled as a higher mutual

mating propensity between these types as compared with the mutual mating propensity of

the A females with other male types ($m_{AA} > m_{AB}$). On the other hand, if B-type males mate

more often than other males whatever the female, this intrasexual competition is modelled

by a higher marginal mating propensity of such males (see below).

By modelling the mating process as a differential mutual mating propensity among

different types of mating pairs, it is possible to express the mean change in mating

phenotypes as the information gained due to non-random mating (Carvajal-Rodríguez

2018b). Expressing the patterns obtained from mate choice and competition in terms of the

information captured in the mutual mating propensity models, permitted to identify the

necessary and sufficient conditions for describing random mating as the zero information

model (see below).

Thus, the mating information-based framework provides a formal approach for developing

a set of hypotheses about the causes (intra sexual competition and/or mate choice) and the

patterns they may provoke (sexual selection and/or assortative mating). In addition, data-

based evidence can be used for ranking each hypothesis and perform multi-model-based

inference (Link and Barker 2006; Burnham et al. 2011; Aho et al. 2014).

In the following sections I proceed as follows:

1.- Given the population frequencies for some discrete trait I define the multinomial

saturated mating model in terms of the mutual mating propensity parameters. The

maximum-likelihood estimates of these parameters are the pair total indices (PTI) as

defined in (Rolán-Alvarez and Caballero 2000). Once the saturated model is defined I

obtain the three necessary and sufficient conditions for random mating. Afterwards, by

relaxing these conditions it is possible to generate models for which differential marginal

mating propensity may produce female or male sexual selection without assortative mating,

or on the contrary, models for which some mutual mating propensities represent mate

choice that may produce assortative mating and frequency dependent sexual selection. I

obtain the maximum likelihood estimates for the parameters of these models.

2.- Relying in the previous section, it is possible to generate several mutual mating

propensity models and apply information criteria for selecting the best candidate ones and

estimating the mating parameter values based on the most supported models. I developed a

software called InfoMating to do so.

3.- Finally, I demonstrate the methodology by analysing simulated and real data.

2. Mutual mating propensity models

Consider a female trait with k_1 different phenotypes and a male trait with k_2 phenotypes, the

total number of possible mating types is $K = k_1 \times k_2$. Let have a sample of n' matings from

which n'_{ij} correspond to i-type females that mated with j-type males. If the probability of the

mating $i{\times}j$ is q'_{ij}, then the logarithm of the multinomial likelihood function of the sample is

$$lnL = C + \sum_{ij}^{K} n'_{ij} \ln(q'_{ij})$$

where C is the multinomial coefficient which is constant given the sample. As it is well-

known, the maximum likelihood estimator of the multinomial probability of the mating $i{\times}j$

is n'_{ij} / n'.

2.1 Saturated non-random mating model M_{sat}

Let the population under study have n_{1i} females of type i from a total of n_1 females and n_{2j}

males of type j from a total of n_2 males. Therefore, the population frequency of females of

type i is $p_{1i} = n_{1i} / n_1$ and the population frequency of males of type j is $p_{2j} = n_{2j} / n_2$.

The mating probability between types i and j can be expressed as $q'_{ij} = m_{ij}q_{ij}$ (Carvajal-

Rodríguez 2018b) where q_{ij} is the product of the female and male population frequencies of

each type ($q_{ij} = p_{1i} \times p_{2j}$) and m_{ij} is the normalized mutual mating propensity that refers to the

expected number of matings given an encounter between females of type i and males of

type j. The values m_{ij} are normalized so that $\Sigma q'_{ij} = 1$.

Under this multinomial model, the log-likelihood of the sample is

$$lnL_{sat} = C + \sum_{ij}^{K} n'_{ij} \ln(m_{ij}q_{ij})$$ (1)

This model is saturated (M_{sat}) because it has as many parameters as independent mating-

type frequencies, $P_{sat} = K - 1$. The female and male population frequencies, p_1 and p_2, are

considered to be known or alternatively, they need to be estimated in all the models.

Therefore, for model comparison, the population frequencies can be ignored when counting

the number of parameters involved in each model.

The maximum likelihood estimate (MLE) of m_{ij} is $(n'_{ij} / n')/q_{ij} = PTI_{i,j}$ where $PTI_{i,j}$ is the

pair total index i.e. the frequency of the observed pair types divided by their expected

frequency under random mating (Rolán-Alvarez and Caballero 2000).

In this work I am interested in the estimation of the mutual mating propensity parameters

(from hereafter mutual-propensity parameters) for various intrasexual competition and mate

choice models. From that point of view, is convenient to express the maximum likelihood

estimator in a different way which I call λ-notation.

2.2 λ-notation

Let consider the non-normalized parameters m'_{ij} so that $m_{ij} = m'_{ij} / M$ where $M =$

$\sum_{i,j} q_{ij} m'_{ij}$ is the mean mutual-propensity. The MLE of m'_{ij} under M_{sat} is simply $M \times PTI_{i,j}$

i.e. $M \times (n'_{ij} / n') / q_{ij}$ that can be conveniently rearranged as $(n'_{ij} / q_{ij}) / (n' / M)$. Because the

mating parameters are normalized, it is possible, without loss of generality, to set just one

of the m'_{ij} to an arbitrary value of 1. Thus, let set $m'_{k1k2} = 1$ and note (details in Appendix A)

that in such case $n' / M = n'_{k1k2} / q_{k1k2}$. Therefore, the MLE of the parameters of the

saturated model can be expressed as

$$\widehat{m}'_{ij} = \frac{\lambda(m'_{ij})}{\lambda(1)}$$

where

$$\lambda(\theta) = \frac{\sum_t^A n'_t}{\sum_{t1t2}^A q_{t1t2}} \qquad\qquad (2)$$

i.e., the function λ of a mating parameter θ is the sum of the counts of mating types having

mutual propensity θ divided by the sum of the expected frequency of the θ mating types

under random mating.

Thus, $\lambda(m'_{ij})$ expresses the sum of the observed matings with mating parameter m'_{ij} divided

by the product of the population frequencies from each partner type. Similarly, $\lambda(1)$

corresponds to the sum of the observed matings having unity mating parameter divided by

the corresponding products of population frequencies.

As already mentioned, the most parameterized model is the saturated model that has K-1

parameters so, when divided by the mean mutual propensity M, the estimates $\lambda(m'_{ij})$ /

$(M\lambda(1))$ are the corresponding pair total indices (PTI_{ij}).

The model M_{sat} is the most complex model that can be fitted to the available data. The

principle of parsimony suggests to consider reduced special cases of this saturated model.

The most reduced is the random mating model that can be obtained by applying three

necessary and sufficient conditions to the saturated model. By relaxing these conditions it is

possible to express mutual propensity models that may produce sexual selection, assortative

mating or both. Following, I computed the ML estimates of different classes of reduced

models that require less fixed parameters.

2.3 Random mating model M_0

The random model M_0 corresponds to the simplest, most reduced model, which is nested

with all the other (is a particular case of any other model) while it is not possible to derive

any simplified version from it. When mating is at random, the mating probability between

types i and j is $q'_{ij} = q_{ij} = p_{1i} \times p_{2j}$. Under this model, the information indices would be zero

(Carvajal-Rodríguez 2018b). This zero-information model is a particular case of the

saturated one when the mutual propensity is equal for every mating type. The number of

independent mating parameters is $P_0 = 0$.

The log-likelihood of the sample of mating is

$$lnL_0 = C + \sum_{ij}^{K} n'_{ij} \ln(q_{ij}) \qquad (3)$$

Now, let's define the marginal propensity m_{Fem_i} for a female of type i as

$$m_{Fem\_i} = \sum_{j}^{males} p_{2j} \frac{m'_{ij}}{M} \quad = \sum_{j}^{males} p_{2j} m_{ij} \qquad (4)$$

Similarly for a male of type j

$$m_{Male\_j} = \sum_{i}^{females} p_{1i} \frac{m'_{ij}}{M} = \sum_{i}^{females} p_{1i} m_{ij}$$

Then, the model M_0 model corresponds to M_{sat} subjected to the following restrictions:

i)   Equal female marginals: $m_{Fem\_i} = m_{Fem\_j} \forall i, j$ (5)

ii)   Equal male marginals: $m_{Male\_i} = m_{Male\_j} \forall i, j$

iii)   Multiplicativity: $m_{ij} = m_{Fem\_i} \times m_{Male\_j} \forall i, j$

It is useful to express M_0 in terms of the three restrictions because by relaxing some of

them, different classes of models emerge. Namely, the marginal propensity of a given male

phenotype corresponds to the average mutual mating propensity of that male so, if different

male phenotypes have different marginal propensities, this represents differential

competition or mating vigour among males that may provoke a sexual selection mating

pattern. Similarly, if some matings m_{ij} has different value than the rest of matings, this mate

choice behaviour may produce assortative mating with or without sexual selection.

Therefore, by relaxing some of the conditions in (5), it is possible to control the kind of

causes that produce the different non-random mating patterns. In fact, there are three

general classes of models that can be combined. The two first classes correspond to

relaxing first or second condition and involve intrasexual competition in females or males,

provoking female or male sexual selection respectively. Provided that the third condition is

maintained these models cannot produce assortative mating pattern (see below). The third

class corresponds to relaxing the third condition and involve mate choice, that may provoke

just assortative mating, or both assortative mating and sexual selection, the latter depending

on the population frequencies (Fig. 1).

Fig 1. Mating models defined by intrasexual competition or mate choice, and their effect after
relaxing some of the conditions imposed to the random mating model M_0.

2.4 Intrasexual competition models

This class of model corresponds to relaxing the first and/or second conditions in M_0 while

maintaining the condition of multiplicativity (5-iii). The maintenance of the third condition

implies that the mutual-propensity of a mating pair (i,j) is the product of the marginal

female (m_{Fem}) and male (m_{Male}) propensities. Under this condition there should not be

deviation from random mating when comparing the observed and expected frequencies

within mates and the assortative mating pattern should not be observed (Carvajal-

Rodríguez 2018b). I distinguished models that generate a sexual selection pattern in just

one sex or in both.

2.4.1 Intra-female competition

Relaxing condition (5-*i*) implies that at least one female marginal propensity, say female of

type *A*, is different from the rest of female types i.e. $m_{Fem\_A} \neq m_{Fem\_B}$ with $B \neq A$. On the

other side, the marginal propensity of males should be the same which means that there is

no intra-male competition, they mate equal in average.

Therefore, a model with intra-female competition is obtained by defining every mutual-

propensity involving a female of type *i*, by an absolute (unnormalized) mating parameter a_i

as follows

$m'_{11} = m'_{12} = ... = m'_{1k2} = a_1$

$m'_{21} = m'_{22} = ... = m'_{2k2} = a_2$
.
.
.
$m'_{(k1-1)1} = m'_{k12} = ... = m'_{k1k2} = a_{k1-1}$

$m'_{k11} = m'_{k12} = \ldots = m'_{k1k2} = 1$ (6)

with $a_i > 0 \ \forall h$.

Note that the relationships among the parameters will not be altered when dividing them by

a_{k1} so that $a_{k1} = 1$. Under this model it can be as much as k_1-1 free mating parameters.

The normalization factor is the mean mutual-propensity $M = \sum m'_{ij}q_{ij}$. When computing the

female and male marginal propensities (4) it is seen that

$m_{Fem\_1} = a_1/M; \ m_{Fem\_2} = a_2/M \ldots; \ m_{Fem\_k1} = 1/M$

$m_{Male\_1} = m_{Male\_2} = m_{Male\_3} = \ldots = m_{Male\_k2} = M/M = 1$

The model (6) has equal male marginal propensity and is multiplicative. The MLE of the

model parameters is

$\hat{a}_i = \dfrac{\lambda(a_i)}{\lambda(1)}$ (7)

where $\lambda(\theta)$ is defined as in (2). Thus, $\lambda(a_i)$ expresses the sum of the observed matings

having mating parameter a_i, divided by the sum of the product of the population

frequencies from each partner type. Similarly, $\lambda(1)$ corresponds to the sum of the observed

matings having unity mating parameter divided by the sum of the corresponding products

of population frequencies (details in Appendix A).

2.4.2 Intra-male competition

Relaxing condition (5-*ii*) implies that at least one male marginal propensity, say male of

type A, is different from the rest of male types i.e. $m_{Male\_A} \neq m_{Male\_B}$ with $B \neq A$. On the

other side, the marginal propensity of females should be the same which means that there is

no intra-female competition, they mate equal in average. The corresponding model can be

obtained just by interchanging rows with columns in (6). Noting the parameters as b_j

instead of a_i, the maximum likelihood estimate is

$$\hat{b}_j = \frac{\lambda(b_j)}{\lambda(1)} \qquad (8).$$

2.4.3 Intra-female and male competition

Obviously, by relaxing conditions (5-*i*) and (5-*ii*) the marginal propensities will be different

within females and males. The corresponding model combines models (6) and (8) and has

as much as $(k_1-1) \times (k_2-1)$ parameters in the most parameterized case, and a minimum of two

(female and male) for the less parameterized one, in order to maintain the multiplicativity

condition (5-*iii*). This type of model may produce a pattern of sexual selection in both sexes

without assortative mating. By notational convenience, I fix the category k_1 in females and

k_2 in males as having unitary parameters. Therefore

$\qquad m'_{ij} = a_i b_j, \; i<k_1, \; j<k_2; \; m'_{ik2} = a_i, \; i<k_1; \; m'_{k1j} = b_j; \; j <k_2, \; m'_{k1k2} = 1; \; \text{with } a_i > 0, \; b_j > 0$

$\forall i, j.$

This model is multiplicative (see Appendix A) and the parameters MLE are

$$\hat{a}_i = \left(\frac{p_{1k1}}{p_{1i}}\right) \frac{\sum_{j=1}^{k2} n'_{ij}}{\sum_{j=1}^{k2} n'_{k1j}} = \frac{\lambda(a_i + \sum_j a_i b_j)}{\lambda_{fem}(1)} \qquad (9)$$

$$\hat{b}_j = \left(\frac{p_{2k2}}{p_{2j}}\right)\frac{\sum_{i=1}^{k1} n'_{ij}}{\sum_{i=1}^{k1} n'_{ik2}} = \frac{\lambda(b_j + \sum_i a_i b_j)}{\lambda_{male}(1)}$$

where $\lambda(a_i + a_i b_1 + ... a_i b_j + ...)$ stands for the sum of observed mating types having mating

parameter a_i or $a_i b_1$, etc... divided by the sum of their expected random mating frequencies

(in our case the sum of the expected frequencies of matings with mating parameter a_i or $a_i b_j$

for every male type, is p_{1i}), and $\lambda_{sex}(1)$ refers to the sum of cases that contribute with 1 to

the mutual-propensity by the side of such sex divided by the corresponding sum of

expected frequencies (e.g. in a multiplicative model as the above, every mating $k_1 \times j$ has

contribution 1 by the female side and the sum of frequencies is p_{1k1}). Formulae (9) is

similar to (7) and (8). Note that (9) becomes (7) by fixing every b_j as 1 while it becomes (8)

by fixing every a_i as 1.

The percentage of sexual selection information corresponding to each sex, J_{S1} and J_{S2} in

(Carvajal-Rodríguez 2018b), would depend on the population frequencies and on the

mating parameter values.

2.5 Mate choice models

Mate choice or intersexual discrimination models correspond to the class of non-

multiplicative models, i.e. they can be obtained by relaxing the condition (5-iii) and may

produce assortative mating patterns (positive or negative). If the female marginal

propensities are equal and the same is true for the males (conditions 5-i and 5-ii hold) there

would not be sexual selection neither in females nor males, and the model may produce

only assortative mating patterns. However, this cannot be guaranteed in general because the

occurrence of the sexual selection pattern is frequency dependent under non-multiplicative

models (see below).

Let consider a model where the unnormalized mutual-propensities are

$m'_{ii} = a_i > 0 \ \forall i \in \min\{k_1, k_2\}$ and $m'_{ij} = 1$ for $i \neq j$.

Thus, the homotype ($i \times i$) mutual-propensities are parameterized while the heterotype are

not. This model is non-multiplicative in general, because the contribution of the type i to

the mutual-propensity is distinct in m_{ii} that in m_{ij} or in m_{ji} (although with an even number of

types a multiplicative model can be obtained by setting $m'_{ii} = 1 / m'_{jj}$).

By recalling the definition of marginal propensities in (4), the condition for equal female

marginal $m_{Fem\_i} = m_{Fem\_j}$ is

$p_{2i}(a_i - 1) = p_{2j}(a_j - 1)$ (10)

and in males

$p_{1i}(a_i - 1) = p_{1j}(a_j - 1)$

In general, depending on the conditions in (10), the mate choice models have double effect

i.e. they produce assortative pattern jointly with sexual selection in at least one sex.

The maximum likelihood estimate for the model parameters is

$\hat{a}_i = \dfrac{\lambda(a_i)}{\lambda(1)}$ (11)

Note that the homotype mating parameter may imply higher mutual propensity than the

heterotype ($a_i > 1$, positive assortative mating) or viceversa, the homotype has lower mutual

propensity ($a_i < 1$, negative assortative). The number of different parameters ranges from 1

($a_1 = a_2 = ... = a_i$) to H; where $H = min\{k_1, k_2\}$ corresponds to the maximum possible

number of different homotype matings.

It is also possible to define mate choice models with the heterotype mutual-propensities

parameterized instead of the homotype ones (see Appendix A for details).

2.6 Models with intrasexual competition and mate choice parameters

I have shown that mate choice models may generate both kind of patterns, assortative

mating and sexual selection, depending on the within sex population frequencies. While it

is not possible to assure that the mate choice model produce no sexual selection, it is

possible to combine the previous models to ensure that there are parameters directly linked

to intra-sexual competition and parameters directly linked to mate choice. These combined

models have the property that when the mate choice parameter is set to 1, there is only a

known sexual selection effect as caused by the competition parameter (female, male, or

both). When the mate choice parameter is added, the assortative mating pattern appears and

an extra effect of frequency-dependent sexual selection may be added to that of the original

competition parameter.

2.6.1 Models with male competition and mate choice: independent parameters

Consider the model $m'_{i1} = \alpha$; $m'_{ii} = c$ for $i \neq 1$ and $m'_{ij} = 1$ otherwise; with $i \leq k_1, j \leq k_2$. An

example of this kind of model can be seen in Fig. 2.

$$\begin{pmatrix} \alpha & 1 & 1 & 1 & 1 \\ \alpha & c & 1 & 1 & 1 \\ \alpha & 1 & c & 1 & 1 \\ \alpha & 1 & 1 & c & 1 \\ \alpha & 1 & 1 & 1 & c \end{pmatrix}$$

Fig 2. An example of male competition and mate choice independent parameters model with 5×5
mating types. α is the male competition parameter and c is the choice parameter.

For the particular case of $\alpha \neq 1$, $c = 1$; the model has within male competition that

corresponds to the marginal propensity α of the type-1 male compared with the other males,

so, a male sexual pattern may be generated. On the contrary, the female marginal

propensities are equal so there is no female competition. Considering mate choice and the

assortative mating pattern, when $c = 1$ the model is multiplicative so assortative mating

should not be produced, indeed, the pair sexual isolation statistics (PSI) are equal (see

Appendix A for details) so the joint isolation index (I_{PSI}) is 0 as corresponds to the absence

of assortative mating (see also Carvajal-Rodríguez 2018b).

However, by taking $c \neq 1$ a new component is added to the male sexual selection. The

parameter c corresponds to mate choice and produces positive ($c > 1$) or negative ($c < 1$)

assortative mating. The joint isolation index is a function of the parameter c and the

population frequencies. Female sexual selection may also emerge depending on the value

of c and the population frequencies.

The MLEs of both parameters are

$$\hat{\alpha} = \frac{\lambda(\alpha)}{\lambda(1)}$$

$$\hat{c} = \frac{\lambda(c)}{\lambda(1)}$$

A variant of the above model can be generated by changing the c parameter from the main

diagonal to the anti-diagonal. Similarly female sexual competition linked to the α-

parameter is obtained by transposing the matrix of the model so that the first column

converts in the first row and so on.

2.6.2 Models with male competition and mate choice: compound parameters

Consider the model $m'_{11} = c\alpha$; $i > 1$: $m'_{i1} = \alpha$; $m'_{ii} = c$; and $m'_{ij} = 1$ otherwise; with $i \le k_1$, $j \le$

k_2. An example of this model can be seen in Fig. 3.

$$\begin{pmatrix} \alpha c & 1 & 1 & 1 & 1 \\ \alpha & c & 1 & 1 & 1 \\ \alpha & 1 & c & 1 & 1 \\ \alpha & 1 & 1 & c & 1 \\ \alpha & 1 & 1 & 1 & c \end{pmatrix}$$

Fig 3. An example of male competition and mate choice compound parameters model with 5×5

mating types. α is the male competition parameter and c is the choice parameter.

With $c = 1$ the model is the same as the previous one. When $c \neq 1$ is added, the mate choice

parameter provokes an extra effect of sexual selection in males and females, plus assortative

mating. The MLE of α and c are

359
$$\hat{\alpha} = \frac{\sum_{i=1}^{k1} n'_{i1} / (c q_{11} + \sum_{i>1} q_{i1})}{\lambda(1)} = \frac{\lambda_{c.11-}(\alpha)}{\lambda(1)}$$

Where $\lambda_{c.11-}(\alpha)$ indicates that for mating types with mating parameter α, the expected

frequency indexed as 11 (i.e. q_{11}) is weighted by c. Similarly,

362
$$\hat{c} = \frac{\sum_{i=1}^{k} n'_{ii} / (\alpha q_{11} + \sum_{i>1} q_{ii})}{\lambda(1)} = \frac{\lambda_{\alpha.11-}(c)}{\lambda(1)}$$

Where $k = \min\{k_1, k_2\}$ and $\lambda_{\alpha.11-}(c)$ indicates that for the mating types with mating parameter

c, the expected frequency indexed as 11 is weighted by α.

The above estimates are dependent one on each other, so, for obtaining the estimates of this

compound parameter model, I have used a numerical bounded Nelder-Mead simplex

algorithm with restriction $\alpha > 0$, $c > 0$ (Press 2002; Singer and Singer 2004; Gao and Han

2012).

2.6.3 General model with male competition and mate choice parameters

The general model with male competition and mate choice parameters is $m'_{11} = c_1\alpha$; $i > 1$:

$m'_{i1} = \alpha$; $m'_{ii} = c_k$; and $m'_{ij} = 1$ otherwise; with $i \leq k_1$, $j \leq k_2$. A particular case of this model

can be seen in Fig. 4.

$$
\begin{pmatrix}
\alpha c_1 & 1 & 1 & 1 & 1 \\
\alpha & c_1 & 1 & 1 & 1 \\
\alpha & 1 & c_3 & 1 & 1 \\
\alpha & 1 & 1 & c_4 & 1 \\
\alpha & 1 & 1 & 1 & c_5
\end{pmatrix}
$$

Fig 4. Male sexual selection and mate choice compound model with 5 × 5 mating types. α is the

male sexual selection parameter and c_k's are the choice parameters with $c_2 = c_1$.

Note that for distinguishing the competition and mate choice parameters, it is necessary that

at least one c_k parameter is equal to c_1 (as in Figs. 3 and 4) or that $c_1 = 1$ as in Fig. 2,

otherwise the m_{11} parameter does not distinguishes competition and choice. Therefore, the

model in Fig. 4 has H parameters with $H = \min\{k_1, k_2\}$ from which, $H - 1$ are choice

parameters (c) plus one male competition parameter α. The MLE are

$$\hat{\alpha} = \frac{\sum_{i=1}^{k1} n'_{i1} / (c_1 q_{11} + \sum_{i>1} q_{i1})}{\lambda(1)} = \frac{\lambda_{c1.11-}(\propto)}{\lambda(1)}$$

$$\hat{c}_1 = \frac{n'_{11}/(\alpha q_{11})}{\lambda(1)} = \frac{\lambda_{\propto.11-}(c_1)}{\lambda(1)}$$

$$\hat{c}_{k>1} = \frac{n'_{kk}/(q_{kk})}{\lambda(1)} = \frac{\lambda(c_k)}{\lambda(1)}$$

The model parameters $c_{k>1}$ can be estimated directly from the sample; on the contrary, the α

and c_1 estimates are dependent on each other, so, for obtaining these estimates, I used a

numerical bounded Nelder-Mead simplex algorithm with restriction $\alpha > 0$, $c_1 > 0$ (Press

2002; Singer and Singer 2004; Gao and Han 2012).

Previous models were simplified versions of the general model. For example the model in

Fig. 2 is the general model with restrictions $c_1 = 1$; $c_2 = c_3 =... = c_k = c$. Also, the model in

Fig. 3 corresponds to $c_1 = c_2 = c_3 =... = c_k = c$. Another particular case that could be defined

is $c_1 = c$; $c_2 = c_3 =... = c_k = 1$. In the latter, the MLE of the parameters can again be expressed

as a quotient of lambdas similar to the compound parameter case

$$\hat{\alpha} = \frac{\lambda_{c1.11-}(\propto)}{\lambda(1)}$$

$$\hat{c} = \frac{\lambda_{\propto.11-}(c)}{\lambda(1)}$$

It is also possible to define other general model with the mate choice parameters in the anti-

diagonal. Using the λ notation, the estimates follow the same formulae as defined for the

general model with the choice parameters in the main diagonal.

Concerning models with female competition and mate choice, they are obtained just by

transposing the matrices of the mating parameters.

2.7 General double effect models

The mating parameters $m_{ij} = \theta_{ij}$ with the restriction that at least some is equal to one, permit

to generate any particular model. In general, these models produce patterns of sexual

selection and assortative mating with each parameter possibly linked to the occurrence of

both (see Appendix A). The MLE is

407           $$\hat{\theta}_{ij} = \frac{\lambda(\theta_{ij})}{\lambda(1)} \qquad (12)$$

The most parameterized model of this kind is the saturated, with *K*-1 parameters. In such

case, as already mentioned, the estimates in (12) are the corresponding pair total indices

(PTI).

All the above derived MLE formulae have been verified by numerical approximation using

the bounded Nelder-Mead simplex algorithm (Press 2002; Singer and Singer 2004; Gao and

Han 2012).The set of described models jointly with their expected effects are summarized

in Table 1.

Table 1. Mutual mating propensity models as defined by different parameters in a case

with two different types at each sex ($k_1 = k_2 = 2$). The unnormalized m'_{ij} values not

explicitly given are assumed to be 1.

Name (abbreviation)	Model	MLE	Effect
Random (M_0)	$m'_{ij} = 1 \forall i,j$		Random mating
Competition Multiplicative Models			
Female competition (S_{Fem}-1P)	$m'_{11} = m'_{12} = a$	$\hat{a} = \lambda(a)/\lambda(1)$	Fem sexual selection
Male competition (S_{Male}-1P)	$m'_{11} = m'_{21} = b$	$\hat{b} = \lambda(b)/\lambda(1)$	Male sexual selection
Female and male competition ($S2$-2P)	$m'_{11} = ab$		2-sex sexual selection
	$m'_{12} = a$	$\hat{a} = \lambda_{fem}(a)/\lambda_{fem}(1)$	
	$m'_{21} = b$	$\hat{b} = \lambda_{male}(b)/\lambda_{male}(1)$	
Mate Choice Models			
One-parameter (C-1P)	$m'_{11} = m'_{22} = a$	$\hat{a} = \lambda(a)/\lambda(1)$	Assortative mating + sex sel (freqdep)
General mate choice (C-HP)	$m'_{11} = a_1, m'_{22} = a_2$	$\hat{a}_i = \lambda(a_i)/\lambda(1)$	Assortative mating + sex sel (freqdep)
Competition and Mate Choice			
2 independent parameters ($S_{Fem}C$-2P)	$m'_{1j} = \alpha;$ $m'_{jj} = c; j > 1$	$\hat{\alpha} = \lambda(\alpha)/\lambda(1)$ $\hat{c} = \lambda(c)/\lambda(1)$	α-sexual selection in one sex + mate choice effect c-assortative mating
2 parameters (1 compound: $S_{Fem}C$-2Pc)	$m'_{11} = c\alpha; j > 1:$ $m'_{1j} = \alpha; m'_{jj} = c;$	$\hat{\alpha} = \lambda_{c.11\text{-}}(\alpha)/\lambda(1)$ $\hat{c} = \lambda_{\alpha.11\text{-}}(c)/\lambda(1)$	α-sexual selection in one sex + mate choice effect c-assortative mating
H parameters (1 compound: $S_{Fem}C$-HPc)	$m'_{11} = c_1\alpha; j > 1:$ $m'_{1j} = \alpha; m'_{jj} = c_k;$	$\hat{\alpha} = \lambda_{c1.11\text{-}}(\alpha)/\lambda(1)$ $\hat{c}_1 = \lambda_{\alpha.11\text{-}}(c_1)/\lambda(1)$ $\hat{c}_{k \neq 1} = \lambda(c_k)/\lambda(1)$	α-sexual selection in one sex + mate choice effect c_k-assortative mating
General Double Models (D-xP)			Assortative mating

+ sex sel (freq dep)

Saturated (M_{sat}) $\qquad m'_{ij} = a_{ij}; m'_{k1k2} = 1 \qquad \hat{a}_i = \lambda(a_i)/\lambda(1)$

419 k_1: number of female categories; k_2: number of male categories; $H = min\{k_1,k_2\}$; sexsel (freqdep): frequency
420 dependent sexual selection.

423 3. Model Selection and multimodel inference

Information-based model selection and multi-model inference can be applied for describing

uncertainty in a set of models and performing inference on the parameters of interest

(Burnham et al. 2011; Grueber et al. 2011; Barker and Link 2015; Claeskens 2016).

There are several information criteria at hand, although trusting on a single form of

information criterion is unlikely to be universally successful (Liu and Yang 2011; Vrieze

2012; Brewer et al. 2016; Aho et al. 2017; Dziak et al. 2019). In the present work, two

Kullback- Leibler divergence-based measures plus the so called Bayesian information

criterion are considered.

3.1. Information criteria

The Akaike information criterion (AIC) provides the link between the Kullback-Leibler

divergence and the maximized log-likelihood of a given model (Akaike 1973). Here I use

the sample-corrected version AICc, because it is asymptotically equivalent and may work

better for small sample size

$\qquad \text{AIC}_c = -2ln(L) + 2P_m + (2P_m(P_m+1))/(n' - P_m -1)$

where L is the maximum likelihood of the model, P_m the total number of estimated mating

parameters and n' is the number of matings.

There is also a version for the symmetric K-L divergence (Jeffrey's) called the KICc

criterion (Cavanaugh 2004; Keerativibool 2014). Because mate choice models can be

described by the informational flow in the mating phenotypes, in the form of the Jeffrey's

divergence (Carvajal-Rodríguez 2018b), it seems adequate considering KICc criterion

besides the AICc one

$KIC_c = -2ln(L) + n'\ln(n' / (n' - P_m)) + P_2$

with $P_2 = n'[(n' - P_m)(2P_m + 3)-2] / [(n' - P_m -2)(n' - P_m)]$

Finally, the Bayesian information criterion (BIC Schwarz 1978) permits an approximation

to the Bayes factor applied for model comparison (Wagenmakers 2007)

$BIC = -2ln(L) + P_m\ln(n')$

3.2 Overdispersion

In the context of model selection, data overdispersion, i.e. greater observed variance than

expected, could generate the selection of overly complex models. The simplest approach to

estimate overdispersion is by computing a single variance inflation factor (v). This inflation

factor is the observed variation divided by the expected under the model with the highest

likelihood (M_c), other than the saturated, among the proposed ones (Richards 2008;

Symonds and Moussalli 2011). It can be asymptotically approximated by the deviance i.e.

twice the difference between the log-likelihood of the saturated (M_{sat}) and the M_c model,

divided by the difference in the number of parameters ($P_{Msat} - P_{Mc}$) between both models

$v = 2[\ln(L_{Msat}) - \ln(L_{Mc})] / df$

where $df = P_{Msat} - P_{Mc}$.

If $1 \leq v \leq 4$ this indicate overdispersion, while if higher than 4-6 this may indicate poor

model structure and the construction of the set of models should be reconsidered (Burnham

and Anderson 2002). For v values around 1 to 4, quasi-likelihood theory provides a way to

analyse over dispersed data (Anderson et al. 1994; Richards 2008). The quasi-likelihood is

the likelihood divided by an estimate of v. The quasi-likelihood version of the various

information criteria, namely QAICc, QKIC$_c$ (Kim et al. 2014) and QBIC, is obtained

simply by replacing the likelihood with the quasi-likelihood in the corresponding formula.

In such cases, the number of parameters is increased by one and the model variance is

multiplied by v (see below). When the quasi-likelihood version is used, it must be done for

all models and criteria.

3.3. Model weights

Let IC be any information criterion. For a particular criterion and for any set of R models

there is a minimum criterion value e.g. AIC$_{cmin}$, BIC$_{min}$, etc. Thus, the models can be ranked

regarding the difference with that minimum

$\Delta_i = IC_i - IC_{min}$, for $i = 1, 2, …, R$

where IC$_i$ refers to any specific information criterion for the model i.

Models can also be ranked by their weights from higher to lower. The weight w_i refers to

the strength of evidence for that model (Burnham et al. 2011; Claeskens 2016)

$w_i = l_i / \Sigma l_j$ for $j = 1, 2, \ldots, R$

where $l_i = \exp(-0.5\Delta_i)$ is the relative likelihood of each model given the data.

3.4 Multi-model inference

Multi-model-based inference estimate the parameters of interest based on a group of the

most credible models instead of on a best-fit single model (Burnham and Anderson 2002;

Burnham et al. 2011; Symonds and Moussalli 2011). The multi-model inference is

performed as a model averaged prediction for the parameters that are variables in the best

model.

Before performing the average, the different models should be translated to the same scale

of mutual-propensity. For example, a model like $m'_{11} = 2$, $m'_{12} = m'_{21} = m'_{22} = 1$, (there is at

least one value fixed to 1) is not in the same scale that $m'_{11} = 2$, $m'_{12} = m'_{21} = m'_{22} = 0.5$.

Without loss of generality, the latter can be transformed into an equivalent model $m'_{11} = 4$,

$m'_{12} = m'_{21} = m'_{22} = 1$, which is now in the same scale that the first model.

The averaged parameter estimates were computed as a weighted mean where the weights

are the strength of evidence for each model as obtained under a given information criterion.

The parameters were averaged only over the models for which they appear as a variable.

Because the weights need to sum up to 1, it was necessary renormalize them by dividing by

the accumulated weight in the confidence subset.

Therefore, for each parameter m included in the confidence subset R_s the average was

computed as

$$\hat{m} = \frac{\sum_i^{Rs} w_i \hat{m}_i}{\sum_i^{Rs} w_i}$$

This way of performing the model averaged prediction is called natural averaging

(Symonds and Moussalli 2011).

Finally, the reliability of each parameter estimate was measured as the unconditional

standard error

$$Se(\widehat{m}) = \sum_i^{Rs} w_i \sqrt{vV(\hat{m}_i) + (\hat{m}_i - \hat{m})^2}$$

where $V(\hat{m}_i) = V(m_i |\text{model } i) = V(q') = q'(1-q') / n'$ is the model standard error squared and

v is the variance inflation factor.

The use of the sum of weights to estimate variable importance in regression models has

been criticized because multicollinearity among the predictor variables and the imprecision

of the weight measures (Galipaud et al. 2014; Cade 2015; Galipaud et al. 2017).

However, the mutual propensity parameters do not belong to a regression model and their

average is performed in the same scale and with comparable units. Therefore, under the

mutual mating propensity setting, the multimodel inference would work well as it was

confirmed by Monte Carlo simulation (next section).

4. Simulations

To test how well the exposed methodology is able to distinguish the different classes of

models and estimate the mating parameters, I used the sampling with replacement

algorithm in the program MateSim (Carvajal-Rodríguez 2018a) for generating mating

tables by Monte Carlo simulation (see Appendix B for detailed explanation).

The simulated mating tables correspond to one-sex competition and mate choice models.

The resulting mating tables were consequence of the mating pattern and the sampling

process. The obtained mating tables consisted in two types of information (Fig. B1 in

Appendix B). First, the population frequencies (premating individuals) which were

generated randomly for each simulation run. Second, the sample, that was 500 mating pairs

($n' = 500$) for a hypothetical trait with two classes at each sex. Because the simulated

species has high population size ($n = 10,000$) the mating process was represented as a

sampling with replacement, and the population frequencies were constant over the mating

season. The minimum phenotype frequency (MPF) allowed was 0.1.

Concerning the mating pairs, the number of occurrences for each mating type $i \times j$ was

obtained as

$$Q(i,j) = n' \times p_{1i} \times p_{2j} \times m_{ij}$$

where n' is the sample size, p_{1i} is the female population frequency for the phenotype i, p_{2j} is

the male population frequency for the phenotype j, and m_{ij} is the normalized mutual-

propensity of the mating pair.

The mutual-propensities assayed corresponded to random mating, $m'_{11}=m'_{22}=m'_{12}=m'_{21}=$

1(M_0 in Table 2); female competition ($\alpha = 2$) and mate choice ($c = 3$), with independent

parameters $m'_{11}=m'_{12}=2$, $m'_{22}=3$, $m'_{21}=1$ (SfC Table 2) and with compound parameters,

$m'_{11}=6$, $m'_{12}=2$, $m'_{22}=3$, $m'_{21}=1$ (SfCc Table 2); and male competition ($\alpha = 2$) and mate

choice ($c = 3$), with independent parameters $m'_{11}=m'_{21}=2$, $m'_{22}=3$, $m'_{12}=1$ (SmC Table 2)

and with compound parameters, $m'_{11} = 6$, $m'_{21} = 2$, $m'_{22} = 3$, $m'_{12} = 1$ (SmCc Table 2). The

parameter estimates were averaged over 1000 runs for each case.

The multimodel inference results appear in Table 2. The random mating was perfectly

estimated by the three IC methods. The competition plus mate choice parameter estimates

were fairly good under the three criteria. The estimates were slightly better under AICc and

slightly less accurate under BIC.

Table 2. Average (standard error) parameter estimates for different models under

sample size 500.

Model		m'_{11}	m'_{12}	m_{21}	m_{22}
M_0	**Expected**	1	1	1	1
	AICc	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
	KICc	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
	BIC	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
SfC	**Expected**	2	2	1	3
	AICc	1.94 ± 0.0017	1.99 ± 0.0009	1.0 ± 0.0000	2.95 ± 0.0025
	KICc	1.94 ± 0.0024	1.99 ± 0.0020	1.0 ± 0.0000	2.95 ± 0.0029
	BIC	1.90 ± 0.0060	1.94 ± 0.006	1.0 ± 0.0000	2.90 ± 0.0074
SfCc	**Expected**	6	2	1	3

	AICc	5.93 ± 0.0044	2.0 ± 0.0001	1.0 ± 0.0000	2.97 ± 0.0027
	KICc	5.92 ± 0.0051	2.0 ± 0.0001	1.0 ± 0.0000	2.97 ± 0.0028
	BIC	5.87 ± 0.0086	2.0 ± 0.0017	1.0 ± 0.0000	2.96 ± 0.0044
SmC	**Expected**	**2**	**1**	**2**	**3**
	AICc	1.94 ± 0.0020	1.0 ± 0.0000	1.99 ± 0.0013	2.94 ± 0.0030
	KICc	1.93 ± 0.0032	1.0 ± 0.0000	1.98 ± 0.0029	2.93 ± 0.0037
	BIC	1.90 ± 0.0062	1.00 ± 0.0000	1.93 ± 0.0065	2.88 ± 0.0080
SmCc	**Expected**	**6**	**1**	**2**	**3**
	AICc	5.93 ± 0.0046	1.0 ± 0.0000	2.0 ± 0.0001	2.97 ± 0.0029
	KICc	5.92 ± 0.0052	1.0 ± 0.0000	2.0 ± 0.0001	2.97 ± 0.0029
	BIC	5.87 ± 0.0085	1.0 ± 0.0000	2.0 ± 0.0010	2.97 ± 0.0037

551 M_0: Random mating model. SfC: female competition and mate choice with independent parameters. . SfCc:

552 female competition and mate choice with compound parameters. SmC: male competition and mate choice

553 with independent parameters. . SmCc: male competition and mate choice with compound parameters.

5. Example of application

Littorina saxatilis is a marine gastropod mollusc adapted to different shore habitats in

Galician rocky shores. There are two different ecotypes, an exposed-to-wave (smooth un-

banded, SU), and a non-exposed (rough banded, RB) ecotype. Several experimental studies

have shown that these ecotypes have evolved local adaptation at small spatial scale. For

example, stronger waves on the lower shore may provoke that the SU ecotype becomes

sexually mature at smaller size than the upper-shore (RB) ecotype. In addition, in some

areas of the mid-shore habitat, the two ecotypes occasionally mate, producing apparently

fertile intermediate morphological forms that are called hybrids (HY) (Rolan-Alvarez et al.

2015a).

Sexual isolation (positive assortative mating) between RB and SU morphs was observed in

wild mating pairs in the mid-shore zone, likewise size-assortative mating in all shore levels

(Cruz et al. 2001). It is assumed that the size is the key trait causing the increase of sexual

isolation in this model system, being the males the choosy sex in this species (Rolan-

Alvarez 2007).

Here, I reanalysed a *L. saxatilis* data set (Cruz et al. 2001) for estimating the mutual-

propensity parameters between the RB, SU and HY morphs in the mid-shore habitat (Table

3).

Table 3. The population frequencies by sex and the sample of matings from Cruz et al.

(2001) data.

	Total	RB	HY	SU
Female freqs	1254	0.22	0.11	0.67
Male freqs	1080	0.26	0.12	0.62
Matings			**Males**	
		RB	**HY**	**SU**
Females				
RB		19	9	13
HY		6	4	10
SU		6	7	80

First, I computed the information partition (Carvajal-Rodríguez 2018b) that indicated

significant assortative mating from the Chi-square test ($J_{\mathrm{PSI}}p$ value < 0.0000001) while no

significant sexual selection was detected. However, the randomization test was not

significant in any case, possible due to the low sample size within the mating classes.

Then, I proceeded with the model estimation and initially assayed only the subset of models

with male and/or female intrasexual competition plus the saturated (M_{sat}) and random

mating (M_0) models. The estimate of overdispersion was high (7.20) indicating poor

structure of the set of models regarding the data. The three information criteria gave similar

output with the M_0 as the best fit model. The multimodel estimates of the mutual-

propensities were just one in every case as expected from random mating. Because in the

simulation study the AICc criterion gave the best estimates I will rely on this criterion from

now on.

The next step was to study only models with choice parameter plus the saturated (M_{sat}) and

random mating (M_0) models. The overdispersion was 4.65 that still indicates somewhat

poor model structure. The best fit model for AICc and KICc criteria was a choice model

with one parameter, but was M_0 for BIC. The multimodel inference under AICc gave a

clear pattern of positive assortative mating, that was higher for the RB $\times$ RB mating

($m'_{RBRB} = 3$), intermediate for HY $\times$ HY ($m'_{HYHY} = 2.3$) and slightly lower for SU $\times$ SU

($m'_{SUSU} = 2$).

Then, I considered the previous competition and choice models and added new models

having separated competition and choice parameters. The overdispersion was 3.4 that is an

acceptable value for multinomial models and can be corrected by using quasi-likelihoods

(see the overdispersion section). Now, even BIC selected a choice model, the same as

KICc, while AICc selected a compound parameters model with female competition and

choice. The estimates from this model were a RB female competition of $\alpha = 1.7$ and choice

$c = 2.4$. The multi-model estimates gave positive asssortative mating, $m'_{RBRB} = 3$, $m'_{HYHY} = $

3, $m'_{SUSU} = 2$ and sexual selection favouring RB females.

Finally, I considered all the previous models plus models having parameters with double

effect. This implies a total of 35 models including M_0 and M_{sat}. The overdispersion was 2.5.

The best model was the same for the three criteria and it was a double effect model with 2

parameters $a = 0.2$ and $b = 2$ distributed as indicated in Fig. 5. Approximately, the same

model was obtained using the multi-model estimates.

$$\begin{pmatrix} 2 & 2 & 1 \\ 1 & 2 & 1 \\ 0.2 & 1 & 1 \end{pmatrix}$$

Fig 5. Model D-2P-Rep3: Double two parameter model (a =0.2, b =2) producing female and male

sexual selection plus positive assortative mating.

It is also possible to focus only on the models with separated parameters for competition

and choice. The best fit model from this subgroup involves female competition. Recall that

in *Littorina saxatilis* the choosy sex are the males, so I considered that the competitive

advantage from the side of the females is explained by the males preferring a given kind of

females. The best fit model is $S_{Fem}C$-2Pc (see Table 1) with RB female competitive

advantage of 1.7 more times matings than the other females and a choice parameter of 2.4.

The qualitative pattern obtained from this models is similar to that in Fig.5; the RB females

(first row) are preferred and there is a choice for within ecotype mating. The combination

of competition and choice provokes that the mating RB×RB is the preferred by RB males;

the matings RB×HY and HY×HY are preferred by HY males, and finally, it seems that the

SU males don't discriminate very much regarding the female ecotype.

6. Discussion

6.1 Simulations

I have simulated mating tables corresponding to random mating, intrasexual competition

and choice models. The random mating pattern was perfectly assessed. For the other

models, the competition and choice parameters were estimated quite accurately. Not

surprisingly, BIC was slightly more conservative, while AICc presented slightly more

accurate estimates in most cases. The KICc criterion performed similar to the best AICc

and BIC cases. In general, the estimation was accurate and even in the worst cases (extreme

phenotypic frequencies), the mean estimates were closer to the real value than to random

mating.

6.2 Example

Previous studies in the Galician *L. saxatilis* hybrid zone showed that mate choice favours

within-morph pairs. The estimates obtained by multimodel inference support the positive

assortative mating for the ecotype. This seems the main factor acting in the system since the

tests for sexual selection are not significant neither the set of sexual selection models are

preferred against the random mating.

In addition, another result emerged from the analysis: The RB females are preferred in

general i.e. RB male with SU female has less mutual preference than SU male with RB

female ($m_{SURB} < m_{RBSU}$). This pattern may be favoured by the physical difficulty for the

mating involving bigger RB males with the smaller SU females, and could be related with

the somehow more frequent occurrence of mating pairs having female bigger than the male

(a typical trend in gastropods, E. Rolán-Alvarez personal communication).

In addition to the mating pattern depicted by the multi-model approach, the estimates of the

mutual-propensities were also obtained. Testing the reliability of these estimates is,

however, out of the scope of the present manuscript, and it was left for future work.

6.3 General

The advantages of model selection and multimodel inference in evolutionary ecology has

been widely discussed, jointly with the pros and cons of applying any information criteria

(Link and Barker 2006; Burnham et al. 2011; Aho et al. 2014; Barker and Link 2015; Aho

et al. 2017; Dziak et al. 2019)or the reliability of the obtained estimates (Galipaud et al.

2014; Cade 2015; Giam and Olden 2016; Galipaud et al. 2017).

Multimodel inference has been however, rarely utilized to study the mating patterns that

may emerge from mate choice and intrasexual competition. Here, by developing general

models that incorporate competition and mate choice, and providing their maximum

likelihood estimates, I am proposing a standardized methodology for model selection and

multimodel inference of the mating parameters producing the sexual selection and

assortative mating patterns.

The set of *a priori* models permits to perform an *a posteriori* quantification of the data-

based evidence and provide confidence sets on plausible non-trivial models while letting

multi-model inference of the parameter values. The approach was implemented by allowing

three different information criteria. Under the scenarios assayed, they performed similarly

for simulated and real data.

The proposed framework should ideally work under scenarios where the availability of

individuals is not affected by the matings that have already occurred, as expected for

polygamous species, or even for monogamous species, when the number of available

individuals is higher that the mating pairs. However, in the case of monogamous species

with low population size, the population frequencies may be altered during the reproductive

season and so, the distribution of mutual-propensities might not define the same sexual

selection pattern over the season. This could happen because sexual selection is frequency

dependent. On the contrary, the assortative mating effects are not frequency dependent, and

should not be affected by the variation of the population frequencies (Carvajal-Rodríguez

2019).

The developed methodology has been fully implemented in a program called InfoMating

available at http://acraaj.webs6.uvigo.es/InfoMating/Infomating.htm or upon request to the

author.

Acknowledgement

I would like to thank Carlos Canchaya and Emilio Rolán-Alvarez for their valuable

comments on the manuscript. E. R-A also provided the data for the example. This work was

supported by Xunta de Galicia (Grupo de Referencia Competitiva, ED431C2016-037),

Ministerio de Economía y Competitividad (CGL2016-75482-P) and by Fondos FEDER

("Unha maneira de facer Europa").

Appendix A) Mutual Mating Propensity Models

Saturated non-random mating model: λ notation.

Consider the total number of possible mating types $K = k_1 \times k_2$ and the saturated

multinomial model for the K-1 free mating parameters m'_{ij}.

The log-likelihood function is

695
$$lnL_{sat} = C + \sum_{ij}^{K-1} n'_{ij} \ln(m'_{ij}q_{ij}) + n'_{k1k2}\ln\left(a(1 - \sum_{ij}^{K-1} q_{ij})\right) - n'\ln(M)$$

696   where n' is the number of matings in the sample and n'_{ij} is the number of matings between i-

697   type females and j-type males. I have fixed the parameter m'_{k1k2} to a.

698   Compute the first derivative of the likelihood with respect to a

699
$$\frac{dlnL_{sat}}{da} = \frac{n'_{k1k2}}{a} - \frac{n}{M}q_{k1k2}$$

then by taking $a = 1$ and equating to 0 we get

$n/M = n'_{k1k2} / q_{k1k2} \equiv \lambda(1)$

that corresponds to the number of observed matings having unity mating parameter divided

by the corresponding product of population frequencies. Under the saturated model there is

only one (for convenience m'_{k1k2}) mating parameter having unitary value and so the number

of observed matings is n'_{k1k2} and the product of the corresponding population frequencies is

$p_{1k1} \times p_{2k2} = q_{k1k2}$.

Now, let find the m'_{ij} parameter value that maximizes the likelihood

708
$$\frac{dlnL_{sat}}{dm'_{ij}} = \frac{n'_{ij}}{m'_{ij}} - \frac{n}{M}q_{ij} = 0$$

709
$$\widehat{m'_{ij}} = \frac{n'_{ij}/q_{ij}}{n/M} \equiv \frac{\lambda(m'_{ij})}{\lambda(1)}$$

710   The λ notation can be generalized for any set A of mating types having the same value of

711   propensity θ as follows

712
$$\lambda(\theta) = \frac{\sum_t^A x_t}{\sum_{t1t2}^A q_{t1t2}}$$

where x_t represents the number of mating pairs having absolute (non-normalized) mating

parameter θ and q_{t1t2} is the product of the population frequencies p_{1t1} and p_{2t2} i.e. the

expected frequency of the θ mating types under random mating.

Intrafemale competition models

The model is

$m'_{ij} = a_i$ with $a_i > 0 \ \forall \ i < k_1$ and $m'_{k1j} = a = 1 \ \forall \ j.$

There are k_1-1 independent parameters. Note that the parameters m'_{k1j} have been fixed to a

= 1. The log-likelihood function is

722
$$lnL = C + \sum_{i<k1}^{k1-1}\sum_{j=1}^{k2} n'_{ij}\ln(a_iq_{ij}) + \sum_{j=1}^{k2} n'_{k1j}\ln(aq_{k1j}) - nln(M)$$

723   Now, assume that the parameter a is not fixed and compute the first derivative of the

724   likelihood with respect to a

725
$$\frac{dlnL}{da} = \frac{\sum_j^{k2} n'_{k1j}}{a} - \frac{n}{M} \sum_j^{k2} q_{k1j} = 0$$

726   then by taking $a = 1$ and equating to 0 we get

727     $n/M = \sum_j n'_{k1j} / \sum_j q_{k1j} \equiv \lambda(1)$

728   Now find the a_i parameter value that maximizes the likelihood

729
$$\frac{dlnL}{da_i} = \frac{\sum_j^{k2} n'_{ij}}{a_i} - \frac{n}{M} \sum_j^{k2} q_{ij} = 0$$

730   Solving for a_i

731
$$\hat{a}_\iota = \frac{\sum_j^{k2} n'_{ij} / \sum_j^{k2} q_{ij}}{\frac{n}{M}} \equiv \frac{\lambda(a_i)}{\lambda(1)}$$

The formula expressed as the quotient of lambdas is valid for any number h of different

parameters, $1 \leq h < k_1$. In the particular case of having only one parameter the sum of

observed matings having propensity a_1, implies $\sum\sum n'_{ij}$ where the first summation is for all

the female types except females of type k_1, and the second is over all male types. The sum

of the product of frequencies is $1 - p_{1k1}$.

As before, $\lambda(1)$ also corresponds to the sum of the observed matings having expected

propensity 1 divided by the sum of the corresponding products of population frequencies.

The model for male sexual selection is solved in a similar way.

Intrasexual competition in both sexes

The model is

$m'_{ij} = a_i b_j, i < k_1, j < k_2; m'_{ik2} = a_i, i < k_1; m'_{k1j} = b_j, j < k_2; m'_{k1k2} = 1$

with $a_i > 0, b_j > 0 \; \forall \; i, j$.

It is easy to see that is multiplicative. Let $A = \sum_i^{k1-1} a_i p_{1i} + a p_{1k1}$ and $B =$

$\sum_j^{k2-1} b_j p_{2j} + b p_{2k2}$.

The mean mutual mating propensity is

$M = \sum_{i,j} q_{ij} m'_{ij} = \sum_i^{k1} \sum_j^{k2} a_i b_j p_{1i} p_{2j} = AB$

with $a_{k1} = a$ and $b_{k2} = b$.

The marginal propensity for *i*-type females is

$m_{Fem\_i} = a_i \sum_j^{males} p_{2j} \frac{b_j}{M} = a_i \frac{B}{M}$

Similarly, the marginal for *j*-type males

$m_{Male\_j} = b_j \frac{A}{M}$

with $a_{k1} = a$ and $b_{k2} = b$.

The condition 5-iii) for a multiplicative model implies that $m_{ij} = M_{\text{Fem\_i}} \times M_{\text{Male\_j}}$. In

addition, $m_{ij} = a_i b_j/M$ that jointly with the multiplicative condition requires $a_i b_j/M = M_{\text{Fem\_i}}$

$\times M_{\text{Male\_j}} = a_i B b_j A /M^2$ solving for M we get $M = AB$ which we have already seen it is true.

The log-likelihood function

759
$$lnL = C + \sum_{i=1}^{k1}\sum_{j=1}^{k2} n'_{ij}\ln(a_i b_j q_{ij}) - n\ln(M)$$

760 with $a_{k1} = a = 1$ and $b_{k2} = b = 1$.

761 Consider the derivatives

762
$$\frac{dA}{da_i} = p_{1i} \; ; \frac{dB}{db_j} = p_{2j}; \frac{dM}{da_i} = p_{1i}B; \frac{dM}{db_j} = p_{2j}A$$

763 Now by taking the derivative of the log-likelihood with respect to a_i or b_j and equating to 0

764 we get the estimates

765
$$\widehat{a_\iota} = \frac{\sum_j^{k2} n'_{ij} / \sum_j^{k2} q_{ij}}{\frac{n}{M}B} = \frac{\lambda(a_i + \sum_j a_i b_j)}{\lambda_{fem}(1)}$$

766
$$\widehat{b_\jmath} = \frac{\sum_i^{k1} n'_{ij} / \sum_i^{ki} q_{ij}}{\frac{n}{M}A} = \frac{\lambda(b_j + \sum_i a_i b_j)}{\lambda_{male}(1)}$$

767 Where

768
$$\frac{n}{M}B = \frac{\sum_j^{k2} n'_{k1j}}{p_{1k1}} \equiv \lambda_{fem}(1)$$

769
$$\frac{n}{M}A = \frac{\sum_i^{k1} n'_{ik2}}{p_{2k2}} \equiv \lambda_{male}(1)$$

Mate choice models with parameterized heterotypes

Consider models in which the homotype mating has absolute propensity of 1 while the

different heterotypes have absolute value of a_{ij}. The maximum likelihood estimate is

$$\hat{a}_{h1h2} = \frac{\lambda(a_{h1h2})}{\lambda(1)}$$

The number of parameters in this type of model is $K - min\{k_1, k_2\} - \Sigma_S(C_s-1)$ where the

sum is over the set of different heterotype matings and C_s is the cardinality of each set.

Double effect models

The following models generate a double pattern of sexual selection and assortative mating

even when the population frequencies are uniform.

Double effect models producing sexual selection in one sex under uniform frequencies

A simple approach consists to build a new model by setting $m'_{ii} = 1$ and $m'_{jj} = 1 + c$. Then, if

we desire assortative mating jointly with sexual selection only in females we additionally

set $m'_{ij} = 1 - c$; on the contrary, if we desire selection only in males we set $m'_{ji} = 1 - c$ with -

$1 < c < 1$. If the frequencies are not uniform the model generates assortative mating jointly

with sexual selection in both sexes.

In the case of the model with $m'_{ij} = 1 - c$ (female sexual selection if frequencies are

uniform) the maximum likelihood estimate of c is one of the roots of the quadratic

$$(x_{jj} - x_{ij} + n'D) - c[x_{ij} + x_{jj} + D(x_{jj} - x_{ij})] - c^2D[n' - (x_{ij} + x_{jj})] = 0$$

where $D = q_{ij} - q_{jj}$ and $n' = \sum x_{ij}$ is the number of matings (sample size).

If the frequencies are uniform and $k_1 = k_2$, i.e. $p_{1i} = p_{1j} = p_{2i} = p_{2j} \ \forall \ i, j$ then

$$\hat{c} = \frac{x_{jj} - x_{ij}}{x_{jj} + x_{ij}}$$

The case for male sexual selection is obtained simply by interchanging x_{ij} by x_{ji} and q_{ij} by

q_{ji} in the formulas.

The above model has only one parameter c; we can introduce a more complex two

parameter model, $M_{(a,c)}$ by setting $m'_{ii} = a$, $m'_{jj} = 1 + c$ and $m'_{ij} = 1 - c$, for female sexual

selection (or $m'_{ji} = 1 - c$ for male sexual selection). For obtaining the MLE of this two

parameter double model, with restrictions $a > 0$, $c < |1|$, I have used a numerical bounded

Nelder-Mead simplex algorithm (Press 2002; Singer and Singer 2004; Gao and Han 2012).

Double effect models with sexual selection in both sexes under uniform frequencies

To get assortative mating jointly with sexual selection in both sexes under uniform

frequencies, we just need to combine the above uniform one parameter models of each sex,

so that $m'_{ii} = 1$, $m'_{jj} = 1 + c$ and $m'_{ij} = m'_{ji} = 1 - c$.

The maximum likelihood estimate of c involves the solution of the quadratic

$[x_{jj} - x_s + n'D_2] - c[x_{jj} + x_s + D_2(x_{jj} - x_s)] + c^2 D_2[x_{jj} + x_s - n'] = 0$

where $x_s = x_{ij} + x_{ji}$ and $D_2 = q_{ij} + q_{ji} - q_{jj}$.

General double effect models

We can also define a set of general models where any propensity m'_{ij} has parameter θ_{ij} with

at least one propensity having value of 1. The MLE of the parameters of this kind of model

is

$\hat{\theta}_{ij} = \frac{\lambda(\theta_{ij})}{\lambda(1)}$

where $\lambda(\theta_{ij})$ is defined as in (A2).

The simplest model defined in this way is

$m'_{ii} = a$ and $m'_{ij} = m'_{ji} = m'_{jj} = 1$,

which produces assortative mating and sexual selection in both sexes.

Consider as an example of this model, the case with $k_1 = k_2 = 2$ so that $0 < p_{11} < 1$; $0 < p_{21} <$

1; $m'_{11} = a$ and $m'_{12} = m'_{21} = m'_{22} = 1$. The mean mating propensity is $M = q_{11}(a - 1) + 1$. The

absolute marginal propensity for the first female type $m'_{Fem\_1} = a\, p_{21} + 1 - p_{21} = p_{21}(a - 1)$

$+1$, and for the second female type $m'_{Fem\_2} = 1$. Similarly the male marginals are $m'_{Male\_1} =$

$p_{11}(a - 1) + 1$ and $m'_{Male\_2} = 1$.

Recall that the condition for the sexual selection pattern within a given sex is that the

marginal mating propensities are different which here is true for both sexes provided that a

$\neq 1$. Regarding the assortative mating pattern it can be proved that the joint isolation index

(I_{PSI}) is 0 only if $a = 1$. However, it is sufficient to prove that the model is not multiplicative

(Carvajal-Rodríguez 2018b). Consider that the model is multiplicative, this implies, m'_{12} / M

$= (m'_{Fem\_1} / M) \times (m'_{Male\_2} / M)$ that given the model values becomes

$$M = (m'_{Fem\_1}) \times (m'_{Male\_2})$$

which can be true only when $p_{11} = 1$ and so it is false by definition.

The estimate of a under this model is $\lambda(a) / \lambda(1)$.

The most parameterized model that can be defined in this way has K-1 free parameters and

coincides with the saturated model so that the estimates are the corresponding pair total

indices (PTI_{ij}).

Moreover, note that if no mutual propensity is fixed to 1 then $\lambda(1) = (n - A) / (1 - P) = n$

where A = number of observations having value 1 = 0 and P = product of population

frequencies of the involved types having mutual propensity 1 = 0. Therefore the estimate of

θ_{ij} can also expressed as $\lambda(\theta_{ij}) / n$ which is the observed frequency of mating pairs (i, j)

divided by the expected frequency by random mating which is the definition of the pair

total index PTI_{ij} (K-1 are free and one PTI is dependent on the others).

All the above derived MLE formulae have been checked by a numerical bounded Nelder-

Mead simplex algorithm (Press 2002; Singer and Singer 2004; Gao and Han 2012).

Appendix B) Monte Carlo simulation of mating tables

The mating tables for the simulation experiments were generated by the program MateSim

(Carvajal-Rodríguez 2018a) available at http://acraaj.webs.uvigo.es/MateSim/matesim.htm.

The number of replicates for each case was 1000. For each run the program first generated

the number of premating males and females from a given population size. For example, if

the population size consisted in n_1 (= 5,000) females and n_2 (= 5,000) males, the program

got $n_{1A} = n_1 \times U$ females of the A type and $n_{1B} = n_1 - n_{1A}$ females of the B type. Where U is

a value sampled from the standard uniform distribution. The premating males were

obtained similarly. Then, the female population frequencies were $p_{1i} = n_{1i} / n_1$, and $p_{2i} = n_{2i}$

/ n_2 for the male ones. Finally, a sample of n' (= 500) matings was obtained, where the

number of counts for each mating type $i \times j$ was

856            $Q(i,j) = n' \times p_{1i} \times p_{2j} \times m'_{ij} / M$

857   where m'_{ij} are the mutual-propensity parameters as defined for each kind of model, and $M =$

858    $\sum p_{1i} \times p_{2j} \times m'_{ij}$.

859   The format of the obtained tables was the same as the JMating (Carvajal-Rodriguez and

860   Rolan-Alvarez 2006) input files (Fig. B1).

```
# format number
0
# num of types
2
#premating male numbers
1626        3374
#premating female numbers
917     4083
# matings by rows (females)
30      62
133     275
```

862 **Fig. B1.** Example of a table generated by the simulations. The format is the same as for the JMating
863 software.

References

Aho, K., D. Derryberry, and T. Peterson. 2014. Model selection for ecologists: the
worldviews of AIC and BIC. Ecology 95:631-636.
Aho, K., D. Derryberry, and T. Peterson. 2017. A graphical framework for model selection
criteria and significance tests: refutation, confirmation and ecology. Methods in
Ecology and Evolution 8:47-56.
Akaike, H. 1973. Information theory and an extension of the maximum likelihood
principle. Pp. 267-281 *in* B. N. Petrov, and F. Csaki, eds. Second International
Symposium on Information Theory, Budapest: Akademiai Kiado.
Anderson, D. R., K. P. Burnham, and G. C. White. 1994. AIC Model Selection in
Overdispersed Capture-Recapture Data. Ecology 75:1780-1793.
Barker, R. J. and W. A. Link. 2015. Truth, models, model sets, AIC, and multimodel
inference: A Bayesian perspective. The Journal of Wildlife Management 79:730-
738.
Brewer, M. J., A. Butler, and S. L. Cooksley. 2016. The relative performance of AIC,
AICC and BIC in the presence of unobserved heterogeneity. Methods in Ecology
and Evolution 7:679-692.
Burnham, K. P. and D. R. Anderson. 2002. Model selection and multimodel inference: a
practical information-theoretic approach. Springer-Verlag, New York, NY.
Burnham, K. P., D. R. Anderson, and K. P. Huyvaert. 2011. AIC model selection and
multimodel inference in behavioral ecology: some background, observations, and
comparisons. Behavioral Ecology and Sociobiology 65:23-35.
Cade, B. S. 2015. Model averaging and muddled multimodel inferences. Ecology 96:2370-
2382.

Carvajal-Rodríguez, A. 2018a. MateSim: Monte Carlo simulation for the generation of
mating tables. Biosystems 171:26-30.
Carvajal-Rodríguez, A. 2018b. Non-random mating and information theory. Theoretical
Population Biology 120:103-113.
Carvajal-Rodríguez, A. 2019. A generalization of the informational view of non-random
mating: Models with variable population frequencies. Theoretical Population
Biology 125:67-74.
Carvajal-Rodriguez, A. and E. Rolan-Alvarez. 2006. JMATING: a software for the analysis
of sexual selection and sexual isolation effects from mating frequency data. BMC
Evol Biol 6:40.
Casares, P., M. C. Carracedo, B. del Rio, R. Piñeiro, L. Garcia-Florez, and A. R. Barros.
1998. Disentangling the Effects of Mating Propensity and Mating Choice in
Drosophila. Evolution 52:126-133.
Cavanaugh, J. E. 2004. Criteria for linear model selection based on Kullback's symmetric
divergence. Australian & New Zealand Journal of Statistics 46:257-274.
Claeskens, G. 2016. Statistical Model Choice. Annual Review of Statistics and Its
Application 3:233-256.
Cruz, R., E. Rolán-Álvarez, and C. García. 2001. Sexual selection on phenotypic traits in a
hybrid zone of Littorina saxatilis (Olivi). Journal of Evolutionary Biology 14:773-
785.
Dziak, J. J., D. L. Coffman, S. T. Lanza, R. Li, and L. S. Jermiin. 2019. Sensitivity and
specificity of information criteria. bioRxiv:449751.
Galipaud, M., M. A. F. Gillingham, M. David, F. X. Dechaume-Moncharmont, and R. B.
O'Hara. 2014. Ecologists overestimate the importance of predictor variables in
model averaging: a plea for cautious interpretations. Methods in Ecology and
Evolution 5:983-991.
Galipaud, M., M. A. F. Gillingham, and F.-X. Dechaume-Moncharmont. 2017. A farewell
to the sum of Akaike weights: The benefits of alternative metrics for variable
importance estimations in model selection. Methods in Ecology and Evolution
8:1668-1678.
Gao, F. and L. Han. 2012. Implementing the Nelder-Mead simplex algorithm with adaptive
parameters. Computational Optimization and Applications 51:259-277.
Gavrilets, S. 2004. Fitness landscapes and the origin of species. Princeton University Press,
Princeton, N.J.
Giam, X. and J. D. Olden. 2016. Quantifying variable importance in a multimodel inference
framework. Methods in Ecology and Evolution 7:388-397.
Grueber, C. E., S. Nakagawa, R. J. Laws, and I. G. Jamieson. 2011. Multimodel inference
in ecology and evolution: challenges and solutions. Journal of Evolutionary Biology
24:699-711.
Hartl, D. L. and A. G. Clark. 1997. Principles of Population Genetics. Sinauer Associates,
Inc., Sunderland, MA.
Jennions, M. D. and M. Petrie. 1997. Variation in mate choice and mating preferences: a
review of causes and consequences. Biological Reviews 72:283.
Keerativibool, W. 2014. Unifying the Derivations of Kullback Information Criterion and
Corrected Versions. Thailand Statistician 12:37-53.

Kim, H.-J., J. E. Cavanaugh, T. A. Dallas, and S. A. Foré. 2014. Model selection criteria for
overdispersed data and their application to the characterization of a host-parasite
relationship. Environmental and ecological statistics 21:329-350.
Knoppien, P. 1985. Rare male mating advantage: a review. Biological Reviews 60:81-117.
Lewontin, R., D. Kirk, and J. Crow. 1968. Selective mating, assortative mating, and
inbreeding: definitions and implications. Eugen Q 15:141-143.
Link, W. A. and R. J. Barker. 2006. Model weights and the fundations of multimodel
inference. Ecology 87:2626-2635.
Liu, W. and Y. Yang. 2011. Parametric or nonparametric? A parametricness index for
model selection. The Annals of Statistics 39:2074-2102.
Merrell, D. J. 1950. Measurement of sexual isolation and selective mating. Evolution:326-
331.
Press, W. H. 2002. Numerical recipes in C++ : the art of scientific computing. Cambridge
University Press, Cambridge.
Richards, S. A. 2008. Dealing with overdispersed count data in applied ecology. Journal of
Applied Ecology 45:218-227.
Rolan-Alvarez, E. 2007. Sympatric speciation as a by-product of ecological adaptation in
the Galician Littorina saxatilis hybrid zone. Journal of Molluscan Studies 73:1-10.
Rolan-Alvarez, E., C. Austin, and E. G. Boulding. 2015a. The contribution of the genus
Littorina to the field of evolutionary ecology. Oceanography and Marine Biology:
an Annual Review 53:157-214.
Rolán-Alvarez, E. and A. Caballero. 2000. Estimating sexual selection and sexual isolation
effects from mating frequencies. Evolution 54:30-36.
Rolan-Alvarez, E., A. Carvajal-Rodriguez, A. de Coo, B. Cortes, D. Estevez, M. Ferreira,
R. Gonzalez, and A. D. Briscoe. 2015b. The scale-of-choice effect and how
estimates of assortative mating in the wild can be biased due to heterogeneous
samples. Evolution 69:1845-1857.
Schwarz, G. 1978. Estimating the dimension of a model. The Annals of Statistics 6:461-
464.
Singer, S. and S. Singer. 2004. Efficient Implementation of the Nelder–Mead Search
Algorithm. Applied Numerical Analysis & Computational Mathematics 1:524-534.
Spieth, H. T. and J. M. Ringo. 1983. Mating behavior and sexual isolation in Drosophila.
Genetics and biology of Drosophila.
Symonds, M. R. E. and A. Moussalli. 2011. A brief guide to model selection, multimodel
inference and model averaging in behavioural ecology using Akaike's information
criterion. Behavioral Ecology and Sociobiology 65:13-21.
Vrieze, S. I. 2012. Model selection and psychological theory: a discussion of the
differences between the Akaike information criterion (AIC) and the Bayesian
information criterion (BIC). Psychological methods 17:228.
Wagenmakers, E.-J. 2007. A practical solution to the pervasive problems of p values.
Psychonomic bulletin & review 14:779-804.

975