Separate the wheat from the chaff: genomic analysis of local adaptation in

the red coral *Corallium rubrum* [1]

1. We have been informed by F. Allendorf that another article had a very similar title: "Waples, R. S. 1998.

Separating the wheat from the chaff: Patterns of genetic differentiation in high gene flow species. Journal of

Heredity 89:438-450."

AUTHORS

Pratlong M.[1,2]•, Haguenauer A.[1], Brener K.[3], Mitta G.[3], Toulza E.[3], Garrabou J.[4,5],

Bensoussan N.[6], Pontarotti P.[2,7,8]* & Aurelle D[1,5,9]*•

AFFILIATIONS

1. Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE, Marseille, France
2. Aix Marseille Univ, CNRS, Centrale Marseille, I2M, Marseille, France, Equipe Evolution
Biologique et Modélisation, Marseille, France
3. Perpignan Via Domitia Univ., IHPE UMR 5244, CNRS, IFREMER, Montpellier Univ.,
Perpignan, France
4. Institute of Marine Sciences (ICM-CSIC), Barcelona, 08003, Spain
5. Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO UM 110, 13288, Marseille,
France
6. IPSO FACTO, SCOPArl, Pole Océanologie, Marseille, 13001, France
7. Aix Marseille Univ, IRD, APHM, Microbe , Evolution, PHylogénie, Infection IHU
Méditerranée Infection,Marseille France
8. CNRS
9 Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire
naturelle, CNRS, Sorbonne Université, EPHE, 57 rue Cuvier, 75005 Paris, France
27
28   * These authors jointly supervised this work.

29   • Corresponding authors: marine.pratlong@gmail.com / didier.aurelle@univ-amu.fr
30

A colony of red coral, *Corallium rubrum***, near Marseille. Photo: F. Zuberer / OSU**

Pythéas / CNRS

ABSTRACT

Genomic data allow an in-depth and renewed study of local adaptation. The red coral

(*Corallium rubrum*, Cnidaria) is a highly genetically structured species and a promising

model for the study of adaptive processes along an environmental gradient. Here, we used

RAD-Sequencing in order to study the vertical genetic structure of this species and to search

for signals of local adaptation to depth and thermal regime in the red coral. Previous studies

have shown different thermotolerance levels according to depth in this species which could

correspond to genetic or environmental differences. We designed a sampling scheme with six

pairs of 'shallow vs deep' populations distributed in three geographical regions as replicates.

Our results showed significant differentiation among locations and among sites separated by

around 20 m depth. The tests of association between genetics and environment allowed the

identification of candidate loci under selection but with a potentially high rate of false

positive. We discuss the methodological obstacles and biases encountered for the detection of

selected loci in such a strongly genetically structured species. On this basis, we also discuss

the significance of the candidate loci for local adaptation detected in each geographical region

and the evolution of red coral populations along environmental gradients.

INTRODUCTION

The study of the mechanisms of adaptation by species to their local environment is of great

interest in evolutionary biology. The interaction between environmental conditions, biological

traits and evolutionary factors (selection, drift, migration and mutation) will shape the relative

importance of genetic and plastic responses for each species facing heterogeneous

environmental conditions. If selection is predominant, and if the environmental gradient is

persistent for an extended period of time, each local population exposed to local selection

could become genetically adapted to the corresponding local environment (Kawecki and

Ebert, 2004; Gagnaire and Gaggiotti, 2016). An organism can also cope with local

environmental conditions via plasticity or acclimatization, whereby a given genotype

develops during its lifetime morphological or physiological responses (DeWitt *et al.*, 1998;

Pigliucci, 2001). Although particular situations favoring local adaptation or acclimatization

are documented, it is often difficult to disentangle the effects of these two mechanisms and

establish their relative contributions to adaptability (Palumbi *et al.*, 2014). In addition,

understanding these mechanisms has a fundamental interest in the current context of climate

change for improving predictive models and proposing management strategies (Mumby et al.,

2011; Gagnaire and Gaggiotti, 2016).

The evolution of adaptive processes depends on the interaction between different processes,

including local selection and gene flow. Gene flow can hinder local adaptation through the

input in a population of potentially maladapted individuals (migration load; Lenormand,

2002). Conversely, several theoretical studies have shown that gene flow can counteract the

effects of genetic drift and promote local adaptation (Hastings and Rohlf, 1974; Felsenstein,

1975; Slatkin and Maruyama, 1975; Nagylaki, 1978; Alleaume-Benharira *et al.*, 2006). The

use of high throughput sequencing renewed the study of local adaptation. Various *bottom – up*

approaches are now available to study local adaptation through the identification signals of

selection along the genome (Barrett and Hoekstra, 2011). In the marine realm, such studies

have been conducted at very large scale on highly dispersive teleost species (Bradbury *et al.*,

2010; Limborg *et al.*, 2012; Wang *et al.*, 2013; Milano *et al.*, 2014; Bernardi *et al.*, 2016; Guo

et al., 2016), and on benthic invertebrates with a highly dispersive, planctonic larvae stage

(Chu *et al.*, 2014; Bay and Palumbi, 2014; Araneda *et al.*, 2016; Benestan *et al.*, 2016).

8   4

Marine species with strong genetic structure are less frequent than more dispersive ones, and

genomic studies of local adaptation in such species are still scarce (see Bongaerts *et al.*, 2017

for a recent example). The study of local adaptation in a context of strong genetic structure

may also be difficult from a methodological point of view: high average F_{ST} values can lead to

a high number of false positives in outlier tests for the detection of selection by the

corresponding increase in the variance of F_{ST} values (Bierne *et al.*, 2013; Hoban *et al.*, 2016).

Furthermore, in a context of high average genomic differentiation, it could be difficult to

identify selected loci with a higher differentiation than expected under the neutral model.

Finally, if genetic drift is strong, it can generate outlier loci with apparent correlation with an

environmental variable outside any selective effect (Kawecki and Ebert, 2004; Hofer *et al.*,

2009; Coop *et al.*, 2010). Therefore the empirical study of local adaptation in such situation

remains often challenging and with few empirical data in the marine realm.

Marine coastal environments offer particularly interesting conditions for studies of local

adaptation, because of the gradual changes in environmental conditions along coastline at

small scale, the more or less gradual vertical changes from shallow to deep water and the

patchy distribution of contrasted habitats at different scales (Sanford and Kelly, 2011;

Lundgren *et al.*, 2013; Wrange *et al.*, 2014). This interest promoted studies of local adaptation

in coastal ecosystems (Ayre, 1995; Ulstrup and Van Oppen, 2003; Smith et al., 2007; Sherman

and Ayre, 2008; Barshis *et al.*, 2010; Bongaerts *et al.*, 2011; Barshis *et al.*, 2013; Lundgren *et*

al., 2013; Kersting *et al.*, 2013; Haguenauer *et al.*, 2013; Ziegler *et al.*, 2014; Palumbi *et al.*,

2014; Bay and Palumbi, 2014; Ledoux *et al.*, 2015; Pivotto *et al.*, 2015; Jin *et al.*, 2016;

Bongaerts *et al.*, 2017). Studying the genetic basis of local adaptation and the connectivity

between habitats, could also give some information on the response to climate change (e.g.

Bongaerts *et al.*, 2017). In this context genome scans are powerful approaches to explore

10   5

adaptive processes in natural populations (Manel *et al.*, 2016).

The red coral (*Corallium rubrum*) is an asymbiotic (without *Symbiodinium*) temperate

octocoral distributed from 5 to 1016 m depth in the Mediterranean sea and the near Atlantic

(Boavida *et al.*, 2016; Knittweis *et al.*, 2016). It is a sessile and long-living species (more than

100 years), with low growth and recruitments rates (Marschal *et al.*, 2004; Santangelo *et al.*,

2012). This species is included in two international conventions (annex III of the Bern

Convention and annex V of the European Union Habitats directive), and its harvesting is

regulated by national legislations (Ledoux *et al.*, 2013). The study of a few microsatellite loci

has demonstrated a strong genetic structure in this species (Ledoux *et al.*, 2010a; Ledoux, *et*

al., 2010b). The shallowest populations, above the seasonal thermocline, are exposed to high

maximum temperatures and to frequent and intense thermal fluctuations in summer

(Haguenauer *et al.*, 2013). The intensity and frequency of extreme thermal events decrease

with depth, and the deepest populations are exposed to stable thermal regimes. Since the

observation of mass mortality events affecting this species during thermal anomalies in 1999

and 2003, the thermotolerance of the red coral has been intensively studied in the region of

Marseille (France; Garrabou *et al.*, 2001, 2009). Common garden experiments highlighted

differences in polyp activities, calcification rate, necrosis rate and expression of HSP70

between shallow and deep individuals (10 or 20 m compared to 40 m depth) facing thermal

stress (Torrents *et al.,* 2008 ; Ledoux *et al.* 2015; Haguenauer *et al.,* 2013). Transcriptomes of

individuals from 5 and 40 m were compared and several genes were detected as differentially

expressed without the application of any stress (Pratlong *et al.*, 2015). These results suggested

the possibility of local adaptation to depth in this species, but the possibility of environmental

effects could not be excluded.

Together, these studies highlighted phenotypic differences in thermotolerance levels between

individuals from different depths in Marseille, with shallower individuals more tolerant than

deeper ones. Nevertheless we still do not know if these differences are the result of local

adaptation or of individual acclimatization, or both. Previous works on this species enabled us

to have a precise idea of the geographic scale at which local adaptation may occur, and were

useful to optimize our sampling design. Because populations from different regions may have

evolved similar responses to thermal stress, through similar or different genetic basis, it is

interesting to investigate local adaptation in pairs of 'shallow vs deep' populations exposed to

contrasted thermal regimes in distinct geographical regions (Jones *et al.*, 2012; Hoban *et al.*,

2016). Finally, the study of the genetic structure of this species would be useful to better

understand the potential role of deeper populations in reseeding shallower ones following

disturbances (Bongaerts *et al.*, 2017).

Here we applied Restriction site Associated DNA sequencing (RAD-Seq) to individuals from

pairs of 'shallow *vs.* deep' populations in three geographical regions of the Mediterranean Sea.

The goal of this study was to characterize the neutral and adaptive genomic variation in this

species and to test the possibility of local adaptation to depth through a genome scan

approach. Our results enable us to discuss the neutral genetic structure of the red coral. Then

we highlight the methodological obstacles expected in the detection of local adaptation in this

context. Finally, we discuss the robustness of the candidates of local adaptation detected in

each geographical region.

MATERIAL AND METHODS

Sampling and DNA extraction

Corallium rubrum colonies were collected by scuba diving at two depths of two sites in three

geographical regions (Marseille, Banyuls, Corsica) between February and August 2013

(Fig. S1, Table 1). Red coral populations from these three regions correspond to different

genetic clusters according to microsatellites (Ledoux *et al.*, 2010b) and RAD-Seq (see

results). The two depths of each site presented contrasted thermal regimes with higher mean,

maximum and standard deviation of temperature at shallower depths (surveys from March

2012 to October 2014; Table 2). Samples from the two depths at each site will be referred as

shallow and deep. The three geographical regions presented different annual variations of

temperatures between the two studied depths: a difference of 3.8 °C between the maximum

observed at the two depths in Marseille, 1.7 °C in Corsica and 0.5 °C in Banyuls (Table 2).

Thirty individuals per site and depth were collected (total 360 individuals), preserved in 95 %

ethanol and stored at -20 °C until DNA extraction. Total genomic DNA was extracted

according to the protocol of Sambrook *et al.* (1989), followed by a purification using Qiagen

DNeasy blood and tissue spin columns (Qiagen). Genomic DNA concentration was quantified

using a Qubit 2.0 Fluorometer (Life Technologies).

RAD-Sequencing

Twelve RAD libraries were prepared according to the protocol described in Etter *et al.* (2011),

with small modifications. Briefly, 1 μg of genomic DNA for each sample was digested using

high-fidelity PstI during 60 min at 37 °C. P1 adapters, with 4-6 bp individual barcodes were

then ligated to each sample using 0.5 μL of T4 DNA ligase (NEB), 0.5 μL of rATP 100 mM

(Promega), 1 μL of DTT 500 mM (Promega), 1 μL of 10X T4 ligase buffer (NEB) and

incubated during 60 min at 22 °C, 10 min at 65 °C and 1 min at 64 °C. Individual samples

were pooled by 32 (generally by location), sheared, size selected and P2-barcoded. Final PCR

for RAD-tags enrichment were performed with 16 cycles and primers dimers were removed

during a final AMPure Beads Purification (Agencourt). Libraries were sequenced on an

Illumina HiSeq2000 using 100 bp single-end reads, at the Biology Institute of Lille (IBL,

UMR 8199 CNRS) and at the MGX sequencing platform in Montpellier (France).

The STACKS pipeline (Catchen *et al.*, 2011, 2013) was used for the loci *de novo* assembly

and genotyping. Quality filtering and demultiplexing were performed with the

process_radtags module with default parameters which enables to remove any read with

uncalled base and to perform a phred-33 quality filtering of raw reads. Exact-matching RAD

loci (putative orthologous tags) were individually assembled using *ustacks* with a minimum

depth of coverage of five reads per allele (m = 5) and a maximum of five nucleotide

mismatches between allele (M = 5). These parameters were optimized during preliminary

runs. *Cstacks* was used to build a catalog of consensus loci from all individuals, with five

mismatches allowed between individuals at the same locus (n = 5). Matches of individual

RAD loci to the catalog of loci were searched using *sstacks*. Finally, the *population* module

was used to obtain the loci that were successfully genotyped in at least 75 % of individuals

from all populations. We observed an increase in the number of SNPs from position 86 bp to

91 bp and we removed these positions from the analysis which were due to sequencing

problems. In order to filter for poor-quality SNPs and artifacts due paralogous sequences, we

used VCFtools (Danecek *et al.*, 2011) to remove SNPs that were not at the Hardy-Weinberg

equilibrium within at least one of the 12 populations with a p-value threshold of 0.01. SNPs

with a minor allele frequencies below 0.01 were removed using VCFTools. Individuals with

more than 30 % of missing genotypes were discarded. Finally, only the first SNP of each

RAD locus was kept for further analysis. The whole dataset has been previously used for the

study of sex determinism in *C. rubrum* (Pratlong *et al.*, 2017); we develop here the study of

genetic structure and local adaptation.

Diversity and neutral genetic structure

Global F_{IS} over alleles and gene diversity (expected heterozygosity) parameters were

estimated using GENEPOP and ARLEQUIN v.3.5 (Rousset, 2008; Excoffier and Lischer,

2010). Differences in diversity levels between regions were tested with a Kruskal-Wallis test

by using the package agricolae in R (R Core Team, 2016). The *C. rubrum* genetic structure

was first analyzed by principal component analysis (PCA) using the R package adegenet

(Jombart, 2008) . This analysis was performed on the total dataset (12 populations) and inside

each of the three studied geographical regions (four populations in each). The dataset was

centered (they have been modified to a mean of zero) and missing data were replaced by the

mean allele frequency for each locus (http://adegenet.r-forge.r-project.org/files/tutorial-

basics.pdf). In a second step, we performed a Bayesian population clustering implemented in

the program STRUCTURE v.2.3.4 (Pritchard *et al.*, 2000; Falush *et al.*, 2003, 2007; Hubisz *et*

al., 2009). We performed ten independent replicates from $K = 1$ to 10 with a burn-in of 50 000

and a number of MCMC iterations after burn-in of 100 000, with the model allowing for

admixture and correlated allele frequencies. We calculated the ΔK statistic of Evanno *et al.*

(2005) to help in the choice of the most appropriate number of genetic clusters but we also

considered different K values. We used CLUMPAK to summarize the STRUCTURE results

from the ten independent runs (Kopelman *et al.*, 2015). The global and pairwise populations

F_{ST} and exact tests for population differentiation were computed with GENEPOP 4.0.10

(Rousset, 2008). The correlation between the spatial distance between the two depths of the

same site and the corresponding population pairwise F_{ST} was tested with the correlation test of

Spearman implemented in R (R Core Team, 2016). Finally, we conducted an analysis of

molecular variance (AMOVA) in ARLEQUIN v.3.5 (Excoffier and Lischer, 2010) with

10 000 permutations. The hierarchy for this analysis was chosen to follow the three

geographical regions of our samples (Marseille, Corsica and Banyuls). This choice was

justified by the PCA on the overall dataset. Finally, we performed the PCA and F_{ST} calculation

using a dataset comprising only putatively neutral SNPs (without the SNPs detected as

outliers by F_{ST} outlier methods, see below).

Detection of local adaptation

In order to search for loci potentially involved in local adaptation, we first used BayeScEnv

(Villemereuil and Gaggiotti, 2015). This method identifies F_{ST} outlier loci that show a

relationship between genetic differentiation and environmental differentiation. Runs were

performed using default parameters, except the number of pilot runs that was set at 40. The

maximal temperature recorded in each site was used as environmental variable (Table 2). We

tested other descriptors of the thermal regime and we got similar results (data not shown). The

convergence of runs was checked with the Gelman and Rubin's diagnostic using the R

package coda (Plummer *et al.*, 2006).

Second, we searched for F_{ST} outliers among red coral populations using ARLEQUIN v.3.5

(Hofer *et al.*, 2009; Excoffier and Lischer, 2010). Because hierarchical genetic structures are

known to lead to a high number of false positives in the search of outlier loci (Hofer *et al.*,

2009), we performed this analysis independently in the three geographical regions in order to

down a level in the structure. With this method, a distribution of F_{ST} across loci as a function

of heterozygosity between populations is obtained by performing simulations under a

hierarchical island model (two depths in one site and two sites in one geographical region).

Outliers were identified as loci being in the tails of the generated distribution ($p < 0.01$).

Outliers detected by ARLEQUIN could be false positives or the result of a selective pressure

independent of depth. Therefore, we selected among these candidate loci, those linked with

depth differentiation by searching, inside each geographical regions, loci with significant

differences in genotypic frequencies between depths according to a Chi² test. Significant

differences were identified by using a false discovery rate of 0.05 (Benjamini and Hochberg,

1995).

Finally, we used the R package pcadapt to search for outliers loci by taking into account

population structure and individual admixture (Luu *et al.*, 2017). This method is

recommended in cases of hierarchical genetic structure for a better control of the false

positive rate. By identifying outliers loci linked with a particular principal component,

pcadapt enabled us to focus on candidates linked with our biological question. From the

pcadapt analyses, we selected outliers candidates linked with the relevant principal

components with a q-value cutoff of 0.01.

Functional annotation and enrichment tests

The RAD tags were aligned on the red coral transcriptome (Pratlong *et al.*, 2015) using the

Burrows-Wheeler Alignment Tool (BWA; Li and Durbin, 2009). Blast2GO was used for the

annotation of resulting contigs and functional enrichment analysis (Conesa *et al.*, 2005). First,

a blastp was first performed on the NCBI nr database with an e-value threshold of 10^{-10}

(Altschul *et al.*, 1990). Then, Blast2GO retrieved Gene Ontology (GO) terms associated with

the obtained BLAST hits. Finally, in order to identify function potentially over-represented in

outliers, we performed an enrichment analysis using a Fisher's exact test corrected using a

false discovery rate of 0.05 (Benjamini and Hochberg, 1995).

RESULTS

RAD-Sequencing and genotyping

An average of 191 ± 21 millions of reads by library was obtained after sequencing. After the

demultiplexing and cleaning processes of the STACKS's *process_radtags* module, an average

of 180 ± 22 millions of reads by library was obtained. From these reads, we were able to

assemble 138 810 unique consensus RAD-tags present in at least 75 % of our 360

individuals. After all quality filter steps (Table 3), 27 461 SNPs were available. Finally, we

removed six individuals presenting more than 30 % of missing data (one individual from the

MEJ40 population, two from the BANN40 population, two from the GAL20 population and

one from the GAL40 population). Our final dataset used for further analysis consisted in 359

individuals genotyped on 27 461 SNPs.

Genetic diversity

Multilocus values of the F_{IS} ranged between 0.005 (ELV12) and 0.065 (BANS40) (Table 4).

Gene diversity varied from 0.12 (GAL20) to 0.18 (all populations of Marseille) (Table 4).

There was no significant difference in nucleotide diversity between regions (Kruskal-Wallis

test p = 0.17), but there was a significant difference in gene diversity (p = 0.01) with post-hoc

comparisons indicating a higher diversity in Marseille compared to Banyuls and Corsica , nut

no difference between Banyuls and Corsica.

Population structure analysis

The positioning of individuals with respect to the first two principal components reflected the

geographical, and, partly, the depth origin of the individuals (Fig. 1A). Individuals from

Marseille and from Banyuls formed two clear and homogeneous groups while individuals

from the two sites of Corsica formed two different groups with an important distance between

them on the second axis. The first PCA axis explained 7.28 % of the total genotypic variance

and separated individuals from Marseille from individuals from Banyuls and Corsica. The

second axis explained 4.4 % of the total genetic variance and separated individuals from the

Porto site in Corsica from other individuals. The fifth axis of the PCA separated all

individuals according to their sex, independently from their geographical origin, as observed

in a previous study (Pratlong et al., 2017). This axis can be explained by loci diagnostic of

sex, heterozygous in females and homozygous in males (see Pratlong et al., 2017).

Concerning PCA inside geographical regions, individuals from the two sites of Corsica and

Marseille (north and south) were separated along the first axis (13.41 % and 6.74 % of the

total genetic variance respectively; Fig. 1B and 1C). The second axis (2.77 % of the total

genetic variance) separated populations from the two depths of the two sites of Marseille. In

Corsica and Banyuls, no PCA axis showed clear association with depth. Individuals from the

two depths of the Galeria (GAL) site of Corsica were separated along the second axis (4.02 %

of the total genetic variance) but this was not the case for individuals from the two depths of

the Porto site (POR). Individuals from Banyuls showed much less structure than individuals

from Marseille and Corsica (Fig. 1D). The first axis (2.99 % of the total genetic variance)

separated individuals from the two sites (north and south). The second axis (2.46 % of the

total genetic variance) separated individuals according to their sex (Pratlong *et al.*, 2017). The

PCA on the overall dataset and inside each geographical region gave similar results when only

putatively neutral SNPs were considered (Fig. S2).

The delta(K) criterion (Evanno *et al.*, 2005) indicated K = 2 as the most informative number

of clusters for the STRUCTURE analysis. We present here the results for K =2 to K =4 which

captured the main information (Fig. 2). In all cases, all clusters corresponded to the main

geographical boundaries and the two depths of each site always clustered together. For K = 2,

a clear separation between the Marseille regions and the Corsica / Banyuls regions was

observed, confirming the separation of populations along the first PCA axis (Fig. 1). The

clustering at K = 3 separated the three geographical regions in 7/10 replicates, and the

remaining replicates grouped either one or the other Corsican sites with Banyuls populations

(Fig. S3). Finally, K = 4 separated the two Corsican sites.

The overall multilocus F_{ST} of the total dataset was 0.13. Pairwise F_{ST} values ranged from 0.01

(BANS20 vs BANS40, BANS20 vs BANN40 and BANN40 vs BANS40) to 0.24 (ELV12 vs

GAL20 and FIG8 vs GAL20; Table 5). The exact test of genetic differentiation was highly

significant for all pairwise comparisons (p < 0.001), even for the Banyuls and Galeria

populations separated by 10 m depth (F_{ST} = 0.022 for BANN20 vs BANN40, F_{ST} = 0.012 for

BANS20 vs BANS40 and F_{ST} = 0.10 for GAL20 vs GAL40). Considering the F_{ST} between

depths, high F_{ST} values can be observed for different loci and different samples comparisons

(Fig. S4). The average F_{ST} between the two depths of the same site was 0.04 in Marseille, 0.08

in Corsica and 0.02 in Banyuls (0.04 for the total dataset). Considering only putatively neutral

loci (see below for outliers loci), the overall F_{ST} of the total dataset was 0.12 and pairwise F_{ST}

values ranged from 0.01 (BANS20 vs BANS40, BANS20 vs BANN40 and BANN40 vs

BANS40) to 0.23 (ELV12 vs GAL20) (Table S1). There was no correlation between the

distance between two depths of the same site and the corresponding population pairwise F_{ST}

(p = 0.75). We obtained a similar result (p = 1) if we removed the four populations of

Marseille whose sampling sites for the two considered depths were not exactly the same

(653 m horizontal distance between FIG8 and MOR40 and 995 m between ELV12 and

MEJ40). Finally, the F_{ST} between the two shallow sites inside a geographical region was in all

three cases higher than those between the two deep sites of the same region (0.10 vs 0.058 in

Marseille, 0.20 vs 0.14 in Corsica and 0.025 vs 0.014 in Banyuls. This difference was

significant in each of the three regions (p < 1.10^{-16} with a t-test in all cases). In a similar way,

the F_{ST} between two shallow sites of two different geographical regions were in all cases

higher than those between the two corresponding deep sites, except for the comparisons

between the Porto sites and the Banyuls sites (p = 0.38 and p = 0.02 for the POR/BANN and

POR/BANS comparisons respectively; p < 1.10^{-16} for the other comparisons).

The AMOVA indicated a similar percentage of the molecular variance explicated by

differences among group and within groups (7.8 and 7.07 % respectively) and approximately

85 % of variance explicated by differences within populations (Table 6). There was significant

genetic differentiation at the three studied levels (F_{ST} = 0.15, F_{SC} = 0.08, F_{CT} = 0.08 ; p < 0.001

in the three cases).

Outliers SNPs

We identified 82 outliers with BayeScEnv. However, we noticed that all these outliers seemed

to be driven by the divergence between particular populations, with one allele being always

fixed in one or several populations without logical association with depth. ARLEQUIN

detected 563 loci potentially under selection in Marseille, 869 in Corsica and 397 in Banyuls.

Among these SNPs, all corresponded to a signal of divergent selection in Marseille and

Banyuls, 207 of the 869 candidate loci corresponded to a signal of balanced selection in

Corsica and the remaining 662 loci corresponded to a signal of divergent selection.

Considering only these outliers, the overall F_{ST} of the total dataset was 0.25 and pairwise F_{ST}

values ranged from 0.02 (BANS20 vs BANS40) to 0.42 (GAL20 vs POR40) (Table S2). The

207 loci potentially under balanced selection in Corsica were linked with sex differentiation

and were not further analyzed here (Pratlong *et al*. 2017). Eight outlier SNPs were detected

both in Marseille and Banyuls, 12 both in Marseille and Corsica and 12 both in Corsica and

Banyuls. No SNP was detected as potentially under divergent selection and common in the

32   16

three regions. The complementary Chi² test of homogeneity of genotypic frequencies

between depths inside each region detected 162 candidate loci in Marseille, 1 371 in Corsica

and 3 in Banyuls. Among these loci, 35, 248 and 2 where also respectively detected with the

ARLEQUIN analysis. The numbers of outlier loci were correlated with the variance and the

average of F_{ST} values inside each geographical regions (correlation coefficient of 0.97). The

second axis of the PCA using the Marseille individuals showed apparent association with

depth and appeared to be influenced by the variation of the candidates for local adaptation to

depth detected by ARLEQUIN: 51 % of these loci were in the top 1 % of the contributions to

this second axis, and 86 % were in the top 5 %.

Because the Marseille region is the only one presenting a principal component linked with

depth (Fig. 1C), the pcadapt results obtained for the Corsica et Banyuls regions have poor

biological relevance for our biological question. We chose thus to present only the pcadapt

results obtained for the Marseille region. Pcadapt detected 58 outliers loci linked with the

second PCA axis, the one which was linked to depth differentiation. All these candidates were

detected by the ARLEQUIN analyses and 20 were also common with the Chi² test presented

above.

Functional annotation

Among the 27 461 analyzed RAD-tags, 6 376 had hits on the red coral transcriptome

(23.2 %). Concerning SNPs detected as outliers by ARLEQUIN and contributing to the depth

divergence, 8 on the 35 detected in Marseille had hits on the transcriptome, 46 on the 248

detected in Corsica and 2 on the 2 detected in Banyuls (Table S3). We did not observed any

GO term enriched in coding regions among candidates SNPs, nor any functional enrichment

in these outliers.

DISCUSSION

Genetic diversity and structure

Our results confirm at a genomic scale the high genetic structure of the red coral, which was observed with a small number of microsatellite loci (Costantini *et al.*, 2007; Ledoux *et al.*, 2010b). These results could be the consequence of reduced mean larval dispersal distance, despite a quite long pelagic larval duration estimated in aquarium (from 16 to 42 days; Martínez-Quintana *et al.*, 2015). Genetic incompatibilities could also contribute to the observed differentiation at least for some loci (Kulmuni and Westram, 2017). Our analysis of the genetic structure of the red coral revealed several clusters mainly corresponding to the geographical distributions of this species. A high differentiation was observed here with PCA and STRUCTURE between the two sites of Corsica separated by around 22 km. This pattern of genetic structure could be explained by a putative barrier to gene flow (through currents or lack of suitable habitats) between the two Corsican sites, or it could also correspond to an historical separation of these populations: two lineages could then be present in Corsica, with one being related to Banyuls populations.

We reported here a significant vertical genetic structure between the two depths of the same site (populations separated by less than 20 m). This differentiation was also observed when outlier loci were removed indicating that it is also shaped by neutral processes (migration / drift) as suggested previously with microsatellites (e.g. Ledoux *et al.*, 2010a). In a study of the vertical genetic structure of red coral in two western Mediterranean sites (Cap de Creus, Spain and Portofino, Italia), Costantini *et al.* (2011) observed a drop in connectivity around 40 – 50 m depth, with genetic diversity declining with depth, but our sampling scheme did not allow us to test this hypotheses. The study of the vertical genetic structure in corals is important in

the context of climate change. As deeper populations may be less affected by climate change,

they could possibly reseed shallower populations (Bongaerts *et al.*, 2017). Nevertheless this

possibility of reseeding depends on the connectivity or potential barriers between depths. A

lack of connectivity could erroneously be inferred in cases of cryptic species (Pante *et al.*,

2015). Contrary to Prada *et al.* (2008) who showed the existence of two cryptic lineages at

two different depths in a tropical octocoral, the populations of red coral from the two studied

depths clearly correspond here to the same species, and the differentiation between depth was

lower than the differentiation between sites and regions. This vertical structure may be the

result of both inherent life history traits and environmental variables. Weinberg (1979) has

described a negative geotropism for the planulae of *C. rubrum,* and Martínez-Quintana *et al.*

(2015) demonstrated that this was an active behavior. Depending on the orientation of the

substrate, this can limit the connectivity between depths. The seasonal stratification during

larval emission (which can occur from June to September; Haguenauer A., pers. comm.) could

also limit dispersal. According to these hypotheses the larval behavior and oceanographic

factors should lead to the genome-wide neutral differentiation between depths. This

differentiation is probably also shaped by drift induced by the small effective size of red coral

populations (Ledoux et al., 2010a).

The horizontal genetic differentiation between the two shallow sites was higher than those

between the two corresponding deep sites, inside and between geographical region. This

suggests a higher connectivity or lower rate of genetic drift for deep populations compared to

shallow ones. The repeated colonization of shallow depths from deeper ones or the higher

harvesting pressure on shallow populations compared to deep one could enhance genetic drift

as well (Rossi *et al.*, 2008; Cannas *et al.*, 2016). We did not observe here a reduction in gene

diversity for shallow populations, connectivity differences then seem to be more probable in

explaining the observed differences of genetic differentiation. Interestingly, Rossi *et al.* (2008)

observed a higher frequency of patches of red coral below 50 m compared to above 50 m: if

such pattern is present in the area and depths considered here, then it could increase gene flow

through stepping stones migration. The observed vertical and horizontal genetic structure

could indicate reduced recolonization abilities following disturbances such as mortality event

induced by heat waves. Nevertheless the observed genetic structure could also be shaped by

colonization history and monopolization effect (Orsini *et al.*, 2013). In this case, a disturbance

leading to free habitats would facilitate recolonization from other populations.

Potential biases in the search of outlier loci

We observed high F_{ST} values for different loci and different sample comparisons, and not only

between depths (Fig. S4). The methods used here to identify selected loci will most likely

detect loci with strong effects (Pritchard and Di Rienzo, 2010; Gagnaire and Gaggiotti, 2016),

and it is highly improbable to observe such a high number of selected loci. Both hierarchical

genetic structure and high levels of differentiation are known to lead to a high number of false

positives in genomic studies of local adaptation, as observed here (Bierne *et al.*, 2013; Hoban

et al. 2016).

The markers density obtained with RAD-Seq may be also insufficient to detect a RAD-tag in

linkage disequilibrium with a selected locus (Lowry *et al.*, 2017; McKinney *et al.*, 2017;

Catchen *et al.*, 2017). With a genome size of about 500 Mb for the red coral (Ganot *et al.*,

2016), we expected a SNP sampling of 1 for 18 kb after the SNPs filtering steps (55 SNPs per

Mb). Nevertheless, a potential high linkage disequilibrium may be present in here because of

drift, and we thus expected that our RAD-tags to at least detect a signal of genetic adaptation,

which remains to be confirmed. Approaches dedicated to the study of polygenic adaptation

(Daub *et al.*, 2013) or to the genomic distribution of F_{ST} or nucleotide diversity (Hohenlohe *et*

al., 2010) could be interesting here used, but a reference genome is still lacking for the red

coral. On a more theoretical point of view, outlier loci could be linked to intrinsic genetic

incompatibilities whose allelic frequencies coupled with environmental barriers (Bierne *et al.*,

2011). The frequency of genetic incompatiblities in marine populations is largely unknown

but probably under-estimated (Plough *et al.*, 2016). Even if not directly linked to local

adaptation, such loci are important factors in the evolution of red coral populations.

Local adaptation to depth in the red coral

We focused on candidate loci meeting the following criteria: i) detection with ARLEQUIN

and pcadapt, ii) significant differentiation between depth, iii) function relevant to the

adaptation to thermal regime. These loci are the most relevant as factors of local adaptation.

The absence of candidate SNPs common to the three geographical regions could indicate that

the adaptation to comparable shallow environmental pressures in these independent regions

are based on different genetic pathways, or on non-genetic mechanisms (Putnam and Gates,

2015). However, most candidate loci should be in linkage disequilibrium with selected loci,

and such association can easily be lost between distant locations through recombination.

Differences in the strength of selective pressure in the three regions could also explain the

differences in the detected loci. In Marseille we evidenced a clear signal of differentiation

between depths according to multivariate and outlier loci analyses. This detection of a signal

of local adaptation in the Marseille region is consistent with the observations from studies of

thermotolerance differences in this region (Torrents *et al.*, 2008; Haguenauer *et al.*, 2013;

Ledoux *et al.*, 2015; Pratlong *et al.*, 2015). In the case of the Marseille region there are then

strong evidences of the existence of adaptive differentiation at a scale of few tens of meters

only. Concerning Corsican populations, Ledoux *et al*. (2015) reported no phenotypic signal of

local adaptation after reciprocal transplant experiment. Here, the most promising candidate for

the adaptation to thermal regime, was an homologous to an allene oxide synthase-

lipoxygenase which is known to be involved in the response to thermal stress in octocorals

(Lõhelaid *et al.*, 2015). This indirect argument would support the presence of local adaptation

in this area as well, but more experimental analyzes will be necessary to confirm the

involvement of this function in adaptation to thermal stress in this species. Finally, the

detection of a reduced number of candidate loci (two) in the Banyuls region would be

consistent with the weaker selective pressure here (see above, Table 2).

Previously, we have sequenced the transcriptome of individuals from the two depths of the

Marseille site studied here in Marseille (Pratlong *et al.*, 2015). Several genes were

differentially expressed between individuals from the two depths outside thermal stress

conditions. Some of these genes, such as those from the Tumor Necrosis Factors Receptor

Associated Factors (TRAF) family, have been identified as involved in the response to

thermal stress in the hexacoral *Acropora hyacinthus* (Barshis et al., 2013). However, none of

these differentially expressed genes were identified in our RAD-Seq study. Additionally the

differences of expression may result from acclimatization, and are not necessarily adaptive.

The use of a reference genome would be useful here as well to study the potential link

between candidate SNPs and genes location and function (Manel *et al.*, 2016).

CONCLUSION

To our knowledge, the red coral presents among the highest levels of differentiation among

studies of local adaptation thought genome scans approaches in marine environment

(Bradbury *et al.*, 2010; Limborg *et al.*, 2012; Wang *et al.*, 2013; Chu *et al.*, 2014; Milano *et*

al., 2014; Bay and Palumbi, 2014; Bernardi *et al.*, 2016; Araneda *et al.*, 2016; Guo *et al.*,

2016; Benestan *et al.*, 2016; Bongaerts *et al.*, 2017). This study enabled us to empirically

emphasize the limitations in the detection and the interpretation of signals of local adaptation

using usual statistical methods in this strongly structured species. Both neutral an adaptive

divergence highlighted here demonstrate the genetically singularity of shallow populations of

the red coral, especially in the Marseille region were the shallowest populations of this species

are found. Together, the strong genetic structure we observed between shallow populations,

the low dispersal abilities of the red coral and the local adaptation of these individuals to the

highly variable thermal conditions they experiment, raise strong concerns about the evolution

of shallow populations and the possibility of loss of adaptive variations in case of mortality

events. Extending the genomic study initiated here would be useful to study the evolution of

this species in heterogeneous and changing environments. Whatever their origin (genetic or

environmental), the different thermotolerance levels observed between depths and populations

in the red coral should also be taken into account in future studies of adaptive evolution in this

species. These results would also be useful for the management of this harvested species

which is present in two international conservation conventions.

Acknowledgements:

This work is a contribution to the Labex OT-Med (n° ANR-11-LABX-0061) funded by the

French Government "Investissements d'Avenir" program of the French National Research

Agency (ANR) through the A*MIDEX project (n° ANR-11-IDEX-0001-02). This project has

been funded by the ADACNI program of the French National Research Agency (ANR)

(project n°ANR-12-ADAP-0016; http://adacni.imbe.fr). We thank the ECCOREV Research

Federation (FR 3098) for the financial support of part of this study. The project leading to this

publication has received funding from European FEDER Fund under project 1166-39417. We

thank Nicolas Fernandez and Béatrice Loriod from the Marseille TGML platform for their

invaluable help and advice with the preparation of the RAD libraries; the team of the MGX

platform for the sequencing of the RAD libraries. The authors thank the UMR 8199 LIGAN-

PM Genomics platform (Lille, France, especially Véronique Dhennin) which belongs to the

'Federation de Recherche' 3508 Labex EGID (European Genomics Institute for Diabetes;

ANR-10-LABX-46) and was supported by the ANR Equipex 2010 session (ANR-10-EQPX-

07-01; 'LIGAN-PM'). The LIGAN-PM Genomics platform (Lille, France) is also supported

by the FEDER and the Region Nord-Pas-de-Calais-Picardie. We thank the molecular biology

service of the IMBE, the informatic service of the Pytheas Institute, and Frédéric Zuberer of

the Pytheas Institute for his support in sampling. We thank the Scandola Natural Reserve,

especially Jean-Marie Dominici. We thank Manuela Carenzi, Pierre-Alexandre Gagnaire,

François Bonhomme and Nicolas Bierne for stimulating discussions.

References

Alleaume-Benharira M, Pen IR, Ronce O (2006). Geographical patterns of adaptation within a species' range: interactions between drift and gene flow. *J Evol Biol* **19**: 203–215.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403–410.

Araneda C, Larraín MA, Hecht B, Narum S (2016). Adaptive genetic variation distinguishes Chilean blue mussels (*Mytilus chilensis*) from different marine environments. *Ecol Evol* **6**: 3632–3644.

Ayre DJ (1995). Localized adaptation of sea anemone clones: evidence from transplantation over two spatial scales. *J Anim Ecol*: 186–196.

Barrett RD, Hoekstra HE (2011). Molecular spandrels: tests of adaptation at the genetic level. *Nat Rev Genet* **12**: 767–780.

Barshis DJ, Ladner JT, Oliver TA, Seneca FO, Traylor-Knowles N, Palumbi SR (2013). Genomic basis for coral resilience to climate change. *Proc Natl Acad Sci* **110**: 1387–1392.

Barshis DJ, Stillman JH, Gates RD, Toonen RJ, Smith LW, Birkeland C (2010). Protein expression and genetic structure of the coral Porites lobata in an environmentally extreme Samoan back reef: does host genotype limit phenotypic plasticity? *Mol Ecol* **19**: 1705–1720.

Bay RA, Palumbi SR (2014). Multilocus Adaptation Associated with Heat Resistance in Reef-Building Corals. *Curr Biol* **24**: 2952–2956.

Benestan L, Quinn BK, Maaroufi H, Laporte M, Rochette R, Bernatchez L (2016). Seascape genomics provides evidence for thermal adaptation and current-mediated population structure in American lobster (*Homarus americanus*). *Mol Ecol* **25(20)**: 5073-5092.

Benjamini Y, Hochberg Y (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B Methodol* **57**: 289–300.

Bernardi G, Azzurro E, Golani D, Miller MR (2016). Genomic signatures of rapid adaptive evolution in the bluespotted cornetfish, a Mediterranean Lessepsian invader. *Mol Ecol* **25(14)** : 3384-3396.

Bierne N, Roze D, Welch JJ (2013). Pervasive selection or is it…? why are FST outliers sometimes so frequent? *Mol Ecol* **22**: 2061–2064.

Bierne N, Welch J, Loire E, Bonhomme F, David P (2011). The coupling hypothesis: why genome scans may fail to map local adaptation genes. *Mol Ecol* **20**: 2044–2072.

Boavida J, Paulo D, Aurelle D, Arnaud-Haond S, Marschal C, Reed J, *et al.* (2016). A well-kept treasure at depth: precious red coral rediscovered in Atlantic deep coral gardens (SW Portugal) after 300 years. *PloS One* **11**: e0147228.

Bongaerts P, Riginos C, Brunner R, Englebert N, Smith SR, Hoegh-Guldberg O (2017). Deep reefs are not universal refuges: Reseeding potential varies among coral species. *Sci Adv* **3**: e1602373.

Bongaerts P, Riginos C, Hay KB, Oppen MJ van, Hoegh-Guldberg O, Dove S (2011). Adaptive divergence in a scleractinian coral: physiological adaptation of Seriatopora hystrix to shallow and deep reef habitats. *BMC Evol Biol* **11**: 303.

Bradbury IR, Hubert S, Higgins B, Borza T, Bowman S, Paterson IG, *et al.* (2010). Parallel adaptive evolution of Atlantic cod on both sides of the Atlantic Ocean in response to temperature. *Proc R Soc Lond B Biol Sci* **277**: 3725–3734.

Cannas R, Sacco F, Cau A, Cuccu D, Follesa MC, Cau A (2016). Genetic monitoring of deep-water exploited banks of the precious Sardinia coral *Corallium rubrum* (L., 1758): useful data for a sustainable management. *Aquat Conserv Mar Freshw Ecosyst* **26**: 236–250.

Catchen JM, Amores A, Hohenlohe P, Cresko W, Postlethwait JH (2011). Stacks: building and genotyping loci de novo from short-read sequences. *G3 Genes Genomes Genet* **1**: 171–182.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013). Stacks: an analysis tool set for population genomics. *Mol Ecol* **22**: 3124–3140.

Catchen JM, Hohenlohe PA, Bernatchez L, Funk WC, Andrews KR, Allendorf FW (2017). Unbroken: RADseq remains a powerful tool for understanding the genetics of adaptation in natural populations. *Mol Ecol Resour* **17(3)**: 362-365.

Chu ND, Kaluziak ST, Trussell GC, Vollmer SV (2014). Phylogenomic analyses reveal latitudinal population structure and polymorphisms in heat stress genes in the North Atlantic snail Nucella lapillus. *Mol Ecol* **23**: 1863–1873.

Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M (2005). Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* **21**: 3674–3676.

Coop G, Witonsky D, Rienzo AD, Pritchard JK (2010). Using Environmental Correlations to Identify Loci Underlying Local Adaptation. *Genetics* **185**: 1411–1423.

Costantini F, Fauvelot C, Abbiati M (2007). Genetic structuring of the temperate gorgonian coral (*Corallium rubrum*) across the western Mediterranean Sea revealed by microsatellites and nuclear sequences. *Mol Ecol* **16**: 5168–5182.

Costantini F, Rossi S, Pintus E, Cerrano C, Gili J-M, Abbiati M (2011). Low connectivity and declining genetic variability along a depth gradient in Corallium rubrum populations. *Coral Reefs* **30**: 991–1003.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, *et al.* (2011). The variant call format and VCFtools. *Bioinformatics* **27**: 2156–2158.

Daub JT, Hofer T, Cutivet E, Dupanloup I, Quintana-Murci L, Robinson-Rechavi M, *et al.* (2013). Evidence for polygenic adaptation to pathogens in the human genome. *Mol Biol Evol* **30(7)**: 1544-1558.

DeWitt TJ, Sih A, Wilson DS (1998). Costs and limits of phenotypic plasticity. *Trends Ecol Evol* **13**: 77–81.

Doney SC, Ruckelshaus M, Duffy JE, Barry JP, Chan F, English CA, *et al.* (2012). Climate change impacts on marine ecosystems. *An Rev Mar Sci* **4(1)** : 11-37.

Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA (2011). SNP discovery and genotyping for evolutionary genetics using RAD sequencing. In *Molecular methods for evolutionary genetics*, Humana press. pp 157-178.

Evanno G, Regnaut S, Goudet J (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* **14**: 2611–2620.

Excoffier L, Lischer HEL (2010). Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* **10**: 564–567.

Falush D, Stephens M, Pritchard JK (2003). Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* **164**: 1567–1587.

Falush D, Stephens M, Pritchard JK (2007). Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Mol Ecol Notes* **7**: 574–578.

Felsenstein J (1975). Genetic Drift in Clines Which Are Maintained by Migration and Natural-Selection. *Genetics* **81**: 191–207.

Gagnaire P-A, Gaggiotti OE (2016). Detecting polygenic selection in marine populations by combining population genomics and quantitative genetics approaches. *Curr Zool* **62(6)**: 603-616.

Ganot P, Fritz M, Rausch T, Aurelle D, Haguenauer A, Romans P, *et al.* (2016). Calcification in Corallium rubrum: dissecting the process using genomics and transcriptomics together with scalpels and scissors. 13th International Coral Reef Symposium, Honolulu, Hawai'i

Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P, Cigliano M, *et al.* (2009). Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Glob Change Biol* **15**: 1090–1103.

Garrabou J, Perez T, Sartoretto S, Harmelin JG (2001). Mass mortality event in red coral Corallium rubrum populations in the Provence region (France, NW Mediterranean). *Mar Ecol Prog Ser* **217**: 263–272.

Guo B, Li Z, Merilä J (2016). Population genomic evidence for adaptive differentiation in the

Baltic Sea herring. *Mol Ecol* **25**: 2833–2852.

Haguenauer A, Zuberer F, Ledoux J-B, Aurelle D (2013). Adaptive abilities of the Mediterranean red coral *Corallium rubrum* in a heterogeneous and changing environment: from population to functional genetics. *J Exp Mar Biol Ecol* **449**: 349–357.

Hastings A, Rohlf FJ (1974). Gene Flow: Effect in Stochastic Models of Differentiation. *Am Nat* **108**: 701–705.

Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, *et al.* (2016). Finding the genomic basis of local adaptation: pitfalls, practical solutions, and future directions. *Am Nat* **188**: 379–397.

Hofer T, Ray N, Wegmann D, Excoffier L (2009). Large allele frequency differences between human continental groups are more likely to have occurred by drift during range expansions than by selection. *Ann Hum Genet* **73**: 95–108.

Hohenlohe PA, Bassham S, Etter PD, Stiffler N, Johnson EA, Cresko WA (2010). Population Genomics of Parallel Adaptation in Threespine Stickleback using Sequenced RAD Tags. *PLoS Genet* **6**: e1000862.

Hubisz MJ, Falush D, Stephens M, Pritchard JK (2009). Inferring weak population structure with the assistance of sample group information. *Mol Ecol Resour* **9**: 1322–1332.

Jin YK, Lundgren P, Lutz A, Raina J-B, Howells EJ, Paley AS, *et al.* (2016). Genetic markers for antioxidant capacity in a reef-building coral. *Sci Adv* **2**: e1500842.

Jombart T (2008). adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**: 1403–1405.

Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, Johnson J, *et al.* (2012). The genomic basis of adaptive evolution in threespine sticklebacks. *Nature* **484**: 55–61.

Kawecki TJ, Ebert D (2004). Conceptual issues in local adaptation. *Ecol Lett* **7**: 1225–1241.

Kersting DK, Bensoussan N, Linares C (2013). Long-Term Responses of the Endemic Reef-Builder *Cladocora caespitosa* to Mediterranean Warming. *PLoS ONE* **8**: e70820.

Knittweis L, Aguilar R, Alvarez H, Borg J, Evans J, Garcia S, *et al.* (2016). New depth record of the precious red coral *Corallium rubrum* for the Mediterranean. *antiquity* **7**: 8.

Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I (2015). Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Mol Ecol Resour* **15**: 1179–1191.

Kulmuni J, Westram AM (2017). Intrinsic incompatibilities evolving as a by-product of divergent ecological selection: Considering them in empirical studies on divergence with gene flow. *Mol Ecol* **26**: 3093–3103.

Ledoux J-B, Aurelle D, Bensoussan N, Marschal C, Féral J-P, Garrabou J (2015). Potential for adaptive evolution at species range margins: contrasting interactions between red coral populations and their environment in a changing ocean. *Ecol Evol* **5**: 1178–1192.

Ledoux J-B, Aurelle D., Féral J-P, Garrabou J (2013). Molecular forensics in the precious Mediterranean red coral, *Corallium rubrum*: testing DNA extraction and microsatellite genotyping using dried colonies. *Cons Gen Res* **5(2)** : 327-330.

Ledoux J-B, Garrabou J, Bianchimani O, Drap P, Féral J-P, Aurelle D (2010a). Fine-scale genetic structure and inferences on population biology in the threatened Mediterranean red coral, *Corallium rubrum*. *Mol Ecol* **19**: 4204–4216.

Ledoux J-B, Mokhtar-Jamaï K, Roby C, Féral J-P, Garrabou J, Aurelle D (2010b). Genetic survey of shallow populations of the Mediterranean red coral [*Corallium rubrum* (Linnaeus, 1758)]: new insights into evolutionary processes shaping nuclear diversity and implications for conservation. *Mol Ecol* **19**: 675–690.

Lenormand T (2002). Gene flow and the limits to natural selection. *Trends Ecol Evol* **17**: 183–189.

Li H, Durbin R (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**: 1754–1760.

Limborg MT, Helyar SJ, de Bruyn M, Taylor MI, Nielsen EE, Ogden ROB, *et al.* (2012). Environmental selection on transcriptome-derived SNPs in a high gene flow marine fish, the Atlantic herring (*Clupea harengus*). *Mol Ecol* **21**: 3686–3703.

Lõhelaid H, Teder T, Samel N (2015). Lipoxygenase-allene oxide synthase pathway in octocoral thermal stress response. *Coral Reefs* **34**: 143–154.

Lowry DB, Hoban S, Kelley JL, Lotterhos KE, Reed LK, Antolin MF, *et al.* (2017). Breaking RAD: an evaluation of the utility of restriction site-associated DNA sequencing for genome scans of adaptation. *Mol Ecol Resour* **17(2)** : 142-152.

Lundgren P, Vera JC, Peplow L, Manel S, Oppen MJ van (2013). Genotype – environment correlations in corals from the Great Barrier Reef. *BMC Genet* **14**: 9.

Luu K, Bazin E, Blum MGB (2017). pcadapt: an R package to perform genome scans for selection based on principal component analysis. *Mol Ecol Resour* **17**: 67–77.

Manel S, Perrier C, Pratlong M, Abi-Rached L, Paganini J, Pontarotti P, *et al.* (2016). Genomic resources and their influence on the detection of the signal of positive selection in genome scans. *Mol Ecol* **25**: 170–184.

Marschal C, Garrabou J, Harmelin JG, Pichon M (2004). A new method for measuring growth and age in the precious red coral *Corallium rubrum* (L.). *Coral Reefs* **23**: 423–432.

Martínez-Quintana A, Bramanti L, Viladrich N, Rossi S, Guizien K (2015). Quantification of larval traits driving connectivity: the case of *Corallium rubrum* (L. 1758). *Mar Biol* **162**: 309–318.

McKinney GJ, Larson WA, Seeb LW, Seeb JE (2017). RADseq provides unprecedented insights into molecular ecology and evolutionary genetics: comment on Breaking RAD by Lowry et al.(2016). *Mol Ecol Resour* **17**: 356–361.

Milano I, Babbucci M, Cariani A, Atanassova M, Bekkevold D, Carvalho GR, *et al.* (2014). Outlier SNP markers reveal fine-scale genetic structuring across European hake populations (*Merluccius merluccius*). *Mol Ecol* **23**: 118–135.

Mumby PJ, Elliott IA, Eakin CM, Skirving W, Paris CB, Edwards HJ, *et al.* (2011). Reserve design for uncertain responses of coral reefs to climate change. *Ecol Lett* **14**: 132–140.

Nagylaki T (1978). Random Genetic Drift in a Cline. *Proc Natl Acad Sci U S A* **75**: 423–426.

Orsini L, Vanoverbeke J, Swillen I, Mergeay J, Meester L (2013). Drivers of population genetic differentiation in the wild: isolation by dispersal limitation, isolation by adaptation and isolation by colonization. *Mol Ecol* **22**: 5983–5999.

Palumbi SR, Barshis DJ, Traylor-Knowles N, Bay RA (2014). Mechanisms of Reef Coral Resistance to Future Climate Change. *Science* **344(6186)** : 895-898.

Pante E, Puillandre N, Viricel A, Arnaud-Haond S, Aurelle D, Castelin M, *et al.* (2015). Species are hypotheses: avoid connectivity assessments based on pillars of sand. *Mol Ecol* **24**: 525–544.

Pigliucci M (2001). *Phenotypic plasticity: beyond nature and nurture*. JHU Press.

Pivotto ID, Nerini D, Masmoudi M, Kara H, Chaoui L, Aurelle D (2015). Highly contrasted responses of Mediterranean octocorals to climate change along a depth gradient. *R Soc Open Sci* **2**: 140493.

Plough L, Shin G, Hedgecock D (2016). Genetic inviability is a major driver of type III survivorship in experimental families of a highly fecund marine bivalve. *Mol Ecol* **25**: 895–910.

Plummer M, Best N, Cowles K, Vines K (2006). CODA: Convergence diagnosis and output analysis for MCMC. *R News* **6**: 7–11.

Prada C, Schizas NV, Yoshioka PM (2008). Phenotypic plasticity or speciation? A case from a clonal marine organism. *BMC Evol Biol* **8**: 1.

Pratlong M, Haguenauer A, Chabrol O, Klopp C, Pontarotti P, Aurelle D (2015). The red coral (*Corallium rubrum*) transcriptome: a new resource for population genetics and local adaptation studies. *Mol Ecol Resour* **15**: 1205–1215.

Pratlong M, Haguenauer A, Chenesseau S, Brener K, Mitta G, Toulza E, *et al.* (2017). Evidence for a genetic sex determination in Cnidaria, the Mediterranean red coral (*Corallium rubrum*). *R Soc Open Sci* **4**.

Pritchard JK, Di Rienzo A (2010). Adaptation–not by sweeps alone. *Nat Rev Genet* **11**: 665–667.

Pritchard JK, Stephens M, Donnelly P (2000). Inference of population structure using multilocus genotype data. *Genetics* **155**: 945–959.

Putnam HM, Gates RD (2015). Preconditioning in the reef-building coral *Pocillopora damicornis* and the potential for trans-generational acclimatization in coral larvae under future climate change conditions. *J Exp Biol* **218**: 2365–2372.

R Core Team (2016). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing: Vienna, Austria.

Rossi S, Tsounis G, Orejas C, Padrón T, Gili J-M, Bramanti L, *et al.* (2008). Survey of deep-dwelling red coral (*Corallium rubrum*) populations at Cap de Creus (NW Mediterranean). *Mar Biol* **154**: 533–545.

Rousset F (2008). genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Mol Ecol Resour* **8**: 103–106.

Sambrook J, Fritsch EF, Maniatis T, others (1989). *Molecular cloning*. Cold spring harbor laboratory press New York.

Sanford E, Kelly MW (2011). Local Adaptation in Marine Invertebrates. *Annu Rev Mar Sci* **3**: 509–535.

Santangelo G, Bramanti L, Rossi S, Tsounis G, Vielmini I, Lott C, *et al.* (2012). Patterns of variation in recruitment and post-recruitment processes of the Mediterranean precious gorgonian coral *Corallium rubrum*. *J Exp Mar Biol Ecol* **411**: 7–13.

Sherman CDH, Ayre DJ (2008). Fine-scale adaptation in a clonal sea anemone. *Evol Int J Org Evol* **62**: 1373–1380.

Slatkin M, Maruyama T (1975). Genetic drift in a cline. *Genetics* **81**: 209–222.

Smith LW, Barshis D, Birkeland C (2007). Phenotypic plasticity for skeletal growth, density and calcification of Porites lobata in response to habitat type. *Coral Reefs* **26**: 559–567.

Szulkin M, Gagnaire P-A, Bierne N, Charmantier A (2016). Population genomic footprints of fine-scale differentiation between habitats in Mediterranean blue tits. *Mol Ecol* **25**: 542–558.

Torrents O, Tambutté E, Caminiti N, Garrabou J (2008). Upper thermal thresholds of shallow vs. deep populations of the precious Mediterranean red coral *Corallium rubrum* (L.): Assessing the potential effects of warming in the NW Mediterranean. *J Exp Mar Biol Ecol* **357**: 7–19.

Ulstrup KE, Van Oppen MJH (2003). Geographic and habitat partitioning of genetically distinct zooxanthellae (*Symbiodinium*) in Acropora corals on the Great Barrier Reef. *Mol Ecol* **12**: 3477–3484.

Villemereuil P, Gaggiotti OE (2015). A new FST-based method to uncover local adaptation using environmental variables. *Methods Ecol Evol* **6**: 1248–1258.

Wang L, Liu S, Zhuang Z, Guo L, Meng Z, Lin H (2013). Population genetic studies revealed local adaptation in a high gene-flow marine fish, the small yellow croaker (*Larimichthys polyactis*). *PloS One* **8**: e83493.

Weinberg S (1979). The light-dependent behaviour of planula larvae of *Eunicella singularis* and *Corallium rubrum* and its implication for octocorallian ecology. *Bijdr Tot Dierkd* **49**: 16–30.

Wrange A-L, André C, Lundh T, Lind U, Blomberg A, Jonsson PJ, *et al.* (2014). Importance of plasticity and local adaptation for coping with changing salinity in coastal areas: a test case with barnacles in the Baltic Sea. *BMC Evol Biol* **14**: 156.

Ziegler M, Roder CM, Büchel C, Voolstra CR (2014). Limits to physiological plasticity of the coral Pocillopora verrucosa from the central Red Sea. *Coral Reefs* **33**: 1115–1129.

Figure 1. Principal component analysis (Axes 1 and 2) of A) the 12 red coral populations (n = 354 individuals, 27 461 SNPs), B) the four red coral populations from Marseille (n =119 individuals, 27 461 SNPs), C) the four red coral populations from Corsica (n =117 individuals, 27 461 SNPs), D) the four red coral populations from Banyuls (n =118 individuals, 27 461 SNPs). Triangles and circles correspond to shallow and deep individuals respectively.

Figure 2. Results from Bayesian individual clustering with STRUCTURE for K = 2 to K = 4. For K = 2 and K = 4, all ten replicates produced the same

structure. For K = 3, the major mode presented here was the result of 7/10 replicates. Minor modes are presented in Fig. S3.

561   Table 1. Characteristics of red coral sampling sites.

Population	Geographic region	Site	Depth (m)	GPS	GPS
FIG8	Marseille	Marseille South	8	43° 12.330'N	5° 26.790'E
MOR40	Marseille	Marseille South	40	43° 12.060'N	5° 27.100'E
ELV12	Marseille	Marseille North	12	43° 19.780'N	5° 14.210'E
MEJ40	Marseille	Marseille North	40	43° 19.700'N	5° 13.480'E
BANN20	Banyuls	Banyuls North	25	42° 26.890'N	3° 10.330'E
BANN40	Banyuls	Banyuls North	35	42° 26.890'N	3° 10.330'E
BANS20	Banyuls	Banyuls South	26	42° 26.390'N	3° 10.790'E
BANS40	Banyuls	Banyuls South	36	42° 26.390'N	3° 10.790'E
POR20	Corsica	Porto	21	42° 16.292'N	8° 41.255'E
POR40	Corsica	Porto	33	42° 16.292'N	8° 41.255'E
GAL20	Corsica	Galeria	26	42° 28.210'N	8° 38.950'E
GAL40	Corsica	Galeria	36	42° 28.210'N	8° 38.950'E

564 Table 2. Temperatures (in °C) characteristics of the sampling sites from March 2012 to

565 October 2014.

	Depth (m)	Minimum	Maximum	Mean	Standard Deviation
FIG8	8	12.63	26.92	17.03	3.52
MOR40	40	12.73	23.06	15.40	2.11
ELV12	12	11.81	26.70	16.60	3.24
MEJ40	40	11.86	22.87	15.29	2.18
BANN20	25	12.22	24.29	17.20	2.63
BANN40	35	9.41	23.83	14.49	2.45
BANS20	26	12.22	24.29	17.20	2.63
BANS40	36	9.41	23.83	14.49	2.45
POR20	21	12.51	25.91	17.51	3.41
POR40	33	12.56	23.83	16.26	2.45
GAL20	26	12.46	25.09	17.13	3.13
GAL40	36	12.56	23.83	16.26	2.45

568   Table 3. Counts of SNP loci after each filtering step.

Step	Number of SNPs	Software
After assembly raw data	138 810	Stacks (Catchen *et al.*, 2011, 2013)
Excluding loci not in within population HWE	86 520	VCFtools (Danecek *et al.*, 2011)
MAF 1 %	56 844	VCFtools (Danecek *et al.*, 2011)
One SNPs per RAD-tag	27 461	

74   37

571 Table 4. Summary statistics of genetic diversity. The nucleotide diversities (SNP level),

572 numbers of private alleles and Tajima's D are based on the dataset with all SNPs for each

573 RAD-loci, excluding those with the lowest quality, and those with minor allele frequencies

574 below 0.01 (total 56 844 SNPs). Measures of F_{IS} and gene diversity were based on the final

575 dataset comprising with one SNP per locus (total 27 461 SNPs).

Population	Sample size	Nucleotide diversity	Number of private alleles	F_{IS}	Gene diversity
BANN20	30	0.499	0	0.018	0.15
BANN40	28	0.475	0	0.012	0.15
BANS20	30	0.496	0	0.019	0.15
BANS40	30	0.492	0	0.065	0.13
ELV12	30	0.5	0.001	0.005	0.17
MEJ40	29	0.496	0	0.053	0.18
FIG8	30	0.502	0.001	0.005	0.18
MOR40	30	0.498	0.001	0.036	0.18
GAL20	28	0.489	0.002	0.013	0.09
GAL40	29	0.478	0.001	0.019	0.13
POR20	30	0.494	0	0.023	0.13
POR40	30	0.500	0.006	0.009	0.15

578   Table 5. Pairwise F_{ST} estimates. All comparisons were highly significant. Intra-region

579   comparisons are highlighted.

	BANN20	BANN40	BANS20	BANS40	ELV12	MEJ40	FIG8	MOR40	GAL20	GAL40	POR20	POR40
BANN20	-											
BANN40	0.02	-										
BANS20	0.03	0.01	-									
BANS40	0.03	0.01	0.01	-								
ELV12	0.13	0.13	0.13	0.13	-							
MEJ40	0.11	0.11	0.11	0.11	0.03	-						
FIG8	0.14	0.13	0.13	0.13	0.10	0.08	-					
MOR40	0.11	0.10	0.10	0.10	0.08	0.06	0.05	-				
GAL20	0.18	0.17	0.17	0.17	0.24	0.22	0.24	0.21	-			
GAL40	0.11	0.10	0.10	0.10	0.19	0.16	0.18	0.15	0.10	-		
POR20	0.14	0.13	0.13	0.13	0.17	0.16	0.17	0.14	0.20	0.13	-	
POR40	0.14	0.13	0.13	0.13	0.17	0.15	0.17	0.14	0.21	0.14	0.05	-

581   Table 6. Percent of the variation explained by grouping populations according to their

582   geographical region on the analysis of molecular variance (performed with ARLEQUIN).

Source of variation	d.f.	Percentage of variation
Among groups	2	7.80
Among populations within groups	9	7.07
Within populations	696	85.13

Supplementary material

Figure S1. Location of the sampling sites of the red coral among the three studied

geographical regions.

Figure S2. Principal component analysis (Axes 1 and 2), using only putative neutral SNPs, of

the A) 12 red coral populations (n = 354 individuals, 25 669 SNPs), B) four red coral

populations from Marseille (n =119 individuals, 26 898 SNPs), C) four red coral populations

from Corsica (n =117 individuals, 26 592 SNPs), D) four red coral populations from Banyuls

(n =118 individuals, 27 069 SNPs).

.

Figure S3. Results from Bayesian individual clustering with STRUCTURE for K = 3. The three figures correspond to major and minor modes detected.

Figure S4. Joint distribution of between-depths F_{ST} in the three geographical regions.

617 Table S1. Pairwise F_{ST} estimates using only putatively neutral SNPs. All comparisons were
618 highly significants. Intra-region comparisons are highlighted.
619

	BANN20	BANN40	BANS20	BANS40	ELV12	MEJ40	FIG8	MOR40	GAL20	GAL40	POR20	POR40
BANN20	-											
BANN40	0.02	-										
BANS20	0.02	0.01	-									
BANS40	0.02	0.01	0.01	-								
ELV12	0.13	0.12	0.12	0.12	-							
MEJ40	0.11	0.10	0.10	0.10	0.03	-						
FIG8	0.13	0.12	0.12	0.12	0.09	0.07	-					
MOR40	0.10	0.10	0.09	0.09	0.07	0.05	0.04	-				
GAL20	0.17	0.16	0.16	0.16	0.23	0.21	0.22	0.20	-			
GAL40	0.11	0.09	0.09	0.09	0.18	0.16	0.18	0.15	0.10	-		
POR20	0.13	0.12	0.12	0.12	0.17	0.15	0.16	0.14	0.17	0.11	-	
POR40	0.13	0.12	0.12	0.12	0.16	0.15	0.16	0.14	0.18	0.12	0.04	-

620
621 Table S2. Pairwise F_{ST} estimates using only outlier SNPs from the ARLEQUIN analysis. All
622 comparisons were highly significants. Intra-region comparisons are highlighted.
623

	BANN20	BANN40	BANS20	BANS40	ELV12	MEJ40	FIG8	MOR40	GAL20	GAL40	POR20	POR40
BANN20	-											
BANN40	0.04	-										
BANS20	0.06	0.03	-									
BANS40	0.06	0.03	0.02	-								
ELV12	0.19	0.18	0.19	0.18	-							
MEJ40	0.16	0.14	0.15	0.15	0.04	-						
FIG8	0.19	0.17	0.18	0.18	0.23	0.19	-					
MOR40	0.14	0.12	0.13	0.13	0.19	0.14	0.08	-				
GAL20	0.27	0.27	0.28	0.27	0.37	0.35	0.36	0.32	-			
GAL40	0.15	0.13	0.15	0.13	0.26	0.24	0.26	0.22	0.17	-		
POR20	0.23	0.21	0.21	0.21	0.26	0.22	0.22	0.18	0.41	0.30	-	
POR40	0.24	0.23	0.24	0.23	0.26	0.23	0.23	0.19	0.42	0.31	0.07	-

625
626

628   Table S3. Results of the annotation analysis of candidates for local adaptation in the three

629   geographical regions.

	evalue	Description		GO	Region
Contig_16793	1.1E-117	LON peptidase N-terminal domain and RING finger partial		ubiquitin-protein transferase activity zinc ion binding metal ion binding proteolysis ATP-dependent peptidase activity protein ubiquitination	Marseille
Contig_20016	0	Chromodomain-helicase-DNA-binding 1-like			
Contig_23068	2.8E-117	RNA-directed DNA polymerase from mobile element jockey-like		nucleic acid phosphodiester bond hydrolysis RNA-directed DNA polymerase activity endonuclease activity RNA-dependent DNA biosynthetic process	
Contig_38936	4.1E-23	E3 ubiquitin-ligase DZIP		nucleic acid binding zinc ion binding	
Contig_44372	3.1E-61	PREDICTED : uncharacterized protein LOC107346707		binding	
Contig_47492	7.8E-98	No description		metal ion binding oxidoreductase activity metabolic process oxidation-reduction process	
Contig_7346	2.3E-150	FAM46C-like			
Contig_10570	4.8E-37	PREDICTED : uncharacterized protein			Corsica
Contig_10731	2.8E-68	allene oxide synthase-lipoxygenase		oxidoreductase activity, acting on single donors with incorporation of molecular oxygen, incorporation of two atoms of oxygen	

				metal ion binding fatty acid metabolic process
Contig_11268	2.3E-90	neuronal acetylcholine receptor subunit alpha-9		integral component of membrane plasma membrane part extracellular ligand-gated ion channel activity ion transport synaptic transmission response to stimulus biological regulation
Contig_11731	0	E3 ubiquitin-ligase RNF213		binding
Contig_12920	5.5E-149	Stonin-2 isoform X2		cytoplasmic part vesicle-mediated transport single-organism process intracellular transport
Contig_13771	4.0E-151	A-kinase anchor mitochondrial		hemopoiesis cell differentiation
Contig_16202	6.5E-65	PIN2 TERF1-interacting telomerase inhibitor 1		Nucleus chromosome intracellular organelle part nucleic acid binding regulation of cellular process
Contig_16843	0	Succinate-semialdehyde mitochondrial		aldehyde dehydrogenase (NAD) activity succinate-semialdehyde dehydrogenase (NAD+) activity succinate-semialdehyde dehydrogenase [NAD(P)+] activity glycosylceramide metabolic process multicellular organism development glutamine family amino acid metabolic

				process gamma-aminobutyric acid catabolic process dicarboxylic acid metabolic process oxidation-reduction process	
Contig_16868	4.3E-24	glioma tumor suppressor candidate region gene 1-like		Membrane integral component of membrane	
Contig_17255	7.0E-53	Nanos 1		zinc ion binding RNA binding	
Contig_18282	1.3E-44	centromere K		nucleus	
Contig_19099	0	epithelial growth factor receptor substrate 15-like partial		Calcium ion binding	
Contig_19611	2.0E-134	DDB1- and CUL4-associated factor 5		mitochondrion Cul4-RING E3 ubiquitin ligase complex	
Contig_24102	1.6E-62	nuclease HARBI1			
Contig_24221	0	tRNA (guianine(26)-N(2))-diethyltransferase		nucleus mitochondrion tRNA (guanine-N2-)-methyltransferase activity binding tRNA N2-guanine methylation	
Contig_31623	0	No description		binding	
Contig_32690	1.4E-163	ubiquitin carboxyl-terminal hydrolase isozyme L5		nucleus cytoplasm thiol-dependent ubiquitin-specific protease activity binding ubiquitin-dependent protein catabolic	

				process protein deubiquitination regulation of proteolysis forebrain development regulation of cellular protein metabolic process
Contig_33144	6.6E-56	Dok-7		protein kinase binding positive regulation of protein tyrosine kinase activity insulin receptor binding
Contig_35407	3.8E-11	hypothetical protein AC249_AIPGENE14243		
Contig_36059	5.9E-47	cytosolic non-specific dipeptisase		exopeptidase activity metabolic process
Contig_36102	2.8E-135	nucleolar complex 4 homolog		Nuclear part
Contig_37478	8.9E-28	fibroblast growth factor receptor 3		membrane cell part protein kinase activity phosphorylation positive regulation of phosphorylation regulation of primary metabolic process
Contig_38721	2.3E-41	L-seryl-tRNA(Sec) kinase		phosphorylation kinase activity
Contig_38739	2.0E-56	Dr1		Ada2/Gcn5/Ada3 transcription activator complex DNA binding transcription corepressor activity TBP-class protein binding protein heterodimerization activity negative regulation of transcription from

				RNA polymerase II promoter transcription, DNA-templated histone H3 acetylation
Contig_39033	7.1E-155	homeodomain transcription factor 1 isoform X1		intracellular membrane-bounded organelle
Contig_40626	1.0E-51	RNA-directed DNA polymerase from mobile element jockey-like		nucleic acid phosphodiester bond hydrolysis RNA-directed DNA polymerase activity RNA binding endonuclease activity RNA-dependent DNA biosynthetic process
Contig_41259	9.9E-45	serine protease 23-like		proteolysis serine-type endopeptidase activity serine-type peptidase activity peptidase activity
Contig_41360	8.9E-134	tonsoku		nuclear lumen protein complex DNA metabolic process cellular response to DNA damage stimulus single-organism metabolic process single-organism cellular process
Contig_41417	3.3E-27	eukaryotic translation initiator factor 4E-binding 1		cytosol eukaryotic initiation factor 4E binding translational initiation signal transduction negative regulation of translational initiation
Contig_41500	1.8E-128	Cytoplasmic 1-like		cytoskeleton

				plasma membrane focal adhesion dense body	
Contig_42135	0	No description		membrane exopeptidase activity	
Contig_42435	3.8E-84	Tensin-partial	3	focal adhesion actin binding cell-substrate junction assembly fibroblast migration	
Contig_43083	2.0E-47	No description			
Contig_45623	0	kinesin KIF16B isoform X2		early endosome cytosol kinesin complex phosphatidylinositol-3,4,5-trisphosphate binding ATP-dependent microtubule motor activity, plus-end-directed phosphatidylinositol-3-phosphate binding phosphatidylinositol-3,4-bisphosphate binding phosphatidylinositol-3,5-bisphosphate binding formation of primary germ layer regulation of receptor recycling Golgi to endosome transport microtubule-based movement epidermal growth factor receptor signaling pathway endoderm development	

				fibroblast growth factor receptor signaling pathway cytoskeleton-dependent intracellular transport receptor catabolic process early endosome to late endosome transport	
Contig_45771	8.3E-94	No description		membrane dopamine neurotransmitter receptor activity G-protein coupled receptor signaling pathway	
Contig_47280	2.0E-168	No description		regulation of Rho protein signal transduction metabolic process Rho guanyl-nucleotide exchange factor activity transferase activity positive regulation of GTPase activity	
Contig_47344	0	allene oxide synthase-lipoxygenase		oxidoreductase activity, acting on single donors with incorporation of molecular oxygen, incorporation of two atoms of oxygen metal ion binding fatty acid metabolic process	
Contig_5993	1.5E-95	GA-binding subunit beta-partial		nucleus protein homodimerization activity transcription regulatory region DNA binding protein heterodimerization activity	

				transcription, DNA-templated positive regulation of transcription from RNA polymerase II promoter	
Contig_7742	0	Indole-3-acetaldehyde oxidase-like		oxidoreductase activity ion binding single-organism metabolic process	
Contig_8936	5.2E-105	drebrin isoform X1		actin filament binding synapse assembly ruffle assembly actin binding neuron projection morphogenesis receptor-mediated endocytosis intracellular	
Contig_8963	3.4E-169	tubuline delta chain		Intracellular part	
Contig_26377	6.7E-138	ribokinase isoform X1	2	ribokinase activity D-ribose metabolic process carbohydrate phosphorylation	Banyuls