Not so proportional after all?

Thomas M.H. Hope [1] *, Karl Friston [1], Cathy J. Price [1], Alex P. Leff [2,3],

Pia Rotshtein [4] & Howard Bowman [4,5]

1. Wellcome Centre for Human Neuroimaging, University College London, UK.
2. Institute of Cognitive Neuroscience, University College London, UK.
3. Centre for Neurorehabilitation, University College London, UK.
4. School of Psychology, University of Birmingham, UK.
5. School of Computing, University of Kent, UK.

* Corresponding author: Dr Thomas M.H. Hope, Wellcome Centre for Human Neuroimaging, Institute of Neurology, University College London, 12 Queen Square, London, WC1N 3AR, UK. Phone: +44 (0)20 3448 4345. E-mail: t.hope@ucl.ac.uk

References: 31

Keywords: proportional recovery, stroke, methods, statistics.

ABSTRACT

Background

The proportional recovery rule asserts that most stroke survivors recover a fixed proportion of lost function. Reports claiming that the rule can be used to predict recovery, extraordinarily accurately, are rapidly accumulating. Here, we show that the proportional recovery rule may not be as powerful as it seems.

Methods

We provide a formal analysis of the relationship between baseline scores and subsequent change, to highlight the shortcomings of the proportional recovery rule. These problems are illustrated using simulations in which outcomes are generated by different types of recovery processes.

Findings

The correlation between baseline scores and subsequent change is a function of: (a) the correlation between baseline scores and outcomes; and (b) the ratio of the standard deviations of outcomes to baseline scores. In this sense, the two correlations are always coupled. When outcomes are less variable than baselines, the correlation between baselines and change can be extreme regardless of the correlation between baselines and outcomes. This creates a misleadingly optimistic impression of how predictable outcomes really are. This scenario appears to be common in empirical studies of the proportional recovery rule, so we argue that future studies should at least report both correlations. Yet even when both are reported, our simulations suggest that they cannot distinguish proportional recovery from recovery which is not proportional at all.

Interpretation

Analyses relating baselines to change are a minefield: our formal analysis, which identifies when these relationships may be spurious, applies just as consistently outside the area of stroke as it does within it. As applied to recovery from stroke, these analyses do not reliably distinguish proportional recovery from other types of recovery, and will often yield inflated effect sizes even if recovery really is proportional. The implication is that proportional recovery rule is not as predictive, of real recovery, as recent empirical studies suggest.

Funding

Wellcome, the Medical Research Council and the Stroke Association.

1. INTRODUCTION

Stroke clinicians and researchers have long known that the severity of patients' initial symptoms after stroke is related to their long term outcomes[1]. Recent studies have suggested that this relationship is a great deal stronger than previously thought: that most patients recover a fixed proportion of lost function after stroke. Evidence in support of this 'proportional recovery rule' is rapidly accumulating, as summarised by Stinear in a recent review[2]. In five separate studies since 2015[3-7], researchers used the Fugl-Meyer test to assess stroke patients' upper limb motor function within two weeks after stroke onset, and then again either three or six months post-stroke. The results were consistent with earlier observations[8,9] that most patients recovered around 70% of lost upper limb function. Taken together, these studies report analyses involving over 500 patients, spread across different countries with different approaches to rehabilitation, regardless of the patients' ages at stroke onset, stroke type, sex, or therapy dose[2]. And there is increasing evidence that the proportional recovery rule also captures recovery from other post-stroke impairments, including impairments of lower limb function[10], attention[11,12], and language[12,13]: one recent study suggests that proportional recovery applies quite generally across cognitive domains[14]. Indeed, even rats appear to recover proportionally after stroke[15].

These results are especially compelling because of the extraordinary accuracy with which the proportional recovery rule appears to predict patients' recoveries. Most proportional recovery studies report that the rule explains more (and sometimes much more) than 80% of the variance in empirical recovery. The 2015 study by Winters and colleagues[3] reports that recovery predicted using the rule could explain a staggering 95% of the variance in the empirical recoveries of 146 stroke patients. When predicting behavioural performance in humans, these effect sizes are unprecedented. Like many of its counterparts[5,10,12], this study also reported a group of 'non-fitters', who did not make the predicted recovery. But if non-fitters can be distinguished from fitters at the acute stage, as the authors' results also suggest, the implication is that we can predict most patients' recovery near-perfectly, at least during the first few months after stroke onset. Significantly, this is true entirely irrespective of the patients' experiences of therapy, which in turn implies that those therapeutic interventions are ineffective. Stroke researchers are used to thinking of recovery as a complex, multi-factorial process[16-20]. If the proportional recovery rule is really as powerful as it seems, then much of what we thought we knew about recovery from stroke will need to be revisited[2,11,21].

In what follows, we argue that the proportional recovery rule may not be as powerful as it seems. At least as currently employed, the rule can yield effect sizes whose magnitude is misleadingly large, whether or not recovery is really proportional at all. This is because these analyses are all potentially confounded by mathematical coupling.

2. MATHEMATICAL COUPLING

Analyses relating baseline scores to subsequent change pose well-known challenges, which have been discussed by statisticians for decades[22,23]. Chief among these challenges, is the problem of 'mathematical coupling': we contend that this problem is endemic in studies of the proportional recovery rule, and that its severity has been underestimated. Mathematical coupling refers to

situations in which the same variable is employed to calculate both the independent and the dependent variables in an analysis. In analyses relating baselines to change, this happens because change is defined as 'outcome scores minus baseline scores': baseline scores are both the independent variable, and a component of the dependent variable.

For the sake of brevity in what follows, we define 'baselines' = X, 'outcomes' = Y, and 'change' (or recovery) = Δ: i.e. delta, or outcomes minus baselines. The 'correlation between baselines and outcomes' is r(X, Y), and the 'correlation between baselines and change' is r(X, Δ). Finally, we define the 'variability ratio' as the ratio of the standard deviation of Y to the standard deviation of X (σY/σX). X and Y are construed as lists of scores, with each entry being the performance of a single patient at the specified time point. We assume that performance improves as the numbers get bigger, which in turn means that r(X, Δ) will be negative if recovery is proportional to lost function, because in this case those with the most severe initial symptoms (i.e. the smallest X) will also recover the most. One can equally formulate the relationship with 'lost function' (i.e. maximum X minus actual X) instead of 'baseline score', but while this makes r(X, Δ) positive if recovery is proportional to lost function, it is otherwise entirely equivalent.

It has long been known that mathematical coupling can induce large, spurious effects in analyses relating baselines to change[24]. For example, when baselines (X) and outcomes (Y) are independent, random variables drawn from the standard normal distribution, and change (Δ) is Y-X, it has been shown that: r(X, Δ) ≈ -0.71[24]. In other words, there is a strong correlation between baselines and change, even though baselines and outcomes are entirely unrelated: r(X, Y) ≈ 0.

Another expression of this canonical example of mathematical coupling, which may be more germane to researchers interested in post-stroke prognoses, is that r(X, Δ) in this case delivers a misleading impression of how predictable Y really is. Following the approach taken in some empirical studies of the proportional recovery rule, we can derive predicted change (pΔ) from the least-squares line defined by r(X, Δ)[*], such that |r(X, Δ)| = |r(pΔ, Δ)|: i.e. the magnitudes of the correlations between: (a) baselines and change; and (b) predicted and empirical change, are the same[†] (see Supplementary Appendix A, proposition 8, bullet point 1). Correlations between predicted and empirical data are a popular measure of predictive accuracy: the stronger the correlation, the better the predictions. Since change (or recovery) is precisely the difference between baselines and outcomes, a result suggesting we can predict change accurately naturally encourages the expectation that we can predict outcomes *equally* accurately – by simply adding 'known baselines' to 'accurately predicted change' to produce 'accurately predicted outcomes'. Or more formally, the expectation is that: |r(X + pΔ, Y)| ≈ |r(X, Δ)|.

However, as might be obvious from our canonical example of mathematical coupling, this is a fallacy: in fact, |r(X + pΔ, Y)| = |r(X, Y)|. No matter how accurately we can seemingly predict recovery, the correlation between outcomes calculated using those predictions, and empirical outcomes, always has the same magnitude as the correlation between baselines and outcomes (see Supplementary Appendix A, proposition 8, bullet 2). In our canonical example, the impression that

---

[*] This is not a properly out-of-sample derivation, but predictions derived more properly, for example via cross-validation, will converge with that least-squares line in large samples.
[†] The emphasis on magnitudes is necessary here because r(X, Δ) may be positive or negative, but r(pΔ, Δ) should always be positive.

we can predict change relatively well (because r(pΔ, Δ) ≈ 0.71), is spurious because we cannot use those predictions to predict outcomes even nearly as well (because r(X, Y) ≈ 0).

3. IS MATHEMATICAL COUPLING LESS IMPORTANT IN PRACTICE THAN IN THEORY?

Krakauer and Marshall recently made the case that spurious effects associated with mathematical coupling are likely to be less significant for the proportional recovery rule in practice, than they might be in theory[21]. This is because empirical baselines and outcomes are dependent, repeated measures from the same patients, whereas in the canonical example, baselines and outcomes were independent random variables. The implication is that our canonical example of mathematical coupling is not a proper null hypothesis for analyses of proportional recovery.

Krakauer and Marshall use True Score theory [25], to advance a more plausible alternative, in which 'true' baselines become 'true' outcomes via some 'true' recovery trajectory, and observed baselines and outcomes are just the true states plus measurement noise. From this perspective, the null hypothesis is that baselines and outcomes are dependent, but linked by random recovery trajectories, and under this hypothesis, r(X, Δ) ≈ 0 regardless of the magnitude of recovery. Only when the variance of the measurement noise is large, so the argument goes, do baselines and outcomes start to behave as if they were independent, with the consequent emergence of spurious correlations between observed baselines and observed change. Since the most popular measurement tool in the literature on proportional recovery is highly reliable, it follows that the effects reported in these studies are unlikely to be spurious[21].

One potential weakness in this argument is that it only relates to assessments of hemiparesis after stroke using the Fugl-Meyer scale, which does appear to be remarkably reliable[26]. But the proportional recovery rule is perhaps particularly exciting because it also appears to apply quite generally, across cognitive domains, and none of the measurement tools employed in those wider studies is as reliable as the Fugl-Meyer scale. For example, while inter-rater and test-retest scores on the upper extremity subsection of the Fugl-Meyer scale are highly correlated (r > 0.95[26]), this correlation seems considerably weaker for scores on the letter cancellation task[27] used to apply the proportional recovery rule to visuospatial neglect[11]. In short, in every new domain to which the proportional recovery rule is applied, Krakauer and Marshall's argument needs to be made again.

Whether or not those arguments can be made, we contend that there are many more – and more likely – situations in which r(X, Δ) may be misleadingly strong, regardless of measurement noise. This is the focus of the next section.

4. RELATING THE CORRELATIONS BETWEEN BASELINES, OUTCOMES AND CHANGE

For any baselines (X) and outcomes (Y), defined as described previously, it can be shown that:

$$r(X, \Delta) = \frac{\sigma_Y \, . \, r(X,Y) - \sigma_X \, . \, r(X,X)}{\sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \, . \, \sigma_X \, . \, \sigma_Y . r(X,Y)}} \quad \text{(Equation 1)}$$

A formal proof of Equation 1 is provided in Supplementary Appendix A (see, proposition 4 and theorem 1); its core consequence is that $r(X, \Delta)$ is a (somewhat complex) function of: (a) $r(X, Y)$, and (b) $\sigma Y/\sigma X$ (and $r(X, X)$, which is 1). To illustrate that relationship, we ran a series of simulations in which both $r(X, Y)$ and $\sigma Y/\sigma X$ were varied independently: a Matlab script for this analysis is provided in Supplementary Appendix B, and the results are illustrated in Figure 1.

--Insert Figure 1--

Our canonical example of mathematical coupling (i.e., when X and Y are independent standard normal variables, and $r(X, \Delta) \approx -0.71$) is consistent with Equation 1, and corresponds to the red sphere marked in Figure 1. More generally, the figure illustrates that there are two principle regimes or modes of interaction between $r(X, Y)$ and $r(X, \Delta)$. When Y is more variable than X, the two correlations are more closely related, rising and falling together. However, when Y is less variable than X, this relationship ceases to hold, and $r(X, \Delta)$ can be extreme regardless of $r(X, Y)$. To see why this is so, consider an example in which baselines and outcomes are connected by proportional recovery: with no measurement noise, $r(X, Y) = 1$ and $r(X, \Delta) = -1$ (Figure 2a). When we shuffle the outcomes data, $r(X, Y) \approx 0$, because we have expressly removed the relationship between X and Y, but $r(X, \Delta)$ remains relatively strong ($r = -0.96$; see Figure 2b).

--Insert Figure 2--

Crucially, we expect outcomes to be less variable than baseline scores if most patients recover, and there are ceiling effects in the scale used to measure that recovery – as there always are in practice. Indeed, even without ceiling effects, we expect to be in this region of the surface if the patients with the most severe initial impairments also recover the most, as the proportional recovery rule predicts. In other words, even if recovery really is proportional, $r(X, \Delta)$ may be much more extreme than $r(X, Y)$, regardless of measurement noise, and regardless of whether we assume that something like Krakauer and Marshall's[21] True Score analysis holds. This is because, as the variability of Y decreases relative to that of X, $r(X, Y-X)$ tends toward $r(X, -X)$: see proposition 9 in Supplementary Appendix A.

5. RE-EXAMINING STUDIES OF PROPORTIONAL RECOVERY AFTER STROKE

The relationships described previously, between $r(X, Y)$, $r(X, \Delta)$ and the variability ratio of Y to X, merit a re-examination of the proportional recovery rule for stroke. The only study in this literature that reports individual patients' behavioural data is that by Zarahn and colleagues[9], who reported recovery from hemiparesis in 30 stroke patients. When measured across the whole sample, $r(X, Y) = 0.80$ and $r(X, \Delta) = -0.49$; this changes to $r(X, Y) = 0.75$ and $r(X, \Delta) = -0.95$, after removing the 7

patients (23%) who did not fit the rule. Calculating predicted recovery (pΔ) from the least-squares line for r(X, Δ), we get r(pΔ, Δ) = 0.95, but r(X + pΔ, Y) = 0.75: this confirms our reasoning in the first section on mathematical coupling, that we cannot predict Y as well as is implied by the effect size of r(pΔ, Δ). By removing those 7 non-fitters, Zarahn and colleagues dramatically increased the apparent predictability of recovery while actually reducing the apparent predictability of outcomes; they also reduced the variability ratio for these data, from 0.88 in the full sample to 0.36 in the reduced sample of fitters. Notably, the residuals or errors for both correlations (r(X, Y) and r(X, Δ)) are identical (see Figure 3), and in fact this is always true, as proved in Supplementary Appendix A, proposition 10. Since r(X, Y) and r(X, Δ) assign different effect sizes to the same errors, we contend that r(X, Δ) is misleadingly strong in this case.

--Insert Figure 3--

We can also use Equation 1 to reinterpret other studies in this literature, which do not report individual patient data. One example is the first study to apply the proportional recovery rule to aphasia after stroke, by Lazar and colleagues[13]. Baseline and follow-up scores in this study had standard deviations of 7.7 and 3.7 respectively (from their Table 1), so the variability ratio = 3.7/7.7 = 0.48. And the authors also reported that baselines alone explained 81% of the variance in recovery: i.e. R^2 = 0.81, so r(X, Δ) ≈ -0.9. Allowing for rounding in these figures, Equation 1 suggests that r(X, Y) is at most 0.78, but could also be zero: i.e. when outcomes and baselines are uncorrelated. We do not doubt that they were correlated in this study, but we do contend that r(X, Y) should be reported in cases like this.

A similar point emerges from the recent study of proportional recovery in rats by Jeffers and colleagues[15], who report that initial scores on their task of interest explained 49.9% of the variance in recovery of the 182 rats they classified as 'fitters' (from a total sample of 593) to the proportional recovery rule. These authors also report standard deviations of the baseline and outcomes variables of 4 and 3.2 respectively, defining a variability ratio of 0.8. By our equation 1, this implies that the correlation between baselines and outcomes in these data was either much stronger (>0.95) or considerably weaker (~0.29) than their correlation between baselines and change. Here again, we have learned rather less about the predictability of outcomes than the headline effect size might suggest.

Several authors prefer to report inter-quartile ranges (IQR) for their data instead of standard deviations. Accepting some room for error, we can estimate variability ratios from those IQRs. This lets us employ Equation 1 to reinterpret more studies, such as that by Winters and colleagues[3]. In Table 1 of their manuscript, these authors report median and inter-quartile ranges for Fugl-Meyer scores assigned to the 146 patients deemed to fit the proportional recovery rule (from a total sample of 211). From those inter-quartile ranges, we estimate a variability ratio of: 6 (follow-up) / 38 (baseline) = 0.158. By Equation 1, when the ratio is this small, r(X, Δ) < -0.98 entirely regardless of r(X, Y). This is a stronger correlation than that reported (r = -0.97), likely reflecting divergence between the IQR and standard deviations of their data. Nevertheless, as in the previous studies, our analysis suggests that their headline effect size tells us less about outcomes than expected, because r(X, Y) is not really constrained by the reported effect size for r(X, Δ). And the same issues arise from

the recent study of proportional recovery from lower-limb impairments, by Veerbeck and colleagues[28]. Here again, the variability ratio that we estimate from the inter-quartile ranges in Table 1 (0.438) implies that $r(X, \Delta)$ could be at least as strong as that reported ($R^2 = 0.768$, so $r \approx -0.88$) entirely regardless of $r(X, Y)$.

Finally, while we have emphasised that $r(X, \Delta)$ can be misleading if its magnitude is extreme relative to that of $r(X, Y)$, the reverse is also true. For example, the recent study by Feng and colleagues is also the only one in this literature that focuses on outcomes instead of recovery: i.e. reporting $r(X, Y)$ rather than $r(X, \Delta)$ [6]. In Table 1 of their manuscript, these authors report that the R^2 for 'outcome prediction' from 'initial motor impairment' was 0.64 for their 'combined group' (n = 76). Given the variability ratio which they also report in the same table (23.4/19.5 = 1.2: i.e. outcomes were 20% more variable than baselines), we can use Equation 1 to calculate that: $r(X, \Delta) = -0.05$. These authors only report proportional recovery in a subset of their sample, but it appears that their full sample was actually quite consistent with the null hypothesis, of random recovery, implied by Krakauer and Marshall's True Score analysis[21].

In light of our formal analysis, it is worth noting that all of the studies referenced in this section, except the last, had variability ratios < 1: i.e. outcomes were less variable than baselines. This is the predicted pattern if recovery is really proportional, but it is also the pattern that will tend to encourage much more extreme $r(X, \Delta)$ than $r(X, Y)$: even if patients really do recover proportionally, there is a risk that $r(X, \Delta)$ can deliver an inflated impression of how well the rule predicts their recovery. We found reason to suspect inflation in every study (apart from that by Feng and colleagues), which reported enough information for us to make the judgement. The most common reason why we could not make a judgement was when baselines were only related to recovery through multivariable models which included baseline scores[4,11,12]. But inflation in one variable's effect size will inflate the multivariable model's effect size as well: while we cannot estimate the magnitude of that inflation with any confidence for any particular model, we can predict that it's also there in those multivariable models.

As a consequence, we would argue that studies in this field should at a minimum report both $r(X, Y)$ and $r(X, \Delta)$, since the latter alone can encourage an inflated sense of how predictable outcomes really are. However, even when both of these correlations are reported, we contend that analyses like this cannot convincingly demonstrate that recovery is really proportional: this is the focus of the next section.

6. **HOW DO PATIENTS REALLY RECOVER?**

The principal consequence of our discussion so far, is that bivariate correlation (or regression) analyses may be a misleading way to test the proportional recovery rule, and that studies should at a minimum report both $r(X, Y)$ and $r(X, \Delta)$. However, even when both are reported, we cannot necessarily infer that recovery really is proportional. To illustrate this point, we report the results of a series of simulations in which these correlations are calculated from synthetic data, generated by different types of recovery process. All of the processes assume that a True Score analysis is essentially correct: i.e. that baselines and outcomes are dependent, and that each is observed with some measurement noise. And all of the analyses also assume that we have very large samples to

work with (N = 1,000); the distortions induced by small samples are well documented elsewhere[29], and will not concern us here.

Our simulations vary: (a) the magnitude of measurement noise; (b) the real variation in participants' recovery trajectories (i.e. 'recovery noise'); (c) the presence or absence of boundaries on the measurement scale (i.e. ceiling and floor effects); and (d) whether or not 'non-fitters' are removed from the analysis: i.e. those whose 'recovery' is less than expected. We also manipulate the real relationship between baselines and recovery, considering three different types or regimes: (i) random recovery; (ii) constant recovery (i.e. patients recover a fixed amount); and (iii) proportional recovery.

Each simulation starts by defining a random vector of 'true' initial states, drawn at random from the range 0-65 (i.e. scores indicating impairment on the upper extremity subsection of the Fugl-Meyer scale). Observed baselines are those true states plus measurement noise. Recovery, calculated using the selected regime, is then added to the true baselines to produce true outcomes, which are also observed with measurement noise. In each simulation, we calculate both $r(X, Y)$ and $r(X, \Delta)$. Each simulation is repeated 100 times, and results are mean values across those repeated simulations. The Matlab script required to run these simulations is included in Supplementary Appendix C.

Figure 4 displays the results: i.e. $r(X, Y)$ and $r(X, \Delta)$ with increasing recovery noise, with each configuration of recovery mechanism and other constraints. As expected given Krakauer and Marshall's True Score analysis[21], when measurements are noisy (Figure 3, row 1), mathematical coupling dominates and all of the recovery regimes yield largely the same results. Notably though, these results are all also distinguishable from empirical data, in that $r(X, Y) \approx 0$; whereas we expect this will rarely be true of baseline scores and outcomes measuring recovery after stroke. When measurements are more reliable (Figure 3, row 2), the results are more promising, in that proportional recovery yields strong $r(X, \Delta)$, like the effects reported in the literature, and also trajectories of $r(X, \Delta)$ with increasing recovery noise, which are dissimilar to the equivalent trajectories driven by different recovery mechanisms. In this case, one can reasonably hope to distinguish proportional recovery from other types of recovery, using $r(X, \Delta)$ alone.

However, the simulations in rows 1 and 2 of Figure 4 make the unrealistic assumption that the scale used to measure X and Y is infinitely long. In practice, all assessment tools employ a finite scale, and the Fugl-Meyer scale in particular is known to exert ceiling effects on patients with mild-to-moderate hemiparesis[26]. When we include these ceiling effects (Figure 4, row 3), the constant recovery regime produces results that are similar to those derived by proportional recovery: correlation coefficients cease to be an effective way to identify how our patients actually recover. And these similarities persist when we also remove non-fitters prior to calculating the correlations (Figure 4, last row), with the principal effect being to preserve extremely strong $r(X, \Delta)$ despite significant, real variability in the simulated recovery trajectories.

--Insert Figure 4--

These results demonstrate that, with more realistic constraints on our generative models of empirical data, r(X, Y) and r(X, Δ) become less useful in distinguishing how patients are really recovering. Moreover, they show that extreme r(X, Δ) are potentially consistent with quite dramatic, real variability in patients' recoveries: for example, in our final simulation of proportional recovery (row 4, column 2), r(X, Δ) ≈ -0.94 even when the standard deviation of the recovery noise is >0.35: since the mean proportion is 70% in these simulations, this means that the 95% confidence intervals on the proportion of lost function actually recovered include both 0% and 100%.

7. Discussion

The proportional recovery rule is attractive party because it formalises the popular, clinical intuition that initial symptom severity is by far the most, and perhaps the only, important predictor of recovery after stroke. The rule is also striking because it implies that recovery is simple and consistent across patients (non-fitters notwithstanding), and because that implication appears to be justified by an increasing number of extremely strong empirical results[3,7,9-14,21,28,30]. That incredible, empirical power has even begun to encourage the search for some deeper, general recovery mechanism which cuts across cognitive domains[21]. However, our results suggest that the proportional recovery rule may not be as powerful as it seems.

We are not claiming that the proportional recovery rule is necessarily wrong. Our analysis suggests that the empirical results to date do not confirm that the rule holds or how well, but those results are still broadly consistent with the predictions that the rule makes. Perhaps just as importantly, the rule is also falsifiable, as illustrated by a recent study of recovery from hemiparesis and aphasia during the first week after stroke onset, which found no evidence of proportional recovery[31]. Our concern is that estimates of the rule's predictive power – those headline effect sizes for r(X, Δ) or r(pΔ, Δ) – deliver a misleadingly optimistic impression of how predictable outcomes really are given baseline scores alone. And a misleadingly optimistic impression here encourages misleadingly pessimistic assessments elsewhere, for example of the likely impacts of therapeutic interventions for stroke survivors. If baseline scores really do predict recovery near-perfectly, those interventions may appear to be redundant, but as that predictive power decreases, so increases the variance, in outcomes, which interventions may purport to explain.

Much of the work in this area has focused on the distinction between fitters and non-fitters to the proportional recovery rule; distinguishing the groups via analyses of the residuals or errors of the least-squares line for r(X, Δ). This is a circular definition, but it defines groups which can then be distinguished in non-circular ways[6]. Notably, the residuals for r(X, Y) and r(X, Δ) are always the same (see Figure 2, and Supplementary Appendix A); so the fitters and non-fitters in a given sample will also be the same regardless of whether the dependent variable is outcomes or change. It follows that our concerns about analyses involving change have no bearing on the validity of work to distinguish fitters from non-fitters. That being said, extreme r(X, Δ) for patients classified as fitters will naturally encourage the assumption that, if they can be identified at the acute stage, we can predict those fitters' outcomes near-perfectly too. Our analysis suggests that this assumption is wrong: those outcomes were never more predictable, from baselines alone, than is implied by the relationship between baselines and outcomes. And indeed those outcomes are likely rather less predictable than the headline effect sizes in this literature suggest.

In summary, our results suggest that recovery from stroke might not be as proportional as it seems. Strong correlations between baseline scores and subsequent change, or between predicted and empirical change, can drive a misleadingly optimistic impression of how predictable outcomes really are. This situation appears to be common, both in our simulations and in the empirical literature on the proportional recovery rule. Moreover, even when reported together – as we suggest they should be (together with the variability ratio) – these analyses cannot distinguish proportional recovery from recovery which is either not consistently proportional, or not proportional at all. Quite how to make this distinction with confidence, remains an open question: this is a subject for future work.

8. A PARELLEL CHALLENGE?

During the final stages of our preparation of this manuscript, we discovered that another researcher, Robinson Kundert, was also working on a parallel challenge to the proportional recovery rule, an initial draft of which was distributed as a 'pre-paper' at a meeting. None of the authors attended the meeting, but we learned of it through colleagues who did attend. Our understanding is that the principal focus of this challenge is the consistency (or lack thereof) of the proportion of lost function that patients recover: in this sense, that work may be seen as complementary to our focus on predictive power. We look forward to studying this challenge in detail after it is published.

ACKNOWLEDGMENTS

This study was supported for the Medical Research Council (MR/M023672/1, MR/K022563/1), the Wellcome Trust (091593/Z/10/Z), and the Stroke Association (TSA PDF 2017/02). The funders had no participation in the design and results of this study.

REFERENCES

1.   Jongbloed L. Prediction of function after stroke: a critical review. *Stroke; a journal of cerebral circulation* 1986; **17**(4): 765-76.
2.   Stinear CM. Prediction of motor recovery after stroke: advances in biomarkers. *The Lancet Neurology*; **16**(10): 826-36.
3.   Winters C, Wegen EEH, Daffertshofer A, Kwakkel G. Generalizability of the Proportional Recovery Model for the Upper Extremity After an Ischemic Stroke. *Neurorehabilitation and neural repair* 2015; **29**.
4.   Buch ER, Rizk S, Nicolo P, Cohen LG, Schnider A, Guggisberg AG. Predicting motor improvement after stroke with clinical assessment and diffusion tensor imaging. *Neurology* 2016; **86**(20): 1924-5.
5.   Byblow WD, Stinear CM. Proportional upper limb recovery after stroke is predicated upon corticospinal tract integrity. *Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation*; **8**(2): 429-30.
6.   Feng W, Wang J, Chhatbar PY, et al. Corticospinal tract lesion load: An imaging biomarker for stroke motor outcomes. *Annals of neurology* 2015; **78**(6): 860-70.

7.   Stinear CM, Byblow WD, Ackerley SJ, Smith M-C, Borges VM, Barber PA. Proportional Motor Recovery After Stroke. *Implications for Trial Design* 2017; **48**(3): 795-8.

8.   Prabhakaran S, Zarahn E, Riley C, et al. Inter-individual variability in the capacity for motor recovery after ischemic stroke. *Neurorehabilitation and neural repair* 2007; **22**.

9.   Zarahn E, Alon L, Ryan SL, et al. Prediction of motor recovery using initial impairment and fMRI 48 h poststroke. *Cereb Cortex (New York, NY 1991)* 2011; **21**.

10.   Smith M-C, Byblow WD, Barber PA, Stinear CM. Proportional Recovery From Lower Limb Motor Impairment After Stroke. *Stroke; a journal of cerebral circulation* 2017.

11.   Winters C, Wegen EEHv, Daffertshofer A, Kwakkel G. Generalizability of the Maximum Proportional Recovery Rule to Visuospatial Neglect Early Poststroke. *Neurorehabilitation and neural repair* 2017; **31**(4): 334-42.

12.   Marchi NA, Ptak R, Di Pietro M, Schnider A, Guggisberg AG. Principles of proportional recovery after stroke generalize to neglect and aphasia. *European Journal of Neurology* 2017; **24**(8): 1084-7.

13.   Lazar RM, Minzer B, Antoniello D, Festa JR, Krakauer JW, Marshall RS. Improvement in aphasia scores after stroke is well predicted by initial severity. *Stroke; a journal of cerebral circulation* 2010; **41**(7): 1485-8.

14.   Ramsey LE, Siegel JS, Lang CE, Strube M, Shulman GL, Corbetta M. Behavioural clusters and predictors of performance during recovery from stroke. *Nature human behaviour* 2017; **1**: 0038.

15.   Jeffers MS, Karthikeyan S, Corbett D. Does Stroke Rehabilitation Really Matter? Part A: Proportional Stroke Recovery in the Rat. *Neurorehabilitation and neural repair* 2018; **32**(1): 3-6.

16.   Plowman E, Hentz B, Ellis C, Jr. Post-stroke aphasia prognosis: a review of patient-related and stroke-related factors. *Journal of evaluation in clinical practice* 2012; **18**(3): 689-94.

17.   Kwakkel G, Wagenaar RC, Kollen BJ, Lankhorst GJ. Predicting disability in stroke--a critical review of the literature. *Age and ageing* 1996; **25**(6): 479-89.

18.   Jorgensen HS, Nakayama H, Raaschou HO, Vive-Larsen J, Stoier M, Olsen TS. Outcome and time course of recovery in stroke. Part I: Outcome. The Copenhagen Stroke Study. *Archives of physical medicine and rehabilitation* 1995; **76**(5): 399-405.

19.   Allen CM. Predicting recovery after acute stroke. *British journal of hospital medicine* 1984; **31**(6): 428, 30, 32-4.

20.   Millikan CH, Moersch FP. Factors that influence prognosis in acute focal cerebrovascular lesions. *AMA archives of neurology and psychiatry* 1953; **70**(5): 558-62.

21.   Krakauer JW, Marshall RS. The proportional recovery rule for stroke revisited. *Annals of neurology* 2015; **78**(6): 845-7.

22.   Oldham PD. A note on the analysis of repeated measurements of the same subjects. *Journal of Chronic Diseases* 1962; **15**(10): 969-77.

23.   Blomqvist N. On the relation between change and initial value. *Journal of the American Statistical Association* 1977; **72**(360a): 746-9.

24.   Tu YK, Gilthorpe MS. Revisiting the relation between change and initial value: a review and evaluation. *Statistics in medicine* 2007; **26**(2): 443-57.

25.   Traub RE. Classical test theory in historical perspective. *Educational Measurement: Issues and Practice* 1997; **16**(4): 8-14.

26.   Gladstone DJ, Danells CJ, Black SE. The Fugl-Meyer Assessment of Motor Recovery after Stroke: A Critical Review of Its Measurement Properties. *Neurorehabilitation and neural repair* 2002; **16**(3): 232-40.

27.   Pradhan B, Nagendra HR. Normative data for the letter-cancellation task in school children. *International Journal of Yoga* 2008; **1**(2): 72-5.

28.   Veerbeek JM, Winters C, van Wegen EEH, Kwakkel G. Is the proportional recovery rule applicable to the lower limb after a first-ever ischemic stroke? *PLOS ONE* 2018; **13**(1): e0189279.

29.   Button KS, Ioannidis JPA, Mokrysz C, et al. Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* 2013; **14**(5): 365-76.

30.   Prabhakaran S, Zarahn E, Riley C, et al. Inter-individual variability in the capacity for motor recovery after ischemic stroke. *Neurorehabilitation and neural repair* 2008; **22**(1): 64-71.

31.   Dunn LE, Schweber AB, Manson DK, et al. Variability in Motor and Language Recovery during the Acute Stroke Period. *Cerebrovascular diseases extra* 2016; **6**(1): 12-21.

Figure 1

Figure 1: The relationship between r(X, Y), r(X, Δ) and the variability ratio of Y to X. Note that the x-axis is log-transformed to ensure symmetry around 1 (i.e. when X and Y are equally variable, and log(σY/σX) = 0). The two major regimes of Equation 1 are also marked in red. In Regime 1, the imprint of baselines (X) dominates, and r(X, Δ) tends toward r(X, -X) (i.e. -1). In Regime 2, the correlations of interest are more tightly related, and r(X, Δ) ≈ r(X, Y). And there is a transition between the two regimes when the variability ratio is not dramatically skewed one way or the other – though even this transition allows for misleadingly strong r(X, Δ), as evidenced by the fact that our canonical example of mathematical coupling occurs in this region. This is indicated by the red mark, which corresponds to the situation where X and Y are independent random normal variables: i.e. r(X, Y) ≈ 0; log variability ratio = 0; and r(X, Δ) ≈ -0.71. Proposition 7 in Supplementary Appendix A provides a justification for unambiguously using a ratio of standard deviations in this figure, rather than σY and σX as separate axes.

Figure 2: Illustrating how |r(X, Δ)| > |r(X, Y)| when X is more variable than Y. In both panels, X is a vector of 20 numbers drawn from a random uniform distribution in the range 1-65. In the left panel, Y is calculated using the proportional recovery rule: i.e. Y = X + ((66-X) x 0.7). In this case the correlation between X and Y is 1 and the correlation between X and change is -1. In the right panel, Y is randomly shuffled: X and Y are now uncorrelated, but the correlation between X and change remains extremely strong (r = -0.96).

Figure 3: (Left) Least squares linear fits for analyses relating baselines to (upper) outcomes and (lower) recovery, using the fitters' data reported by Zarahn and colleagues[9]. (Middle) Plots of residuals relative to each least squares line, against the fitted values in each case. (Right) A scatter plot of the residuals from the model relating baselines to change, against the residuals from the model relating baselines to outcomes: the two sets of residuals are the same.

Figure 4

Figure 4: Simulation results. Plotting r(X, Y) and r(X, Δ) as a function of increasing true variability in recovery trajectories, where recovery is: (a) random; (b) proportional; or (c) constant. The top row displays results when the variance of measurement noise is large relative to the variance of baselines. The second row presents results when measurement noise is minimal. The third row presents results when measurement noise is minimal but there are ceiling effects on the measurement scale. And the last row presents results when measurement noise is minimal, ceiling effects are applied, and 'non-fitters' are removed: i.e. the ~19% (this is the figure in [3]) of the sample whose actual recovery deviates most strongly and negatively from the least squares line defined by r(X, Δ).

Supplementary Appendix A: formal relationships between the correlations

We present a very simple and general formulation of the proportional recovery concept. We assume two variables X' and Y' corresponding to performance at initial test (X') and at second test (Y'). These will be represented as column vectors, with each entry being the performance of a single patient and vector lengths being $N \in \mathbb{N}$. Performance improves as numbers get bigger, up to a maximum, denoted Max, which corresponds to no discernible deficit. Severity is measured as difference from maximum, i.e. $Max - X'$.

The two variables (X' and Y') could be specialised to more detailed formulations: e.g., true score theory or with an explicit modelling of measurement or state error. However, this would not impact any of the derivations or inferences that follow. Indeed, the results that we present would hold even in the complete absence of measurement noise, which has been considered the main concern for the validity of quantifications of proportional recovery.

Demeaning

Without loss of generality, we work with demeaned variables. That is, where over-lining denotes mean, we define new variables as,

$$X = X' - \bar{X}'$$

$$Y = Y' - \bar{Y}'$$

This also means that recovery, i.e. $Y - X$, will be demeaned, since, using proposition 1, the following holds.

$$Y - X = (Y' - \bar{Y}') - (X' - \bar{X}') = (Y' - X') - (\bar{Y}' - \bar{X}') = (Y' - X') - \overline{(Y' - X')}$$

Proposition 1

Let V and W be vectors of the same length, denoted N. Then, the following holds,

$$\bar{V} + \bar{W} = \overline{(V + W)}$$

with $\bar{V} - \bar{W} = \overline{(V - W)}$ as a trivial consequence.

Proof

By distributivity of multiplication through addition and associativity of addition, the following holds.

$$\bar{V} + \bar{W} = \left(\frac{1}{N}\sum_{i=1}^{N} V_i\right) + \left(\frac{1}{N}\sum_{i=1}^{N} W_i\right) = \frac{1}{N}\left(\sum_{i=1}^{N} V_i + \sum_{i=1}^{N} W_i\right) = \frac{1}{N}\left(\sum_{i=1}^{N}(V_i + W_i)\right) = \overline{(V + W)}$$

QED

Correlations

There are two basic correlations we are interested in, (1) the correlation between initial performance and performance at second test, i.e. $r(X, Y)$, and (2) the correlation between initial performance and recovery, i.e. $r(X, Y - X) = r(X, \Delta)$. The latter of these is the key relationship, and we would expect this to be a negative correlation; that is, as initial performance is smaller (i.e. further from Max), the larger is recovery. (One could also formulate the correlation as $r((Max - X), Y - X)$, which would flip the correlation to positive, but the two approaches are equivalent).

Our main correlations are defined as follows,

$$r(X, Y) = \frac{\sum_{i=1}^{N} X_i \cdot Y_i}{(N-1) \cdot \sigma_X \cdot \sigma_Y}$$

$$r(X, (Y-X)) = \frac{\sum_{i=1}^{N} (X_i \cdot (Y_i - X_i))}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}}$$

Standard Deviation of a Difference

We need a straightforward result on the standard deviation of a difference.

Proposition 2

$$\sigma_{(A-B)} = \sqrt{\sigma_A^2 + \sigma_B^2 - 2 \cdot cov(A, B)}$$

Proof

The result is a direct consequence of the following standard result from probability theory, e.g. see Ross, S. M. (2014). *Introduction to probability and statistics for engineers and scientists*. Academic Press.,

$$\sigma_{(A-B)}^2 = \sigma_A^2 + \sigma_B^2 - 2 \cdot cov(A, B)$$

Key Results

The following proposition enables us to express the key correlation, $r(X, (Y-X))$, in terms of covariance of its constituent variables.

Proposition 3

$$r(X, (Y-X)) = \frac{cov(X, Y) - cov(X, X)}{\sigma_X \cdot \sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \cdot cov(X, Y)}}$$

Proof

Using distributivity of multiplication through addition, associativity of addition, the definition of covariance and proposition 2, we can reason as follows.

$$r(X, (Y-X)) = \frac{\sum_{i=1}^{N} (X_i \cdot (Y_i - X_i))}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}} = \frac{\sum_{i=1}^{N} (X_i Y_i - X_i X_i)}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}} = \frac{\sum_{i=1}^{N} (X_i Y_i) - \sum_{i=1}^{N} (X_i X_i)}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}}$$

$$= \frac{cov(X, Y) - cov(X, X)}{\sigma_X \cdot \sigma_{(Y-X)}} = \frac{cov(X, Y) - cov(X, X)}{\sigma_X \cdot \sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \cdot cov(X, Y)}}$$

QED

It is straightforward to adapt proposition 3 to be fully in terms of correlations.

Proposition 4

$$r(X, (Y-X)) = \frac{\sigma_Y \cdot r(X, Y) - \sigma_X \cdot r(X, X)}{\sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \cdot \sigma_X \cdot \sigma_Y \cdot r(X, Y)}}$$

Proof

Straightforward from proposition 3 and definition of correlations, which gives the relationship $cov(A, B) = \sigma_A . \sigma_B . r(A, B)$. QED

Scale Invariance

The next set of propositions justifies working with a standardised X variable.

Lemma 1

$$\forall c \in \mathbb{R} \cdot |c|. \sigma_A = \sigma_{(c.A)}$$

Proof

Using distributivity of a multiplicative constant through averaging, $\sqrt{d^2} = |d|$ and distributivity of square root through multiplication, we can reason as follows.

$$\sigma_{(c.A)} = \sqrt{\frac{\sum_{i=1}^{N} (c.A_i - \overline{c.A})^2}{N-1}} = \sqrt{\frac{\sum_{i=1}^{N} (c.A_i - c.\bar{A})^2}{N-1}} = |c| . \sqrt{\frac{\sum_{i=1}^{N} (A_i - \bar{A})^2}{N-1}} = |c|. \sigma_A$$

QED

Proposition 5 (Invariance to scaling)

The absolute magnitude of a correlation is not changed by scaling either variable by a constant, i.e.

$$\forall c \in \mathbb{R} \cdot r(A, B) = sign(c).r(c.A, B) = sign(c).r(A, c.B)$$

where $sign(d) = if \ (d < 0) \ then -1 \ else +1$.

Proof

For any $c \in \mathbb{R}$, using distributivity of multiplication through mean and addition, and lemma 1, the following holds,

$$r(c.A, B) = \frac{\sum_{i=1}^{N} (c.A_i - \overline{c.A})(B_i - \bar{B})}{(N-1).\sigma_{(c.A)} \sigma_B} = \frac{\sum_{i=1}^{N} (c.A_i - c.\bar{A})(B_i - \bar{B})}{(N-1).\sigma_{(c.A)} \sigma_B}$$

$$= \frac{c.\sum_{i=1}^{N} (A_i - \bar{A})(B_i - \bar{B})}{(N-1).|c|.\sigma_A.\sigma_B} = \frac{sign(c).\sum_{i=1}^{N} (A_i - \bar{A})(B_i - \bar{B})}{(N-1).\sigma_A.\sigma_B} = sign(c).r(A, B)$$

Then, one can multiply both sides by $sign(c)$ to obtain $r(A, B) = sign(c).r(c.A, B)$. Additionally, as correlations are symmetric, $sign(c).r(c.B, A) = sign(c).r(A, c.B)$, and the full result follows.

QED

Corollary 1

$$\forall c \in \mathbb{R} \cdot r(A, B) = r(c.A, c.B)$$

Proof

Follows from twice applying proposition 5, and that $sign(c)^2 = +1$. QED

Proposition 6

$$\forall c \in \mathbb{R} \cdot r(X, (Y - X)) = r(c.X, (c.Y - c.X))$$

Proof

We can use distributivity of multiplication through subtraction and corollary 1 to give us the following.

$$r(c.X, (c.Y - c.X)) = r(c.X, c.(Y - X)) = r(X, (Y - X))$$

QED

It follows from proposition 6 that we can work with a standardised X variable, since,

$$r(X/\sigma_X, (Y/\sigma_X - X/\sigma_X)) = r(X, (Y - X))$$

Proposition 7 (Sufficiency of variability ratio)

Assume two pairs of variables: X_1, Y_1 and X_2, Y_2, such that, $r(X_1, Y_1) = r(X_2, Y_2)$, then,

$$\frac{\sigma_{Y_1}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{X_2}} \implies r(X_1, (Y_1 - X_1)) = r(X_2, (Y_2 - X_2))$$

Proof

The proof has two parts.

1) We consider the implications of equality of ratio of standard deviations. Firstly, we note that,

$$\frac{\sigma_{Y_1}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{X_2}} \iff \frac{\sigma_{X_2}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{Y_1}} \quad (eqn\ ratios)$$

Secondly, using eqn ratios, we can argue as follows,

$$\frac{\sigma_{Y_1}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{X_2}} \iff \left(\sigma_{Y_2} = \frac{\sigma_{X_2}}{\sigma_{X_1}}\sigma_{Y_1} \land \sigma_{X_2} = \frac{\sigma_{Y_2}}{\sigma_{Y_1}}\sigma_{X_1}\right) \iff \left(\sigma_{Y_2} = \frac{\sigma_{X_2}}{\sigma_{X_1}}\sigma_{Y_1} \land \sigma_{X_2} = \frac{\sigma_{X_2}}{\sigma_{X_1}}\sigma_{X_1}\right)$$

$$\implies \left(\exists d \in \mathbb{R} \cdot \sigma_{Y_2} = d.\sigma_{Y_1} \land \sigma_{X_2} = d.\sigma_{X_1}\right)$$

2) Using 4, the fact that $r(X_1, Y_1) = r(X_2, Y_2)$, the property just derived in part 1), with $d = \frac{\sigma_{X_2}}{\sigma_{X_1}}$ and rules of square roots, we can reason as follows,

$$r(X_2, (Y_2 - X_2)) = \frac{\sigma_{Y_2}.r(X_2, Y_2) - \sigma_{X_2}.r(X_2, X_2)}{\sqrt{\sigma_{Y_2}^2 + \sigma_{X_2}^2 - 2.\sigma_{X_2}.\sigma_{Y_2}.r(X_2, Y_2)}} = \frac{\sigma_{Y_2}.r(X_1, Y_1) - \sigma_{X_2}.r(X_1, X_1)}{\sqrt{\sigma_{Y_2}^2 + \sigma_{X_2}^2 - 2.\sigma_{X_2}.\sigma_{Y_2}.r(X_1, Y_1)}}$$

$$= \frac{d.\sigma_{Y_1}.r(X_1, Y_1) - d.\sigma_{X_1}.r(X_1, X_1)}{\sqrt{d^2.\sigma_{Y_1}^2 + d^2.\sigma_{X_1}^2 - 2.d.\sigma_{X_1}.d.\sigma_{Y_1}.r(X_1, Y_1)}} = \frac{d.(\sigma_{Y_1}.r(X_1, Y_1) - \sigma_{X_1}.r(X_1, X_1))}{d.\sqrt{\sigma_{Y_1}^2 + \sigma_{X_1}^2 - 2.\sigma_{X_1}.\sigma_{Y_1}.r(X_1, Y_1)}}$$

$$= r(X_1, (Y_1 - X_1)).$$

QED

Proposition 8

If $\Delta = Y - X$ and $p\Delta = X.\beta$, where $\beta \in \mathbb{R}$, then,

1) $r(p\Delta, \Delta) = sign(\beta).r(X, \Delta)$; and
2) $r(X + p\Delta, Y) = sign(1 + \beta).r(X, Y)$.

Proof

Both results are easy consequences of proposition 5.

1) $r(p\Delta, \Delta) = r(X.\beta, \Delta) = sign(\beta).r(X, \Delta).$

2) $r(X + p\Delta, Y) = r((X + (X.\beta)), Y) = r((X.(1 + \beta)), Y) = sign(1 + \beta).r(X, Y) = r(X, Y).$

QED

Main Findings

Theorem 1:

Since X will be standardised, we can adapt the finding in proposition 4, to give us the key relationship we need,

$$r\big(X, (Y - X)\big) = \frac{\sigma_Y \,.r(X,Y) - r(X,X)}{\sqrt{\sigma_Y^2 + 1 - 2\,.\sigma_Y.r(X,Y)}} \qquad \text{(eqn } Imprint)$$

Proof

Immediate from proposition 4.   QED

Theorem 1 shows clearly that $r(X, (Y - X))$ is fully defined by the correlations $r(X,Y)$ and $r(X,X)$, along with the variability of Y. The correlation of X with itself, i.e. $r(X,X)$, is a prominent aspect of this equation, which drives its oddities. $r(X,X)$ reflects the coupling in the equation that arises because X appears in both the terms being correlated in $r(X, (Y - X))$. $r(X,X)$ is of course a constant, i.e. 1 for any X, so in fact, σ_Y and $r(X,Y)$, are the only variables; accordingly, their size determines the extent to which the imprint of X in $Y - X$ drives $r(X, (Y - X))$.

This leads to the key observation that, as σ_Y gets smaller, $r(X, (Y - X))$ tends towards $-r(X,X)$, which equals -1. In other words, as the variability of Y decreases, the imprint of X becomes increasingly prominent. This is shown in the next proposition.

Proposition 9

$$r\big(X, (Y - X)\big) \longrightarrow -r(X,X) = -1, \qquad \text{as } \sigma_Y \longrightarrow 0$$

Proof

The right hand side of equation *Imprint*, has five constituent terms, two in the numerator and three in the denominator. Of these five, three are products with the standard deviation of Y, i.e. σ_Y. Assuming all else is constant, as σ_Y reduces, the absolute value of each of these three terms reduces towards zero. The rate of reduction is different amongst the three, but they will all decrease. Accordingly, as σ_Y decreases, $r\big(X, (Y - X)\big)$ becomes increasingly determined by the two terms not involving σ_Y, and thus, it tends towards $-\frac{r(X,X)}{\sqrt{+1}} = -r(X,X) = -1$.
QED

Equality of Residuals

An important finding of section 5 of the main text, is that the residuals resulting from regressing Y onto X are the same as regressing Y-X onto X. We show in this section, that this equality of residuals is necessarily the case.

We focus on the following two equations,

Eqn 1) $Y = \tilde{X}.\beta_1 + \varepsilon_1$

Eqn 2) $Y - X = \tilde{X}.\beta_2 + \varepsilon_2$

where $\tilde{X}$ is the $N \times 2$ matrix, with first column being X and second being the $N \times 1$ vector of ones (which provides the intercept term); β_1 and β_2 are 2×1 vectors of parameters and Y, X, ε_1 and ε_2 are $N \times 1$ vectors. As in the rest of this document, Y and X are our (demeaned) initial and outcome variables, while ε_1 and ε_2 are our residual error terms.

Proposition 10

If we assume that β_1 and β_2 are fit with ordinary least squares, with ε_1 and ε_2 the associated residuals, then, $\varepsilon_1 = \varepsilon_2$.

Proof

Under ordinary least squares, the parameters are set as follows.

$$\beta_1 = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T Y \quad \text{(Eqn 3)}$$

$$\beta_2 = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T (Y - X) \quad \text{(Eqn 4)}$$

We start with the second of these, and using left distributivity of matrices, and then substituting Eqn 3, we obtain the following.

$$\beta_2 = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T (Y - X) = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T Y - (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T X = \beta_1 - (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T X$$

Using the fact that the variable X is demeaned, we can now evaluate the main term here as follows,

$$\beta_2 = \beta_1 - (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T X = \beta_1 - \begin{pmatrix} X^2 & \Sigma X \\ \Sigma X & N \end{pmatrix}^{-1} \begin{pmatrix} X^2 \\ \Sigma X \end{pmatrix} = \beta_1 - \frac{1}{A} \begin{pmatrix} N & -\Sigma X \\ -\Sigma X & X^2 \end{pmatrix} \begin{pmatrix} X^2 \\ \Sigma X \end{pmatrix}$$

where X^2 is the dot product of X with itself, ΣX is the sum of the vector X, and $A = NX^2 - \Sigma X \Sigma X$ is the determinant of the matrix being inverted. From here we can derive the following,

$$\beta_2 = \beta_1 - \frac{1}{A} \begin{pmatrix} NX^2 - \Sigma X \Sigma X \\ -\Sigma X.X^2 + X^2.\Sigma X \end{pmatrix} = \beta_1 - \frac{1}{A} \begin{pmatrix} A \\ 0 \end{pmatrix} = \beta_1 - \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

We can then substitute this equality for β_2 in eqn 2 and re-arrange to obtain,

$$Y - X = \tilde{X}\beta_2 + \varepsilon_2 = \tilde{X}\left(\beta_1 - \begin{pmatrix} 1 \\ 0 \end{pmatrix}\right) + \varepsilon_2 = \tilde{X}\beta_1 - X + \varepsilon_2$$

It follows straightforwardly from here that,

$$Y - \tilde{X}\beta_1 = \varepsilon_2$$

i.e. $\varepsilon_1 = \varepsilon_2$, as required. QED

Proposition 10 shows that the residuals resulting from fitting equations 1 and 2 will be the same. A consequence of this is that the error variability will be the same. As a result of this, the factor that determines whether more variance is explained when regressing Y onto X or when regressing $Y - X$

onto X, is the variance available to explain. That is, the relative variance of Y and $Y - X$ drive the R^2 values of these two regressions. This then implicates the variance of Y and X and in fact their covariance (which impacts the variance of $Y - X$).

More precisely, we can state the following.

1) If $\sigma^2_{(Y-X)}$ is big relative to σ^2_Y, then regressing $Y - X$ onto X will explain more variability than regressing Y onto X.

2) If $\sigma^2_{(Y-X)}$ is small relative to σ^2_Y, then regressing $Y - X$ onto X will explain less variability than regressing Y onto X.

Supplementary Appendix B: illustrating the relationship between the correlations

```matlab
% This function illustrates the relationship
function [r_XY,std_Y,r2,r3] = CheckEqn1()

noise = [0.01:0.01:1,2:100]; % controls r(X, Y)
scale = [0.01:0.01:1,2:100]; % controls sigma_Y/sigma_X
X = single(randn(1000,1));
for j=1:length(noise)
    Y = X + single(randn(1000,1).*noise(j)); %Y is X plus noise
    Y = zscore(Y); % then scale to X so the actual scaling is consistent
    for k=1:length(scale)
        Yl = Y.*scale(k); % rescale to control the variability ratio
        r_XY(j,k) = corr(X,Y); % calculate the correlation with outcomes

        r2(j,k) = corr(X,Yl-X); % calculate the correlation with change
        std_Y(j,k) = std(Yl)./std(X); % record the variability ratio
        r3(j,k) = eqn_r_X_XminusY(r_XY(j),std_Y(j,k)); % check Equation 1
    end
end

% display the resulting surface (Figure 1)
figure,surf(log(std_Y),r_XY,r3,'edgecolor','none')
lighting flat
l = light('Position',[50 100 100]);
l = light('Position',[50 100 -50]);
l = light('Position',[50 -100 -50]);
l = light('Position',[-50 -15 29]);
l = light('Position',[-50 -15 -29]);
l = light('Position',[-50 15 -29]);
l = light('Position',[50 15 -29]);
l = light('Position',[50 15 -50]);
shading interp
xlabel('log ( sigmaY / sigmaX )')
ylabel('r(X, Y)')
zlabel('r(X, Y-X)')

% confirm that equation 1 does actually match 'empirical' r(X,Y-X)
figure,scatter(r2(:),r3(:))
xlabel('Empirical coefficients')
ylabel('Derived coefficients')

end

% This function implements Equation 1
function res = eqn_r_X_XminusY(r_XY,std_Y)

res = (((r_XY.*std_Y) - 1) ./ sqrt(1 + (std_Y).^2 - (2*(r_XY.*std_Y)))));

end
```

Supplementary Appendix C: simulating recovery

```matlab
function out = mprr_sim_pub(measure_noises, squash, nonfit)

% measure_noises = 300 in the 'high noise' case, and otherwise 3

% squash = true, if ceiling effects are imposed

% nonfit = true, if the 19 percent whose residuals
% to r(X, delta) are most extremely negative, are removed. The
% 19 percent figure comes from the study by Winters and
% colleagues, 2015

sample_size = 1000; % big sample
recover_noises = 0:0.01:1; % true variation in recovery
repeats = 100;
z=0; % flag to zscore everything before correlating

for r=1:repeats
    for s=1:length(sample_size)
        ss = sample_size(s);
        for m=1:length(measure_noises)
            mm = measure_noises(m);
            for rec=1:length(recover_noises)
                rr = recover_noises(rec);

                rX = randi(66,ss,1) - 1; % true baseline
                initial_std = std(rX);
                epsilon = randn(ss,1).*mm; % baselines measurement noise
                epsilon2 = randn(ss,1).*mm; % outcome measurement noise
                X = rX + epsilon; % observed baseline

                % random recovery
                rR_rand = randn(ss,1).*(rr.* initial_std .*10);
                Y_rand = rX + rR_rand + epsilon2; % observed outcome
                if(squash>0)
                    Y_rand(Y_rand>66) = 66;
                    Y_rand(Y_rand<0) = 0;
                end

                % proportional recovery
                rR_prop = (66 - rX).*(0.7+(randn(ss,1).*rr)); % 70% of lost function
                Y_prop = rX + rR_prop + epsilon2;% observed outcome
                if(squash>0)
                    Y_prop(Y_prop>66) = 66;
                    Y_prop(Y_prop<0) = 0;
                    %Y_prop = Y_sort(idx);
                end

                % constant recovery
                rR_const = 66.*(0.7+(randn(ss,1).*rr)); % 70% of 50
                Y_const = rX + rR_const + epsilon2;% observed outcome
                if(squash>0)
                    Y_const(Y_const>66) = 66;
                    Y_const(Y_const<0) = 0;
                end
                if(z>0)
                    X=zscore(X);
                    Y_rand=zscore(Y_rand);
                    Y_prop=zscore(Y_prop);
                    Y_invprop=zscore(Y_invprop);
                    Y_const=zscore(Y_const);
                end
                if(nonfit>0) % remove the bottom 19% of recoverers.
                    mdl = fitlm(X,Y_const);
                    [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
                    s_const = (1:sample_size)';
                    s_const = ~ismember(s_const, idx(1:ceil(sample_size*0.19)));
                    mdl = fitlm(X,Y_rand);
                    [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
                    s_rand = (1:sample_size)';
```

```matlab
                    s_rand = ~ismember(s_rand, idx(1:ceil(sample_size*0.19)));
                    mdl = fitlm(X,Y_prop);
                    [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
                    s_prop = (1:sample_size)';
                    s_prop = ~ismember(s_prop, idx(1:ceil(sample_size*0.19)));
                    mdl = fitlm(X,Y_invprop);
                    [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
                    s_invprop = (1:sample_size)';
                    s_invprop = ~ismember(s_invprop, idx(1:ceil(sample_size*0.19)));
                else
                    s_const = (1:sample_size)';
                    s_rand = (1:sample_size)';
                    s_prop = (1:sample_size)';
                    s_invprop = (1:sample_size)';
                end

                % run the correlations
                r_rand(r,rec,:) = [corr(X(s_rand),[Y_rand(s_rand), Y_rand(s_rand) -
X(s_rand)]),log(var(Y_rand(s_rand))/var(X(s_rand)))];
                r_prop(r,rec,:) = [corr(X(s_prop),[Y_prop(s_prop), Y_prop(s_prop) -
X(s_prop)]),log(var(Y_prop(s_prop))/var(X(s_prop)))];
                r_const(r,rec,:) = [corr(X(s_const),[Y_const(s_const), Y_const(s_const) -
X(s_const)]),log(var(Y_const(s_const))/var(X(s_const)))];
            end
        end
    end
end
r_rand=squeeze(nanmean(r_rand,1));
r_prop=squeeze(nanmean(r_prop,1));
r_const=squeeze(nanmean(r_const,1));
out = [r_rand,r_prop,r_const];
```