Recovery after stroke: not so proportional after all?

Thomas M.H. Hope [1] *, Karl Friston [1], Cathy J. Price [1], Alex P. Leff [2,3],

Pia Rotshtein [4] & Howard Bowman [4,5]

1. Wellcome Centre for Human Neuroimaging, University College London, UK.
2. Institute of Cognitive Neuroscience, University College London, UK.
3. Dept. of Brain Repair and Rehabilitation, Institute of Neurology, University College London,
UK.
4. School of Psychology, University of Birmingham, UK.
5. School of Computing, University of Kent, UK.

* Corresponding author: Dr Thomas M.H. Hope, Wellcome Centre for Human Neuroimaging, Institute
of Neurology, University College London, 12 Queen Square, London, WC1N 3AR, UK. Phone: +44 (0)20
3448 4345. E-mail: t.hope@ucl.ac.uk

References: 29

Words: (abstract): 253; (body): 3,497

Keywords: proportional recovery, stroke, methods, statistics.

Author contributions: T.M.H.H, P.R. and H.B. conceived the analysis. T.M.H.H., H.B. and K.F.
developed the core argument; H.B. derived the mathematical proofs; and T.M.H.H. developed the
simulations. C.J. and A.P.L. supported the interpretation of the results. T.M.H.H. lead the writing of
the manuscript, though all authors contributed to this.

ABSTRACT

Background

The proportional recovery rule asserts that most stroke survivors recover a fixed proportion of lost

function. Reports that the rule can be used to predict recovery, extraordinarily accurately, are

rapidly accumulating. Here, we show that the rule may not be as powerful as it seems.

Methods

We provide a formal analysis of the relationship between baseline scores (X), outcomes (Y) and

recovery (Y-X), to highlight the shortcomings of the proportional recovery rule, and illustrate those

problems with simulations in which synthetic recovery data are derived from different types of

recovery processes.

Findings

When the correlation between baseline scores and recovery is stronger than that between baselines

scores and outcomes, the former can create an inflated impression of how predictable outcomes

really are given baseline scores. This often happens when outcomes are less variable than baseline

scores, as is common in empirical studies of recovery after stroke. Moreover, we cannot use the

results of these correlations to distinguish proportional recovery from recovery which is either not

consistently proportional, or not proportional at all.

Interpretation

Analyses relating baseline scores to subsequent change are a minefield: our formal analysis applies

as consistently outside the area of stroke as it does within it. One implication of our analysis is that

the proportional recovery rule is not as predictive of real recovery after stroke as recent empirical

studies suggest. Another is that different analytical methods will be required to ascertain whether

recovery is even proportional at all.

Funding

Wellcome, the Medical Research Council and the Stroke Association.

Research in context

Evidence before this study

Reports supporting the proportional recovery rule are rapidly accumulating. Five separate studies since 2015 have applied the rule to recovery from hemiparesis after stroke, confirming prior observations that most patients recover a fixed proportion of lost function. This rule appears to predict recovery extremely accurately, regardless of the patients' ages at stroke onset, stroke type, sex, or therapy dose. And there is accumulating evidence that the rule predicts recovery from other post-stroke impairments, including impairments of lower limb function, aphasia and visuospatial neglect. If the proportional recovery rule is as powerful as these studies imply, recovery from stroke is a surprisingly simple and predictable process.

Added value of this study

Our results suggest that empirical analyses relating baseline scores to subsequent change can and often do engender a misleadingly optimistic impression of how predictable outcomes really are, from baseline scores. In other words, we show that the proportional recovery rule is not as predictive as it seems – and indeed that the empirical results to date do not demonstrate that recovery is proportional.

Implications of all available evidence

The emergence – and apparently extraordinary predictive power – of the proportional recovery rule has encouraged a shift in emphasis from outcomes to recovery, as the focus of research and of clinical practice in acute stroke. Our analysis suggests that this may be a mistake. Our results further suggest that the empirical analyses in this area do not reliably distinguish proportional recovery from other types of recovery, raising the question of how future analyses might make that distinction more confidently.

1. INTRODUCTION

Stroke clinicians and researchers have long known that the severity of patients' initial symptoms after stroke is related to their long term outcomes[1]. Recent studies have suggested that this relationship is stronger than previously thought: that most patients recover a fixed proportion of lost function during the first few months after stroke. Evidence in support of this 'proportional recovery rule' is rapidly accumulating, as recently summarised by Stinear[2]. In five separate studies since 2015[3-7], researchers used the Fugl-Meyer test to assess stroke patients' upper limb motor function within two weeks after stroke onset, and then again either three or six months post-stroke. The results were consistent with earlier observations[8,9] that most patients recovered around 70% of lost function. Taken together, these studies report highly consistent recovery in over 500 patients, spread across different countries with different approaches to rehabilitation, regardless of the patients' ages at stroke onset, stroke type, sex, or therapy dose[2]. And there is increasing evidence that the rule also captures recovery from post-stroke impairments of lower limb function[10], attention[11,12], and language[12,13], and may even apply generally across cognitive domains[14]. Even rats appear to recover proportionally after stroke[15].

These results are striking because of the extraordinary accuracy with which the proportional recovery rule appears to predict patients' recoveries. Most of these studies report that the rule explains more (and sometimes much more) than 80% of the variance in empirical recovery: when predicting behavioural performance in humans, these effect sizes are unprecedented. In 2015, Winters and colleagues[3] reported that recovery predicted using the rule could explain a staggering 95% of the variance in the recovery of 146 stroke patients. Like many of its counterparts[7,10,12], this study also reported a group of (65) 'non-fitters', who did not make the predicted recovery. But if non-fitters can be distinguished at the acute stage, as the authors' results also suggest, the implication is that we can predict most patients' recovery near-perfectly, given baseline score alone. Stroke researchers are used to thinking of recovery as a complex, multi-factorial process[16,17]. If the proportional recovery rule is really as powerful as it seems, much of what we thought we knew about recovery from stroke will need to be revisited[2,11,18].

In what follows, we argue that the proportional recovery rule is not as powerful as it seems, because current analyses in this domain are all potentially confounded by mathematical coupling.

2. MATHEMATICAL COUPLING

Mathematical coupling[19,20] refers to situations in which the same variable is employed to calculate

both the independent and the dependent variables in an analysis. In analyses relating baselines to

change, change is defined as 'outcome scores minus baseline scores': baseline scores are both the

independent variable, and a component of the dependent variable. For the sake of brevity in what

follows, we define 'baselines' = X, 'outcomes' = Y, and 'change' (or recovery) = Δ: i.e. delta, or

outcomes minus baselines. The 'correlation between baselines and outcomes' is $r(X, Y)$, and the

'correlation between baselines and change' is $r(X, Δ)$. Finally, we define the 'variability ratio' as the

ratio of the standard deviation (σ) of Y to the standard deviation of X ($σ_Y/σ_X$). X and Y are construed

as lists of scores, with each entry being the performance of a single patient at the specified time

point. We assume that performance improves as the numbers increase, so $r(X, Δ)$ will be negative if

recovery is proportional to lost function (because those who lose the most have smaller X and larger

Δ). One can equally formulate the relationship with 'lost function' (maximum X minus actual X)

instead of 'baseline score', but while this makes $r(X, Δ)$ positive if recovery is proportional to lost

function, it is otherwise entirely equivalent.

It has long been known that, when X and Y are independent, random variables drawn from

the standard normal distribution, $r(X, Δ) ≈ -0.71$[21]. In this canonical example of mathematical

coupling, there is a strong correlation between baselines and change, even though baselines and

outcomes are entirely unrelated: $r(X, Y) ≈ 0$.

Another expression of this example, which is more germane to our current concerns, is that

in this case, $r(X, Δ)$ makes Y appear more predictable than it really is. A strong correlation between

two variables typically implies that we can use each to predict the other: in this case, that we can use

X to predict Δ. In the limit as our sample grows, out-of-sample predicted Δ (pΔ) should tend toward

the least-squares line defined by $r(X, Δ)$, so the correlation between pΔ and empirical Δ should have

the same magnitude as $r(X, Δ)$ (see Supplementary Appendix A, proposition 8, bullet point 1).

Correlations between predicted and empirical data are a popular measure of predictive accuracy:

the stronger the correlation, the better the predictions. Since change is precisely the difference

between baselines and outcomes, a result suggesting we can predict change accurately naturally

encourages the expectation that we can predict outcomes *equally* accurately, by adding 'known

baselines' to 'accurately predicted change' to produce 'accurately predicted outcomes'. The

expectation is that the correlation between these predicted outcomes and empirical outcomes will

be as strong as the correlation between predicted and empirical change, or more formally: $|r(X + pΔ,$

$Y)| ≈ |r(X, Δ)|$.

However, as might be obvious from our canonical example of mathematical coupling, this

expectation is a fallacy: in fact, $|r(X + p\Delta, Y)| = |r(X, Y)|$. No matter how accurately we can seemingly

predict recovery, the correlation between outcomes calculated using those predictions, and

empirical outcomes, has the same strength as the correlation between baselines and outcomes (see

Supplementary Appendix A, proposition 8, bullet 2). In our canonical example, the impression that

we can predict change relatively well (because $r(p\Delta, \Delta) \approx 0.71$), is spurious because we cannot use

those predictions to predict outcomes even nearly as well (because $r(X, Y) \approx 0$).

3. MATHEMATICAL COUPLING AND MEASUREMENT NOISE

Krakauer and Marshall recently argued that our canonical example of mathematical coupling is not a

proper null hypothesis for empirical analyses of recovery after stroke[18]. This is because the canonical

example employs independent baselines and outcomes, whereas empirical baselines and outcomes

are dependent; repeated measures from the same patients. Using True Score theory[22], these

authors advance a more plausible alternative, in which 'true' baselines become 'true' outcomes via

some 'true' recovery trajectory, and observed baselines and outcomes are just the true states plus

measurement noise. From this perspective, the null hypothesis is that baselines are linked to

outcomes via random recovery trajectories, and under this hypothesis, it is argued that $r(X, \Delta) \approx 0$

regardless of the magnitude of recovery: i.e. the spurious effects described previously do not

emerge. Only with extreme measurement noise, so the argument goes, will dependent measures

behave like independent measures, with the consequent emergence of spurious $r(X, \Delta)$. Since the

most popular measurement tool in empirical studies of proportional recovery, the Fugl-Meyer scale,

is reliable[23], it follows that those empirical results are probably not spurious[18].

One weakness in this argument is that it needs to be made again for every new assessment

tool employed to measure proportional recovery. For example, neither the subset of the Western

Aphasia Battery[24] used to measure proportional recovery from post-stroke language impairments[13],

nor the letter cancellation task[25] used to measure proportional recovery from visuospatial neglect[11],

are as reliable as the Fugl-Meyer scale. But whether or not those arguments can be made, we

contend that there are many more – and more likely – situations in which $r(X, \Delta)$ may be

misleadingly strong, regardless of measurement noise.

4. MATHEMATICAL COUPLING REGARDLESS OF MEASUREMENT NOISE

171   For any baselines (X) and outcomes (Y), it can be shown that:

$$r(X, \Delta) = \frac{\sigma_Y \cdot r(X,Y) - \sigma_X}{\sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \cdot \sigma_X \cdot \sigma_Y \cdot r(X,Y)}} \quad \text{(Equation 1)}$$

175   A formal proof of Equation 1 is provided in Supplementary Appendix A (see, proposition 4

176   and theorem 1); its core consequence is that r(X, Δ) is a (somewhat complex) function of r(X, Y) and

177   σ_Y/σ_X. To illustrate that function, we ran a series of simulations in which r(X, Y) and σ_Y/σ_X were varied

178   independently: a Matlab script for this is in Supplementary Appendix B, and the results are

179   illustrated in Figure 1.

181   --Insert Figure 1 and Table 1--

Our canonical example of mathematical coupling (i.e., when r(X, Y) = 0, σ_Y/σ_X = 1 and r(X, Δ) ≈

-0.71) is consistent with equation 1, and corresponds to the red sphere marked in figure 1. One way

to understand this surface more generally, is to consider what different variability ratios do to the

correlation between X and Y-X (i.e. Δ): see Table 1. When Y is more variable than X, Y contributes

more variance to Y-X and r(X, Y-X) tends toward r(X, Y): this is Regime 2. In contrast, when Y is less

variable than X, X contributes more variance to Y-X, and r(X, Y-X) tends toward r(X, -X), the

autocorrelation between X and itself: this is Regime 1 (also see proposition 9 in Supplementary

Appendix A). As an illustration of Regime 1, consider a set of synthetic baselines and outcomes

connected by proportional recovery. With no measurement noise, r(X, Y) = 1 and r(X, Δ) = -1 (Figure

2a); but with shuffled outcomes data, r(X, Y) ≈ 0, but r(X, Δ) = -0.96 (Figure 2b). Indeed, r(X, Δ) will be

extreme in any data where σ_Y is significantly smaller than σ_x.

195   --Insert Figure 2--

5.   RE-EXAMINING STUDIES OF PROPORTIONAL RECOVERY AFTER STROKE

These relationships between r(X, Y), r(X, Δ) and σ_Y/σ_X merit a re-examination of the proportional

recovery rule for stroke. The only study in this literature that reports individual patients' behavioural

data is that by Zarahn and colleagues[9], who reported recovery from hemiparesis in 30 stroke

patients. When measured across the whole sample, r(X, Y) = 0.80 and r(X, Δ) = -0.49; this changes to

r(X, Y) = 0.75 and r(X, Δ) = -0.95, after removing the 7 patients (23%) who did not fit the rule.

Calculating predicted recovery (pΔ) from the least-squares line for r(X, Δ), we get r(pΔ, Δ) = 0.95, but

r(X + pΔ, Y) = 0.75. By removing those 7 non-fitters, Zarahn and colleagues dramatically increased

the apparent predictability of recovery, while actually reducing the apparent predictability of

outcomes – and reducing the variability ratio, from 0.88 in the full sample to 0.36 in the reduced

sample of fitters. Notably, the residuals or errors for both correlations (r(X, Y) and r(X, Δ)) are

identical (see Figure 3), and in fact this is always true (see Supplementary Appendix A, proposition

10). Since r(X, Y) and r(X, Δ) assign different effect sizes to the same errors, r(X, Δ) is misleadingly

strong.

212   --Insert Figure 3--

We can also use equation 1 to reinterpret studies which do not report individual patient

data. One example is the first study to apply the proportional recovery rule to aphasia after stroke[13].

Here, baseline scores explained 81% of the variance in recovery (r(X, Δ) ≈ -0.9), and the variability

ratio was 3.7 (outcomes) / 7.7 (baselines) = 0.48. By equation 1, r(X, Y) was at most 0.78 here, but

could also have been zero; without a report of r(X, Y), we cannot know how predictable Y really was,

given X. Similar concerns emerge from the recent measurement of proportional recovery in 182 rats

(from a total sample of 593)[15]. Here, the variability ratio was 3.2 (outcomes) / 4 (baseline) = 0.8,

baseline scores explained 49.9% of the variance in recovery (r(X, Δ) ≈ -0.71). By equation 1, this

implies that r(X, Y) was either much stronger (>0.95) or considerably weaker (~0.29) than their r(X,

Δ). Here again, we have learned less about the predictability of outcomes than the headline effect

size suggests.

Some studies report inter-quartile ranges (IQR) instead of standard deviations for their data;

accepting some room for error, we can still estimate variability ratios from those IQRs. The study by

Winters and colleagues[3] is an example, which reports proportional recovery in 146 patients (from a

total sample of 211). Here, r(X, Δ) = -0.97 and σ_Y/σ_X = 6 (follow-up) / 38 (baseline) = 0.158. By

equation 1, this small ratio means that r(X, Δ) will be at least as extreme as that reported regardless

of the r(X, Y). And the same issues arise in the recent study of proportional recovery from lower-limb

impairments, by Veerbeck and colleagues[26]. Here, σ_Y/σ_X = 0.438 and r(X, Δ) ≈ -0.88; equation 1

implies that r(X, Δ) could be at least that strong regardless of r(X, Y). The headline effect sizes in

these studies do not tell us how predictable outcomes really were, given baseline scores.

Finally, while r(X, Δ) can be misleading if it is extreme relative to r(X, Y), the reverse is also

true. For example, the only study in this literature to employ outcomes as the dependent variable,

rather than recovery, reports that r(X, Y) ≈ 0.8 and σ_Y/σ_X = 23.4 (follow-up) /19.5 (baseline) = 1.2 in

their 'combined' group of 76 patients. By equation 1, r(X, Δ) = -0.05 in this case. These authors only

claimed that a subset of their sample recovered proportionally, but their full sample seems quite

consistent with the null hypothesis, of random recovery, implied by a True Score analysis of recovery

data[18].

Notably, most of the studies referenced in this section reported outcomes scores which

were less variable than baseline scores. This is the predicted pattern if recovery is proportional to

lost function, because in this case those who lose the most will also recover the most, but this

scenario also encourages extreme r(X, Δ) regardless of r(X, Y) (i.e. Regime 1 from Figure 1). Even if

patients really do recover proportionally, r(X, Δ) can encourage an inflated impression of how well

the rule predicts their recovery. We found potentially misleading results in all of the studies that

reported enough information for us to make the judgement. The most common reason why we

could not make a judgement was when baselines were only related to recovery through

multivariable models[4,11,12]. But inflation in one variable's effect size will also inflate the multivariable

model's effect size: we cannot estimate the magnitude of that inflation confidently for any particular

multivariable model, but we can predict that it's also there in those models.

As a consequence, we suggest that future studies of proportional recovery should at least

report both r(X, Y) and r(X, Δ). However, even when both are reported, we contend that analyses like

this cannot convincingly demonstrate that recovery is proportional: this is the focus of the next

section.

6. **HOW DO PATIENTS REALLY RECOVER?**

To illustrate why r(X, Y) and r(X, Δ) cannot convincingly identify proportional recovery, we ran a

series of simulations in which these correlations were calculated from synthetic data, generated by

different types of recovery process. All of the simulations assume that baselines and outcomes are

dependent, and that each is observed with some measurement noise, and all also assume our

samples are large (N = 1,000); the distortions induced by small samples are well documented

elsewhere[27], and will not concern us here.

Our simulations vary: (a) the magnitude of measurement noise; (b) the real variation in

participants' recovery trajectories (i.e. 'recovery noise'); (c) the presence or absence of boundaries

on the measurement scale (i.e. ceiling and floor effects); and (d) whether or not 'non-fitters' (whose

'recovery' is less than expected) are removed from the analysis. Three different types of recovery

are considered: (i) random recovery; (ii) constant recovery (i.e. patients recover a fixed amount); and

(iii) proportional recovery. Results ($r(X, Y)$ and $r(X, \Delta)$) are mean values from 100 repetitions of the

same simulations: the Matlab script required to run them is in Supplementary Appendix C.

Figure 4 displays the results: $r(X, Y)$ and $r(X, \Delta)$ with increasing recovery noise, with each

configuration of recovery mechanism and other constraints. As expected given our discussion in

section 3, when measurements are noisy (Figure 4, row 1), mathematical coupling dominates and all

of the recovery regimes yield similar results. With more reliable measurements (Figure 3, row 2), the

results are more promising, in that one can reasonably hope to use these correlations to distinguish

proportional recovery from other types of recovery. However, these simulations make the

assumption that the scale used to measure performance is infinitely long. In practice, these scales

are always finite, and the Fugl-Meyer scale in particular is known to exert ceiling effects on patients

with mild-to-moderate hemiparesis[23]. When we include ceiling effects (Figure 4, row 3), the results

derived from constant recovery are similar to those derived from proportional recovery. These

similarities persist when we also remove non-fitters prior to calculating the correlations (Figure 4,

last row), which acts to preserve strong $r(X, \Delta)$ despite significant recovery noise. Indeed, in our final

simulation of proportional recovery (row 4, column 2), $r(X, \Delta) \approx -0.94$ even when the standard

deviation of the recovery noise is >0.35. Since the mean proportion is 70% here (0.7), the 95%

confidence intervals (mean ± two standard deviations) on the proportion of lost function actually

recovered include both 0% and 100%.

288   --Insert Figure 4--

These results demonstrate that, with more realistic constraints on our generative models of

empirical data, $r(X, Y)$ and $r(X, \Delta)$ cannot distinguish proportional recovery from recovery which is

either not consistently proportional, or is not proportional at all.

7. Discussion

The proportional recovery rule is attractive partly because it formalises the popular, clinical intuition

that initial symptom severity is by far the most, and perhaps the only, important predictor of recovery

after stroke. The rule is also striking because it implies that recovery is simple and consistent across

patients (non-fitters notwithstanding), and because that implication appears to be justified by strong

empirical results[2]. That incredible, empirical power has even begun to encourage the search for some

deeper, general recovery mechanism which cuts across cognitive domains[18]. But our results suggest

that the rule may not be as powerful as it seems.

We are not claiming that the rule is wrong. Our analysis suggests that the empirical results to

date do not confirm that it holds or how well, but those results are still consistent with the rule's

predictions. The rule is also falsifiable: a recent study of recovery during the first week after stroke

onset found no evidence of proportional recovery[28]. Nevertheless, we contend that estimates of the

rule's predictive power – those headline effect sizes for $r(X, \Delta)$ or $r(p\Delta, \Delta)$ – deliver a misleadingly

optimistic impression of how predictable outcomes really are. And a misleadingly optimistic

impression here encourages misleadingly pessimistic assessments elsewhere: if baseline scores really

do predict many patients' recoveries near-perfectly, therapeutic interventions will appear to be

redundant for those patients.

Much of the work in this area has focused on the distinction between fitters and non-fitters

to the proportional recovery rule; distinguishing the groups via analyses of the residuals for $r(X, \Delta)$.

This is a circular definition, but it defines groups which can then be distinguished in non-circular ways[5].

Since the residuals for $r(X, Y)$ and $r(X, \Delta)$ are always the same (see Supplementary Appendix A), it

follows that our concerns about effect sizes have no bearing on the validity of work to distinguish

fitters from non-fitters. Nevertheless, extreme $r(X, \Delta)$ for fitters will naturally encourage the

assumption that we can at least predict those fitters' outcomes near-perfectly. Our analysis suggests

that this assumption is wrong.

One open question concerns the implications of our formal analysis beyond the domain of

stroke. Analyses relating baseline scores to subsequent change have been employed in many different

fields[21]; as far as we know, none has employed a formal account like that proposed here. Prior work

in this area has tended to focus on distinguishing significant relationships, rather than measuring their

strength (as we have done here) [20,21,29]; but effect sizes define p-values for a given sample size, so

there is every reason to hope that our equation 1 could be used to address those hypothesis-testing

questions as well.

In summary, our results suggest that recovery from stroke might not be as proportional as it

seems. Strong correlations between baseline scores (or predicted change) and subsequent change can

drive a misleadingly optimistic impression of how predictable outcomes really are. This situation appears to be common, both in our simulations and in the empirical literature on the proportional recovery rule. Moreover, these analyses cannot distinguish proportional recovery from recovery which is either not consistently proportional, or not proportional at all. Quite how to make this distinction with confidence, remains an open question: this is a subject for future work.

8. A PARELLEL CHALLENGE?

As this manuscript approached completion, we discovered that another researcher, Robinson Kundert, had distributed a 'pre-paper', describing a parallel challenge to the proportional recovery rule, at a meeting. None of the authors attended that meeting, but we learned of it through colleagues who did attend. Our understanding is that this challenge is focused on the consistency (or lack thereof) of the proportion of lost function that patients recover. In this sense, that work may be complementary to our focus on predictive power. We look forward to studying this challenge after it is published.

ACKNOWLEDGMENTS

This study was supported for the Medical Research Council (MR/M023672/1, MR/K022563/1), the Wellcome Trust (091593/Z/10/Z), and the Stroke Association (TSA PDF 2017/02). The funders had no participation in the design and results of this study.

REFERENCES

1. Jongbloed L. Prediction of function after stroke: a critical review. *Stroke; a journal of cerebral circulation* 1986; **17**(4): 765-76.
2. Stinear CM. Prediction of motor recovery after stroke: advances in biomarkers. *The Lancet Neurology*; **16**(10): 826-36.
3. Winters C, Wegen EEH, Daffertshofer A, Kwakkel G. Generalizability of the Proportional Recovery Model for the Upper Extremity After an Ischemic Stroke. *Neurorehabilitation and neural repair* 2015; **29**.
4. Buch ER, Rizk S, Nicolo P, Cohen LG, Schnider A, Guggisberg AG. Predicting motor improvement after stroke with clinical assessment and diffusion tensor imaging. *Neurology* 2016; **86**(20): 1924-5.
5. Feng W, Wang J, Chhatbar PY, et al. Corticospinal tract lesion load: An imaging biomarker for stroke motor outcomes. *Annals of neurology* 2015; **78**(6): 860-70.

6.   Stinear CM, Byblow WD, Ackerley SJ, Smith M-C, Borges VM, Barber PA. Proportional Motor
Recovery After Stroke. *Implications for Trial Design* 2017; **48**(3): 795-8.
7.   Byblow WD, Stinear CM, Barber PA, Petoe MA, Ackerley SJ. Proportional recovery after stroke
depends on corticomotor integrity. *Annals of neurology* 2015; **78**(6): 848-59.
8.   Prabhakaran S, Zarahn E, Riley C, et al. Inter-individual variability in the capacity for motor
recovery after ischemic stroke. *Neurorehabilitation and neural repair* 2007; **22**.
9.   Zarahn E, Alon L, Ryan SL, et al. Prediction of motor recovery using initial impairment and fMRI
48 h poststroke. *Cereb Cortex (New York, NY 1991)* 2011; **21**.
10.   Smith M-C, Byblow WD, Barber PA, Stinear CM. Proportional Recovery From Lower Limb
Motor Impairment After Stroke. *Stroke; a journal of cerebral circulation* 2017.
11.   Winters C, Wegen EEHv, Daffertshofer A, Kwakkel G. Generalizability of the Maximum
Proportional Recovery Rule to Visuospatial Neglect Early Poststroke. *Neurorehabilitation and neural*
repair 2017; **31**(4): 334-42.
12.   Marchi NA, Ptak R, Di Pietro M, Schnider A, Guggisberg AG. Principles of proportional recovery
after stroke generalize to neglect and aphasia. *European Journal of Neurology* 2017; **24**(8): 1084-7.
13.   Lazar RM, Minzer B, Antoniello D, Festa JR, Krakauer JW, Marshall RS. Improvement in aphasia
scores after stroke is well predicted by initial severity. *Stroke; a journal of cerebral circulation* 2010;
41(7): 1485-8.
14.   Ramsey LE, Siegel JS, Lang CE, Strube M, Shulman GL, Corbetta M. Behavioural clusters and
predictors of performance during recovery from stroke. *Nature human behaviour* 2017; **1**: 0038.
15.   Jeffers MS, Karthikeyan S, Corbett D. Does Stroke Rehabilitation Really Matter? Part A:
Proportional Stroke Recovery in the Rat. *Neurorehabilitation and neural repair* 2018; **32**(1): 3-6.
16.   Plowman E, Hentz B, Ellis C, Jr. Post-stroke aphasia prognosis: a review of patient-related and
stroke-related factors. *Journal of evaluation in clinical practice* 2012; **18**(3): 689-94.
17.   Jorgensen HS, Nakayama H, Raaschou HO, Vive-Larsen J, Stoier M, Olsen TS. Outcome and
time course of recovery in stroke. Part I: Outcome. The Copenhagen Stroke Study. *Archives of physical*
medicine and rehabilitation 1995; **76**(5): 399-405.
18.   Krakauer JW, Marshall RS. The proportional recovery rule for stroke revisited. *Annals of*
neurology 2015; **78**(6): 845-7.
19.   Oldham PD. A note on the analysis of repeated measurements of the same subjects. *Journal*
of Chronic Diseases 1962; **15**(10): 969-77.
20.   Blomqvist N. On the relation between change and initial value. *Journal of the American*
Statistical Association 1977; **72**(360a): 746-9.
21.   Tu YK, Gilthorpe MS. Revisiting the relation between change and initial value: a review and
evaluation. *Statistics in medicine* 2007; **26**(2): 443-57.
22.   Traub RE. Classical test theory in historical perspective. *Educational Measurement: Issues and*
Practice 1997; **16**(4): 8-14.
23.   Gladstone DJ, Danells CJ, Black SE. The Fugl-Meyer Assessment of Motor Recovery after
Stroke: A Critical Review of Its Measurement Properties. *Neurorehabilitation and neural repair* 2002;
16(3): 232-40.
24.   Shewan CM, Kertesz A. Reliability and validity characteristics of the Western Aphasia Battery
(WAB). *The Journal of speech and hearing disorders* 1980; **45**(3): 308-24.
25.   Pradhan B, Nagendra HR. Normative data for the letter-cancellation task in school children.
International Journal of Yoga 2008; **1**(2): 72-5.
26.   Veerbeek JM, Winters C, van Wegen EEH, Kwakkel G. Is the proportional recovery rule
applicable to the lower limb after a first-ever ischemic stroke? *PLOS ONE* 2018; **13**(1): e0189279.
27.   Button KS, Ioannidis JPA, Mokrysz C, et al. Power failure: why small sample size undermines
the reliability of neuroscience. *Nat Rev Neurosci* 2013; **14**(5): 365-76.
28.   Dunn LE, Schweber AB, Manson DK, et al. Variability in Motor and Language Recovery during
the Acute Stroke Period. *Cerebrovascular diseases extra* 2016; **6**(1): 12-21.

29.   Hayes RJ. Methods for assessing whether change depends on initial value. *Statistics in*
medicine 1988; **7**(9): 915-27.

Figure 1

Figure 1: The relationship between r(X, Y), r(X, Δ) and the variability ratio of Y to X. Note that the x-axis is log-transformed to ensure symmetry around 1 (i.e. when X and Y are equally variable, and $\log(\sigma_Y/\sigma_X) = 0$). The two major regimes of Equation 1 are also marked in red. In Regime 1, the imprint of baselines (X) dominates, and r(X, Δ) tends toward r(X, -X) (i.e. -1). In Regime 2, the correlations of interest are more tightly related, and r(X, Δ) ≈ r(X, Y). And there is a transition between the two regimes when the variability ratio is not dramatically skewed one way or the other – though even this transition allows for misleadingly strong r(X, Δ), as evidenced by the fact that our canonical example of mathematical coupling occurs in this region. This is indicated by the red mark, which corresponds to the situation where X and Y are independent random normal variables: i.e. r(X, Y) ≈ 0; log variability ratio = 0; and r(X, Δ) ≈ -0.71. Proposition 7 in Supplementary Appendix A provides a justification for unambiguously using a ratio of standard deviations in this figure, rather than σ_Y and σ_X as separate axes.

Figure 2: Illustrating how |r(X, Δ)| > |r(X, Y)| when X is more variable than Y. In both panels, X is a vector of 20 numbers drawn from a random uniform distribution in the range 1-65. In the left panel, Y is calculated using the proportional recovery rule: i.e. Y = X + ((66-X) x 0.7). In this case, the correlation between X and Y is 1 and the correlation between X and change is -1. In the right panel, Y is randomly shuffled: X and Y are now uncorrelated, but the correlation between X and change remains extremely strong (r = -0.96).

442   Figure 3

444   **Figure 3:** (Left) Least squares linear fits for analyses relating baselines to (upper) outcomes and

445   (lower) recovery, using the fitters' data reported by Zarahn and colleagues[9]. (Middle) Plots of

446   residuals relative to each least squares line, against the fitted values in each case. (Right) A scatter

447   plot of the residuals from the model relating baselines to change, against the residuals from the

448   model relating baselines to outcomes: the two sets of residuals are the same.

Figure 4

Figure 4: Simulation results. Plotting r(X, Y) and r(X, Δ) as a function of increasing true variability in recovery trajectories, where recovery is: (a) random; (b) proportional; or (c) constant. The top row displays results when the variance of measurement noise is large relative to the variance of baselines. The second row presents results when measurement noise is minimal. The third row presents results when measurement noise is minimal but there are ceiling effects on the measurement scale. And the last row presents results when measurement noise is minimal, ceiling effects are applied, and 'non-fitters' are removed: i.e. the ~19% (this is the figure in [3]) of the sample whose actual recovery deviates most strongly and negatively from the least squares line defined by r(X, Δ).

465 Table 1

REGIME	VARIABILITY OF Y (σ_Y)	VARIABILITY OF X (σ_X)	Δ [= Y-X]	r(X, Δ) [=r(X, Y-X)]
1	Smaller	Larger	Y-X ≈ -X	r(X, Y-X) ≈ r(X, -X) = -1
2	Larger	Smaller	Y-X ≈ Y	r(X, Y-X) ≈ r(X, Y)

Supplementary Appendix A: formal relationships between the correlations

We present a very simple and general formulation of the proportional recovery concept. We assume two variables X' and Y' corresponding to performance at initial test (X') and at second test (Y'). These will be represented as column vectors, with each entry being the performance of a single patient and vector lengths being $N \in \mathbb{N}$. Performance improves as numbers get bigger, up to a maximum, denoted Max, which corresponds to no discernible deficit. Severity is measured as difference from maximum, i.e. $Max - X'$.

The two variables (X' and Y') could be specialised to more detailed formulations: e.g., true score theory or with an explicit modelling of measurement or state error. However, this would not impact any of the derivations or inferences that follow. Indeed, the results that we present would hold even in the complete absence of measurement noise, which has been considered the main concern for the validity of quantifications of proportional recovery.

Demeaning

Without loss of generality, we work with demeaned variables. That is, where over-lining denotes mean, we define new variables as,

$$X = X' - \bar{X}'$$

$$Y = Y' - \bar{Y}'$$

This also means that recovery, i.e. $Y - X$, will be demeaned, since, using proposition 1, the following holds.

$$Y - X = (Y' - \bar{Y}') - (X' - \bar{X}') = (Y' - X') - (\bar{Y}' - \bar{X}') = (Y' - X') - \overline{(Y' - X')}$$

Proposition 1

Let V and W be vectors of the same length, denoted N. Then, the following holds,

$$\bar{V} + \bar{W} = \overline{(V + W)}$$

with $\bar{V} - \bar{W} = \overline{(V - W)}$ as a trivial consequence.

Proof

By distributivity of multiplication through addition and associativity of addition, the following holds.

$$\bar{V} + \bar{W} = \left(\frac{1}{N}\sum_{i=1}^{N}V_i\right) + \left(\frac{1}{N}\sum_{i=1}^{N}W_i\right) = \frac{1}{N}\left(\sum_{i=1}^{N}V_i + \sum_{i=1}^{N}W_i\right) = \frac{1}{N}\left(\sum_{i=1}^{N}(V_i + W_i)\right) = \overline{(V + W)}$$

QED

Correlations

There are two basic correlations we are interested in, (1) the correlation between initial performance and performance at second test, i.e. $r(X, Y)$, and (2) the correlation between initial performance and recovery, i.e. $r(X, Y - X) = r(X, \Delta)$. The latter of these is the key relationship, and we would expect this to be a negative correlation; that is, as initial performance is smaller (i.e. further from Max), the larger is recovery. (One could also formulate the correlation as $r((Max - X), Y - X)$, which would flip the correlation to positive, but the two approaches are equivalent).

503   Our main correlations are defined as follows,

504
$$r(X,Y) = \frac{\sum_{i=1}^{N} X_i \cdot Y_i}{(N-1) \cdot \sigma_X \cdot \sigma_Y}$$

505
$$r(X,(Y-X)) = \frac{\sum_{i=1}^{N} (X_i \cdot (Y_i - X_i))}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}}$$

506   **Standard Deviation of a Difference**

507   We need a straightforward result on the standard deviation of a difference.

508   **Proposition 2**

509
$$\sigma_{(A-B)} = \sqrt{\sigma_A^2 + \sigma_B^2 - 2 \cdot cov(A,B)}$$

510   **Proof**

511   The result is a direct consequence of the following standard result from probability theory, e.g. see
512   Ross, S. M. (2014). *Introduction to probability and statistics for engineers and scientists*. Academic
513   Press.,

514
$$\sigma_{(A-B)}^2 = \sigma_A^2 + \sigma_B^2 - 2 \cdot cov(A,B)$$

516   **Key Results**

517   The following proposition enables us to express the key correlation, $r(X,(Y-X))$, in terms of
518   covariance of its constituent variables.

519   **Proposition 3**

520
$$r(X,(Y-X)) = \frac{cov(X,Y) - cov(X,X)}{\sigma_X \cdot \sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \cdot cov(X,Y)}}$$

521   **Proof**

522   Using distributivity of multiplication through addition, associativity of addition, the definition of
523   covariance and proposition 2, we can reason as follows.

524
$$r(X,(Y-X)) = \frac{\sum_{i=1}^{N} (X_i \cdot (Y_i - X_i))}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}} = \frac{\sum_{i=1}^{N} (X_i\, Y_i - X_i\, X_i)}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}} = \frac{\sum_{i=1}^{N} (X_i\, Y_i) - \sum_{i=1}^{N} (X_i\, X_i)}{(N-1) \cdot \sigma_X \cdot \sigma_{(Y-X)}}$$

525
$$= \frac{cov(X,Y) - cov(X,X)}{\sigma_X \cdot \sigma_{(Y-X)}} = \frac{cov(X,Y) - cov(X,X)}{\sigma_X \cdot \sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \cdot cov(X,Y)}}$$

526   QED

527   It is straightforward to adapt proposition 3 to be fully in terms of correlations.

528   **Proposition 4**

529
$$r(X,(Y-X)) = \frac{\sigma_Y \cdot r(X,Y) - \sigma_X \cdot r(X,X)}{\sqrt{\sigma_Y^2 + \sigma_X^2 - 2 \cdot \sigma_X \cdot \sigma_Y \cdot r(X,Y)}}$$

Proof

Straightforward from proposition 3 and definition of correlations, which gives the relationship
$cov(A, B) = \sigma_A . \sigma_B . r(A, B)$. QED

Scale Invariance

The next set of propositions justifies working with a standardised X variable.

Lemma 1

536
$$\forall c \in \mathbb{R} \cdot |c| . \sigma_A = \sigma_{(c.A)}$$

537 **Proof**

538 Using distributivity of a multiplicative constant through averaging, $\sqrt{d^2} = |d|$ and distributivity of
539 square root through multiplication, we can reason as follows.

540
$$\sigma_{(c.A)} = \sqrt{\frac{\sum_{i=1}^{N} (c.A_i - \overline{c.A})^2}{N-1}} = \sqrt{\frac{\sum_{i=1}^{N} (c.A_i - c.\bar{A})^2}{N-1}} = |c| . \sqrt{\frac{\sum_{i=1}^{N} (A_i - \bar{A})^2}{N-1}} = |c| . \sigma_A$$

541 QED

542 **Proposition 5 (Invariance to scaling)**

543 The absolute magnitude of a correlation is not changed by scaling either variable by a constant, i.e.

544
$$\forall c \in \mathbb{R} \cdot r(A, B) = sign(c) . r(c.A, B) = sign(c) . r(A, c.B)$$

545 where $sign(d) = if\ (d < 0)\ then - 1\ else + 1$.

546 **Proof**

547 For any $c \in \mathbb{R}$, using distributivity of multiplication through mean and addition, and lemma 1, the
548 following holds,

549
$$r(c.A, B) = \frac{\sum_{i=1}^{N} (c.A_i - \overline{c.A})(B_i - \bar{B})}{(N-1) . \sigma_{(c.A)} \sigma_B} = \frac{\sum_{i=1}^{N} (c.A_i - c.\bar{A})(B_i - \bar{B})}{(N-1) . \sigma_{(c.A)} \sigma_B}$$

550
$$= \frac{c . \sum_{i=1}^{N} (A_i - \bar{A})(B_i - \bar{B})}{(N-1) . |c| . \sigma_A . \sigma_B} = \frac{sign(c) . \sum_{i=1}^{N} (A_i - \bar{A})(B_i - \bar{B})}{(N-1) . \sigma_A . \sigma_B} = sign(c) . r(A, B)$$

551 Then, one can multiply both sides by $sign(c)$ to obtain $r(A, B) = sign(c) . r(c.A, B)$. Additionally,
552 as correlations are symmetric, $sign(c) . r(c.B, A) = sign(c) . r(A, c.B)$, and the full result follows.

553 QED

554 **Corollary 1**

555
$$\forall c \in \mathbb{R} \cdot r(A, B) = r(c.A, c.B)$$

556 **Proof**

557 Follows from twice applying proposition 5, and that $sign(c)^2 = +1$. QED

558 **Proposition 6**

559
$$\forall c \in \mathbb{R} \cdot r(X, (Y - X)) = r(c.X, (c.Y - c.X))$$

560 **Proof**

561 We can use distributivity of multiplication through subtraction and corollary 1 to give us the
562 following.

$$r\big(c.X,(c.Y - c.X)\big) = r\big(c.X, c.(Y - X)\big) = r(X,(Y - X))$$

564 QED

565 It follows from proposition 6 that we can work with a standardised X variable, since,

$$r(X/\sigma_X,(Y/\sigma_X - X/\sigma_X)) = r(X,(Y - X))$$

567 **Proposition 7 (Sufficiency of variability ratio)**

568 Assume two pairs of variables: X_1, Y_1 and X_2, Y_2, such that, $r(X_1, Y_1) = r(X_2, Y_2)$, then,

$$\frac{\sigma_{Y_1}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{X_2}} \implies r\big(X_1,(Y_1 - X_1)\big) = r\big(X_2,(Y_2 - X_2)\big)$$

570 **Proof**

571 The proof has two parts.

572 1) We consider the implications of equality of ratio of standard deviations. Firstly, we note that,

$$\frac{\sigma_{Y_1}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{X_2}} \iff \frac{\sigma_{X_2}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{Y_1}} \quad (eqn\ ratios)$$

574 Secondly, using eqn ratios, we can argue as follows,

$$\frac{\sigma_{Y_1}}{\sigma_{X_1}} = \frac{\sigma_{Y_2}}{\sigma_{X_2}} \iff \left(\sigma_{Y_2} = \frac{\sigma_{X_2}}{\sigma_{X_1}}\sigma_{Y_1} \wedge \sigma_{X_2} = \frac{\sigma_{Y_2}}{\sigma_{Y_1}}\sigma_{X_1}\right) \iff \left(\sigma_{Y_2} = \frac{\sigma_{X_2}}{\sigma_{X_1}}\sigma_{Y_1} \wedge \sigma_{X_2} = \frac{\sigma_{X_2}}{\sigma_{X_1}}\sigma_{X_1}\right)$$

$$\implies \left(\exists d \in \mathbb{R} \cdot \sigma_{Y_2} = d.\sigma_{Y_1} \wedge \sigma_{X_2} = d.\sigma_{X_1}\right)$$

577 2) Using 4, the fact that $r(X_1, Y_1) = r(X_2, Y_2)$, the property just derived in part 1), with $d = \frac{\sigma_{X_2}}{\sigma_{X_1}}$ and
578 rules of square roots, we can reason as follows,

$$r\big(X_2,(Y_2 - X_2)\big) = \frac{\sigma_{Y_2}.r(X_2,Y_2) - \sigma_{X_2}.r(X_2,X_2)}{\sqrt{\sigma_{Y_2}^2 + \sigma_{X_2}^2 - 2.\sigma_{X_2}.\sigma_{Y_2}.r(X_2,Y_2)}} = \frac{\sigma_{Y_2}.r(X_1,Y_1) - \sigma_{X_2}.r(X_1,X_1)}{\sqrt{\sigma_{Y_2}^2 + \sigma_{X_2}^2 - 2.\sigma_{X_2}.\sigma_{Y_2}.r(X_1,Y_1)}}$$

$$= \frac{d.\sigma_{Y_1}.r(X_1,Y_1) - d.\sigma_{X_1}.r(X_1,X_1)}{\sqrt{d^2.\sigma_{Y_1}^2 + d^2.\sigma_{X_1}^2 - 2.d.\sigma_{X_1}.d.\sigma_{Y_1}.r(X_1,Y_1)}} = \frac{d.(\sigma_{Y_1}.r(X_1,Y_1) - \sigma_{X_1}.r(X_1,X_1))}{d.\sqrt{\sigma_{Y_1}^2 + \sigma_{X_1}^2 - 2.\sigma_{X_1}.\sigma_{Y_1}.r(X_1,Y_1)}}$$

$$= r\big(X_1,(Y_1 - X_1)\big).$$

QED

Proposition 8

If $\Delta = Y - X$ and $p\Delta = X.\beta$, where $\beta \in \mathbb{R}$, then,

1) $r(p\Delta, \Delta) = sign(\beta).r(X,\Delta)$; and
2) $r(X + p\Delta, Y) = sign(1 + \beta).r(X,Y)$.

Proof

Both results are easy consequences of proposition 5.

1) $r(p\Delta, \Delta) = r(X.\beta, \Delta) = sign(\beta).r(X, \Delta).$

2) $r(X + p\Delta, Y) = r((X + (X.\beta)), Y) = r((X.(1 + \beta)), Y) = sign(1 + \beta).r(X, Y) = r(X, Y).$

QED

Main Findings

Theorem 1:

Since X will be standardised, we can adapt the finding in proposition 4, to give us the key
relationship we need,

596
$$r\big(X, (Y - X)\big) = \frac{\sigma_Y . r(X, Y) - r(X, X)}{\sqrt{\sigma_Y^2 + 1 - 2 . \sigma_Y . r(X, Y)}} \qquad (\text{eqn } Imprint)$$

Proof

Immediate from proposition 4.   QED

Theorem 1 shows clearly that $r(X, (Y - X))$ is fully defined by the correlations $r(X, Y)$ and $r(X, X)$,
along with the variability of Y. The correlation of X with itself, i.e. $r(X, X)$, is a prominent aspect of
this equation, which drives its oddities. $r(X, X)$ reflects the coupling in the equation that arises
because X appears in both the terms being correlated in $r(X, (Y - X))$. $r(X, X)$ is of course a
constant, i.e. 1 for any X, so in fact, σ_Y and $r(X, Y)$, are the only variables; accordingly, their size
determines the extent to which the imprint of X in $Y - X$ drives $r(X, (Y - X))$.

This leads to the key observation that, as σ_Y gets smaller, $r(X, (Y - X))$ tends towards $-r(X, X)$,
which equals -1. In other words, as the variability of Y decreases, the imprint of X becomes
increasingly prominent. This is shown in the next proposition.

Proposition 9

609
$$r\big(X, (Y - X)\big) \longrightarrow -r(X, X) = -1, \qquad \text{as } \sigma_Y \longrightarrow 0$$

Proof

The right hand side of equation *Imprint*, has five constituent terms, two in the numerator and three
in the denominator. Of these five, three are products with the standard deviation of Y, i.e. σ_Y.
Assuming all else is constant, as σ_Y reduces, the absolute value of each of these three terms reduces
towards zero. The rate of reduction is different amongst the three, but they will all decrease.
Accordingly, as σ_Y decreases, $r\big(X, (Y - X)\big)$ becomes increasingly determined by the two terms not
involving σ_Y, and thus, it tends towards $-\frac{r(X,X)}{\sqrt{+1}} = -r(X, X) = -1$.
QED

Equality of Residuals

An important finding of section 5 of the main text, is that the residuals resulting from regressing Y
onto X are the same as regressing Y-X onto X. We show in this section, that this equality of residuals
is necessarily the case.

We focus on the following two equations,

Eqn 1) $Y = \tilde{X}.\beta_1 + \varepsilon_1$

Eqn 2) $Y - X = \tilde{X}.\beta_2 + \varepsilon_2$

where $\tilde{X}$ is the $N \times 2$ matrix, with first column being X and second being the $N \times 1$ vector of ones
(which provides the intercept term); β_1 and β_2 are 2×1 vectors of parameters and Y, X, ε_1 and ε_2
are $N \times 1$ vectors. As in the rest of this document, Y and X are our (demeaned) initial and outcome
variables, while ε_1 and ε_2 are our residual error terms.

Proposition 10

631 If we assume that β_1 and β_2 are fit with ordinary least squares, with ε_1 and ε_2 the associated
632 residuals, then, $\varepsilon_1 = \varepsilon_2$.

Proof

634 Under ordinary least squares, the parameters are set as follows.

$$\beta_1 = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T Y \quad \text{(Eqn 3)}$$

$$\beta_2 = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T (Y - X) \quad \text{(Eqn 4)}$$

637 We start with the second of these, and using left distributivity of matrices, and then substituting Eqn
638 3, we obtain the following.

$$\beta_2 = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T (Y - X) = (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T Y - (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T X = \beta_1 - (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T X$$

640 Using the fact that the variable X is demeaned, we can now evaluate the main term here as follows,

$$\beta_2 = \beta_1 - (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T X = \beta_1 - \begin{pmatrix} X^2 & \Sigma X \\ \Sigma X & N \end{pmatrix}^{-1} \begin{pmatrix} X^2 \\ \Sigma X \end{pmatrix} = \beta_1 - \frac{1}{A} \begin{pmatrix} N & -\Sigma X \\ -\Sigma X & X^2 \end{pmatrix} \begin{pmatrix} X^2 \\ \Sigma X \end{pmatrix}$$

642 where X^2 is the dot product of X with itself, ΣX is the sum of the vector X, and $A = NX^2 - \Sigma X \Sigma X$ is
643 the determinant of the matrix being inverted. From here we can derive the following,

$$\beta_2 = \beta_1 - \frac{1}{A} \begin{pmatrix} NX^2 - \Sigma X \Sigma X \\ -\Sigma X.X^2 + X^2.\Sigma X \end{pmatrix} = \beta_1 - \frac{1}{A} \begin{pmatrix} A \\ 0 \end{pmatrix} = \beta_1 - \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

645 We can then substitute this equality for β_2 in eqn 2 and re-arrange to obtain,

$$Y - X = \tilde{X}\beta_2 + \varepsilon_2 = \tilde{X}\left(\beta_1 - \begin{pmatrix} 1 \\ 0 \end{pmatrix}\right) + \varepsilon_2 = \tilde{X}\beta_1 - X + \varepsilon_2$$

647 It follows straightforwardly from here that,

$$Y - \tilde{X}\beta_1 = \varepsilon_2$$

i.e. $\varepsilon_1 = \varepsilon_2$, as required.   QED

Proposition 10 shows that the residuals resulting from fitting equations 1 and 2 will be the same. A
consequence of this is that the error variability will be the same. As a result of this, the factor that
determines whether more variance is explained when regressing Y onto X or when regressing $Y - X$

onto X, is the variance available to explain. That is, the relative variance of Y and $Y - X$ drive the R^2

values of these two regressions. This then implicates the variance of Y and X and in fact their

covariance (which impacts the variance of $Y - X$).

More precisely, we can state the following.

1) If $\sigma^2_{(Y-X)}$ is big relative to σ^2_Y, then regressing $Y - X$ onto X will explain more variability than

regressing Y onto X.

2) If $\sigma^2_{(Y-X)}$ is small relative to σ^2_Y, then regressing $Y - X$ onto X will explain less variability than

regressing Y onto X.

Supplementary Appendix B: illustrating the relationship between the correlations

```matlab
% This function illustrates the relationship
function [r_XY,std_Y,r2,r3] = CheckEqn1()

noise = [0.01:0.01:1,2:100]; % controls r(X, Y)
scale = [0.01:0.01:1,2:100]; % controls sigma_Y/sigma_X
X = single(randn(1000,1));
for j=1:length(noise)
    Y = X + single(randn(1000,1).*noise(j)); %Y is X plus noise
    Y = zscore(Y); % then scale to X so the actual scaling is consistent
    for k=1:length(scale)
        Yl = Y.*scale(k); % rescale to control the variability ratio
        r_XY(j,k) = corr(X,Y); % calculate the correlation with outcomes

        r2(j,k) = corr(X,Yl-X); % calculate the correlation with change
        std_Y(j,k) = std(Yl)./std(X); % record the variability ratio
        r3(j,k) = eqn_r_X_XminusY(r_XY(j),std_Y(j,k)); % check Equation 1
    end
end

% display the resulting surface (Figure 1)
figure,surf(log(std_Y),r_XY,r3,'edgecolor','none')
lighting flat
l = light('Position',[50 100 100]);
l = light('Position',[50 100 -50]);
l = light('Position',[50 -100 -50]);
l = light('Position',[-50 -15 29]);
l = light('Position',[-50 -15 -29]);
l = light('Position',[-50 15 -29]);
l = light('Position',[50 15 -29]);
l = light('Position',[50 15 -50]);
shading interp
xlabel('log ( sigmaY / sigmaX )')
ylabel('r(X, Y)')
zlabel('r(X, Y-X)')

% confirm that equation 1 does actually match 'empirical' r(X,Y-X)
figure,scatter(r2(:),r3(:))
xlabel('Empirical coefficients')
ylabel('Derived coefficients')

end

% This function implements Equation 1
function res = eqn_r_X_XminusY(r_XY,std_Y)

res = (((r_XY.*std_Y) - 1) ./ sqrt(1 + (std_Y).^2 - (2*(r_XY.*std_Y)))));

end
```

 Supplementary Appendix C: simulating recovery

```matlab
720    function out = mprr_sim_pub(measure_noises, squash, nonfit)
721
722    % measure_noises = 300 in the 'high noise' case, and otherwise 3
723
724    % squash = true, if ceiling effects are imposed
725
726    % nonfit = true, if the 19 percent whose residuals
727    % to r(X, delta) are most extremely negative, are removed. The
728    % 19 percent figure comes from the study by Winters and
729    % colleagues, 2015
730
731    sample_size = 1000; % big sample
732    recover_noises = 0:0.01:1; % true variation in recovery
733    repeats = 100;
734    z=0; % flag to zscore everything before correlating
735
736    for r=1:repeats
737        for s=1:length(sample_size)
738            ss = sample_size(s);
739            for m=1:length(measure_noises)
740                mm = measure_noises(m);
741                for rec=1:length(recover_noises)
742                    rr = recover_noises(rec);
743
744                    rX = randi(66,ss,1) - 1; % true baseline
745                    initial_std = std(rX);
746                    epsilon = randn(ss,1).*mm; % baselines measurement noise
747                    epsilon2 = randn(ss,1).*mm; % outcome measurement noise
748                    X = rX + epsilon; % observed baseline
749
750                    % random recovery
751                    rR_rand = randn(ss,1).*(rr.* initial_std .*10);
752                    Y_rand = rX + rR_rand + epsilon2; % observed outcome
753                    if(squash>0)
754                        Y_rand(Y_rand>66) = 66;
755                        Y_rand(Y_rand<0) = 0;
756                    end
757
758                    % proportional recovery
759                    rR_prop = (66 - rX).*(0.7+(randn(ss,1).*rr)); % 70% of lost function
760                    Y_prop = rX + rR_prop + epsilon2;% observed outcome
761                    if(squash>0)
762                        Y_prop(Y_prop>66) = 66;
763                        Y_prop(Y_prop<0) = 0;
764                        %Y_prop = Y_sort(idx);
765                    end
766
767                    % constant recovery
768                    rR_const = 66.*(0.7+(randn(ss,1).*rr)); % 70% of 50
769                    Y_const = rX + rR_const + epsilon2;% observed outcome
770                    if(squash>0)
771                        Y_const(Y_const>66) = 66;
772                        Y_const(Y_const<0) = 0;
773                    end
774                    if(z>0)
775                        X=zscore(X);
776                        Y_rand=zscore(Y_rand);
777                        Y_prop=zscore(Y_prop);
778                        Y_invprop=zscore(Y_invprop);
779                        Y_const=zscore(Y_const);
780                    end
781                    if(nonfit>0) % remove the bottom 19% of recoverers.
782                        mdl = fitlm(X,Y_const);
783                        [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
784                        s_const = (1:sample_size)';
785                        s_const = ~ismember(s_const, idx(1:ceil(sample_size*0.19)));
786                        mdl = fitlm(X,Y_rand);
787                        [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
788                        s_rand = (1:sample_size)';
```

```matlab
                s_rand = ~ismember(s_rand, idx(1:ceil(sample_size*0.19)));
                mdl = fitlm(X,Y_prop);
                [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
                s_prop = (1:sample_size)';
                s_prop = ~ismember(s_prop, idx(1:ceil(sample_size*0.19)));
                mdl = fitlm(X,Y_invprop);
                [resid,idx] = sort(table2array(mdl.Residuals(:,1)));
                s_invprop = (1:sample_size)';
                s_invprop = ~ismember(s_invprop, idx(1:ceil(sample_size*0.19)));
            else
                s_const = (1:sample_size)';
                s_rand = (1:sample_size)';
                s_prop = (1:sample_size)';
                s_invprop = (1:sample_size)';
            end

            % run the correlations
            r_rand(r,rec,:) = [corr(X(s_rand),[Y_rand(s_rand), Y_rand(s_rand) -
X(s_rand)]),log(var(Y_rand(s_rand))/var(X(s_rand)))];
                r_prop(r,rec,:) = [corr(X(s_prop),[Y_prop(s_prop), Y_prop(s_prop) -
X(s_prop)]),log(var(Y_prop(s_prop))/var(X(s_prop)))];
                r_const(r,rec,:) = [corr(X(s_const),[Y_const(s_const), Y_const(s_const) -
X(s_const)]),log(var(Y_const(s_const))/var(X(s_const)))];
            end
        end
    end
end
r_rand=squeeze(nanmean(r_rand,1));
r_prop=squeeze(nanmean(r_prop,1));
r_const=squeeze(nanmean(r_const,1));
out = [r_rand,r_prop,r_const];
```