Maternal pluripotency factors prime the zygotic genome to respond to intercellular signals

George E. Gentsch[1,*], Thomas Spruce[1,2], Nick D. L. Owens[1,3], and James C. Smith[1,*]

[1] Developmental Biology Laboratory, The Francis Crick Institute, 1 Midland Road, London, NW1 1AT, United Kingdom

[2] Present address: Centre for Genomic Regulation (CRG), The Barcelona Institute for Science and Technology, Dr Aiguader 88, 08003 Barcelona, Spain

[3] Present address: Epigenetics of Stem Cells, Department of Stem Cell and Developmental Biology, Pasteur Institute, CNRS UMR3738, 25 Rue du Dr Roux, 75015 Paris, France

* Corresponding Authors
George E. Gentsch and James C. Smith

The ability of cells to respond to intercellular signals, and the repertoire of possible responses to those signals, is defined by their 'competence', a long-standing developmental concept[1], that is key to tissue formation and tissue regeneration. Apart from a few cell type-specific transcription factors[2], little is known about the role of the chromatin landscape in defining competence. Here, we investigate the emergence of competence in *Xenopus tropicalis,* which, like most multicellular organisms, begins development with a transcriptionally silent genome. We first identified the earliest functional regulatory DNA sequences, and inferred from them the critical sequence-specific activators of the zygotic genome. Of these, we showed that the maternal pluripotency factors Pou5f3 and Sox3 specifically open thousands of compacted chromatin sites by recognising their canonical DNA binding sequences. This pioneering activity initiates local chromatin remodelling to facilitate poised or active transcription, including the post-translational marking of displaced nucleosomes and chromatin looping with promoters. These remodelled chromatin sites are then capable of responding to intercellular signalling by recruiting the signal mediators that convert signals into tissue-specific expression of developmentally critical genes. As well as shedding light on genome activation and competence, our findings suggest that transcriptional responses to signals can be re-awakened by overexpressing the appropriate pioneer factors—a much sought-after objective in cell replacement therapies.

Zygotic genome activation (ZGA) in the *Xenopus* embryo occurs in a defined spatio-temporal manner around the mid-blastula transition (MBT) and depends on maternal gene products and inductive signalling between cells[3] (Extended Data Fig. 1a). Thus, spatially restricted Wnt, Nodal and BMP signals trigger the nuclear accumulation of their signal mediators β-catenin, Smad2 and Smad1, respectively, which in turn induce tissue-specific transcriptional responses that control the formation of the body axes and the formation of the three germ layers (ectoderm, mesoderm and endoderm) from pluripotent tissue[4]. The same signals are used subsequently in later development to elicit different responses, and we wondered whether the chromatin landscape might underlie these differences in competence.

To identify chromatin factors likely to confer competence during ZGA, we first examined the DNA motifs present at transcriptionally engaged *cis*-regulatory modules (CRMs) from the 32-cell to the late gastrula stage, by mapping focal RNA polymerase II (RNAPII) depositions on a genome-wide scale (Fig. 1a). RNAPII recruitment to CRMs proved to be dynamic, with the largest changes occurring between the 1,000-cell stage and MBT (Fig. 1a). In particular, pre-MBT recruitment coincided with the enrichment of canonical recognition motifs of the FOXH, POU, SOX and T domain transcription factor (TF) families (Fig. 1b; Extended Data Fig. 1b). Using this approach some of the RNAPII-engaged intronic or gene proximal CRMs were not detectable because they were masked by RNAPII-mediated gene transcription. Thus, MBT-specific CRMs were further characterised for chromatin accessibility (using a novel DNase-Seq method; see exemplar comparison with other chromatin features in Extended Data Fig. 1c) and for the enhancer-associated histone mark H3K4me1. We detected ~20,000 accessible CRMs occupied by RNAPII and flanked by H3K4me1$^+$ nucleosomes (Fig. 1c). The enriched DNA motifs among accessible H3K4me1$^+$/RNAPII$^+$ CRMs were then correlated with the maternally inherited and translated sequence-specific factors identified by egg-staged mass spectrometry[5] and pre-MBT ribosome footprinting[6] (Fig. 1d; Extended Data Fig. 1d; Supplementary Table 1). It proved that the binding preference of the most frequently translated maternal TFs and signal mediators matched the most significantly enriched DNA motifs (Fig. 1e). These were POU-SOX (Pou5f3-Sox3 heterodimer), Krüppel ZF (Sp/Klf), POU (Pou5f3), SOX (Sox3), bZIP (Max), FOXH (Foxh1), ETS (Ets2), NFY (NFYa/b/c), SMAD (Smad1/2), T (mVegT, a maternal VegT isoform), TCF (Tcf/β-catenin), basic helix-span-helix domain (bHSH; Tfap2), and OTX (Otx1). Notably, the enrichment level of these motifs were independent of CRM sequence conservation (based on phastCons and phyloP scoring) detected among vertebrates (Fig. 1c,e).

Figure 1. CRM-enriched DNA motifs match binding preferences of maternal TFs most frequently translated before MBT {dimension of figure: 1 column wide}

a,b, RNAPII recruitment to CRMs from 32-cell to late gastrula stage. **a**, Snapshots and Spearman correlations R_s. **b**, Z-score of enriched DNA motifs. **c**, MBT-staged map of DNase-probed chromatin accessibility, RNAPII binding and H3K4me1 marking grouped by sequence

conservation levels (phastCons, phyloP) and sorted by statistical significance of DHSs. **d**,
Maternal protein concentrations[5] versus ribosome footprint levels at 8-cell stage[6]. Most
frequently translated representatives of various TF families are labelled. **e**, Matching canonical
CRM-enriched DNA motifs (sorted by statistical significance) with frequently translated
sequence-specific factors. **f**, Graphical illustration of protein levels and nuclear localisations
of selected proteins based on our and previously published result[7-12]. Shaded boxes indicate
periods of non-nuclear protein distribution. Arrows indicate tissue movements. Abbreviations:
An, animal; D, dorsal; MZ, marginal zone; V, ventral; Vg, vegetal.
85
Based on this orthogonal correlation, we selected Pou5f3, Sox3 and mVegT as factors that
might specify responsiveness to Wnt, Nodal and BMP signalling. Sox3 and presumably also
Pou5f3 (based on the spatial distribution of its transcripts[13]) are detected ubiquitously, while
mVegT is restricted to the vegetal hemisphere[9] (Fig. 1f; Extended Data Fig. 1g). With respect
to signal mediators, nuclear β-catenin accumulates dorsally (first detected at the 32-cell
stage[8]) before spreading around the forming blastopore[11] (Fig. 1f). Signal-induced nuclear
accumulation of Smad1 and Smad2 is first detected around MBT with Smad1 being
preferentially ventral and Smad2 being vegetal and within the marginal zone[11,12] (Fig. 1f;
Extended Data Fig. 1e).
95
Next, informed by the onset of their nuclear accumulation (Extended Data Fig. 1a), we
generated genome-wide chromatin profiles across MBT to ask whether the recruitment of β-
catenin, Smad1 and Smad2 might be influenced by co-expressed maternal TFs like Sox3,
mVegT and Foxh1[14] (Extended Data Fig. 2 and 3a). As an outgroup control, we selected the
binding profiles of several zygotic T-box TFs Eomes[15], zVegT[15], Xbra[15] and Tbx6 (Extended
Data Fig. 2 and 3b), which collectively regulate the neuro-mesodermal cell lineage during and
beyond gastrulation[16].
103
Maternal TFs and signal mediators shared many chromatin characteristics: (1) DNA
occupancy levels followed a log-normal distribution with only a few hundred RNAPII-
transcribed zygotic genes receiving high and super-enhancer-like[17] input (Extended Data Fig.
3a-c). (2) From the 1,000-cell to the late gastrula stage, frequent chromatin recruitment
occurred on average to ~800 super-enhancers which had zygotic genes nearby (<5kb) that
are primarily involved in early embryonic processes (Extended Data Fig. 3d). (3) Among all
zygotic genes, promoter-proximal regions were most consistently bound (Extended Data Fig.
3e). All these characteristics were also observed for chromatin accessibility and RNAPII
engagement at CRMs (Extended Data Fig. 3c,e). The comparison of DNA occupancy levels
among and between signal mediators and TFs suggests that the recruitment of both depends
on factor-intrinsic and developmental context-dependent elements (Extended Data Fig. 4a,b).
The elements associated with developmental context were largely revealed by the binding

pattern of sequence-nonspecific RNAPII at CRMs (Extended Data Fig. 4c). The discovery and co-clustering of enriched DNA motifs at occupied CRMs indicate that factor-intrinsic elements include the sequence-specificity of DNA binding domains, oligomerisation tendencies (e.g. Xbra binding palindromic motifs due to its propensity to form homodimers[18]) and protein-protein interactions, while the developmental context reflects co-expressed TFs (Fig. 2a; Extended Data Fig. 4d,e).

To demonstrate the importance of developmental context on chromatin engagement, we ectopically expressed HA-tagged MyoD (MyoD-HA) in animal cap cells and profiled the chromatin for MyoD-HA at the early gastrula stage, as well as for endogenous Sox3 and RNAPII (Fig. 2b). MyoD was chosen because its canonical E-box recognition motif is normally not enriched at the early gastrula stage (Fig. 2a), while Sox3 and RNAPII were selected because they are ubiquitously expressed and represent sequence-specific and nonspecific DNA binding factors, respectively. As predicted, the E-box motif of MyoD-HA emerged as a context-dependent motif of Sox3 and RNAPII binding, while MyoD-HA binding itself was influenced by the endogenous TF expression profile (Fig. 2c,e; Extended Data Fig. 5a,b). MyoD-HA co-recruited Sox3 to MyoD gene targets like *actc1* and *myl1* (Fig. 2d; Extended Data Fig. 5c,d). However, this so-called opportunistic recruitment to non-canonical binding sites, such as MyoD-HA to Sox3 gene targets (e.g. *otx2*, *sox2*), did not affect transcription in animal caps (Extended Data Fig. 5c-e).

The influence of TF co-expression on DNA occupancy was further substantiated by profiling chromatin for Sox3 in different anterior-posterior regions of the central nervous system (CNS; Fig. 2f). The analysis of enriched DNA motifs suggested that Sox3 binding was influenced by differentially expressed homeodomain TFs such as orthodenticle homeobox (OTX) in the brain (head) and caudal homeobox (CDX) in the spinal cord (trunk, tailbud) (Fig. 2h; Extended Data Fig. 5f-h). This was particularly apparent in Sox3 binding to the colinear *HoxD* cluster defining anterior-posterior cell identity (Fig. 2g). Similar influences on chromatin engagement were observed within anterior and posterior mesoderm marked by Eomes in gastrula embryos and Xbra/Tbx6 in gastrula and early tailbud embryos, respectively (Fig. 2h; Extended Data Fig. 5f). On a temporal scale, the influence of FOXH and POU motif-recognising TFs in recruiting Sox3 and T-box TF was more pronounced early than late in development (Fig. 2h; Extended Data Fig. 5f). Interestingly, the analysis of enriched DNA motifs suggests that maternal TFs affect the recruitment of signal mediators, but (with the exception of Foxh1, which interacts directly with Smad2[19]) the reverse is much less the case (compare highlighted orange and black boxes in Fig. 2a; Extended Data Fig. 4d,e).

Figure 2. Developmental context influences chromatin engagement {dimension of figure:

1 column wide}

a, Significantly enriched DNA motifs at CRMs occupied by indicated proteins covering
developmental process of embryonic pluripotency and neuromesodermal (NM) bipotency. **b-**
e, MyoD-enforced recruitment of Sox3 and RNAPII. **b**, Experimental design. **c**, PCA of
chromatin engagement. **d**, Snapshot of chromatin co-recruitment. **e**, Significantly enriched
motifs. **f-h**, CNS-specific Sox3 binding along the anterior-posterior axis (head, trunk and bud).
f, Experimental design. **g**, Snapshot of Sox3 binding to *HoxD* cluster. **h**, Spearman correlation
R_s of TF binding. Below, heat map of significantly enriched DNA motifs.

To ask whether signal-mediated ZGA requires maternal TFs, as suggested by the observed

chromatin dynamics, we next compared the effects of loss of maternal Sox3/PouV (Pou5f3.2

and Pou5f3.3; mPouV) or VegT (mVegT), and canonical Wnt, Nodal or BMP signal

transduction, on zygotic transcription from MBT to late blastula and early gastrula stage. The

high translation frequencies (Fig. 1d; Extended Data Fig. 1d) coupled with steady or even

transient protein levels around MBT (Fig. 1f), suggest that maternal TFs and signal mediators

have short half-lives. Consistent with this idea, the injection of morpholinos (MOs) blocking

the translation of maternal transcripts such as *Sox3* or *mVegT* was effective in reducing protein

levels (Extended Data Fig. 1f,i). We used previously validated MOs to knock down mPouV[20]

and β-catenin[21]. The nuclear accumulation of Smad1 and Smad2 in response to BMP and

Nodal signalling was inhibited with the small molecules LDN193189 and SB431542,

respectively. The morphological defects of these loss-of-function (LOF) treatments ranged

from undetectable (Sox3), to weak (mVegT, BMP), to moderate (mPouV), to severe

(mPouV/Sox3, Nodal, β-catenin) (Extended Data Fig. 6a,c,e). Moderate and severe defects

affected gastrulation, while weak ones were only obvious later. All phenotypes were either in

line with previous publications (β-catenin[21], Nodal[22], BMP[23]) or could be rescued on a

morphological (mPouV/Sox3; Extended Data Fig. 6b) or transcriptional level (mVegT,

Extended Data Fig. 6d) by co-injecting cognate mRNA. Interestingly, the role of maternal Sox3

could only be detected by knocking it down together with mPouV. In contrast to single Sox3

or mPouV LOF, double LOF embryos failed to close the blastopore (Extended Data Fig. 6a;

Supplementary Video 1).

Transcriptome analysis of TF and signal LOF embryos was confined to the 3,318 zygotically

expressed genes for which ≥two-fold reductions in exonic and/or intronic transcript level could

be detected following injection of the RNAPII inhibitor α-amanitin (Supplementary Table 2).

About 86% of these genes had a maternal contribution of ≥1 per 10 million transcripts

(Extended Data Fig. 7a). α-amanitin prevented any transcriptional changes and thus blocked

the gastrulation movements that are normally initiated by ZGA (Extended Data Fig. 6f,g). The

LOF-mediated reduction of gene activation ranged from ~1% for BMP to ~24% for

mPouV/Sox3 (Fig. 3a). We note that the additional loss of Sox3 in mPouV LOF embryos
further reduced transcription of developmentally relevant genes, including those that are
expressed ubiquitously, like *mir427*, or those that are activated in a subset of mPouV/Sox3[+]
cells, like *ventx* and *Xbra* (Extended Data Fig. 7b,c).

Spatial analysis of reduced gene activation confirmed that many of the 784 mPouV/Sox3-
dependent genes showed enriched expression along the animal-vegetal or dorso-ventral axis
(Fig. 3a). More specifically, comparison with signal LOFs revealed that 201 of the genes
tissue-specifically induced by Wnt, Nodal or BMP also depended on mPouV/Sox3 (Fig. 3b,c;
Extended Data Fig. 7d,f). Likewise, 142 of the 189 mVegT-dependent genes were activated
in response to signalling within the vegetal hemisphere, such as *foxa1*, *nodal5/6* and *sox17a/b*
(Extended Data Fig. 7d,e). Loss of mVegT and mPouV/Sox3 jointly affected 49 genes, 42 of
which also depended on Wnt, Nodal or BMP signalling (Extended Data Fig. 7f).

Remarkably, the requirement for particular maternal TFs varied even among related genes
with similar expression patterns that are activated by the same signals. For instance, *Xbra*
and *eomes* are both activated by Nodal signalling in the marginal zone between the animal
and vegetal hemispheres, but only *Xbra* requires mPouV/Sox3 for its signal responsiveness
(Fig. 3c). Significantly, DNA occupancy levels did not explain the difference in gene regulation
between *Xbra* and *eomes*, because both gene loci showed high levels of Sox3 binding (see
Fig. 4c). Identical dependencies were found for Wnt-responsive genes like *foxb1* and *zic1*
(Extended Data Fig. 7g)*.*

We confirmed these mPouV/Sox3-dependent signal inductions by treating control and
mPouV/Sox3-depleted naïve (animal cap) tissue with Nodal (Activin) and Wnt (CHR99021)
agonists (Fig. 3c; Extended Data Fig. 7h). Interestingly, mPouV/Sox3 could trigger basal low-
level expression of (especially) *Xbra* and *foxb1* in the absence of signalling. Together, our
selected maternal TFs and signals activated 1,213 of the 3,318 (~37%) zygotic genes. These
included ~58% of those expressed animally, ~88% of the vegetal genes, ~96% of the dorsal
genes and 90% of those expressed ventrally, while many ubiquitously expressed genes
remained unaffected (Fig. 3a). Thus, mPouV and Sox3 tend to affect biological functions for
which tissue-specific expression is essential, such as main body axis formation,
regionalisation and germ layer segregation (Extended Data Fig. 8b). We also note that LOF
of mPouV/Sox3 also caused significant increases in transcript levels of active genes encoding
gamete-specific biological processes, suggesting that reprogramming towards embryonic
pluripotency was compromised (Extended Data Fig. 8a,b).

Figure 3. Signal-induced ZGA depends on maternal TFs {dimension of figure: 1 column wide}

a, Regional expression of genes downregulated by indicated LOFs. Regional expression refers to transcript level ratios along the animal-vegetal (An:Vg) or dorso-ventral (D:V) axis in unperturbed early gastrula embryos[24]. Numbers and percentages of downregulated genes are

either specific to regional expression or reflect overall statistics. **b**, Scatterplot of transcript level changes caused by LOF of mPouV/Sox3 or Nodal signalling. Dots are coloured according to An:Vg transcript level ratios. Numbers refer to some more developmentally relevant genes (see Extended Data Fig. 5h). **c**, Venn diagram of downregulated genes. **d**, Early gastrula-staged WMISH for *Xbra* and *eomes* under indicated LOFs. Scale bar, 0.5 mm. **e**, Quantification of *Xbra* and *eomes* transcript levels in control and mPouV/Sox3 (mPS) LOF animal caps with or without Nodal signalling. Error bars, mean+SD (n=2). Two-tailed Student's t-test: 1, p=0.014; 2, p=0.005; 3, p=0.126; and 4, p=0.019.

To discover how maternal TFs identify genes to become competent to respond to signals, we compared various chromatin features between control and mPouV/Sox3 LOF embryos at MBT (Fig. 4a). Genome-wide analysis showed that mPouV/Sox3 LOF strongly reduced chromatin accessibility as measured by the significant loss of cleavages in 6,738 of the 16,637 (~41%) DNase hypersensitive sites (DHSs) (Fig. 4b; Extended Data Fig. 9a-c). Sorting of DHSs according to the significance of lost accessibility indicates that accessibility depends on the ability of mPouV and Sox3 to recognise their canonical motifs (Fig. 4b; Extended Data Fig. 9d). By comparison, unaffected DHSs rarely contained canonical mPouV/Sox3 motifs, and were strongly enriched for promoter-centric motifs[25] of the Krüppel ZF, bZIP and NFY protein family (Fig. 4b). Indeed, promoters (+/-0.5 kb from a transcription start site, TSS) were less frequently located within mPouV/Sox3-affected (~4%) than unaffected DHSs (~29%). By contrast, at the extreme end of affected loci (e.g. *Xbra*, *foxb1*, *cdc25b* and *zic1*) entire super-enhancers became accessible through the activity of mPouV/Sox3 (Fig. 4c, Extended Data Fig. 10d).

The loss of accessibility at distal DHSs triggered further profound changes to chromatin *in situ* (see arrowheads in Fig. 4c and Extended Data Fig. 10d for strongly affected CRMs). Chromatin looping from affected DHSs to promoters (Fig. 4c; Extended Data Fig. 10a-d) was significantly reduced and so was the deposition of DHS-flanking H3K4me1 and co-recruitment of RNAPII and signal mediators Smad2 and β-catenin (Fig. 4b,c; Extended Data Fig. 9d). At the transcriptional level, a reduction of non-coding and coding RNA also coincided with the affected DHSs (Fig. 4b; Extended Data Figs. 9d and 10d). Importantly, the differential chromatin profiling explained how tissue competence can be established and why, for instance, Nodal-induced transcription of *Xbra* was strongly affected by the loss of mPouV/Sox3 and that of *Eomes* less so. Thus, in contrast to those of *eomes*, all promoter-tied CRMs of *Xbra* contain canonical POU/SOX motifs, so they were not accessible to Smad2 in mPouV/Sox3 LOF embryos (Fig. 4c,d). Thus, Smad2 interactions with critical Nodal responsive regulatory elements of *eomes* remained intact, but those of *Xbra* were

Figure 4. Maternal pluripotency factors pioneer silent zygotic genome to control its

response to signalling {dimension of figure: 1 column wide}
a, Approach to reveal effect of mPouV/Sox3 on chromatin landscape at MBT. **b**, Heat map of
chromatin and RNA profiles at DHSs sorted by significantly reduced DNase cleavages under
mPouV/Sox3 LOF. [#], β-catenin binding profile generated at late blastula stage rather than
early blastula stage (Sox3) or MBT (all others). To the right, co-aligned map of DNA motifs,
distance of DHSs to nearest TSSs and correlation with mPouV/Sox3-dependent ZGA. **c**,
Superimposed line plots showing the LOF-induced changes to various chromatin features
including chromatin contacts (capture-C) with promoters (viewpoints) of *Xbra* and *eomes*. For
statistical significance check heat map below each profile. Header, POU/SOX motif
occurrences at DHSs and sequence conservation heat maps (phastCons, phyloP). Arrowhead
denotes strongly affected chromatin site. **c**, Model of pioneer-initiated chromatin remodelling
to facilitate the transcriptional response to signals during ZGA.
283
284 impeded by closed chromatin. mPouV/Sox3-dependent tissue competence also applied to
285 Wnt signalling with β-catenin failing to access Wnt responsive regulatory elements of *foxb1* in
286 the absence of mPouV/Sox3 (Extended Data Fig. 10d).
287
Our results allow us to propose a model (Fig. 4d) of pioneer-initiated chromatin remodelling
that facilitates the transcriptional response to inductive signals and thus defines competence
during ZGA. First, the maternal TFs mPouV and Sox3 are recruited to silent CRMs either by
recognising canonical DNA motifs or in a more opportunistic manner, presumably through the
activity of other pioneers. Based on *in vitro* experiments[26], pioneering activity may require only
partial motifs to be displayed on nucleosome surfaces to initiate access to compacted
chromatin. Subsequent displacement of nucleosomes may be driven by the higher affinity of
TFs for free DNA. In contrast, opportunistic binding occurs independent of sequence
preference and is influenced by developmental context such as the co-expression of
sequence-specific TFs as shown for the recruitment of Sox3 in various tissues. However, while
pioneering activity triggers extensive local chromatin remodelling (e.g. H3K4me1 marking of
displaced nucleosomes and the chromatin looping with promoters), opportunistic binding
seems to have much less of an effect on chromatin.
301
We suggest that pioneer-initiated chromatin remodelling, or priming, occurs in every
embryonic cell before the onset of Wnt, Nodal or BMP signalling to allow the recruitment of
signal mediators. This is for three reasons. First, the nuclear accumulation of mPouV and
Sox3 occurs ubiquitously and precedes that of signal mediators. Second, mPouV/Sox3 LOF
affects chromatin similarly at zygotic genes with (e.g. *foxb1*) or without (e.g. *cdc25b*) tissue-
specific expression. And finally, dissected animal caps replicate the *in vivo* response of the
marginal zone tissue to Nodal and Wnt signalling.
309

Our work has focussed on the pioneer activity of mPouV and Sox3, and, based on the recognition motifs enriched at active CRMs before MBT, we suggest that only a few more maternal TF families, including those of T-box, FoxH and Krüppel-like ZF families, are involved in the acquisition of competence at the ZGA. Indeed, many of these TF family members are relatively abundant and rapidly turned over, suggesting strong and dynamic TF activity[27]. Most of these TFs are already present in the *Xenopus* egg, which confers pluripotency to somatic cells when they are injected into its cytoplasm[28]. Their activity generates a signature of accessible DNA motifs at the MBT that is conserved among vertebrates and most reminiscent of the chromatin footprints in embryonic stem cells[25].

The findings presented here and elsewhere[29] reveal close connections between ZGA and pluripotency, and we anticipate that competence is controlled in human embryonic stem cells in a manner resembling that in the early *Xenopus* embryo. The approach described here can be applied to any other cell type to discover the molecular basis of its competence. Interestingly, various tumours are associated with the mis-expression of pioneer factors[30] and thus may display different competencies from the surrounding host tissue. Conversely, knowing which pioneers are required to trigger a specific signal response will increase the success of engineering patient-specific tissue for transplantation therapy.

Methods

***Xenopus* manipulation**. Standard procedures were used for ovulation, fertilisation, and manipulation and incubation of embryos[31,32]. Briefly, frogs were obtained from Nasco (Wisconsin, USA). Ovulation was induced by injecting serum gonadotropin (Intervet) and chorionic gonadotropin (Intervet) into the dorsal lymph sac of mature female frogs. Eggs were fertilised *in vitro* with sperm solution including 90% Leibovitz's L-15 medium (Gibco) and 10% fetal calf serum (Gibco). After 10 min, fertilised eggs were de-jellied with 2.2% L-cysteine (Sigma) equilibrated to pH 8.0. *X. tropicalis* embryos were cultured in 5% Marc's Modified Ringer's solution[32] (MMR) at 21ºC-28ºC. *X. laevis* embryos were cultured in 10% Normal Amphibian Medium[32] (NAM) at 14ºC-22ºC. Embryos were staged as previously described[33]. All *Xenopus* work complied fully with the UK Animals (Scientific Procedures) Act 1986 as implemented by the Francis Crick Institute.

Nucleic acid injections and treatments of embryos. Microinjections were performed using calibrated needles and embryos equilibrated in 4% Ficoll PM-400 (Sigma) in 5% MMR or 10% NAM. Microinjection needles were generated from borosilicate glass capillaries (Harvard

Apparatus, GC120-15) using the micropipette puller Sutter p97. Maximally two nanolitres of

morpholino (MO) and/or mRNA were injected into de-jellied embryos at the 1-cell, 2-cell or 4-

cell stage using the microinjector Narishige IM-300. Injections for the mVegT LOF and rescue

were targeted to the vegetal hemisphere. All other injections were targeted to the animal

hemisphere. Embryos were transferred to fresh 5% MMR or 10% NAM (without Ficoll) once

they reached about stage 8.

The following mRNA amounts were used for ectopic expression: 80 pg and 400 pg *MyoD-HA*

mRNA into *X. tropicalis* and *X. laevis* embryos, respectively; and 400 pg *Sox3-HA* into *X.*

laevis embryos.

The following MO amounts and treatments were used for LOF experiments: 5 ng *Pou5f3.2*

MO, 5 ng *Pou5f3.3* MO and 5 ng standard control MO (mPouV LOF); 5 ng *Pou5f3.2* MO, 5 ng

Pou5f3.3 MO and 5 ng *Sox3* MO (mPouV/Sox3 LOF); 5 ng *β-catenin* MO (Wnt LOF); 10 ng

m*VegT* MO (mVegT LOF); standard control MO (5-10 ng according to the dose used for the

β-catenin or mVegT LOF experiment); and 30 pg α-amanitin (Sigma). To block Nodal (Nodal

LOF) or BMP (BMP LOF) signalling, embryos were treated with 100 μM SB431542 or 10 μM

LDN19318 from the 8-cell stage onwards. Control embryos were treated accordingly with

DMSO, in which these antagonists were dissolved.

For the rescue of mVegT LOF, 25-50 pg z*VegT* mRNA were injected at the 2- or 4-cell stage.

For the rescue of mPouV/Sox3 LOF, 75 pg *Pou5f3.3* mRNA and 75 pg *Sox2* mRNA were

injected at the 2- or 4-cell stage. For both rescue experiment, 150-300 pg *mCherry* mRNA

was co-injected to trace mRNA expression in embryonic cells.

Morpholino (MO) oligonucleotide sequences. MOs were designed and produced by

Genetools (Oregon, USA) to block the translation of the following gene transcripts:

β-catenin MO (5'-TTTCAAC<u>A</u>GTTTCCAAAGAACCAGG-3', underlined base was changed

from the original MO[21]), *Pou5f3.2* MO (5'-GCTGTTGGCTGTACAT<u>A</u>GTGTC-3', underlined

base was changed from the original MO[20]), *Pou5f3.3* MO[20] (5'-

GTACAGAACGGGTTGGTCCATGTTC-3'), *Sox3* MO (5'-

GTCTGTGTCCAACATGCTATACATC-3'), *Tbx6* MO (5'-

TACATTGGGTGCAGGGACCCTCTCA-3') and m*VegT* MO[34] (5'-

TGTGTTCCTGACAGCAGTTTCTCAT-3'; see Extended Data Fig. 1h).

RNA constructs. All constructs are products of Gateway (Invitrogen) cloning. First, the coding

sequences of interest were amplified from either plasmid using *Pfu*Turbo DNA polymerase

(Agilent) or cDNA using Phusion HF DNA polymerase (NEB), and unidirectionally cloned into

the pENTR/D-TOPO vector (Invitrogen). Next, these clones were LR-recombined with

Gateway-compatible pCS2+ vectors containing no tag or a C-terminal 3xHA tag (see

Extended Data Fig. 5c). The following forward and reverse primers were used for PCR

amplification: *X. laevis myoD1a* from *MyoD*2-24 pBluescript plasmid[35], 5'-

CACCATGGAGCTGTTGCCCCCAC-3', 5'-TAAGACGTGATAGATGGTGCTG-3'; *X. laevis*

sox3 from IMAGE:443920 clone, 5'-CACCATGTATAGCATGTTGGACAC-3', 5'-

TATGTGAGTGAGCGGTACAGTG-3'; *X. laevis sox2* from stage 11 cDNA, 5'-

CACCATGTACAGCATGATGGAGACCGA-3', 5'-TCACATGTGCGACAGAGGCAGC-3'; *X.*

laevis pou5f3.3 from stage 11 cDNA, 5'-CACCATGGACCAGCCCATATTGTACA-3', 5'-

TCAGCCGGTCAGGACCCCCA-3'; *X. tropicalis zVegT* from stage 11 cDNA, 5'-

CACCATGCACTCTCTGCCGGATGTA-3',   5'-TTACCAACAGCTGTATGGAAAGA-3'.

pCS2+8N*mCherry*[36] (#34936) was obtained from Addgene.

***In vitro* transcription of capped mRNA.** About 10 µg plasmid was linearised by restriction

digestion and purified using the QIAquick PCR purification kit (Qiagen). Capped mRNA was

produced from ~1 µg linearised plasmid using mMessage mMachine SP6 (Ambion) according

to the manufacturer's instructions. The following restrictions enzymes (NEB) were used for

linearising the pCS2+ constructs: *Asp*781 (*myoD1a*, *sox3*), *Pvu*II (*sox2*), *Psp*OMI (*pou5f3.3*,

z*VegT*) and *Not*I (*mCherry*).

Generation of digoxigenin-labelled RNA probes. DNA template for generating the *foxb1 in*

situ hybridisation probe was amplified from *X. tropicalis* stage 19-20 cDNA. KAPA HiFi

HotStart polymerase and the following PCR cycling conditions were used: 98ºC for 45 sec

followed by 40 cycles (98ºC for 10 sec, 63ºC for 10 sec, 72ºC for 15 sec) and final elongation

step of 20 sec at 72ºC. Primers were 5'-GTCAGCGCCTATGGAGTACC-3' and 5'-

AACACTGGAGATGCCATGC-3'. Fresh PCR product was zero-blunt cloned into the pCRII-

TOPO vector (Invitrogen). The identity and direction of insert was verified by restriction digest

and Sanger sequencing. Plasmids were linearised by restriction digestion and purified using

the QIAquick PCR purification kit (Qiagen). All *in situ* hybridisation probes were transcribed

from ~1 µg linearised plasmid using 1x digoxigenin-11-UTP (Roche), 40 U RiboLock RNase

inhibitor (Thermo Fisher Scientific), 1x transcription buffer (Roche) and T7 RNA polymerase

(Roche) at 37ºC for 2 h. The probe was treated with 2 U Turbo DNase (Ambion) to remove

the DNA template and purified by spin-column chromatography (Clontech). RNA was diluted

to 10 ng/µl (10x stock) with hybridisation buffer (50% formamide, 5x SSC, 1x Denhardt's, 10

mM EDTA, 1 mg/ml torula RNA, 100 µg/ml heparin, 0.1% Tween-20 and 0.1% CHAPS) and

stored at -20ºC. The following plasmids and restriction enzymes (NEB) were used for plasmid

linearisation: *X. laevis eomes* pCRII-TOPO[15], *Bam*HI; *X. tropicalis foxb1* pCRII-TOPO, *Bam*HI;

X. tropicalis mir427 pCS108 (IMAGE:7545411)[37], *Sal*I; *X. laevis ventx2.1* pBluescript SK-[38],

*Eco*RI; *X. laevis Xbra* pSP73[39], *Bgl*II; and *X. tropicalis zic1* pCMV-SPORT6 (IMAGE:7668846),

*Sal*I.

Animal cap assay. These assays were carried out for the ectopic expression of mRNA

constructs in *X. laevis* and for the MO-mediated LOF in *X. tropicalis*. All animal caps were

dissected at the blastula stage (stage 8 to 9) and cultured in 75% NAM containing 0.1% bovine

serum albumin (Sigma). Dissections were carried out with 13 µm loop electrodes (Protech

International, 13-Y1) connected to a MC-2010 micro cautery instrument (Protech International)

operating at power level 2. As illustrated in Extended Data Fig. 5c, control (uninjected) and

MyoD-HA or Sox3-HA expressing animal caps were cultured at 20ºC until sibling embryos

reached early neurula stage (stage 13). For the experiment shown in Fig. 3e and Extended

Data Fig. 7h, animal caps of control and mPouV/Sox3 LOF embryos were cultured with or

without 10 ng/ml recombinant human activin A (R&D Systems, 338-AC) (Nodal agonist) or 50

µM CHIR99021 (Tocris, 4423) (canonical Wnt agonist) at 22ºC for ~2 h until sibling embryos

reached early gastrula stage (stage 10^+).

Extraction of total RNA. 10-15 embryos (or 15-20 animal caps) were homogenised in 800

(400) µl TRIzol reagent (Ambion) by vortexing. The homogenate was either snap-frozen in

liquid nitrogen and stored at -80ºC or processed immediately. For phase separation, the

homogenate together with 0.2x volume of chloroform was transferred to pre-spun 1.5-ml

Phase Lock Gel Heavy (PLGH) microcentrifuge tubes, shaken vigorously for 15 sec, left on

the bench for 2 min and spun at ~16,000 g (4ºC) for 5 min. The upper phase was mixed well

with one volume of 95-100% ethanol and spun through the columns of Zymo RNA Clean and

Concentrator 5 or 25 (Zymo Research) at ~12,000 g for 30 sec. Next, the manufacturer's

instructions were followed for the recovery of total RNA (>17 nt) with minor modifications. First,

the flow-through of the first spin was re-applied to the column. Second, the RNA was treated

in-column with 3 U Turbo DNase (Ambion). Third, the RNA was eluted twice with 25 µl

molecular-grade water. The concentration was determined on the NanoDrop 1000

spectrophotometer.

Reverse transcription (RT). About 0.2-1 µg total RNA was denatured at 75ºC for 5 min before

setting up the RT reaction including 40 U M-MLV RT, RNase H minus, point mutant (Promega)

500 µM of each dNTP (NEB), and 10 µM random hexamers (Sigma) in a final volume of 10

µl. The following incubation settings were used: 25ºC for 15 min, 37ºC for 15 min, 55ºC for 45

min, 85ºC for 15 min and 4ºC for indefinite. The reaction was diluted with 90-190 µl molecular-
grade water before using 1-2.5 µl for each qPCR reaction.
459
qPCR primer sequences. The PCR primers were designed to hybridize at ~60ºC (T_m) and to
generate 75-125 bp amplicons. Where possible, primers were checked for their specificity *in*
silico via the UCSC Genome Browser. All primer pairs were tested to produce one amplicon
of the correct size through diagnostic PCR using RedTaq DNA polymerase (Sigma).
Exceptions were previously designed primer pairs for *X. laevis actc1*, *odc1* and *otx2*, all of
which generate amplicons of 200-250 bp.
466
Primers (forward, reverse) for *X. laevis* (L) or *X. tropicalis* (T) cDNA: *actc1* (L, 5'-
TCCTGTACGCTTCTGGTCGTA-3', 5'-TCTCAAAGTCCAAAGCCACATA-3'); *eomes* (T, 5'-
ATTGACCACAACCCATTTGC-3', 5'-TCTAGGAGAATCCGCAGGAG-3'); *foxb1* (T, 5'-
CCGGATGCAGAGTGACAAC-3', 5'-GAGGCTTCTGGTCGCTGTAG-3'); *gata4* (T, 5'-
TCCTCTGATCAAGCCTCAAAG-3', 5'-ACGGCGCCAGAGTGTAGTAG-3'); *myl1* (L, 5'-
GCACCTGCATTTGATCTGTC-3', 5'-TTGCTGTCTCCTGTCCTGTC-3'); *nodal5* (T, 5'-
TTCAAGCCAACAAACCACGC-3', 5'-CGCATCTTCACCGGTATGCA-3'); *nodal6* (T, 5'-
CGCATTTCGTTGTGAGGGGT-3', 5'-TCGTACAGCTTGACCAAACTCT-3'); *odc1* (L, 5'-
GCCATTGTGAAGACTCTCTCC-3', 5'-TTCGGGTGATTCCTTGCCAC-3'); *odc1* (T, 5'-
GGGCAAAAGAGCTTAATGTGG-3', 5'-CATCGTGCATCTGAGACAGC-3'); *otx2* (L, 5'-
TTGAACCAGACCTGGACT-3', 5'-CGGGATGGATTTGTTGCA-3'); *rpl8* (T, 5'-
CTGGCTCCAAGAAGGTCATC-3', 5'-CAGGATGGGTTTGTCAATACG-3'); *sox2* (L, 5'-
ACATCACATGGACGGTTGTC-3', 5'-TCTCTCGGTTTCTGCTCTCG-3'); *sox17b* (T, 5'-
GTTTATGGTGTGGGCAAAGG-3', 5'-TCAGCGATTTCCATGATTTG-3'); *tbx6* (L, 5'-
GGGGCACCACGACTCTTAC-3', 5'-CCCAGAATCCTCTGTGAAGG-3'); and *Xbra* (T, 5'-
CCTGTGGATGAGGTTCAAGG-3', 5'-CACGCTCACCTTTAGAACTGG-3').
483
Primers (forward, reverse) for *X. tropicalis* genomic DNA: *β-actin* -0.5kb (5'-
TTGTACGCAACACAATGCAG-3', 5'- AAATGTCAGGACCAAATGCAG-3'); *dlc* -3.2kb (5'-
CATCCTGCAATTAACTGTCTGC-3', 5'-AGGTGTGACAAATGGCACTG-3'); *gdf3* -0.1kb (5'-
TGGATGAGCACAGAGAGGTG-3', 5'-GTCAGGAGGGAGGTGTCAAC-3'); *hoxd1* +3.2kb (5'-
TGTTGTAGATGCTGATGCTTATCG-3', 5'-AACAGAAAATCAAAGGCTTGCA-3'); *mespa* -
1.3kb (5'-CTCCTTTGCCCTCTGAAATG-3', 5'-CTTCAGGGAATTGGCTTGTG-3'); *msgn1* -
0.1kb (5'-CCAGTCCATTTTCCATGTTG-3', 5'-GGGGAGGCCCTTTTATACAG-3'); *myf5* -
1.1kb (5'-AAAGAGTTCCTGCACCTTGG-3', 5'-TCCCAAATCATCAGCACAAC-3'); *nodal6* -
0.1kb (5'-ACATGAGTTTGGCCTTGGTC-3', 5'-AGGTAACCCCTGAGCTGTCA-3'); *pax3*
5'UTR (5'-CAGATGCCAGAGGAGAGAGC-3', 5'-GGAACTCTCTGGGTCCCAAC-3'); *pax3*

+2.7kb (5'-ATGAGAATGTGAGCGACACG-3', 5'-AAACCCAAGGTGAGCAAGTG-3'); *sia1* -

0.2kb (5'-TGGCTTTTAAGTTCGTCTTGG-3', 5'-CAAAGGGCTTCTCTTTGTGC-3'); *sox3*

TSS (5'-GATGGGGCATTAATGGACAG-3', 5'-CACGGGAGAAAATGAAGCTG-3'); *sox17b* -

0.1kb (5'-GTTAGACTCCGCTCCAGACG-3', 5'-TCTCCACCTAAGGGGAAACC-3'); *wnt8a* -

5.7kb (5'-GGAAACTGCCTGTGGAAATG-3', 5'- ACAAAAGCCCACAGGAGATG-3'); *Xbra* -

4kb (5'-CAATGGATGCCTCTCTGGAC-3', 5'-TGGGTTTAGAATGGCAAAGG-3'); and *Xbra*

TSS (5'-CACACCTTGGGTTTTGTTCC-3', 5'-TTCCTCAAGAGAGGGGCTTC-3').

Whole mount *in situ* hybridisation (WMISH). WMISH was conducted using digoxigenin-

labelled RNA probes. It was based on previously published protocols[32,40]. *X. tropicalis*

embryos were fixed in MEMFA (1x MEM and 3.7% formaldehyde) at room temperature for 1

h. The embryos were then washed once in 1x PBS and two to three times in ethanol. Fixed

and dehydrated embryos were kept at -20ºC for ≥24 h to ensure proper dehydration before

starting hybridisation. Dehydrated embryos were washed once more in ethanol before

rehydrating them in two steps to PBT (1x PBS and 0.1% Tween-20). Embryos were treated

with 5 µg/ml proteinase K (Ambion) in PBT for 6 to 8 min, washed briefly in PBT, fixed again

in MEMFA for 20 min and washed three times in PBT. Embryos were transferred into baskets,

which are kept in an 8x8 microcentrifuge tube holder sitting inside a 10x10 slot plastic box

filled with PBT. Baskets were built by replacing the round bottom of 2-ml microcentrifuge tubes

with a Sefar Nitex mesh. This container system was used to readily process several batches

of embryos at once. These baskets were maximally loaded with 40 to 50 *X. tropicalis* embryos.

The microcentrifuge tube holder was used to transfer all baskets at once and to submerge

embryos into subsequent buffers of the WMISH protocol. Next, the embryos were incubated

in 500 µl hybridisation buffer (50% formamide, 5x SSC, 1x Denhardt's, 10 mM EDTA, 1 mg/ml

torula RNA, 100 µg/ml heparin, 0.1% Tween-20 and 0.1% CHAPS) for 2 h in a hybridisation

oven set to 60ºC. After this pre-hybridisation step, the embryos were transferred into 500 µl

digoxigenin-labelled probe (1 ng/µl) preheated to 60ºC and further incubated overnight at

60ºC. The pre-hybridisation buffer was kept at 60ºC. The next day embryos were transferred

back into the pre-hybridisation buffer and incubated at 60ºC for 10 min. Subsequently, they

were washed three times in 2x SSC/0.1% Tween-20 at 60ºC for 10 min, twice in 0.2x

SSC/0.1% Tween-20 at 60ºC for 20 min and twice in 1x maleic acid buffer (MAB) at room

temperature for 5 min. Next, the embryos were treated with blocking solution (2% Boehringer

Mannheim blocking reagent in 1x MAB) at room temperature for 30 min, and incubated in

antibody solution (10% lamb or goat serum, 2% Boehringer Mannheim blocking reagent, 1x

MAB and 1:2,000 Fab fragments from polyclonal anti-digoxigenin antibodies conjugated to

alkaline phosphatase) at room temperature for 4 h. The embryos were then washed four times

in 1x MAB for 10 min before leaving them in 1x MAB overnight at 4ºC. On the final day of the

WMISH protocol, the embryos were washed another three times in 1x MAB for 20 min and

equilibrated to working conditions of alkaline phosphatase (AP) for a total of 10 min by

submerging embryos twice into AP buffer (50 mM $MgCl_2$, 100 mM NaCl, 100 mM Tris-HCl pH

9.5 and 1% Tween-20). At this stage, the embryos were transferred to 5-ml glass vials for

monitoring the progression of the AP-driven colorimetric reaction. Any residual AP buffer was

discarded before adding 700 µl staining solution (AP buffer, 340 µg/ml nitro-blue tetrazolium

chloride, 175 µg/ml 5-bromo-4-chloro-3'-indolyphosphate). The colorimetric reaction was

developed at room temperature in the dark. Once the staining was clear and intense enough,

the colour reaction was stopped by two washes in 1x MAB. To stabilise and preserve

morphological features, the embryos were fixed with Bouin's fixative without picric acid (9%

formaldehyde and 5% glacial acetic acid) at room temperature for 30 min. Next, the embryos

were washed twice in 70% ethanol/PBT to remove the fixative and residual chromogens. After

rehydration to PBT in two steps, the embryos were treated with weak Curis solution (1% H_2O_2,

0.5x SSC and 5% formamide) at 4ºC in the dark overnight. Finally, the embryos were washed

twice in PBS before imaging them in PBS on a thick agarose dish by light microscopy.

Antibody generation. Affinity-purified rabbit polyclonal antibodies were generated by sdix

(Delaware, USA) against different amino acid (aa) sequences of *X. tropicalis* Tbx6 avoiding

its highly conserved DNA binding domain (aa 95-280) among T-box proteins: aa 1-94 (#4596),

aa 304-403 (#5061) and aa 418-517 (#5107). Antibody #5107 did not work (data not shown).

Antibodies #4596 and #5061 were suitable for immunoprecipitation and Western blotting to

detect endogenous Tbx6 in both *X. tropicalis* and *X. laevis* embryos (Extended Data Fig. 2f-

h). Antibody #4596 complied with ENCODE guidelines[41] and thus was suitable for ChIP

(Extended Data Fig. 2i).

Immunoprecipitation (IP). For each IP, 40 de-jellied embryos were collected at the

developmental stage of interest. They were homogenised in 400 µl ice-cold IP buffer (50 mM

Tris-HCl pH 7.5, 150 mM NaCl, 5 mM $MgCl_2$, 1 mM EDTA and 0.25 % Igepal CA-630)

supplemented with an EDTA-free protease inhibitor cocktail (Roche). The extract was kept on

ice for 5 min before adding 400 µl 1,1,2-trichloro-1,2,2-trifluoroethane (Sigma). The extract

was mixed with Freon by vigorously inverting the tube and cleared by centrifugation at 16,000

g (4ºC) for 5 min. An aliquot of 20 µl supernatant was saved as input. The remaining

supernatant was incubated with 0.5 µl antibody (anti-Sox3[42], anti-VegT[9] or anti-Tbx6; all rabbit

polyclonal antibodies) at 4ºC for 1 h. Subsequently, each embryonic extract was

supplemented with 20 µl washed magnetic beads conjugated to sheep anti-rabbit IgG

(Dynabeads M280, Invitrogen) and kept on a tube rotator at 4ºC for another 2 h. Next, the

beads were washed three times with IP buffer. The immunoprecipitate was eluted off the

beads with 20 µl 1x SDS loading buffer (50 mM Tris-HCl pH 6.8, 2% SDS, 6% glycerol and

5% β-mercaptoethanol). Western blotting was carried out with TrueBlot (Rockland

Immunochemicals) and normal horseradish peroxidase (HRP)-conjugated secondary

antibodies for IP and input samples, respectively.

Western blotting. Protein samples were denatured in 1x SDS loading buffer (final

concentration) at 70ºC for 5 min. Denatured samples were run alongside a standard protein

ladder into pre-cast gradient gels (Any kD Mini-PROTEAN TGX, BioRad) at constant 200 V

for 1 h. Size separated proteins were immediately transferred onto hydrophobic Immobilon-P

PVDF transfer membranes (Millipore) at constant 100 V for 30 min using standard protein

electrophoresis equipment. The membranes were blocked with 5% milk powder in PBS/0.1%

Tween-20 (MPBTw) at room temperature for 30 min. Next, the membranes were incubated at

room temperature for 1 h with the primary antibodies diluted in MPBTw: 1:5,000 of mouse

monoclonal anti-α-tubulin (Sigma, T5168), 1:2,000 rabbit polyclonal anti-Sox3[42], 1:2,000

rabbit polyclonal anti-VegT[9] or 1:2,000 rabbit polyclonal anti-Tbx6 (#5061). The membranes

were washed three times with PBST for 10 min before applying 1:1,000 TrueBlot HRP-

conjugated anti-rabbit IgG (Rockland Immunochemicals, 18-8816-31) or 1:2,000 normal goat

HRP-conjugated anti-mouse IgG (Thermo Fisher Scientific, 31430) in MPBTw at room

temperature. After 1 h, the membranes were washed three times in PBTw for 10 min and once

in PBS for 5 min. Finally, the membranes were treated with UptiLight US HRP WB reagent

(interchim) for 1 min in the dark. HRP-linked chemiluminescence was detected on the

ChemiDoc XRS+ system (BioRad).

Whole-mount immunohistochemistry (WMIHC). Embryos were fixed with MEMFA (0.1 M

MOPS, 2 mM EGTA, 1 mM $MgSO_4$ and 3.7% formaldehyde) in capped 5-ml glass vials at

room temperature for 2 h, dehydrated and stored in absolute ethanol at -20ºC for ≥24 h. For

sagittal sections, fixation was stopped after 30 min. Embryos were placed on a piece of

flattened Bostik Blu-Tack in a large Petri dish filled with PBS to bisect them with a scalpel.

Following a three-step rehydration (50%, 75% and 100% PBS) the embryos were bleached

with weak Curis solution (1% H_2O_2, 0.5x SSC and 5% formamide) on a light box at room

temperature for 2 h or in the dark at 4ºC overnight. Bleached embryos were washed three

times with PBS/0.3% Triton X-100 (PBT) and pre-incubated in blocking solution (PBS, 20%

goat or donkey serum, 2% BSA and 0.1% Triton X-100) at 4ºC for 6 h. The embryos were

then transferred to 2-ml round-bottom microcentrifuge tubes before discarding all remaining

blocking solution. The embryos were incubated at 4ºC for 1-3 days in 50 µl blocking solution

containing the primary antibody at the following dilutions: 1:1,000 rabbit polyclonal anti-Sox3[42],

1:500 rabbit polyclonal anti phospho-Smad1/5/9 (Cell Signalling, #9511) or 1:500 goat

polyclonal Smad2/3 (R&D Systems, AF3797). Afterwards, the embryos were transferred back to capped 5-ml glass vials, washed three times with RIPA buffer (50 mM HEPES pH 7.5, 0.5 M LiCl, 1 mM EDTA, 1% Igepal CA-630 and 0.7% sodium deoxycholate) for 1 h and once with PBT for 5 min. Next, the embryos were incubated with the secondary HRP-conjugated antibody diluted in blocking solution at 4ºC overnight: 1:400 goat anti-rabbit IgG-HRP (ThermoFisher Scientific, G-21234) or 1:200 donkey anti-goat IgG-HRP (Santa Cruz Biotechnology, sc-2020). The washes of the previous day were repeated before pre-incubating the embryos in 400 µl inactive 3,3'-diaminobenzidine tetrahydrochloride with cobalt (DAB with metal enhancer; Sigma). After 1-2 min, the inactive solution was replaced with the H_2O_2-activated DAB solution. The HRP reaction was stopped after 40 sec by washing the embryos several times with PBT. In Extended Data Fig. 1g, embryos were dehydrated in three steps to absolute methanol and cleared with Murray's clear (2:1 benzyl benzoate/benzyl alcohol) on a glass depression slide.

Deep sequencing and quality filter. All deep sequencing libraries were quality controlled: The DNA yield and fragment size distribution were determined by fluorometry and Chip-based capillary or polyacrylamide gel electrophoresis, respectively. Libraries were sequenced on the Illumina platforms GAIIx or HiSeq by the Advanced Sequencing Facility of the Francis Crick Institute to produce single or paired-end reads of at least 40 bases. Next-generation capture-C libraries were sequenced on MiSeq with a read length of 150 bases. Sequencing samples and read alignment results are summarised in Supplementary Table 3. The metrics of paired-end alignments such as insert size mean and standard deviation were determined by Picard (CollectInsertSizeMetrics) from the Broad Institute (USA).

Chromatin immunoprecipitation (ChIP). ChIP was carried out as detailed previously[43]. Briefly, de-jellied *X. tropicalis* embryos were treated with 1% formaldehyde (Sigma, F8775) in 1% MMR at room temperature for 15-45 min to cross-link chromatin proteins to nearby genomic DNA. Duration of fixation was determined empirically and depended mainly on the developmental stage and antibody epitopes[43]. Fixation was terminated by rinsing embryos three times with ice-cold 1% MMR. Where required, post-fixation embryos were dissected to select specific anatomical regions in ice-cold 1% MMR. Fixed embryos were homogenised in CEWB1 (10 mM Tris-HCl pH 8.0, 150 mM NaCl, 1 mM EDTA, 1% Igepal CA-630, 0.25% sodium deoxycholate and 0.1% SDS) supplemented with 0.5 mM DTT, protease inhibitors and, if using phospho-specific antibodies, phosphatase blockers (0.5 mM orthovanadate and 2.5 mM NaF). To solubilise yolk platelets and separate them from the nuclei, the homogenate was left on ice for 5 min and then centrifuged at 1,000 g (4ºC) for 5 min. Homogenisation and centrifugation was repeated once before resuspending the nuclei containing pellet in 1-3 ml

CEWB1. Nuclear chromatin was solubilised and fragmented by isothermal focused or

microtip-mediated sonication. The solution of fragmented chromatin was cleared by

centrifuging at 16,000 g (4ºC) for 5 min. Where required, ~1% of the cleared chromatin extract

was set aside for the input sample (negative control for ChIP). ChIP-grade antibodies were

used to recognise specific chromatin features and to enrich these by coupling the antibody-

chromatin complex to protein G magnetic beads and extensive washing. These steps were

carried out at 4ºC. The beads were washed twice in CEWB1, twice in WB2 (10 mM Tris-HCl

pH 8.0, 500 mM NaCl, 1 mM EDTA, 1% Igepal CA-630, 0.25% sodium deoxycholate and 0.1%

SDS), twice in WB3 (10 mM Tris-HCl pH 8.0, 250 mM LiCl, 1 mM EDTA, 1% Igepal CA-630

and 1% sodium deoxycholate) and once in TEN (10 mM Tris-HCl pH 8.0, 150 mM NaCl and

1 mM EDTA). ChIP was eluted off the beads twice with 100 µl SDS elution buffer (50 mM Tris-

HCl pH 8.0, 1 mM EDTA and 1% SDS) at 65ºC. ChIP eluates were pooled before reversing

DNA-protein cross-links. Input (filled up to 200 µl with SDS elution buffer) and ChIP samples

were supplemented with 10 µl 5 M NaCl and incubated at 65ºC for 6-16 h. Samples were

treated with proteinase K and RNase to remove any proteins and RNA from the co-

immunoprecipitated DNA fragments. The DNA was purified with phenol:chloroform:isoamyl

alcohol (25:24:1, pH 7.9) using 1.5-ml Phase Lock Gel Heavy (PLGH) microcentrifuge tubes

for phase separation and precipitated with 1/70 volume of 5 M NaCl, 2 volumes of absolute

ethanol and 15 µg GlycoBlue (Ambion). After centrifugation, the DNA pellet was air-dried and

dissolved in 11 µl elution buffer (10 mM Tris-HCl pH 8.5). The DNA concentration was

determined on a fluorometer using high-sensitivity reagents for double-stranded DNA (10

pg/µl to 100 ng/µl).

The following antibodies were used for ChIP: rabbit polyclonal anti-β-catenin (Santa-Cruz, H-

102), rabbit polyclonal anti-H3K4me1 (Abcam, ab8895), rabbit polyclonal anti-HA (Abcam,

ab9110), mouse monoclonal anti-RNAPII (Covance, MMS-126R), rabbit polyclonal anti-

phospho-Smad1/5/9 (Cell Signalling, #9511), goat polyclonal anti-Smad2/3 (R&D Systems,

AF3797), rabbit polyclonal anti-Sox3[42], rabbit polyclonal anti-Tbx6 (#4596) and rabbit

polyclonal anti-VegT[9].

Quantitative PCR (qPCR). DNA was amplified in technical duplicates with SYBR Green I

master mix (Roche) on a Light Cycler 480 II (Roche) cycling 55-times between 94ºC, 60ºC

and 72ºC with each temperature step running for 10 sec and switching at +4.8ºC/sec and -

2.5ºC/sec. At the end qPCR reactions were heated from 65ºC to 97ºC with a gradual increase

of 0.11ºC/sec (melting curve) to ensure only fluorescence was collected from one specific

amplicon. ChIP-qPCR results were based on absolute quantification using an 8-point standard

curve of three-fold dilutions of ~1% input DNA (Extended Data Fig. 2b-e,i). RT-qPCR results

were normalised to the housekeeping gene *odc1* (Fig. 3e; Extended Data Fig. 5d and 7h) or

rpl8 (Extended Data Fig. 6d), and scaled relative to control embryos using the $2^{-\Delta\Delta Ct}$ method[44].

The threshold cycle (C_t) was derived from the maximum acceleration of SYBR fluorescence

(second derivative maximum method).

ChIP-Seq library preparation. 0.5 to 10 ng ChIP DNA or 5 ng input DNA were used to

prepare single (only Tbx6 ChIP-Seq) or indexed paired-end libraries as previously

described[43,45,46].

Post-sequencing analysis of ChIP-Seq. Single reads of maximal 50 bases were processed

using trim_galore v0.4.2 (Babraham Institute, UK) to trim off low-quality bases (default Phred

score of 20, i.e. error probability was 0.01) and adapter contamination from the 3' end.

Processed reads were aligned to the *X. tropicalis* genome assembly v7.1 (and v9.1 for some

ChIP-Seq data) running Bowtie2 v2.2.9[47] with default settings (Supplementary Table 3).

Alignments were converted to the HOMER's tag density format[48] with redundant reads being

removed (makeTagDirectory -single -tbp 1 -unique -mapq 10 -fragLength 175 -totalReads all).

Only uniquely aligned reads (i.e. MAPQ ≥10) were processed. We pooled all input alignments

from various developmental stages. This created a comprehensive mappability profile that

covered ~400 million unique base pair positions. All chromatin profiles were position-adjusted

and normalised to the effective total of 1 million aligned reads including multimappers (counts

per million aligned reads, CPM). Agreeing biological replicates according to ENCODE

guidelines[41] were subsequently merged. For stage 10^+ β-catenin and Smad2 ChIP-Seq,

external datasets[20,49,50] were used as biological replicates. HOMER's peak caller was used to

identify transcription factor binding sites by virtue of ChIP-mediated read enrichments

(hereafter called peaks): findpeaks -style factor -minDist 175 -fragLength 175 -

inputFragLength 175 -fdr 0.001 -gsize 1.435e9 -F 3 -L 1 -C 0.97. This means that both ChIP

and input alignments were extended 3' to 175 bp for the detection of significant (0.1% FDR)

peaks being separated by ≥175 bp. The effective size of the *X. tropicalis* genome assembly

v7.1 was set to 1.435 billion bp, an estimate obtained from the mappability profile. These

peaks showed equal or higher tag density than the surrounding 10 kb, ≥3-fold more tags than

the input and ≥0.97 unique tag positions relative to the expected number of tags. To detect

focal RNAPII recruitment to putative CRMs and avoid calling peaks within broad regions of

RNAPII reflecting transcript elongation, the threshold of focal ratio and local enrichment within

10 kb was elevated to 0.6 and 3 (-L 3), respectively. To further eliminate any false positive

peaks, we removed any peaks with <0.5 (TFs including signal mediators) or <1 (RNAPII) CPM

and those falling into blacklisted regions showing equivocal mappability due to genome

assembly errors, gaps or simple/tandem repeats. Regions of equivocal mappability were

identified by a two-fold lower (poor) or three-fold higher (excessive) read coverage than the
average detected in 400-bp windows sliding at 200-bp intervals through normalised ChIP input
and DNase-digested naked genomic DNA. All identified regions ≤800 bp apart were
subsequently merged. Gap coordinates were obtained from the Francis Crick UCSC genome
browser (http://genomes.crick.ac.uk). Simple repeats were masked with RepeatMasker v4.0.6
using the crossmatch search engine v1.090518 and the following settings: RepeatMasker -
species "xenopus silurana tropicalis" -s -xsmall. Tandem repeats were masked with Kent's
trfBig wrapper script of the Tandem Repeat Finder v4.09[51] using the following settings: weight
for match, 2; weight for mismatch, 7; delta, 7; matching probability, 80; indel probability, 10;
minimal alignment score, 50; maximum period size, 2,000; and longest tandem repeat array
(-l), 2 [million bp]. Multi-genome sequence conservation tracks (phastCons and phyloP) for *X.*
tropicalis genome assembly v9.1 were obtained from Xenbase[52]. The following 11 vertebrate
species were used to evaluate sequence similarity: *X. tropicalis*, *X. laevis*, *Nanorana parkeri*
(High Himalaya Frog), *Fugu rubripes* (Japanese Pufferfish), *Chrysemys picta* (Painted Turtle),
Gallus gallus (Chicken), *Anolis carolinensis* (Green Anole lizard), *Monodelphis domestica*
(Gray Short-tailed Opossum), *Canis lupus familiaris* (dog), mouse and human. CRMs with a
phastCons average ≥0.4 were considered 'conserved'.

Detecting zygotic genes by means of RNAPII chromatin profiles. RNAPII and input tag
densities were calculated across the gene body in 10 bins of equal size. Gene annotations
v7.1 were altered based on a few known zygotic isoforms[37] and some corrections obtained
from assembling total and poly(A) RNA[37] from stage 6 to stage 12.5 *de novo* as outlined
elsewhere[53]. A few genes had previously been annotated as gene clusters due to assembly
uncertainties. We reduced the annotation of polycistronic *mir427* to the minus arm
(scaffold_3b:3516900-3523400) and only monitored *nodal3.5* and *nodal5.3* within their
respective gene clusters. Gene bodies with <40% mappability were removed. Here, the
threshold of mappability per bin was set at 10% of the input read density averaged across all
gene bodies in use. Subsequently, enrichment values were only obtained for all mappable
bins by dividing read densities of RNAPII and input. Further, we restricted the analysis to
genes for which ≥3 transcripts per million (TPM) could be detected on average over 3
consecutive time points (i.e. over the developmental time of 1 h) of a high-resolution profile of
total RNA[37] from fertilisation to after gastrulation (stage 13). Genes were considered activated
at a certain developmental stage if ≥0.1 TPM were detected over 3 consecutive time points
within a 2 h window of the developmental stage in question and either of the following RNAPII
enrichments covering ≥80% of the mappable gene body were reached: (1) 2.6-fold, (2) 1.8-
fold and 1.4-fold at the next or previous stage, (3) 1.4-fold and 1.8-fold at the next or previous
stage, or (4) 1.4-fold over 3 consecutive stages.

Poly(A) RNA-Seq. Libraries were made from ~1 µg total RNA by following the low sample protocol of the TruSeq RNA sample preparation guide version 2 with a few modifications. First, 1 µl cDNA purified after second strand synthesis was quantified on a fluorometer using high-sensitivity reagents for double-stranded DNA (10 pg/µl to 100 ng/µl). By this stage, the yield was ~10 ng. Second, the numbers of PCR cycles were adjusted to the detected yield of cDNA to avoid products of over-amplification such as chimera fragments: 7 (~20 ng), 8 (~10 ng), 9 (~5 ng) and 12 (~1 ng).

RNA-Seq read alignment. Paired-end reads were aligned to the *X. tropicalis* genome assembly v7.1 using STAR v2.5.3a[54] with default settings and a revised version of gene models v7.2[37] to improve mapping accuracy across splice junctions. The alignments were sorted by read name using the sort function of samtools v1.3.1. Exon and intron counts (-t 'exon;intron') were extracted from unstranded (-s 0) alignment files using VERSE v0.1.5[55] in featureCounts (default) mode (-z 0). Intron coordinates were adjusted to exclude any overlap with exon annotation. For visualisation, genomic BAM files of biological replicates were merged using samtools v1.3.1[56] and converted to the bigWig format. These genome tracks were normalised to the wigsum of 1 billion excluding any reads with mapping quality <10 using the python script bam2wig.py from RSeQC v2.6.4[57].

Differential gene expression analysis. Differential expression analysis was performed with both raw exon and intron counts excluding those belonging to ribosomal and mitochondrial RNA using the Bioconductor/R package DESeq2 v1.14.1[58]. In an effort to find genes with consistent fold changes over time, p-values were generated according to a likelihood ratio test reflecting the probability of rejecting the reduced (~ developmental stage) over the full (~ developmental stage + condition) model. Resulting p-values were adjusted to obtain false discovery rates (FDR) according to the Benjamini-Hochburg procedure with thresholds on Cook's distances and independent filtering being switched off. Equally, regional expression datasets[24] without time series were subjected to likelihood ratio tests with reduced (~1) and full (~ condition) models for statistical analysis. Fold changes of intronic and exonic transcript levels were calculated for each developmental stage and condition using the mean of DESeq2-normalised read counts from biological replicates. Both intronic and exonic datasets were filtered for ≥10 DESeq2-normalised read counts detected at least at one developmental stage in all uninjected or DMSO-treated samples. Gene-specific fold changes were removed at developmental stages that yielded <10 normalised read counts in corresponding control samples. Next, the means of intronic and exonic fold changes were calculated across developmental stages. The whole dataset was cut down to 3,318 genes for which at least 50%

reductions (FDR <10%) in exonic (default) or intronic counts could be detected in α-amanitin-

injected embryos. Regional expression was based on exonic read counts by default unless

the intronic fold changes were significantly (FDR <10%) larger than the exonic fold changes

(Supplementary Table 2).

Analysing ribosome footprinting and mass-spectrometry data. The ribosome footprinting

reads[6] were trimmed 5' by 8 bases and 3' by as many bases overlapping with the adapter

sequence 5'-TCGTATGCCGTCTTCTGCTTG-3' from the 5' end. All trimmed reads between

27 to 32 bases in length were aligned first to ribosomal RNA as listed in the SILVA rRNA

database[59] using Bowtie v1.0.1[60] with the following parameters: --seedlen 25 (seed length) --

seedmms 1 (number of mismatches allowed in the seed) --un (unaligned reads were

reported). All non-aligned (rRNA-depleted) reads were mapped to the gene model 6.0 of *X.*

laevis using Tophat2 v2.0.10[61] --no-novel-juncs (spliced reads must match splice junctions of

gene model 6.0) --no-novel-indels (indel detetion disabled) --segment-length 15 (minimal

length of read fragment to be aligned) –GTF v6.0.gff3 --prefilter-multihits (reads aligned first

to whole genome to exclude reads aligning >10 times) --max-multihits 10. Alignment files were

converted to the HOMER[48] tag density format before retrieving reads for each CDS per

kilobase and one million mapped reads (rpkm) using HOMER's perl script analyzeRNA.pl.

This read count table was merged with a published list of estimated protein concentrations

(nM) in the *X. laevis* egg using mass-spectrometry[5] (Supplementary Table 1). Chromatin-

associated proteins regulating RNAPII-mediated transcription were filtered based on human

(version 2.0, 09/2014) and Xenbase-released gene ontology (GO) associations. The genes

associated with any of the following GO terms were verified with UniProt[62] whether chromatin

binding is supported by functional evidence: chromatin (GO:0000785), DNA binding

(GO:0003677), DNA-templated regulation of transcription (GO:0006355), sequence-specific

DNA binding and TF activity (GO:0003700) and DNA replication (GO:0006260). We separated

specific TFs from all other chromatin binding proteins to form three categories: (1) sequence-

specific DNA binding factors, (2) other RNAPII-regulating factors and (3) remaining genes,

whereby genes associated with DNA repair (GO:0006281) and RNAPIII (GO:0006383) were

moved to the third category.

SPRI-based enrichment of small DNase-digested fragments (DNase-Seq). In our hands,

the high yolk content in vegetal blastomeres made it impossible to employ ATAC-Seq on whole

Xenopus embryos before the onset of gastrulation. Therefore, DNase-Seq was adapted to

early *Xenopus* embryos with a novel approach to select small DNase-digested DNA

fragments. Ultracentrifugation- or gel electrophoresis-mediated size selection[63,64] was

replaced by two rounds of high molecular DNA depletion by solid phase reverse immobilisation

(SPRI). Wide-bore pipette tips were used for the resuspensions and the transfers of samples

from the second homogenisation step until after SPRI to minimise fragmentation of high

molecular DNA. Approximately 250 de-jellied mid-blastula embryos were collected in 2-ml

round-bottom microcentrifuge tubes and homogenised in 2 ml ice-cold LB-DNase buffer (15

mM Tris-HCl pH 8.0, 15 mM NaCl, 60 mM KCl, 1 mM EDTA, 0.5 mM EGTA and 0.5 mM

spermidine) supplemented with 0.05% Igepal CA-630. This lysate was left on ice for 3 min

before centrifuging at 1,000 g (4ºC) for 2 min. The supernatant was discarded without

disrupting the pellet. The pellet was gently resuspended in 2 ml ice-cold LB-DNase buffer

(without Igepal CA-630) before centrifuging again at 1,000 g (4ºC) for 2 min. After discarding

the supernatant, the pellet was resuspended at room temperature in 600 µl LB-DNase buffer

(without Igepal CA-630) with 6 mM $CaCl_2$. The sample was distributed equally to two 1.5-ml

microcentrifuge tubes, one for probing chromatin accessibility with DNase and the other one

for creating a reference profile of DNase-digested naked genomic DNA. Approximately 0.1 U

recombinant DNase I (Roche) was added to one aliquot. Both samples were incubated at

37ºC for 8 min before adding 300 µl STOP buffer (0.1% SDS, 100 mM NaCl, 100 mM EDTA

and 50 mM Tris-HCl pH 8.0) including 80 µg RNase A (Thermo Fisher Scientific), 333 nM

spermine and 1 µM spermidine. The tubes were inverted gently to mix samples before

incubating them at 55ºC for 15 min. Next, 200 µg proteinase K (Ambion) were added and the

tubes were inverted gently to mix the samples again. After 2 h at 55ºC, the digests were

transferred to pre-spun 1.5-ml Phase Lock Gel Heavy (PLGH) microcentrifuge tubes. 600 µl

phenol:chloroform:isoamylalcohol (25:24:1, pH 7.9) were added to the digests. The tubes

were shaken gently and then centrifuged at 1,500 g (room temperature) for 4 min. The top

phase was transferred to a fresh 2-ml microcentrifuge tube and mixed with 60 µl 3 M sodium

acetate (pH 5.2) and 1.2 ml of absolute ethanol to precipitate the genomic DNA. The

precipitation was stored at -20ºC overnight at least and centrifuged at 16,000 g (4ºC) for 30

min. The supernatant was discarded and the DNA pellet washed by adding 500 µl ice-cold

80% ethanol and centrifuging at 16,000 g (4ºC) for 3 min. The ethanol was discarded and the

DNA pellet was air-dried at room temperature for 10 min. After that the DNA pellet was left on

ice to dissolve in 27 µl elution buffer (10 mM Tris-HCl pH 8.5) for 20 min. To remove any

residual RNA, 10 µg RNase A were added to the DNA. 1 µl was used to determine the DNA

fragment size distribution by gel electrophoresis. On a 0.6% TAE agarose gel, a smear of very

high molecular DNA was visible as expected from previous DNase experiments[64,65]. Genomic

DNA to the amount of 20 untreated mid-blastula embryos was digested with 0.3 U DNase I at

37ºC for 5 min following the same steps as described above, which generated a low molecular

smear of DNA fragments. Next, 70 µl SPRI beads (Agencourt AMPure XP, Beckman Coulter)

per DNase sample were equilibrated to room temperature for 10 min. 22.5 µl SPRI beads

(0.9x of the volume) were added to the DNA sample without pipetting up and down achieving

a final polyethylene glycol (PEG) concentration of ~9.5%. After 3 min, by which time high-
molecular DNA causes beads to coalesce, the tubes were clipped into a magnetic stand for
microcentrifuge tubes. After 3 min or until the beads were separated from the supernatant, the
latter was transferred to a 96-well microplate (350-µl round wells with V-shaped bases). 47.5
µl elution buffer and 43 µl SPRI beads were added sequentially and mixed gently by slowly
pipetting up and down. After 3 min, the plate was transferred to a magnetic stand for 96-well
plates. Once the beads have completely separated from the suspension, the supernatant was
transferred to a pre-spun PLGH tube. 162 µl elution buffer and 300 µl
phenol:chloroform:isoamylalcohol (25:24:1, pH 7.9) were added. The tubes were shaken
gently and then centrifuged at 1,500 g (room temperature) for 4 min. The top phase was
transferred to a 1.5-ml low-retention microcentrifuge tube and mixed with 30 µl 3 M sodium
acetate (pH 5.2) and 900 µl absolute ethanol to precipitate the DNA as outlined above. The
DNA pellet was dissolved in 12 µl elution buffer. The concentrations of the DNA samples were
determined on a fluorometer using high sensitivity reagents for dsDNA (10 pg/µl to 100 ng/µl).
Libraries were generated as previously outlined[43] except that all cleaning steps were executed
with 0.2x more SPRI beads.

Post-sequencing analysis of DNase-Seq. Single and paired-end reads of maximal 50 bases
were processed using trim_galore v0.4.2 from the Babraham Institute (UK) to trim off low-
quality bases (default Phred score of 20, i.e. error probability was 0.01) and adapter
contamination from the 3' end. Processed reads were aligned to the *X. tropicalis* genome v7.1
and v9.1 using Bowtie2 v2.2.9[47] with default settings apart from -X (fragment length), which
was reduced to 250 bp for paired-end reads. Alignments were sorted by genomic coordinates
and only those with a quality score of ≥10 were retained using samtools v1.3.1[56]. Duplicates
were removed using Picard (MarkDuplicates) from the Broad Institute (USA). Paired-end
alignments were dissociated using hex flags (-f 0x40 or 0x80) of samtools view. Single
alignments were converted to HOMER's tag density format[48] (makeTagDirectory -single -
unique -fragLength 100 -totalReads all). DNase hypersensitive sites (DHSs) were identified
using the following or otherwise default parameters of HOMER's peak calling: findpeaks -style
factor -minDist 100 -fragLength 100 -inputFragLength 100 -fdr 0.001 -gsize 1.435e9 -F 3 -L 1
-C 0.97. This means that alignments of DNase-digested chromatin fragments (or naked
genomic DNA fragments considered here as 'input') were extended 3' to 100 bp from the
DNase cleavage site to detect significant (0.1% FDR) DNase-mediated read enrichments
(hereafter called peaks) being separated by ≥100 bp. The effective size of the *X. tropicalis*
genome assembly v7.1 was set to 1.435 billion bp, an estimate obtained from the mappability
of ChIP input reads. These peaks showed equal or higher tag density than the surrounding 10
kb, at least three-fold more tags than the input and ≥0.97 unique tag positions relative to the

expected number of tags. Peaks falling into blacklisted regions (see Post-sequencing analysis

of ChIP-Seq) were removed. DNase-probed chromatin accessibility profiles were position-

adjusted and normalised to the effective total of 1 million aligned reads including

multimappers.

Next-generation capture-C. About 500 mid-blastula embryos per condition were collected

in 9-ml capped glass vials and briefly washed once with 1% MMR. The embryos were then

fixed with 1% formaldehyde (Sigma, F8775) in 9 ml 1% MMR at room temperature for 40 min.

The fixation reaction was terminated by briefly rinsing the embryos three times with ice-cold

1% MMR. The embryos were aliquoted into round-bottom 2-ml microcentrifuge tubes in

batches of ~250 embryos (filling the volume of ~250 μl water). The supernatant was removed

before equilibrating embryos in 250 μl ice-cold HEG buffer (50 mM HEPES pH 7.5, 1 mM

EDTA and 20% glycerol). Once the embryos settled to the bottom of the tube as much liquid

as possible was discarded. The aliquots were snap-frozen in liquid nitrogen and stored at -

80ºC.

For each 3C experiment, 10 ml chromatin extraction buffer CEB-3C (10 mM Tris-HCl pH 8.0,

150 mM NaCl, 1 mM EDTA, 1% Igepal CA-630, 0.25% sodium deoxycholate and 0.2% SDS)

were supplemented with a mini protease inhibitor tablet (Roche) and 0.05 mM DTT (CEB-

3C*). Both aliquots of ~250 embryos of each condition were homogenised in 2 ml ice-cold

CEB-3C*. The tubes were kept on ice for 5 min before centrifuging at 1,000 g (4ºC) for 2 min.

The supernatants were discarded and the pellet resuspended in 0.5 ml ice-cold CEB-3C*. The

resuspensions of each condition were pooled and then equally divided to two 50-ml conical

tubes and each filled up with CEB-3C (without protease inhibitors and DTT) to 50 ml. The

embryonic extracts were incubated at 37ºC for 1 h in a hybridisation oven with the tubes

rotating inside a hybridisation bottle. The tubes were then centrifuged at 1,000 g (room

temperature) for 5 min. The supernatants were discarded and the pellets resuspended in 50

ml double distilled water. The tubes were centrifuged again at 1,000 g (room temperature) for

5 min. The supernatants were discarded and each pellet was resuspended in 300 μl double

distilled water. The resuspensions containing cross-linked nuclei of each condition were

pooled in a 1.5-ml microcentrifuge tube. The samples were then digested with 400 U *Dpn*II

(NEB) in a total volume of 800 μl containing 1x *Dpn*II reaction buffer. The digest was incubated

overnight in a thermomixer set to 37ºC and 1,400 rpm. Of note, any residual yolk platelets in

the solution did not interfere with the restriction digest. The next day, the digest was

supplemented with 200 U *Dpn*II and incubated at 37ºC and 1,400 rpm for another 6-8 h. After

that, *Dpn*II was heat-inactivated at 65ºC for 20 min.

To verify the degree of chromatin digestion, an aliquot of 50 µl from each condition was
transferred into separate 1.5-ml microcentrifuge tubes and processed as follows: First, any
remaining RNA was degraded by incubating the aliquot with 20 µg RNase A (Thermo Fisher
Scientific) at 37ºC for 30 min. Second, proteins were digested by incubating the aliquot mixed
with 50 µl SDS elution buffer (50 mM Tris pH 8.0, 1 mM EDTA and 1% SDS) and 20 µg
proteinase K (Ambion) at 65ºC in a hybridisation oven overnight. Third, the DNA was purified
and ethanol precipitated: 300 µl TE pH 8.0 were added to the digest before transferring it to
pre-spun 1.5-ml Phase Lock Gel Heavy (PLGH) microcentrifuge tubes. The diluted digests
were mixed vigorously with 400 µl (1 volume) phenol:chloroform:isoamylalcohol (25:24:1, pH
7.9) before centrifuging at 16,000 g (room temperature) for 1 min. The top (aqueous) phases
were transferred into 1.5-ml microcentrifuge tubes and mixed with 40 µl sodium acetate (pH
5.2) and 800 µl absolute ethanol and 15 µg GlycoBlue (Ambion). The DNA precipitations were
kept at -80ºC for 30 min before centrifuging at 16,000 g (4ºC) for 30 min. The supernatants
were discarded without disturbing the DNA pellets which were subsequently washed once
with 400 µl ice-cold 80% ethanol. The tubes were centrifuged at 16,000 g (4ºC) for 30 min
before discarding the supernatants. The DNA pellets were air-dried for 10 min and
resuspended in 10 µl elution buffer (10 mM Tris-HCl pH 8.5). The DNA concentrations were
determined with a fluorometer and the DNA fragment distributions were visualised on a 0.6%
TAE agarose gel. The digestion of cross-linked chromatin with *Dpn*II caused extensive
fragmentation of the genomic DNA such that no high molecular DNA fragment bands (>10 kb)
were visible.

The chromatin digests were processed for proximity ligation by adding 240 U T4 DNA ligase
(Thermo Fisher Scientific) in a total volume of 1.2 ml containing 1x T4 DNA ligase buffer
including 5 mM ATP. The ligation reactions were incubated at 16ºC and 1,400 rpm for ≥16 h
before centrifuging at 16,000 g (4ºC) for 1 min. The supernatants were discarded. 200 µl
elution buffer and 40 µg RNase A were added to the pellets to degrade any residual RNA at
37ºC and 1,400 rpm for 30 min. Proteins were digested by adding 200 µl SDS elution buffer
and 160 µg proteinase K and incubating at 65ºC in a hybridisation oven overnight. The digest
was transferred to pre-spun PLGH tubes and processed as described in the previous
paragraph to purify and precipitate ligated genomic DNA. The air-dried DNA pellet was
dissolved in 20 µl elution buffer and quality controlled as above. Proximity ligation of *Dpn*II-
digested chromatin fragments massively increased the size of DNA fragments to the range of
high molecular weight (>10 kb) representing the chromatin conformation capture (3C) library.
The concentration was measured on a fluorometer with broad range concentration (5-500
ng/µl) reagents.

About 10-15 µg of the 3C libraries were diluted with elution buffer to the total volume of 130 µl and transferred to a designated glass vial (microTUBE) for isothermal focused sonication (Covaris). The following settings of the focused ultrasonicator Covaris S220 were used to shear the libraries to an average DNA fragment length of ~200 bp: duty cycle, 10%; intensity, 5; cycles per burst, 200; duration, 60 sec in frequency sweeping mode. Sonication was run with 6 cycles in batch mode. The degree of DNA fragmentation was verified by gel electrophoresis and the DNA concentration was measured on a fluorometer with broad range concentration reagents.

Approximately 2-4 µg of the DNA fragments were converted into next-generation paired-end libraries using the enzymes of the KAPA Hyper Prep Kit (Kapa Biosystems) and TruSeq adaptors (Illumina). The end-repair and A-tailing reactions were set up according to the manufacturer's instructions and incubated at 20ºC for 1 h followed by 30 min at 65ºC. End-repaired and A-tailed DNA fragments were ligated to 150 pmol TruSeq adaptors of index 6 and 12, respectively, at 20ºC for 1 h. The DNA was purified using 0.8x volume of SPRI beads and amplified in a total volume of 100 µl using 50 pmol of each TruSeq PCR primer, KAPA HiFi HotStart ReadyMix and the following PCR conditions: 98ºC for 45 sec followed by 3 cycles (98ºC for 15 sec, 98ºC for 30 sec and 72ºC for 30 sec) and a final elongation step of 1 min at 72ºC. The lid temperature was set to 105ºC. The PCR reactions were cleaned up with 100 µl SPRI beads and eluted in 14 µl elution buffer. The integrity of the library DNA was verified on a Chip-based capillary electrophoresis system, an 8% TBE polyacrylamide gel or a 2% TAE agarose gel. DNA concentration was measured on a fluorometer with broad range concentration reagents. The DNA yield was 2-4 µg.

To capture the genomic regions of interest (viewpoints), 5'-biotinylated oligonucleotides of 120 bases (hereafter called probe; Supplementary Table 4) were designed for each viewpoint as follows: The viewpoints were *Dpn*II fragments of 300-3,300 bp which were overlapping gene promoters or located <1 kb from them. The probe matched the terminal sequence of the same DNA fragment strand including the *Dpn*II restriction site GATC. The probe sequence was examined using BLAT to check whether it is unique and did not partially match any other genomic regions. Furthermore, CENSOR v4.2.29[66] and RepeatMasker v4.0.6[67] programs were run to discard any probes that contained repeats. Because of these design restrictions, six of the selected 30 gene promoters were captured with only one probe. All the probes were purchased as desalted oligomers (4 nmol) from IDT (ultramer technology) and re-constituted in molecular-grade water to 10 µM. The probes were mixed in equimolar quantities and diluted to 2.9 nM such that 4.5 µl contained 13 fmol of oligomers.

The capture was performed with the SeqCap Hybridization and Wash Kit (Roche) and SeqCap
HE-Oligo Kit A (Roche) as follows. Exactly 1 µg of each TruSeq library with index 6 and 12
was mixed with 1 nmol TruSeq universal blocking oligonucleotide and 0.5 nmol blocking
oligonucleotides specific to TruSeq index 6 and 12 in a 1.5-ml low-retention microcentrifuge
tube. Of note, Cot-1 DNA commonly used to mask repetitive DNA in hybridisation reactions
proved to be not required for capturing the genomic DNA of interest. The mixture of library
and blocking oligonucleotides was dried in a vacuum centrifuge before adding 7.5 µl 2x
hybridisation buffer (vial 5) and 3 µl hybridisation component A (vial 6). This mixture was
vortexed for 10 sec, centrifuged at 16,000 g for 10 sec and incubated at 95ºC for 10 min. In
the meantime, 4.5 µl (13 fmol) of the equimolar probe mixture were transferred to a PCR tube
and incubated at 47ºC in a PCR machine with the volume and the lid temperature set to 15 µl
and 57ºC, respectively. Upon denaturation, the libraries and blocking oligonucleotides were
added to the probes without removing either tubes from the heating block or the PCR machine.
The hybridisation mixture was incubated at 47ºC for 64-72 h. The wash buffers were prepared
according to the manufacturer's instructions to make 1x working solutions for one single
capture experiment. The stringent wash buffer and wash buffer I were pre-warmed to 47ºC
and 50 µl M270 streptavidin beads (Invitrogen) were transferred into a 1.5-ml low-retention
microcentrifuge tube to let them equilibrate to room temperature for 10 min. The magnetic
beads were washed twice with 200 µl bead wash buffer. Immediately after the final wash, the
hybridisation mixture was directly added to the beads and vortexed for 10 sec. The sample
was incubated for 45 min in a thermomixer set to 47ºC and 1,100 rpm. The beads were
washed by adding 100 µl pre-warmed wash buffer I and vortexing for 10 sec. The tube was
placed into a magnetic rack. Once the liquid was clear, the supernatant was discarded, and
200 µl pre-warmed stringent wash buffer were added to the beads. In an effort to avoid any
temperature drop, it is important to work quickly according to the manufacturer's instructions.
The beads in stringent wash buffer were incubated in a thermomixer set to 47ºC and 1,100
rpm for 5 min. The wash with pre-warmed stringent wash buffer was repeated once. Next, 200
µl pre-warmed wash buffer I was added to the beads and vortexed at 1,400 rpm (room
temperature) for 2 min. After removing the respective wash buffer, beads were vortexed at
1,400 rpm in 200 µl wash buffer II for 1 min and 200 µl wash buffer III for 30 sec. After the final
the final wash, as much liquid as possible was removed and the beads were resuspended in
40 µl molecular-grade water.
1036
The captured DNA was directly amplified from the beads using KAPA HiFi HotStart master
mix and 25 pmol of each TruSeq PCR primer in two separate 50-µl PCR reactions. The PCR
conditions were the same as outlined above except that 14 cycles were used for amplification.
The PCR reactions were pooled and the DNA was purified using 100 µl SPRI beads. The DNA

was eluted with 11 µl elution buffer. The eluted DNA was subjected to another round of probe-

mediated capture with the hybridisation timespan reduced to 24 h. Furthermore, after washing

the beads, any NG capture-C fragments were amplified using only 10 cycles of PCR.

1044

Post-sequencing analysis of next-generation capture-C reads. The analysis was carried

out in accordance with the original description of the method[68] with some modifications.

Paired-end reads were processed using trim_galore v0.4.2 from the Babraham Institute (UK)

to trim off low-quality bases (default Phred score of 20, i.e. error probability was 0.01) and

adapter contamination from the 3' end. Only complete mate pairs were processed further to

reconstruct single reads from overlapping paired-end sequences using FLASH v1.2.11[69] with

interleaved output settings for non-extended reads (flash --interleaved-output --max-overlap

150). FLASH reads were split in *silico* at *Dpn*II restriction sites using a designated perl script

(https://github.com/telenius/captureC/releases) before aligning them to the *X. tropicalis*

genome assembly 7.1 using Bowtie2 v2.2.9[47]. The alignment was run with default settings and

one thread only to keep the order of the reads. The view function of samtools v1.3.1[56] was

used to retain alignments with a quality score of ≥10. The alignments were analysed further

using a suite of perl scripts (https://github.com/Hughes-Genome-Group/captureC/releases)

modified to process both chromosome and scaffold coordinates. Viewpoint coordinates

included a 1-kb proximity exclusion range. Restriction fragments were classified as capture,

proximity-exclusion or reporter. PCR duplicates were removed. The interaction map was

based on the number of unique paired-end reads per restriction fragment. Windows of 2 kb

incrementing by 200 bp were used to consolidate interactions, which were normalised to

10,000 interactions per viewpoint.

1064

Analysis of genomic profiles, DNA motifs and super-enhancers. Unless otherwise stated,

Spearman correlations (R_s), relative tag densities, PCAs and enrichment of DNA motifs were

determined using a limited number of detected peaks per chromatin feature and

developmental stage: top 2,000 peaks (Fig. 2a; Extended Data Fig. 4a-e), top 5,000 peaks

(Extended Data Fig. 1b) and top 10,000 peaks (Fig. 2c,e,h; Extended Data Fig. 5a,b,f,g). R_s

and PCA were calculated using DiffBind v2.4.8[70] in R (Fig. 1a and 2c,h; Extended Data Fig.

4a-c and 5a). Z-scores of motif enrichments at RNAPII peaks were calculated using ISMARA

v1.2.2[71] (https://ismara.unibas.ch/fcgi/mara) in expert mode (Fig. 1b). Heat maps of relative

tag densities were hierarchically clustered (Ward's method) except for those in Extended Data

Fig. 5b, Fig. 4b and 9d, where genomic coordinates were sorted upon significant changes in

DNA occupancy or chromatin accessibility (see figure legends). In Extended Data Fig. 4d,

cluster leafs were additionally ordered with the Gruvaeus-Wainer's algorithm. The matrices of

normalised tag densities, p-values and motif instances were generated with HOMER as

follows: annotatePeaks.pl -size 2000 (or 200 for Extended Data Fig. 4d) -hist 25 (bin size) -

ghist. In heat maps all genomic profiles were scaled to the means of the highest bin values of

the top 100 (or 1,000; Fig. 4b and Extended Data Fig. 9d) peaks. In addition, to visualise low

RNAPII binding and RNA expression, their top means were divided by 5 or 500, respectively.

Super-enhancers were assembled by stitching together peaks that are ≤25 kb apart:

findPeaks -style super -fragLength 175 -inputFragLength 175 -fdr 0.001 -gsize 1.435e9 -F 3 -

L 1 -localSize 10000 -C 0.97 -superSlope 1 -minDist 25000. Subsequently, super-enhancers

≤5 kb from TSS apart were associated with zygotic genes defined by RNAPII coverage (see

Detecting zygotic genes by means of RNAPII chromatin profiles).

1087

1088 **Analysis of enriched gene ontology (GO) terms.** Over-represented GO terms were found

1089 by applying hypergeometric tests of the Bioconductor/R package GOstats v2.40.0[72] on gene

1090 lists. The gene universe was associated with GO terms by means of BLAST2GO[73] as

1091 previously outlined[46,74].

1092

Differential analysis of chromatin features. The significance of differential chromatin

accessibility between control and mPouV/Sox3 LOF embryos was assessed at DHSs using

the Bioconductor/R package DiffBind v2.4.8[70] and the DESeq2[58] algorithm. The sizes of all

DHSs were fixed to 500 bp. Only the very 5' end of each aligned DNase-Seq read, which

represents the DNase cleavage site, was considered for this analysis. Read counts from

mapping DNase-digested naked genomic DNA were subtracted from chromatin accessibility

readings at each DHS. Any changes to DNase-mediated cleavages at DHSs with FDR ≤10%

were determined as statistically significant (Extended Data Fig. 9b). For visualisation, the

orientations of DHSs were aligned to the direction of transcription of the closest zygotic gene.

Matrices of normalised and raw tag counts were generated at DHSs for various chromatin

features and RNA (Fig. 4b and Extended Data Fig. 9d). Their rows were sorted according to

the statistical significance of differential chromatin accessibility. Raw tag counts were

processed to estimate p-values of chromatin changes across DHSs using DESeq2 v1.14.1[58].

Read count dispersions were locally fitted. The shrinkage of $\log_2$ fold changes, threshold on

Cook's distances and independent filtering were all switched off. For the comparison of NG

capture-C interactions with promoters at affected and unaffected DHSs, the size of the DHSs

was widened to 1.2 kb and filtered for ≥5 reporter fragments (Extended Data Fig. 10c).

Normalised line and bar graph tracks of control and mPouV/Sox3 LOF embryos were

superimposed in the IGV genome browser v2.3.92[75] (Fig. 4c and Extended Data Fig. 10d).

1112

Software for computational analysis and graphical illustration. Post-alignment analysis and graphical illustration were performed using samtools v1.3.1[56], bedtools v2.25.0[76], HOMER v4.8.3[48] and tools in R v3.4.1 / Bioconductor v3.5, Perl v5.18.2 or Python v2.7.12.

Data availability. Raw and processed sequencing data for this study is available at the Gene Expression Omnibus (GSE113186).

External next-generation datasets. The following external datasets were used for this study: 5mC at stage 9, and ChIP for H3K4me3 and H3K27me3 at stage 9 (GSE67974), β-catenin ChIP at stage 10$^+$ (GSE72657), Smad2 ChIP at stage 10$^+$ (GSE30146 and GSE53654), Foxh1 ChIP at stage 8 and 10$^+$ (GSE85273 and GSE53654), total RNA recorded at 30-min intervals from unfertilised eggs to 23.5 hpf (GSE65785), ribosome footprinting from stage 3, 9 and 12$^+$ (GSE52809), and regional expression along the animal-vegetal and dorso-ventral axis at stage 10$^+$ (GSE81458).

Statistical analysis. No statistical method was used for determining sample size. We followed the literature to select the appropriate sample size. The experiments were not randomized. Due to the nature of experiments, the authors were not blinded to allocation during experiments and results analysis. Only viable embryos and embryonic tissues were included for analysis. Due to space constraints, data from the trials of establishing and optimising protocols (ChIP-Seq, DNase-Seq and next-generation capture-C) were excluded. Frequencies of shown morphological phenotypes and WMISH patterns are included in every image. Significances of normally or non-normally distributed data points were calculated using a paired Student's t-test (one-tailed or two-tailed) or paired Wilcoxon test (two-tailed), respectively. The effect size r was estimated from the standard normal deviate of the Wilcoxon p-value (p) as previously described[80], $r_{effect}=Z/sqrt(N)$, where $Z=qnorm(1-p/2)$ is the standardised Z-score and N is the number of observations. Significance of motif enrichments or GO terms are based on hypergeometric tests. The significance of transcriptome-wide differential expression was adjusted for multiple comparisons using the Benjamini-Hochburg procedure.

Code availability. Code for the analyses described in this paper is available from the authors upon request.

References

1. Waddington, C. H. Organisers and Genes. Cambridge University Press (1940).
2. Sagner, A. & Briscoe, J. Morphogen interpretation: concentration, time, competence, and

signaling dynamics. *Wiley Interdiscip Rev Dev Biol* **6**, e271 (2017).
3.   Langley, A. R., Smith, J. C., Stemple, D. L. & Harvey, S. A. New insights into the maternal to
zygotic transition. *Development* **141,** 3834–3841 (2014).
4.   Kimelman, D. Mesoderm induction: from caps to chips. *Nat Rev Genet* **7,** 360–372 (2006).
5.   Wühr, M. *et al.* Deep proteomics of the Xenopus laevis egg using an mRNA-derived reference
database. *Curr Biol* **24,** 1467–1475 (2014).
6.   Subtelny, A. O., Eichhorn, S. W., Chen, G. R., Sive, H. & Bartel, D. P. Poly(A)-tail profiling
reveals an embryonic switch in translational control. *Nature* **508,** 66–71 (2014).
7.   Zhang, C., Basta, T., Fawcett, S. R. & Klymkowsky, M. W. SOX7 is an immediate-early target
of VegT and regulates Nodal-related gene expression in Xenopus. *Dev Biol* **278,** 526–541
(2005).
8.   Larabell, C. A. *et al.* Establishment of the dorso-ventral axis in Xenopus embryos is presaged
by early asymmetries in beta-catenin that are modulated by the Wnt signaling pathway. *J Cell*
Biol **136,** 1123–1136 (1997).
9.   Stennard, F., Zorn, A. M., Ryan, K., Garrett, N. & Gurdon, J. B. Differential expression of VegT
and Antipodean protein isoforms in Xenopus. *Mech Dev* **86,** 87–98 (1999).
10.   Peshkin, L. *et al.* On the Relationship of Protein and mRNA Dynamics in Vertebrate Embryonic
Development. *Dev Cell* **35,** 383–394 (2015).
11.   Schohl, A. & Fagotto, F. Beta-catenin, MAPK and Smad signaling during early Xenopus
development. *Development* **129,** 37–52 (2002).
12.   Faure, S., Lee, M. A., Keller, T., Dijke, ten, P. & Whitman, M. Endogenous patterns of TGFbeta
superfamily signaling during early Xenopus development. *Development* **127,** 2917–2931 (2000).
13.   De Domenico, E., Owens, N. D. L., Grant, I. M., Gomes-Faria, R. & Gilchrist, M. J. Molecular
asymmetry in the 8-cell stage Xenopus tropicalis embryo described by single blastomere
transcript sequencing. *Dev Biol* **408,** 252–268 (2015).
14.   Charney, R. M. *et al.* Foxh1 Occupies cis-Regulatory Modules Prior to Dynamic Transcription
Factor Interactions Controlling the Mesendoderm Gene Program. *Dev Cell* **40,** 595–607.e4
(2017).
15.   Gentsch, G. E. *et al.* In vivo T-box transcription factor profiling reveals joint regulation of
embryonic neuromesodermal bipotency. *Cell Rep* **4,** 1185–1196 (2013).
16.   Gentsch, G. E., Monteiro, R. S. & Smith, J. C. Cooperation Between T-Box Factors Regulates
the Continuous Segregation of Germ Layers During Vertebrate Embryogenesis. *Curr Top Dev*
Biol **122,** 117–159 (2017).
17.   Whyte, W. A. *et al.* Master transcription factors and mediator establish super-enhancers at key
cell identity genes. *Cell* **153,** 307–319 (2013).
18.   Müller, C. W. & Herrmann, B. G. Crystallographic structure of the T domain-DNA complex of the
Brachyury transcription factor. *Nature* **389,** 884–888 (1997).
19.   Chen, X., Rubock, M. J. & Whitman, M. A transcriptional partner for MAD proteins in TGF-beta
signalling. *Nature* **383,** 691–696 (1996).
20.   Chiu, W. T. *et al.* Genome-wide view of TGFβ/Foxh1 regulation of the early mesendoderm
program. *Development* **141**, 4537–4547 (2014).
21.   Heasman, J., Kofron, M. & Wylie, C. βCatenin Signaling Activity Dissected in the Early Xenopus
Embryo: A Novel Antisense Approach. *Dev Biol* **222,** 124–134 (2000).
22.   Skirkanich, J., Luxardi, G., Yang, J., Kodjabachian, L. & Klein, P. S. An essential role for
transcription before the MBT in Xenopus laevis. *Dev Biol* **357,** 478–491 (2011).
23.   Reversade, B., Kuroda, H., Lee, H., Mays, A. & De Robertis, E. M. Depletion of Bmp2, Bmp4,
Bmp7 and Spemann organizer signals induces massive brain formation in Xenopus embryos.
Development **132,** 3381–3392 (2005).
24.   Blitz, I. L. *et al.* A catalog of Xenopus tropicalis transcription factors and their regional expression
in the early gastrula stage embryo. *Dev Biol* **426,** 409–417 (2017).
25.   Neph, S. *et al.* An expansive human regulatory lexicon encoded in transcription factor footprints.
Nature **489,** 83–90 (2012).
26.   Soufi, A. *et al.* Pioneer transcription factors target partial DNA motifs on nucleosomes to initiate
reprogramming. *Cell* **161,** 555–568 (2015).
27.   Gurdon, J. B. Cell Fate Determination by Transcription Factors. *Curr Top Dev Biol* **116,** 445–
454 (2016).
28.   Pasque, V., Jullien, J., Miyamoto, K., Halley-Stott, R. P. & Gurdon, J. B. Epigenetic factors
influencing resistance to nuclear reprogramming. *Trends Genet* **27,** 516–525 (2011).
29.   Lee, M. T. *et al.* Nanog, Pou5f1 and SoxB1 activate zygotic gene expression during the
maternal-to-zygotic transition. *Nature* **503,** 360–364 (2013).

30.   Iwafuchi-Doi, M. & Zaret, K. S. Cell fate control by pioneer transcription factors. *Development*
1211   **143,** 1833–1837 (2016).
1212   31.   Khokha, M. *et al.* Techniques and probes for the study of Xenopus tropicalis development. *Dev*
1213   *Dyn* **225,** 499–510 (2002).
1214   32.   Sive, H., Grainger, R. & Harland, R. Early development of Xenopus laevis: A laboratory manual.
1215   Cold Spring Harbor Press (2000).
1216   33.   Nieuwkoop, P. & Faber, J. Normal table of Xenopus laevis (Daudin): a systematical and
1217   chronological survey of the development from the fertilized egg till the end of metamorphosis.
1218   Garland (1994).
1219   34.   Rana, A., Collart, C., Gilchrist, M. & Smith, J. Defining synphenotype groups in Xenopus
1220   tropicalis by use of antisense morpholino oligonucleotides. *PLoS Genet* **2,** e193 (2006).
1221   35.   Hopwood, N., Pluck, A. & Gurdon, J. MyoD expression in the forming somites is an early
1222   response to mesoderm induction in Xenopus embryos. *Embo J* **8,** 3409–3417 (1989).
1223   36.   Gökirmak, T. *et al.* Localization and substrate selectivity of sea urchin multidrug (MDR) efflux
1224   transporters. *J Biol Chem* **287,** 43876–43883 (2012).
1225   37.   Owens, N. D. L. *et al.* Measuring Absolute RNA Copy Numbers at High Temporal Resolution
1226   Reveals Transcriptome Kinetics in Development. *Cell Rep* **14,** 632–647 (2016).
1227   38.   Ladher, R., Mohun, T. J., Smith, J. C. & Snape, A. M. Xom: a Xenopus homeobox gene that
1228   mediates the early effects of BMP-4. *Development* **122,** 2385–2394 (1996).
1229   39.   Smith, J., Price, B., Green, J., Weigel, D. & Herrmann, B. Expression of a Xenopus homolog of
1230   Brachyury (T) is an immediate-early response to mesoderm induction. *Cell* **67,** 79–87 (1991).
1231   40.   Monsoro-Burq, A. H. A Rapid Protocol for Whole-Mount In Situ Hybridization on Xenopus
1232   Embryos. *Cold Spring Harb Protoc*, pdb.prot4809 (2007).
1233   41.   Landt, S. G. *et al.* ChIP-seq guidelines and practices of the ENCODE and modENCODE
1234   consortia. *Genome Res* **22,** 1813–1831 (2012).
1235   42.   Zhang, C., Basta, T., Jensen, E. D. & Klymkowsky, M. W. The beta-catenin/VegT-regulated
1236   early zygotic gene Xnr5 is a direct target of SOX3 regulation. *Development* **130,** 5609–5624
1237   (2003).
1238   43.   Gentsch, G. E. & Smith, J. C. Efficient Preparation of High-Complexity ChIP-Seq Profiles from
1239   Early Xenopus Embryos. *Methods Mol Biol* **1507,** 23–42 (2017).
1240   44.   Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-Time
1241   Quantitative PCR and the 2CT Method. *Methods* **25,** 402–408 (2001).
1242   45.   Gentsch, G. E. & Smith, J. C. Investigating physical chromatin associations across the Xenopus
1243   genome by chromatin immunoprecipitation. *Cold Spring Harb Protoc* **2014,** (2014).
1244   46.   Gentsch, G. E., Patrushev, I. & Smith, J. C. Genome-wide snapshot of chromatin regulators and
1245   states in Xenopus embryos by ChIP-Seq. *J Vis Exp* **96,** e52535 (2015).
1246   47.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9,**
1247   357–359 (2012).
1248   48.   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-
1249   regulatory elements required for macrophage and B cell identities. *Mol Cell* **38,** 576–589 (2010).
1250   49.   Nakamura, Y., de Paiva Alves, E., Veenstra, G. J. C. & Hoppler, S. Tissue- and stage-specific
1251   Wnt target gene expression is controlled subsequent to β-catenin recruitment to cis-regulatory
1252   modules. *Development* **143,** 1914–1925 (2016).
1253   50.   Yoon, S.-J., Wills, A. E., Chuong, E., Gupta, R. & Baker, J. C. HEB and E2A function as
1254   SMAD/FOXH1 cofactors. *Genes Dev* **25,** 1654–1661 (2011).
1255   51.   Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res*
1256   **27,** 573–580 (1999).
1257   52.   Karpinka, J. B. *et al.* Xenbase, the Xenopus model organism database; new virtualized system,
1258   data types and genomes. *Nucleic Acids Res* **43,** D756–63 (2015).
1259   53.   Pertea, M., Kim, D., Pertea, G. M., Leek, J. T. & Salzberg, S. L. Transcript-level expression
1260   analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc* **11,** 1650–
1261   1667 (2016).
1262   54.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29,** 15–21 (2013).
1263   55.   Zhu, Q., Fisher, S. A., Shallcross, J. & Kim, J. VERSE: a versatile and efficient RNA-Seq read
1264   counting tool. *bioRxiv*, 053306 (2016).
1265   56.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–2079
1266   (2009).
1267   57.   Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq experiments. *Bioinformatics*
1268   **28,** 2184–2185 (2012).
58.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15,** 550 (2014).

59. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing
and web-based tools. *Nucleic Acids Res* **41,** D590–6 (2013).

60. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. *Genome Biol* **10,** R25 (2009).

61. Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of insertions,
deletions and gene fusions. *Genome Biol* **14,** R36 (2013).

62. UniProt Consortium. UniProt: a hub for protein information. *Nucleic Acids Res* **43,** D204–12
(2015).

63. Vierstra, J., Wang, H., John, S., Sandstrom, R. & Stamatoyannopoulos, J. A. Coupling
transcription factor occupancy to nucleosome architecture with DNase-FLASH. *Nat Methods* **11,**
66–72 (2014).

64. He, H. H. *et al.* Refined DNase-seq protocol and data analysis reveals intrinsic bias in
transcription factor footprint identification. *Nat Methods* **11,** 73–78 (2014).

65. Song, L. & Crawford, G. E. DNase-seq: a high-resolution technique for mapping active gene
regulatory elements across the genome from mammalian cells. *Cold Spring Harb Protoc*,
pdb.prot5384 (2010).

66. Kohany, O., Gentles, A. J., Hankus, L. & Jurka, J. Annotation, submission and screening of
repetitive elements in Repbase: RepbaseSubmitter and Censor. *BMC Bioinformatics* **7,** 474
(2006).

67. Smit, A., Hubley, R. & Green, P. RepeatMasker Open-4.0. 2013-2015.

68. Davies, J. O. J. *et al.* Multiplexed analysis of chromosome conformation at vastly improved
sensitivity. *Nat Methods* **13,** 74–80 (2016).

69. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome
assemblies. *Bioinformatics* **27,** 2957–2963 (2011).

70. Stark, R. & Brown, G. DiffBind: Differential binding analysis of ChIP-Seq peak data. 1–33 (2018).

71. Balwierz, P. J. *et al.* ISMARA: automated modeling of genomic signals as a democracy of
regulatory motifs. *Genome Res* **24,** 869–884 (2014).

72. Falcon, S. & Gentleman, R. Using GOstats to test gene lists for GO term association.
Bioinformatics **23,** 257–258 (2007).

73. Conesa, A. *et al.* Blast2GO: a universal tool for annotation, visualization and analysis in
functional genomics research. *Bioinformatics* **21,** 3674–3676 (2005).

74. Collart, C. *et al.* High-resolution analysis of gene activity during the Xenopus mid-blastula
transition. *Development* **141,** 1927–1939 (2014).

75. Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29,** 24–26 (2011).

76. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26,** 841–842 (2010).

77. Rosenthal, R. Meta-Analytic Procedures for Social Research. SAGE (1991).

78. Hontelez, S. *et al.* Embryonic transcription is controlled by maternally defined chromatin state.
Nat Commun **6,** 10148 (2015).

79. Beck, C. W. & Slack, J. M. Analysis of the developing Xenopus tail bud reveals separate phases
of gene expression during determination and outgrowth. *Mech Dev* **72,** 41–52 (1998).

1312
1313

Acknowledgments We thank Abdul Sesay, Leena Bhaw, Harsha Jani, Deborah Jackson and

Meena Anissi for deep sequencing; Mike Klymkowsky, Manolis Gialitakis and John Gurdon

for supplying antibodies; Stefan Hoppler and Yukio Nakamara for discussing ChIP protocol

for β-catenin; Ying Wang for providing sequence conservation tracks; Mareike Thompson,

Rita Monteiro and Clara Collart for critical reading of the manuscript; and the Smith lab for

discussions and advice. G.E.G and J.C.S. were supported by the Medical Research Council

(program number U117597140) and are now supported by the Francis Crick Institute, which

receives its core funding from Cancer Research UK (FC001-157), the UK Medical Research

Council (FC001-157), and the Wellcome Trust (FC001-157).

Extended Data

Extended Data Figure 1. Identification of most influential TFs on ZGA

a, Timeline of the maternal-to-zygotic transition during early *Xenopus* embryogenesis. Examples of maternal TFs Sox3, mPouV and mVegT being nuclear before the signal mediators β-catenin, Smad2 and Smad1 are based on our and previously published result[7-12] (also see Fig. 1f). **b**, Enrichment (hypergeometric p-value and coverage) of DNA motifs at RNAPII+ CRMs from 32-cell to 1000-cell stage (pre-MBT) and from MBT to late gastrula stage (MBT+). **c**, Snapshot of DNase-probed chromatin accessibility (biological replicates #1 and #2) and various other chromatin features (Sox3, RNAPII, H3K4me1, poly(A) RNA from this study and H3K4me3, H3K27me3, 5-methylcytosine [5mC] from a previously published study[78]) at the *sox2* locus (promoter-proximal and downstream distal regulatory region) around MBT to validate our novel DNase-Seq method. Negative control of DNase-Seq, DNase-treated naked genomic DNA (gDNA). **d**, Ribosome footprinting[6] of selected maternal and zygotic TFs across MBT. **e**, WMIHC of Smad1 and Smad2 on bisected early gastrula embryos. **f**, Western blotting of Sox3 in standard control and Sox3 morphants (Sox3 LOF) at blastula stage. α-tubulin, loading control. **g**, WMIHC of Sox3 in control and Sox3 LOF embryos at 64-cell and blastula stage. Nuclear accumulation of Sox3 detected in both the animal (an) and vegetal (veg) hemisphere of control embryos. Scale bar, 0.5 mm. **h**, Ribosome footprinting mapped to the *VegT* locus showing the post-MBT switch of translation from the maternal (m) to the zygotic (z) *VegT* transcript. Design of MO to block translation of the maternal transcript

Extended Data Figure 2. Validation of ChIP-grade antibodies

a, Sox3 immunoprecipitation (IP). IgG H and L, detected IgG heavy and light chains of the IP antibody. **b-e,i**, ChIP-qPCR for Sox3, β-catenin, Smad1, Smad2 and Tbx6 at indicated developmental stages. Mean+SD (n=2-3). One-tailed Student's t-test (comparing to IgG control): *, p ≤0.1 and ≥2-fold enrichment relative to lowest DNA recovery (% input) with ChIP antibody (dotted line). **f**, Tbx6 IP using *X. tropicalis* and *X. laevis* embryonic extract from late gastrula embryos. Antibody #4596 was chosen for subsequent IP and ChIP experiments, and #5061 for Western blotting. **g**, Western blotting (WB) of Tbx6 in standard control and Tbx6 morphants at early tailbud stage. α-tubulin, loading control. **h**, Relative expression levels of the transcript (RT-qPCR) and its protein (IP/WB) from early blastula to early tailbud stage. α-tubulin, loading control.

 Extended Data Figure 3. Super-enhancers of ZGA are associated with embryonic and
tissue-specific gene functions
a, Dynamic chromatin engagement of endogenous Sox3, Foxh1[14], VegT, β-catenin, Smad2,
Smad1 and RNAPII to super-enhancers of the *siamois* and *ventx* gene cluster from 32-cell to
early gastrula stage. Illustrated sagittal sections (dorsal side is right) show the nuclear
localisation of the selected TFs with arrows pointing to any tissue movements. Arrowheads,

promoter and enhancer activity based on the peak-like recruitment of RNAPII and/or TFs.
Rectangles, transcriptional engagement based on the broad distribution of RNAPII and the
occurrence of intronic RNA. **b**, Chromatin landscape of endogenous Sox3, Eomes, zVegT,
Xbra and Tbx6 to super-enhancer of the *mesp* gene cluster at late gastrula and/or early tailbud
stage. Illustrated sagittal sections (dorsal side is right) show the nuclear localisation of
selected TFs in late gastrula embryo and at the caudal end of early tailbud embryo. **c**, Super-
enhancer-like binding correlates with high levels of chromatin accessibility and DNA
occupancy of TFs, signal mediators and RNAPII. Grey box explains following plots. They show
normalised levels of chromatin accessibility or DNA occupancy (±20 kb from TSSs) for each
zygotic gene (detected by RNAPII profiling from 32-cell to late gastrula stage, see Methods)
and the cumulative frequency of detected super-enhancers. Genes with super-enhancer-like
binding <5kb from TSSs are highlighted in red. **d**, Significantly enriched biological processes
associated with the gene function of nearby super-enhancers detected at indicated
developmental stages. **e**, Meta-summary of chromatin accessibility and DNA occupancy levels
of TFs and signal mediators at zygotic genes (mean±SD; $\log_{10}$ scale).

Extended Data Figure 4. Endogenous chromatin dynamics in cells of embryonic
pluripotency and neuro-mesodermal bipotency
a-e, Endogenous dynamics of chromatin engagement between TFs (Sox3, mVegT, Foxh1[14],
Eomes[15], zVegT[15], Xbra[15] and Tbx6) and signal mediators (β-catenin, Smad1, Smad2)
covering developmental process of embryonic pluripotency and neuromesodermal (NM)
bipotency. **a**, Spearman correlation R_s of DNA occupancy levels. **b**, PCA of chromatin
engagement. **c**, PCA of RNAPII recruitment levels at CRMs occupied by TFs and signal
mediators. r, biological replicates. **d**, DNA occupancy-assisted clusters of CRMs co-aligned

 with enriched DNA motifs detected in peak centres (±50 bp) and flanking sequences (each 50
1399 bp). **e**, Coverage and enrichment of DNA motifs.

Extended Data Figure 5. Developmental context influences chromatin engagement
a-e, MyoD-enforced change of the development context in early gastrula embryos and its
influence on chromatin recruitment of Sox3 and RNAPII. **a**, Spearman correlation R_s of DNA
occupancy levels. **b**, Map of DNA occupancies and enriched DNA motifs sorted by significantly
differential (pΔ) Sox3 binding. **c**, Animal cap assay to measure MyoD-enforced transcription.
d, RT-qPCR results from the animal cap assay. Error bars, mean+SD (n=2). **e**, Opportunistic
recruitment of MyoD-HA to Sox3 target gene *sox2*. **f-h**, Spatio-temporal co-enrichment of DNA
motifs at CRMs occupied by Sox3, Eomes[15], Xbra[15] and/or Tbx6 at indicated developmental
stages reflect developmental context. **f**, Coverage and enrichment of DNA motifs. **g**,
Hierarchical clustering of DNA occupancies separated by time and space. **h**, Sox3-expressing
cells of the posterior neural tube originate from the chordoneural hinge (CNH) and posterior
wall of the neurenteric canal (PNC) expressing Xbra and Xbra/Tbx6, respectively. Both neural
tube and PNC are also exposed to the expression of Cdx such as Cdx4[79].

Extended Data Figure 6. Maternal TFs and signals are critical to ZGA-dependent tissue movements of gastrulation

Numbers attached to any embryo images refer to the count of embryos detected with the displayed phenotype among all embryos analysed. **a**, Morphological phenotypes caused by single and combined LOFs of Sox3 and mPouV when control embryos reached mid-gastrula and mid-tailbud stage. **b**, Phenotypical rescue of mPouV/Sox3 LOF embryos by the co-injection of both *X. laevis Pou5f3.3* and *Sox2* mRNA alongside *mCherry* mRNA as a tracer. **c**, Morphological phenotype caused by the LOF of mVegT at late tailbud stage. **d**, Rescue of reduced *nodal5*, *nodal6*, *sox17b* and *gata4* transcript levels in mVegT LOF embryos by the injection of *X. tropicalis* z*VegT* mRNA as determined by RT-qPCR at blastula stage. Error bars, mean+SD (n=5). Two-tailed Student's t-test: *, p <0.02. **e**, Morphological phenotypes caused by the single LOF of canonical Wnt, Nodal or BMP signalling when control embryos reached late gastrula and mid-tailbud stage. **f**, Morphological phenotype and transcriptome analysis of embryos injected with α-amanitin. PCA of the poly(A) RNA transcriptome of control and α-amanitin-injected embryos at indicated developmental stages in biological triplicates (#1-3).

Extended Data Figure 7. Signal-induced tissue-specific gene expression depends on maternal TFs

a, Detection of 3,318 zygotic genes with reduced transcript levels (≥50%) in α-amanitin-injected embryos of stage 8+, 9+ and 10+ (Extended Data Fig. 6f). These genes were used as reference for all other LOFs. Dots in scatterplot are coloured according to the maternal contribution[37] to the transcript level of each of these zygotic genes. List of some developmentally relevant genes whose numbers are used in **b** and **d** and Fig. 3b. **b**, Scatterplot of transcript fold changes (FCs) caused by mPouV and mPouV/Sox3 LOFs with dots coloured according to the ratio of FCs. **c**,**g**, Early gastrula-staged WMISH: *mir427*, animal view; *ventx*, lateral view, ventral side facing right; *Xbra*, dorsal view; *foxb1* and *zic1*, vegetal view, dorsal side facing top. Numbers in the right bottom corner of each image refer to the count of embryos detected with the displayed WMISH staining among all embryos analysed

per condition and *in situ* probe. Scale bars, 0.5 mm. **d**, Transcriptional comparison of zygotic
genes between indicated LOFs of maternal TFs or signals. Dots are coloured according to
regional expression along the animal-vegetal (An:Vg) or dorso-ventral (D:V) axis. **e**,**f**, Venn
diagrams of genes downregulated by indicated LOFs. **h**, Quantification of *foxb1* transcript
levels in control and mPouV/Sox3 LOF animal caps with or without canonical Wnt signalling.
Error bars, mean+SD (n=2). Two-tailed Student's t-test: 1, p=0.050; and 2, p=0.047.

**Extended Data Figure 8. Selected maternal TFs and signals regulate gene signature of
earliest tissue regionalisation and animal organogenesis**
a, Percentage of genes misregulated by indicated LOFs. **b**, Enrichment analysis of key GO
term-related biological processes among zygotic genes misregulated by indicated LOFs. Size
of bubble reflects the statistical significance (hypergeometric p-value) of enrichment while the
colour indicates the number of affected genes.

Extended Data Figure 9. Pioneering activity of mPouV and Sox3 initiate extensive chromatin remodelling.

a, Heat map of Spearman correlations R_s among indicated biological replicates of MBT-staged DNase-Seq on extracted chromatin and naked genomic DNA (gDNA). **b**, Double logarithmic Volcano plot of DNase cleavage changes at DHSs between mPouV/Sox3 LOF and uninjected embryos. **c**, Violin plot of DNase cleavages comparing uninjected and mPouV/Sox3 LOF embryos at all and affected (FDR <10%) DHSs. Wilcoxon test: *, p <2.2x10^{-16}. **d**, Map of various chromatin features, expressed RNA and enriched DNA motifs at DHSs sorted by significantly reduced DNase cleavages under mPouV/Sox3 LOF. #, β-catenin binding profile generated at late blastula stage rather than early blastula stage (Sox3) or MBT (all others). Top row, log-scaled meta-profiles (mean±SD) for chromatin features and RNA followed by meta-profile of DNA motif instances. Bottom row, normalised read coverage levels (and differential p-values, pΔ) followed by co-aligned map of DNA motifs.

Extended Data Figure 10. mPouV and Sox3 stabilise chromatin contacts with promoters
a,**b**, Performance of next-generation capture-C using novel 3C (chromatin conformation capture) protocol adapted to early *Xenopus* embryos. **a**, Bar graph of normalised counts (mean+SD; n=6) of non-redundant capture-reporter FLASH[69] reads for each viewpoint captured with 1 or 2 probes. **b**, Summary of FLASH reads from sequencing 3C libraries after two rounds of hybridisation-based capture. **c**, Violin plot of promoter-proximal interactions with unaffected and affected DHS for both all and affected (FDR <10%) chromatin contacts. Wilcoxon test and effect size estimates: 1, $p=5\times10^{-5}$ and $r_{effect}=0.12$ (small effect); 2, $p=5\times10^{-30}$ and $r_{effect}=0.34$ (medium effect); and 3, $p=5\times10^{-10}$ and $r_{effect}=0.78$ (large effect). **d**, Superimposed line plots showing the LOF-induced changes to various chromatin features at MBT including chromatin contacts with promoters (viewpoints) of *foxb1*, *cdc25b* and *zic1*. #, β-catenin binding profile generated at late blastula stage rather than early blastula stage (Sox3) or MBT (all others). RNA profile split into low and high expression windows. Low expression profile shows pervasively transcribed non-coding RNA. For statistical significance check heat map below each profile. Header, POU/SOX motif occurrences at DHSs and sequence conservation heat maps (phastCons, phyloP). Arrowheads denote strongly affected chromatin sites.

1483 **Supplementary Table 1**
1484 Maternal protein concentrations versus ribosome densities (translation) across the mid-
1485 blastula transition
1486
1487 **Supplementary Table 2**
1488 Transcriptional effect of maternal transcription factors and intercellular signals on zygotic
1489 genome activation
1490
1491 **Supplementary Table 3**
1492 Summary of deep sequencing and read alignments
1493
1494 **Supplementary Table 4**
1495 Genomic coordinates of selected capture-C viewpoints and probe DNA sequences used for
1496 hybridisation-based capture
1497
1498 **Supplementary Video 1**
1499 Gastrulation defects caused by the loss-of-function of maternal pluripotency factors