1   Genetic analyses led to the discovery of a super-active mutant of the RNA

2   polymerase I

Tommy Darrière[1], Michael Pilsl[2], Marie-Kerguelen Sarthou[1], Adrien Chauvier[1], Titouan

Genty[1], Sylvain Audibert[1], Christophe Dez[1], Isabelle Léger-Silvestre[1], Christophe Normand[1],

Anthony K. Henras[1], Marta Kwapisz[1], Olga Calvo[3], Carlos Fernández-Tornero[4], Herbert

Tschochner[2] and Olivier Gadal[1*]

[1] Laboratoire de Biologie Moléculaire Eucaryote, Centre de Biologie Intégrative (CBI),

Université de Toulouse, CNRS, UPS, Toulouse, France

[2]Universität Regensburg, Biochemie-Zentrum Regensburg (BZR), Lehrstuhl Biochemie III,

Regensburg, Germany

[3]Instituto de Biología Funcional y Genómica, IBFG-CSIC, Universidad de Salamanca,

Salamanca, Spain

[4]Centro de Investigaciones Biológicas, CSIC, Madrid, Spain

* Corresponding author

Email: olivier.gadal@ibcg.biotoul.fr (OG)

Abstract:

Most transcriptional activity of exponentially growing cells is carried out by the RNA

Polymerase I (Pol I), which produces a ribosomal RNA (rRNA) precursor. In budding yeast,

Pol I is a multimeric enzyme with 14 subunits. Among them, Rpa49 forms with Rpa34 a Pol

I-specific heterodimer (homologous to PAF53/CAST heterodimer in human Pol I), which

might be responsible for the specific functions of the Pol I. Previous studies provided insight

in the involvement of Rpa49 in initiation, elongation, docking and releasing of Rrn3, an

essential Pol I transcription factor. Here, we took advantage of the spontaneous occurrence of

extragenic suppressors of the growth defect of the *rpa49* null mutant to better understand the

activity of Pol I. Combining genetic approaches, biochemical analysis of rRNA synthesis and

investigation of the transcription rate at the individual gene scale, we characterized mutated

residues of the Pol I as novel extragenic suppressors of the growth defect caused by the

absence of Rpa49. When mapped on the Pol I structure, most of these mutations cluster

within the jaw-lobe module, at an interface formed by the lobe in Rpa135 and the jaw made

up of regions of Rpa190 and Rpa12. *In vivo,* the suppressor allele *RPA135-F301S* restores

normal rRNA synthesis and increases Pol I density on rDNA genes when Rpa49 is absent.

Growth of the Rpa135-F301S mutant is impaired when combined with exosome mutation

rrp6Δ and it massively accumulates pre-rRNA. Moreover, Pol I bearing Rpa135-F301S is a

hyper-active RNA polymerase in an *in vitro* tailed-template assay. We conclude that wild-

type RNA polymerase I can be engineered to produce more rRNA *in vivo* and *in vitro*. We

propose that the mutated area undergoes a conformational change that supports the DNA

insertion into the cleft of the enzyme resulting in a super-active form of Pol I.

Author summary:

The nuclear genome of eukaryotic cells is transcribed by three RNA polymerases. RNA

polymerase I (Pol I) is a multimeric enzyme specialized in the synthesis of ribosomal RNA.

Deregulation of the Pol I function is linked to the etiology of a broad range of human

diseases. Understanding the Pol I activity and regulation represents therefore a major

challenge. We chose the budding yeast *Saccharomyces cerevisiae* as a model, because Pol I

transcription apparatus is genetically amenable in this organism. Analyses of phenotypic

consequences of deletion/truncation of Pol I subunits-coding genes in yeast indeed provided

insights into the activity and regulation of the enzyme. Here, we characterized mutations in

Pol I that can alleviate the growth defect caused by the absence of Rpa49, one of the subunits

composing this multi-protein enzyme. We mapped these mutations on the Pol I structure and

found that they all cluster in a well-described structural element, the jaw-lobe module.

Combining genetic and biochemical approaches, we showed that Pol I bearing one of these

mutations in the Rpa135 subunit is able to produce more ribosomal RNA *in vivo* and *in vitro*.

We propose that this super-activity is explained by structural rearrangement of the Pol I

jaw/lobe interface.

1 Introduction

The nuclear genome of eukaryotic cells is transcribed by three RNA polymerases [1]. RNA

polymerase II (Pol II) transcribes most of the genome and is responsible for all messenger

RNA production. RNA polymerases III and I are specialized in the synthesis of a limited

number of transcripts. RNA polymerase III (Pol III) produces small structured RNAs,

including tRNAs and the 5S ribosomal RNA. RNA polymerase I (Pol I) produces a single

transcript, the large polycistronic precursor (35S pre-rRNA in yeast; 47S in human), which

constitutes the first step of ribosome biogenesis. Pre-rRNA is then processed by multiple

successive steps into the mature rRNAs (25S, 18S, and 5.8S in yeast; 28S, 18S and 5.8S in

human). Despite producing a single transcript, Pol I is by far the most active eukaryotic RNA

polymerase, responsible for up to 60% of the total transcriptional activity in exponentially

growing cells [2]. The strongly transcribed rRNA genes can be visualized using the DNA

spread method developed by Miller *et al,* 1969, in which the 35S rRNA genes (rDNA) exhibit

a "Christmas tree" configuration, with up to 120 polymerases per transcribed gene [3].

The full subunit composition and structural data are now available for the three nuclear RNA

polymerases of the budding yeast *Saccharomyces cerevisiae* [4,5][6,7]. Pol I contains a core

of shared or homologous subunits that are largely conserved in eukaryotes and archaea, as for

the other two nuclear RNA polymerases [8]. The two largest subunits (Rpa190 and Rpa135)

form the DNA-binding cleft that carries the catalytic center. Rpb5, Rpb6, Rpb8, Rpb10, and

Rpb12 are shared with Pol II and Pol III, whereas Rpc40 and Rpc19 are only shared with Pol

III. This nine-subunit core is associated with the stalk, a structure formed in Pol I by the

heterodimeric complex Rpa43/Rpa14, which is involved in docking the essential Rrn3

initiation transcription factor to the enzyme [9–12]. The Pol I-Rrn3 complex interacts with

promoter bound factors, the core factor (CF), forming the initially transcribing complex (ITC)

[13–16]. Additionally, Pol I and Pol III contain subunits that are functionally and structurally

related to Pol II-specific basal transcription factors, called the "Built-in Transcription Factors" [17–19]. Their presence in Pol I and Pol III results in a higher number of subunits, from 12 subunits in Pol II, to 14 and 17 for Pol I and Pol III respectively, and correlates with substantial transcript production from a few genes [8]. The heterodimer formed by Rpa34 and the N-terminal domain of Rpa49 (Rpa49Nt) in Pol I (equivalent to Rpc53 and Rpc37 in Pol III) is related to the basal transcription factor TFIIF, and stimulates endogenous transcript cleavage activity [18,20,21]. Rpc34 in Pol III and the Rpa49 C-terminal domain (Rpa49Ct) bear a tandem winged helix domain similar to TFIIE, also named A49tWH [18,20]. Rpa49Ct binds upstream DNA [15,22] and is involved in initiation and elongation [18,23]. Finally, Rpa12 in Pol I and Rpc11 in Pol III harbour a C-terminal domain involved in stimulating endogenous transcript cleavage activity, similar to that of TFIIS for Pol II [17,24].

Yeast genetic studies of Pol III and Pol I "Built-in Transcription Factors" have revealed striking differences, despite their clear similarities. Each Pol III subunit is essential, but none of the Pol I "Built-in Transcription Factors" is required for cell growth. Lack of Rpa34 or invalidation of Rpa49Nt, by removing the TFIIF-like heterodimer, has no effect on growth *in vivo* [18,25,26]. In contrast, full or C-terminal deletion of *RPA49* leads to a strong growth defect at all temperatures, which is more severe below 25°C [18,26,27]. Full deletion of *RPA12* leads to a strong growth defect at 25 and 30°C, and is lethal at higher temperatures [28]. Lack of the C-terminal extension of Rpa12 abolishes stimulation of intrinsic cleavage, without any detectable growth defect [17,24]. Finally, yeast strains carrying the triple deletion of *RPA49*, *RPA34* and *RPA12* are viable, but accumulate the growth defects associated with each of the single mutants [25].

Pol I is functional in the absence of Rpa49, but shows well-documented initiation and elongation defects, both *in vivo* and *in vitro* [23,26,27,29–31]. Restoration of active rRNA synthesis, in the absence of Rpa49, has been used to identify factors involved in initiation and

elongation, such as Hmo1 and Spt5 [26,30,32]. Here, we made use of the spontaneous

occurrence of extragenic suppressors of the growth defect of the *rpa49* null mutant [27] to

better understand the activity of Pol I. We showed that the suppressing phenotype was caused

by specific point mutations in the two largest Pol I subunits, Rpa190 and Rpa135. We

identified a small area around Phe301 in subunit Rpa135, at an interface formed by the lobe in

Rpa135 and the jaw made up of regions of Rpa190 and Rpa12, where most mutations cluster.

Characterizing the Rpa135-F301S allele, we showed in an *in vitro* assay, that such Pol I

mutant is more active than the wild-type enzyme. *In vivo*, overproduction of rRNA by Pol I

bearing the Rpa135-F301S mutation was observed in backgrounds where the nuclear

exosome activity is impaired by *RRP6* deletion.

Results

Isolation of extragenic suppressor mutants of the growth defect in absence of Rpa49

We characterized extragenic suppressors of the *RPA49* deletion to better understand how cell growth is achieved in the absence of Rpa49. *RPA49* full-deletion mutants show a strong growth defect at 30°C and are unable to grow at 25°C. However, spontaneous suppressors have been previously observed [27]. We quantified the frequency of occurrence of individual clones able to grow at 25°C. There was a low frequency of colony occurrence, comparable with the spontaneous mutation rate of a single control gene (*CAN1*; $< 5.10^{-6}$). We isolated more suppressors after irradiating the cells with UV light. UV irradiation, resulting in a survival rate of about 50%, increased the frequency of suppressor mutations by approximately 10-fold. We identified clones that grew at 25°C after three days and selected individual colonies, called SGR for Suppressor of Growth Defect of *RPA49* deletion, with various growth rates. We ranked SGR from 1 to 186 based on their growth rates at 25°C; *SGR1* had a growth rate comparable to the wild-type (WT) condition (Fig. 1A). We crossed the 186 SGR clones with a strain of the opposite mating type bearing the deletion of *RPA49* to obtain diploid cells homozygous for the *RPA49* deletion and heterozygous for each suppressor. The restoration of growth of the diploids at 25°C showed that all suppressor phenotypes obtained were fully or partially dominant. We focused on the most efficient suppressor clones, SGR1 and SGR2, and performed tetrad analysis to follow segregation of the observed suppression phenotype. Each suppressor phenotype was linked to a single locus in the genome and SGR mutants had no strong growth defect (SGR1 in Fig. 1A). We used global genomic mapping of SGR1 and SGR2, derived from "genetic interaction mapping" (GIM) methods [33] (Materials and Methods; S1 Fig), and found a genomic linkage with genes encoding the two largest Pol I subunits: *RPA135* for SGR1 and *RPA190* for SGR2 (S1 Fig). Sequencing of the genomic DNA revealed that SGR1 bears a double mutation, whereas SGR2 bears a single one

(*RPA135-I218T/R379K* and *RPA190-A1557V* alleles, respectively). Furthermore, we

identified an additional mutant, SGR3, in *RPA135* (*RPA135-R305L)*. The heterogeneity of the

growth induced by strong UV mutagenesis prevented suppressor cloning from the 183 other

SGR clones.

We next used the dominant phenotype of these suppressors to isolate more alleles, which

suppress the deletion phenotype of *RPA49* in *RPA190* and *RPA135*. We constructed a library

of randomly generated mutants (see Materials and Methods) by propagating plasmids bearing

WT *RPA135* or *RPA190* in a mutagenic *E. coli* strain. After phenotypic selection of *rpa49Δ*

mutants bearing a mutagenized Rpa190 or Rpa135 subunit at 25°C, each plasmid bearing a

suppressor allele was extracted, sequenced, and re-transformed into yeast to confirm the

suppressor phenotype. We thus isolated nine novel alleles of *RPA190* and thirteen of *RPA135*

that were able to restore growth of *rpa49* deletion mutant at 25°C (S1 Table). We evaluated

the suppression strength based on growth restoration relative to WT at 25°C, as for the SGR

strains. Suppressor alleles obtained by mutagenesis of *RPA190* or *RPA135*, more effective

than SGR1, 2, or 3 were identified (S1 Table). In conclusion, we identified 22 novel alleles of

genes coding for the two largest Pol I subunits as extragenic suppressors of the *rpa49Δ*-

associated growth defect.

***rpa190* and *rpa135* mutant alleles can bypass the need of *RPA49* for optimal growth**

The growth of the strains bearing one of six suppressor alleles (*RPA190-E1274K*, *RPA190-*

C1493R, *RPA190-L1262P*, *RPA135-R379G*, *RPA135-Y252H*, and *RPA135-F301S*) was

evaluated by a 10-fold dilution test (Fig. 1B), showing significant suppression by all in the

absence of Rpa49. In previous genetic studies, other genetic backgrounds that alleviate the

growth defect of *rpa49Δ* at 25°C were isolated: *rpa43-35,326* [26], decreased rDNA copy

number [29], Hmo1 over-expression [30], or Spt5 truncations [32]. For all these mutants,

rRNA synthesis was only partially restored in the absence of Rpa49 and significant

transcription defects remained. Here, we focused on the *RPA135-F301S* allele, the most

effective growth suppressor of the *RPA49* deletion: the *rpa49Δ RPA135-F301S* double mutant

grew almost as well at 25°C as the WT strain (Fig. 1B).

We sought further insight into the effect of the suppressors by integrating the *RPA135-F301S*

point mutation into the endogenous gene in three genetic backgrounds: WT, *rpa49Δ* (full

deletion), or *rpa49ΔCt*. Note that yeast bearing *rpa49ΔCt* or *rpa49Δ* full-deletion have a

similar growth defect, but have different Pol I subunits composition [26,27]. In the absence of

Rpa49, Rpa34 does not associate with transcribing Pol I while in strains bearing the *rpa49ΔCt*

allele, Rpa34 and Rpa49Nt remain associated with the polymerase [26,27]. The growth rate

was determined in each of these yeast strains at 30°C, in the presence or absence of *RPA135-*

F301S. The suppressor allele *RPA135-F301S* had no effect on growth in the WT strain

(doubling time of 102 min). The doubling time was 180 min for the *rpa49Δ* strain and

RPA135-F301S restored growth to a doubling time of 135 min. We observed similar

suppression on the *rpa49ΔCt* background.

Most suppressors mutations are clustered in the jaw-lobe module

Structural data are now available for Pol I in an inactive form [4,5], in complex with Rrn3

[12,23,34], associated with other initiation factors [13–15], in elongating forms [22,35], and

in the paused state [36](Fig. 2A). We mapped Rpa135 and Rpa190 residues that suppress the

growth defect of the mutant strain *rpa49Δ* onto the structure of WT Pol I in which the full

structure of Rpa49 was determined [15] (Fig. 2B). Most of the suppressor mutations, which

provided growth recovery (S1 table), appeared to be clustered at a specific interface between

the two largest subunits, Rpa190 and Rpa135 (Fig. 2B), between the lobe (Rpa135 - salmon)

and the jaw (Rpa190 - blue). In *RPA135*, we found five suppressor mutations, which modify a

small region of 60 residues within the lobe domain (S2 Fig). Note that in this region, three

amino acids "DSF" (D299, S300, F301), which are conserved among eukaryotic species, are

all mutated in suppressors (S2 Fig). Substituted residues likely result in destabilization of this

interface, suggesting a specific rearrangement of the interface lobe/jaw in each mutant. The

jaw is also characterized by the presence of a *β*-strand in the structure of Rpa12 (Rpa12 -

yellow: residues 46-51, Fig. 2B) that, along with four *β*-strands in Rpa190, forms a five-

stranded anti-parallel *β*-sheet. The Rpa12 *β*-strand faces the Rpa135 lobe domain (residue 252

to 315 of Rpa135 - salmon), in which six independent mutations were found, including

RPA135-F301S.

To evaluate the implication of Rpa12 in suppression, we then tested whether mutated alleles

of *RPA12* could behave as suppressors. We generated a library of randomly mutagenized

RPA12, and screen for *RPA12* alleles able to correct *rpa49Δ* growth defect at 25°C. Two

dominant alleles (*RPA12-S6L* and *RPA12-T49A*) indeed efficiently suppressed the growth

defect of *rpa49Δ* and of *rpa49ΔCt* (Fig. 2D, not shown here for *rpa49ΔCt*). *RPA12-S6L* and

RPA12-T49A obtained by random mutagenesis are specifically located in the "hotspot" at the

jaw/lobe interface. Threonine 49 of Rpa12 is located on the *β*-strand (Fig. 2C), facing residues

D299, S300, and F301 of Rpa135, and Rpa190-E1274 (Fig. 2B). The second mutation,

RPA12-S6L, is located in the N-terminal domain of Rpa12 (Fig. 2C).

In conclusion, all point mutations in Rpa190, Rpa135, and Rpa12 detected in the hotspot

domain of the jaw/lobe interface substitute for *RPA49 in vivo*.

Pol I bearing Rpa135-F301S or Rpa12-S6L restores efficient rRNA synthesis and Pol I

occupancy on rRNA genes in the absence of Rpa49Ct *in vivo*

We used yeast mutant cells with a low (about 25 copies, +/- 3 copies) and stabilized (*fob1Δ)*

number of rDNA repeats to better associate the growth phenotype of *RPA135-F301S* or

RPA12-S6L allele to rRNA synthesis activity and Pol I density on transcribed genes *in vivo*.

This genetic background is the best suited to study variations in the number of polymerase

molecules per rRNA gene because it has a low number of rDNA copies, almost all in the

active state with a very high Pol I loading rate [29,37,38]. We generated five strains in this

low copy background (bearing single mutations; *rpa49ΔCt*, *RPA135-F301S*, *RPA12-S6L* and

double mutants combining *rpa49ΔCt* with *RPA135-F301S* or *RPA12-S6L* alleles) and

determined their doubling time (Fig. 3A) and *de novo* synthesis of rRNA (Fig. 3B). Note that

rDNA copies number is similar between strains, as indicated by chromosome XII size in

pulse-field gel electrophoresis (S3 Fig). The presence of the *RPA135-F301S* allele in Pol I

effectively compensated the growth defect caused by the absence of C-terminal part of Rpa49

in this background.

Labelling of the nascent rRNA was performed using a 2-min pulse with ^{3}H adenine. We

performed the labelling in three independent cultures because of heterogeneity due to random

occurrence of suppressors in cell cultures of the *rpa49ΔCt* mutant. When compared to a WT

strain, RNA precursors synthesis was reduced approximately five-fold for *rpa49ΔCter*, even

under permissive conditions (30°C) (compare Fig. 3B, lane 1 to lanes 2 to 4). Pol I activity in

the presence of *RPA135-F301S*, with or without Rpa49Ct, was similar to that of the WT

enzyme (Fig. 3B, lane 5 and 6). Similarly, *RPA12-S6L* partially restored Pol I activity in the

absence of Rpa49Ct (compare Fig. 3B, lanes 7 and 8). Thus, *RPA135-F301S* and *RPA12-S6L*

appeared to largely restore rRNA production in the absence of C-terminal part of Rpa49.

To get insight in Pol I activity in suppressors strains, we combined the rRNA synthesis

quantification with the analysis of the Pol I distribution along the rRNA genes. We evaluated

Pol I density on transcribed genes by performing Miller spreads, the only technique that

currently allows the counting of individual Pol I molecules on a single rRNA genes [3,29].

Using Miller spreads, we previously showed that full deletion of *rpa49* resulted in a three-fold

decrease of Pol I density per gene [3,29]. We show here that strain expressing the *rpa49ΔCt*

allele results in a four-fold decrease of Pol I density per gene, with about 21 Pol I detected per

gene, as compared to about 91 detectable in WT condition (Fig. 2C). Expression of the

RPA135-F301S or *RPA12-S6L* allele, WT for *RPA49,* had no detectable influence on Pol I

density (Fig. 3C, *RPA135-F301S* and not shown). In contrast to strain *rpa49ΔCt*, double

mutant *rpa49ΔCt RPA135-F301S* or *rpa49ΔCt RPA12-S6L* showed significantly higher Pol I

occupancy (46 and 43 respectively instead of 21 Pol I molecules per gene). Using ChIP, we

reproduced that Pol I occupancy in absence of Rpa49Ct is drastically reduced [26]. We

confirmed Miller spread quantification, in which *RPA135-F301S* significantly increased Pol I

occupancy in absence of Rpa49Ct, although not to WT level (Fig. 3D).

Overall, these results show that the presence of the *RPA135-F301S*, or to a lesser extend

RPA12-S6L allele, in a strain lacking C-terminal part of Rpa49 restores rRNA synthesis to

WT levels. However, Pol I density on rRNA genes is only partly restored, indicative of an

improved transcription initiation, or increased stability of elongating Pol I in absence of

Rpa49Ct.

Genetic interplay between suppressors alleles and Pol I domains

Extensive genetic characterization of Pol I subunits together with recent structural analysis

have provided insight in their involvement in catalytic steps (initiation, pause release or

termination). To investigate suppression mechanism, we then decided to explore which

domains or subunits of Pol I are required for the suppression to occur. We tested deletion of

RPA14, *RPA34* or *RPA12*, and of *RPA190* alleles (*rpa190Δloop*) coding for Rpa190 lacking

specific domain. The structure of Rpa190 revealed the presence of an extended loop inside the

DNA-binding cleft folded in a "expander/DNA mimicking loop" conformation when Pol I is

in an inactive, dimeric form [4,5]. This loop is inserted in the jaw domain of Rpa190 (Fig.

2B), in the vicinity of the mutation hotspot. A small deletion of this Rpa190 domain (1361-

1390) resulted in a slight slow-growth phenotype [4]. We generated a larger deletion allele,

rpa190Δloop (deletion of residues 1342-1411 of Rpa190), and observed no associated growth

defect (Fig. 4A). We were unable to generate a viable double mutant when combining this

mutation with the *rpa49* full deletion. Thus, the DNA-mimicking loop is required for Pol I

activity in the absence of Rpa49. We next tested whether deletion of this loop influences the

suppression by the *RPA135-F301S* allele. Note that the *rpa190Δloop* combined with *RPA135-*

F301S had no growth phenotype. There was no difference in the growth of the *rpa49Δ*

RPA135-F301S double mutant and that of the triple mutant *rpa49Δ RPA135-F301S*

rpa190Δloop (Fig. 4A). Thus, the expander/DNA mimicking loop of Rpa190 is not required

for suppression, but is required for the viability of the *rpa49* deletion mutant.

Rpa34 forms a heterodimer with Rpa49Nt, and Rpa14 is essential in absence of Rpa49. We

then introduced *RPA135-F301S* in yeast strains lacking either Rpa34 or Rpa14 (Fig. 4B and

C). Growth of *RPA135-F301S/rpa34Δ* and *RPA135-F301S/rpa14Δ* double mutants were not

different from that of the single mutants. However, *RPA135-F301S* suppressed the growth

defect of the viable double mutant, *rpa34Δ rpa49Δ* (Fig. 4B). The double deletion mutant

lacking both Rpa49 and Rpa14 was not viable [25]. Introduction of the suppressor *RPA135-*

F301S, by genetic crossing, resulted in a triple mutant (*rpa14Δ rpa49Δ RPA135-F301S*) that

could grow, but slower than WT (Fig. 4C). We conclude that *RPA135-F301S* does not require

Rpa14 or Rpa34 for the suppression to occur.

We next evaluated which part of Rpa12 subunit was required for the suppression to occur

(Fig. 5). First, we evaluated growth of *rpa12* alleles when combined with *rpa49* deletion. The

C-terminal region of Rpa12 (TFIIS-like) is inserted towards the active center of Pol I to

stimulate intrinsic cleavage activity but is displaced during productive initiation and

elongation steps. C-terminal deletion of Rpa12 resulted in normal growth [24] (Fig. 5A- lane

2), although the *rpa12ΔCt* allele is unable to stimulate cleavage activity *in vitro* [17]. Full

deletion of *RPA12* led to a heterogeneous growth phenotype when propagated at 30°C. To

overcome this heterogeneity, we constructed a strain with *RPA12* under the control of the

regulatable pGAL promoter. Depletion of Rpa12 on glucose containing medium, like full

RPA12 deletion, resulted in a slight growth defect at 25°C, which was stronger at 30°C [28]

(Fig. 5A-lane 3). In contrast, *RPA49* deletion resulted in a growth defect at 30°C, which was

stronger at 25°C [27] (Fig. 5A-lane 4). Combining *rpa12ΔCt* with *rpa49Δ* resulted in a mild

synergistic phenotype, with a stronger growth defect at both 25°C and 30°C (Fig. 5A - lane

5). The double mutant lacking both full Rpa12 and Rpa49 subunits was viable, but had a

major growth defect [25] (Fig. 5A - lane 6).

Secondly, in double *rpa12/rpa49* mutants, we tested the expression of the suppressors alleles,

which were isolated (*RPA12-S6L*, *RPA12-T49A* or *RPA135-F301S*). We explored whether the

C-terminal extension of Rpa12 was necessary for suppression of the *rpa49Δ* phenotype. We

introduced the Rpa12 C-terminal truncation into the strain bearing both *rpa49Δ* and

suppressor alleles *RPA12-S6L* or *RPA12-T49A*. *RPA12-S6L* and *RPA12-S6L-ΔCt* resulted in

similar suppression of the *rpa49Δ* growth defect (Fig. 5B). *RPA12-T49A* or *RPA12-T49A-ΔCt*

also suppressed equivalently (data not shown). We then introduced the *RPA135-F301S* allele

in strains lacking Rpa49 with or without the entire Rpa12 subunit and assessed the

suppression phenotype at 25°C (Fig. 5C). The growth defect of *rpa49Δ* was completely

suppressed by the *RPA135-F301S* allele when Rpa12 was expressed (Fig. 5C, left panel),

whereas suppression mediated by *RPA135-F301S* was not detected in the absence of Rpa12

(Fig. 5C, middle panel). After 10 days, *rpa49Δ rpa12Δ* double mutant behaved exactly the

same with or without the *RPA135-F301S* allele, demonstrating that suppressor allele has no

effect in absence of Rpa12 (Fig. 5C, right panel).

In conclusion, we show that *RPA135-F301S* suppression of *rpa49Δ*-associated growth defect

does not require Rpa190 DNA mimicking loop, Rpa34, or Rpa14. Rpa12 C-terminal portion

involved in stimulating cleavage activity is also not required for suppression. In contrast,

Rpa12 N-terminal domain is required for the suppression to occur.

In vitro* characterization of RNA polymerase I bearing *RPA135-F301S

In vitro, C-terminal part of Rpa49 is essential in promoter-dependent transcription assay [23].

Our *in vivo* analysis suggests that in *RPA135-F301S* mutant background, C-terminal part of

Rpa49 is not required for rRNA synthesis. Our hypothesis is that Rpa135-F301S partly

compensates the requirement for C-terminal part of Rpa49 in initiation. We used the

promoter-dependent *in vitro* transcription system and tailed-template system to assess this

hypothesis (Fig. 6A and B). Results were strikingly different when using promoter-dependent

or tailed-template systems. After depletion of Rpa49, purified Pol I lacks both Rpa34 and

Rpa49 subunits and is the so-called Pol A* complex [23,25,31]. Pol I lacking subunits

Rpa49/Rpa34 (Pol A*, Fig. 6A lane 2) was almost inactive in promoter-dependent assay

when compared to wild-type Pol I (WT) or Pol I bearing Rpa135-F301S (Fig. 6A, lane 1 and

3). Note that addition of recombinant Rpa34/Rpa49, Rpa49Ct alone or Rpa34/Rpa49N-ter

stimulated transcription by Pol A* [23]. Ruling out our hypothesis, Pol I bearing Rpa135-

F301S did not restore promoter-dependent activity of RNA Pol I lacking Rpa49 (Fig. 6A lane

4). We next tested RNA synthesis in a tailed-template system. Pol I lacking Rpa34/Rpa49 was

partly deficient in tailed template assay (Fig. 6B, lane 2) [23]. Interestingly, in this promoter-

independent assay, we clearly observed that the RNA synthesis by the polymerase bearing

Rpa135-F301S was increased compared to one with the WT polymerase (Fig. 6B compare

lane 1 to 3). Moreover, Pol I bearing Rpa135-F301S fully restored tailed template production

of RNA Pol I lacking Rpa49 (Fig. 6B, lane 4). We conclude that in the *in vitro* promoter-

dependent transcription assay, *RPA135-F301S* suppressor does not correct initiation defect

due to the absence of Rpa49. However, in presence of Rpa135-F301S, a more efficient

polymerase is engineered, able to produce more ribosomal RNAs from DNA tailed template.

***In vivo* characterization of RNA polymerase I bearing Rpa135-F301S**

In vitro, Pol I bearing Rpa135-F301S is over-producing RNA compared to WT. However *in*

vivo, we could not reveal increased production of rRNA (2 min pulse labelling, see Fig. 3B).

Pre-rRNAs which are not properly folded into pre-ribosome are targeted to degradation by the

3' to 5' exoribonucleolytic activity of the exosome [39]. We hypothesized that overproduced

rRNAs in *RPA135-F301S* mutant background could be targeted by the nuclear exosome**.**

Rrp6, part of the nuclear exosome complex, was deleted in a WT strain and in a strain bearing

RPA135-F301S mutation. We observed a strong synergistic growth defect in strain bearing

both *RPA135-F301S* and the deletion of *RRP6* (S5 Fig). Northern blot analysis (Fig. 7A)

showed that accumulation in *RPA135-F301S* single mutant was indistinguishable from the

WT for all RNA probed. *rrp6Δ* single mutant accumulates 23S and 35S (pre-)rRNA [40,41]

and in correlation with the growth defect of the double mutant *RPA135-F301S rrp6Δ*, we

could observe a 2-fold increase in 35S and 23S (pre-)rRNA accumulation as compared to

rrp6Δ. Accumulation of 35S and 23S could indicate an increase of RNA production, or a

defect in early maturation steps. We decided to directly assess over-expression of (pre-)rRNA

using short *in vivo* labelling experiments (40 seconds). As previously reported with very short

pulse labelling, accumulation of 20S rRNA is barely detectable, while 27SA and 35S are

already accumulated [42]. We could detect a strong accumulation of newly synthesized (pre-)

rRNA in the double mutant *RPA135-F301S rrp6Δ* compared to WT, *RPA135-F301S or rrp6Δ*

strains. Note that increased background signal could indicate an accumulation of partially

degraded, abortive transcripts or elongating rRNA transcript of various sizes. These results

suggests that rRNA are over-expressed in strain bearing *RPA135-F301S,* but are quickly

decayed by Rrp6. We confirmed this observation by evaluating ongoing transcription

independently of decay machinery using high-resolution transcriptional run-on (TRO)

analysis (Fig. 7C). Indeed, TRO assay make use of 10% sarkosyl, which permeabilizes cell

membranes, reversibly blocks elongating polymerases and inhibits RNAse activity [43–46].

Permeabilized cells are then incubated with $[\alpha^{32}P]$-UTP to resume transcription.

Neosynthesized radiolabeled RNAs are extracted, and used to probe slot-blots loaded with

single strand DNA fragments complementary to rDNA locus. Using incorporation of $[\alpha^{32}P]$-

UTP in the 5S rRNA transcribed by RNA polymerase III as internal control, TRO revealed a

three-fold increase of rRNA transcription in cells bearing *RPA135-F301S* allele, irrespective

of Rrp6 presence (Fig. 7C). These results confirmed that Pol I bearing Rpa135-F301S is over-

producing RNA compared to WT and that over-produced RNAs are targeted for degradation

by the exosome. All together, we concluded that Pol I bearing Rpa135-F301S is a hyper-

active RNA polymerase *in vivo*.

1   Discussion

2   Here, we characterized extragenic suppressors of the growth defect of the *rpa49* null mutant

3   to better understand the activity of Pol I. We showed that altering a very specific area of Pol I

4   resulted in an enzyme with modified catalytic properties sufficient to restore wild-type growth

5   in absence of Rpa49.

7   Suppressor mutations are not at the Rrn3-Pol I stalk interface

Our previous studies suggested the specific involvement of Rpa49 in the association and

dissociation of initiation factor Rrn3 from the Pol I stalk [26,29]). Here, we show that

genetically modified polymerases lacking Rpa49 or Rpa49Ct, with a single modified residue

in Rpa190, Rpa135, or Rpa12, at a position diametrically to the position that binds to Rrn3,

can initiate transcription and that strains harbouring them grow normally. Moreover, mutant

Pol I with Rpa135-F301S does not restore promoter dependent activity in absence of Rpa49.

We propose that, independently of the important interplay between Rpa49 and Rrn3 during

initiation, Rpa135-F301S can stimulate Pol I activity.

17   A novel role of Rpa12 subunit

The Rpa12 subunit is involved in stimulating the intrinsic cleavage activity of Pol I through a

TFIIS-like domain at its C-terminus. Purified Pol I with Rpa12 lacking the C-terminal domain

has no cleavage activity [17]. Furthermore, the C-terminal domain of Rpa12 can contact the

active site of the polymerase in the inactive conformation and is retrieved in both initiation

competent and elongating forms of the polymerase. However, the cleavage activity is re-

activated when Pol I is paused [36]. Direct evidence that cleavage is not involved in

suppression of the growth defect came from the experiments showing a fully functional

suppressor phenotype for *RPA12ΔCt-S6L*, which lacks the domain required for stimulating

cleavage.

The N-terminal domain of Rpa12, at the surface of Pol I, is involved in the recruitment of the

largest subunit, Rpa190 [24] and is required for docking this subunit to the enzyme. A linker

region of Rpa12 connects its N-terminal module (equivalent to the N-terminal domain in the

Pol II subunit Rpb9) at the surface of Pol I to its mobile C-terminal region (TFIIS-like) and is

therefore indirectly required for cleavage. *In vitro*, purified Pol I that lacks Rpa12 has less

activity than WT Pol I in promoter-dependent transcription assays (data not shown).

Mutations in other Pol I domains, such as deletions in the Rpa190-DNA mimicking loop,

Rpa34 or Rpa14, did not influence suppression of the *rpa49* deletion growth defect by the

RPA135-F301S allele. In contrast, the Rpa12 linker was absolutely required for efficient

suppression. Accordingly, *RPA135-F301S* allele was unable to restore efficient growth when

Rpa12 was absent. Thus, Rpa12 and *RPA135-F301S* likely cooperate in the super-active Pol I

enzyme. Note that rearrangement of Rpa12 have recently be shown to correlate with

dissociation of Rpa49/Rpa34 heterodimer from Pol I, confirming the tight interplay between

those subunits [47].

Modification of the jaw/lobe interface may facilitate DNA cleft closure

Pol I undergoes major conformational changes during the transcription cycle, mainly affecting

the width of the DNA-binding cleft [48]. During the initiation of transcription, the cleft

aperture narrows from a semi-open configuration, as seen in cryo-EM structures of the

enzyme bound to Rrn3 [12,23,34], to a fully closed conformation observed in transcribing

complexes [13,15] (Fig. 8). This allows gripping of the transcription bubble inside the cleft

(Fig. 8A-B). Following Rrn3 release, DNA binding is further secured, by the Rpa49-linker,

which crosses the cleft from the lobe to the clamp, passing over the downstream DNA, and by

the Rpa49Ct, which binds the upstream DNA in the vicinity of the clamp [15,22]. Therefore,

Rpa49Ct is anchored in a position securing cleft closure (Fig. 8B).

Cleft closure is achieved by the relative movement of two structural units, located on opposite

sides of the cleft, pivot with respect to each other using five hinges [4]. The unit consisting of

the shelf and clamp modules appears to be rigid, whereas the unit comprising the core and

lobe modules, which is in the vicinity of the mutated residues involved in suppression of the

rpa49Δ growth defect, undergoes internal rearrangements (S1 Movie). The most prominent

reorganization within this latter unit affects the Rpa190 jaw domain, the outer rim of which

shifts away from the DNA by approximately 3.7 Å, using the lobe/jaw interface as a hinge

(Fig. 8B; blue arrow). This movement also involves the linker region of Rpa12, which

contains a *β*-strand (residues 46-50) that completes a four-stranded *β*-sheet in the Rpa190 jaw

domain. As a result, a short α-helix within the Rpa12 linker region shifts by approximately

3.0 Å (Fig. 8B; yellow arrow).

Rearrangements in the jaw made up of regions of Rpa190 and Rpa12 are likely essential to

allow pivoting of the shelf-clamp unit against the core-lobe unit. Without such motion, cleft

closure would be impossible (S1 Movie). Structural analysis suggests that the C-terminal

domain of Rpa49 and its linker domain are involved in securing the closed cleft conformation

[15,22]. Cleft closure is likely destabilized in the *rpa49Δ* mutant (Fig. 8C). We propose that

Rpa135-F301S or Rpa12-S6L favors DNA capture by increasing the flexibility of the

lobe/jaw/Rpa12 interface of Pol I relative to that of the WT polymerase, facilitating cleft

closure in the absence of Rpa49 (Fig. 8D, red arrow). As shown *in vitro,* mutant Pol I with

Rpa135-F301S is super-active. *In vivo*, the increased accumulation of pre-rRNA, detectable in

absence of the nuclear exosome (*rrp6Δ)*, is in agreement with the *in vitro* super-activity. We

propose that Pol I bearing Rpa135-F301S might facilitate cleft closure, in addition to secure

cleft closure by Rpa49. Such mutant enzyme could capture DNA more efficiently than WT

polymerase (Fig. 8E). Alternatively, the mutations could also favour a 'closed cleft'

conformation in the presence of DNA with little impact on flexibility.

Pol I cleft closure is a limiting step in catalytic cycle

Here, we show that Pol I can be engineered to generate more rRNA. A super-active mutant of

Pol II was already characterized: point mutation in trigger loop (Rpb1-E1103G) leads to

increase RNA polymerization rate, affecting pausing and transcriptional fidelity [49,50].

Despite a very similar active site organization, an analogous mutation in the Pol I trigger loop

resulted in a very different outcome, as it reduces the elongation rate [51]. Such observations

led to the conclusion that Pol I catalytic cycle is very different than the one of Pol II, with

different rate limiting steps [51–53]. We show here that mutations away of the active center

can lead to a super-active form of Pol I. This difference can stem from wide-open

configuration of the DNA-binding cleft in Pol I compared to other polymerases. We propose

that Pol I cleft opening and closure is a limiting step in the Pol I catalytic cycle.

Materials and Methods

Construction of plasmids and yeast strains

The oligonucleotides used in this study are listed in S4 Table. Plasmids and details of the cloning steps are described in S3 Table. Randomly mutagenized *RPA190* and *RPA135* libraries were obtained by transformation and amplification of pVV190 and pNOY80, respectively, into XL1-red strains according to the manufacturer's guidelines (XL1-Red Competent Cells, from Agilent Technologies). Yeast strains are listed in S2Table, and were constructed by meiotic crossing and DNA transformation [54][55]. The yeast media and genetic techniques were described previously [56]. yCNOD226-1a was obtained from yCNOD223-2a by switching the *KAN-MX* to the *NAT-MX* marker under the control of the MF(ALPHA)2/YGL089C promoter (alphaNAT-MX4), which allowed selection of MATα haploid cells [33]. OGT9-6a is an offspring of yCNOD226-1a crossed with BY4741. OGT8-11a is an offspring of BY4742 crossed with Y1196. Strains OGT9-6a and OGT8-11a were plated on rich media and UV irradiated (5W/m^2 during 5 second), resulting in 50% survival. LH514D and LH11D are suppressor clones of the growth defect selected from UV-irradiated OGT9-6a grown at 25°C. AH29R is a suppressor clone of the growth defect selected from UV-irradiated OGT8-11a grown at 25°C. Genetic interaction mapping (GIM) analysis of the *RPA49* deletion mutant was performed as described previously [33]. Microarray data were normalized using MATLAB (MathWorks, Inc., Natick, MA) as previously described [29]. OGT15-7b is an offspring of LH514D with BY4741, followed by homologous recombination using PCR-amplified fragments generated with oligos 1716 and 1717 and pCR4-HIS3 as template. yTD16-1a was first transformed with plasmid pCJPF4-GAL49-1. Then, *RPA135* was tagged by homologous recombination using PCR-amplified fragments generated with oligos 835 and 836 and genomic DNA of strain *RPA135-TAP* or yTD6-6c, generating yTD27-1 and yTD28-1a, respectively. Strain yTD25-1a bears a C-terminal deletion of *RPA49*

generated by homologous recombination using PCR-amplified fragments generated with oligos 208 and 1515 and pFA6-KAN-MX6 as template. yTD11-1a was derived from strain yTD25-1a after switching to HPH-MX by homologous recombination using pUC19-HPH cut by *Bam*HI. C-terminal deletion of *RPA49 in* yTD29-1a and yTD30-1a were generated by homologous recombination using PCR-amplified fragments generated with oligos 649 and 650 and yTD11-1 genomic DNA as template, transformed into yTD27-1 and yTD28-1a, respectively. Genomic allelic insertion of *RPA12-S6L* in yTD31-1a and yTD23-1a was performed by homologous recombination using PCR-amplified fragments generated with oligos 1556 and 1557 and pRS316-A12-S6L-KAN as template, transformed into yTD27-1 and yTD29-1a, respectively.

TGT135-3b was obtained after sporulation of y27138 transformed by pNOY80. TGT135-3b and OGT15-9d were mated to generate TGT12. yTD2-3b and yTD2-3d are offspring of TGT12 transformed with pGL135_33. yTD6-6c and yTD6-6b were generated by homologous recombination using pTD2_6c_135TAP cut with *Xho*I-*Nsi*I, transformed into yTD2-3b and yTD2-3d, respectively. yTD48-1a was generated by deletion of the Rpa190 DNA-mimicking loop using homologous recombination with PCR-amplified fragments generated using oligos 1189 and 1194 and genomic DNA of strain SCOC2260 as template, transformed into BY4741.

yTD51-2c, yTD51-8a, and yTD51-5a are offspring of yTD48-1a mated with yTD37-3a. yTD36-2b is an offspring of yCN224-1a mated with yTD40-1a. yTD37-3d and yTD37-7d are offspring of yCN224-1a mated with yTD41-1a. yTD38-3d is an offspring of yCN225-1a mated with yTD40-1a and yTD39-8a is an offspring of yCN225-1a mated with yTD41-1a.

Strain yTD53-1a was constructed by homologous recombination using a PCR-amplified fragment generated with oligos 1634 and 1635 and pFA6a-KanMX6-GAL::3HA as template.

OGT30-1a and OGT30-3a are offspring of yTD53-1a mated with yTD6-6b. yTD40-1a and

yTD41-1a were generated by homologous recombination using PCR-amplified fragments

generated with oligos 700 and 1679 and pFA6a-HA-KlURA3 as template, transformed into

OGT30-3a and OGT30-1a respectively, switching *RPA135-TAP*-tag to untagged *RPA135*.

Strain yMKS8-1a was constructed by homologous recombination using a PCR-amplified

fragment generated with oligos 1711 and 1713 and pFA6a-KanMX6-GAL::3HA as template,

transformed into strain yCD2-2a. yMKS9-9d is offspring of yMKS8-1a mated with yTD6-6c.

Mapping extragenic suppressors allele by genetic linkage

Extragenic suppressors allele SGR were mapped using GIM methods, in which a strain is

crossed with the entire pool of haploid deletion strains [57]. Here, deletions are used to

evaluate linkage to SGR locus. Due to the genetic suppression, Individual deletions

genetically linked to SGR are counter-selected in *rpa49Δ* background. Genetic mapping of

SGR locus is based on strong genetic linkage of counter-selected deletion alleles (see S1 Fig),

evaluated using micro-array of deletion bar-code (ArrayExpress accession E-MTAB-7831)

[58].

In vivo **labelling and RNA extraction and analysis**

Metabolic labelling of pre-rRNA was performed as previously described [59] with the

following modifications. Strains were pre-grown in synthetic glucose-containing medium

lacking adenine at 30°C to an OD_{600} of 0.8 at. One-milliliter cultures were labeled with 50

μCi [8-^{3}H] adenine (NET06300 PerkinElmer) for 2 min. Cells were collected by

centrifugation and the pellets were frozen in liquid nitrogen. RNA was then extracted as

previously described [60] and precipitated with ethanol. For high molecular weight RNA

analysis, 20% of the RNA was glyoxal denatured and resolved on a 1.2% agarose gel. Low

molecular weight RNAs were resolved on 8% polyacrylamide/8.3 M urea gels.

Miller spreads experiments and analysis

Chromatin spreading was mainly performed as described previously with minor modifications [61]. Carbon-coated grids were rendered hydrophilic by glow discharge instead of ethanol treatment. Negatively stained chromatin was obtained by short incubation with heavy metal followed by quick drying of the sample. Images were obtained using a JEOL JEM-1400 HC electron microscope (40 to 120 kV) with an Orius camera (11Mpixels). The position of the RNA polymerase I molecules and the rDNA fiber were determined by visual inspection of micrographs using Image J (http://rsb.info.nih.gov/ij/). Digital images were processed by software programs Image J and Adobe Photoshop® (v. CS6).

In vitro **promoter-dependent and tailed template transcription assays**

In vitro promoter-dependent transcription reactions were performed as previously described [23,62] with some modifications. Briefly, 1.5 ml reaction tubes (Sarstedt safety seal) were placed on ice. Template (0.5–1 µl; 50–100 ng DNA) was added, corresponding to a final concentration of 5–10 nM per transcription reaction (25-µl reaction volume). Core factor (1–2 µl; 0.5 to 1 pmol/µl; final concentration 20–40 nM) and 1–3 µl Pol I (final concentration 4–12 nM) were added to each tube. Then, 20 mM HEPES/KOH pH 7.8 was added to a final volume of 12.5 µl. Transcription was started by adding 12.5 µl 2X transcription buffer. The samples were incubated at 24°C for 30 min at 400 rpm in a thermomixer.

nM of tailed templates were used in a total volume of 25 µl. The transcription was performed as described in promoter-dependent transcription reactions: Tailed templates are PCR-amplified fragments generated with oligos 1834 and 1835 and "pUC19tail_g-_601_elongated" as PCR-template, treated with Nb.BsmI (NEB) to generate a nick. Nicked site allows non-specific initiation of Pol I without the addition of initiation factors.

Transcription was stopped by adding 200 µl Proteinase K buffer (0.5 mg/ml Proteinase K in

0.3 M NaCl, 10 mM Tris/HCl pH 7.5, 5 mM EDTA, and 0.6% SDS) to the supernatant. The

samples were incubated at 30°C for 15 min at 400 rpm in a thermomixer. Ethanol (700 µl)

p.a. was added and the tubes mixed. Nucleic acids were precipitated at -20°C overnight or for

30 min at -80°C. The samples were centrifuged for 10 min at 12,000g and the supernatant

removed. The precipitate was washed with 0.15 ml 70% ethanol. After centrifugation, the

supernatant was removed and the pellets dried at 95°C for 2 min. RNA in the pellet was

dissolved in 12 µl 80% formamide, 0.1 M TRIS-Borate-EDTA (TBE), 0.02% bromophenol

blue, and 0.02% xylene cyanol. Samples were heated for 2 min with vigorous shaking at 95°C

and briefly centrifuged. After loading on a 6% polyacrylamide gel containing 7M urea and 1X

TBE, RNAs were separated by applying 25 watts for 30–40 min. The gel was rinsed in water

for 10 min and dried for 30 min at 80°C using a vacuum dryer. Radiolabelled transcripts were

visualised using a PhosphoImager.

RNA extractions and Northern Hybridizations.

RNA extractions and Northern hybridizations were performed as previously described [60].

For high molecular weight RNA analysis, 3µg of total RNA were glyoxal denatured, resolved

on a 1.2% agarose gel and transferred to a nylon membrane. The sequences of

oligonucleotides used to detect the RNA species: 35S rRNA, 25S rRNA, 20S rRNA, 18S

rRNA, 27S rRNA, *PGK1* mRNA and *SCR1* ncRNA are respectively 774, 1829, **1833**, 892,

1830, 1831 and 1832 are reported in S4 Table.

Chromatin immunoprecipitation (ChIP). ChIPs were performed essentially as described

previously [63]. At least three independent cultures of each yeast strain were grown to

exponential phase OD_{600} = 0.4-0.8 in YPD (yeast extract-peptone-glucose) at 30°C, and cross-

linked for 10 min at RT by the addition of formaldehyde (F1635, SIGMA) to a final

concentration of 1.2%. Adding glycine quenched the cross-linking reaction. Cells were

broken in lysis buffer (50 mM HEPES [pH 7.5], 150 mM NaCl, 1 mM EDTA, 1% Triton X-

100, 0.1% deoxycholate Na, 0.1% SDS, 1 mM AEBSF [Euromedex] and cOmplete EDTA-

free [Roche]) with glass beads (diameter, 0.425 to 0.6 mm; SIGMA) in a Precellys 24

homogenizer (bertin technologies) for 3 min, 6000 rpm at 4°C. Extracted chromatin was

sonicated to obtain 300-500 bp DNA fragments. 100 μg of sonicated chromatin (protein

content measured using BCA Protein Assay kit, ref# 23225, Thermo Scientific) was

immunoprecipitated for 4h at 21°C on Pan mouse Dynabeads (Invitrogen, 200 μl).

Immunoprecipitated DNA was purified and quantified by real-time qPCR using iTaq

universal SYBR®Green Supermix (Bio-Rad) and the ViiA7 AB Applied Biosystems (Life

Technologies). Primer pairs used for amplification are listed in S4 table. Signals were

analysed with Quant Studio Real Time PCR Software v1.1 and are expressed as percentage of

input DNA. Bars on the graph show the median of values normalised to wild-type value.

Error bars correspond to standard deviations of at least three independent cultures for each

strain.

Pulse-field gel electrophoresis (PFGE). PFGE is a technique that resolves chromosome-

sized DNA molecules in an agarose gel. Over-night YPD cultures at 30°C were harvested in

0.1% azide Na and kept on ice. Cells were washed twice in cold 0.05 M EDTA [pH 8.0].

~2.10^8 cells were resuspended in 1ml of freshly prepared NZ buffer (Citrate Phosphate buffer

33 mM, ETDA 0.05 M [pH 8.0], Sorbitol 1.2 M). Cells were pellets (8000 rpm, 1 min, 4°C)

and resuspended in 500 μL of Zymolyase buffer (NZ buffer with 20T zymolyase) and 500 μL

LMP 2% (SeaKem LE agarose, Lonza, ref# 50001). Homogenized pellets were poured into

the plug molds (~100 μl/plug) and incubated for 30 min in humid chamber at 37°C then 10

min at 4°C. Solidified plugs were transferred in 14 ml tubes with round bottom and incubate

with 400 μl/plug of PK buffer (EDTA 0.125 M [pH 9.5], Sarkosyl 1%, PK 1 mg/ml (Roche,

ref# 03115801001) and incubated 1-2 hours at 37°C with gentle agitation. PK buffer was changed and plugs were incubated O/N with gentle agitation. De-proteinated plugs were washed with 10 ml TE followed by washing with 10 ml TE + 1 mM AEBSF (Euromedex) for 2 hours. AEBSF was washed out twice with 10 ml of 0.05 M EDTA [pH 8.0] for 30 minutes. Plugs were stored at 4°C till usage in TE or 0.05 M EDTA [pH 8.0]. Plugs were loaded on 0.8% agarose gel in 1×TAE buffer (Certified Megabase agarose, Bio-Rad, ref# 161-3109) and run in 1×TAE buffer for 50h in CHEF-DRIII System with chiller unit (Bio-Rad) with following parameters: 3V/cm, initial S/time 250 sec, final S/time 900 sec, angle 120. Gel was stained with ethidium bromide (10 mg/ml in water). Gel was processed for Southern blotting: DNA was transferred on nylon membrane by passive transfer in 10×SSC buffer (Amersham Hybond XL, GE Healthcare). Membranes were UV-cross-linked and hybridized overnight with ^{32}P-labeled single-stranded DNA probes at 42 °C. Blots were washed twice with 2×SSC and 1% SDS for 20 min and once with 0.5×SSC and 1% SDS for 20 min at 37 °C. rDNA locus (chromosome XII) was detected with 18S specific probe (CATGGCTTAATCTTTGAGAC). This protocol was kindly provided by B. Pardo (IGH, Montpellier, France), and is extensively described in [64].

Transcriptional run-on analysis

TRO was performed as previously described [39,43]. Slot blots were loaded with single-stranded 80-mers DNA oligonucleotides: 1855 (NTS2), 1857 (5'ETS), 1859 (18S.2), 1860 (25S.1), 1861 (3'ETS), 1862 (NTS1), 1863 (5S US) and 1864 (5S DS).

1 Acknowledgements

We would like to thank the imaging platform of Toulouse TRI for their assistance. We thank

Margaux Cescato, Lilian Guillot and Jorge Perez-Fernandez for strain and plasmid

constructions. YCp50-26 carrying *RPA49* was kindly provided by P. Thuriaux. We also thank

Melanie Panarotto for the Miller spreads analysis. This work benefited from exchanges with

all unit members, including Anthony Henras, Frederic Beckouët and Lise Dauban. We thank

Stephanie Balor and Vanessa Soldan from the METI platform for the TEM acquisitions.

References

1. Chambon P (1975) Eukaryotic nuclear RNA polymerases. Annu Rev Biochem 44: 613–638. doi:10.1146/annurev.bi.44.070175.003145.

2. Warner JR (1999) The economics of ribosome biosynthesis in yeast. Trends Biochem Sci 24: 437–440.

3. Miller OL, Beatty BR (1969) Visualization of nucleolar genes. Science 164: 955–957.

4. Fernández-Tornero C, Moreno-Morcillo M, Rashid UJ, Taylor NMI, Ruiz FM, et al. (2013) Crystal structure of the 14-subunit RNA polymerase I. Nature 502: 644–649. doi:10.1038/nature12636.

5. Engel C, Sainsbury S, Cheung AC, Kostrewa D, Cramer P (2013) RNA polymerase I structure and transcription regulation. Nature 502: 650–655. doi:10.1038/nature12712.

6. Hoffmann NA, Jakobi AJ, Moreno-Morcillo M, Glatt S, Kosinski J, et al. (2015) Molecular structures of unbound and transcribing RNA polymerase III. Nature 528: 231–236. doi:10.1038/nature16143.

7. Cramer P, Bushnell DA, Kornberg RD (2001) Structural basis of transcription: RNA polymerase II at 2.8 angstrom resolution. Science 292: 1863–1876. doi:10.1126/science.1059493.

8. Werner F (2008) Structural evolution of multisubunit RNA polymerases. Trends Microbiol 16: 247–250. doi:10.1016/j.tim.2008.03.008.

9. Peyroche G, Milkereit P, Bischler N, Tschochner H, Schultz P, et al. (2000) The recruitment of RNA polymerase I on rDNA is mediated by the interaction of the A43 subunit with Rrn3. EMBO J 19: 5473–5482. doi:10.1093/emboj/19.20.5473.

10. Yamamoto RT, Nogi Y, Dodd JA, Nomura M (1996) RRN3 gene of Saccharomyces cerevisiae encodes an essential RNA polymerase I transcription factor which interacts with the polymerase independently of DNA template. EMBO J 15: 3964–3973.

11. Blattner C, Jennebach S, Herzog F, Mayer A, Cheung ACM, et al. (2011) Molecular basis of Rrn3-regulated RNA polymerase I initiation and cell growth. Genes Dev 25: 2093–2105. doi:10.1101/gad.17363311.

12. Torreira E, Louro JA, Pazos I, González-Polo N, Gil-Carton D, et al. (2017) The dynamic assembly of distinct RNA polymerase I complexes modulates rDNA transcription. Elife 6. doi:10.7554/eLife.20832.

13. Engel C, Gubbey T, Neyer S, Sainsbury S, Oberthuer C, et al. (2017) Structural basis of RNA polymerase I transcription initiation. Cell 169: 120–131.e22. doi:10.1016/j.cell.2017.03.003.

14. Sadian Y, Tafur L, Kosinski J, Jakobi AJ, Wetzel R, et al. (2017) Structural insights into transcription initiation by yeast RNA polymerase I. EMBO J 36: 2698–2709. doi:10.15252/embj.201796958.

15. Han Y, Yan C, Nguyen THD, Jackobel AJ, Ivanov I, et al. (2017) Structural mechanism of ATP-independent transcription initiation by RNA polymerase I. Elife 6. doi:10.7554/eLife.27414.

16. Keener J, Josaitis CA, Dodd JA, Nomura M (1998) Reconstitution of yeast RNA polymerase I transcription in vitro from purified components. TATA-binding protein is not required for basal transcription. J Biol Chem 273: 33795–33802.

17. Kuhn C-D, Geiger SR, Baumli S, Gartmann M, Gerber J, et al. (2007) Functional architecture of RNA polymerase I. Cell 131: 1260–1272. doi:10.1016/j.cell.2007.10.051.

18. Geiger SR, Lorenzen K, Schreieck A, Hanecker P, Kostrewa D, et al. (2010) RNA polymerase I contains a TFIIF-related DNA-binding subcomplex. Mol Cell 39: 583–

594. doi:10.1016/j.molcel.2010.07.028.

19. Fernández-Tornero C, Böttcher B, Rashid UJ, Steuerwald U, Flörchinger B, et al. (2010) Conformational flexibility of RNA polymerase III during transcriptional elongation. EMBO J 29: 3762–3772. doi:10.1038/emboj.2010.266.

20. Landrieux E, Alic N, Ducrot C, Acker J, Riva M, et al. (2006) A subcomplex of RNA polymerase III subunits involved in transcription termination and reinitiation. EMBO J 25: 118–128. doi:10.1038/sj.emboj.7600915.

21. Wu C-C, Lin Y-C, Chen H-T (2011) The TFIIF-like Rpc37/53 dimer lies at the center of a protein network to connect TFIIIC, Bdp1, and the RNA polymerase III active center. Mol Cell Biol 31: 2715–2728. doi:10.1128/MCB.05151-11.

22. Tafur L, Sadian Y, Hoffmann NA, Jakobi AJ, Wetzel R, et al. (2016) Molecular structures of transcribing RNA polymerase I. Mol Cell 64: 1135–1143. doi:10.1016/j.molcel.2016.11.013.

23. Pilsl M, Crucifix C, Papai G, Krupp F, Steinbauer R, et al. (2016) Structure of the initiation-competent RNA polymerase I and its implication for transcription. Nat Commun 7: 12126. doi:10.1038/ncomms12126.

24. Van Mullem V, Landrieux E, Vandenhaute J, Thuriaux P (2002) Rpa12p, a conserved RNA polymerase I subunit with two functional domains. Mol Microbiol 43: 1105–1113.

25. Gadal O, Mariotte-Labarre S, Chedin S, Quemeneur E, Carles C, et al. (1997) A34.5, a nonessential component of yeast RNA polymerase I, cooperates with subunit A14 and DNA topoisomerase I to produce a functional rRNA synthesis machine. Mol Cell Biol 17: 1787–1795.

26. Beckouet F, Labarre-Mariotte S, Albert B, Imazawa Y, Werner M, et al. (2008) Two RNA polymerase I subunits control the binding and release of Rrn3 during transcription. Mol Cell Biol 28: 1596–1605. doi:10.1128/MCB.01464-07.

27. Liljelund P, Mariotte S, Buhler JM, Sentenac A (1992) Characterization and mutagenesis of the gene encoding the A49 subunit of RNA polymerase A in Saccharomyces cerevisiae. Proc Natl Acad Sci USA 89: 9302–9305.

28. Nogi Y, Yano R, Dodd J, Carles C, Nomura M (1993) Gene RRN4 in Saccharomyces cerevisiae encodes the A12.2 subunit of RNA polymerase I and is essential only at high temperatures. Mol Cell Biol 13: 114–122.

29. Albert B, Léger-Silvestre I, Normand C, Ostermaier MK, Pérez-Fernández J, et al. (2011) RNA polymerase I-specific subunits promote polymerase clustering to enhance the rRNA gene transcription cycle. J Cell Biol 192: 277–293. doi:10.1083/jcb.201006040.

30. Gadal O, Labarre S, Boschiero C, Thuriaux P (2002) Hmo1, an HMG-box protein, belongs to the yeast ribosomal DNA transcription system. EMBO J 21: 5498–5507.

31. Huet J, Buhler JM, Sentenac A, Fromageot P (1975) Dissociation of two polypeptide chains from yeast RNA polymerase A. Proc Natl Acad Sci USA 72: 3034–3038.

32. Viktorovskaya OV, Appling FD, Schneider DA (2011) Yeast transcription elongation factor Spt5 associates with RNA polymerase I and RNA polymerase II directly. J Biol Chem 286: 18825–18833. doi:10.1074/jbc.M110.202119.

33. Decourty L, Saveanu C, Zemam K, Hantraye F, Frachon E, et al. (2008) Linking functionally related genes by sensitive and quantitative characterization of genetic interaction profiles. Proc Natl Acad Sci USA 105: 5821–5826. doi:10.1073/pnas.0710533105.

34. Engel C, Plitzko J, Cramer P (2016) RNA polymerase I-Rrn3 complex at 4.8 Å resolution. Nat Commun 7: 12129. doi:10.1038/ncomms12129.

35. Neyer S, Kunz M, Geiss C, Hantsche M, Hodirnau V-V, et al. (2016) Structure of RNA polymerase I transcribing ribosomal DNA genes. Nature 540: 607–610.

doi:10.1038/nature20561.

36. Sanz-Murillo M, Xu J, Belogurov GA, Calvo O, Gil-Carton D, et al. (2018) Structural basis of RNA polymerase I stalling at UV light-induced DNA damage. Proc Natl Acad Sci USA. doi:10.1073/pnas.1802626115.

37. Cioci F, Vu L, Eliason K, Oakes M, Siddiqi IN, et al. (2003) Silencing in yeast rDNA chromatin: reciprocal relationship in gene expression between RNA polymerase I and II. Mol Cell 12: 135–145.

38. Takeuchi Y, Horiuchi T, Kobayashi T (2003) Transcription-dependent recombination and the role of fork collision in yeast rDNA. Genes Dev 17: 1497–1506. doi:10.1101/gad.1085403.

39. Wery M, Ruidant S, Schillewaert S, Leporé N, Lafontaine DLJ (2009) The nuclear poly(A) polymerase and Exosome cofactor Trf5 is recruited cotranscriptionally to nucleolar surveillance. RNA 15: 406–419. doi:10.1261/rna.1402709.

40. LaCava J, Houseley J, Saveanu C, Petfalski E, Thompson E, et al. (2005) RNA degradation by the exosome is promoted by a nuclear polyadenylation complex. Cell 121: 713–724. doi:10.1016/j.cell.2005.04.029.

41. Houseley J, Tollervey D (2006) Yeast Trf5p is a nuclear poly(A) polymerase. EMBO Rep 7: 205–211. doi:10.1038/sj.embor.7400612.

42. Kos M, Tollervey D (2010) Yeast pre-rRNA processing and modification occur cotranscriptionally. Mol Cell 37: 809–820. doi:10.1016/j.molcel.2010.02.024.

43. Prescott EM, Osheim YN, Jones HS, Alen CM, Roan JG, et al. (2004) Transcriptional termination by RNA polymerase I requires the small subunit Rpa12p. Proc Natl Acad Sci USA 101: 6068–6073. doi:10.1073/pnas.0401393101.

44. Smale ST (2009) Nuclear run-on assay. Cold Spring Harb Protoc 2009: pdb.prot5329. doi:10.1101/pdb.prot5329.

45. Green MH, Buss J, Gariglio P (1975) Activation of nuclear RNA polymerase by sarkosyl. Eur J Biochem 53: 217–225. doi:10.1111/j.1432-1033.1975.tb04060.x.

46. Wery M, Gautier C, Descrimes M, Yoda M, Vennin-Rendos H, et al. (2018) Native elongating transcript sequencing reveals global anti-correlation between sense and antisense nascent transcription in fission yeast. RNA 24: 196–208. doi:10.1261/rna.063446.117.

47. Tafur L, Sadian Y, Hanske J, Wetzel R, Weis F, et al. (2019) The cryo-EM structure of a 12-subunit variant of RNA polymerase I reveals dissociation of the A49-A34.5 heterodimer and rearrangement of subunit A12.2. Elife 8. doi:10.7554/eLife.43204.

48. Fernández-Tornero C (2018) RNA polymerase I activation and hibernation: unique mechanisms for unique genes. Transcription: 1–7. doi:10.1080/21541264.2017.1416267.

49. Malagon F, Kireeva ML, Shafer BK, Lubkowska L, Kashlev M, et al. (2006) Mutations in the Saccharomyces cerevisiae RPB1 gene conferring hypersensitivity to 6-azauracil. Genetics 172: 2201–2209. doi:10.1534/genetics.105.052415.

50. Kaplan CD, Larsson K-M, Kornberg RD (2008) The RNA polymerase II trigger loop functions in substrate selection and is directly targeted by alpha-amanitin. Mol Cell 30: 547–556. doi:10.1016/j.molcel.2008.04.023.

51. Viktorovskaya OV, Engel KL, French SL, Cui P, Vandeventer PJ, et al. (2013) Divergent contributions of conserved active site residues to transcription by eukaryotic RNA polymerases I and II. Cell Rep 4: 974–984. doi:10.1016/j.celrep.2013.07.044.

52. Viktorovskaya OV, Schneider DA (2015) Functional divergence of eukaryotic RNA polymerases: unique properties of RNA polymerase I suit its cellular role. Gene 556: 19–26. doi:10.1016/j.gene.2014.10.035.

53. Appling FD, Schneider DA, Lucius AL (2017) Multisubunit RNA polymerase cleavage

factors modulate the kinetics and energetics of nucleotide incorporation: an RNA polymerase I case study. Biochemistry 56: 5654–5662. doi:10.1021/acs.biochem.7b00370.

54. Schiestl RH, Gietz RD (1989) High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. Curr Genet 16: 339–346.

55. Sambrook J, Fritsch EF, Maniatis T (1989) Molecular cloning: a laboratory manual. Molecular cloning: a laboratory manual. Available: https://www.cabdirect.org/cabdirect/abstract/19901616061.

56. Sherman F, Fink GR, Hicks JBN (1986) Methods in Yeast Genetics. A Laboratory Course Manual. New York: Cold Spring Harbor Laboratory.

57. Decourty L, Saveanu C, Zemam K, Hantraye F, Frachon E, et al. (2008) Linking functionally related genes by sensitive and quantitative characterization of genetic interaction profiles. Proc Natl Acad Sci U S A 105: 5821–5826.

58. Kolesnikov N, Hastings E, Keays M, Melnichuk O, Tang YA, et al. (2015) ArrayExpress update--simplifying data submissions. Nucleic Acids Res 43: D1113-6. doi:10.1093/nar/gku1057.

59. Hermann-Le Denmat S, Werner M, Sentenac A, Thuriaux P (1994) Suppression of yeast RNA polymerase III mutations by FHL1, a gene coding for a fork head protein involved in rRNA processing. Mol Cell Biol 14: 2905–2913.

60. Beltrame M, Tollervey D (1992) Identification and functional analysis of two U3 binding sites on yeast pre-ribosomal RNA. EMBO J 11: 1531–1542.

61. Osheim YN, French SL, Sikes ML, Beyer AL (2009) Electron microscope visualization of RNA transcription and processing in Saccharomyces cerevisiae by Miller chromatin spreading. Methods Mol Biol 464: 55–69. doi:10.1007/978-1-60327-461-6_4.

62. Tschochner H (1996) A novel RNA polymerase I-dependent RNase activity that shortens nascent transcripts from the 3' end. Proc Natl Acad Sci USA 93: 12914–12919.

63. Kwapisz M, Ruault M, van Dijk E, Gourvennec S, Descrimes M, et al. (2015) Expression of Subtelomeric lncRNAs Links Telomeres Dynamics to RNA Decay in S. cerevisiae. Noncoding RNA 1: 94–126. doi:10.3390/ncrna1020094.

64. Hage AE, Houseley J (2013) Resolution of budding yeast chromosomes using pulsed-field gel electrophoresis. Methods Mol Biol 1054: 195–207. doi:10.1007/978-1-62703-565-1_13.

Figure legends

Figure 1. Alleles of *RPA190* and *RPA135* suppress the growth defect of the *rpa49Δ*

mutant at various levels. (A) The *SGR1* mutant restores growth of the *rpa49Δ* mutant. Ten-

fold serial dilutions of wild-type (WT), *rpa49Δ* single mutant, *SGR1* single mutant, and

SGR1/rpa49Δ double mutant strains were spotted on rich media to assess growth at 30 and

25°C for three days. (B) Ten-fold dilutions of WT and *rpa49Δ* compared to *rpa49Δ* carrying

various plasmids: pGL190_3 (*RPA190-E1274K*), pGL190_11 (*RPA190-C1493R*),

pGL190_23 (*RPA190-L1262P*), pGL135_6prim (*RPA135-R379G*), pGL135_54 (*RPA135-*

Y252H), or pGL135_33 (*RPA135-F301S*). Growth was evaluated after three days at 25°C.

The strains and plasmids used are listed in S2-S3 Tables, respectively.

Figure 2. Mapping of the modified residues in Rpa190, Rpa135 on the structure of Pol I

and isolation of Rpa12 alleles. (A) Two different views of the initially transcribing complex

model and its 14 different subunits (PDB 5W66[15]). (B) Most mutated suppressor residues

are clustered at the interface between the jaw (Rpa190, blue) and lobe (Rpa135, salmon)

modules of Pol I. Note that residues 46- 51 in Rpa12 (yellow) are part of this interface. (C)

Zoom views of the areas containing the modified residues (Rpa190-N863, -S1259, -L1262, -

E1274, -C1493; Rpa135-Y252, D299, S300, F301, R305; and Rpa12-S6, T49) shown in

panel B. The figure was prepared with Pymol using the crystal structure of Pol I PDB 4C3I

[4]). (D) Ten-fold dilutions of the *rpa49Δ* mutant carrying various plasmids: an empty

pRS316 plasmid (-) YCp50-26 bearing *RPA49* (*RPA49*), pRS316-A12-S6L (*RPA12-S6L),* or

pRS316-A12-T49A (*RPA12-T49A)*. Growth was evaluated after three days at 25°C or two

days at 30°C.

Figure 3. *RPA135-F301S* and *RPA12-S6L* alleles restore growth and rRNA synthesis,

and modulate Pol I occupancy of rDNA genes in the absence of Rpa49Ct. (A) Doubling

times of WT, *rpa49ΔCt*, *RPA135-F301S*, *rpa49ΔCt/RPA135-F301S* double mutant, *RPA12-*

S6L, and the *rpa49ΔCt/RPA12-S6L* double mutant in a low rDNA copy number background

(see S2 Table). (B) *In vivo* labelling of newly synthesized RNAs. WT (lane 1), *rpa49ΔCt*

(lanes 2-4), *RPA135-F301S* (lane 5), the *rpa49ΔCt/RPA135-F301S* double mutant (lane 6),

RPA12-S6L (Lane 7), the *rpa49ΔCt/RPA12-S6L* double mutant (lane 8) were grown to an

OD_{600} of 0.8. Cells were then pulse-labeled with $[8-^{3}H]$ adenine for 2 min. Samples were

collected, and total RNA extracted and separated by gel electrophoresis. (C) Representative

Miller spreads of WT, *rpa49ΔCt*, *RPA135-F301S*, *rpa49ΔCt/RPA135-F301S*, and

rpa49ΔCt/RPA12-S6L double mutant. Panels on the right of each micrograph show

interpretive tracing of the genes. Polymerases that appear on the gene are shown on the

tracing by black dots. The number of polymerases counted on the genes is indicated below. N

represents the number of individual spread genes used for quantification (see Materials and

Methods). Scale bar = 200 nm. (D) ChIP analysis of Pol I occupancy at rDNA. Strains

bearing WT Pol I, expressing Rpa49 lacking its C-terminal part (*rpa49ΔCt*), bearing

suppressor mutation (*RPA135-F301S*) or double mutant, were subjected to ChIP experiments

using TAP tagged Rpa135, as described in Materials and Methods. Experiments were

reproduced three times, representative Pol I occupancy relative to WT level are shown.

Position of qPCR amplicons are depicted on rDNA unit. ** Marks significantly different

values (p-values > 0.01) in student's test of *rpa49ΔCt* with WT, *RPA135-F301S* and double

rpa49ΔCt RPA135-F301S.

Figure 4. Rpa14, Rpa34, and the DNA mimicking loop of Rpa190 are not required for

suppression. Deletion of the DNA mimicking loop of Rpa190 (A) or *RPA34* (B) does not

modulate the suppression activity of *RPA135-F301S*. (C) *RPA135-F301S* suppresses the

synthetic lethality between *rpa14Δ* and *rpa49Δ*. Ten-fold serial dilutions were performed and

growth evaluated after three days at 25°C.

Figure 5. *RPA12* alleles can modulate the *rpa49Δ*-associated growth defect. (A) Growth of

the double mutants: *rpa49Δ rpa12ΔCt,* or *rpa49Δ* combined with full depletion of *rpa12*.

Depletion of Rpa12 was achieved using a *pGAL-RPA12* construct on glucose containing

medium (strain OGT30-1c). Ten-fold serial dilutions of OGT30-1c bearing pRS316-A12

(WT), pRS316-A12-DCt expressing Rpa12 bearing a C-terminal deletion of residues 65-125

(*rpa12ΔCt*), or an empty plasmid pRS316 (-*rpa12*) were seeded onto media. The growth of

rpa49Δ combined with *RPA12* depletion was tested using strain OGT30-3c bearing pRS316-

A12 (*rpa49Δ*), pRS316-A12-DCt (*rpa12ΔCt rpa49Δ*), or an empty plasmid pRS316 (-*rpa12*

rpa49Δ). Growth was assessed after four days at 25°C or 30°C. (B) The C terminus of Rpa12

is not required for suppression. Ten-fold serial dilutions of OGT30-1c (*RPA49*-WT), bearing

pRS316-A12 (WT) or pRS316-A12-DCter (*rpa12ΔCt*), and OGT30-3c (*rpa49Δ*), bearing

pRS316-A12 (WT), pRS316-A12-DCter (*rpa12ΔCt*), pRS316-A12-S6L (*RPA12-S6L*), or

pTD10 (*RPA12-S6L-ΔCt*) were seeded onto media. Growth was assessed after four days at

25°C. (C) Suppression activity of *RPA135-F301S* is abolished in the absence of Rpa12.

RPA12, under a regulatable promoter (pGAL) was either expressed (+*RPA12; left panel*) on

galactose containing medium or repressed (-*RPA12; right panel*) on glucose containing

medium. Ten-fold serial dilutions of *RPA49*, *rpa49Δ*, *RPA49*, *RPA135-F301S*, or *rpa49Δ*

RPA135-F301S were grown at 25°C. Depletion of *RPA12* abolishes the suppression activity

of *RPA135-F301S* (compare the left to the middle and right panels). Extended incubation

(right panel, 10 days) was used to detect growth of the double mutant -*RPA12 rpa49Δ* on

plates.

Figure 6. *In vitro* transcription assays of WT Pol I and Pol I mutants. WT, Pol A* (lacking

Rpa34 and Rpa49), Pol I bearing Rpa135-F301S, and Pol A* bearing Rpa135-F301S were

affinity-purified and 5 nM of each enzyme was used in either promoter dependent (A) or

tailed template assay (B). Promoter-dependent assays were performed in the presence of 70

nM Rrn3 and 20 nM CF. Radiolabelled transcripts were separated on a denaturing

polyacrylamide/urea gel and detected using a PhosphorImager. Note that upper radiolabeled

bands in the experiment analyzing promoter-dependent transcription are due to nonspecific

background labelling. Experiments were reproduced at least three times, with different Pol I

concentration; representative experiments is shown.

Figure 7. *RPA135-F301S* led to over-production of rRNA *in vivo.* (A) WT, *RPA135-F301S*, *rrp6Δ*

and *RPA135-F301SΔ rrp6Δ* strains were grown to mid-log phase in glucose containing media. Cell

samples were collected and total RNAs were extracted, separated by gel electrophoresis and

transferred to a nylon membrane. The accumulation of the different (pre-) rRNAs was then analyzed

by northern blot using different probes (see Materials and Methods). (B) *In vivo* labelling of newly-

synthesized RNAs. WT, *RPA135-F301S*, *rrp6Δ* and *RPA135-F301SΔ rrp6Δ* strains were grown to an

OD_{600} of 0.8. Cells were then pulse-labeled with [8-^{3}H] adenine for 40 seconds. Samples were

collected, and total RNA extracted and separated by gel electrophoresis. Newly-synthetized RNA are

revealed by autoradiography, loading control was performed by northern blot (*PGK1* mRNA probe –

1831) on the same membrane. (C) High-resolution transcriptional run-on (TRO) analysis of WT,

RPA135-F301S, *rrp6Δ* and *RPA135-F301SΔ rrp6Δ* strains. Nascent transcripts were labelled, and

revealed using antisens oligonucleotides immobilized on slot-blot as described in Materials and

Methods. Each experiment was performed twice; a representative example is shown in the lower left

panel. NTS2, Pol I (mean of 5'ETS, 18S.2, 25S.1, 3' ETS) are quantified relative to 5S signal in the

lower right panel. Yeast rDNA unit is represented in the upper panel, with the position of the

corresponding antisense oligonucleotides used.

Figure 8. Schematic representation of Pol I. (A) Free monomeric Pol I with mobile

Rpa49Ct and linker. (B) Initially transcribing complex (ITC) upon insertion of melted DNA

in the presence of Rpa49 (purple). Rpa49Ct interacts with upstream DNA and the Rpa49-

linker is folded, closing the cleft (black anchor). Movements of Rpa12 and the jaw with

respect to the lobe are indicated with arrows. (C) Pol I lacking Rpa49 is likely defective in

stabilizing the closed conformation in the DNA-binding cleft, resulting in a looser gripping of

DNA inside the cleft (red asterisks). (D) Suppressor mutations (green) facilitate movement of

the jaw/lobe interface and gripping of the DNA by the Pol I enzyme (red arrow), in the

absence of Rpa49. (E) Combination of the presence of Rpa49 and a suppressor mutation

(green dot) results in a super-active Pol I compared to the WT enzyme.

Supporting information captions

S1 Fig: Genetic interaction mapping (GIM) to identify suppressor mutations in SGR1

and SGR2. (A) Schematic representation of GIM interaction assay. Enrichment ratios

between control and SGR are used as read-out to map genetic interactions. (B) Relative

enrichment of each barcode (black cross) along chromosome (in red) is used to map genetic

linkage. Green curve represents mean in a sliding windows of 20 barcodes. Local maximum

in such curve was used to identify locus of interest: RPA49 as positive control (upper panel,

chr 14); RPA135 in SGR1 (middle panel, chr 16) ; RPA190 in SGR2 (lower panel, chr 15).

S2 Fig: Mapping of the mutated residues in Rpa135. (A) Structural domains of Rpa135 are

depicted [4,5], including the lobe domain in which mutations are clustered. (B) Sequences

alignment of positions 251-310 of Rpa135 from *S. cerevisiae* (NP_015335.1), compared with

S. pombe (NP_595819.2), *H. sapiens* (NP_061887.2) , *M. musculus* (NP_033112.2), *D. rerio*

(NP_956812.2), *D. melanogaster* (NP_476708.1) and *C. elegans* (NP_492476.1). Color code

from light to dark blue indicates residue conservations. Mutated residues at position 252 (Y to

H), 299 (D to G), 300 (S to F), 301 (F to S or L) and 305 (R to L) are depicted in red.

S3 Fig: Analysis of the rDNA size by pulse field gel electrophoresis. Chromosomes from

indicated strains (same as in Fig. 3D, two clones) were separated on the agarose gel and

analysed by Southern blot using rDNA-specific probe.

S4 Fig: Promoter dependent *in vitro* transcription assays of Pol A* (lacking Rpa34 and

Rpa49) and Pol A* bearing Rpa135-F301S are complemented with recombinant A34.5/A49

heterodimer. Promoter-dependent assays were performed as in fig. 6, with recombinant

A49/A34.5 protein, described in [23].

S5 Fig: Ten-fold serial dilutions of *WT, RPA135-F301S, rrp6Δ* and *RPA135-F301S -*

rrp6Δ grown at 25°C for 3 days.

S1 Table: List of 24 individual suppressor mutations of the growth defect of *rpa49Δ* strain in

the Rpa190, Rpa135, and Rpa12 subunits.

S2 Table: Yeast strains used in this study.

S3 Table: Plasmids used in this study.

S4 Table: Oligonucleotides used in this study.

S1 Movie: Conformational changes in Pol I upon initiation. The movie starts with the closed

cleft conformation (PDB 5W66 [15]), in which melted DNA occupies the cleft and then

changes to the intermediate cleft conformation observed in monomeric Pol I (PDB 5M3M

[35]). Relevant structural regions have been variously colored and labeled, and residues

mutated in this report are shown in red.

A

B

Figure 1

Figure 2

A

Strain	Doubling time (min)
WT	99 +/- 2
rpa49ΔCt	292 +/- 11
RPA135-F301S	98 +/- 2
rpa49ΔCt RPA135-F301S	118 +/-3
RPA12-S6L	100 +/-1
rpa49ΔCt RPA12-S6L	173 +/- 12

B

WT rpa49ΔCt RPA135-F301S RPA135-F301S rpa49ΔCt RPA12-S6L RPA12-S6L rpa49ΔCt

35S
27S
20S

1 2 3 4 5 6 7 8

C

WT rpa49ΔCt rpa49ΔCt RPA135-F301S RPA135-F301S rpa49ΔCt RPA12-S6L

91 +/- 13   21 +/- 5   46 +/- 10   93 +/- 17   43 +/- 10
N=42   N=31   N=61   N=36   N=49

D

Pol I

NTS2 PROM 5'ETS 18S ITS1 5.8S ITS2 25S 3'ETS NTS1

5'ETS 18S 25S

% IP / TOT/ WT

5'ETS

☐ WT
☐ RPA135-F301S
■ rpa49ΔCt
▨ rpa49ΔCt RPA135-F301S

Figure 3

Figure 4

Figure 5

A

Promoter-dependent assay

WT   Pol A*   Pol I -F301S   Pol A* - F301S

← Promoter transcript

1   2   3   4

Pol I

Promoter

B

Tailed template assay

WT   Pol A*   Pol I -F301S   Pol A* - F301S

← Tailed template transcript

1   2   3   4

Pol I

Figure 6

Figure 7

Figure 8