Effective population sizes and adaptive genetic variation in a captive bird population

Giridhar Athrey[1,2‡], Nikolas Faust[1], Anne-Sophie Charlotte Hieke[1], I. Lehr Brisbin[3]

[1]Department of Poultry Science, Texas A&M University, 2472 TAMU, College Station, TX,

77843.

[2]Faculty of Ecology and Evolutionary Biology, Texas A&M University, 2472 TAMU, College

Station, TX, 77843.

[3]University of Georgia/Savannah River Ecology Lab, Drawer E, Aiken, SC, 29802.

[‡]Corresponding author: Giridhar Athrey, Department of Poultry Science, Texas A&M

University, 2472 TAMU, College Station, TX, 77843. Email: giri.athrey@tamu.edu

Abstract

Captive populations are considered a key component of *ex situ* conservation programs. Research

on multiple taxa have shown the differential success of maintaining demographic versus genetic

stability and viability in captive populations. In typical captive populations, usually founded by

few or related individuals, genetic diversity can be lost and inbreeding can accumulate rapidly,

calling into question their ultimate utility for release into the wild. Furthermore, domestication

selection for survival in captive conditions is another concern. Therefore, it is crucial to

understand the dynamics of population sizes, particularly the effective population size, and

genetic diversity at non-neutral, at adaptive loci in captive populations.

In this study, we assessed effective population sizes and genetic variation at both neutral

microsatellite markers, as well as SNP variants from the MHC-B locus of a captive Red

Junglefowl population. This population is a rare instance of a well-documented history in

captivity, and also following a realistic chain-of-custody, unlike captive lab populations. Our

analysis of 27 individuals that comprise the entirety of one captive population. Our results show

very low neutral and adaptive genetic variation, as well as low effective sizes, which are

surprising in spite of the known demographic history. Finally, our study also shows the divergent

impacts of small effective size and inbreeding in captive populations on microsatellite versus

adaptive genetic variation in the MHC-B locus. Our study provides insights into the difficulties

of maintaining adaptive genetic variation in small captive populations.

Introduction

Genetics of captive populations

The management of demographic security and genetic diversity are among the central

considerations in conservation (1986a, b). Captive populations have long played a key role in

conservation as a management tool to ensure demographic security (Hedrick 1992). The earliest

work in conservation of endangered species focused on breeding in captivity to increase

population sizes and intended for eventual release back into the wild. However demographic

security and the maintenance of genetic diversity can sometimes be at odds, as genetic diversity

is derived from the effective population size (N_e). One issue in captive populations is the

differential reproductive success of some breeding pairs that are better suited to captive

conditions, leading to selection for survival in captive environments. This phenomenon can

further reduce the N_e due to founder effects (Nei *et al.* 1975; Newman & Pilson 1997). From the

perspective of the reintroduction of the offspring of such captive breeders back into the wild, this

may be a less than ideal situation, as these captive-adapted offspring may not carry the adaptive

genetic variation required for survival in their natural habitat. Historically, captive individuals

were raised for the genetic support of threatened wild populations, the genetic compatibility

(prevent outbreeding depression) and genetic variation (ability to adapt to natural environment).

However, these objectives can be materialized only if genetic diversity can be sustained, and if

genetic drift and inbreeding can be limited.

The relationship between population bottlenecks and reduction in effective population sizes and

genetic diversity has been reported from various vertebrate taxa (Athrey *et al.* 2012; Athrey *et al.*

2011; Emami-Khoyi *et al.* 2017; Leberg 1992; Lovatt & Hoelzel 2014). The loss of genetic

diversity due to drift and inbreeding in captive populations has also been well

documented (Willoughby *et al.* 2015; Willoughby *et al.* 2017) and the drawbacks of using

closed captive breeding populations for reintroduction have been described (Lynch & O'Hely

2001). These range from the rapid decline of fitness upon reintroduction (Araki *et al.* 2007)to a

range of consequences for genetic diversity (Witzenberger & Hochkirch 2011). Witzenberger et

al. (2011) also suggest the need for assessment of genetic diversity before *ex situ* conservation

programs. The consensus of these studies suggest that considerations of genetic diversity of

captive populations may be as crucial as maintaining demographic stability, but is perhaps much

more challenging than the maintenance of demographic stability, because non-relative matings

are not entirely avoidable. While some studies have focused on assessing genetic diversity, in

many instances the history of the founding individuals is not well known, or the studbook data

may be incomplete (Witzenberger & Hochkirch 2011). Additionally, inbreeding estimates may

assume that founders are unrelated (Ruiz-Lopez *et al.* 2009). For example, Alcaide et al.

(2009)compared wild versus captive and reintroduced Kestrel, but in that instance, it is not clear

how many generations the kestrels had bred in captivity. In another case study using

experimental captive breeding, Willoughby et al. (Willoughby *et al.* 2017) report on white-

footed mice populations that were bred in captivity for 20 generations, and were able to observe

mitigating effects on genetic diversity using a very deliberate mating protocol (mean-kinship

method). However, in many instances, there may be no choice available as to the source of

individuals obtained and used in captive breeding programs. Furthermore, in extreme cases,

small numbers of breeding individuals may have to be maintained in captivity for decades until

suitable habitat and molecular/technological tools to enable ex situ applications become

applicable. From that perspective, it is essential to have realistic estimates of how much genetic

diversity is maintained, and whether estimates of genetic diversity at neutral markers are

informative about variation at loci crucial for survival in the wild.

In this study, we used a captive population of the Red Junglefowl (*Gallus gallus murghi*) to

characterize the impacts of un-managed captive breeding over a period of 57 years (38-57

generations). The study population called the Richardson Strain (RRJF) has a well-documented

history since its import to the United States in 1961. We investigated the consequences of captive

breeding on genetic diversity at neutral and adaptive genetic markers experienced by this

population over the last 50 years.

Demographic history of Richardson's Red Junglefowl (RRJF)

The study population of Red Junglefowl was initially imported from Northern India in 1961 as

part of the U.S. Fish and Wildlife's Foreign Game Introduction Program (FGIP) and documented

in USFWS Special Reports (Bump 1964). More recently, the history of the population in the

United States was documented in detail in the graduate thesis of Tomas Condon (Condon 2012).

Briefly, the birds that were imported to the United States were captured in the Indian states of

Bihar and Uttar Pradesh (now Uttarakhand), between 1959-1961. A total of 119 survived the

capture and shipping to the United States. These birds were placed in hatcheries across nine

different states, and over the next decade, approximately 10,000 birds were released into U.S.

habitats for hunting, until the FGIP program ended. Over the next four decades, almost all the

descendant RJF a) failed to survive in their new habitats, b) became incorporated into private

aviaries, or c) remained in small breeding populations maintained by dedicated researchers and

aviculturists. All the surviving RJF of known ancestry (estimated around 100-150) were derived

from a small number of birds. These birds have a documented chain of custody. In 1969, I. Lehr

Brisbin (of the Savannah River Ecology Lab) received five birds from the South Carolina

Department of Wildlife Resources, to which 26 were added in 1970. By 1971, this flock

numbered only eight adult birds (four of each sex). While more hatchlings were added to this in

mid-1971, the population remained at under 15 individuals, until they were transferred to Mr.

Isaac Richardson of Alabama. Mr. Richardson started his colony with 12 individuals (8 males).

Between 1972-2010, the colony was sustained by Mr. Richardson, hence giving this population

the name Richardson's Red Junglefowl (RRJF). Over this time, the population expanded and was

distributed to a few others, but at any time, the breeding colony never exceeded 20 individuals

(Condon 2012). Based on behavioral and morphological studies of Red Junglefowl, it has been

argued that the RRJF represent a pure and unique population of Red Junglefowl (Brisbin &

Peterson 2007), whose preservation is critical in the face of hybridization with domestic chickens

in the native range of Junglefowl (Brisbin 1995; Brisbin & Peterson 2007).

The research colony maintained at Texas A&M University (TAMU) is one of the three known

remaining populations that trace their ancestry back to the survivors of Mr. Richardson's flock.

The TAMU colony was established with two males and three females and expanded to 27

individuals at the time of the study. All available individuals were included in this present study.

Genetic diversity and effective population sizes

As the management of genetic diversity is one of the primary intentions of captive breeding

programs, it is important to assess the consequences of captive breeding over multiple

generations on genetic diversity. Other authors have shown reductions in genetic diversity in

captive populations, and these typically result from differential survival and increased genetic

drift, but also a form of (unintentional) domestication selection (Briscoe *et al.* 1992;

Montgomery *et al.* 2010). In these instances, it is important to characterize what the effective

population size is, as this one parameter can drive various other genetic parameter estimates. The

effective population size (N_e) can be defined as the size of an ideal population that experiences

the same amount of genetic drift or inbreeding as the actual population (Wright 1931, 1948).

While a number of approaches have been developed to estimate N_e - for example, temporal

estimates based on two or more samples, methods based on linkage disequilibrium, and

coalescent approaches (Wang *et al.* 2016), due to our access to a single point in time, as well as

our interest in understanding recent effective population size, we focused only on methods for

contemporary or recent Ne estimates. In this study, we evaluated genetic diversity, estimated the

effective population size (Ne), and analyzed linkage disequilibrium using two types of markers -

namely microsatellite markers and SNP markers.

Materials and methods

Study Population

A total of 27 birds bred in captivity at the Texas A&M Poultry Research Center were genotyped

in this study. The population represents three generations. The parental cohort was comprised of

2 males and three females in January 2016 and had expanded to 27 individuals by the second

half of 2017. At the time of sampling, the population included original parents (used to establish

colony), in addition to two subsequent generations that were hatched in Texas. All individuals

available were included in this study.

Molecular Markers

Venal blood was sampled from all individuals by puncturing the brachial vein and stored in

Longmire Buffer until further processing. Genomic DNA was isolated from whole blood using

the DNeasy Blood and Tissue kit (Qiagen Inc). For microsatellite genotyping, a total of 18

microsatellite markers were randomly selected (from Tadano *et al.*(2007)) and screened for the

presence of polymorphic loci. Of the screened loci, three loci were monomorphic, or were not

amplified by PCR consistently and were not used further (full list and primer sequences

presented in S1). To generate genotype data from microsatellite loci, we used the fluorescently

labeled M13 method (Schuelke 2000), which allows greater flexibility of fluorescent dye tagging

for multiplexing. PCR thermoprofile was optimized based on published T_m for each primer pair

(and considering M13 sequence), and amplification was performed in a 25µl reaction using NEB

Taq Polymerase with reaction buffer (New England Biolabs) and NEB dNTPs (New England

Biolabs).

Multiplexed samples were prepared for fragment analysis length analysis on the ABI 3730

capillary analyzer, and sent to the DNA sequencing facility on Science Hill (Yale University,

New Haven, CT) for fragment analysis. For SNP analyses, equimolar DNA isolates were

submitted to the molecular genetics lab at Hy-Line International for genotyping using a custom

SNP panel (Fulton *et al.* 2016; Nguyen-Phuc *et al.* 2016). The SNP panel was developed as an

assay for the Major Histocompatibility Complex - B locus (MHC-B), based on the

KASP^{TM} chemistry (LGC group, UK) where each allele is identified by its fluorescent label

(VIC or FAM). A total of 90 known SNP loci are genotyped using this panel and covers the

region between basepairs 30,189 to 240,933 on chromosome 16. While microsatellite loci came

from multiple chromosomes, all the SNP loci are from a single gene-rich microchromosome, and

as is typical for SNP datasets, the markers represent a mix of intronic/intergenic (putatively

neutral) and coding regions.

Genetic Analyses

The raw data was downloaded into the software Geneious (Biomatters, New Zealand) and

analyzed using the microsatellite plugin. Allele bins, based on peak data from across all samples,

were created for each locus. Following this, every individual biological sample was processed

through the peak calling step. Once the automatic peak calls were obtained, every allele call was

manually verified to check for errors. The final allele calls were exported into the MS Excel-

based tool, GENALEX v6.5 (Peakall & Smouse 2012). The data were then exported to the

GENEPOP format for estimation of the effective population size. The SNP calls were generated

(by Hy-Line International) using the Kraken software (LGC group, UK).   IUPAC base calls

were first re-coded manually into 2-letter genotypes, and then imported into GENALEX and

recoded into a numeric format for further analysis.

Estimation of expected and unbiased expected heterozygosity (H_e and uH_e), and measures of the

inbreeding coefficient F_{IS} were both calculated within GENALEX, based on 999 permutations of

the data. These parameters were estimated for both the microsatellite and SNP data

independently.

Effective Population Size

Next, to estimate the effective population size of this captive-bred population, we used three

different estimators that are known to estimate contemporary or recent effective population sizes

- namely the heterozygosity excess (HE) method (Zhdanova & Pudovkin 2008), the linkage

disequilibrium (LD) method (Waples & Do 2008)and the molecular coancestry (MC)

method (Nomura 2008). All three N_e estimates were generated using the software program

NeEstimator, version 2 (Do *et al.* 2014). These estimators are expected to estimate the

population size based on shared alleles (molecular coancestry) or the signal of genetic drift -

either due to allele frequency differences among parents (Heterozygote excess) or due to the

linkage disequilibrium among markers. In this case, we also know which assumptions were

violated with some certainty. For example, this captive population has been closed to

immigration and not subject to selection (except unintentional domestication selection in

captivity), but the assumption of random mating is likely to be violated, as we expect this

population to have an increased frequency of mating among relatives over the last five decades.

While it is known that most real natural populations may violate one or more assumptions of an

ideal population, the departure from random mating is expected to result in underestimates of

N_e (Waples *et al.* 2014). Similarly, the molecular coancestry estimate is expected to be biased

downward in inbred populations (Nomura 2008). Considering the differential consequences of

these assumptions for our captive population, we estimated N_e using the three methods

mentioned above. The analyses were carried out separately for the microsatellite data, and for the

SNP data. Furthermore, we were particularly interested in determining the N_e as obtained from

markers from a single chromosome. By definition, a chromosome is a single linkage group, and

linkage of loci over a single chromosome can be expected to be stabler over evolutionary time,

compared to markers sampled from across the genome. However, estimates of contemporary N_e

are expected to be biased downward by the length of the chromosome, and also the number of

chromosomes (Waples *et al.* 2016) primarily due to physically linked loci, and recombination

frequencies being lower on short chromosomes. To our knowledge, this is the first study where a

high density of markers from a single chromosome was used to explore the population history.

Haplotype and Linkage Analyses

To understand the effects of captive breeding on the number and structure of unique haplotypes

in the captive population, we first phased the SNP genotype data to identify the unique

haplotypes present in the population. We used the program PHASE v2 (Stephens *et al.* 2001) to

phase the SNP loci into unique haplotype sequences. We performed phasing and estimation of

haplotype frequencies from five replicate runs of the program PHASE. Each replicate run started

with a different seed and was comprised of 1000 iterations, with a thinning interval of five and a

burn-in interval of 100. Following the completion of these runs, the number of unique haplotypes

and their frequencies were recorded and checked across replicate runs. As the number of

haplotypes and their frequencies was very consistent across runs, we determined that increasing

the run length was not necessary (Stephens *et al.* 2001).The list of unique haplotypes was then

used to construct a haplotype network using the 'APE' package on the R statistical

platform (Paradis *et al.* 2017). This approach constructs a matrix of Kimura-2 parameter genetic

distance (Kimura 1980) based on nucleotide differences among the haplotypes and plots a

network representing differences among haplotypes.

Finally, to examine linkage structure across chromosome 16 based on the 90 SNP loci, we used

the program HaploView (Barrett *et al.* 2005).This program summarizes estimated Minor Allele

Frequency (MAF) for each locus and generates haplotype and linkage information for locus

pairs, which are then represented in a graphical format. Higher values of linkage disequilibrium

(LD) suggest stronger association among loci, and in turn, showing low recombination between

loci.

Results

Genetic Diversity

For the microsatellite data, 93.3% of the loci were polymorphic, and the three estimates of

heterozygosity (observed, expected, and unbiased) based on microsatellite markers were very

similar to each other, ranging from 0.359 to 0.371 to 0.379, respectively. For the SNP dataset,

only 37.8% of the loci were polymorphic, which translated into heterozygosity estimates of

0.107, 0.096, and 0.98 (observed, estimated, and unbiased, respectively, Table 1). Estimates of F,

the fixation index/inbreeding coefficient, were also divergent between the microsatellite (0.065)

and SNP datasets (-0.137). Positive values of F are typically indicative of more significant

inbreeding than expected, whereas negative values suggest more outbreeding than expected. This

latter result is potentially a consequence of the low observed and expected heterozygosities for

the SNP dataset, arising from the small proportion of SNP loci that were polymorphic.

Effective Population Size estimates

The N_e estimates were in the low single digits for the microsatellite marker dataset, except for

the heterozygote excess method. For the single chromosome SNP dataset, both the LD and MC

estimates were 1 or lower with narrow 95% intervals (Table 2), whereas the H_E estimator was in

the low single digits (N_{eHE}=4.8, 95% CI=3.1-11). If these estimates are compared against the

total local population (Texas A&M), the N_e/N_c ratios range from about 0.14-0.27 for

microsatellite markers, whereas they range from 0.007-0.17 for the SNP based estimates.

Differences between neutral microsatellite markers and SNP markers

Given the known history of the study population, the low N_e estimates were not unexpected, but

the differences between the microsatellite markers from multiple locations in the genome, and

the single chromosome SNP markers were notable, and reveal different population histories. Both the MC and the LD methods estimated N_e to be equal to or lower than one, based on the SNP dataset. Whether these differences are the result of different recombination frequencies for marker types or actual biological lineages for unique haplotypes is challenging to differentiate. However, identification of unique haplotypes (next section) partly answers this question.

Unique Haplotypes and Linkage

Analysis of linkage disequilibrium based on the SNP loci on chromosome 16 revealed very high D' values across all pairwise comparisons. Linkage disequilibrium (D') was estimated to have a value of 1, indicating little or no recombination between marker pairs (Figure 1). Such high values indicate that as a consequence of captive breeding, individuals in this population have inherited virtually identical chromosomes (chromosome 16 at least), a phenomenon that would be indistinguishable from the absence of recombination.

Analysis of haplotypes in the study population using the program PHASE revealed five unique haplotypes (Figure 2), and of these only three were found at frequencies higher than 10%, with the most common haplotype represented in 70% of individuals (haplotype 1). The next two most common haplotypes were found at approximately 14% frequency each. These haplotypes suggest at best three unique haplotype lineages contributing to the population and presumably do not reflect novel haplotypes emerged by mutations since the initial population contraction.

Discussion

Genetic Diversity

We found relatively low genetic diversity based on heterozygosity, among both marker types,

but estimates of heterozygosity were higher using microsatellite markers compared to SNP loci.

This is not a surprising result, as microsatellites may not be highly correlated with underlying

genomic diversity (Vali *et al.* 2008), for example, due to ascertainment bias for polymorphic

loci. Tokarska et al. (2009) have also shown that in species or populations with low genetic

diversity, heterozygous SNP loci may be more informative about population history and

structure. In our case, the SNP marker panel included loci from both coding and noncoding

regions, but all these markers came from a single chromosome. While it is difficult in our study

to determine how the SNP loci may be representative of whole genome diversity, there is no

particular reason to believe that chromosome 16 would be experiencing selection or drift in ways

that are inconsistent with the rest of the genome. Therefore, we believe that the measures of

genomic diversity based on the SNP panel to be more informative about genome-wide patterns

of diversity, while the microsatellite-based estimates are consistent with and relevant to our

understanding of neutral diversity.

The second takeaway of such a comparison between microsatellite markers and SNP markers is

the potential implications of such disparity for other endangered or threatened species of

conservation interest. Several studies have used microsatellite markers to assess genetic variation

following population contractions (Athrey *et al.* 2011; Bouzat *et al.* 1998; Hammerly *et al.* 2013;

Hoelzel *et al.* 2002; Johnson & Dunn 2006; Miao *et al.* 2016; Wisely *et al.* 2002). If

microsatellite-based measures of genetic diversity are higher than SNP-based whole-genome

genetic diversity**,** as observed in the current study, it would be a potentially worrying trend from

the perspective of measuring the genetic viability of wild populations. Due to the relatively low

cost and accessibility of microsatellite loci, the usage of these markers has become ubiquitous in

population and conservation genetics, and in the assessment of genetic diversity. If high

microsatellite variability does not correlate positively with genome-wide diversity, then

management decisions need to consider the potential limitations of such data.

Effective Size estimates

We found low (single digit) N_e estimates across all estimators. On the one hand this is consistent

with the known demographic histories, but on the other hand, this may also be explained by

biases arising from sample size or other assumptions. We expect the N_e estimates to be biased

downward due to the small sample size in this study, but these values are not unusual in

endangered populations. Several of the N_e estimators expect biased estimates at sample sizes

under N=50. However, the sample size in this study included 100% of available individuals,

which in this case was the total census population. Another source of bias, especially for the LD

method, as reported by Waples *et al.* (2016), is the number of length of chromosomes. In our

case, the SNP data were generated from chromosome 16 - one of the smaller micro-

chromosomes, which has an estimated length of 48-80cM (Burt & Cheng 1998; Groenen *et al.*

2000). We expect both the smaller sample size and the use of single chromosome SNP markers

to be contributing to low N_e estimates using the LD, and perhaps the MC method.

Due to our exact knowledge of the census size, the N_e/N_c ratios are informative about both the

processes of marker inheritance, as well as the performance of the estimators themselves.

Secondly, all the three estimators used in this study are expected to represent the previous or the

most recent parental generations. Therefore, for both the LD and MC methods, we believe that

the estimates obtained here represent the founding event of the Texas population in 2016. On the

other hand, as changes in heterozygosity occur more gradually over time, and the accumulation

of inbreeding may take several generations, the HE estimator may be more representative of a

founder event not in the immediate past, but bounded by the initial founding of the U.S.

populations in 1961.

Haplotype diversity and linkage

We observed only three major haplotypes in the population, and it will be valuable to understand

how these haplotypes are maintained in subsequent generations. We believe that the low

haplotype diversity is a consequence of the history of captive breeding among a small number of

individuals, rather than a reflection of the source populations. Haplotype diversity in other wild

junglefowl populations has been reported to be quite high. For example, Nguyen-Phuc et al.

(2016)reported 313 unique haplotypes from 199 wild-caught individuals from Vietnam, which

indicates that the progenitors of the RRJF came from a similarly diverse population, which have

since lost genetic diversity in captivity. It is also worth noting that the genetic bottleneck and the

resulting reduction in genetic diversity in only 50 years perhaps mirror the early genetic history

during selection for domestication, except for periodic gene flow from wild individuals. It would

appear that such immigration was necessary to maintain the high levels of genetic diversity that

is found in domestic chickens today (Blackburn 2012; Fulton *et al.* 2016; Granevitze *et al.*

2007).

We observed high linkage disequilibrium among loci in the SNP dataset - an estimate that is

known to be inflated when using small sample sizes, and very likely to be contributing to the

high D' values observed (Ardlie *et al.* 2002; England *et al.* 2006). Taken together with the reduced haplotype diversity in this population, however, the linkage disequilibrium estimates are biologically plausible.

Selection for tameness and success in captive environments is inevitable in captive breeding populations (Frankham *et al.* 2002; Frankham *et al.* 1986). The genetic consequences of domestication have received much attention in recent years (Wright *et al.* 2010) and the genetic architecture of traits - mainly driven by linkage and pleiotropy are considered to be crucial in the expression of specific phenotypes in domesticated varieties. For example, Wright et al. (2010)showed the presence of strong linkage blocks, as well as low heterozygosity regions associated with selective sweeps in domesticated varieties of chicken. The haplotype diversity and linkage patterns we observed in our captive population appear to be consistent with what might occur early during domestication.

Conclusions

Our study showed the dynamics of genetic diversity and effective population sizes in a small captive breeding bird population. Furthermore, our study was able to relate effective population sizes to the known demographic history of the captive population. We found that the demographic history corresponds to a great extent with the genetic estimates. Furthermore, the low genetic diversity, and haplotype diversity both indicate that these populations are indeed pure, and perhaps unmixed with domestic varieties, since their initial importation to the United States. Our study shows the rapid loss of genetic diversity in captive populations in only a few generations (37-50), which can be particularly concerning from the standpoint of *ex situ*

conservation goals. Finally, we showed how captive breeding populations, even in the absence of

any intentional selection, can affect linkage structure across a set of adaptive genetic loci, and

potentially reduce the fitness and adaptability of captive-born individuals upon release back into

the wild.

Acknowledgements

The authors thank Dr. Town Peterson, and Shawna Peer for comments on an earlier draft of the

manuscript. This research was funded by undergraduate research grants from Texas A&M

University. We also thank Dr. Janet Fulton of Hy-Line International for her support of this

project.

References

Alcaide M, Serrano D, Tella J, Negro J (2009) Strong philopatry derived from capture-recapture

records does not lead to fine-scale genetic differentiation in lesser kestrels. *Journal of*

Animal Ecology **78**, 468-475.

Araki H, Cooper B, Blouin MS (2007) Genetic effects of captive breeding cause a rapid,

cumulative fitness decline in the wild. *Science* **318**, 100-103.

Ardlie KG, Kruglyak L, Seielstad M (2002) Patterns of linkage disequilibrium in the human

genome. *Nature Reviews Genetics* **3**, 299-309.

Athrey G, Barr KR, Lance RF, Leberg PL (2012) Birds in space and time: genetic changes

accompanying anthropogenic habitat fragmentation in the endangered black-capped vireo

(Vireo atricapilla). *Evolutionary Applications* **5**, 540-552.

Athrey G, Lindsay DL, Lance RF, Leberg PL (2011) Crumbling diversity: comparison of historical archived and contemporary natural populations indicate reduced genetic diversity and increasing genetic differentiation in the golden-cheeked warbler. *Conservation Genetics* **12**, 1345-1355.

Barrett JC, Fry B, Maller J, Daly M (2005) Haploview: analysis and visualization of LD and haplotype maps. . *Bioinformatics* **21**, 263-265.

Blackburn HD (2012) Genetic Selection and Conservation of Genetic Diversity. *Reproduction in Domestic Animals* **47**, 249-254.

Bouzat JL, Cheng HH, Lewin HA*, et al.* (1998) Genetic evaluation of a demographic bottleneck in the Greater Prairie Chicken. *Conservation Biology* **12**, 836-843.

Brisbin IL (1995) CONSERVATION OF THE WILD ANCESTORS OF DOMESTIC-ANIMALS. *Conservation Biology* **9**, 1327-1328.

Brisbin IL, Peterson AT (2007) Playing chicken with red junglefowl: identifying phenotypic markers of genetic purity in Gallus gallus. *Animal Conservation* **10**, 429-435.

Briscoe DA, Malpica JM, Robertson A*, et al.* (1992) RAPID LOSS OF GENETIC-VARIATION IN LARGE CAPTIVE POPULATIONS OF DROSOPHILA FLIES - IMPLICATIONS FOR THE GENETIC MANAGEMENT OF CAPTIVE POPULATIONS. *Conservation Biology* **6**, 416-425.

Bump G (1964) Summary of foreign game bird propagation and liberations, 1960-1963. UWFWS.

Burt D, Cheng H (1998) The chicken gene map. *ILAR Journal* **39**, 229-236.

Condon T (2012) *Morphological Detection of Genetic Introgression in Red Junglefowl (Gallus Gallus)*, Georgia Southern University.

Do C, Waples RS, Peel D, *et al.* (2014) NEESTIMATOR v2: re-implementation of software for the estimation of contemporary effective population size (N-e) from genetic data. *Molecular Ecology Resources* **14**, 209-214.

Emami-Khoyi A, Paterson A, Hartley D (2017) Mitogenomics data reveal effective population size, historical bottlenecks, and the effects of hunting on New Zealand fur seals (Arctocephalus forsteri). In: *Mitochondrial DNA A DNA Mapp Seq Analysis*, pp. 1-13.

England PR, Cornuet JM, Berthier P, Tallmon DA, Luikart G (2006) Estimating effective population size from linkage disequilibrium: severe bias in small samples. *Conservation Genetics* **7**, 303-308.

Frankham R, Ballou J, Briscoe DA (2002) Genetic management of captive populations. In: *Introduction to Conservation Genetics*. Cambridge University Press.

Frankham R, Hemmer H, Ryder OA, *et al.* (1986) SELECTION IN CAPTIVE POPULATIONS. *Zoo Biology* **5**, 127-138.

Fulton JE, McCarron AM, Lund AR, *et al.* (2016) A high-density SNP panel reveals extensive diversity, frequent recombination and multiple recombination hotspots within the chicken major histocompatibility complex B region between BG2 and CD1A1. *Genetics Selection Evolution* **48**.

Granevitze Z, Hillel J, Chen GH, *et al.* (2007) Genetic diversity within chicken populations from different continents and management histories. *Animal Genetics* **38**, 576-583.

Groenen MAM, Cheng HH, Bumstead N, *et al.* (2000) A consensus linkage map of the chicken genome. *Genome Research* **10**, 137-147.

Hammerly SC, Morrow ME, Johnson JA (2013) A comparison of pedigree- and DNA-based measures for identifying inbreeding depression in the critically endangered Attwater's Prairie-chicken. *Molecular Ecology* **22**, 5313-5328.

Hedrick P (1992) Rapid genetic deterioration in captive populations: Causes and conservation implications. In: *Monographiae Biologicae*, pp. 45-68. Springer Netherlands.

Hoelzel AR, Fleischer RC, Campagna C, Le Boeuf BJ, Alvord G (2002) Impact of a population bottleneck on symmetry and genetic diversity in the northern elephant seal. *Journal of Evolutionary Biology* **15**, 567-575.

Johnson JA, Dunn PO (2006) Low genetic variation in the heath hen prior to extinction and implications for the conservation of prairie-chicken populations. *Conservation Genetics* **7**, 37-48.

Kimura M (1980) A SIMPLE METHOD FOR ESTIMATING EVOLUTIONARY RATES OF BASE SUBSTITUTIONS THROUGH COMPARATIVE STUDIES OF NUCLEOTIDE-SEQUENCES. *Journal of Molecular Evolution* **16**, 111-120.

Leberg PL (1992) EFFECTS OF POPULATION BOTTLENECKS ON GENETIC DIVERSITY AS MEASURED BY ALLOZYME ELECTROPHORESIS. *Evolution* **46**, 477-494.

Lovatt FM, Hoelzel AR (2014) Impact on Reindeer (Rangifer tarandus) Genetic Diversity from Two Parallel Population Bottlenecks Founded from a Common Source. *Evolutionary Biology* **41**, 240-250.

Lynch M, O'Hely M (2001) Captive breeding and genetic fitness of natural populations. *Conservation Genetics* **2**, 363-378.

Miao YC, Zhang ZJ, Su JR (2016) Low genetic diversity in the endangered Taxus yunnanensis

following a population bottleneck, a low effective population size and increased

inbreeding. *Silvae Genetica* **65**, 59-66.

Montgomery ME, Woodworth LM, England PR, Briscoe DA, Frankham R (2010) Widespread

selective sweeps affecting microsatellites in Drosophila populations adapting to captivity:

Implications for captive breeding programs. *Biological Conservation* **143**, 1842-1849.

Nei M, Mauyama T, Chakraborty R (1975) THE BOTTLENECK EFFECT AND GENETIC

VARIABILITY IN POPULATIONS. *Evolution* **29**, 1-10.

Newman D, Pilson D (1997) Increased probability of extinction due to decreased genetic

effective population size: Experimental populations of Clarkia pulchella. *Evolution* **51**,

354-362.

Nguyen-Phuc H, Fulton JE, Berres ME (2016) Genetic variation of major histocompatibility

complex (MHC) in wild Red Junglefowl (Gallus gallus). *Poultry Science* **95**, 400-411.

Nomura T (2008) Estimation of effective number of breeders from molecular coancestry of

single cohort sample. *Evolutionary Applications* **1**, 462-474.

Paradis E, Blomberg S, Bolker B*, et al.* (2017) Package 'ape'.

Peakall R, Smouse PE (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic

software for teaching and research-an update. *Bioinformatics* **28**, 2537-2539.

Ralls K, Ballou J (1986a) Captive breeding programs for populations with a small number of

founders. *Trends in Ecology & Evolution* **1**, 19-22.

Ralls K, Ballou J (1986b) Rapid genetic deterioration in captive populations: Causes and

conservation implications. *Zoo Biology* **5**, 81-86.

Ruiz-Lopez MJ, Roldan ERS, Espeso G, Gomendio M (2009) Pedigrees and microsatellites

among endangered ungulates: what do they tell us? *Molecular Ecology* **18**, 1352-1364.

Schuelke M (2000) An economic method for the fluorescent labeling of PCR fragments. *Nature*

Biotechnology **18**, 233-234.

Stephens M, Smith NJ, Donnelly P (2001) A new statistical method for haplotype reconstruction

from population data. *American Journal of Human Genetics* **68**, 978-989.

Tadano R, Nishibori M, Nagasaka N, Tsudzuki M (2007) Assessing genetic diversity and

population structure for commercial chicken lines based on forty Microsatellite analyses.

Poultry Science **86**, 2301-2308.

Tokarska M, Marshall T, Kowalczyk R*, et al.* (2009) Effectiveness of microsatellite and SNP

markers for parentage and identity analysis in species with low genetic diversity: the case

of European bison. *Heredity* **103**, 326-332.

Vali U, Einarsson A, Waits L, Ellegren H (2008) To what extent do microsatellite markers

reflect genome-wide genetic diversity in natural populations? *Molecular Ecology* **17**,

3808-3817.

Wang J, Santiago E, Caballero A (2016) Prediction and estimation of effective population size.

Heredity **117**, 193-206.

Waples RK, Larson WA, Waples RS (2016) Estimating contemporary effective population size

in non-model species using linkage disequilibrium across thousands of loci. *Heredity*

117, 233-240.

Waples RS, Antao T, Luikart G (2014) Effects of Overlapping Generations on Linkage

Disequilibrium Estimates of Effective Population Size. *Genetics* **197**, 769-U603.

Waples RS, Do C (2008) LDNE: a program for estimating effective population size from data on

linkage disequilibrium. *Molecular Ecology Resources* **8**, 753-756.

Willoughby JR, Fernandez NB, Lamb MC*, et al.* (2015) The impacts of inbreeding, drift and

selection on genetic diversity in captive breeding populations. *Molecular Ecology* **24**, 98-

110.

Willoughby JR, Ivy JA, Lacy RC, Doyle JM, DeWoody JA (2017) Inbreeding and selection

shape genomic diversity in captive populations: Implications for the conservation of

endangered species. *Plos One* **12**.

Wisely SM, Buskirk SW, Fleming MA, McDonald DB, Ostrander EA (2002) Genetic diversity

and fitness in black-footed ferrets before and during a bottleneck. *Journal of Heredity* **93**,

231-237.

Witzenberger KA, Hochkirch A (2011) Ex situ conservation genetics: a review of molecular

studies on the genetic consequences of captive breeding programmes for endangered

animal species. *Biodiversity and Conservation* **20**, 1843-1861.

Wright D, Rubin CJ, Barrio AM*, et al.* (2010) The genetic architecture of domestication in the

chicken: effects of pleiotropy and linkage. *Molecular Ecology* **19**, 5140-5156.

Wright S (1931) Evolution in Mendelian populations. *Genetics* **16**, 0097-0159.

Wright S (1948) ON THE ROLES OF DIRECTED AND RANDOM CHANGES IN GENE

FREQUENCY IN THE GENETICS OF POPULATIONS. *Evolution* **2**, 279-294.

Zhdanova OL, Pudovkin AI (2008) Nb_HetEx: A Program to Estimate the Effective Number of

Breeders. *Journal of Heredity* **99**, 694-695.

Tables

Table 1: Estimates of genetic diversity for both microsatellite and SNP datasets are presented.

For each marker, genetic diversity measures are shown along with their standard errors. The

columns present the number of alleles (Na), effective number of alleles (Nae), observed

heterozygosity (Ho), expected heterozygosity (He), the unbiased expected heterozygosity (uHe),

and the fixation index F.

Marker Type		Na	Ne	Ho	He	uHe	F
Microsatellite	Mean	2.33	1.74	0.36	0.37	0.38	0.07
	SE	0.16	0.14	0.06	0.05	0.05	0.08
SNP	Mean	1.38	1.13	0.11	0.1	0.1	-0.14
	SE	0.05	0.02	0.01	0.01	0.01	0.02

Table 2: Estimates of effective population size Ne are presented for microsatellite and SNP

datasets. Ne estimates from three estimators are shown along with their respective 95%

confidence intervals. Ne was estimated using the Heterozygote Excess (NeHE), Linkage

Disequilibrium (NeLD), and the Molecular Coancstry (NeMC) methods.

	NeHE	95% CI	NeLD	95% CI	NeMC	95% CI
Microsatellite	inf	7.3-inf	7.3	3.6-12.5	4	1.1-8.8
SNP	4.8	3.1-11	0.2	0.1-0.2	1	1-1

Figure 1: A linkage map based on the 90 SNP loci on chromosome 16, generated from genotype

data for the study population. The map shows pairwise estimates of linkage disequilibrium

between markers. Red colored blocks suggest high linkage disequilibrium (D') values of 1,

implying little to no recombination between markers.

Figure 2: A haplotype network of the the five (phased) haplotypes detected in the study

population. Only three of the five unique haplotypes were found at a frequently of over 10% in

the population (I, III, and V).