Identification of a novel subfamily of bacterial AAT-fold basic amino acid decarboxylases

and functional characterization of its first representative: *Pseudomonas aeruginosa* LdcA

Diego Carriel-Lopez,[1,2,#] Pierre Simon Garcia,[3,4,#] Florence Castelli,[5] Patricia Lamourette,[5]

François Fenaille,[5] Céline Brochier-Armanet,[3,4,*] Sylvie Elsen[2,] and Irina Gutsche[1,***]**

[1]University of Grenoble Alpes, CNRS, CEA, CNRS, IBS, F-38000 Grenoble, France

[2]Biology of Cancer and Infection, U1036 INSERM, CEA, University of Grenoble Alpes, ERL5261

CNRS, Grenoble, France

[3]Université Lyon 1, CNRS, UMR5558, Laboratoire de Biométrie et Biologie Évolutive, F-69622,

Villeurbanne, France

[4]MMSB Molecular Microbiology and Structural Biochemistry, Institut de Biologie et de Chimie

des Protéines 69367 LYON cedex 07, France

[5]Service de Pharmacologie et Immuno-Analyse (SPI), Laboratoire d'Etude du Métabolisme des

Médicaments, CEA, INRA, Université Paris Saclay, MetaboHUB, F-91191 Gif-sur-Yvette, France

For correspondence. *E-mail celine.brochier-armanet@univ-lyon1.fr; Tel. (33) 426234476; Fax

(33) 472431388; **E-mail sylvie.elsen@cea.fr; Tel. (33) 438783074; Fax (33) 438785058;

***E-mail irina.gutsche@ibs.fr; Tel. (33) 457428766; Fax (33) 438785122.

Short Title: LdcA, a new phylogenetic family of bacterial Ldcs

Key words: bacterial amino acid decarboxylases, polyamines, evolution, lysine decarboxylase,

Pseudomonas aeruginosa

[#]These authors contributed equally to this work

Summary

Polyamines are small amino-acid derived polycations capable of binding negatively charged

macromolecules. Bacterial polyamines are structurally and functionally diverse, and are mainly

produced biosynthetically by PLP-dependent amino acid decarboxylases referred to as LAOdcs

(Lysine-Arginine-Ornithine decarboxylases). In a phylogenetically limited group of bacteria,

LAOdcs are also induced in response to acid stress. Here, we performed an exhaustive

phylogenetic analysis of the AAT-fold LAOdcs which showcased the ancestral nature of their

short forms in *Cyanobacteria* and *Firmicutes*, and emergence of distinct subfamilies of long

LAOdcs in *Proteobacteria*. Complex evolutionary history of these subfamilies and their

relationships were disclosed. Moreover, we identified a novel subfamily of lysine

decarboxylases, LdcA, ancestral in *Betaproteobacteria* and *Pseudomonadaceae*. Thus, we

analyzed expression of LdcA from *Pseudomonas aeruginosa*, and uncovered its role in

promoting growth and reducing persistence of this multidrug resistant human pathogen during

carbenicillin treatment, intimately linked to cadaverine production. Finally, we documented a

certain redundancy in the function of the three main polyamines - cadaverine, putrescine and

spermidine – in *P. aeruginosa* by demonstrating the link between their intracellular level, as well

as the capacity of putrescine and spermidine to complement the growth phenotype of the *ldcA*

mutant.

Introduction

Polyamines are small amino acid-derived molecules with two or more amino groups separated

by alkyl chains (Lightfoot & Hall, 2014, Tabor & Tabor, 1964). They perform essential functions

in all living organisms by participating in DNA replication, gene expression and protein synthesis,

and are generally described as growth factors (Lightfoot & Hall, 2014, Michael, 2016b). At

physiological pH, polyamines behave as polycations and can interact with negatively charged

macromolecules such as nucleic acids, membrane phospholipids and proteins (Tabor & Tabor,

1964, Tabor & Tabor, 1985). They were proposed to bind and structurally modify RNA thereby

acting at the level of translation (Igarashi, 2006, Igarashi & Kashiwagi, 2015). Strengthening this

hypothesis, "polyamine modulons" were identified in *Escherichia coli* and more recently in

eukaryotes (Igarashi, 2006, Igarashi & Kashiwagi, 2015). As a possible consequence, bacterial

polyamines were shown to participate in expression of proteins essential for growth fitness and

viability but also in processes such as biofilm formation, antibiotic resistance and virulence

(Michael, 2016a, Di Martino *et al.*, 2013, Karatan & Michael, 2013, Kwon & Lu, 2006, Shah &

Swiatlo, 2008).

The triamine spermidine (Spd), essential in eukaryotes and archaea (Michael, 2016b,

Lightfoot & Hall, 2014, Michael, 2016a), and its diamine precursor putrescine (Put) are

extensively studied because these two polyamines are the only ones produced in all eukaryotes

(Michael, 2016a, Michael, 2016b). In bacteria, the polyamine repertoire and hence their roles

are the most diverse. In particular, a third widespread bacterial polyamine, cadaverine (Cad),

appears increasingly important for the physiology of *Proteobacteria*. Cad is recognized as an

important player in enterobacterial acid stress response wherein it decreases porin permeability

to protons and alkalinizes the medium (Dela Vega & Delcour, 1996, Zhao & Houry, 2010). During

oxidative stress response, Cad is also capable of scavenging reactive oxygen species (ROS) in

Vibrio vulnificus (Kang *et al.*, 2007). In some Negativicutes, *Selenomonas ruminantium* and

Veillonella sp., Cad was shown to be incorporated in the peptidoglycan and essential for its

stability (Kamio *et al.*, 1986, Kamio & Nakamura, 1987). Finally, this polyamine is involved in iron

uptake and is required for the synthesis of hydroxamate-type siderophores in different bacterial

species such as *Streptomyces coelicolor* (Burrell *et al.*, 2012).

Polyamine biosynthesis depends on the activity of basic amino acid decarboxylases using

Lysine, Arginine or Ornithine as specific substrate to produce Cad, Agmatine (Agm) or Put,

respectively. These enzymes can therefore be generally referred to as LAOdcs (Lysine-Arginine-

Ornithine decarboxylases). In most eukaryotes, Put synthesis is carried out by an Odc. An

alternative pathway for Put biosynthesis found in bacteria and plants involves decarboxylation

of arginine by arginine decarboxylases (Adc); these enzymes produce Agm which is further

converted into Put by agmatine iminohydrolase/deiminase and N-carbamoylputrescine

amidohydrolase (Michael, 2016b, Michael, 2016a). Interestingly, plants (Bunsupa *et al.*, 2012,

Lee & Cho, 2001) and some bacteria exemplified by *S. ruminantium* and *V. vulnificus* (Lee *et al.*,

2007, Takatsuka *et al.*, 2000) possess a bifunctional Odc/Ldc capable of synthesizing both Put

and Cad. Spd is formed by spermidine synthase (SpdSyn) through aminopropylation of Put,

using an aminopropyl group released by decarboxylation of S-adenosylmethionine (Michael,

2016a, Michael, 2016b). In addition, *E. coli* SpdSyn transforms Cad into aminopropyl-Cad, a Spd

analogue sharing its growth stimulating properties (Kim *et al.*, 2016).

Two major structural protein super-families are responsible for polyamine biosynthesis

through pyridoxal-5-phosphate (PLP)-dependent LAOdcs: the alanine racemase fold (AR-fold)

super-family and the aspartate aminotransferase fold (AAT-fold) super-family (Eliot & Kirsch,

2004). Phylogenetic studies of the LAOdcs of the AR-fold super-family revealed that these

enzymes are widespread throughout the three domains of life (Lee *et al.*, 2007, Burrell *et al.*,

2010) while the AAT-fold LAOdcs are exclusive to Bacteria and Archaea. Bacterial AAT-fold

LAOdcs can be divided in two types according to the presence or absence of a particular CheY-

like response regulator receiver domain, known as the "wing domain", necessary for the

formation of higher-order oligomers (Burrell *et al.*, 2010, Kanjee *et al.*, 2011b). The short form

referred as wing-less LAOdc was found in *Firmicutes*, *Cyanobacteria* and *Actinobacteria* phyla,

and a few "wing-less" AAT-fold decarboxylases from *Firmicutes* and *Actinobacteria* were shown

to have an Adc activity, required in particular for biofilm formation in *Bacillus subtilis* (Burrell *et*

al., 2010). The long, wing domain-containing form appeared featured by *Proteobacteria* (Kanjee

et al., 2011b).

AAT-fold decarboxylases with the wing domain have been intensively studied in

Enterobacteria since the early 1940s (Gale & Van Heyningen, 1942, Gale & Epps, 1944)

because of the link between enterobacterial pathogenicity for humans and their capacity to

withstand acid stress thanks to the crucial role of the inducible LAOdcs. Consequently, the

current understanding of the AAT-fold LAOdc is based on analyses of a very limited number of

bacterial species, *i.e.* mostly *Enterobacteria E. coli*, *Salmonella typhimurium*, *Vibrio cholerae* and

V. vulnificus. At a specific acidic pH, expression of the decarboxylase gene is induced by an

excess of the target amino acid uptaken by a dedicated inner membrane antiporter (Kanjee &

Houry, 2013). The enzyme transforms the amino acid substrate into the corresponding

polyamine upon consumption of a proton and production of a CO_2 molecule. In association with

the polyamine excretion by the antiporter, this reaction results in an efficient buffering of the

intracellular medium and the extracellular surroundings. These inducible stress response

decarboxylases are distinguished from "biosynthetic" enzymes that are involved only in

polyamine biosynthesis. *E. coli* encodes two biosynthetic decarboxylases (LdcC: Lys->Cad and

OdcC/SpeC: Orn->Put) responsible for Cad and Put biosynthesis respectively, and three acid

stress-inducible decarboxylases (LdcI: Lys->Cad, AdcI/AdiA: Arg->Agm and OdcI/SpeF: Orn->Put)

which together constitute a very robust acid stress response system that allows the bacterium

to survive upon acid stress as low as pH 2.0 (Zhao & Houry, 2010, Kanjee *et al.*, 2011a, Kanjee

& Houry, 2013). The importance of LdcI was also demonstrated in *S. typhimurium, V. cholerae*

and *V. vulnificus*, where it promotes growth and survival under acidic conditions but also

confers protection from oxidative stress insults (Merrell & Camilli, 2000, Kang *et al.*, 2007, Viala

et al., 2011). It should be noted that LdcI-encoding genes are often designated as *cadA* as

originally proposed upon identification of this gene in *E. coli* (Tabor *et al.*, 1980).

E. coli LdcI but not LdcC interacts with the AAA+ ATPase RavA to assemble into a huge

macromolecular cage (Snider *et al.*, 2006, El Bakkouri *et al.*, 2010, Malet *et al.*, 2014, Kandiah *et*

al., 2016). One of the functions of this mysterious complex is to protect LdcI from inhibition by

the stringent response alarmone ppGpp, thus enabling the bacterium to efficiently cope with

both acid and nutrient stress simultaneously (El Bakkouri *et al.*, 2010, Kanjee *et al.*, 2011a,

Malet *et al.*, 2014). While investigating why RavA binds only LdcI but not LdcC, we documented

numerous inconsistencies in annotation of enterobacterial *ldcI* and *ldcC* genes and realized that

each of these two families appeared to have a distinct genetic context (Kandiah *et al.*, 2016).

Thus, spurred on by the limited nature of the previous studies, we set out to perform an

extensive phylogenetic analysis of the AAT-fold LAOdcs in circa 4,500 complete prokaryotic

proteomes to decipher the evolutionary history of these proteins and their functional evolution.

In the present study, we revealed the ancestral nature of the wing-less LAOdcs in *Cyanobacteria*

and *Firmicutes*, and the complex evolutionary history of long AAT-fold LAOdcs in *Proteobacteria*,

leading to the emergence of distinct subfamilies. Moreover, we disclosed a novel subfamily of

enzymes, clearly distinct from the well-known LdcI, LdcC, AdcI, OdcI and OdcC families, but

more related to Ldc and Adc. Excitingly, this novel evolutionary subfamily has been overlooked

in previous phylogenetic analyses in spite of its wide distribution in *Betaproteobacteria* and

Pseudomonadaceae, implying that it deserves a thorough characterization and a functional

comparison with the known AAT-fold long LAOdc. The only previously mentioned LAOdc from

these taxa is the lysine decarboxylase LdcA from a major multidrug resistant opportunistic

human pathogen *Pseudomonas aeruginosa* (Chou *et al.*, 2010). Thus, here we went beyond the

initial characterization of the *P. aeruginosa* LdcA (Chou *et al.*, 2010), and further analyzed its

expression, regulation, and function in the light of the available knowledge in particular on *E.*

coli LdcI and LdcC. This combined phylogenetic and functional study revealed that LdcA belongs

to a novel subgroup of the long AAT-fold LAOdcs, and that its function is linked to Cad

production and to the general polyamine metabolism rather than to stress response.

Results

Evolutionary history of bacterial LAOdcs

An in-depth survey of 4,466 prokaryotic proteomes representing 1,904 species revealed 4,090

protein sequences belonging to the AAT-fold LAOdcs (13 of which were unannotated or

annotated as pseudo-genes) (Table S1). Representatives of this super-family are mainly present

in *Proteobacteria*, *Firmicutes*, *Actinobacteria* and *Cyanobacteria* (Fig. 1A). The corresponding

Maximum Likelihood (ML) tree could be divided in three parts (Fig. 1B). Cluster I encompasses

sequences devoid of the wing domain (short AAA-fold LAOdc), found in *Firmicutes*,

Cyanobacteria and *Actinobacteria* (Bootstrap value (BV) = 99%). Cluster II encloses nearly all

proteobacterial sequences and a few sequences from Firmicutes (BV = 99%), all containing a

wing domain. Cluster III branches in-between Cluster I and Cluster II; it is composed of a mix of

long and short AAT-fold LAOdc sequences from unrelated taxonomic groups (*Proteobacteria*,

Actinobacteria, *Bacteroidetes*, and *Firmicutes*), which strongly suggests that their taxonomic

distribution results from horizontal gene transfers among these lineages.

Within Cluster I, all cyanobacterial sequences group together albeit with a weak support

(Bootstrap value, BV < 90%) and appear widely distributed in this phylum (Fig. S1A).

Furthermore, the phylogeny of these sequences is overall consistent with a reference phylogeny

of *Cyanobacteria* based on ribosomal proteins (Fig. S1A and S1B). This indicates that a gene

coding for a wing-less AAT-fold LAOdc was ancestral and had been mainly transmitted vertically

in *Cyanobacteria*. Genomic context of LAOdc in Cyanobacteria is not conserved even in

relatively close species (Fig. S1C). Similarly, a large group of *Firmicutes* sequences is observed in

the tree (Fig. 1). Comparison of the taxonomic distribution and the phylogeny of these

sequences with a reference phylogeny of *Firmicutes* suggests the ancestral presence of a gene

coding for a short AAT-fold LAOdc in this phylum (Fig. S2A and S2B). Furthermore, most of

Firmicutes harbor two LAOdc copies (A and B) that can be distinguished by their genomic

context (Fig. S2C), with exception of the most early-branching lineages (a single copy in

Natranaerobiaceae, gene absence in *Halobacteroidaceae* and *Halanaerobiaceae*). The

paralogue A, ancestral in *Firmicutes*, presents a well-conserved genomic context (thymidylate

kinase and DNA polymerase III subunit delta). In contrast, the paralogue B appeared at the

emergence of *Clostridia* and *Bacilli,* likely as the result of a gene duplication event, and its

genetic context is not conserved. A loss of both paralogues can be inferred at the emergence of

Lactobacillales followed by a reacquisition in some *Streptococcaceae* and *Carnobacteriaceae* by

horizontal gene transfer from different *Firmicutes* donors (Fig. S2A). Finally, Cluster I

encompassed a group of sequences from *Actinobacteria*. These sequences emerged from the

Firmicutes group, suggesting that they have been acquired by horizontal gene transfer from

Firmicutes.

Analysis of proteobacterial LAOdcs composing the Cluster II of the tree revealed two

monophyletic groups, Odc (BV/Posterior probabilities PP = 100%/1) and LAdc (BV/PP = 100%/1)

(Fig. 2A and 2B). The LAdc group is further split into LdcI/LdcC (BV/PP = 100%/1), Adc (BV/PP =

93%/1), and a hitherto non-identified subfamily (BV/PP =100%/1), that will be referred as LdcA

in the name of its only previously mentioned member, LdcA from *P. aeruginosa* (Chou *et al.*,

2010). It should be emphasized that although the monophyly of OdcI/OdcC, LdcI/LdcC, Adc and

LdcA groups is well supported, the relationships between LdcI/LdcC, Adc and LdcA are

unresolved (BV/PP < 80%/0.67).

Comparison of LAOdc phylogeny, taxonomic distribution and genomic contexts with a

reference phylogeny of *Proteobacteria* suggests that OdcI could be ancestral in

Bradyrhizobiaceae, *Xanthobacteriaceae*, *Methylobacteriaceae* and *Beijerinckiaceae*

(*Alphaproteobacteria*) and *Vibrionaceae*, *Pasteurellaceae* and *Enterobacteriaceae*

(*Gammaproteobacteria*) (Fig. 3A). Interestingly, a few horizontal gene transfers of *odcI* occurred

from *Proteobacteria* to *Firmicutes*, especially to some *Lactobacillus, Staphylococcus lugdunensis*

and *Megasphaera elsdenii*. In contrast, *odcC* seems more recent and appears to result from a

gene duplication of *odcI* that occurred in *Enterobacteriaceae*, before the divergence of *Sodalis*

(BV/PP = 75%/1, Fig. 3B). Genes coding for OdcI and OdcC are clearly distinguished by their

context (Fig. 4). More precisely, *odcI* is predominantly upstream from a gene (*potE*) coding for

putrescine-ornithine antiporter, consistently with the function of OdcI that converts Orn to Put.

In contrast, OdcC coding gene is located in the vicinity of Fe^{2+}-trafficking protein (*yggX*) and a

lytic murein transglycosylase (*mltC*) genes.

LdcI appears to be ancestral in some gammaproteobacterial lineages (*Francisellaceae*,

Aeromonadaceae, *Vibrionaceae* and *Enterobacteriaceae*, Fig. 3A). Moreover, similar to

odcI/odcC, *ldcC* apparently derived from *ldcI*, and more precisely from a gene duplication that

occurred, as for *odcI*, just before the emergence of *Sodalis* (BV/PP = 100%/1, Figure 3B). The *ldcI*

gene, also called *cadA*, forms a *cadBA* operon together with the lysine-cadaverine antiporter-

encoding the *cadB* gene. In *Enterobacteriaceae*, this operon is known to be regulated by the

transcriptional factor CadC integrating three external signals – low pH, high lysine and low Cad

levels (Kuper & Jung, 2005, Fritz *et al.*, 2009). Our analysis confirmed the conserved genomic

organization of the *cadCBA* system (Fig. 4) (Zhao & Houry, 2010) allowing *Enterobacteria* to

face acid and oxidative stresses. The *ldcC* genomic context seems to be also strongly conserved

(Fig. 4) with genes coding for hydroxymyristol acyl carrier protein dehydratase, UDP-N-

acetylglucosamine acetyltransferase, tetraacyldisaccharide-1-P synthase, ribonuclease HII, DNA

polymerase III alpha subunit, and acetyl-CoA carboxylase 2C alpha subunit upstream and

putative lyase and tRNA(Ile)-lysidine synthetase genes downstream.

In sharp contrast with OdcI/OdcC and LdcI/LdcC, Adc group presents a patchy taxonomic

distribution and their relationships are at odds with species phylogeny, with sequences from

different classes of *Proteobacteria* being intermixed on the tree (Fig. 3). This indicates that the

Adc subfamily was heavily impacted by horizontal gene transfers. Accordingly, it cannot be

inferred as ancestral in any taxonomic group. The genomic context of *adc* is not conserved (Fig.

4), precluding identification of potential functional partners. Interestingly, the *Burkholderia* sp.

AIU 395 LAOdc that locates in the Adc group was shown to possess a lysine decarboxylase and

oxidase activity (Sugawara *et al.*, 2014).

As specified above, beside OdcI/C, LdcI/C and Adc, a fourth group of LAOdc sequences

emerged in the tree. This group contains LdcA from *P. aeruginosa*, shown to possess a lysine

decarboxylase activity (Chou *et al.*, 2010). Homologues of LdcA are found in other

representatives of *Pseudomonadaceae*, as well as in nearly all *Betaproteobacteria*. Both groups

of sequences are well separated on the tree and widely distributed in these two taxa, and the

relationships among betaproteobacterial sequences are globally in agreement with the species

taxonomy. This suggests that an LdcA homologue was present in the ancestor of

Pseudomonadaceae and in the ancestor of *Betaproteobacteria* and has been conserved along

the diversification of these two taxa. Worthy of note, *ldcA* was secondly lost in *Neisseriaceae*

and *Nitrosomonadaceae*. The genetic environment of *ldcA* is not conserved (Fig. 4). To our

knowledge, the only publication describing a member of the LdcA subfamily concerns *P.*

aeruginosa, a highly versatile bacterium that efficiently grows on arginine but not on lysine

(Fothergill & Guest, 1977, Rahman & Clarke, 1980). In this organism exhibiting tightly

interconnected lysine and arginine catabolism networks (Chou *et al.*, 2010, Madhuri Indurthi *et*

al., 2016), the *PA1818* gene was identified as a part of the ArgR regulon upon growth on excess

arginine, but was surprisingly found to code for a lysine decarboxylase, and not for an arginine

decarboxylase as one would have logically supposed, and therefore called *ldcA* (Chou *et al.*,

2010, Madhuri Indurthi *et al.*, 2016). *P. aeruginosa* is a major cause of gram-negative infections,

especially in patients with compromised and weakened immune system. This opportunistic

pathogen is also a well-identified threat for patients suffering from Cystic Fibrosis (CF), because

the chronic respiratory infections associated to host inflammatory responses lead to pulmonary

tissue destruction and lung failure. The occurrence and persistence of *P. aeruginosa* in the CF

patients' lungs, whose secretions were shown to be acidified and to become oxidative, hints to

a possible role of LdcA in promoting bacterial fitness. Therefore, in the following sections, we

chose to deepen the present knowledge on expression, regulation and functional

characterization of *P. aeruginosa* LdcA.

ldcA expression is growth-phase dependent

To determine the role of the LdcA protein in *P. aeruginosa*, the prerequisite was to know when

the protein is produced by the bacterium and what are the mechanisms controlling its

production. Thus, we first analyzed the expression of its gene by creating a transcriptional

fusion between the *ldcA* promoter (P*ldcA*) and the reporter *lacZ* gene. The fusion was then

integrated into the chromosome of PAO1 (see Materials and Methods for further details), the

first *P. aeruginosa* sequenced strain (Stover *et al.*, 2000) considered as a reference strain and

widely studied. We then measured the β-galactosidase activity of the PAO1::P$_{ldcA}$-*lacZ* strain

grown in a minimal medium P (MMP) containing glutamate as a carbon source and lysine or

arginine (20mM) as additives. Unlike lysine, arginine was able to induce the expression of *ldcA* in

PAO1::P_{ldcA}-*lacZ* (not shown), in agreement with published data identifying ArgR as a positive

regulator of *ldcA* expression in the PAO1 strain (Chou *et al.*, 2010, Lu *et al.*, 2004).

The pattern of *ldcA* expression was followed during the growth both in minimal and rich

media. In MMP medium supplemented with glutamate and arginine (MMP-GR), the β-

galactosidase activity increased slightly but continuously along the growth and reached a

maximum in the stationary phase (Fig. 5A). Expression of *ldcA* in LB rich medium followed the

same pattern of expression (Fig. 5B), that was paralleled by an increase in LdcA protein amount

assessed by western blot (not shown). The expression level obtained in LB was two-fold lower

that the one measured in MMP-GR. Addition of 20 mM arginine to LB did not change the growth

rate of the bacteria but the *ldcA* promoter activity increased during the transition to the

stationary phase to reach a level similar to the one measured in MMP-GR (Fig. 5B), indicating a

probable limiting concentration of the amino acid in LB at late growth.

ldcA *expression differs from that of* ldcC *and* ldcI

The *ldcA* genomic context being different from that of *ldcI* and *ldcC* genes (Fig. 4), we wondered

if *ldcA* was also regulated differently than these genes; this knowledge could further provide

information on its role in *P. aeruginosa* physiology. Even if *E. coli* LdcC is called "constitutive

Lysine Decarboxylase", the corresponding gene was shown to be induced in stationary phase in

LB medium by RpoS, the sigma factor of stationary phase (Kikuchi *et al.*, 1998). However, *ldcA*

expression was not affected in a *rpoS* mutant background (data not shown), in agreement with

previous transcriptomics analyses demonstrating that *ldcA* is not part of the RpoS

regulon(Schuster *et al.*, 2004). To compare with LdcI, the "inducible Lysine Decarboxylase",

despite the absence of a CadC homologue in *P. aeruginosa,* we assessed if acid stress could

activate the expression of *ldcA* in a CadC-independent manner. Hence, we induced acid stress

by decreasing the pH of the medium to a value of 5 during the exponential phase of growth and

documented an absence of effect of this treatment on *ldcA* expression (Fig. S4A). In addition,

ldcI from *V. vulnificus* was reported to be induced by SoxR upon H_2O_2 stress (Kim *et al.*, 2006).

In *P. aeruginosa*, SoxR is not a key player in the oxidative stress response, but H_2O_2 activates the

global regulator OxyR that orchestrates the defense against ROS (Ochsner *et al.*, 2000). Yet,

addition of 1 mM of H_2O_2 in the culture did not affect *ldcA* expression (Fig. S4A). To conclude, *P.*

aeruginosa does not overexpress *ldcA* to respond to low pH or oxidative stress conditions.

Low pH survival is not affected in absence of LdcA

To clarify the function of LdcA in *P. aeruginosa,* a mutant deleted of the *ldcA* gene, as well as a

complemented strain in which one copy of *ldcA* driven by its own promoter was inserted in the

chromosome, were engineered (see Materials and Methods). Using Biolog system, a Phenotype

MicroArray analysis was carried out to test a large number of stress conditions in systematical

and reproducible manner. A potential role of LdcA in detoxifying and protecting *P. aeruginosa*

during acid, alkaline and oxidative stress, antibiotic treatment and toxic molecules causing DNA

damage, nitrosative stress, and membrane destabilization was assessed. After monitoring the

growth of *P. aeruginosa* in the different conditions in minimal medium during 24 hours (see

Materials and Methods), the "Area Under the Curve" (AUC) in each condition was calculated

and compared between the strains. Analysis of the growth fitness of wild-type, mutant and

complemented strains indicated that *P. aeruginosa* could grow optimally in a pH range from 5 to

10 without considerable effect on the metabolism and biomass growth. At pH 4 to 5, the

bacterial growth started to be strongly inhibited and the strains were unable to grow below pH

4 (Fig. S4B). This set of experiments did not show any significant difference between the mutant

and the wild-type and complemented strains, pointing to a non-involvement of LdcA in survival

at low pH. Similarly, no significant effect of *ldcA* absence on resistance against antibiotics,

oxidative and toxic agents could be detected (not shown). Hence LdcA seems not to be

important for any stress response.

The Cad pool generated by LdcA impacts persistence phenotype and polyamine content

In our quest for deciphering the importance of LdcA in bacterial physiology, we analyzed the

role of its product Cad shown to play a fundamental role in *P. aeruginosa* virulence. Indeed, this

polyamine was reported to be involved in the persistence of the bacterium which is of

importance for its eradication by antibiotics; specifically, Cad production was shown to lead to a

reduction of the dormant cells that form an antibiotic-tolerant subpopulation in MH medium

(Manuel *et al.*, 2010).

Therefore, to assess the role of LdcA in persistence, we first confirmed the impact of *ldcA*

mutation on Cad production in this rich MH medium. To do so, we quantified the intracellular

Cad amounts in the bacterial strains by liquid chromatography coupled to high resolution mass

spectrometry (LC/HRMS) during early-, mid- and late-exponential growth phases. Cad was

completely absent in the *ldcA* mutant and complementation with a wild-type *ldcA* copy restored

the metabolite level in the strain, indicating that, in PAO1, this polyamine is produced

exclusively through the enzymatic activity of LdcA in this growth condition (Fig. 6). Moreover,

the wild-type strain showed an impressive increase (around 25-fold) of Cad concentration

during the growth, probably reflecting the *ldcA* expression pattern in rich medium (increase

during exponential growth, Fig. 5B).

Then, the impact of *ldcA* on the number of persisters during carbenicillin treatment was

assessed. As anticipated, *ldcA* mutant showed a number of persisters significantly higher

compared to the wild-type PAO1 and complemented strains (Fig. S5), confirming the

importance of LdcA activity in the persistence phenotype.

Interestingly, in parallel to Cad, we also quantified the intracellular concentrations of two

other polyamines, Put and Spd, in the wild-type strain and *ldcA* mutant (Fig. 6). While in the

wild-type PAO1, the amount of Cad is clearly growth-phase dependent, the amounts of Put and

Spd were found to be abundant and constant in the MH medium. Remarkably, inactivation of

ldcA affected not only the production of Cad but also the amount of intracellular Put and Spd

which were reduced by around two-fold in the *ldcA* mutant at late exponential growth. This

could either indicate a higher turnover of polyamine metabolism in the *ldcA* mutant or a

reduced production of Put and Spd as a compensation for the absence of Cad.

Polyamines are important for growth fitness

As mentioned above, LdcA synthesis is induced in minimal MMP medium in presence of

arginine. Therefore, we monitored the growth of the *ldcA* mutant in this condition in presence

of either lysine, the substrate of LdcA, or Cad, the product of its enzymatic transformation. The

observed clear reduction of growth of the *ldcA* mutant highlighted the metabolic role of the

enzyme, whereas wild-type growth was restored in the complemented strain (Fig. 7A). Addition

of exogenous Cad was sufficient to restore a normal growth in the mutant indicating that the

limiting factor was indeed the polyamine product (Fig. 7B). Strikingly, growth was also restored

when Put (Fig. 7C) or Spd (Fig. 7D) were added to the growth medium at the same

concentrations, indicating that these polyamines can substitute for Cad as a growth factor.

Discussion

The present exhaustive phylogenetic analysis of the AAT-fold LAOdcs super-family

reveals that wing-less enzymes were ancestral in *Firmicutes*, in agreement with earlier reports

(Sekowska *et al.*, 1998, Burrell *et al.*, 2010). This analysis highlights also the ancestral presence

of wing-less AAT-fold LAOdc in *Cyanobacteria*. Yet, no LAOdc activity of a cyanobacterial AAT-

fold enzyme has been documented despite their purification from *Anabaena variabilis* and

Nostoc punctiforme (Burrell *et al.*, 2010). We show that wing domain-containing AAT-fold LAOdc

emerged during the diversification of *Proteobacteria*, suggesting that short AAT-fold LAOdc

could be more ancient and that the acquisition of the CheY-like wing domain occurred likely

secondarily, as previously proposed (Burrell *et al.*, 2010, Kanjee *et al.*, 2011a). The absence of

AR-fold decarboxylases in *Firmicutes* implies that their AAT-fold decarboxylases would be the

only route for polyamine biosynthesis and thus emphasizes the importance of these enzymes in

this phylum. The *Firmicutes* AAT-fold LAOdc correspond to two paralogues (A and B) present in

most species and easily distinguishable by their genomic context. These two copies result from a

duplication event that occurred very early during diversification of the phylum, probably in the

ancestor of Clostridia and Bacilli (Fig. S1). Paralogue A encompasses the *B. subtilis yaaO*

(Sekowska *et al.*, 1998), while paralogue B contains the *B. subtilis speA* gene coding for an

arginine decarboxylase (NP_389346.1) (Burrell *et al.*, 2010). Two other *Firmicutes* proteins

corresponding to paralogue B sequences (from *Clostridium difficile* YP_001087362.1 and *S.*

ruminantium WP_014425426.1) (Fig. S2B) were also shown to possess an arginine

decarboxylase activity (Burrell *et al.*, 2010, Liao *et al.*, 2008), which led to a hypothesis that

paralogue B AAT-fold decarboxylases would be Adcs (Burrell *et al.*, 2010). The function of

paralogue A awaits further investigation and, to our knowledge, the only study focused on the

yaaO gene (paralogue A) concluded that in *B. subtilis* this protein had no effect on polyamine

production (Sekowska *et al.*, 1998).

Regarding the evolutionary history of the wing domain-containing LAOdcs, our analysis

reveals that proteobacterial LAOdcs form two monophyletic groups, Odc and LAdc. Thus, lysine

and arginine decarboxylases appear more closely related to each other than to ornithine

decarboxylases, although the relationships among the two Ldc families (LdcI/C and LdcA) and

the Adc family are not resolved (PP <0.5) (Fig. 2B). Furthermore, we showed that the

proteobacterial biosynthetic OdcC and LdcC emerged from inducible Odc and Ldc (*i.e.* OdcI and

LcdI, respectively) through two independent gene duplication events that occurred in

Enterobacteriaceae, after the divergence of *Sodalis*. Given that both duplication events seem to

occur in the same branch of the phylogenetic tree, it is tempting to hypothesize that they are

linked, and that a functional connection between both biosynthetic subgroups may have

existed. The emergence of biosynthetic enzymes from inducible ones may appear contra-

intuitive at the first glance, but may reflect an expansion and a diversification of polyamine

functions in these lineages. This would also explain why *Enterobacteriaceae* possess also a

constitutive pathway of polyamine synthesis through an AR-fold Adc.

The exhaustive phylogenetic analysis of AAT-fold decarboxylases discloses multiple cases

of horizontal gene transfers (e.g. within Cluster III, from *Firmicutes* to *Firmicutes* and to

Actinobacteria within Cluster I, but also from *Proteobacteria* to *Firmicutes* within cluster II).

Interestingly, the two wing-less AAT-fold LAOdc coding genes found in *L. saerimneri* 30a were

proposed to result from the horizontal gene transfer of an acid stress inducible Odc from

Enterobacteriaceae, followed by a gene duplication event (Romano *et al.*, 2013, Romano *et al.*,

2014). One of the two resulting paralogues is thought to have kept the original function

([WP 009553942.1](#)), while the other acquired the capacity to use lysine as substrate

([EKW98991.1](#)). Instead, our analysis points to a different scenario. In fact, the two copies

present in *L. saerimneri* branch in two different parts of the OdcI tree (Fig. S3), meaning that

they have different origins, and result likely from two independent horizontal gene transfer

events of OdcI coding genes, whereupon one of these two acquired genes shifted secondarily

towards the capacity to use lysine instead of ornithine. Interestingly, the lysine decarboxylating

copy is present in various *Lactobacilli*, while the copy using ornithine as substrate is specific to *L.*

saerimneri (Fig. S3), suggesting that the former was acquired first. Remarkably, these two

enzymes rely on the same antiporter capable to exchange both ornithine/Put and lysine/Cad

pairs, resulting in a unique three-component decarboxylation system involved in acid stress

response {Romano, 2013 #52}. This case of substrate specificity shift is probably not an

exception, as exemplified by *Burkholderia* sp. AIU 395 in which an AAT-fold enzyme using lysine

as substrate branches within the Adc cluster in phylogenetic trees (Fig. 2A). Beside substrate

shift, existence of dual specificity has been most extensively documented in the case of AR-fold

LOdcs, exemplified by bifunctional enzymes of *S. ruminantium* (Takatsuka *et al.*, 2000) and *V.*

vulnificus (Lee *et al.*, 2007). In particular, crystal structures of the *V. vulnificus* enzyme in

complex with either Put or Cad revealed that the dual substrate specificity is based on a bridging

water molecule necessary for the binding of a shorter Put ligand in addition to the longer Cad. A

similar dual substrate specificity mechanism may also exist in the case of the AAT-fold LAOdc

enzymes although it has not been documented up to now.

Altogether, our data indicate that functional changes affecting gene regulation, substrate

fixation, and cellular function occurred several times during the evolution of AAT-fold enzymes.

In the light of these observations, one may wonder if the so-called Adc, LdcC, LdcI, OdcC, and

OdcI clusters defined according to phylogenetic criteria correspond indeed to homogeneous

functional groups and thus if phylogenetic criteria/sequence similarity based measures are good

predictors of the function of the AAT-fold enzymes. The very restricted number of

experimentally characterized enzymes calls for caution and for the urgent need for additional

experimental data.

One of the main results of the presented phylogenetic analysis is the identification of a

novel large family of decarboxylases, called LdcA, ancestral in *Betaproteobacteria* and

Pseudomonadaceae. Thus, the second part of this work was dedicated to functional

characterization of the LdcA from a major human opportunistic pathogen *P. aeruginosa*.

The *ldcA* gene belongs to the core genome of *P. aeruginosa*, comprising at least 4,000

conserved genes (Hilker *et al.*, 2015, Valot *et al.*, 2015). Similarly to *ldcI*, in *P. aeruginosa*,

Pseudomonas resinovorans, Pseudomonas denitrificans and *Pseudomonas knackmusii*, *ldcA* is

organized in a gene cluster with a gene coding for a homologue of the CadB antiporter,

although *cadB* (*PA1819* in *P. aeruginosa*) appears downstream, and not upstream, of the lysine

decarboxylase-encoding gene. The presence of a dedicated Lys/Cad transporter could be

important from a physiological standpoint because CadB is involved not only in

substrate/product exchange but also in the generation of proton motive force

(Soksawatmaekhin *et al.*, 2004). Remarkably, the proximity of *cadB* and *ldcA* is an exception in

the *ldcA* subfamily (Fig. 4), which does not rule out a hypothesis that LdcA could have another

function in other taxa. The whole of our data suggests that LdcI and *P. aeruginosa* LdcA are

different in terms of regulation and function. Indeed, we clearly show that LdcA is not evolved

for acid or oxidative stress response, and its expression is triggered by neither of these stresses;

instead it is controlled by ArgR and QS. More similar to LdcC, *P. aeruginosa* LdcA, and certainly

the other members of this novel lysine decarboxylase subfamily, are biosynthetic enzymes

responsible for the Cad production by the bacterium.

Importantly, LdcA is the only Cad producing enzyme in *P. aeruginosa* PAO1, as

demonstrated by the measurement of the intracellular contents of Cad (Fig. 6). This conclusion

was unexpected considering that the product of another gene, *PA4115*, was previously reported

to be responsible for 25% of Cad production in overnight cultures grown in the same medium as

the one used in our assays, and therefore proposed to be an Ldc (Manuel *et al.*, 2010). After

carboxypenicillin treatment, the *PA4115* mutant exhibited an increased number of persisters

that was significantly reduced upon addition of exogenous Cad, further supporting the

hypothesis of PA4115 being an Ldc (Manuel *et al.*, 2010). Yet, our careful inspection of its

sequence revealed that PA4115 belongs to the family of Lonely Guy (LOG) proteins because it

contains a highly conserved PGGxGTxxE motif and a nucleotide-binding Rossmann fold.

Remarkably, LOG proteins were shown to be often mis-annotated as lysine decarboxylase

enzymes, yet without support by biochemical or functional data, whereas they actually possess

a cytokinin-specific phosphoribohydrolase activity (Dzurova *et al.*, 2015, Seo & Kim, 2017) or a

pyrimidine/purine nucleotide 5'-monophosphate nucleosidase activity (Sevin *et al.*, 2017).

Recently, PPnN (or YghD) from *E. coli*, a close homologue sharing 56% identity with PA4115, has

been shown to catalyze the hydrolysis of N-glycosidic bond of AMP, GMP, IMP, CMP, dTMP and

UMP to form ribose 5-phosphate and the corresponding free base. Hence, it is quite likely that

PA4115 catalyzes the same reaction and plays a role in maintaining the nucleotide pool

homeostasis by degrading excess nucleotides in *P. aeruginosa* (Sevin *et al.*, 2017). Therefore,

the reduced production of Cad in *PA4115* mutant needs to be confirmed because the

relationship between PA4115 and LdcA activity is not clear and could involve indirect causes.

The persistence phenotype observed in *ldcA* mutants is consistent with the beneficial

effect of LdcA on growth fitness. Indeed, recent research on bacterial persistence uncovered

intracellular ATP concentrations as being one of the major factors affecting the amount of

persister cells, and demonstrated that ATP levels are sufficient to predict bacterial tolerance to

antibiotics (Conlon *et al.*, 2016, Shan *et al.*, 2017). Considering that Cad produced by LdcA is

metabolized and used up by the Krebs cycle to create ATP and that the activity of the lysine/Cad

antiporter generates proton motive force essential for ATP synthesis (Soksawatmaekhin *et al.*,

2004), one would expect that in *P. aeruginosa* the *ldcA* mutation may lead to a decrease in ATP

levels, which in turn would result in an increase of persisters' population. This hypothesis could

be challenged by blocking the Cad degradation pathway or the CadB antiporter activity.

The capacity of Put and Spd to complement the growth phenotype of the *ldcA* mutant

and boost *P. aeruginosa* cultures in minimal medium suggests that in *P. aeruginosa* the three

major polyamines share certain properties. In the same lines, Cad was shown to substitute for

Put and Spd as growth factors in *E. coli* cells depleted of these two polyamines (Igarashi *et al.*,

1986). The growth phenotype described in our work is observed under specific growth

conditions. It reveals the importance of Cad when growing *P. aeruginosa* in minimal medium

and highlights a certain redundancy in the function of polyamines. It remains to be investigated

whether the phenotype is linked to a regulatory effect of Cad or to its anabolism. Recent

literature about *Eikenella corrodens* has pointed out the importance of an AAT-fold Ldc

(belonging to LdcI group) as a virulence factor against eukaryotic cells that acts through

depletion of essential lysine (Lohinai *et al.*, 2015). Therefore the potential role of LdcA in the

virulence of *P. aeruginosa* may warrant further investigation. In the present study, we observed

that the absence of the enzyme did not affect T3SS-dependent cytotoxicity or the mobility

relying on flagellum and Type IV pili (not shown). However, it would also be relevant to probe

the importance of LdcA during mouse infection, where the proper functioning of *P. aeruginosa*

metabolism is primordial for virulence.

Our study reveals that the *ldcA* gene is relatively ancient in *Proteobacteria*, being

ancestral in *Betaproteobacteria* and in *Pseudomonadaceae*, two taxa that cover a wide range of

ecological niches. Information about regulation and function of an enzyme of the previously

unknown LdcA subfamily enables a step further towards understanding of the evolution and the

importance of Cad metabolism in bacteria.

Experimental procedures

Phylogeny: Dataset assembly

Functionally characterized sequences of AAT-fold LAOdcs were retrieved from NCBI: LdcI

(NP_418555.1), LdcC (NP_414728.1), Adc (NP_418541.1), OdcC (NP_417440.1), and OdcI

(NP_415220.1) from *E. coli* str. K-12 substr., MG1655 and LdcA (NP_250509.1) from *P.*

aeruginosa PAO1. These sequences were used as seeds to query a local database containing

4,466 complete proteomes of prokaryotes downloaded from the NCBI

(ftp://ftp.ncbi.nlm.nih.gov) with the BLASTP 2.2.6 software (Altschul *et al.*, 1997) using default

parameters. Homologues of LAODc were retrieved and aligned using MAFFT v7 (Katoh &

Standley, 2013). The resulting multiple alignment was used to build a HMM profile with the

HMMbuild program from the HMMER v3.1b1 package (McClure MA, 1996). This profile was

then used to query the local database of complete proteomes with the HMMsearch program.

Sequences with e-values lower than 2.2e-13 were retrieved. Finally, the search for potential

unannotated sequences was performed using TBLASTN 2.2.6 on genomic sequences

corresponding to the 4,466 complete proteomes. This led to the identification of 4,090

homologous sequences.

Phylogeny: Phylogenetic inference

To limit taxonomic redundancy for phylogenetic analysis, a sampling of the retrieved sequences

by selecting randomly one strain per species has been performed. Multiple alignments were

built with MAFFT using the L-INS-i option that allows the construction of accurate multiple

alignments and trimmed with BMGE v1.1 with matrix substitution BLOSUM30 (Criscuolo &

Gribaldo, 2010).

Maximum likelihood trees were inferred with PhyML 3.1 (Guindon *et al.*, 2010). The best

suited evolutionary models were selected using the model test tool implemented in IQ-TREE

v1.4.1 according to the BIC criteria (Nguyen *et al.*, 2015). The robustness of the inferred trees

was assessed using the non-parametric procedure implemented in PhyML (100 replicates of the

original datasets). Bayesian trees were inferred using MrBayes v3.2.6 (Ronquist & Huelsenbeck,

2003). Two runs were launched with four chains for each run (50,000 iterations). The first 25%

of the trees were discarded as burn-in and chain convergence has been checked by analysing

the evolution of the Ln(L) curve and checking the average standard deviation of split frequencies

values.   Figures   of   trees   have   been   generated   using   EvolView   (He   *et   al.*,   2016)

(http://nar.oxfordjournals.org/content/44/W1/W236)   and   iTOL   (Letunic   &   Bork,   2016).

Genomic   context   figures   have   been   generated   using   GeneSpy   (https://lbbe.univ-

lyon1.fr/GeneSpy/).

Reference phylogenies of species containing Ldc, Adc or Odc homologues were inferred

using   ribosomal   proteins   as   suggested   elsewhere   (Ramulu   *et   al.*,   2014).   Ribosomal   protein

sequences from the complete proteomes of these species were identified using the RiboDB

database engine (Jauffrit *et al.*, 2016) and aligned with MAFFT using the L-INS-i option. The

resulting multiple alignments were trimmed as described above and combined to build a large

supermatrix that was used to build maximum likelihood phylogenetic trees.

Bacterial strains, plasmids, and growth conditions

The bacterial strains and plasmids used in this study are listed in Table S2 in the supplemental

material. Bacteria were cultivated aerobically at 37°C in rich Lysogeny Broth (LB) medium, in

Mueller Hinton II (Becton Dickinson) or in minimal medium P (30mM Na_2HPO_4, 14 mM KH_2PO_4,

20mM $(NH4)_2SO_4$, 1 mM $MgSO_4$, 4µM $FeSO_4$, 0.4 µM Pyridoxal-5'-phosphate, pH 7.4) (Haas *et*

al., 1977) containing the indicated carbon and nitrogen sources. *P. aeruginosa* was also cultured

on Pseudomonas Isolation Agar plates (PIA; Difco). When required, antibiotics were added at

the following concentrations (in µg/ml): 100 (ampicillin), 25 (gentamycin), 25 (kanamycin) and

10 (tetracycline) for *E. coli*, 200 (carbenicillin), 200 (gentamycin) and 200 (tetracycline) for *P.*

aeruginosa.

Genetic manipulations

To delete *ldcA* gene, fused uspstream and downstream flanking regions of the gene were

generated by "Splicing by Overlap Extension" (SOE)-PCR using PAO1 genomic DNA as matrix and

appropriate primer pairs. The resulting fragment of 819 bp was cloned into pCR-Blunt II-TOPO

vector, sequenced and then subcloned into *Bam*HI-*Hind*III sites of pEXG2, leading to

pEXG2Δ*ldcA.* The suicide plasmid carries the counter-selectable *sacB* marker from *B. subtilis*

which confers sensitivity to sucrose. The plasmid was mobilized into *P. aeruginosa* strain by

triparental mating, using the conjugative properties of the helper plasmid pRK2013. Co-

integration events were selected on PIA plates containing gentamycin. Single colonies were then

plated on PIA medium containing 5% (w/v) sucrose to select for the loss of plasmid: the

resulting sucrose-resistant strains were checked for antibiotic sensitivity and for *ldcA* (wild-type

or truncated gene) genotype by PCR.

To complement the *ldcA* mutant, a 2785 bp-long fragment encompassing the coding

sequence and 495 bases upstream the ATG was PCR amplified from PAO1 genomic DNA using

appropriate primer pairs. The PCR product was cloned into pCR-Blunt II-TOPO and sequenced.

The *Spe*I restriction fragment was subcloned in mini-CTX1 cut with the same enzyme, leading to

miniCTX-*ldcA*. A mini-CTX derivative was used to construct a *lacZ* reporter vector. PCR

amplification was used to produce the 548 bp-long *ldcA* promoter fragment (-498/+44 relative

to translation initiation) with appropriate primers. After ligation into pCR-Blunt II-TOPO vector

and sequencing, the fragment was sub-cloned into the *Xho*I-*Eco*RI sites of mini-CTX-lacZ. Both

miniCTX-*ldcA* and mini-CTX-*PldcA-lacZ* were introduced into *P. aeruginosa* by triparental

conjugation and the transconjugants were selected on PIA plates containing tetracycline. The

pFLP2 plasmid was then used to excise the FRT cassette as described (Hoang *et al.*, 1998).

Plasmids and primers used in PCR are listed in the Tables S2 and S3, respectively.

β-Galactosidase assays

Bacteria were grown aerobically at 37°C in 100 ml flasks with agitation (300 rpm). At the

indicated OD, β-galactosidase activity was assayed as described (Miller, 1972), with details

reported in (Thibault *et al.*, 2009).

Intracellular metabolite analysis

Strains were first isolated from an overnight solid culture at 37°C on Mueller Hinton II agar 1.5%.

Precultures and cultures were performed on a Minitron II rotary shaker at 220 rpm (Infors HT)

under aerobic conditions (10% of total volume of Erlenmeyer flask). A few bacterial colonies

from the agar plate were precultured overnight at 37°C and an aliquot was withdrawn and

diluted to an OD_{600} of ~0.1 in a fresh culture medium for the culture step. Growth curves were

determined for each strain and used to determine their respective concentration (CFU/ml) and

the OD at which the bacteria should be harvested to correspond to early-, mid-, and late-

exponential phases.

The protocol of the intracellular metabolites sampling was adapted from a previously

described procedure (Aros-Calt *et al.*, 2015). In brief, a 5 ml aliquot of cell culture broth was

taken from the main culture and was filtered in a few seconds using poly(ether sulfone) sterile

membrane disc filters (Supor450, 0.45 μm pore size, PALL) mounted on a Millipore filtration

device (Darmstadt). The bacteria on the filter were quickly washed with 5 ml of 0.6% NaCl

solution maintained at room temperature. The filter was then rapidly transferred to a 50 ml

Falcon tube containing 5 ml of cold 60% ethanol (v/v, ≤ −20°C). The Falcon tube was

subsequently quickly immersed in liquid nitrogen. Following this quenching step, tubes

containing bacteria on filters in the extraction solution were vortexed 10 times on ice to remove

the cells from the filter. Then, a 1 ml aliquot of the cell suspension was transferred to 2 ml tubes

containing 0.1 mm glass beads (Bertin Technologies). Cell disruption was performed by three

cycles in a Precellys 24 homogenizer (Bertin Technologies) for 30 s at 3,800 rpm at ~4°C. The

glass beads and cell debris were separated from the supernatant by centrifugation for 5 min at

4°C and 10,000g. A 400 μL aliquot of the supernatant was withdrawn and further vacuum-dried

using a SpeedVac instrument (Thermo Fisher Scientific) and stored at −80°C until analysis. Dried

extracts were dissolved in an adjusted volume of 95% mobile phase A / 5% mobile phase B to

obtain the equivalent of 1.25 10^7 CFU in 15μl before analysis by LC/HR-MS.

To detect intracellular metabolites, LC/HR-MS experiments were performed using a Dionex

Ultimate chromatographic system (Thermo Fisher Scientific) coupled to an Exactive (Orbitrap)

mass spectrometer from Thermo Fisher Scientific fitted with an electrospray source. The mass spectrometer was externally calibrated before each analysis using the manufacturer's predefined methods and recommended calibration mixture provided by the manufacturer. Chromatographic separation was performed on a Discovery HS F5 PFPP 5 µm, 2.1 × 250 mm column (Sigma) at 30°C. The chromatographic system was equipped with an on-line pre-filter (Thermo Fisher Scientifics). Mobile phases were 100% water (A) and 100% ACN (B), both of which containing 0.1% formic acid. Chromatographic elution was achieved with a flow rate of 250 µl/min. After injection 15 µl of sample, elution started with an isocratic step of 2 min at 5% phase B, followed by a linear gradient from 5 to 100% of phase B in 18 min. These proportions were kept constant for 4 min before returning to 5% of phase B and letting the system equilibrate for 6 min. The column effluent was directly introduced into the electrospray source of the mass spectrometer, and analyses were performed in the positive ion mode. Source parameters were as follows: capillary voltage set at 5 kV; capillary temperature at 300°C; sheath and auxiliary gas (nitrogen) flow rates at 50 and 25 arbitrary units, respectively; mass resolution power of the analyzer set at 50,000 at m/z 200 (full width at half maximum, FWHM) for singly charged ions. The acquisition was achieved from m/z 50 to 250 in the positive ionization mode during the first 12 min of the run. Under these conditions were achieved a good separation and detection (with an average mass accuracy better than 3ppm) of the targeted molecules (under their $[M+H]^+$ form). These species were readily identified in the extracts through the use of the corresponding commercial molecules obtained from Sigma-Aldrich. Extracted ion chromatograms were generated and resulting peaks integrated using the Xcalibur software (version 2.1, Thermo Fisher Scientific) for Spd ($[M+H]^+$ at theoretical m/z 146.1652, retention

time 5.24 min), Put (m/z 89.1073, 3.63 min) and Cad (m/z 103.1230, 3.94 min).

REFERENCES

Altschul, S.F., T.L. Madden, A.A. Schaffer, J. Zhang, Z. Zhang, W. Miller & D.J. Lipman,
(1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search
programs. *Nucleic Acids Res* **25**: 3389-3402.
Aros-Calt, S., B.H. Muller, S. Boudah, C. Ducruix, G. Gervasi, C. Junot & F. Fenaille, (2015)
Annotation of the Staphylococcus aureus Metabolome Using Liquid Chromatography
Coupled to High-Resolution Mass Spectrometry and Application to the Study of
Methicillin Resistance. *J Proteome Res* **14**: 4863-4875.
Bunsupa, S., K. Katayama, E. Ikeura, A. Oikawa, K. Toyooka, K. Saito & M. Yamazaki, (2012)
Lysine decarboxylase catalyzes the first step of quinolizidine alkaloid biosynthesis and
coevolved with alkaloid production in leguminosae. *Plant Cell* **24**: 1202-1216.
Burrell, M., C.C. Hanfrey, L.N. Kinch, K.A. Elliott & A.J. Michael, (2012) Evolution of a novel
lysine decarboxylase in siderophore biosynthesis. *Mol Microbiol* **86**: 485-499.
Burrell, M., C.C. Hanfrey, E.J. Murray, N.R. Stanley-Wall & A.J. Michael, (2010) Evolution and
multiplicity of arginine decarboxylases in polyamine biosynthesis and essential role in
Bacillus subtilis biofilm formation. *J Biol Chem* **285**: 39224-39238.
Chou, H.T., M. Hegazy & C.D. Lu, (2010) L-lysine catabolism is controlled by L-arginine and
ArgR in Pseudomonas aeruginosa PAO1. *J Bacteriol* **192**: 5874-5880.
Conlon, B.P., S.E. Rowe, A.B. Gandt, A.S. Nuxoll, N.P. Donegan, E.A. Zalis, G. Clair, J.N.
Adkins, A.L. Cheung & K. Lewis, (2016) Persister formation in Staphylococcus aureus is
associated with ATP depletion. *Nat Microbiol* **1**: 16051.
Criscuolo, A. & S. Gribaldo, (2010) BMGE (Block Mapping and Gathering with Entropy): a new
software for selection of phylogenetic informative regions from multiple sequence
alignments. *BMC Evol Biol* **10**: 210.
Dela Vega, A.L. & A.H. Delcour, (1996) Polyamines decrease Escherichia coli outer membrane
permeability. *J Bacteriol* **178**: 3715-3721.
Di Martino, M.L., R. Campilongo, M. Casalino, G. Micheli, B. Colonna & G. Prosseda, (2013)
Polyamines: emerging players in bacteria-host interactions. *Int J Med Microbiol* **303**: 484-
491.
Dzurova, L., F. Forneris, S. Savino, P. Galuszka, J. Vrabka & I. Frebort, (2015) The three-
dimensional structure of "Lonely Guy" from Claviceps purpurea provides insights into the
phosphoribohydrolase function of Rossmann fold-containing lysine decarboxylase-like
proteins. *Proteins* **83**: 1539-1546.
El Bakkouri, M., I. Gutsche, U. Kanjee, B. Zhao, M. Yu, G. Goret, G. Schoehn, W.P. Burmeister
& W.A. Houry, (2010) Structure of RavA MoxR AAA+ protein reveals the design
principles of a molecular cage modulating the inducible lysine decarboxylase activity.
Proc Natl Acad Sci U S A **107**: 22499-22504.

Eliot, A.C. & J.F. Kirsch, (2004) Pyridoxal phosphate enzymes: mechanistic, structural, and
evolutionary considerations. *Annu Rev Biochem* **73**: 383-415.
Fothergill, J.C. & J.R. Guest, (1977) Catabolism of L-lysine by Pseudomonas aeruginosa. *J Gen*
Microbiol **99**: 139-155.
Fritz, G., C. Koller, K. Burdack, L. Tetsch, I. Haneburger, K. Jung & U. Gerland, (2009)
Induction kinetics of a conditional pH stress response system in Escherichia coli. *J Mol*
Biol **393**: 272-286.
Gale, E.F. & H.M. Epps, (1944) Studies on bacterial amino-acid decarboxylases: 1. l(+)-lysine
decarboxylase. *Biochem J* **38**: 232-242.
Gale, E.F. & W.E. Van Heyningen, (1942) The effect of the pH and the presence of glucose
during growth on the production of alpha and theta toxins and hyaluronidase by
Clostridium welchii. *Biochem J* **36**: 624-630.
Guindon, S., J.F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk & O. Gascuel, (2010) New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. *Syst Biol* **59**: 307-321.
Haas, D., B.W. Holloway, A. Schambock & T. Leisinger, (1977) The genetic organization of
arginine biosynthesis in Pseudomonas aeruginosa. *Mol Gen Genet* **154**: 7-22.
Hilker, R., A. Munder, J. Klockgether, P.M. Losada, P. Chouvarine, N. Cramer, C.F. Davenport,
S. Dethlefsen, S. Fischer, H. Peng, T. Schonfelder, O. Turk, L. Wiehlmann, F. Wolbeling,
E. Gulbins, A. Goesmann & B. Tummler, (2015) Interclonal gradient of virulence in the
Pseudomonas aeruginosa pangenome from disease and environment. *Environ Microbiol*
17: 29-46.
Hoang, T.T., R.R. Karkhoff-Schweizer, A.J. Kutchma & H.P. Schweizer, (1998) A broad-host-
range Flp-FRT recombination system for site-specific excision of chromosomally-located
DNA sequences: application for isolation of unmarked Pseudomonas aeruginosa mutants.
Gene **212**: 77-86.
Igarashi, K., (2006) [Physiological functions of polyamines and regulation of polyamine content
in cells]. *Yakugaku Zasshi* **126**: 455-471.
Igarashi, K. & K. Kashiwagi, (2015) Modulation of protein synthesis by polyamines. *IUBMB Life*
67: 160-169.
Igarashi, K., K. Kashiwagi, H. Hamasaki, A. Miura, T. Kakegawa, S. Hirose & S. Matsuzaki,
(1986) Formation of a compensatory polyamine by Escherichia coli polyamine-requiring
mutants during growth in the absence of polyamines. *J Bacteriol* **166**: 128-134.
Jauffrit, F., S. Penel, S. Delmotte, C. Rey, D.M. de Vienne, M. Gouy, J.P. Charrier, J.P. Flandrois
& C. Brochier-Armanet, (2016) RiboDB Database: A Comprehensive Resource for
Prokaryotic Systematics. *Mol Biol Evol* **33**: 2170-2172.
Kamio, Y. & K. Nakamura, (1987) Putrescine and cadaverine are constituents of peptidoglycan
in Veillonella alcalescens and Veillonella parvula. *J Bacteriol* **169**: 2881-2884.
Kamio, Y., H. Poso, Y. Terawaki & L. Paulin, (1986) Cadaverine covalently linked to a
peptidoglycan is an essential constituent of the peptidoglycan necessary for the normal
growth in Selenomonas ruminantium. *J Biol Chem* **261**: 6585-6589.
Kandiah, E., D. Carriel, J. Perard, H. Malet, M. Bacia, K. Liu, S.W. Chan, W.A. Houry, S.
Ollagnier de Choudens, S. Elsen & I. Gutsche, (2016) Structural insights into the
Escherichia coli lysine decarboxylases and molecular determinants of interaction with the
AAA+ ATPase RavA. *Sci Rep* **6**: 24601.
Kang, I.H., J.S. Kim, E.J. Kim & J.K. Lee, (2007) Cadaverine protects Vibrio vulnificus from
superoxide stress. *J Microbiol Biotechnol* **17**: 176-179.

Kanjee, U., I. Gutsche, E. Alexopoulos, B. Zhao, M. El Bakkouri, G. Thibault, K. Liu, S.
Ramachandran, J. Snider, E.F. Pai & W.A. Houry, (2011a) Linkage between the bacterial
acid stress and stringent responses: the structure of the inducible lysine decarboxylase.
EMBO J **30**: 931-944.
Kanjee, U., I. Gutsche, S. Ramachandran & W.A. Houry, (2011b) The enzymatic activities of the
Escherichia coli basic aliphatic amino acid decarboxylases exhibit a pH zone of
inhibition. *Biochemistry* **50**: 9388-9398.
Kanjee, U. & W.A. Houry, (2013) Mechanisms of acid resistance in Escherichia coli. *Annu Rev*
Microbiol **67**: 65-81.
Karatan, E. & A.J. Michael, (2013) A wider role for polyamines in biofilm formation. *Biotechnol*
Lett **35**: 1715-1717.
Katoh, K. & D.M. Standley, (2013) MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**: 772-780.
Kikuchi, Y., O. Kurahashi, T. Nagano & Y. Kamio, (1998) RpoS-dependent expression of the
second lysine decarboxylase gene in Escherichia coli. *Biosci Biotechnol Biochem* **62**:
1267-1270.
Kim, J.S., S.H. Choi & J.K. Lee, (2006) Lysine decarboxylase expression by Vibrio vulnificus is
induced by SoxR in response to superoxide stress. *J Bacteriol* **188**: 8586-8592.
Kim, S.H., Y. Wang, M. Khomutov, A. Khomutov, C. Fuqua & A.J. Michael, (2016) The
Essential Role of Spermidine in Growth of Agrobacterium tumefaciens Is Determined by
the 1,3-Diaminopropane Moiety. *ACS Chem Biol* **11**: 491-499.
Kuper, C. & K. Jung, (2005) CadC-mediated activation of the cadBA promoter in Escherichia
coli. *J Mol Microbiol Biotechnol* **10**: 26-39.
Kwon, D.H. & C.D. Lu, (2006) Polyamines induce resistance to cationic peptide,
aminoglycoside, and quinolone antibiotics in Pseudomonas aeruginosa PAO1. *Antimicrob*
Agents Chemother **50**: 1615-1622.
Lee, J., A.J. Michael, D. Martynowski, E.J. Goldsmith & M.A. Phillips, (2007) Phylogenetic
diversity and the structural basis of substrate specificity in the beta/alpha-barrel fold basic
amino acid decarboxylases. *J Biol Chem* **282**: 27115-27125.
Lee, Y.S. & Y.D. Cho, (2001) Identification of essential active-site residues in ornithine
decarboxylase of Nicotiana glutinosa decarboxylating both L-ornithine and L-lysine.
Biochem J **360**: 657-665.
Letunic, I. & P. Bork, (2016) Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**: W242-245.
Liao, S., P. Poonpairoj, K.C. Ko, Y. Takatuska, Y. Yamaguchi, N. Abe, J. Kaneko & Y. Kamio,
(2008) Occurrence of agmatine pathway for putrescine synthesis in Selenomonas
ruminatium. *Biosci Biotechnol Biochem* **72**: 445-455.
Lightfoot, H.L. & J. Hall, (2014) Endogenous polyamine function--the RNA perspective. *Nucleic*
Acids Res **42**: 11275-11290.
Lohinai, Z., B. Keremi, E. Szoko, T. Tabi, C. Szabo, Z. Tulassay, J.C. DiCesare, C.A. Davis,
L.M. Collins & M. Levine, (2015) Biofilm Lysine Decarboxylase, a New Therapeutic
Target for Periodontal Inflammation. *J Periodontol* **86**: 1176-1184.
Lu, C.D., Z. Yang & W. Li, (2004) Transcriptome analysis of the ArgR regulon in Pseudomonas
aeruginosa. *J Bacteriol* **186**: 3855-3861.
Madhuri Indurthi, S., H.T. Chou & C.D. Lu, (2016) Molecular characterization of lysR-lysXE,
gcdR-gcdHG and amaR-amaAB operons for lysine export and catabolism: a
comprehensive lysine catabolic network in Pseudomonas aeruginosa PAO1. *Microbiology*

162: 876-888.

Malet, H., K. Liu, M. El Bakkouri, S.W. Chan, G. Effantin, M. Bacia, W.A. Houry & I. Gutsche,
(2014) Assembly principles of a unique cage formed by hexameric and decameric E. coli
proteins. *Elife* **3**: e03653.

Manuel, J., G.G. Zhanel & T. de Kievit, (2010) Cadaverine suppresses persistence to
carboxypenicillins in Pseudomonas aeruginosa PAO1. *Antimicrob Agents Chemother* **54**:
5173-5179.

McClure MA, S.C., Elton P., (1996) Parameterization studies for the SAM and HMMER
methods of hidden markov model generation. In: Proceedings / International Conference
on Intelligent Systems for Molecular Biology. pp.

Merrell, D.S. & A. Camilli, (2000) Regulation of vibrio cholerae genes required for acid
tolerance by a member of the "ToxR-like" family of transcriptional regulators. *J Bacteriol*
182: 5342-5350.

Michael, A.J., (2016a) Biosynthesis of polyamines and polyamine-containing molecules.
Biochem J **473**: 2315-2329.

Michael, A.J., (2016b) Polyamines in Eukaryotes, Bacteria, and Archaea. *J Biol Chem* **291**:
14896-14903.

Miller, J., (1972) Experiments in Molecular Genetics. In. C.S.H.L. Press (ed). Cold Spring
Harbor, pp.

Nguyen, L.T., H.A. Schmidt, A. von Haeseler & B.Q. Minh, (2015) IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol*
Evol **32**: 268-274.

Ochsner, U.A., M.L. Vasil, E. Alsabbagh, K. Parvatiyar & D.J. Hassett, (2000) Role of the
Pseudomonas aeruginosa oxyR-recG operon in oxidative stress defense and DNA repair:
OxyR-dependent regulation of katB-ankB, ahpB, and ahpC-ahpF. *J Bacteriol* **182**: 4533-
4544.

Rahman, M. & P.H. Clarke, (1980) Genes and enzymes of lysine catabolism in Pseudomonas
aeruginosa. *J Gen Microbiol* **116**: 357-369.

Ramulu, H.G., M. Groussin, E. Talla, R. Planel, V. Daubin & C. Brochier-Armanet, (2014)
Ribosomal proteins: toward a next generation standard for prokaryotic systematics? *Mol*
Phylogenet Evol **75**: 103-117.

Romano, A., V. Ladero, M.A. Alvarez & P.M. Lucas, (2014) Putrescine production via the
ornithine decarboxylation pathway improves the acid stress survival of Lactobacillus
brevis and is part of a horizontally transferred acid resistance locus. *Int J Food Microbiol*
175: 14-19.

Romano, A., H. Trip, J.S. Lolkema & P.M. Lucas, (2013) Three-component lysine/ornithine
decarboxylation system in Lactobacillus saerimneri 30a. *J Bacteriol* **195**: 1249-1254.

Ronquist, F. & J.P. Huelsenbeck, (2003) MrBayes 3: Bayesian phylogenetic inference under
mixed models. *Bioinformatics* **19**: 1572-1574.

Schuster, M., A.C. Hawkins, C.S. Harwood & E.P. Greenberg, (2004) The Pseudomonas
aeruginosa RpoS regulon and its relationship to quorum sensing. *Mol Microbiol* **51**: 973-
985.

Sekowska, A., P. Bertin & A. Danchin, (1998) Characterization of polyamine synthesis pathway
in Bacillus subtilis 168. *Mol Microbiol* **29**: 851-858.

Seo, H. & K.J. Kim, (2017) Structural basis for a novel type of cytokinin-activating protein. *Sci*
Rep **7**: 45985.

Sevin, D.C., T. Fuhrer, N. Zamboni & U. Sauer, (2017) Nontargeted in vitro metabolomics for

826   high-throughput identification of novel enzymes in Escherichia coli. *Nat Methods* **14**:
827   187-194.
828 Shah, P. & E. Swiatlo, (2008) A multifaceted role for polyamines in bacterial pathogens. *Mol*
829   *Microbiol* **68**: 4-16.
830 Shan, Y., A. Brown Gandt, S.E. Rowe, J.P. Deisinger, B.P. Conlon & K. Lewis, (2017) ATP-
831   Dependent Persister Formation in Escherichia coli. *MBio* **8**.
832 Snider, J., I. Gutsche, M. Lin, S. Baby, B. Cox, G. Butland, J. Greenblatt, A. Emili & W.A.
833   Houry, (2006) Formation of a distinctive complex between the inducible bacterial lysine
834   decarboxylase and a novel AAA+ ATPase. *J Biol Chem* **281**: 1532-1546.
835 Soksawatmaekhin, W., A. Kuraishi, K. Sakata, K. Kashiwagi & K. Igarashi, (2004) Excretion
836   and uptake of cadaverine by CadB and its physiological functions in Escherichia coli. *Mol*
837   *Microbiol* **51**: 1401-1412.
838 Stover, C.K., X.Q. Pham, A.L. Erwin, S.D. Mizoguchi, P. Warrener, M.J. Hickey, F.S.
839   Brinkman, W.O. Hufnagle, D.J. Kowalik, M. Lagrou, R.L. Garber, L. Goltry, E.
840   Tolentino, S. Westbrock-Wadman, Y. Yuan, L.L. Brody, S.N. Coulter, K.R. Folger, A.
841   Kas, K. Larbig, R. Lim, K. Smith, D. Spencer, G.K. Wong, Z. Wu, I.T. Paulsen, J. Reizer,
842   M.H. Saier, R.E. Hancock, S. Lory & M.V. Olson, (2000) Complete genome sequence of
843   Pseudomonas aeruginosa PAO1, an opportunistic pathogen. *Nature* **406**: 959-964.
844 Sugawara, A., D. Matsui, N. Takahashi, M. Yamada, Y. Asano & K. Isobe, (2014)
845   Characterization of a pyridoxal-5'-phosphate-dependent l-lysine decarboxylase/oxidase
846   from Burkholderia sp. AIU 395. *J Biosci Bioeng* **118**: 496-501.
847 Tabor, C.W. & H. Tabor, (1985) Polyamines in microorganisms. *Microbiol Rev* **49**: 81-99.
848 Tabor, H., E.W. Hafner & C.W. Tabor, (1980) Construction of an Escherichia coli strain unable
849   to synthesize putrescine, spermidine, or cadaverine: characterization of two genes
850   controlling lysine decarboxylase. *J Bacteriol* **144**: 952-956.
851 Tabor, H. & C.W. Tabor, (1964) Spermidine, Spermine, and Related Amines. *Pharmacol Rev* **16**:
852   245-300.
853 Takatsuka, Y., Y. Yamaguchi, M. Ono & Y. Kamio, (2000) Gene cloning and molecular
854   characterization of lysine decarboxylase from Selenomonas ruminantium delineate its
855   evolutionary relationship to ornithine decarboxylases from eukaryotes. *J Bacteriol* **182**:
856   6732-6741.
857 Thibault, J., E. Faudry, C. Ebel, I. Attree & S. Elsen, (2009) Anti-activator ExsD forms a 1:1
858   complex with ExsA to inhibit transcription of type III secretion operons. *J Biol Chem* **284**:
859   15762-15770.
860 Valot, B., C. Guyeux, J.Y. Rolland, K. Mazouzi, X. Bertrand & D. Hocquet, (2015) What It
861   Takes to Be a Pseudomonas aeruginosa? The Core Genome of the Opportunistic Pathogen
862   Updated. *PLoS One* **10**: e0126468.
863 Viala, J.P., S. Meresse, B. Pocachard, A.A. Guilhon, L. Aussel & F. Barras, (2011) Sensing and
864   adaptation to low pH mediated by inducible amino acid decarboxylases in Salmonella.
865   *PLoS One* **6**: e22397.
866 Zhao, B. & W.A. Houry, (2010) Acid stress response in enteropathogenic gammaproteobacteria:
867   an aptitude for survival. *Biochem Cell Biol* **88**: 301-314.
868
869

Acknowledgments: we thank Rémi Peyraud and Ina Attrée for helpful discussions. This work

has received funding from GRAL (ANR-10-LABX-49-01), the European Union's Horizon 2020

research and innovation programme under grant agreement No 647784, and the ANR-16-CE02-

0005. This work was also supported by CEA and the French Ministry of Research and National

Research Agency as part of the French metabolomics and fluxomics infrastructure (MetaboHUB,

ANR-11-INBS-0010 grant). Diego Carriel was supported by a PhD grant from LABEX GRAL and

Pierre S. Garcia was supported by a PhD grant from ARC1 Santé Rhône-Alpes Auvergne.

Author contributions: DCL, PSG, FC, and PL performed experiments. All authors analyzed data.

IG, SE and CBA designed and supervised the overall study and wrote the paper with contribution

from all authors.

Abbreviated Summary

Bacterial polyamines are involved in many fundamental processes and are mainly synthetized by

dedicated lysine, arginine and ornithine decarboxylases or LAOdcs. Our exhaustive phylogenetic

analysis reveals evolutionary history of LAOdcs and discloses a hitherto overlooked LdcA

subfamily. We show that LdcA is playing an important role in growth and persistence of the

major multidrug resistant human pathogen *P. aeruginosa*, exerted through cadaverine

biosynthesis and concomitant regulation of intracellular levels of putrescine and spermidine.

Figure legends

Fig. 1. LAOdc is a widespread family in Bacteria

A. Taxonomic distribution in Prokaryotes. Ratio corresponds to the number of proteomes containing at least one homologue on the number of proteomes present in our database for each phylum. Phyla represented by at least 10 proteomes, among which more than 20% contained at least one LAOdc homologue are highlighted by colors.

B. Unrooted ML tree of LAOdcs (PhyML, LG+I+G4 according to BIC, 1,117 sequences, 273 amino-acid positions). The tree can be divided in three parts: Cluster I corresponds to wing-less LAOdc mainly from *Firmicutes*, *Actinobacteria* and *Cyanobacteria*; Cluster II encompasses wing domain containing LAOdc mainly from *Proteobacteria*, while Cluster III gathers wing-less and wing domain-containing LAOdc from various and unrelated taxa. The scale bar represents the number of substitutions per site. Numbers at branches correspond to bootstrap values (BV).

Fig. 2. Phylogeny of proteobacterial LAOdc (Cluster II)

A. ML phylogeny of cluster II LAOdc (PhyML, LG+I+G4 according to BIC, 551 sequences, 589 amino-acid positions). LAOdc sequences from *E. coli* and *P. aeruginosa* are indicated with gray arrows. Red circles represent bootstrap values.

B. Bayesian phylogeny of Cluster II LAOdc sequences inferred from a sample of representative species. The tree is rooted with LAOdc sequences from Cluster I and III (see also Fig. 1) (MrBayes, mixed model+G4, 54 sequences, 392 amino acid positions). The scale bar represents the number of substitutions per site. Numbers at branches represent posterior probabilities values.

Fig. 3. Evolutionary scenario of LAOdcs in *Proteobacteria*

A. Phylogeny in *Proteobacteria* based on ribosomal proteins (PhyML, LG+I+G4 according to BIC, 108 sequences, 6,129 amino acid positions). Green: *Deltaproteobacteria*, light green: *Epsilonproteobacteria*, violet: *Alphaproteobacteria*, red: *Betaproteobacteria*, blue: *Gammaproteobacteria*. Red circles: BV (Bootstrap value), blue diamond: ancestral presence, rectangle: at least one genome possessing at least one copy of LAOdc, green rectangle: ancestral presence, red rectangle: secondary acquisition by horizontal gene transfer. Leaves correspond to families.

B. Evolutive scenario of LAOdc in Enterobacteraceae (PhyML, LG+I+G4 according to BIC, 34 sequences, 6333 amino acid positions). Leaves correspond to genius. Red circles: BV (Bootstrap value), blue diamond: ancestral acquisition, rectangle: at least one genome possessing at least one copy of LAOdc, green rectangle: ancestral presence, red rectangle: secondary acquisition by horizontal gene transfer.

Fig. 4. Genomic context of LAOdc genes from Cluster II

Colored arrows correspond to conserved neighbor genes. Black arrows correspond to LAOdc considered gene.

Fig. 5. Factors influencing *ldcA* expression

Activity of *ldcA* promoter fused to *lacZ* reporter gene was assessed during growth in different media and genetic backgrounds. Measurements of the β-galactosidase activity of PAO1*::PldcA-*

lacZ strain grown either in minimal medium P (MMP) containing 20 mM L-glutamate and 20 mM

arginine (A), or in LB containing or not 20 mM arginine (B) were performed at times indicated.

C. β-galactosidase activity of wild-type and indicated mutant CHA strains harboring the *PldcA*-

lacZ fusion and grown in LB. β-galactosidase activity is expressed in Miller Units (left Y-axis) and

presented in the bar graphs. Growth was performed in 125 ml flasks, followed by measure of

OD_{600} (right Y-axis) and plotted on lines. Results are the average of values from three

independent experiments ± standard deviation (SD).

Fig. 6. Intracellular Cad in rich medium is produced by the lysine decarboxylase LdcA

A. Growth curves of the wild-type strain, the *ldcA* mutant and the complemented strain in the

rich Mueller Hinton medium.

B. Intracellular concentration, expressed in area of chromatogram peak, of the three indicated

polyamines at (1) early-, (2) mid- and (3) late-exponential growth phases as indicated.

Fig. 7. Polyamines are important for growth fitness in minimal medium

Growth of the wild-type PAO1 strain, the *ldcA* mutant and the complemented strain in minimal

MMP medium supplemented with 20 mM glutamate, 1mM arginine and 5 mM of (A) lysine, (B)

Cad, (C) Put or (D) Spd in 96-well plates. The experiments are representative of two

experiments.

Supporting informations

Supplemental material and methods

FIG S1. LAOdcs are ancestral in *Cyanobacteria*. (A) Taxonomic distribution of LAOdc in

Cyanobacteria (PhyML, LG+I+G4 according to BIC, 30 sequences, 6394 amino acid positions).

Leaves correspond to genera. Red circles: BV (Bootstrap value), blue diamond: ancestral

acquisition, rectangle: at least one genome possessing at least one copy of LAOdc, green

rectangle: ancestral presence. (B) Phylogeny of LAOdc of *Cyanobacteria* (PhyML, LG+I+G4

according to BIC, 28 sequences, 446 amino-acid positions). (C) Genomic context of LAOdc in

three *Cyanobacteria*. Black arrows: LAOdc gene.

FIG S2. LAOdcs are ancestral in *Firmicutes*. (A) Taxonomic distribution of LAOdc in *Firmicutes*

(PhyML, LG+I+G4 according to BIC, 38 sequences, 6133 amino-acid positions). Brown:

Natranaerobiales and *Halanaerobiales*, green and blue: *Clostridia*, red: *Bacillales*, pink:

Lactobacillales. Red circles: BV (Bootstrap value), blue diamond: ancestral acquisition, rectangle:

at least one genome possessing at least one copy of LAOdc, green rectangle: ancestral presence,

red rectangle: horizontal acquisition. (B) Phylogeny of LAOdc Cluster I (PhyML, LG+I+G4

according to BIC, 504 sequences, 295 amino-acid positions). Leaves color correspond to species

families. Gray leaves represent non-*Firmicutes* sequences. Ring color correspond to paralogue A

(blue) and B (red). (C) Genomic context of LAOdc A and B in *Firmicutes*. Black arrows: LAOdc

gene, colored arrows: conserved neighbor genes.

FIG S3 Evolutionary origin of *Firmicutes* OdcI sequences from *Proteobacteria*. (PhyML, LG+I+G4

according to BIC, 514 sequences, 583 amino-acid positions). Blue leaves: *Proteobacteria*, red

leaves: *Firmicutes*. Gray arrows indicate the position of the two copies of LAOdc of *Lactobacillus*

saerimneri. Pink: OdcC, green: LdcA, red: Adc, blue: OdcI/C. The scale bar indicates the number

of substitutions per site. Numbers at branches correspond to aLRT supports.

FIG S4 Influence of stress on *ldcA* expression and growth fitness of *P. aeruginosa.* (A) Effect of

acid and oxidative stress during growth in rich medium. At T_0, HCl was added to shift the pH of

LB medium from neutral to pH 5 while H_2O_2 was added to a final entration of 1mM. β-

galactosidase activity of PAO1*::PldcA-lacZ* strain was measured at T_0, then monitored 30 and 60

min after treatment. B) Growth of PAO1 in minimal medium at different pHs was evaluated

using Biolog high-throughput system. Each point corresponds to the "Area Under the Curve"

(AUC) measured after 24h of bacterial growth.

FIG S5 LdcA function affects carbenicillin persistence. Percentage of survivors in rich medium

(cation-adjusted Mueller Hinton Broth) after 24h of carbenicillin treatment. Growth was

performed in erlenmeyer flasks at 37°C with agitation (300 RPM). Percentage of survivors was

calculated from CFU counting after 24h of antibiotic treatment at 500µg/ml (8X MIC).

Table S1 AAT-fold LAOdc homologues in prokaryotes.

Table S2 List of bacterial strains and plasmids used in this work.

Table S3 Primers used in this study.

1002

1003

1004

1005
1006

1007

1008

A

Phylum	ratio
Proteobacteria	323/778
Alphaproteobacteria	30/205
Betaproteobacteria	120/135
Gammaproteobacteria	172/342
Deltaproteobacteria	1/56
Epsilonproteobacteria	0/38

Phylum	ratio
Firmicutes	233/307
Cyanobacteria	39/39
Actinobacteria	62/230
Other bacteria	26/366
Archaea	0/166

B

Cluster I

BV = 99 %

Cluster III

BV = 99 %

Cluster II

■ *Proteobacteria*

■ *Firmicutes*

■ *Cyanobacteria*

■ *Actinobacteria*

■ Others

0.5

A

E. coli str. K12 subtr. MG1655
NP_415220.1
OdcI

E. coli str. K12 subtr. MG1655
NP_417440.1
OdcC

E. coli str. K12 subtr. MG1655
NP_418555.1
LdcI

E. coli str. K12 subtr. MG1655
NP_418541.1
Adc

E. coli str. K12 subtr. MG1655
NP_414728.1
LdcC

P. aeruginosa PAO1
NP_250509.1
LdcA

Cluster
- OdcI
- OdcC
- LdcI
- LdcC
- Adc
- LdcA

Taxon
- Proteobacteria
- Firmicutes
- Cyanobacteria
- Actinobacteria
- Others

B

Gloeobacter kilaueensis JS1 WP 051382630 1
Bacillus methanolicus MGA3 WP 003347112 1
Paenibacillus larvae subsp larvae DSM 25430 WP 023482262 1
Natranaerobius thermophilus JWNMWNLF WP 012446521 1
Mahella australiensis 501 BON WP 041643771 1
Clostridium sticklandii DSM 519 WP 013361042 1
Mycobacterium kansasii ATCC 12478 AGZ53084 1
Streptomyces pristinaespiralis HCCB 10218 WP 005307429 1
Amycolatopsis mediterranei U32 YP 003766478 1
Thermincola potens JR WP 013119009 1
Thermosediminibacter oceani DSM 16646 WP 013276919 1
Clostridium sticklandii DSM 519 WP 013360267 1
Clostridium aceticum DSM 1496 WP 044825303 1
Bacillus halodurans C125 WP 010898791 1
Symbiobacterium thermophilum IAM 14863 WP 011197381 1
Alteromonas stellipolaris LMG 21856 WP 057790397 1
Chromobacterium violaceum ATCC 12472 WP 043595977 1
Myxococcus stipitatus DSM 14675 WP 063639755 1
Aequorivita sublithincola DSM 14238 WP 014783579 1
Halomonas campaniensis LS21 WP 038477757 1
Sphingomonas sp MM1 WP 041864870 1
Sinorhizobium medicae WSM419 YP 001313905 1
Lactobacillus saerimneri 30a WP 009553942 1
Lactobacillus saerimneri 30a EKW98991 1
Cedecea neteri SSMD04 WP 038474637 1
Escherichia coli str K12 substr MG1655 NP 415220 1
Enterobacter lignolyticus SCF1 WP 013367100 1
Serratia marcescens B3R3 WP 060558422 1
Serratia marcescens B3R3 WP 060560469 1
Enterobacter lignolyticus SCF1 WP 013364807 1
Escherichia coli str K12 substr MG1655 NP 417440 4
Cedecea neteri SSMD04 WP 038478940 1
Pseudomonas fulva 12X WP 013791358 1
Pseudomonas aeruginosa PAO1 NP 250509 1
Pusillimonas sp T77 WP 013743231 1
Paraburkholderia xenovorans LB400 WP 011487262 1
Sulfuricella denitrificans skB26 WP 009204871 1
Aromatoleum aromaticum EbN1 WP 011238315 1
Sulfuritalea hydrogenivorans sk43H WP 041100603 1
Streptomyces ambofaciens ATCC 23877 WP 053126122 1
Burkholderia sp AIU 395 BAP47521 1
Sphingomonas sp MM1 WP 015457230 1
Paucibacter sp KCTC 42545 WP 058720126 1
Escherichia coli str K12 substr MG1655 NP 418541 2
Vibrio sp EJY3 WP 014234789 1
Legionella fallonii LLAP10 WP 045094554 1
Aeromonas schubertii WL1483 WP 060586925 1
Serratia marcescens B3R3 WP 004931982 1
Escherichia coli str K12 substr MG1655 NP 418555 1
Enterobacter lignolyticus SCF1 WP 013365210 1
Serratia marcescens B3R3 WP 004931974 1
Cedecea neteri SSMD04 WP 038475417 1
Escherichia coli str K12 substr MG1655 NP 414728 1
Enterobacter lignolyticus SCF1 WP 013367518 1

Tree scale: 0.1