Single-trial neural dynamics are dominated by richly varied movements

Simon Musall*, Matthew T. Kaufman*, Ashley L. Juavinett, Steven Gluf, Anne K. Churchland

When experts are immersed in a task, do their brains prioritize task-related activity? Most efforts to understand neural activity during well-learned tasks focus on cognitive computations and specific task-related movements. We wondered whether task-performing animals explore a broader movement landscape, and how this impacts neural activity. We characterized movements using video and other sensors and measured neural activity using widefield and two-photon imaging. Cortex-wide activity was dominated by movements, especially uninstructed movements, reflecting unknown priorities of the animal. Some uninstructed movements were aligned to trial events. Accounting for them revealed that neurons with similar trial-averaged activity often reflected utterly different combinations of cognitive and movement variables. Other movements occurred idiosyncratically, accounting for trial-by-trial fluctuations that are often considered "noise". This held true for extracellular Neuropixels recordings in cortical and subcortical areas. Our observations argue that animals execute expert decisions while performing richly varied, uninstructed movements that profoundly shape neural activity.

Introduction

Cognitive functions, such as perception, attention, and decision-making, are often studied in the context of movements. This is because most cognitive processes will naturally lead to action: pondering where to go or with whom to interact both ultimately lead to movement. Laboratory studies of cognition therefore often rely on the careful quantification of instructed movements (e.g., key press, saccade or spout lick), to track outcomes of cognitive computations. However, studying cognition in the context of movements leads to a well-known challenge: untangling neural activity that is specific to cognition from neural activity related to the instructed reporting movements.

To address this problem, researchers often isolate the time of decision from the time of movement using a delay period[1–6] or prevent the subject from knowing which movement to use until late in the trial[7–9]. A second strategy is to evaluate and account for the extent to which putative cognitive activity is modulated by metrics of the instructed movement: for instance, assessing how decision-related activity co-varies with saccade parameters[2,10,11] or orienting movements[5]. However, instructed movements are only a subset of possible movements that may modulate neural activity.

In the absence of a behavioral task, uninstructed movements, like running on a wheel, can drive considerable neural activity, even in sensory areas[12–18]. Despite this, uninstructed movements, such as hindlimb flexions during a lick task, are usually not considered when analyzing neural data, exposing two

implicit assumptions (Fig. 1A). The first assumption is that uninstructed movements have a negligible impact
on neural activity compared to task-related activity or instructed movements. The second assumption is
that uninstructed movements occur infrequently and at random times, while instructed movements are task-
aligned, occurring at stereotyped moments on each trial. In this case, uninstructed movements would
increase neural variability trial-to-trial but their effects could be removed by trial averaging[19]. Both
assumptions are largely untested, however, in part due to the difficulty in accurately measuring multiple
movement types and systematically relating them to neural activity.

Revealing the impact of uninstructed movements on neural activity is critical for behavioral experiments:
movements may account for considerable trial-to-trial variance and may also mimic or overshadow the
cognitive processes that are commonly studied in trial-averaged data. It is also unclear whether movements
may drive substantial activity in specific areas (such as association or motor areas) but have negligible
effects in others. We therefore leveraged multiple movement sensors and dual video recordings to track a
wide array of movements and measure their impact on neural activity across the dorsal cortex. A linear
encoding model demonstrated that cortex-wide neural variability was dominated by movements, with
uninstructed movements outpacing the predictive power of instructed movements and task variables. Both
instructed and uninstructed movements also accounted for a large degree of task-aligned activity that was
still present after trial-averaging. Our analysis allowed us to separate out such movement-related activity,
thereby recovering task-related dynamics that were otherwise obscured. Taken together, these results
argue that during cognition, animals engage in a diverse array of uninstructed movements, with profound
consequences for neural activity.
54
55 **Results**
56
Cortex-wide imaging during auditory and visual decision-making

We trained mice to report the spatial position of 0.6-s long sequences of auditory click sounds or visual
LED stimuli. Animals grabbed handles to initiate trials; stimuli were presented at randomized times after the
handle grab (Fig. 1B). Each stimulus was presented twice and after a 1s delay, animals reported a decision
and received a water reward for licking the spout corresponding to the stimulus presentation side (Fig. 1C).
Animals were trained on either auditory or visual stimuli and achieved expert performance in their trained
modality (Fig. 1D). This enabled us to analyze cortical activity in the same detection behavior with different
sensory stimuli.

To measure cortex-wide neural dynamics, we used a custom-built widefield macroscope[20]. Mice were
transgenic, expressing the Ca^{2+}-indicator GCaMP6f in excitatory neurons. Fluorescence was measured
through the cleared, intact skull[21] (Fig. 1E). Using excitation light at two different wavelengths, we isolated
Ca^{2+}-dependent fluorescence and corrected for intrinsic signals (e.g., hemodynamic responses)[22,23]. The
imaging data were then aligned to the Allen Mouse Common Coordinate Framework v3 (CCF, Extended

Data Fig. 1). To confirm accurate CCF alignment, we performed retinotopic visual mapping[24,25] in each

animal and found high correspondence between functionally identified visual areas and the CCF (Extended

Data Fig. 2).

Figure 1. Widefield calcium imaging during auditory and visual decision making.

(A) Schematic for the two main questions, addressed in this work. First: How important are uninstructed movements

compared to instructed movements for driving neural activity during a behavioral task (columns)? Second: To what

extent are instructed and uninstructed movement temporally aligned to the task (rows)? Uninstructed movements are

exemplified as 'hindlimb' but numerous movements are considered throughout this work. **(B)** Bottom view of a mouse

in the behavioral setup. **(C)** Single-trial timing of behavior. Mice held the handles for 1s (±0.25s) to trigger the stimulus

sequence. One second after stimulus end, water spouts moved towards the mice so they could report a choice. **(D)**

Visual experts (blue) had high performance with visual but chance performance with auditory stimuli. Auditory experts

(green) showed the opposite. Thin lines show individual animals, thick show averages. Error bars represent

mean±s.e.m. **(E)** Example image of cortical surface after skull clearing. Overlaid white lines show Allen atlas borders.

(F) Cortical activity during different task episodes averaged over 11 mice. Shown are responses when holding the

handles ('Hold'), visual stimulus presentation ('Stim 1&2'), the subsequent delay ('Delay') and the response period

('Response'). In each trial, stimulus onset was pseudo-randomized within a 0.25-s long time window (inset). **(G)** Left:

Traces show average responses in primary visual cortex (V1), hindlimb somatosensory cortex (HL) and secondary

motor cortex (M2) on the right hemisphere during visual (black) or auditory (red) stimulation. Trial averages are double-

aligned to the time of trial initiation (left dashed line) and stimulus onset (gray bars). Right dashed line indicates

response period, fills indicate s.e.m. Right: d' between visual and auditory trials during first visual stimulus and the

subsequent delay period. **(H)** Same as (G) but for correct visual trials on the left versus right side.

Baseline-corrected fluorescence (ΔF/F) revealed significant modulation of neural activity across dorsal

cortex (Fig. 1F, Video S1; average response to visual trials, 22 sessions from 11 mice). During trial initiation

(handle grab), cortical activity was strongest in sensorimotor areas for hind- and forepaw ('Hold'). The first

visual stimulus caused robust activation of visual areas and weaker responses in medial secondary motor

cortex (M2) ('Stim 1'). Activity in anterior cortex increased during the second stimulus presentation ('Stim

2') and the delay ('Delay'). During the response period, neural activity strongly increased throughout dorsal

cortex ('Response'). A comparison of neural activity across conditions confirmed that neural activity was

modulated by whether the stimulus was visual or auditory (Fig. 1G) and presented on the left or the right

(Fig. 1H). Differences across both conditions were mainly restricted to primary and secondary visual areas

and the posteromedial part of M2.

Movements dominate cortical activity

To assess the impact of movements, we built a linear encoding model[26,27] that could take into account a

large array of instructed and uninstructed movements as well as various task-variables to predict changes

in cortex-wide activity over all trials. We designed model predictors in two ways. The first type of predictor

allowed the model to learn an event kernel for different types of behavioral events (Fig. 2A). Some events

were related to movements, such as licking or grabbing a handle, and others to task variables like animal

choice in a given trial or sensory stimulus onset. For uninstructed movements, we leveraged piezo sensors

underneath the animal to track hindpaw movements and video data to detect movements of the nose or

whisker pad (Fig. 2B). The resulting traces were then thresholded to isolate individual movement events

(Fig. 2C). For all events, we used ridge regression to learn a time-varying event kernel that was time-locked

to the behavioral event and predicted related changes in cortex-wide activity. Event kernels perform well

when the behavioral event is associated with a similar neural response on every repetition and can

recapitulate delayed or multiphasic responses (Fig. 2D). Similar approaches are often used for motion-

correction in neuroimaging[28] and have been successful in fitting complex neural responses to sensory or

motor events during cognitive tasks[26,27]. Our second type of predictor was formed from analog

(continuously-valued) variables. Analog predictors perform well when neural responses scale linearly with

the measured variable, but cannot account for delayed or multiphasic responses (Fig. 2E). Our analog

predictors included a measure of pupil diameter (Fig. 2B) but also video data from two cameras observing

the animal's face and body. For video data, we applied singular value decomposition (SVD) to extract the

200 highest-variance video dimensions and used them as analog predictors to provide additional

information on movements that we had not previously considered[18,29].

In such a model, it is crucial that the predictors are sufficiently decorrelated. This was accomplished through

several sources of trial-to-trial variability, including an experimenter-controlled randomized delay between

the animal grabbing the handles and stimulus onset, as well as natural variability in response times and the

number of actions (licking, etc.) performed. To ensure that video regressors were not redundant with other

movement regressors, we removed overlapping information from the video by orthogonalizing the video

regressors to other movement variables (see Methods). We also orthogonalized all uninstructed movement

variables against instructed movement variables to ensure that uninstructed movement variables only

contained information that could not be captured by instructed movements. To reduce computational cost,

we used SVD on the imaging data and ridge regression to fit the model to data components.

Figure 2. A linear model to reveal behavioral correlates of cortical activity.

(A) Two example trials, illustrating different classes of behavioral events. **(B)** Image of facial video data with 3 movement

variables used in the model. **(C)** Absolute averaged motion energy in the whisker-pad over two trials, showing individual

bouts of movement (top trace). Data was thresholded at two standard deviations (dashed line) to infer whisking events.

For each event, we created a time-shifted design matrix (X_{Whisk}) that was used to compute an event kernel (β_{Whisk}) using

ridge regression. **(D)** Average β_{Whisk} maps -0.3, 0.1 and 0.3 seconds relative to whisk onset. Whisking caused different

responses across cortex, with retrosplenial (RS) being most active 0.1 seconds after whisk onset (bottom red trace)

and barrel cortex (BC) after 0.3 seconds (bottom black trace). **(E)** Illustration of an analog predictor (black) fitted to

cortical activity (gray). Analog predictors are well-suited to account for correlated changes in activity that scale linearly

with a behavioral measure instead of assuming a fixed event response structure (1). In contrast to the event kernel

traces in (D), analog predictors cannot account for neural responses that are shifted in time (2) or include additional

response features (3-4).

To evaluate how well the model captured neural activity at different cortical locations, we computed the 10-

fold cross-validated R^2 (cvR^2) for the full model during different epochs during the trial (Fig. 3A). While some

areas were particularly well predicted in specific trial epochs (e.g., V1 during stimulus presentation), there

was high predictive power throughout the cortex during all epochs of the trial. For all data ('Whole trial'), the

model predicted 41.2±0.9% (mean±s.e.m., *n*=22 sessions) of all variance across cortex.

Figure 3. Linear model predicts cortical activity with highly specific model variables.

(A) Maps of cross-validated explained variance for two task epochs and for the whole trial. **(B)** Example weight maps of the event kernel for right visual stimuli, 0.2 and 1 seconds after stimulus onset. Trace on the right shows, for left V1, the average weights learned by the model. Gray lines indicate times of the maps on the left. **(C)** Same as in (B) but for the event kernel corresponding to right handle grabs, 0 and 0.33 seconds after event onset. **(D)** Same as in (C) but for the event kernel corresponding to nose movements. (**E**) Weight map for the analog pupil regressor. **(F)** Cumulative remaining variance for PCs of model's weight matrix for all video regressors. Black line shows session average, gray line individual sessions. On average, >90% of all variance was explained by 8 dimensions (dashed line). **(G)** Widefield maps corresponding to the sparsened top 6 video-weight dimensions for an example session. **(H)** Influence of each behavioral video pixel on widefield data. The opacity and color of the overlay were scaled between the 0th and 99th percentile over all values.

Spatiotemporal dynamics of the event kernels sensibly matched known roles of sensory and motor cortices. For example, pixels in left primary visual cortex (V1) were highly positive in response to a rightward visual stimulus (Fig. 3B) with temporal dynamics that matched stimulus responses when averaging over visual trials (Fig. 1H). Left somatosensory (FL) and primary motor forelimb areas were activated during right handle grabs (Fig. 3C) and bi-phasic responses in the olfactory bulb (OB) were associated with nose movements (Fig. 3D). In contrast, the analog pupil regressor was associated with changes in multiple cortical areas, resembling previously observed effects of arousal[30] (Fig. 3E). To assess the contribution of video data to the model, we performed principal component analysis (PCA) on the model's weight matrix for all video regressors (Fig. 3F). In many sessions, video weights were at least 8-dimensional, indicating that the video contained different movement patterns that accounted for separate neural responses. To

reveal the diversity of relationships between neural responses and video, the first six dimensions of the

cortical response weights are shown for an example session, after sparsening (Fig. 3G). When projecting

model weights onto video pixels (Fig. 3H), we found that several specific areas of the animal's face were

particularly important, especially around the animal's nose, eye, and jaw. This indicates that video

dimensions contributed additional information to the model that was not already contained in other

movement variables (e.g., nose movements).

We next sought to address which variables were most important for the model's success. The simplest way

to gauge importance is to fit a model consisting of a single variable and ask how well it predicts the data

compared to the full model (Fig. 4A, light green versus both circles combined). However, the cvR^2 maps for

different single-variable models (Fig. 4B, top row) were far less spatially precise than was indicated by their

respective event kernels in the full model (Fig. 3, B-D). This is due to overlap in the explanatory power of

any given variable with other model variables that contain related information (Fig. 4A, light green inside

the black circle). For instance, right visual stimulation also predicted neural activity in the somatosensory

mouth area, as well as different parts of anterior cortex. This is because visual stimulation tends to be

followed by animal movements and licking in the response period that causes increased activity in

corresponding areas. Correspondingly, cvR^2 maps for the right handle and nose were almost identical,

indicating that both might both be associated with other animal movements (Fig. 4B, top right). Single-

variable models are therefore an upper bound for the linear information within a given variable. However,

the same variable may contribute little to the full model's overall predictive power if it is largely redundant

with other model variables. To measure each variable's unique contribution to the model, we created

reduced models in which we shuffled the regressor set corresponding to a particular variable. The resulting

loss of predictive power (ΔR^2) relative to the full model provides a lower bound for the amount of unique

information (non-redundant with any other model variable) that a given variable contributed to the model

(Fig. 4A, dark green). In contrast to cvR^2, ΔR^2 maps (Fig. 4B, bottom row) were highly spatially localized

and matched brain areas that were also most modulated in their corresponding event kernels (Fig. 3). This

was consistent in both visual and auditory experts (Extended Data Fig. 3). Control recordings, from animals

expressing GFP instead of a GCaMP indicator, confirmed that this was not explained by hemodynamic

signals or potential motion artifacts (Extended Data Fig. 4).

We then compared cvR^2 and ΔR^2 values (averaged over all pixels) for all model variables. Many variables

individually predicted a large amount of neural variance (Fig. 4C, light green bars). Surprisingly, movement

variables, both instructed and uninstructed (Fig. 4C, right), contained particularly high predictive power

compared to task variables (Fig. 4C, left). Video ('Video') and video motion energy ('Video ME') were the

most predictive model variables, each explaining ~23% of all variance. Differences were even more striking

for ΔR^2 values, which were particularly low for task variables (Fig. 4C, dark green bars). Consider the 'time'

variable, a regressor set designed to capture signal deviations occurring at consistent times in each trial

(similar to an average over all trials). Although the time-only model captured considerable variance (light

green bar), eliminating this variable had a negligible effect on the model's predictive power (dark green bar
~0). This is because other task variables, such as choice or stimulus, could capture time-varying modulation
equally well. In contrast, movement variables made much larger unique contributions.

To compare the importance of all instructed or uninstructed movements (Fig. 1A, columns) relative to task
variables, we repeated the analysis above on groups of variables (Fig. 4D). Both task and instructed
movement groups contained similar predictive power (cvR^2_{Task}=17.6±0.6%; $cvR^2_{Instructed}$=17.3±0.6%) and
unique contributions (ΔR^2_{Task}=2.9±0.2%; $\Delta R^2_{Instructed}$=3.6±0.2%). Uninstructed movements, however, were
more than twice as predictive ($cvR^2_{Uninstructed}$=38.3±0.9%), with a 5-fold higher unique contribution
($\Delta R^2_{Uninstructed}$=17.8±0.6%). To assess the spatial extent of this large difference, we created pixel-wise ΔR^2
maps for every group (Fig. 4E). Throughout cortex, unique contributions of uninstructed movements were
much higher than either instructed movements or the task group. The contribution of uninstructed
movements was particularly prominent in anterior somatosensory and motor cortices, but also clearly
evident in posterior areas such as retrosplenial cortex and even V1.

These results strongly suggest that uninstructed movements are critical for predicting neural activity across
the dorsal cortex. We considered three possible confounds. First, perhaps the task was insufficiently
complex to properly engage cortex. To address this, we tested animals in a more challenging auditory rate
discrimination task that is known to require parietal and frontal cortices[31,32]. Results were very similar
(Extended Data Fig. 5), arguing that our results were not due to insufficient task complexity. Second, we
asked whether the greater predictive power of uninstructed movements might stem from using analog
predictors. Indeed, the high ΔR^2 for video variables shows that analog predictors are particularly effective
in detecting movements in an unsupervised manner. To address this, we built a version of the model
excluding all analog regressors. Again, uninstructed movements were the most predictive group (Extended
Data Fig. 6), demonstrating that their importance does not hinge on the different kinds of model predictors.

Third, we asked whether uninstructed movements might be important because they simply provide
additional information about instructed movements (e.g., a preparatory body movement before licking)
instead of representing self-generated actions that are independent from the temporal structure of the task.
To address this, we computed ΔR^2 for each group at every time point in the trial (Fig. 4F, Video S2). Here,
$\Delta R^2_{Uninstructed}$ was highest in the baseline period when $\Delta R^2_{Instructed}$ was low (black and blue traces).
Conversely, $\Delta R^2_{Uninstructed}$ was reduced at times when $\Delta R^2_{Instructed}$ was high, arguing against a tight relation
between instructed and uninstructed movements. Instead, their continuously high unique contribution
indicates that uninstructed movements mostly represent spontaneous "fidgets" that are independent from
instructed movements and the temporal structure of the task.

Figure 4. Uninstructed movements dominate cortical activity.

(A) Black circle denotes information of a reduced model, lacking one variable. The single variable (green circle) has information that overlaps with the reduced model (light green) and a unique contribution (dark green) that increases the full model's information. **(B)** Top row: Cross-validated explained variance (cvR^2) maps for different single-variable models. Bottom row: Unique model contribution (ΔR^2) maps for the same variables. **(C)** Explained variance for single model variables, averaged across cortical maps. Shown is either cvR^2 (light green) or ΔR^2 (dark green). Bars represent mean ± s.e.m over 22 sessions. Prev.: previous. **(D)** Explained variance for variable groups. Conventions as in (C). **(E)** ΔR^2 map for each variable group. **(F)** ΔR^2 for variable groups at each time-point, averaged across cortex.

The impact of movements on single-trial and trial-averaged neural activity

To determine how strongly uninstructed movements impacted trial-averaged activity, we asked whether movements were aligned to specific times in the task or only occurred at random times (Fig. 1A, rows). We required instructed movements to occur at specific times (e.g., licking during the response period) and therefore expected them to be strongly task-aligned. Conversely, uninstructed movements like hindpaw flexion might occur at different time points trial-to-trial and therefore be mostly task-independent. If movement timing strongly varies across trials, the hindpaw variable would account for trial-to-trial variability but have no impact on a peri-event time histogram (PETH) that reflects the average over many trials.

To determine the task-aligned and task-independent contributions for each movement variable, we computed the increase in explained variance of the task-only model when a given movement variable was added (Fig. 5A). This is similar to computing unique contributions as described above but here indicates the task-independent contribution (dark blue portion) of a given movement variable. Subtracting this task-independent component from the overall explained variance of each single-movement-variable model (complete blue circle in Fig. 5A, light green bars in Fig. 4C) yields the task-aligned contribution (light blue fill). As expected, instructed movements had a considerable task-aligned contribution, especially the lick variable (Fig. 5B, light blue bars, left). Surprisingly, the same was true for most uninstructed movement variables (Fig. 5B, right), which made considerable task-aligned and task-independent contributions. This was even clearer after pooling variables into groups (Fig. 5C). Here, uninstructed movements contained more task-aligned ($cvR^2_{Instructed}=11.0\pm0.5\%$; $cvR^2_{Uninstructed}=14.5\pm0.5\%$) and task-independent ($\Delta R^2_{Instructed}=6.3\pm0.3\%$; $\Delta R^2_{Uninstructed}=23.8\pm0.7\%$) explained variance than instructed movements. For both movement groups, the amount of task-aligned information was substantial: instructed and uninstructed movements captured ~63% and ~83% of all information in the task model, respectively.

These results have important implications for interpreting both trial-averaged and single-trial data (e.g., in area M2, Fig. 5D). First, although a task-only model could, by construction, accurately predict the PETH (Fig. 5D, left, top), it cannot account for most single-trial fluctuations (left, bottom 3 rows). Instructed movements also predicted parts of the PETH, in particular at trial times when instructed movements were required (near the black dashed lines, middle top). As with the task model, the instructed movement model failed to predict most single-trial fluctuations (middle, bottom 3 rows). In contrast, an uninstructed movement model also predicted the PETH with good accuracy (right top) and was the only one to also accurately predict neural activity on single trials (right, bottom 3 rows). This demonstrates that much of the single-trial activity that is often assumed to be noise or reflect random network fluctuations is instead related to uninstructed movements.

Figure 5. Uninstructed movements make considerable task-aligned and task-independent contributions

(A) Black circle denotes task-model information. A movement variable (blue circle) has information that overlaps with the task (task-aligned, light blue) and a unique contribution (dark blue) that is task-independent. **(B)** Explained variance for all movement variables. Shown is task-independent (dark blue) and task-aligned explained variance (light blue). Values averaged across cortex, bars represent mean ± s.e.m over 22 sessions. **(C)** Task-aligned and task-independent explained variance for movement groups. Conventions as in (B). **(D)** Example M2 data from a single animal (gray traces), predicted by different models that were based on a single variable group. PETHs (top row) are mostly well-explained by all three models. Single trials were only well predicted by uninstructed movements (right column, black). **(E)** The M2 PETH is accurately predicted by the full model (top trace, red). Reconstructing the data based on model weights of each variable group allows partitioning the PETH and revealing the groups' respective contributions to the full model (bottom row). Red dashed arrow indicates an example PETH feature that is best-explained by uninstructed movements. **(F)** Same as in (E) but for V1 data.

Importantly, because many movements are task-aligned, some seemingly cognitive features could be
equally well explained by uninstructed movements (Fig. 5D, top row, red dashed arrows). To resolve this
issue, we developed an approach to "partition" the PETH into respective contributions from the task,
instructed and uninstructed movements, based on how much variance each group accounted for (Fig. 5E).
To do so, we fitted the full model to the trial-by-trial data (gray trace). We then split the full model prediction
(red trace) into three components based on the weights for each group, without re-fitting. This provides the
best estimate of the relative contributions of the task (green traces), instructed movements (blue traces)
and uninstructed movements (black traces) to the PETH while explaining the most trial-by-trial neural
variance. In M2, partitioning the PETH revealed that diverse dynamics, such as increased activity during
the delay (bottom right, red dashed arrow), are best accounted for by uninstructed movements. In contrast,
partitioning a PETH in V1 revealed that most features were reassuringly well-explained by the task as they
are most likely due to a neural response to visual stimulation (Fig. 5F). This analysis therefore allows a
better interpretation of trial-averaged PETHs by using trial-by-trial data to separate task-related activity from
the impact of task-aligned movements.

Taken together, these results highlight that most of the activity without trial-averaging is explained by
uninstructed movements, which is critical to interpret trial-to-trial variability. Furthermore, uninstructed
movements can also occur at predictable time points during a learned behavior and significantly impact the
shape of a PETH, sometimes masquerading as activity related to instructed movements or cognitive
variables. This reveals a conundrum that is difficult to resolve: many PETH features can be explained by
either task or movement variables alone (Fig. 5D top row). Our approach offers insight into this issue:
partitioning the PETH into multiple components of neural activity that jointly recreate the complete shape
(Fig. 5E,F).

The impact of movements on single neuron activity

Widefield imaging reflects the pooled activity over many neurons, especially those in superficial layers[22]
and is affected by dendritic and axonal projections[33]. Although we observed substantial area-level specificity
(Figs. 3,4B), we further extended our approach to single neurons. Using 2-photon imaging, we measured
the activity of over 13,000 single neurons in 10 animals expressing either GCaMP6f (7 mice) or GCaMP6s
(3 mice) at depths spanning 150-450μm. We recorded in 5 functionally identified areas, covering large parts
of the dorsal cortex. In anterior cortex, we targeted M2 and functionally identified the anterior lateral motor
cortex (ALM) and medial motor cortex (MM) based on stereotactic coordinates and their averaged neural
responses to licking[3] (Fig. 6A, top). In posterior cortex we targeted V1 (identified by retinotopic mapping),
retrosplenial (RS) and primary somatosensory cortex (S1) (Fig. 6A, bottom). In each area, average
population activity was modulated as expected for sensory, motor, and association areas (Fig. 6B).
322

Figure 6. Movements are important for interpreting single-neuron data.

(A) Example windows in M2 (top) and posterior cortex (bottom). Colors in top window indicate neural response strength during licking. Licking responses match known locations of areas MM and ALM. Colors in bottom window indicate location of visual areas, based on retinotopic mapping. Top right picture shows example field-of-view in ALM with neurons colored randomly. **(B)** Population PETHs from different cortical areas averaged over all trials and recorded neurons. **(C)** Explained variance for single model variables. Shown is either all explained variance (light green) or unique model contribution (dark green). Values averaged over all neurons, bars represent mean ± s.e.m. over 13770 neurons. Prev.: previous. **(D)** Explained variance for variable groups. Conventions as in (C). Left panel shows results for all cells, right panels show results for individual cortical areas. **(E)** Explained variance of variable groups for individual neurons, sorted by full-model performance (light gray trace). Traces above the horizontal axis reflect all explained variance and traces below show unique model contributions (similar to light and dark green bars in D). Colors indicate variable groups. **(F)** Partitioning of the ALM population PETH. Colors show contributions from different variable groups. Summation of all group contributions results in the original PETH (gray traces). **(G)** PETH modulation indices. Histograms show MIs for each variable group. Dashed lines show MI values for example cells in H. **(H)** PETH partitioning of single neurons. Boxes show example cells, most strongly modulated by the task (green), uninstructed movements (black), instructed movements (blue) or a combination of all three (red). Original PETHs in gray.

We then applied our linear model to the 2-photon data, now aiming to predict single cell activity. In the single-cell data, as in the widefield data, the model had high predictive power, capturing 42.8±0.1% (mean±s.e.m., *n*=13770 neurons) of the variance across all neurons. Again, individual movement variables outperformed task variables (Fig. 6C, light green bars) and made larger unique contributions (dark green bars). Given the known causal role of ALM for licking[4,6], one might expect that the licking variable would be particularly important for predicting single cell activity. Instead, in agreement with our widefield results, almost all movement variables made large contributions. Video-based regressors were again far more powerful than other model variables. The same was true when analyzing cortical areas separately

(Extended Data Fig. 7). When grouping variables, we again found that uninstructed movements had much

higher predictive power (cvR$^2_{Task}$=13.9±0.1%; cvR$^2_{Instructed}$=10.1±0.1%; cvR$^2_{Uninstructed}$=38.8±0.1%) and

made larger unique contributions (ΔR$^2_{Task}$=2.8±0.1%; ΔR$^2_{Instructed}$=1.0±0.1%; ΔR$^2_{Uninstructed}$=25.7±0.1%) (Fig.

6D, left). Across areas, full model performance varied from cvR^2=45.1±0.1% in ALM to cvR^2=30.0±0.5% in

V1 (black dashed lines) potentially due to differences in signal-to-noise ratio across mice. Despite this, the

difference between variable groups was highly consistent (Fig. 6D, right). These differences were not driven

by outliers but found in almost every recorded neuron (Fig. 6E). Uninstructed movements were consistently

the strongest predictors of single neuron activity (Fig. 6E, top black trace) and had the greatest unique

contribution to the model (Fig. 6E, bottom black trace). The above results also held true when controlling

for potential artifacts due to imaging plane motion by excluding frames with substantial motion (Extended

Data Fig. 8).

We again leveraged our ability to partition PETHs into different sources to gain a deeper insight into the

functional tuning of single neurons. As with widefield data, partitioning a PETH into different components

revealed individual contributions from task, instructed and uninstructed movement variables at different

times during the trial. This is exemplified using data averaged over all ALM neurons (Fig. 6F). Average

population activity rises consistently over the trial (gray trace), which, at first glance, could be interpreted

as a single cognitive process that is progressively increasing. However, the partitioning analysis revealed

that this rise reflects the contribution of several components, each with its own dynamics. Task-related

activity explained the early rise during the stimulus presentation. It then plateaued during the delay (top

green trace). Next, the uninstructed movement contribution increased during the delay (middle black trace)

and plateaued before the response period. Finally, the instructed movement contribution increased ~350

ms before the response period. The activity increases contributed by each group are distinct in time and

together create average neural activity that smoothly rises over the entire second half of the trial.

We then examined the PETHs of individual cells. To identify cells that were mostly modulated by one

variable group versus another, we computed three modulation indices (MIs) for each cell, specifying the

extent to which its PETH was best explained by the task, instructed or uninstructed movements (Fig. 6G).

Each MI ranges from 0 to 1, with high MI values indicating stronger PETH modulation due to the variable

group of interest. Individual cells at the extremes of the MI distributions exemplify strong modulation by task

(Fig. 6H, cell 1), uninstructed movements (cell 2) or instructed movements (cell 3). Importantly, the impact

of uninstructed movements could not have been inferred from the shape of the PETH alone: while cell 3

was clearly modulated by an instructed movement (licking in the response period), the PETHs of cell 1 and

cell 2 exhibited similar temporal dynamics. However, partitioning the PETHs revealed that the activity of

these cells reflects very different processes. Notably, this distinction would not have been possible without

measuring and analyzing uninstructed movements. PETH partitioning therefore provides additional insight

for interpreting the trial average and improves isolation of neurons of interest. Aside from identifying neurons

that were purely modulated by a single variable group, we also found that in many cells, we could isolate

task-related dynamics that were otherwise overshadowed by the impact of movements on the PETH (cell
4, Extended Data Fig. 9).

Lastly, we tested whether our approach can be extended to other kinds of data by analyzing
electrophysiological recordings from Neuropixels probes in visual cortex (VC), superior colliculus (SC) and
the midbrain reticular nucleus (MRN) (Fig. 7A). Here, two head-fixed mice were presented with visual
looming stimuli and we used PETH partitioning to separate trial-averaged data into stimulus and movement
components (Fig. 7B). Consistent with our results from V1 (Fig. 5F), trial-averaged neural responses were
well-explained by stimulus predictors with minimal movement contributions. In contrast, across all neurons,
trial-by-trial cvR^2 was far higher for movement variables than for visual stimuli, demonstrating that moment-
by-moment neural activity is mostly related to movements (Fig. 7C). This can be seen in an example neuron
that exhibited slow fluctuations in firing rate during the recording session (~40 minutes, Fig. 7D), which
were not explained by visual stimuli but were highly predictable when using movement regressors. The
same was also true when comparing cvR^2 and ΔR^2 for stimuli and movements in different cortical and
subcortical areas (Fig. 7E). In all recorded brain areas, movements were far more important for predicting
neural activity, indicating that the dominance of movements in driving neural activity is not limited to cortex
but potentially a brain-wide phenomenon.

Figure 7. Uninstructed movements predict single-neuron activity in cortical and subcortical areas
(A) Coronal slice, -3.8 mm from bregma, showing the location of the Neuropixels probe in an example recording. We recorded single-
cell activity from neurons in visual cortex (VC), superior colliculus (SC) and the midbrain reticular nucleus (MRN). **(B)** Population PETH
averaged over all neurons and visual stimuli. Gray traces show recorded data. Top red trace shows model reconstruction. Bottom:
PETH partitioning of modeled data into stimulus (left, green trace) and movement (right black trace) components. **(C)** Explained
variance for all recorded neurons using either the full model (gray) or a movement- (blue) or stimulus-only model (green). **(D)** Activity
of a single example neuron over the entire session (~40 minutes). Gray trace is recorded activity, black is the cross-validated model
reconstruction. **(E)** Explained variance for stimulus or movement variables in different brain areas. Shown is either all explained
variance (light green) or unique model contribution (dark green). Bars represent mean ± s.e.m. over 80, 67 and 85 neurons for VC,
SC and MRN, respectively.

Discussion

Our results demonstrate that cortex-wide neural activity is dominated by movements. By including a wide array of uninstructed movements, our model predicted single-trial neural activity with high accuracy. This was true for thousands of individual neurons, recorded with three different methods, spanning multiple cortical areas and depths. Both instructed and uninstructed movements also had large task-aligned contributions, affecting trial-averaged neural activity. By partitioning PETHs, we separated movement-related from task-related contributions, providing a new tool to unmask truly task-related dynamics and isolate the impact of movements on cortical computations.

Movement-related modulations of neural activity can either reflect the movement itself (due to movement generation, efference copy, or sensory feedback), or changes in internal state that correlate with movements[12,17,34]. Here, we found that movement kernels were highly specific in both cortical areas and temporal dynamics (Fig. 3), indicating that they were probably not strongly reflective of internal state changes. State changes (e.g., arousal) might instead be best captured by analog regressors like the pupil, as indicated by its broad effects across cortex. Video regressors could, in principle, also provide significant internal state information, which may partially explain their particularly high model contributions. However, video weights usually contained at least 8 dimensions that explained neural activity, indicating that it did not solely reflect a low-dimensional state like arousal (Fig. 3). The combined importance of internal states and specific movements highlights the need for tracking multiple movements when assessing their combined impact on neural activity. Using video recordings to track animal movements is a comparably easy way to achieve this goal, especially with new tools that are available for interpreting video data[35–37], and should therefore become standard practice for physiological recordings.

The vast majority of single neuron activity was well-predicted by uninstructed movements, regardless of brain area or depth (Figs. 6D, 7E, Extended Data Fig. 8). The prevalence of such movement modulation across cortex may explain why apparent task-related activity has been observed in a variety of areas[22,38,39] and highlights the importance of additional controls, like neural inactivation, to establish involvement in a given behavior[40]. However, in areas with established causal relevance, movements must still be considered. For instance, in ALM, which is known to play a causal role in similar tasks to the one studied here[3,6,21], many neurons were well-explained by uninstructed movement variables (Fig. 6H). Our partitioning method can help to better interpret such neural activity by identifying neurons that are best explained by task variables. Moreover, for neurons with mixed selectivity for task and movement variables[41], the model can separate their respective contributions and reveal obscured task-related neural dynamics (Extended Data Fig. 9).

Surprisingly, task variables contained far less predictive power than movement variables, although several areas we studied (V1, A1 and ALM) are known to be involved in similar or simpler detection tasks[21,42]. Importantly, this does not indicate that task-related dynamics are absent from cortical circuits but simply

that movements account for a much larger amount of neural variance. Virtually all cortical neurons we
recorded also exhibited mixed selectivity and were strongly modulated by movements regardless of whether
or not they were also modulated by the task (Fig. 6E). Therefore, even for neurons that are strongly tuned
to task variables, cognitive computations are often embedded within a broader context of movement-related
information.

The profound impact of uninstructed movements on neural activity suggests that movement signals play a
critical role in cortical information processing. Widespread integration of movement signals might be
advantageous for sensory processing[15,43], canceling of self-motion[44], gating of inputs[45] or permitting
distributed associational learning[46,47]. All of these functions build on the idea that the brain is creating an
internal model of the world that is based on the integration of sensory and motor information to predict
future events. Our results highlight the notion that such predictive coding is not limited to sensory areas but
might be a common theme across cortex[48]: while sensory areas integrate motion signals to predict expected
future stimuli[13,49], motor areas equally integrate sensory signals to detect deviations between intended and
actual motor state[50]. The principles of comparing current and predicted feedback also extend to other brain
areas like the ventral tegmental area[51] or the cerebellum[52]. This suggests that robust movement
representations may be found throughout the brain.

References

1. Shadlen, M. N. & Newsome, W. T. Motion perception: seeing and deciding. *Proc Natl Acad Sci U S A* **93**, 628–633 (1996).

2. Maimon, G. & Assad, J. A. A cognitive signal for the proactive timing of action in macaque LIP. *Nat. Neurosci.* **9**, 948–955 (2006).

3. Chen, T.-W., Li, N., Daie, K. & Svoboda, K. A Map of Anticipatory Activity in Mouse Motor Cortex. *Neuron* **94**, 866-879.e4 (2017).

4. Guo, Z. V. *et al.* Maintenance of persistent activity in a frontal thalamocortical loop. *Nature* **545**, 181–186 (2017).

5. Erlich, J. C., Bialek, M. & Brody, C. D. A Cortical Substrate for Memory-Guided Orienting in the Rat. *Neuron* **72**, 330–343 (2011).

6. Li, N., Chen, T.-W., Guo, Z. V., Gerfen, C. R. & Svoboda, K. A motor cortex circuit for motor planning and movement. *Nature* **519**, 51–56 (2015).

7. Horwitz, G. D., Batista, A. P. & Newsome, W. T. Representation of an abstract perceptual decision in macaque superior colliculus. *J. Neurophysiol.* **91**, 2281–2296 (2004).

8. Bisley, J. W., Zaksas, D., Droll, J. A. & Pasternak, T. Activity of neurons in cortical area MT during a memory for motion task. *J. Neurophysiol.* **91**, 286–300 (2004).

9. Gold, J. I. & Shadlen, M. N. The influence of behavioral context on the representation of a perceptual decision in developing oculomotor commands. *J. Neurosci.* **23**, 632–651 (2003).

10. Roitman, J. D. & Shadlen, M. N. Response of neurons in the lateral intraparietal area during a combined visual discrimination reaction time task. *J. Neurosci.* **22**, 9475–9489 (2002).

11. Churchland, A. K., Kiani, R. & Shadlen, M. N. Decision-making with multiple alternatives. *Nat Neurosci* **11**, 693–702 (2008).

12. Niell, C. M. & Stryker, M. P. Modulation of Visual Responses by Behavioral State in Mouse Visual Cortex. *Neuron* **65**, 472–479 (2010).

13. Keller, G. B., Bonhoeffer, T. & Hübener, M. Sensorimotor mismatch signals in primary visual cortex of the behaving mouse. *Neuron* **74**, 809–815 (2012).

14. Ayaz, A., Staeuble, A., Saleem, A. B. & Helmchen, F. Layer-specific integration of locomotion and concurrent wall touching in mouse barrel cortex. *bioRxiv* 265165 (2018). doi:10.1101/265165

15. Saleem, A. B., Ayaz, A., Jeffery, K. J., Harris, K. D. & Carandini, M. Integration of visual motion and locomotion in mouse visual cortex. *Nat. Neurosci.* **16**, 1864–1869 (2013).

16. Schneider, D. M., Nelson, A. & Mooney, R. A synaptic and circuit basis for corollary
discharge in the auditory cortex. *Nature* **513**, 189–194 (2014).
17. Vinck, M., Batista-Brito, R., Knoblich, U. & Cardin, J. A. Arousal and Locomotion Make
Distinct Contributions to Cortical Activity Patterns and Visual Encoding. *Neuron* **86**, 740–754
(2015).
18. Stringer, C. *et al.* Spontaneous behaviors drive multidimensional, brain-wide population
activity. *bioRxiv* 306019 (2018). doi:10.1101/306019
19. Bures, J., Petran, M. & Zachar, J. *Electrophysiological Methods in Biological Research*.
(Academic Press, 1967).
20. Ratzlaff, E. H. & Grinvald, A. A tandem-lens epifluorescence macroscope: hundred-fold
brightness advantage for wide-field imaging. *J. Neurosci. Methods* **36**, 127–137 (1991).
21. Guo, Z. V. *et al.* Flow of Cortical Activity Underlying a Tactile Decision in Mice. *Neuron* **81**,
179–194 (2014).
22. Allen, W. E. *et al.* Global Representations of Goal-Directed Behavior in Distinct Cell Types
of Mouse Neocortex. *Neuron* **94**, 891-907.e6 (2017).
23. Wekselblatt, J. B., Flister, E. D., Piscopo, D. M. & Niell, C. M. Large-scale imaging of cortical
dynamics during sensory perception and behavior. *Journal of Neurophysiology* **115**, 2852–
2866 (2016).
24. Marshel, J. H., Garrett, M. E., Nauhaus, I. & Callaway, E. M. Functional Specialization of
Seven Mouse Visual Cortical Areas. *Neuron* **72**, 1040–1054 (2011).
25. Garrett, M. E., Nauhaus, I., Marshel, J. H. & Callaway, E. M. Topography and Areal
Organization of Mouse Visual Cortex. *J. Neurosci.* **34**, 12587–12600 (2014).
26. Runyan, C. A., Piasini, E., Panzeri, S. & Harvey, C. D. Distinct timescales of population
coding across cortex. *Nature* **548**, 92–96 (2017).
27. Scott, B. B. *et al.* Fronto-parietal Cortical Circuits Encode Accumulated Evidence with a
Diversity of Timescales. *Neuron* **95**, 385-398.e5 (2017).
28. Caballero-Gaudes, C. & Reynolds, R. C. Methods for cleaning the BOLD fMRI signal.
NeuroImage **154**, 128–149 (2017).
29. Powell, K., Mathy, A., Duguid, I. & Häusser, M. Synaptic representation of locomotion in
single cerebellar granule cells. *eLife Sciences* **4**, e07290 (2015).
30. Shimaoka, D., Harris, K. D. & Carandini, M. Effects of Arousal on Mouse Sensory Cortex
Depend on Modality. *Cell Reports* **22**, 3160–3167 (2018).
31. Hanks, T. D. *et al.* Distinct relationships of parietal and prefrontal cortices to evidence
accumulation. *Nature* **520**, 220–223 (2015).

32. Erlich, J. C., Brunton, B. W., Duan, C. A., Hanks, T. D. & Brody, C. D. Distinct effects of
prefrontal and parietal cortex inactivations on an accumulation of evidence task in the rat.
eLife Sciences **4**, e05457 (2015).

33. Zhuang, J. *et al.* An extended retinotopic map of mouse cortex. *Elife* **6**, (2017).

34. Polack, P.-O., Friedman, J. & Golshani, P. Cellular mechanisms of brain state–dependent
gain modulation in visual cortex. *Nature Neuroscience* **16**, 1331–1339 (2013).

35. Pereira, T. D. *et al.* Fast animal pose estimation using deep neural networks. *Nature*
Methods **16**, 117 (2019).

36. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. *Nature Neuroscience* **21**, 1281 (2018).

37. Wiltschko, A. B. *et al.* Mapping Sub-Second Structure in Mouse Behavior. *Neuron* **88**, 1121–
1135 (2015).

38. Goard, M. J., Pho, G. N., Woodson, J. & Sur, M. Distinct roles of visual, parietal, and frontal
motor cortices in memory-guided sensorimotor decisions. *eLife* **5**, e13764 (2016).

39. Le Merre, P. *et al.* Reward-Based Learning Drives Rapid Sensory Signals in Medial
Prefrontal Cortex and Dorsal Hippocampus Necessary for Goal-Directed Behavior. *Neuron*
97, 83-91.e5 (2018).

40. Gilad, A., Gallero-Salas, Y., Groos, D. & Helmchen, F. Behavioral Strategy Determines
Frontal or Posterior Location of Short-Term Memory in Neocortex. *Neuron* **99**, 814-828.e7
(2018).

41. Raposo, D., Kaufman, M. T. & Churchland, A. K. A category-free neural population supports
evolving demands during decision-making. *Nat Neurosci* **17**, 1784–1792 (2014).

42. Song, Y.-H. *et al.* A Neural Circuit for Auditory Dominance over Visual Perception. *Neuron*
93, 940-954.e6 (2017).

43. Ayaz, A., Saleem, A. B., Schölvinck, M. L. & Carandini, M. Locomotion controls spatial
integration in mouse visual cortex. *Curr. Biol.* **23**, 890–894 (2013).

44. Sommer, M. A. & Wurtz, R. H. Brain circuits for the internal monitoring of movements. *Annu*
Rev Neurosci **31**, 317 (2008).

45. Schmitt, L. I. *et al.* Thalamic amplification of cortical connectivity sustains attentional control.
Nature **545**, 219–223 (2017).

46. Wang, L., Rangarajan, K. V., Gerfen, C. R. & Krauzlis, R. J. Activation of Striatal Neurons
Causes a Perceptual Decision Bias during Visual Change Detection in Mice. *Neuron* **97**,
1369-1381.e5 (2018).

47. Engel, T. A., Chaisangmongkon, W., Freedman, D. J. & Wang, X.-J. Choice-correlated

activity fluctuations underlie learning of neuronal category representation. *Nature*

Communications **6**, 6454 (2015).

48. Keller, G. B. & Mrsic-Flogel, T. D. Predictive Processing: A Canonical Cortical Computation.

Neuron **100**, 424–435 (2018).

49. Leinweber, M., Ward, D. R., Sobczak, J. M., Attinger, A. & Keller, G. B. A Sensorimotor

Circuit in Mouse Cortex for Visual Flow Predictions. *Neuron* **95**, 1420-1432.e5 (2017).

50. Inoue, M., Uchimura, M. & Kitazawa, S. Error Signals in Motor Cortices Drive Adaptation in

Reaching. *Neuron* **90**, 1114–1126 (2016).

51. Schultz, W. Dopamine neurons and their role in reward mechanisms. *Curr. Opin. Neurobiol.*

7, 191–197 (1997).

52. Wolpert, D. M., Miall, R. C. & Kawato, M. Internal models in the cerebellum. *Trends Cogn.*

Sci. (Regul. Ed.) **2**, 338–347 (1998).

573 **Acknowledgements**

574 We thank Onyekachi Odoemene, Sashank Pisupati and Hien Nguyen for technical assistance and scientific

575 discussions, Hongkui Zeng for providing Ai93 mice, Jason Tucciarone and Fred Marbach for breeding

576 assistance, and Alea Mills and Padmina Shrestha for providing GFP mice. Financial support was received

577 from the Swiss National Science foundation (SM, P2ZHP3_161770), the Pew Charitable Trusts (AKC), the

578 Simons Collaboration on the Global Brain (AKC, MTK) and the NIH (EY R01EY022979).

Methods

Animal Subjects

The Cold Spring Harbor Laboratory Animal Care and Use Committee approved all animal procedures and experiments. Experiments were conducted with male mice from the ages of 8-25 weeks. All mouse strains were of C57BL/6J background and purchased from Jackson Laboratory. Six transgenic strains were used to create the transgenic mice used for imaging: Emx-Cre (JAX 005628), LSL-tTA (JAX 008600), CaMK2α-tTA (JAX 003010), Ai93 (JAX 024103), G6s2 (JAX 024742) and H2B-eGFP (JAX 006069). Ai93 mice as described below were Ai93; Emx-Cre; LSL-tTA; CaMK2α-tTA. G6s2 as described below were CaMK2α-tTA; tetO GCaMP6s.

For widefield imaging we used 11 Ai93 mice (4 mice were negative for CaMK2α-tTA). Two G6s2 and H2B-eGFP mice were used for widefield control experiments. For two-photon imaging, we recorded from 7 Ai93 mice (6 mice were negative for CaMK2α -tTA) and 3 G6s2 mice. All trained mice were housed in groups of two or more under an inverted 12:12-h light-dark regime and trained during their active dark cycle.

Surgical procedures

All surgeries were performed under 1-2% isoflurane in oxygen anesthesia. After induction of anesthesia, 1.2 mg/kg of meloxicam was injected subcutaneously and lidocaine ointment was topically applied to the skin. After making a medial incision, the skin was pushed to the side and fixed in position with tissue adhesive (Vetbond, 3M). We then created an outer wall using dental cement (C&B Metabond, Parkell; Ortho-Jet, Lang Dental) while leaving as much of the skull exposed as possible. A circular headbar was attached to the dental cement. For widefield imaging, after carefully cleaning the exposed skull we applied a layer of cyanoacrylate (Zap-A-Gap CA+, Pacer technology) to clear the bone. After the cyanoacrylate was cured, cortical blood vessels were clearly visible.

For 2-photon imaging, instead of clearing the skull, we performed a circular craniotomy using a biopsy punch. For imaging anterior cortex, a 3-mm wide craniotomy was centered 1.5 mm lateral and 1.5 mm anterior to bregma. For imaging posterior cortex, a 4-mm wide craniotomy was centered 1.7 mm lateral and 2 mm posterior to bregma. We then positioned a circular coverslip of similar size over the cortex and sealed the remaining gap between the bone and glass with tissue adhesive. The coverslip window was then secured to the skull using C&B Metabond (Parkell) and the remaining exposed skull was sealed using dental cement. After surgery, animals were kept on a heating mat for recovery and a daily dose of analgesia (1.2 mg/kg Meloxicam) and antibiotics (2.3 mg/kg Enrofloxacin) were administered subcutaneously for at least 3 days or longer.

For electrophysiology experiments, we used 13-17 week old male mice. Mice were given medicated food

cups (MediGel CPF, Clear H20 74-05-5022) 1-2 days prior to surgery. We performed a small circular

craniotomy over visual cortex using a dental drill. Rather than a circular headbar, a boomerang-shaped

custom Titanium headbar was cemented to the skull, just posterior to the eyes, near Bregma. In addition,

a small ground screw was drilled into the skull over the cerebellum. The probe was mounted onto a 3D

printed piece within an external casing, and affixed to a custom stereotaxic adapter, and lowered into the

brain as previously described[53]. Recordings were performed after the mouse had fully recovered from

surgery (3-4 days).

Behavior

The behavioral setup was based on an Arduino-controlled finite state machine (Bpod r0.5, Sanworks) and

custom Matlab code (2015b, Mathworks) running on a linux PC. Servo motors (Turnigy TGY-306G-HV),

touch sensors and visual stimuli were controlled by microcontrollers (Teensy 3.2, PJRC) running custom

code. Twenty-five mice were trained on a delayed two-alternative forced choice (2AFC) spatial

discrimination task. Mice initiated trials by touching either of two handles with their forepaws. Handles were

mounted on servo motors and were moved out of reach between trials. After one second of holding a

handle, sensory stimuli were presented. Sensory stimuli consisted of either a sequence of auditory clicks,

or repeated presentation of a visual moving bar (3 repetitions, 200 ms each). Auditory stimuli were

presented from either a left or right speaker, and visual stimuli were presented on one of two small LED

displays on the left or right side. The sensory stimulus was presented for 600 ms, there was a 500 ms

pause with no stimulus, and then the stimulus was repeated for another 600 ms. After the second stimulus,

a 1000 ms delay was imposed, then servo motors moved two lick spouts into close proximity of the animal's

mouth. If the animal licked twice to the spout on the same side as the stimulus, he was rewarded with a

drop of water. After one spout was contacted twice, the other spout was moved out of reach to force the

animal to commit to its decision.

Animals were trained over the course of approximately 30 days. After 2-3 days of restricted water access,

animals were head-fixed and received water in the setup. Water was given by presenting a sensory

stimulus, subsequently moving the correct spout close to the animal, then dispensing water automatically.

After several habituation sessions, animals had to touch the handles to trigger the stimulus presentation.

Once animals reliably reached for the handles, the required touch duration was gradually increased up to

1 second. Lastly, the probability for fully self-performed trials, in which both spouts were moved towards

the animal after stimulus presentation, was gradually increased until animals reached stable detection

performance of 80% or higher.

Each animal was trained exclusively on a single modality (6 visual animals, 5 auditory for widefield imaging;

4 auditory for widefield controls; 5 visual animals, 5 auditory animals for 2-photon imaging). Only during

imaging sessions were trials of the untrained modality presented as well. This allowed us to compare neural

activity on trials where animals performed stimulus-guided sensorimotor transformations versus trials where
animal decisions were random. To ensure that detection performance was not overly affected by
presentation of the untrained modality, the trained modality was presented in 75% and the untrained
modality in 25% of all trials.

Behavioral sensors

We used information from several sensors in the behavioral setup to measure different aspects of animal
movement. The handles detected contact with the animal's forepaws, and the lick spouts detected contact
with the tongue, using a grounding circuit. An additional piezo sensor (1740, Adafruit LLC) below the
animal's trunk was used to detect hindpaw and whole-body movements. Sensor data were normalized and
thresholded at 2 standard deviations to extract hindpaw movements. Based on hindpaw events we created
an event-kernel design matrix that was also included in the linear model (see below).

Video monitoring

Two webcams (C920 and B920, Logitech) were used to monitor animal movements. Cameras were
positioned to capture the animal's face (side view) and the body (bottom view). To target particular
behavioral variables of interest, we defined subregions of the video which were then examined in more
detail. These included a region surrounding the eye, the whisker pad and the nose. From the eye region
we extracted changes in pupil diameter using custom Matlab code. To analyze whisker movements, we
computed the absolute temporal derivative averaged over the entire whisker pad. The resulting 1-D trace
was then normalized and thresholded at 2 standard deviations to extract whisking events. Based on
whisking events we created an event-kernel design matrix that was also included in the linear model (see
below). The same approach was used for the nose and pupil diameter.

Widefield imaging

Widefield imaging was done using an inverted tandem-lens macroscope[20] in combination with an sCMOS
camera (Edge 5.5, PCO) running at 60 fps. The top lens had a focal length of 105 mm (DC-Nikkor, Nikon)
and the bottom lens 85 mm (85M-S, Rokinon), resulting in a magnification of 1.24x. The total field of view
was 12.5 x 10.5 mm and the image resolution was 640 x 540 pixels after 4x spatial binning (spatial
resolution: ~20µm/pixel). To capture GCaMP fluorescence, a 525 nm band-pass filter (#86-963, Edmund
optics) was placed in front of the camera. Excitation light was projected on the cortical surface using a 495
nm long-pass dichroic mirror (T495lpxr, Chroma) placed between the two macro lenses. The excitation light
was generated by a collimated blue LED (470 nm, M470L3, Thorlabs) and a collimated violet LED (405 nm,
M405L3, Thorlabs) that were coupled into the same excitation path using a dichroic mirror (#87-063,
Edmund optics). We alternated illumination between the two LEDs from frame to frame, resulting in one set
of frames with blue and the other with violet excitation at 30 fps each. Excitation of GCaMP at 405 nm

results in non-calcium dependent fluorescence[54], allowing us to isolate the true calcium-dependent signal
by rescaling and subtracting frames with violet illumination from the preceding frames with blue illumination.
All subsequent analysis was based on this differential signal at 30 fps.

To ensure that our correction approach effectively removed non-calcium dependent fluorescence, we
performed additional control experiments with two GFP-expressing animals (CAG-H2B-eGFP). Here, the
hemodynamic correction removed ~90% of variance in the data and the model lost most of its predictive
power and spatial specificity (Extended Data Fig. 4). We also imaged two G6s2 mice expressing
GCaMP6s[23]. These animals were expected to exhibit stronger calcium-dependent fluorescence due to
large signal-to-noise ratio (SNR) of the GCaMP6s indicator. Accordingly, the remaining variance after
hemodynamic correction was much higher (Extended Data Fig. 4A-B) and the model predicted even more
variance than in Ai93 mice (Extended Data Fig. 4C-D). The β-weights in modulated areas were also
strongest with GCaMP6s-expressing mice and close to zero in GFP controls (Extended Data Fig. 4F&H).
Together, these controls therefore provide strong evidence that our widefield results were not due to
potential contributions from uncorrected hemodynamic signals.

Two-photon imaging

Two-photon imaging was performed with a resonant-scanning two-photon microscope (Sutter Instruments,
Movable Objective Microscope, configured with the "Janelia" option for collection optics), a Ti:Sapphire
femtosecond pulsed laser (Ultra II, Coherent Inc.), and a 16X 0.8 NA objective (Nikon Instruments). Images
were acquired at 30.9 Hz with an excitation wavelength of 930 nm. All focal planes were between 150-450
μm below the pial surface. The objective height was manually adjusted during the recording in 1-2 μm
increments as often as necessary to maintain the same focal plane.

Preprocessing of neural data

To analyze widefield data, we used SVD to compute the 200 highest-variance dimensions. These
dimensions accounted for at least 90% of the total variance in the data. Using 500 dimensions accounted
for little additional variance (~0.15%), indicating that additional dimensions were mostly capturing recording
noise. SVD returns 'spatial components' U (of size pixels x components), 'temporal components' V^T (of size
components x frames) and singular values S (of size components x components) to scale components to
match the original data. To reduce computational cost, all subsequent analysis was performed on the
product SV^T. SV^T was high-pass filtered above 0.1Hz using a zero-phase, second-order Butterworth filter.
Results of analyses on SV^T were later multiplied with U, to recover results for the original pixel space. All
widefield data was rigidly aligned to the Allen Common Coordinate Framework v3, using four anatomical
landmarks: the left, center, and right points where anterior cortex meets the olfactory bulbs and the medial
point at the base of retrosplenial cortex.

To analyze 2p data we used Suite2P[55] with model-based background subtraction. The algorithm was used
to perform rigid motion correction on the image stack, identify neurons, extract their fluorescence, and
correct for neuropil contamination. ΔF/F traces were produced using the method of Jia et al.[56], skipping the
final filtering step. Using these traces, we produced a matrix of size neurons x time, and treated this similarly
to SV^T above. Finally, we confirmed imaging stability by examining the average firing rate of neurons over
trials. If all neurons varied substantially at the beginning or end of a session, trials containing the unstable
portion were discarded. Recording sessions yielded 188.63±19.28 neurons and contained 378.56±9
performed trials (mean±s.e.m.).

We also used the XY-translation values from the initial motion correction to ask whether motion of the
imaging plane (due to imperfect registration or Z-motion) could contribute to our 2-photon results. To
account for this possibility, we removed all frames that were translated by more than 2 pixels in the X or Y
direction and repeated the analysis (Extended Data Figure 8). In 4983 neurons, we observed negative
unique contributions from the task-group, indicative of model overfitting when removing too many data
frames. We therefore rejected these neurons from further analysis. For the remaining 8787 neurons,
explained variance was highly similar as in our original findings, demonstrating that our results are not
explained by motion of the imaging plane.

To compute trial-averages, imaging data were double-aligned to the time when animals initiated a trial and
to the stimulus onset. After alignment, single trials consisted of 1.8 s of baseline, 0.83 s of handle touch
and 3.3 s following stimulus onset. The randomized additional interval between initiation and stimulus onset
(0 - 0.25 s) was discarded in each trial and the resulting trials of equal length were averaged together.

Neuropixels recordings

To investigate single-neuron responses, a set of linearly expanding (looming, L) or contracting (receding,
R) dots (40 cm/s) were presented above the mouse's head while the mouse was head-restrained but free
to move on a wheel[57]. Stimuli were high or low contrast, and either visual only or with an additional auditory
looming stimulus -- white noise of increasing volume (80 dB at max volume). Eight different types of stimuli
were presented: high and low contrast visual looming (VL), high and low contrast visual receding (VR), high
and low contrast audiovisual looming, and high and low contrast audiovisual receding. Stimuli (20 repeats
of each type) were 0.5 s long and randomly presented, with randomized gaps (more than 4 s) between
stimuli. Results in Fig. 7B were obtained by averaging over all stimulus conditions. A video camera (Basler
AG) in combination with infrared lights was used to record the pupil and face during visual stimulation.
These videos were synchronized with the electrophysiology data for subsequent analysis.

Electrophysiology data was collected with SpikeGLX (Bill Karsh, https://github.com/billkarsh/SpikeGLX).
We recorded from 384 channels spanning ~4 mm of the brain[58]. The data were first median subtracted

across channels and time, sorted with Kilosort spike sorting software[59], and manually curated using phy

(https://github.com/kwikteam/phy). Sorted data were analyzed using custom MATLAB code.

Linear model

The linear model was constructed by combining multiple sets of regressors into a design matrix, to capture

signal modulation by different task or motor events (Fig. 2C). Each regressor set (except for 'analog'

regressors) was structured to capture a time-varying event kernel. To do so, each set included a binary

vector containing a pulse at the time of the relevant event, and numerous copies of this vector each shifted

in time by one frame relative to the original. For sensory stimuli, we created post-event regressor sets

spanning all frames from stimulus onset until the end of the trial. For motor events like licking or whisking,

we created peri-event regressor sets that spanned the frames from 0.5 s before until 1 s after each event.

Lastly, we created whole-trial regressors, covering each frame in a given trial. Whole-trial regressors were

aligned to stimulus onset and contained information about decision variables, such as animal choice or

whether a given trial was rewarded. The model also contained several analog regressors, such as 1-D

regressors for pupil diameter. To capture animal movements, we used SVD to compute the 200 highest

dimensions of video information in both cameras. SVD was performed either on the raw video data ('video')

or the absolute temporal derivative (motion energy; 'video ME'). SVD analysis of behavioral video was the

same as for the widefield data, and we used the product SV^T of temporal components and singular values

as analog regressors in the linear model. We did not use lagged versions of the analog regressors, including

the video regressors.

To use video data regressors, it was important to ensure that their explanatory power did not overlap with

other model variables like licking and whisking that can also be inferred from video data. To accomplish

this, we first created a reduced design matrix X_r, containing all movement regressors as well as times when

spouts or handles were moving or visual stimuli were presented. X_r was ordered so that the motion energy

and video columns were at the end. We then performed a QR decomposition of X_r[60]. The QR decomposition

of a matrix A is $A = QR$, where Q is an orthonormal matrix and R is upper triangular. Columns 1 to j of Q

therefore span the same space as columns 1 to j of A for all j, but all the columns are orthogonal to one

another. Finally, we replaced the motion and video columns of the full design matrix X with the

corresponding columns of Q. This allowed the model to improve the fit to the data using any unique

contributions of the motion and video regressors, while ensuring that the weights given to other regressors

were not altered. Note that the QR decomposition is limited to the original space spanned by X, and

therefore cannot enhance the explanatory power of the video regressors - it can only reduce it.

Subsequently, we used the same approach to orthogonalize all uninstructed movement regressors against

instructed movements. This was done to ensure that all information that could be explained by instructed

movements could not be accounted for by correlated, uninstructed movement regressors. As described

above, this can only reduce the explanatory power of the uninstructed movement regressors.

The following table provides an overview of all model variables and how they were generated:

Variable name	Description	Regressor type	Category
Hindpaw	Piezo sensor below the animal	Analog + Event kernel	Movement
Handles (Left / Right)	Touch events from handle sensors	Event kernel	Movement
Licks (Left / Right)	Lick events from spout sensors	Event kernel	Movement
Pupil	Pupil diameter, extracted from face camera	Analog + Event kernel	Movement
Nose	Nose movements, extracted from face camera	Analog + Event kernel	Movement
Whisking	Whisker movements, extracted from face camera	Analog + Event kernel	Movement
Body	Average motion energy across all body camera pixels	Analog + Event kernel	Movement
Video	Video dimensions from both cameras (SVD)	Analog	Movement
Video ME	Video dimensions from motion energy in both cameras (SVD)	Analog	Movement
Time	All trials	Event kernel	Task
Choice	All leftward choice trials	Event kernel	Task
Previous choice	Every trial after a leftward choice trial	Event kernel	Task

Previous modality	Every trial after a visual trial	Event kernel	Task
Previous success	Every trial after a successful trial	Event kernel	Task
Success	All successful trials	Event kernel	Task
Water given	All frames after a reward was given	Event kernel	Task
Audio (Left / Right)	All frames after a left- or rightward auditory stimulus	Event kernel	Task
Vision (Left / Right)	All frames after a left- or rightward visual stimulus	Event kernel	Task

When a design matrix has columns that are close to linearly dependent (multicollinear), model fits are not
reliable. To test for this, we devised a novel method we call "cumulative subspace angles." The idea is that
for each column of the design matrix, we wish to know how far it lies from the space spanned by the previous
columns (note that pairwise angles do not suffice to determine multicollinearity). Our method works as
follows: (1) the columns of the matrix X were normalized to unit magnitude, (2) a QR decomposition of X
was performed, (3) the absolute value of the elements along the diagonal of R were examined. Each of
these values is the absolute dot product of the original vector with the same vector orthogonalized relative
to all previous vectors. The values range from zero to one, where zero indicates complete degeneracy and
one indicates no multicollinearity at all. Over all experiments, the most collinear regressor received a 0.26,
indicating that it was 15° from the space of all other regressors. The average value was 0.84, corresponding
to a mean angle of 57°.

To avoid overfitting, the model was fit using ridge regression. The regularization penalty was estimated
separately for each column of the widefield data using marginal maximum likelihood estimation[61] with minor
modifications that reduced numerical instability for large regularization parameters.

Variance analysis

Explained variance (cvR^2) was obtained using 10-fold cross-validation. To compute all explained variance
by individual model variables, we created reduced models where all regressors that did not correspond to
a given variable were shuffled in time. The explained variance by each reduced model revealed the
maximum potential predictive power of the corresponding model variable.

To assess unique explained variance by individual variables, we created reduced models for each variable

where only the corresponding regressor set was shuffled in time. The difference in explained variance

between the full and the reduced model yielded the unique contribution ΔR^2 of that model variable. The

same approach was used to compute unique contributions for groups of variables, i.e., 'instructed

movements', 'uninstructed movements' or 'task'. Here, all variables that corresponded to a given group

were shuffled at once.

To compute the 'task-aligned' or 'task-independent' explained variance for each movement variable, we

created a reduced 'task-only' model where all movement variables were shuffled in time. This task-only

model was then compared to other reduced models where all movement variables but one were shuffled.

The difference between the task-only model and this model yielded the task-independent contribution of

that movement variable. The task-aligned contribution was computed as the difference between the total

variance explained by a given variable and its task-independent contribution.

PETH partitioning

Reconstructed trial averages (Figs. 5 and 6) were produced by fitting the full model and averaging the

reconstructed data over all trials. To split the model into the respective contributions of instructed

movements, uninstructed movements and task variables, we reconstructed the data based on that variable

group alone (using the weights from the full model, without re-fitting) and averaging over all trials. To

evaluate the relative impact of each group variable on the perievent time histogram (PETH), we computed

a modulation index (MI), defined as

$$MI_{Group} = \frac{\left(1 + \frac{\Delta Group - \Delta nonGroup}{\Delta Group + \Delta nonGroup} \right)}{2},$$

where ΔGroup and ΔnonGroup denotes the sum of the absolute deviation from zero when reconstructing

the PETH either based on all variables of a given group (ΔGroup) or all other variables (ΔnonGroup). The

MI ranges from 0 (variable group has no impact on PETH) to 1 (PETH is fully explained by variable group).

Intermediate values denote a mixed contribution to the PETH from different group variables (Extended Data

Fig. 9).

Model-based video reconstruction

To better understand how the video related to the neural data, we analyzed the portion of the β-weight

matrix that corresponded to the video regressors. This portion of the matrix was projected back up into the

original video space. The result was of size *p* x *d*, where *p* is the number of video pixels (153,600) and *d* is

the number of dimensions of the widefield data (200). We performed PCA on this matrix, reducing the

number of rows. The top few 'scores' (projections onto the principal components) are low-dimensional

representations of the widefield maps that were most strongly influenced by the video. To choose the

dimensionality, we used the number of dimensions required to account for >90% of the variance (Fig. 3F).

To obtain the widefield maps showing how the video was related to neural activity (Fig. 3G), we projected

the scores back into widefield data pixel space and sparsened them using the varimax rotation. To

determine the influence of each video pixel on the widefield (Fig. 3H), we projected the low-dimensional β-

weights into video pixel space, took the magnitude of the β-weights for each pixel, and multiplied by the

original standard deviation for that pixel (to reverse the Z-scoring step of PCA).

Aberrant cortical activity in Ai93 transgenic animals

Mice with both Emx-Cre and Ai93 transgenes can exhibit aberrant, epileptiform cortical activity patterns,

especially when expressing GCaMP6 during development[62]. To avoid this issue, we raised most of the 11

mice in our widefield data set (6 mice) on a doxycycline-containing diet (DOX), preventing GCaMP6

expression until they were 6 weeks or older. This was also true for all Ai93 mice in our 2-photon data set.

However, 5 mice were raised on standard diet, raising the concern that aberrant activity may have affected

our widefield results.

To test for presence of epileptiform activity, we used the same comparison as Steinmetz et al.[62] on the

cortex-wide average. A peak in cortical activity was flagged as a potential interictal event if it had a width of

60-235 ms and a prominence of 0.03 or higher. These parameters flagged nearly all cases of apparent

interictal events (Extended Data Fig. 10A) and identified four out of the 11 mice in the widefield data set to

exhibit potential epileptiform activity (Extended Data Fig. 10B). None of the identified mice were raised on

DOX.

To ensure that epileptiform activity would not bias our results, we removed flagged events and interpolated

over the resulting gaps (in low-D) with Matlab's built-in autoregressive modeling (fillgaps.m) and a 20-frame

prediction window. The result did not show any perturbations around the former interictal events (Extended

Data Fig. 10C). When comparing modeling results between DOX- and non-DOX-raised mice, predicted

variance was highly similar in all cases (Extended Data Fig. 10D-G). This shows that our results were not

due to epileptiform activity and indicated that it was safe to include all mice in the dataset with this data

processing step.

This was further supported by our additional experiments with GCaMP6s-expressing animals. Consistent

with previous observations[62], we found no epileptiform activity in these mice. Nevertheless, our model

predicted an even larger amount of variance than in Ai93 mice and produced highly-specific maps of unique

contributions from single variables (Extended Data Fig. 4).

Additional References

53. Juavinett, A. L., Bekheet, G. & Churchland, A. K. Chronically-implanted Neuropixels probes enable high yield recordings in freely moving mice. *bioRxiv* 406074 (2018). doi:10.1101/406074

54. Lerner, T. N. *et al.* Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* **162**, 635–647 (2015).

55. Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *bioRxiv* 061507 (2016). doi:10.1101/061507

56. Jia, H., Rochefort, N. L., Chen, X. & Konnerth, A. In vivo two-photon imaging of sensory-evoked dendritic calcium signals in cortical neurons. *Nat Protoc* **6**, 28–35 (2011).

57. Marbach, F. & Zador, A. M. A self-initiated two-alternative forced choice paradigm for head-fixed mice. *bioRxiv* 073783 (2017). doi:10.1101/073783

58. Jun, J. J. *et al.* Fully integrated silicon probes for high-density recording of neural activity. *Nature* **551**, 232–236 (2017).

59. Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M. & D, H. K. Kilosort: realtime spike-sorting for extracellular electrophysiology with hundreds of channels. *bioRxiv* 061481 (2016). doi:10.1101/061481

60. Mumford, J. A., Poline, J.-B. & Poldrack, R. A. Orthogonalization of Regressors in fMRI Models. *PLOS ONE* **10**, e0126255 (2015).

61. Karabatsos, G. Marginal maximum likelihood estimation methods for the tuning parameters of ridge, power ridge, and generalized ridge regression. *Communications in Statistics - Simulation and Computation* (2017).

62. Steinmetz, N. A. *et al.* Aberrant Cortical Activity in Multiple GCaMP6-Expressing Transgenic Mouse Lines. *eNeuro* ENEURO.0207-17.2017 (2017). doi:10.1523/ENEURO.0207-17.2017