Replicate Engineered Virtual Patient Populations as Surrogates for Real Patient-

Level Data

Francis J. Alenghat

Section of Cardiology, Department of Medicine, University of Chicago, Chicago, Illinois, USA

Corresponding Author: Francis J. Alenghat, MD, PhD; University of Chicago; 5841 S. Maryland

Ave, MC 6080; Chicago, IL 60637; USA; Email: falenghat@bsd.uchicago.edu

Short Title: Replicate Engineered Virtual Patient Populations (RE-ViPPs)

Keywords: virtual patients, simulation, open data, clinical trials

Abstract

Objectives: To demonstrate a new method for generating virtual, individual-level data by

testing it on a known clinical trial population.

Design: Virtualization of aggregate data from a clinical trial.

Setting: Virtual

Participants: 936,100 virtual patients

Interventions: None

Main Outcomes Measures: Odds ratios for adverse outcomes in virtual patient populations

compared to clinical trial participants.

Methods: The replicate engineered virtual patient populations (RE-ViPPs) method, based on

aggregate cross-tabulated categorical population data, does not require access to individual-

level data. Using sequential regression combined with randomization, it generates virtual

individual patients to comprise populations that, on average, closely resemble the real

population in question. The method is validated by applying it to aggregated data from the

seminal SPRINT trial, which compared intensive versus standard blood pressure treatment goals

on major adverse cardiovascular events.

Results: The method yields virtual populations, each with 9361 patients, faithfully mimicking

the real SPRINT participants. Multiple logistic regression on 100 such populations shows that

factors with the highest odds ratios for the primary event are, in descending order, past clinical

cardiovascular disease, age ≥ 75, chronic kidney disease, high non-HDL, and smoking history.

Intensive blood pressure treatment, the trial's intervention, had an odds ratio of 0.74 [0.63-

0.87]. On all these measures, the 100 RE-ViPPs mirrored the real SPRINT participants, including

the intensive therapy result (actual SPRINT odds ratio: 0.74 [0.62-0.88]).

Conclusions: Clinical data dissemination has limitations. The most coveted data is descriptive at

the individual level but comes with significant cost, effort, and time. There is potential for

privacy breaches, and the open-data movement has progressed slowly due to data-ownership

concerns. RE-ViPPs closely matched the true SPRINT population. Applied to trials, registries, and

databases, RE-ViPPs could reduce open-data burdens by encouraging dissemination of

aggregate cross-tabulated real data that allow investigators to generate and measure virtual

patients.

Introduction

Patient-level data from clinical trials, registries, and the electronic health record (EHR) is the basis for most clinical research. Obtaining and sharing such data can be challenging, and concerns have been raised about cost, effort, patient privacy, and data ownership [1]. Even journals with strong policies to promote data sharing of published papers have suboptimal adherence by authors [2]. Paying the right amount of attention to who requests individual level data, the purpose of the request, the structure and format of the data, and its de-identification to an extent sufficient to eliminate any chance of deducing identity, can be administratively taxing on the source institution, and often those costs are passed on to the requesting investigator [3-5].

Several entities have taken the initiative to open their data for specific trials and registries, which is a positive step forward. These include a consortium of clinical study sponsors initiating clinicalstudydatarequest.com [6,7], the Yale University Open Data Access (YODA) project [8], Duke's Supporting Open Access Research (SOAR) initiative [9], ImmPort [10], and Project Data Sphere [11]. Some have associated costs in the thousands of dollars whereas others are free. Biolincc is the NHLBI's free source for trial data and has reasonable access [12,13]. Besides clinical trials, sources of data can also include institutional, local, national, and international registries as well as clinical data warehouses driven by EHR data from one or more institutions. Typical time from request to obtaining individual-level de-identified data from one's institutional EHR ranges from weeks to months and costs from hundreds to thousands of dollars [4,5].

However, there are sources of *aggregated* data, based on real clinical populations,

which are currently underutilized for their potential. Public systems such as Center for Disease

Control's WONDER [14], the Healthcare Cost and Utilization Project (HCUPnet) [15], and the

Behavioral Risk Factor Surveillance System Web Enabled Analysis Tool [16] are examples of this

sort of aggregate open data for vital statistics, registries, and survey results, respectively. As for

EHR data, the Informatics for Integrating Biology & the Bedside platform (i2b2) and similar

platforms are deployed by multiple hospital systems to query data in an aggregate fashion at

those institutions; they are mostly promoted as tools for cohort identification for planning

future requests for individual-level data, but they have potential to be more widely used for

research purposes [17-19]. Beyond informing requests for individual level data, these sources of

aggregate data, as long as they can be queried in a richly cross-tabulated fashion, could contain

enough information to approximate real populations. If this can be demonstrated, it would

allow investigators using open data to analyze clinical populations without infringing upon

patient privacy or overly taxing current systems in place, and at the same time, allow

investigators running trials or maintaining registries to continue to have ownership of their

individual-level data.

Here a method is described for creating populations of virtual patients from cross-

tabulated aggregate data. It is applied to the SPRINT trial, a study of 9361 patients randomized

to intensive versus standard blood pressure control, for which the individual-level data

summarized in the original publication [20] was recently 'opened' through NHLBI's Biolincc and

is thus known [21]. SPRINT was chosen because it is a landmark trial with a well-defined

population, and most importantly, provides verified individual-level data that allows for

comparison of the replicate engineered virtual patient populations (RE-ViPPs) generated from

aggregate data with the actual SPRINT population.

Methods

Study data

This is a secondary analysis of SPRINT data, which is provided publicly, in de-identified fashion, by the National Heart, Lung, and Blood Institute (NHLBI) via the Biolincc data repository (https://biolincc.nhlbi.nih.gov/home/). The study was deemed exempt by the University of Chicago IRB and the data was obtained after a signed data use agreement.

In SPRINT, participants were > 50 years old with a systolic blood pressure (SBP) of 130 to 180 mm Hg and had an increased risk of cardiovascular events for at least one of the following four reasons: clinical or subclinical cardiovascular disease, chronic kidney disease (CKD), a 10-year risk of 15% or greater on the basis of the Framingham risk score, or ≥ 75 years old. Patients with diabetes or prior stroke were excluded. Participants were randomized to an SBP target of either less than 140 mm Hg (the standard-treatment group) or less than 120 mm Hg (the intensive-treatment group). The primary composite outcome was myocardial infarction, other acute coronary syndromes, stroke, heart failure, or death from cardiovascular causes, measured over an average of 3.3 years [20].

The SPRINT individual-level data were aggregated by first choosing 9 independent categorical factors, all of which are well-established as influencing cardiovascular outcomes. Of these, eight were categorical in the original data [sex, senior (age≥75), black race, current or former smoker (grouped together in the present study), SBP above the highest tertile at the start of the trial, CKD (eGFR < 60 ml/min/1.73m^2), history of clinical cardiovascular disease, and intensive SBP treatment]. The last (high non-HDL) was determined by calculating the non-HDL

for each patient and identifying those with non-HDL > 160 mg/dL. The dependent variable was

the primary composite outcome, also categorical.

Construction of Replicate Engineered Virtual Patient Population (RE-ViPP)

One hundred RE-ViPPs were created. The starting material for RE-ViPP construction consists of

aggregate data cross-tabulated for every pair of factors to be analyzed. The cross-tabulation is

created by partitioning the entire population into subgroups by the status of one independent

factor and recording, for each subgroup, the prevalence of each of the remaining factors, along

with the rate of the dependent variable (**Supplementary Figure S1**, Steps 1-2). For SPRINT, this

is straightforward, since the individual-level data is known. For EHR, registry, and large trial

databases, this can be done, without giving investigators access to the individual level data, as

long as the database can be queried for two factors at a time. The results of all this partitioning

will yield a cross-tabulation of factor prevalences and outcome rates for each subgroup (as in

Figure 1 for SPRINT).

With this cross-tabulation in hand, to start a RE-ViPP, the exact number of patients with

and without the first independent variable is assigned to match the real population

(**Supplementary Figure S1**, Step 3). In the case of the SPRINT analysis, this was, arbitrarily,

female sex. Then, in those virtual patients with the first factor, the second factor (in SPRINT,

this was senior status) was assigned to the exact percentage measured in the real population

and the same was done for those virtual patients without the first factor. Thus the status of the

first and second chosen factors will match the real population exactly.

Starting with the third factor (**Supplementary Figure S1**, Step 4), linear regression was

performed on the aggregate data using all the previously assigned factors to estimate the

chances of a patient having the new factor as a function of the previously assigned factors. The

β coefficients and y-intercept from this linear regression comprise a formula to calculate a

probability of having the new factor for each virtual patient (Supplementary Materials,

Appendix 1). Significant interactions between pairs of factors are checked in the linear

regression and included if the interaction β is within a factor of 1000 from the other β

coefficients (in SPRINT, no such interactions were found between any pair of the 9 factors).

Once the probability (between 0 and 1) of having the new factor is calculated for each

virtual patient, a random number (between 0 and 1) is generated for each patient

(**Supplementary Figure S1**, Step 5). If the random number is less than the patient's probability

of having the new factor, the patient is assigned as having the new factor. Otherwise, the

virtual patient does not. This repeated for each patient. Once all patients are assigned, the

process repeats for the next factor to be assigned. After sequentially assigning the status for all

the factors for all the virtual patients in the above manner, the outcomes are assigned in the

same manner.

Once all factors and outcomes have been assigned to all virtual patients, the overall rate

of each factor and outcome is measured against the actual population, and the virtual

population is only accepted if within a specified tolerance, for all factors, of the actual

population. If the rate of any factor is outside this tolerance, the random number generation is

repeated for all patients and factors in order to create a new virtual population until a

population is found that is within tolerance for all factors (**Supplementary Figure S1**, Step 6).

For this study of SPRINT, the tolerance was set at ±0.25%. One hundred such acceptable RE-

ViPPs are generated in the above manner.

Analysis of RE-ViPPs

Each of the 100 RE-ViPPs is checked for correlation between the independent variables. It is

then subjected to multiple logistic regression using the nine categorical independent variables

and the primary composite outcome as the dependent variable. Random number generation,

assignment of virtual patient factors, and tolerance testing were conducted in Microsoft Excel

with Macros. The linear and logistic regression analyses were all conducted using R. Figures

were created with ggplot2 in R.

Patient and public involvement

Patients were integral to the SPRINT trial, and patient representatives were involved in the

decision to make the trial data open through the National Heart, Lung, and Blood Institute.

Patients were not involved in the design or implementation of the RE-ViPPs method or in the

interpretation of the results, and results of this study cannot be disseminated directly to study

participants due to the virtual nature of the data. The findings will be disseminated to the

general population through national and international press, with the hope that it will

encourage aggregate open data sharing.

Results

The process for creating RE-ViPPs is presented in the methods section and summarized in

Supplementary Figure S1. The process starts with curating cross-tabulated aggregate data. For

SPRINT, this was obtained by analyzing the individual-level data. However, in its application for

populations where individual-level data is not desired or easily available, the creation of the

cross-tabulated data would come from querying an EHR database (such as via i2b2) or open

aggregate/summary data from registries or clinical trials. **Figure 1** depicts the cross-tabulated

aggregate data from SPRINT.

In the case of RE-ViPPs based on SPRINT, the populations were created based on 9

independent factors and the combined primary outcome event. Each RE-ViPP is composed of

9361 patients, utilizes eight equations (Supplementary Materials, Appendix 1), derived from

linear regression analysis, for estimating the chance of a patient having a factor or outcome,

and requires a total of 74,888 random numbers (9361 virtual patients x 8 random numbers per

virtual patient). The tolerance was set at 0.25%, meaning that the prevalence of each factor and

the primary event rate all had to be within this absolute percentage from the true rate in

SPRINT. If this was not true for a given virtual population, all 9361 virtual patients were

regenerated until the criterion was met. The "SPRINT RE-ViPP Generator" is available for

download (Supplementary Materials).

One hundred RE-ViPPs were created in this randomized fashion, constrained by the

chosen tolerance (each of these RE-ViPPs is available for download in Supplementary

Materials). The rate of each included demographic factor (sex, age≥75, black race), clinical

factor (prior/current smoking, SBP above the highest tertile, non-HDL>160 mg/dL, CKD defined

by eGFR>60, past history of clinical CVD), trial factor (intensive vs standard BP therapy), and the

primary outcome (composite myocardial infarction, other acute coronary syndromes, stroke,

heart failure, or cardiovascular death), as required by the design, all adhere very closely to the

SPRINT rates (**Figure 2**).

Also across all subgroups, the prevalence of each demographic and clinical factor in each

of the 100 virtual patient populations clusters around the actual rates in SPRINT (**Figure 3**). The

virtual patient population is randomized well to intensive and standard BP therapy, just as the

real SPRINT patients, and primary event rates also match across all subgroups (**Figure 3**). It is

worth noting that the SPRINT population has higher female and lower senior prevalence

amongst black participants, lower rates of smoking and higher non-HDL amongst women, more

frequent CKD amongst seniors, and higher event rates in seniors, those with CKD, those with

past clinical CVD, and those randomized to non-intensive treatment. All these features are

preserved in all of the virtual patient populations.

Beyond subgroup characteristics, the RE-ViPPs can be analyzed on the individual level

similar to a multivariate analysis performed on real populations. Correlation matrices show no

strong correlations between the analyzed factors in the RE-ViPPs, just as in the real SPRINT

population (**Supplementary Figure S2**)**.**

Multiple logistic regression performed on each of the 100 virtual populations show that

the odds ratios (ORs) for each factor closely approximates the ORs in SPRINT (**Figure 4**). The

confidence intervals of the RE-ViPP ORs are also similar to SPRINT. A volcano plot shows the OR

plotted against p values for each factor in each RE-ViPP, again demonstrating close adherence

to SPRINT in terms of both magnitude and significance of the findings (**Figure 5**). The risk factors

with the highest odds ratio for the primary event are, in descending order, past clinical

cardiovascular disease, age ≥ 75, chronic kidney disease, high non-HDL, and smoking history.

Factors associated with fewer events are female sex and intensive blood pressure treatment.

Just as with the true SPRINT data, neither black race nor high initial SBP had confidence

intervals that clear the line of neutrality.

Discussion

The approach described here was chosen to demonstrate that meaningful information can be gleaned from certain aggregated data to simulate individual patients. The method extends beyond what is traditionally done with summary information and it can approximate the results from the true individual-level data. It is also distinct from prior methods that rely on individual-level data to create virtual patients for the purposes of physiological experiment simulation or clinical trial simulation [22,23]. It is very important to emphasize that the method cannot be done with any form of aggregate data—the source data must be cross-tabulated by the studied factors and outcomes, and, as such, all the factors should be categorical, either nominal or ordinal. In the current example with SPRINT, the variables were binary, but this process could be extended to incorporate a limited number of non-binary categorical variables, in which case the cross-tabulation should account for all values of those variables. Another important consideration is the quality of the source data. In a clinical trial such as SPRINT, it is likely very reliable, but in EHR databases, variables such as race and smoking status can be variably and inconsistently reported, and diagnoses and outcomes may rely on ICD codes which primarily serve a billing/documentation function and may not be subject to rigorous independent verification [24,25]. Another limitation is that although the method checks and includes any pairwise interaction between variables that is large and significant, it does not address 3-way or more complex interactions. There were no important interactions between the variables chosen for this SPRINT study, and for other studies this limitation could be addressed by checking for such complex interactions during the linear regression analyses and/or by choosing, when possible, sufficiently distinct variables to analyze.

The method is reminiscent of imputation, the process by which missing data for a

variable is replaced by a reasonable prediction, often using regression models, based on the

extant values for that variable as a function of the other variables in the dataset [26,27]. This

has perhaps taken its most robust form in multiple imputation of chained equations [28,29].

The method here is distinct from multiple imputation in several ways, but an analogy would be

that every patient's status for all the variables beyond the first two are treated as missing, and

they are imputed sequentially based on logistic regression analysis of the percentage

distributions in the aggregate data.

The current study is meant to serve as a practical proof-of-principle example. Data

known at the individual level was used to highlight the method and show that the RE-ViPPs

result matches that of the real population. In practice, this method would be most useful in

situations when the individual-level data is either impossible or difficult to obtain. These are

common situations. In many cases, there are concerns about data ownership or patient privacy

[30]. In others, it can also involve excess time, effort, and cost [3,31]. Still in others, the issue

may be one of data standardization, where a study would ideally be done across datasets or

platforms, but the data exist in incongruent formats in each platform [32]. Also, the sheer bulk

of individual-level data may be prohibitive, particularly when there is a desire to combine data

from multiple EHRs across the world. With these applications in mind, the "SPRINT RE-ViPP

Generator" is provided in the Supplementary Materials and can be customized to other sets of

aggregate data and variables.

There are many recent high profile examples, not only in healthcare, of bulk individual-

level data being hacked or leaked with major fallout [33,34]. In all these situations, keeping the

individual level data undisclosed while providing the requisite summary statistics could allow

other investigators to study the data while abrogating these concerns. One way this could be

done is to keep the individual level data behind a firewall but allow conditional queries to

extract summary data – this is what the i2b2 platform and others do for EHR data warehouses,

what Center for Disease Control's WONDER provides for vital statistics and other limited clinical

information, and what the Behavioral Risk Factor Surveillance System Web Enabled Analysis

Tool does for its large national health survey results [14,16,17,19]. Another approach would be

to have the "data owners" provide the richly cross-tabulated results across as many categorical

variables as possible as a small static file (like **Figure 1**) and make it open to interested

investigators; this approach would probably be more feasible and require less digital

infrastructure for individual clinical trial results in which full data releases are not planned. This

could also allay concerns about the provenance of the source data in secondary analysis, as the

shared information would be more succinct and verifiable. In all the above scenarios, it is worth

emphasizing that none of the analyzed individual patients derived from this method are real.

Patients in the real population are well represented by the virtual patients, but as digital

versions of creation, the virtual patients are nameless composites only [35], with no

opportunity to deduce identity.

This method should allow us to analyze trial, registry, and routinely collected clinical

information over large populations to find potential connections between heterogeneous

clinical and demographic factors in a novel way that does not directly access patient-level data.

The method could reduce barriers that currently impede access and use of open data. It would

thereby stimulate more meaningful use of extant information, ultimately to generate new

hypotheses for further investigation in real individuals and to identify previously unrecognized

clinical relationships.

Acknowledgements

Author Contributions: Dr. Alenghat had full access to the aggregate data, preformed the

analysis, and takes responsibility for its accuracy

Competing Interests: none

Funding: This work was supported, in part, by the National Heart, Lung, and Blood Institute,

grant number HL116600.

Data sharing statement: Access to SPRINT data is available from the National Heart, Lung and

Blood Institute via Biolinnc. Access to the 100 RE-ViPPs is provided in the Supplementary

Materials.

References

1. Rockhold F, Nisen P, Freeman A. Data Sharing at a Crossroads. N Engl J Med. 2016;375(12):1115-7. PMID: 27653563.

2. Naudet F, Sakarovitch C, Janiaud P, Cristea I, Fanelli D, Moher D, Ioannidis JPA. Data sharing and reanalysis of randomized controlled trials in leading biomedical journals with a full data sharing policy: survey of studies published in The BMJ and PLOS Medicine. BMJ. 2018;360:k400. PMID: 29440066; PMCID: PMC5809812.

3. Sebaa A, Chikh F, Nouicer A, Tari A. Medical Big Data Warehouse: Architecture and System Design, a Case Study: Improving Healthcare Resources Distribution. J Med Syst. 2018;42(4):59. PMID: 29460090.

4. University of Chicago Clinical Research Data Warehouse (CRDW) [3/28/2018]. Available from: http://cri.uchicago.edu/crdw/.

5. UCSF Request Clinical Data for Research [3/28/2018]. Available from: https://data.ucsf.edu/data-assets/clinical-research.

6. Vaduganathan M, Nagarur A, Qamar A, Patel RB, Navar AM, Peterson ED, Bhatt DL, Fonarow GC, Yancy CW, Butler J. Availability and Use of Shared Data From Cardiometabolic Clinical Trials. Circulation. 2018;137(9):938-47. PMID: 29133600; PMCID: PMC5828960.

7. Nisen P, Rockhold F. Access to patient-level data from GlaxoSmithKline clinical trials. N Engl J Med. 2013;369(5):475-8. PMID: 23902490.

8. Krumholz HM, Waldstreicher J. The Yale Open Data Access (YODA) Project--A Mechanism for Data Sharing. N Engl J Med. 2016;375(5):403-5. PMID: 27518657.

9. Pencina MJ, Louzao DM, McCourt BJ, Adams MR, Tayyabkhan RH, Ronco P, Peterson ED. Supporting open access to clinical trial data for researchers: The Duke Clinical Research Institute-Bristol-Myers Squibb Supporting Open Access to Researchers Initiative. Am Heart J. 2016;172:64-9. PMID: 26856217.

10. Bhattacharya S, Andorf S, Gomes L, Dunn P, Schaefer H, Pontius J, Berger P, Desborough V, Smith T, Campbell J, Thomson E, Monteiro R, Guimaraes P, Walters B, Wiser J, Butte AJ. ImmPort: disseminating data to the public for the future of immunology. Immunol Res. 2014;58(2-3):234-9. PMID: 24791905.

11.   Green AK, Reeder-Hayes KE, Corty RW, Basch E, Milowsky MI, Dusetzina SB, Bennett AV, Wood
WA. The project data sphere initiative: accelerating cancer research by sharing data. Oncologist.
2015;20(5):464-e20. PMID: 25876994; PMCID: PMC4425388.
12.   Ross JS, Ritchie JD, Finn E, Desai NR, Lehman RL, Krumholz HM, Gross CP. Data sharing through
an NIH central database repository: a cross-sectional survey of BioLINCC users. BMJ Open.
2016;6(9):e012769. PMID: 27670522; PMCID: PMC5051517.
13.   Coady SA, Wagner E. Sharing individual level data from observational studies and clinical trials: a
perspective from NHLBI. Trials. 2013;14:201. PMID: 23837497; PMCID: PMC3750470.
14.   Friede A, Rosen DH, Reid JA. CDC WONDER: a cooperative processing architecture for public
health. J Am Med Inform Assoc. 1994;1(4):303-12. PMID: 7719813; PMCID: PMC116209.
15.   Weiss AJ, Elixhauser A, Steiner C. Readmissions to U.S. Hospitals by Procedure, 2010: Statistical
Brief #154. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville (MD)2006.
16.   Fisher MA, Ma ZQ. Multiple chronic conditions: diabetes associated with comorbidity and
shared risk factors using CDC WEAT and SAS analytic tools. J Prim Care Community Health.
2014;5(2):112-21. PMID: 24327591.
17.   Klann JG, Buck MD, Brown J, Hadley M, Elmore R, Weber GM, Murphy SN. Query Health:
standards-based, cross-platform population health surveillance. J Am Med Inform Assoc.
2014;21(4):650-6. PMID: 24699371; PMCID: 4078284.
18.   Murphy SN, Gainer V, Mendis M, Churchill S, Kohane I. Strategies for maintaining patient privacy
in i2b2. J Am Med Inform Assoc. 2011;18 Suppl 1:i103-8. PMID: 21984588; PMCID:
PMC3241166.
19.   Murphy SN, Weber G, Mendis M, Gainer V, Chueh HC, Churchill S, Kohane I. Serving the
enterprise and beyond with informatics for integrating biology and the bedside (i2b2). J Am Med
Inform Assoc. 2010;17(2):124-30. PMID: 20190053; PMCID: 3000779.
20.   Group SR, Wright JT, Jr., Williamson JD, Whelton PK, Snyder JK, Sink KM, Rocco MV, Reboussin
DM, Rahman M, Oparil S, Lewis CE, Kimmel PL, Johnson KC, Goff DC, Jr., Fine LJ, Cutler JA,
Cushman WC, Cheung AK, Ambrosius WT. A Randomized Trial of Intensive versus Standard
Blood-Pressure Control. N Engl J Med. 2015;373(22):2103-16. PMID: 26551272; PMCID:
PMC4689591.
21.   Burns NS, Miller PW. Learning What We Didn't Know - The SPRINT Data Analysis Challenge. N
Engl J Med. 2017;376(23):2205-7. PMID: 28445656.

22. Teutonico D, Musuamba F, Maas HJ, Facius A, Yang S, Danhof M, Della Pasqua O. Generating

Virtual Patients by Multivariate and Discrete Re-Sampling Techniques. Pharm Res.

2015;32(10):3228-37. PMID: 25994981; PMCID: PMC4577546.

23. Resalat N, El Youssef J, Tyler N, Castle J, Jacobs PG. A statistical virtual patient population for the

glucoregulatory system in type 1 diabetes with integrated exercise model. PLoS One.

2019;14(7):e0217301. PMID: 31344037; PMCID: PMC6657828 following competing interests:

JRC and PGJ have a financial interest in Pacific Diabetes Technologies Inc., a company that may

have a commercial interest in the results of this research and technology.

24. Williams DJ, Shah SS, Myers A, Hall M, Auger K, Queen MA, Jerardi KE, McClain L, Wiggleton C,

Tieder JS. Identifying pediatric community-acquired pneumonia hospitalizations: Accuracy of

administrative billing codes. JAMA Pediatr. 2013;167(9):851-8. PMID: 23896966; PMCID:

PMC3907952.

25. O'Malley KJ, Cook KF, Price MD, Wildes KR, Hurdle JF, Ashton CM. Measuring diagnoses: ICD

code accuracy. Health Serv Res. 2005;40(5 Pt 2):1620-39. PMID: 16178999; PMCID:

PMC1361216.

26. Rubin DB. Basic Ideas of Multiple Imputation for Nonresponse. Survey Methodology.

1986;12(1):37-47.

27. Rubin DB, Schenker N. Multiple Imputation for Interval Estimation from Simple Random Samples

with Ignorable Nonresponse. J Am Stat Assoc. 1986;81(394):366-74. PMID:

WOS:A1986C648000012.

28. Ambler G, Omar RZ, Royston P. A comparison of imputation techniques for handling missing

predictor values in a risk model with a binary outcome. Stat Methods Med Res. 2007;16(3):277-

98. PMID: 17621472.

29. Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations: what is it

and how does it work? Int J Methods Psychiatr Res. 2011;20(1):40-9. PMID: 21499542; PMCID:

PMC3074241.

30. Kostkova P, Brewer H, de Lusignan S, Fottrell E, Goldacre B, Hart G, Koczan P, Knight P, Marsolier

C, McKendry RA, Ross E, Sasse A, Sullivan R, Chaytor S, Stevenson O, Velho R, Tooke J. Who

Owns the Data? Open Data for Healthcare. Front Public Health. 2016;4:7. PMID: 26925395;

PMCID: PMC4756607.

31. Ishikawa KB. Medical Big Data for Research Use: Current Status and Related Issues. Japan Med

Assoc J. 2016;59(2-3):110-24. PMID: 28299245; PMCID: PMC5333614.

32.   Hauser RG, Quine DB, Ryder A. LabRS: A Rosetta stone for retrospective standardization of

clinical laboratory test results. J Am Med Inform Assoc. 2018;25(2):121-6. PMID: 28505339.

33.   The Equifax Data Breach: What to Do [3/29/2018]. Available from:

https://www.consumer.ftc.gov/blog/2017/09/equifax-data-breach-what-do.

34.   Patient Data Landed Online After a Series of Missteps 2011 [3/30/2018]. Available from:

https://www.nytimes.com/2011/10/06/us/stanford-hospital-patient-data-breach-is-

detailed.html.

35.   Shelley MW. Frankenstein, or, The modern Prometheus : the 1818 text: Oxford ; New York :

Oxford University Press, 1998.; 1998.

Figure Legends

Figure 1. Cross-tabulated aggregate data from SPRINT. This represents the results from Steps 1-2 outlined in the methods and in Figure S1. The primary event rate and the prevalence of 9 independent factors are shown for subgroups defined by presence or absence of those same factors. The verbal equivalent of the top two rows, for instance, is: "Of the 3332 female participants of SPRINT, 100% were female, 30.01% were senior, 40.82% were black, [etc..], and 4.98% suffered the primary event. Of the 6029 non-female participants of SPRINT, 0% were female, 27.14% were senior, 26.32% were black, [etc..], and 6.57% suffered the primary event." The data shown here is the only starting material required to generate RE-ViPPs.

Figure 2. Patient counts in each RE-ViPP subgroup by demographic factor, clinical factor, trial factor, and primary outcome all adhere closely to SPRINT. One hundred replicate populations, each with 9361 virtual patients, were generated. The rates were, by design, within an absolute 0.25% of the actual SPRINT trial for every factor in every RE-ViPP. The full range of RE-ViPP counts are shown, as are p values from one-sample t-tests comparing the 100 RE-ViPPs to the actual SPRINT participants.

Figure 3. RE-ViPPs match the real SPRINT population. In each plot, a dot represents a single virtual patient population subgroup and horizontal bars are the real SPRINT rates. Red and gray respectively depict the subgroups with and without the corresponding x-axis factor. The prevalence of demographic factors [female sex (A), age≥75 (B), and black race (C)] and baseline clinical factors [prior/current smoking (D), highest tertile SBP (E), non-HDL>160 mg/dL (F), CKD defined by eGFR>60 (G), and past history of clinical CVD (H)] in each of the 100 virtual patient populations clusters around the actual rate in SPRINT across all subgroups. The virtual patient population is randomized well to intensive vs. standard BP therapy (I) across subgroups just as the real SPRINT patients, and primary event rates also match across all subgroups (J). Notable

features of the SPRINT population include higher female and lower senior rates amongst black

participants (A-C), lower rates of smoking and higher non-HDL amongst women (D,F), higher

CKD amongst seniors (G), and higher event rates in seniors, those with CKD, those with past

clinical CVD, and those randomized to non-intensive treatment (J). All these features are

preserved in all of the virtual patient populations.

Figure 4. Multiple logistic regression of RE-ViPPs. Multiple logistic regression performed on

the 100 virtual populations (9361 patients each) show that the Odds Ratio (OR) for each factor

approximate those of SPRINT. Each colored dot is the calculated OR for a virtual patient

population. The OR calculated from the mean beta coefficient for each factor along with the

central 95% of these ORs is shown in black. For comparison, the actual SPRINT OR and 95%

confidence interval for each factor is shown in red.

Figure 5. Multiple logistic regression of RE-ViPPs match SPRINT. Volcano plot shows the OR

plotted against the -log of the p value for each factor (color-coded) in each RE-ViPP and in

SPRINT. Lower panel is a magnified area from upper panel. The magnitude and significance of

the each factor's impact in SPRINT is recapitulated the by RE-ViPPS, as the SPRINT value falls in

the middle of the values for the RE-ViPPs for each respective factor.

Figure 1

		Count (%)	% Female	% Senior	% Black	% Smokers	% Hi-SBP	% Hi-Non-HDL	% CKD	% Clinical CVD	% Intensive	Event Rate (%)
Female	+	3332 (35.6)	100	30.01	40.82	44.87	39.05	31.90	31.75	11.43	50.54	4.98
	-	6029 (64.4)	0	27.14	26.32	61.67	31.28	20.42	26.34	19.59	49.66	6.57
Senior (≥75 years)	+	2636 (28.2)	37.94	100	17.75	52.85	39.23	14.95	44.04	21.85	49.96	9.29
	-	6725 (71.8)	34.68	0	36.86	56.80	32.01	28.25	22.08	14.66	49.98	4.71
Black / African-American	+	2947 (31.5)	46.15	15.88	100	55.55	34.98	27.28	21.85	10.52	49.34	4.99
	-	6414 (68.5)	30.75	33.80	0	55.75	33.61	23.23	31.21	19.52	50.27	6.47
Smoker (Current/Former)	+	5213 (55.7)	28.68	26.72	31.40	100	33.72	23.13	27.58	19.85	50.18	6.85
	-	4148 (44.3)	44.29	29.97	31.58	0	34.45	26.23	29.12	12.70	49.71	4.94
SBP in highest tertile	+	3187 (34.0)	40.82	32.44	32.35	55.16	100	27.11	27.55	15.25	50.39	6.59
	-	6174 (66.0)	32.90	25.95	31.03	55.96	0	23.16	28.64	17.43	49.76	5.70
Non-HDL > 160 mg/dL	+	2294 (24.5)	46.34	17.18	35.05	52.57	37.66	100	23.67	9.85	50.22	6.63
	-	7067 (75.5)	32.11	31.72	30.32	56.70	32.87	0	29.76	18.90	49.89	5.80
CKD (eGFR <60)	+	2646 (28.3)	39.98	43.88	24.34	54.35	33.18	20.52	100	21.01	50.26	8.84
	-	6715 (71.7)	33.86	21.97	34.30	56.22	34.39	26.08	0	14.98	49.86	4.88
Clinical CVD	+	1562 (16.7)	24.39	36.88	19.85	66.26	31.11	14.47	35.60	100	49.87	12.29
	-	7799 (83.3)	37.84	26.41	33.81	53.57	34.63	26.52	26.80	0	49.99	4.74
Intensive Arm	+	4678 (50.0)	36.00	28.15	31.08	55.92	34.33	24.63	28.43	16.65	100	5.19
	-	4683 (50.0)	35.19	28.17	31.88	55.46	33.76	24.39	28.10	16.72	0	6.81

Figure 2

		N: 100 RE-ViPPs				
	N: SPRINT	**Mean**	**SD**	**Minimum**	**Maximum**	**p**
Total	9361	9361	0.0	9361	9361	n/a
Subgroups						
Female	3332	3332.0	0.0	3332	3332	n/a
Senior (≥75 years)	2636	2636.0	0.0	2636	2636	n/a
Black / African-American	2947	2945.9	13.3	2924	2970	0.39
Smoker (Current/Former)	5213	5210.9	12.0	5190	5235	0.08
SBP in highest tertile	3187	3187.7	13.6	3164	3210	0.59
Non-HDL > 160 mg/dL	2294	2292.0	13.4	2271	2317	0.14
CKD (eGFR <60)	2646	2646.6	12.5	2624	2669	0.64
Clinical CVD	1562	1561.9	13.1	1539	1584	0.92
Intensive Arm	4678	4679.4	12.9	4656	4701	0.29
Primary Outcome	562	560.1	11.8	539	583	0.11

Demographic Factors (row group: Female, Senior, Black / African-American)

Clinical Factors (row group: Smoker, SBP in highest tertile, Non-HDL, CKD, Clinical CVD)

Trial Factor (row group: Intensive Arm)

Figure 3

Figure 4

Figure 5