Reverse Engineered Virtual Patient Populations as Surrogates for Real Patient-

Level Data

Francis J. Alenghat

Section of Cardiology, Department of Medicine, University of Chicago, Chicago, Illinois, USA

Email: alenghat@uchicago.edu (FJA)

Short Title: Reverse Engineered Virtual Patient Populations (RE-ViPPs)

Abstract

Dissemination of clinical data for research has limitations. The most coveted data is richly descriptive at the individual level, but acquiring such granularity comes with significant cost, effort, or time. De-identification of individual records is not foolproof, with potential for privacy breaches, especially for "real-world" data derived from electronic health records. Also, the open data movement has progressed slowly for clinical trials, partly due to concerns about data ownership. Here, reverse engineered virtual patient populations (RE-ViPPs) are described, based on aggregate cross-tabulated categorical data from populations. The method does not require end-user access to individual-level data. Rather, using sequential linear regressions and random number generation, it generates virtual individual patients to comprise populations that, on average, closely resemble the real population in question. The method is validated by applying it to aggregated data derived from the seminal SPRINT trial, for which the individual-level data is known. The method yields virtual populations, each with 9361 patients, faithfully mimicking the 9361 real SPRINT participants. Multiple logistic regression on 100 such populations shows that, just as in SPRINT, risk factors with the highest odds ratio for the primary event are, in descending order, past clinical cardiovascular disease, age ≥ 75, chronic kidney disease, high non-HDL, and smoking history. Factors associated with fewer events are female sex and intensive blood pressure treatment (the trial's intervention). Application of RE-ViPPs to trials, registries, and health record databases could reduce the cost, time, ownership, and de-identification burdens hindering open data by encouraging dissemination of aggregate, richly cross-tabulated real data that investigators can use to construct virtual patients and make meaningful conclusions.

Introduction

Patient-level data from clinical trials, registries, and the electronic health record (EHR) is the basis for most clinical research. Obtaining and sharing such data can be challenging, and concerns have been raised about cost, effort, patient privacy, and data ownership [1]. Even journals with strong policies to promote data sharing of published papers have suboptimal adherence by authors [2]. Paying the right amount of attention to who requests individual level data, the purpose of the request, the structure and format of the data, and its de-identification to an extent sufficient to eliminate any chance of deducing identity, can be administratively taxing on the source institution, and often those costs are passed on to the requesting investigator [3-5].

Several entities have taken the initiative to open their data for specific trials and registries, which is a positive step forward. These include a consortium of clinical study sponsors initiating clinicalstudydatarequest.com [6,7], the Yale University Open Data Access (YODA) project [8], Duke's Supporting Open Access Research (SOAR) initiative [9], ImmPort [10], and Project Data Sphere [11]. Some have associated costs in the thousands of dollars whereas others are free. Biolincc is the NHLBI's free source for trial data and has reasonable access [12,13]. Besides clinical trials, sources of data can also include institutional, local, national, and international registries as well as clinical data warehouses driven by EHR data from one or more institutions. Typical time from request to obtaining individual-level de-identified data from one's institutional EHR ranges from weeks to months and costs from hundreds to thousands of dollars [4,5].

However, there are sources of aggregated data, based on real clinical populations,

which are currently underutilized for their potential. Public systems such as Center for Disease

Control's WONDER [14], the Healthcare Cost and Utilization Project (HCUPnet) [15], and the

Behavioral Risk Factor Surveillance System Web Enabled Analysis Tool [16] are examples of this

sort of aggregate open data for vital statistics, registries, and survey results, respectively. As for

EHR data, the i2b2 and similar platforms are deployed by multiple hospital systems to query

data in an aggregate fashion at those institutions; they are mostly promoted as tools for cohort

identification for planning future requests for individual-level data, but they have potential to

be more widely used for research purposes [17-19]. Beyond informing requests for individual

level data, these sources of aggregate data, when provided in a richly cross-tabulated fashion,

could contain enough information to approximate real populations. If this can be

demonstrated, it would allow investigators using open data to analyze without infringing upon

patient privacy or overly taxing current systems in place, and at the same time, allow

investigators running trials or maintaining registries to continue to have ownership of their

individual-level data.

Here a method is described for reverse engineering populations of virtual patients from

cross-tabulated aggregate data. It is applied to the SPRINT trial, a study of 9361 patients

randomized to intensive versus standard blood pressure control, for which the individual-level

data summarized in the original publication [20] was recently 'opened' through NHLBI's Biolincc

and is thus known [21]. This allows for comparison of the reverse engineered virtual patient

populations (RE-ViPPs) generated from aggregate SPRINT data with the actual SPRINT

population.

Methods

Study data

This is a secondary analysis of SPRINT data, which is provided publicly, in de-identified fashion, by the National Heart, Lung, and Blood Institute (NHLBI) via the Biolincc data repository (https://biolincc.nhlbi.nih.gov/home/). The study was deemed exempt by the University of Chicago IRB and the data was obtained after a signed data use agreement.

In SPRINT, participants were > 50 years old with a systolic blood pressure (SBP) of 130 to 180 mm Hg and had an increased risk of cardiovascular events for at least one of the following four reasons: clinical or subclinical cardiovascular disease, chronic kidney disease (CKD), a 10-year risk of 15% or greater on the basis of the Framingham risk score, or ≥ 75 years old. Patients with diabetes or prior stroke were excluded. Participants were randomized to an SBP target of either less than 140 mm Hg (the standard-treatment group) or less than 120 mm Hg (the intensive-treatment group). The primary composite outcome was myocardial infarction, other acute coronary syndromes, stroke, heart failure, or death from cardiovascular causes, measured over an average of 3.3 years [20].

The SPRINT individual-level data were aggregated by first choosing 9 independent categorical factors. Of these, eight were categorical in the original data [sex, senior (age≥75), black race, current or former smoker (grouped together in the present study), highest SBP tertile at the start of the trial, CKD (eGFR < 60 ml/min/1.73m^2), history of clinical cardiovascular disease, and intensive SBP treatment]. The last (high non-HDL) was determined by calculating

the non-HDL for each patient and identifying those with non-HDL > 160 mg/dL. The dependent

variable was the primary composite outcome, also categorical.

Construction of a Reverse Engineered Virtual Patient Population (RE-ViPP)

The starting material for RE-ViPP construction consists of aggregate data cross-tabulated for

every pair of factors to be analyzed. The cross-tabulation is created by partitioning the entire

population into subgroups by the status of one independent factor and recording, for each

subgroup, the prevalence of each of the remaining factors, along with the rate of the

dependent variable (**Fig 1**, Steps 1-2). For SPRINT, this is straightforward, since the individual-

level data is known. For EHR, registry, and large trial databases, this can be done, without giving

investigators access to the individual level data, as long as the database can be queried for

multiple factors at a time.

To start a RE-ViPP, the exact number of patients with and without the first independent

variable is assigned to match the real population (**Fig 1**, Step 3). In the case of the SPRINT

analysis, this was female sex. Then, in those virtual patients with the first factor, the second

factor (in SPRINT, this was senior status) was assigned to the exact percentage measured in the

real population and the same was done for those virtual patients without the first factor. Thus

the status of the first and second chosen factors will match the real population exactly.

Starting with the third factor (**Fig 1**, Step 4), linear regression was performed on the

aggregate data using all the previously assigned factors to estimate the chances of a patient

having the new factor as a function of the previously assigned factors. The β coefficients and y-

intercept from this linear regression comprise a formula to calculate a probability of having the

new factor for each virtual patient (Supplemental Information, Appendix 1). Significant

interactions between pairs of factors are checked in the linear regression and included if the

interaction β is within a factor of 1000 from the other β coefficients (in SPRINT, no such

interactions were found between any pair of the 9 factors).

Once the probability (between 0 and 1) of having the new factor is calculated for each

virtual patient, a random number (between 0 and 1) is generated for each patient (**Fig 1**, Step

5). If the random number is less than the patient's probability of having the new factor, the

patient is assigned as having the new factor. Otherwise, the virtual patient does not. This

repeated for each patient. Once all patients are assigned, the process repeats for the next

factor to be assigned. After sequentially assigning the status for all the factors for all the virtual

patients in the above manner, the outcomes are assigned in the same manner.

Once all factors and outcomes have been assigned to all virtual patients, the overall rate

of each factor and outcome is measured against the actual population, and the virtual

population is only accepted if within a specified tolerance, for all factors, of the actual

population. If the rate of any factor is outside this tolerance, the random number generation is

repeated for all patients and factors in order to create a new virtual population until a

population is found that is within tolerance for all factors (**Fig 1**, Step 6). For this study of

SPRINT, the tolerance was set at ±0.25%. One hundred such acceptable populations are

generated in this manner.

Analysis of RE-ViPPs

Each RE-ViPP is checked for correlation between the independent variables. It is then subjected

to multiple logistic regression using the nine categorical independent variables and the primary

composite outcome as the dependent variable.

Random number generation, assignment of virtual patient factors, and tolerance testing

were conducted in Microsoft Excel with Macros. The linear and logistic regression analyses

were all conducted using R. Figures were created with ggplot2 in R.

Results

The process for creating reverse engineered virtual patient populations (RE-ViPPs) is presented

in the methods section and summarized in **Fig 1**. The process starts with curating cross-

tabulated aggregate data. For SPRINT, this was obtained by analyzing the individual-level data.

However, in its application for populations where individual-level data is not desired or easily

available, the creation of the cross-tabulated data would come from querying an EHR database

(such as via i2b2) or open aggregate/summary data from registries or clinical trials. **Fig 2** depicts

the cross-tabulated aggregate data from SPRINT.

In the case of RE-ViPPs based on SPRINT, the populations were created based on 9

independent factors and the combined primary outcome event. Each RE-ViPP is composed of

9361 patients, utilizes eight equations (Supporting Information, Appendix 1), derived from

linear regression analysis, for estimating the chance of a patient having a factor or outcome,

and requires a total of 74,888 random numbers. The tolerance was set at 0.25%, meaning that

the prevalence of each factor and the primary event rate all had to be within this absolute

percentage from the true rate in SPRINT. If this was not true for a given virtual population, all

9361 patients were regenerated until the criterion was met. The "SPRINT RE-ViPP Generator" is

available for download (Supporting Information, upon publication).

One hundred RE-ViPPs were created in this randomized fashion, constrained by the

chosen tolerance (each of these RE-ViPPs is available for download in Supporting Information,

upon publication). The rate of each included demographic factor (sex, age≥75, black race),

clinical factor (prior/current smoking, highest tertile SBP, non-HDL>160 mg/dL, CKD defined by

eGFR>60, past history of clinical CVD), trial factor (intensive vs standard BP therapy), and the

primary outcome (composite myocardial infarction, other acute coronary syndromes, stroke,

heart failure, or cardiovascular death), as required by the design, all adhere very closely to the

SPRINT rates (**Fig 3**).

Also across all subgroups, the prevalence of each demographic and clinical factor in each

of the 100 virtual patient populations clusters around the actual rates in SPRINT (**Fig 4**). The

virtual patient population is randomized well to intensive and standard BP therapy, just as the

real SPRINT patients, and primary event rates also match across all subgroups (**Fig 4**). It is worth

noting that the SPRINT population has higher female and lower senior rates amongst black

participants, lower rates of smoking and higher non-HDL amongst women, more frequent CKD

amongst seniors, and higher event rates in seniors, those with CKD, those with past clinical

CVD, and those randomized to non-intensive treatment. All these features are preserved in all

of the virtual patient populations.

Beyond subgroup characteristics, however, the RE-ViPPs can be analyzed on the

individual level similar to a multivariate analysis performed on real populations. Correlation

matrices show no strong correlations between the analyzed factors in the RE-ViPPs, just as in

the real SPRINT population (**Fig 5**)**.**

Multiple logistic regression performed on each of the 100 virtual populations show that

the odds ratios (ORs) for each factor approximates the ORs in SPRINT (**Fig 6**). The confidence

intervals of the RE-ViPP ORs are also similar to SPRINT. A volcano plot shows the OR plotted

against p values for each factor in each RE-ViPP, again showing close adherence to SPRINT (**Fig

7). The risk factors with the highest odds ratio for the primary event are, in descending order,

past clinical cardiovascular disease, age ≥ 75, chronic kidney disease, high non-HDL, and

smoking history. Factors associated with fewer events are female sex and intensive blood

pressure treatment. Just as with the true SPRINT data, neither black race nor starting in the

highest tertile of SBP had confidence intervals that clear the line of neutrality.

Discussion

The approach described here was chosen to demonstrate that meaningful information can be

gleaned from certain aggregated data to simulate individual patients. The method extends

beyond what is traditionally done with summary information and it can approximate the results

from the true individual-level data. The method cannot be done with any form of aggregate

data. The data must be cross-tabulated by the studied factors and outcomes, and, as such, all

the factors should be categorical, either nominal or ordinal. In the current example with

SPRINT, the variables were binary, but this process could be extended to incorporate a limited

number of non-binary categorical variables, in which case the cross-tabulation should account

for all values of those variables. Thus, there are important limitations to this method. Another

important limit is the quality of the source data. In a clinical trial such as SPRINT, it is likely very

reliable, but in EHR databases, variables such as race and smoking status can be variably and

inconsistently reported, and diagnoses and outcomes may rely on ICD codes which primarily

serve a billing/documentation function and may not be subject to rigorous independent

verification [22,23]. Another limitation is that although the method checks and includes any

pairwise interaction between variables that is large and significant, it does not address 3-way or

more complex interactions. There were no important interactions between the variables

chosen for this SPRINT study, and for other studies this limitation could be addressed by

checking for such complex interactions during the linear regression analyses and/or by

choosing, when possible, sufficiently distinct variables to analyze.

The method described here is reminiscent of imputation, the process by which missing

data for a variable is replaced by a reasonable prediction, often using regression models, based

on the extant values for that variable as a function of the other variables in the dataset [24,25].

This has perhaps taken its most robust form in multiple imputation of chained equations

[26,27]. The method here is distinct from multiple imputation in several ways, but an analogy

would be that every patient's status for all the variables beyond the first two are treated as

missing, and they are imputed sequentially based on logistic regression analysis of the

percentage distributions in the aggregate data.

The current study is meant to serve as a practical proof-of-principle example. Data

known at the individual level was used to highlight the method and show that the RE-ViPPs

result matches that of the real population. In practice, this method would be most useful in

situations when the individual-level data is either impossible or difficult to obtain. These are

common situations. In many cases, there are concerns about data ownership or patient privacy

[28]. In others, it can also involve excess time, effort, and cost [3,29]. Still in others, the issue

may be one of data standardization, where a study would ideally be done across datasets or

platforms, but the data exist in incongruent formats in each platform [30]. Also, the sheer bulk

of individual-level data may be prohibitive, particularly when there is a desire to combine data

from multiple EHRs across the world. With these applications in mind, the "SPRINT RE-ViPP

Generator" is provided in the Supporting Information (upon publication) and can be customized

to other sets of aggregate data and variables.

There are many recent high profile examples, not only in healthcare, of bulk individual-

level data being hacked or leaked with major fallout [31,32]. In all these situations, keeping the

individual level data undisclosed while providing the requisite summary statistics could allow

other investigators to study the data while abrogating all these concerns. One way this could be

done is to keep the individual level data behind a firewall but allow conditional queries to

extract summary data – this is what the i2b2 platform and others do for EHR data warehouses,

what Center for Disease Control's WONDER provides for vital statistics and other limited clinical

information, and what the Behavioral Risk Factor Surveillance System Web Enabled Analysis

Tool does for its large national health survey results [14,16,17,19]. Another approach would be

to have the "data owners" provide the richly cross-tabulated results across as many categorical

variables as possible as a small static file (like **Fig 2**) and make it open to interested

investigators; this approach would probably be more feasible and require less digital

infrastructure for individual clinical trial results in which full data releases are not planned. This

could also allay concerns about the provenance of the source data in secondary analysis, as the

shared information would be more succinct and verifiable. In all the above scenarios, it is worth

emphasizing that none of the analyzed individual patients derived from this method are real.

Patients in the real population are well represented by the virtual patients, but as digital

versions of creation, the virtual patients are nameless composites only [33], with no

opportunity to deduce identity.

This method should allow us to analyze trial, registry, and routinely collected clinical

information over large populations to find connections between heterogeneous clinical and

demographic factors in a novel way that does not directly access patient-level data. The

method could reduce barriers that currently impede access and use of open data. It would

thereby stimulate more meaningful use of extant information, ultimately to generate new

hypotheses for further investigation and to identify previously unrecognized clinical

relationships.

Acknowledgements

Dr. Alenghat had full access to the aggregate data and takes responsibility for the accuracy of the data analysis. Data were provided by the National Heart, Lung, and Blood Institute via Biolincc.

References

References

1.   Rockhold F, Nisen P, Freeman A. Data Sharing at a Crossroads. N Engl J Med. 2016;375(12):1115-

7. PMID: 27653563.

2.   Naudet F, Sakarovitch C, Janiaud P, Cristea I, Fanelli D, Moher D, Ioannidis JPA. Data sharing and

reanalysis of randomized controlled trials in leading biomedical journals with a full data sharing

policy: survey of studies published in The BMJ and PLOS Medicine. BMJ. 2018;360:k400. PMID:

29440066; PMCID: PMC5809812.

3.   Sebaa A, Chikh F, Nouicer A, Tari A. Medical Big Data Warehouse: Architecture and System

Design, a Case Study: Improving Healthcare Resources Distribution. J Med Syst. 2018;42(4):59.

PMID: 29460090.

4.   University of Chicago Clinical Research Data Warehouse (CRDW) [accessed 3/28/2018]. Available

from: http://cri.uchicago.edu/crdw/.

5.   UCSF Request Clinical Data for Research [accessed 3/28/2018]. Available from:

https://data.ucsf.edu/data-assets/clinical-research.

6.   Vaduganathan M, Nagarur A, Qamar A, Patel RB, Navar AM, Peterson ED, Bhatt DL, Fonarow GC,

Yancy CW, Butler J. Availability and Use of Shared Data From Cardiometabolic Clinical Trials.

Circulation. 2018;137(9):938-47. PMID: 29133600; PMCID: PMC5828960.

7.   Nisen P, Rockhold F. Access to patient-level data from GlaxoSmithKline clinical trials. N Engl J

Med. 2013;369(5):475-8. PMID: 23902490.

8.   Krumholz HM, Waldstreicher J. The Yale Open Data Access (YODA) Project--A Mechanism for

Data Sharing. N Engl J Med. 2016;375(5):403-5. PMID: 27518657.

9.   Pencina MJ, Louzao DM, McCourt BJ, Adams MR, Tayyabkhan RH, Ronco P, Peterson ED.

Supporting open access to clinical trial data for researchers: The Duke Clinical Research

Institute-Bristol-Myers Squibb Supporting Open Access to Researchers Initiative. Am Heart J.

2016;172:64-9. PMID: 26856217.

10.   Bhattacharya S, Andorf S, Gomes L, Dunn P, Schaefer H, Pontius J, Berger P, Desborough V,

Smith T, Campbell J, Thomson E, Monteiro R, Guimaraes P, Walters B, Wiser J, Butte AJ.

ImmPort: disseminating data to the public for the future of immunology. Immunol Res.

2014;58(2-3):234-9. PMID: 24791905.

11.   Green AK, Reeder-Hayes KE, Corty RW, Basch E, Milowsky MI, Dusetzina SB, Bennett AV, Wood

WA. The project data sphere initiative: accelerating cancer research by sharing data. Oncologist.

2015;20(5):464-e20. PMID: 25876994; PMCID: PMC4425388.

12.   Ross JS, Ritchie JD, Finn E, Desai NR, Lehman RL, Krumholz HM, Gross CP. Data sharing through

an NIH central database repository: a cross-sectional survey of BioLINCC users. BMJ Open.

2016;6(9):e012769. PMID: 27670522; PMCID: PMC5051517.

13.   Coady SA, Wagner E. Sharing individual level data from observational studies and clinical trials: a

perspective from NHLBI. Trials. 2013;14:201. PMID: 23837497; PMCID: PMC3750470.

14.   Friede A, Rosen DH, Reid JA. CDC WONDER: a cooperative processing architecture for public

health. J Am Med Inform Assoc. 1994;1(4):303-12. PMID: 7719813; PMCID: PMC116209.

15.   Weiss AJ, Elixhauser A, Steiner C. Readmissions to U.S. Hospitals by Procedure, 2010: Statistical

Brief #154. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville (MD)2006.

16.   Fisher MA, Ma ZQ. Multiple chronic conditions: diabetes associated with comorbidity and

shared risk factors using CDC WEAT and SAS analytic tools. J Prim Care Community Health.

2014;5(2):112-21. PMID: 24327591.

17.   Klann JG, Buck MD, Brown J, Hadley M, Elmore R, Weber GM, Murphy SN. Query Health:

standards-based, cross-platform population health surveillance. J Am Med Inform Assoc.

2014;21(4):650-6. PMID: 24699371; PMCID: 4078284.

18.   Murphy SN, Gainer V, Mendis M, Churchill S, Kohane I. Strategies for maintaining patient privacy

311   in i2b2. J Am Med Inform Assoc. 2011;18 Suppl 1:i103-8. PMID: 21984588; PMCID:

312   PMC3241166.

313   19.   Murphy SN, Weber G, Mendis M, Gainer V, Chueh HC, Churchill S, Kohane I. Serving the

314   enterprise and beyond with informatics for integrating biology and the bedside (i2b2). J Am Med

315   Inform Assoc. 2010;17(2):124-30. PMID: 20190053; PMCID: 3000779.

316   20.   Group SR, Wright JT, Jr., Williamson JD, Whelton PK, Snyder JK, Sink KM, Rocco MV, Reboussin

317   DM, Rahman M, Oparil S, Lewis CE, Kimmel PL, Johnson KC, Goff DC, Jr., Fine LJ, Cutler JA,

318   Cushman WC, Cheung AK, Ambrosius WT. A Randomized Trial of Intensive versus Standard

319   Blood-Pressure Control. N Engl J Med. 2015;373(22):2103-16. PMID: 26551272; PMCID:

320   PMC4689591.

321   21.   Burns NS, Miller PW. Learning What We Didn't Know - The SPRINT Data Analysis Challenge. N

322   Engl J Med. 2017;376(23):2205-7. PMID: 28445656.

323   22.   Williams DJ, Shah SS, Myers A, Hall M, Auger K, Queen MA, Jerardi KE, McClain L, Wiggleton C,

324   Tieder JS. Identifying pediatric community-acquired pneumonia hospitalizations: Accuracy of

325   administrative billing codes. JAMA Pediatr. 2013;167(9):851-8. PMID: 23896966; PMCID:

326   PMC3907952.

327   23.   O'Malley KJ, Cook KF, Price MD, Wildes KR, Hurdle JF, Ashton CM. Measuring diagnoses: ICD

328   code accuracy. Health Serv Res. 2005;40(5 Pt 2):1620-39. PMID: 16178999; PMCID:

329   PMC1361216.

330   24.   Rubin DB. Basic Ideas of Multiple Imputation for Nonresponse. Survey Methodology.

331   1986;12(1):37-47.

25. Rubin DB, Schenker N. Multiple Imputation for Interval Estimation from Simple Random Samples

with Ignorable Nonresponse. J Am Stat Assoc. 1986;81(394):366-74. PMID:

WOS:A1986C648000012.

26. Ambler G, Omar RZ, Royston P. A comparison of imputation techniques for handling missing

predictor values in a risk model with a binary outcome. Stat Methods Med Res. 2007;16(3):277-

98. PMID: 17621472.

27. Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations: what is it

and how does it work? Int J Methods Psychiatr Res. 2011;20(1):40-9. PMID: 21499542; PMCID:

PMC3074241.

28. Kostkova P, Brewer H, de Lusignan S, Fottrell E, Goldacre B, Hart G, Koczan P, Knight P, Marsolier

C, McKendry RA, Ross E, Sasse A, Sullivan R, Chaytor S, Stevenson O, Velho R, Tooke J. Who

Owns the Data? Open Data for Healthcare. Front Public Health. 2016;4:7. PMID: 26925395;

PMCID: PMC4756607.

29. Ishikawa KB. Medical Big Data for Research Use: Current Status and Related Issues. Japan Med

Assoc J. 2016;59(2-3):110-24. PMID: 28299245; PMCID: PMC5333614.

30. Hauser RG, Quine DB, Ryder A. LabRS: A Rosetta stone for retrospective standardization of

clinical laboratory test results. J Am Med Inform Assoc. 2018;25(2):121-6. PMID: 28505339.

31. The Equifax Data Breach: What to Do [accessed 3/29/2018]. Available from:

https://www.consumer.ftc.gov/blog/2017/09/equifax-data-breach-what-do.

32. Patient Data Landed Online After a Series of Missteps 2011 [accessed 3/30/2018]. Available

from: https://www.nytimes.com/2011/10/06/us/stanford-hospital-patient-data-breach-is-

detailed.html.

33. Shelley MW. Frankenstein, or, The modern Prometheus : the 1818 text: Oxford ; New York :

Oxford University Press, 1998.; 1998.

① **Partition Entire Real Population of N Individuals by Factor X_n**

② Measure prevalence (P) of all factors as well as outcome rate (Z) for those **with** and **without** X_n

Repeat **①** and **②** for each factor $X_1..X_n$

③ Start a Virtual Patient Population of N individuals. Assign the exact number of patients with and without Factor X_1 to match the real population. Of those with Factor X_1, assign the exact P_2% to have Factor X_2 and the rest lack Factor X_2. Of those without Factor X_1, assign the exact P_2% to have Factor X_2 and the rest lack Factor X_2. For Factors X_1 & X_2, the virtual population will match the real population exactly.

④ Starting with Factor X_3 and proceeding sequentially through Factor X_n, perform linear regression using $P_1..P_{n-1}$ & $P_1..P_{n-1}$ for each Factor $X_1..X_{n-1}$ as independent variables and P_n & P_n for $X_1..X_{n-1}$ as the dependent variable to determine coefficients $\beta_1...\beta_{n-1}$ and intercept Y. Check and include significant interactions between pairs of factors. Use $\beta_1...\beta_{n-1}$ and Y and already assigned X_1-X_{n-1} in the virtual population to determine the probability, PROB, of having X_n for each virtual patient. Repeat for the outcomes Z and Z using $P_1..P_n$ and $P_1..P_n$ for all Factors as the independent variables.

⑤ Generate a random number, R, between 0 and 1 for each virtual patient. If R<PROB, then assign the patient to have X_n. Otherwise, assign the patient to lack X_n. Do the same for assigning the outcome to each patient.

⑥ Repeat **⑤** by regenerating the N•(n-1) random numbers until the prevalence of each Factor X_1-X_n and the outcome rate are all within an tolerance T of the entire real population. Once this occurs, generation of a single virtual patient population generation is complete. Repeat until the desired number of virtual populations are obtained.

Fig 1. Reverse Engineered Virtual Patient Populations (RE-ViPPs): A step-by-step method. The source material can be created by querying a database of accessible clinical data (Steps 1-2) or be provided by the data owners. Generation of RE-ViPPs (Steps 3-6) relies on assigning variable values to each virtual patient in a sequential manner based on linear regression models, driven by random number generation, and subjected to tolerance limits.

		Count (%)	% Female	% Senior	% Black	% Smokers	% Hi-SBP	% Hi-Non-HDL	% CKD	% Clinical CVD	% Intensive	Event Rate (%)
Female	+	3332 (35.6)	100	30.01	40.82	44.87	39.05	31.90	31.75	11.43	50.54	4.98
	−	6029 (64.4)	0	27.14	26.32	61.67	31.28	20.42	26.34	19.59	49.66	6.57
Senior (≥75 years)	+	2636 (28.2)	37.94	100	17.75	52.85	39.23	14.95	44.04	21.85	49.96	9.29
	−	6725 (71.8)	34.68	0	36.86	56.80	32.01	28.25	22.08	14.66	49.98	4.71
Black / African-American	+	2947 (31.5)	46.15	15.88	100	55.55	34.98	27.28	21.85	10.52	49.34	4.99
	−	6414 (68.5)	30.75	33.80	0	55.75	33.61	23.23	31.21	19.52	50.27	6.47
Smoker (Current/Former)	+	5213 (55.7)	28.68	26.72	31.40	100	33.72	23.13	27.58	19.85	50.18	6.85
	−	4148 (44.3)	44.29	29.97	31.58	0	34.45	26.23	29.12	12.70	49.71	4.94
SBP in highest tertile	+	3187 (34.0)	40.82	32.44	32.35	55.16	100	27.11	27.55	15.25	50.39	6.59
	−	6174 (66.0)	32.90	25.95	31.03	55.96	0	23.16	28.64	17.43	49.76	5.70
Non-HDL > 160 mg/dL	+	2294 (24.5)	46.34	17.18	35.05	52.57	37.66	100	23.67	9.85	50.22	6.63
	−	7067 (75.5)	32.11	31.72	30.32	56.70	32.87	0	29.76	18.90	49.89	5.80
CKD (eGFR <60)	+	2646 (28.3)	39.98	43.88	24.34	54.35	33.18	20.52	100	21.01	50.26	8.84
	−	6715 (71.7)	33.86	21.97	34.30	56.22	34.39	26.08	0	14.98	49.86	4.88
Clinical CVD	+	1562 (16.7)	24.39	36.88	19.85	66.26	31.11	14.47	35.60	100	49.87	12.29
	−	7799 (83.3)	37.84	26.41	33.81	53.57	34.63	26.52	26.80	0	49.99	4.74
Intensive Arm	+	4678 (50.0)	36.00	28.15	31.08	55.92	34.33	24.63	28.43	16.65	100	5.19
	−	4683 (50.0)	35.19	28.17	31.88	55.46	33.76	24.39	28.10	16.72	0	6.81

Fig 2. Cross-tabulated aggregate data from SPRINT. The data here represent the results from Fig 1, Steps 1-2. The primary event rate and the prevalence of nine independent factors are shown for subgroups defined by presence or absence of those same factors. This is the starting material necessary to generate RE-ViPPs.

| | | N: SPRINT | N: 100 RE-ViPPs | | | | p |
			Mean	SD	Minimum	Maximum	
	Total	9361	9361	0.0	9361	9361	n/a
	Subgroups						
Demographic Factors	**Female**	3332	3332.0	0.0	3332	3332	n/a
	Senior (≥75 years)	2636	2636.0	0.0	2636	2636	n/a
	Black / African-American	2947	2945.9	13.3	2924	2970	0.39
Clinical Factors	**Smoker (Current/Former)**	5213	5210.9	12.0	5190	5235	0.08
	SBP in highest tertile	3187	3187.7	13.6	3164	3210	0.59
	Non-HDL > 160 mg/dL	2294	2292.0	13.4	2271	2317	0.14
	CKD (eGFR <60)	2646	2646.6	12.5	2624	2669	0.64
	Clinical CVD	1562	1561.9	13.1	1539	1584	0.92
Trial Factor	**Intensive Arm**	4678	4679.4	12.9	4656	4701	0.29
	Primary Outcome	562	560.1	11.8	539	583	0.11

Fig 3. Patient counts in each RE-ViPP subgroup by demographic factor, clinical factor, trial factor, and primary outcome all adhere closely to SPRINT. One hundred populations, each with 9361 virtual patients, were generated. The rates were, by design, within an absolute 0.25% of the actual SPRINT trial for every factor in every RE-ViPP. The full range of RE-ViPP counts are shown, as are p values from one-sample t-tests comparing the 100 RE-ViPPs to SPRINT.

Fig 4. Reverse engineered virtual patient populations match the real SPRINT population. In each plot, a dot represents a single virtual patient population subgroup and horizontal bars are the real SPRINT rates. Red and gray respectively depict the subgroups with and without the corresponding x-axis factor. The prevalence of demographic factors [female sex (**A**), age≥75 (**B**), and black race (**C**)] and baseline clinical factors [prior/current smoking (**D**), highest tertile SBP (**E**), non-HDL>160 mg/dL (**F**), CKD defined by eGFR>60 (**G**), and past history of clinical CVD (**H**)] in each of the 100 virtual patient populations clusters around the actual rate in SPRINT across all subgroups. The virtual patient population is randomized well to intensive vs. standard BP therapy (**I**) across

subgroups just as the real SPRINT patients, and primary event rates also match across all subgroups (**J**). Notable features of the SPRINT population include higher female and lower senior rates amongst black participants (A-C), lower rates of smoking and higher non-HDL amongst women (D,F), higher CKD amongst seniors (G), and higher event rates in seniors, those with CKD, those with past clinical CVD, and those randomized to non-intensive treatment (J). All these features are preserved in all of the virtual patient populations.

SPRINT

RE-ViPPs

Fig 5. Correlation coefficients of SPRINT and RE-ViPPs. Correlation coefficients are shown for each pair of factors in SPRINT (upper panel). For the RE-ViPPS (lower panel), the average correlations for each pair of factors is reflected by the color scale, and the full range of correlations is provided. The pattern is similar to SPRINT. No coefficient exceeds 0.3 between any pair of factors in any RE-ViPP.

Fig 6. Multiple logistic regression results of RE-ViPPs. Multiple logistic regression performed on the 100 virtual populations (9361 patients each) show that the Odds Ratio (OR) for each factor approximates those of SPRINT. Each colored dot is the calculated OR for a virtual patient population. The OR calculated from the mean β coefficient for each factor along with the central 95% of these ORs is shown in black. For comparison, the actual SPRINT OR and 95% confidence interval for each factor is shown in red.

Fig 7. Multiple logistic regression results of RE-ViPPs. Volcano plot shows the OR plotted against the -log of the p value for each factor (color-coded) in each RE-ViPP and in SPRINT. Lower panel is a magnified area from upper panel. For all factors, the SPRINT value falls in the middle of the values for the RE-ViPPs.