Discovery of a *Streptococcus pneumoniae* serotype 33F capsular polysaccharide locus

that lacks *wcjE* and contains a *wcyO* pseudogene

Sam Manna[a,#], Eileen M. Dunne[a,b], Belinda D. Ortika[a], Casey Pell[a], Mike Kama[c], Fiona

M. Russell[b,d], Tuya Mungun[e], E. Kim Mulholland[a,c,f], Jason Hinds[g,h], Catherine Satzke[a,b,i]

Pneumococcal Research, Murdoch Children's Research Institute, Royal Children's

Hospital, Parkville, VIC, Australia[a]; Department of Paediatrics, The University of

Melbourne, Parkville, VIC, Australia[b]; Ministry of Health and Medical Services, Suva,

Fiji[c]; Centre for International Child Health, Royal Children's Hospital, Melbourne,

Australia[d]; National Center for Communicable Diseases, Ministry of Health, Ulaanbaatar,

Mongolia[e]; Department of Infectious Disease Epidemiology, London School of Hygiene

& Tropical Medicine, London, United Kingdom[f]; Institute for Infection and Immunity,

St. George's, University of London, London, United Kingdom[g]; BUGS Bioscience,

London Bioscience Innovation Centre, London, United Kingdom[h]; Department of

Microbiology and Immunology at the Peter Doherty Institute for Infection and Immunity,

The University of Melbourne, Parkville, VIC, Australia[i]

#Address correspondence to Sam Manna, sam.manna@mcri.edu.au, Address: Murdoch

Children's Research Institute, Royal Children's Hospital, 50 Flemington Road, Parkville

VIC 3052 Australia

Telephone: +61 (3) 9936 6773; Fax: +61 (3) 8341 6212

Running title: A novel pneumococcal serotype 33F capsule locus

## **Abstract**

Objectives:

As part of large on-going vaccine impact studies in Fiji and Mongolia, we identified

25/2750 (0.9%) of nasopharyngeal swabs that were positive for *Streptococcus*

*pneumoniae* contained pneumococci with a divergent 33F capsular polysaccharide locus

(designated '33F-1'). Here, we investigated the 33F-1 capsular polysaccharide locus to

better understand the genetic variation and its impact on serotyping results.

Methods:

Whole genome sequencing was conducted on ten 33F-1 pneumococcal isolates. Initially,

sequence reads were used for molecular serotyping by PneumoCaT. Phenotypic typing of

33F-1 isolates was then performed using the Quellung reaction and latex agglutination.

Genome assemblies were used in phylogenetic analyses of each gene in the capsular

locus to investigate genetic divergence.

Results:

All ten pneumococcal isolates with the 33F-1 *cps* locus typed as 33F by Quellung and

latex agglutination. Unlike the reference 33F capsule locus sequence, DNA microarray

and PneumoCaT analyses found that 33F-1 pneumococci lack the *wcjE* gene, and instead

contain *wcyO* with a frameshift mutation. Phylogenetic analyses found the *cpsA, cpsB,*

*cpsC, cpsD, wchA, wciG* and *glf* genes in the 33F-1 *cps* locus had higher DNA sequence

similarity to homologues from other serotypes than to the 33F reference sequence.

Conclusions:

We have discovered a novel genetic variant of serotype 33F, which lacks *wcjE* and

contains a *wcyO* pseudogene. This has important implications for molecular approaches

measuring vaccine impact and disease surveillance, which continue to increase in

popularity, particularly in low- and middle-income countries, where pneumococcal

serotype diversity is poorly understood.

## 53   **Introduction**

*Streptococcus pneumoniae* (the pneumococcus) is a Gram-positive pathogenic bacterium

and a leading cause of community-acquired pneumonia in young children and the elderly

[1]. Pneumococci are classified by serotype, defined by an antigenically-distinct

polysaccharide capsule. Capsule biosynthesis is encoded by the capsular polysaccharide

(*cps*) locus within the pneumococcal genome. High levels of genetic diversity within this

locus has resulted in over 90 pneumococcal serotypes described to date.

The pneumococcal capsule is the target for currently licensed vaccines, which only

include a subset of serotypes. Although pneumococcal conjugate vaccines (PCVs) have

been successful in reducing carriage and disease caused by the targeted serotypes, a rise

in carriage and disease caused by serotypes not included in these vaccines is commonly

observed (serotype replacement) [2,3]. To precisely monitor vaccine impact and disease

surveillance, accurate tools for pneumococcal serotyping are required.

Molecular approaches to serotyping pneumococci rely on existing knowledge of *cps* loci.

Data on pneumococcal *cps* loci from low- and middle-income countries (LMICs) are

relatively limited, which can impact serotyping results. For example, we recently

described a novel genetic variant of pneumococcal serotype 11A in Fiji. Genetically, the

*cps* locus of these isolates is most closely related to the 11F *cps* locus, with only a few

minor nucleotide changes resulting in the production of 11A capsule [4].

Among the replacing serotypes post-PCV introduction, serotype 33F has become a

concern world-wide. Serotype 33F is commonly reported among the predominant

serotypes not included in PCVs causing invasive disease following vaccine introduction

[5–7]. The increased invasive disease caused by serotype 33F has warranted its inclusion

in two new vaccine formulations, which are in development by Merck [8]. In this study,

we describe a novel 33F *cps* locus identified in Fiji and Mongolia by investigating the

genetic basis of the variation in this locus and the potential impact this may have on

serotyping results.

## Materials and Methods

**Nasopharyngeal swab collection and screening for pneumococci**

Nasopharyngeal swabs were collected from either healthy participants (Fiji) or children

with pneumonia (Mongolia) in accordance with WHO recommendations [9]. Ethical

approval for the study in Fiji was granted from the Fiji National Research ethics review

committee, as well as The University of Melbourne Human research ethics committee.

Ethical approval for the study in Mongolia was granted from the ethics committee

associated with The Ministry of Health in Mongolia and the Royal Children's Hospital.

Written consent for those who participated in these studies was provided by

parents/guardians. Following collection, the swabs were placed in 1 ml skim milk,

tryptone, glucose, and glycerol media [10] and stored at -80°C. Swabs were stored at -

80°C until required. Samples were screened for the presence of pneumococci by

conducting real-time quantitative PCR (qPCR) on DNA extracted from 100 µl aliquots of

the swabs using the pneumococcal *lytA* gene as a target as previously described [11].

**Molecular typing by microarray**

Molecular typing of pneumococci was performed by DNA microarray. An aliquot of the

nasopharyngeal swab was inoculated onto Horse Blood Agar supplemented with

gentamicin (5 µg/ml), to select for pneumococci, and incubated overnight at 37°C with

5% $CO_2$. For plates with α-hemolytic growth, the bacterial growth was collected using 1

ml PBS, pelleted by centrifugation and stored at -30°C until required for DNA extraction.

DNA was extracted from thawed bacterial pellets using the QIAcube HT with the

QIAamp 96 DNA QIAcube HT Kit (Qiagen) with the inclusion of a pre-treatment lysis

step consisting of the addition of 180 µl lysis buffer (20 mM TrisHCl, 2 mM EDTA, 1%

Triton X-100, 2 mg/ml RNase A, 20 mg/ml lysozyme) to the bacterial pellet and

incubated at 37°C for 60 min. The rest of the extraction procedure was as per the kit

instructions. This DNA was then used for microarray as described previously [12]. In

brief, 200 ng of DNA was labelled with Cy3 or Cy5 using the Genomic DNA ULS

Labeling Kit (Agilent Technologies) and incubated at 85°C for 30 min. The labelled

pneumococcal DNA was incubated with Senti-SPv1.5 microarray slides (BUGS

Bioscience) overnight at 65°C rotating at 20 rpm. Microarray slides were washed,

scanned, and analyzed using Agilent microarray scanner and feature extraction software.

Serotype calls were analyzed by Senti-NET software.

**Bacterial isolates**

The *S. pneumoniae* isolates used in this study were purified from ten nasopharyngeal

swabs containing 33F-1 from Fiji and Mongolia on selective media as described above.

Isolates were confirmed as *S. pneumoniae* with microarray and whole genome

sequencing.

**Whole genome sequencing and molecular typing**

For whole genome sequencing, DNA was extracted from pure cultures using the Wizard

SV genomic DNA purification system (Promega) with some modifications. Briefly,

pneumococcal cultures were pre-treated with a lysis solution containing 5 mM EDTA, 3

mg/ml lysozyme and 37.5 µg/ml mutanolysin in TE buffer and incubated at 37°C for 2 h.

Proteinase K was added to a final concentration of 1 mg/ml and samples were incubated

at 55°C for 1 h. Following incubation, 200 µl of nuclear lysis buffer and 5 µl of RNase

(final concentration of 40 µg/ml) were added and samples were incubated at 80°C for 10

min. The rest of the extraction procedure was performed as per the manufacturer's

instructions. Eluted DNA was sequenced in 2 x 300 bp paired end reads on the MiSeq

platform. Using the Geneious 11.0.4 software package [13], sequence reads were

trimmed with BBDuk and *de novo* assembled using SPAdes. The capsule loci were

annotated within Geneious using a database consisting of capsule loci from the 90

serotypes described by Bentley et al. [14]. Sequence reads were also used for molecular

typing with PneumoCaT [15].

**Sequence analysis**

Pairwise alignments were performed in Geneious 11.04 using MUSCLE. Additional

alignments examining the *wcyO* gene for insertions/deletions were conducted using

Clustal Omega. Phylogenetic analyses were performed for each 33F-1 *cps* gene using

MEGA 7 [16]. For each gene, the phylogenetic analysis included a representative 33F-1

sequence as well as homologues from all other serotypes containing that gene as

described by Bentley et al. [14], where GenBank accession numbers are provided. DNA

sequences were aligned using MUSCLE and the alignments were used to generate

maximum likelihood trees based on the Tamura-Nei model. Phylogenetic relationships

were statistically analyzed by bootstrapping (1000 replicates). The 33F-1 *cps* loci have

been deposited in Genbank (accession no. MH256127, MH256128, MH256129,

MH256130, MH256131, MH256132, MH256133, MH256134, MH256135, MH256136).

**Quellung and latex agglutination serotyping**

Quellung serotyping was performed as described previously [17]. A saline suspension of

pneumococci was prepared from an overnight culture on non-selective agar. Using an

inoculation loop, 1 µl was placed on a microscope slide and mixed with 1 µl of antisera

from the Statens Serum Institut (SSI) (http://www.ssi.dk/ssidiagnostica). The sample was

then viewed under the microscope (x400 magnification). A positive reaction was defined

as an enlargement or 'swelling' of cells, with serotype call based on the reaction profile

with each typing sera. For latex agglutination, latex reagents were prepared with SSI

typing sera [18] and testing performed as previously described [19]. The bacterial

suspension and latex reagent (10 µl of each) were mixed on a glass slide. The slide was

then incubated on an orbital shaker for 2 min at ~140 rpm. A positive reaction was

defined by the presence of visible agglutination.

**Results**

As part of ongoing programs in the Asia-Pacific region measuring pneumococcal vaccine

impact, we have collected nasopharyngeal swabs from healthy participants in Fiji, and

children diagnosed with pneumonia in Mongolia. In these studies, we have used DNA

microarray to serotype pneumococci contained within these swabs. DNA microarray uses

15,000 oligonucleotides that are spotted onto glass slides and recognize each capsule

gene from the 90+ serotypes. Labelled pneumococcal DNA is allowed to hybridize to the

oligonucleotides so that pneumococcal serotype can be inferred. From 2750 swabs that

contained pneumococci 25 (0.9%) contained pneumococci that typed as '33F-like'

(hereby referred to as '33F-1'). Ten of these samples were selected and the pneumococci

from these swabs were isolated for further analysis (Table 1).

Microarray reported the *wciG, glf* and *wcjE* genes in the nasopharyngeal swabs

containing these isolates as 'absent/divergent'. The *wcyO* gene was also detected, which

has not been reported in the serotype 33F *cps* locus previously. To investigate the impact

of the divergent 33F-1 *cps* locus on other molecular approaches to serotyping, we

sequenced the genomes of all ten isolates and ran the sequence reads through the

PneumoCaT pipeline [15]. PneumoCaT uses *wcjE* to differentiate 33A from 33F, as this

gene contains a frameshift mutation in 33F, resulting in a lack of WcjE-mediated O-

acetylation of the 33F capsular polysaccharide [20]. Consistent with microarray,

PneumoCaT typed all isolates as 33F and was unable to detect *wcjE*. Phenotypic

serotyping methods (Quellung and latex agglutination) also typed these isolates as 33F.

Following investigation of the 33F-1 *cps* locus, it was evident that not only did all ten

isolates lack *wcjE,* the locus contained *wcyO* at this position. The *wcyO* gene encodes an

acetyltransferase and mediates the same modification as *wcjE* (6-O-acetylation of

galactose) [21]. The *wcyO* open reading frame from all 33F-1 isolates contained a

frameshift mutation. The *wcyO* gene in 33F-1 pneumococci from Fiji had a single T

insertion (position 313) whereas this gene in isolates from Mongolia contained a single A

deletion (position 246) (Fig. 1). These frameshift mutations were also confirmed by

Sanger sequencing and were not present in traditional *wcyO*-containing isolates

(serotypes 34 and 39) from Fiji (Supplementary Fig. S1).

In addition to the differences in *wcjE* and *wcyO,* microarray detected some divergence in

other genes in the 33F-1 *cps* locus compared to the reference 33F sequence. To gain a

better understanding of the relationships of the 33F-1 *cps* genes to homologues from

other serotypes we performed phylogenetic analyses for each gene. In support of the

pairwise alignments (Supplementary Table S1), the 33F-1 *wciB, wciD, wciE, wciF, wzy*

and *wzx* genes clustered with 37/33A/33F sequences (Fig. 2F-K). In contrast, 33F-1 *cpsA,*

*cpsB, cpsC, cpsD* (also known as *wzg, wzh, wze* and *wzd,* respectively) and *wchA*

clustered with serotype 33B sequences (Figures 2A-D), *wciG* with serotype 37 (Fig. 2L),

*glf* with serotypes 34 and 39 (Figure 2M) and *wcyO* with 33C, 34 and 39 (Fig. 2N). All

branches had strong statistical support (>85% bootstrap score from 1000 replicates for all

genes, except *cpsD* with a 67% bootstrap score for the 33F-1/33B branch).

## Discussion

Pneumococcus is a highly successful pathogen in part due to the high level of capsule

diversity, resulting in over 90 serotypes each with unique antigenic properties. Even

small genetic differences in the *cps* locus can have biologically relevant consequences,

impacting opsonophagocytosis efficiency and immune recognition. Serotypes 33F and

33A have the same *cps* locus, except that 33F has a *wcjE* gene containing a frameshift

mutation [22]. Using DNA microarray, we identified a high degree of genetic divergence

in the capsule DNA sequence of some serotypes in Fiji and Mongolia. Here, we

characterized a serotype 33F variant (33F-1) that has the same genes as the canonical 33F

and 33A *cps* loci, except it possesses *wcyO* instead of *wcjE*. Interestingly, in the 33F-1

variants *wcyO* is predicted to encode a truncated protein due to a frameshift mutation.

These frameshift mutations suggest a loss of 6-O-acetylation in 33F-1 capsular

polysaccharide as the truncated protein (110 amino acids and 91 amino acids long in the

Fiji and Mongolia isolates, respectively compared to the full length WcyO protein of 352

amino acids) would unlikely be functional.

The frameshift mutations in isolates from Fiji and Mongolia have both occurred within

homopolymeric regions (Fig. 1, Supplementary Fig. S1). Such regions are prone to

slipped-strand mispairing, whereby errors made during DNA replication can result in the

insertion or deletion of a nucleotide [23]. Therefore, we postulate that the frameshift

mutations in the 33F-1 *wcyO* genes are the result of slipped-strand mispairing events.

This the first report of the *wcyO* acetyltransferase gene in the 33F *cps* locus, and it is also

the first report of a naturally occurring frameshifted allele of *wcyO*. The fact that the

mutation type and location differ between isolates from Fiji and Mongolia demonstrates

this mutation event has occurred on at least two independent occasions. Whether the

mutation of *wcyO* is due to selective pressure to inactivate a disadvantageous gene or due

to a lack of selective advantage conferred by WcyO-mediated acetylation remains to be

investigated. Previously, mutations have been identified in other pneumococcal capsule

acetyltransferase genes including *wciG* [24] and *wcjE* [22,25,26]. Serotype 11E, which

lacks WcjE-mediated acetylation can evade opsonophagocytosis more efficiently

compared to 11A (which possess WcjE-mediated acetylation) [25]. Pneumococci

expressing 33F capsules, which lack WcjE-mediated acetylation, exhibit enhanced

survival during drying compared to serotype 33A (with intact WcjE-mediated

acetylation) [27]. Laboratory constructed *wciG* mutants in serogroup 33 isolates were

more susceptible to opsonophagocytosis, and displayed increased adherence and biofilm

formation, which has yet to be evaluated in wildtype strains [27]. It is plausible that

mutation of *wcyO* in the 33F-1 pneumococci may serve a similar purpose, however this

requires further investigation.

Within the 33F-1 *cps* locus we identified 7/15 genes that exhibit higher DNA sequence

similarity to homologues from other serotypes rather than 33F. Both *glf* and *wcyO* are

similar to sequences from serotypes 34 and 39 (and 33C for *wcyO*) (Fig. 2M and 2N) and

*cpsA, cpsB, cpsC, cpsD* and *wchA* similar to sequences from 33B (Fig. 2A-E).

Recombination of the pneumococcal capsule genes resulting in mosaic *cps* loci such as

that of 33F-1 have been reported previously [28,29]. Although it is difficult to infer the

direction of horizontal transfer of these genes, the mosaic nature of the 33F-1 *cps* locus

would suggest an ancestral 33F-1 *cps* locus was the recipient of these genes.

Interestingly, serogroup 33 related *cps* genes have been identified in two non-

pneumococcal streptococcal strains (*Streptococcus oralis* subsp. *tigurinus* Az_3a and

*Streptococcus mitis* SK269) [30]. The 33D-like *cps* locus in *S. mitis* SK269 exhibited low

similarity to the 33F-1 *cps* locus, including the absence of *wchA* and *wciC* and all other

*cps* gene exhibiting <76% DNA identity (Supplementary Table S1). The *S. oralis* subsp.

*tigurinus* Az_3a *cps* locus possessed the same genes as the 33F-1 locus with variable

DNA identity across the locus (<77% with the 33F-1 *cpsA, cpsB, cpsC, cpsD, wchA* and

*wciB* genes, >96% for *wciC, wciD, wciE, wciF* and *wzy* genes, and 85-90% for *wzx, wciG*

and *glf* genes, Supplementary Table S1). The higher DNA identity of 33F-1 *cps* genes

with homologues from other pneumococcal serotypes suggests the Az_3a *cps* locus may

have evolved independently of the 33F-1 locus. In contrast to 33F-1, the *wcyO* gene in

Az_3a has an intact open reading frame and is most similar to the *wcyO* gene from

pneumococcal serotype 21 (DNA identity score 86.8% with serotype 21 *wcyO* compared

to 74.5% with 33F-1 *wcyO*). The existence of a divergent 33F-1 *cps* locus with a

functional *wcyO* raises interesting questions around why this gene has been inactivated in

33F-1 pneumococci but remains intact in a non-pneumococcal streptococcal species.

This study describes a novel genetic basis for pneumococcal serotype 33F, which we

discovered from nasopharyngeal isolates from Fiji and Mongolia. Serotype 33F is a

replacing serotype in invasive disease following vaccine introduction [5–7]. The public

health importance of 33F is reflected in that it has been included in two upcoming

vaccine formulations (PCV15 and PCV24) [8]. In addition, there is increasing popularity

in molecular serotyping approaches and it is therefore important to identify genetic

variants, which have the potential to impact serotyping results. This is particularly

important for the implementation of these methods in LMICs, where there is limited

understanding of the pneumococcal *cps* loci. The data gained from this study will be used

to update genetic typing tools for more accurate typing of serotype 33F in LMICs.

**References**

[1]   O'Brien KL, Wolfson LJ, Watt JP, Henkle E, Deloria-Knoll M, McCall N, et al.

Burden of disease caused by *Streptococcus pneumoniae* in children younger than 5

years: global estimates. Lancet 2009;374:893–902. doi:10.1016/S0140-

6736(09)61204-6.

[2]   Mulholland K, Satzke C. Serotype replacement after pneumococcal vaccination.

Lancet (London, England) 2012;379:1387; author reply 1388-9.

doi:10.1016/S0140-6736(12)60588-1.

[3]   Weinberger DM, Malley R, Lipsitch M. Serotype replacement in disease after

pneumococcal vaccination. Lancet 2011;378:1962–73. doi:10.1016/S0140-

6736(10)62225-8.

[4]  Manna S, Ortika BD, Dunne EM, Holt KE, Kama M, Russell FM, et al. A novel

genetic variant of *Streptococcus pneumoniae* serotype 11A discovered in Fiji. Clin

Microbiol Infect 2018;24:428.e1-428.e7. doi:10.1016/j.cmi.2017.06.031.

[5]  Balsells E, Guillot L, Nair H, Kyaw MH. Serotype distribution of *Streptococcus*

*pneumoniae* causing invasive disease in children in the post-PCV era: A

systematic review and meta-analysis. PLoS One 2017;12:e0177113.

doi:10.1371/journal.pone.0177113.

[6]  Pilishvili T, Lexau C, Farley MM, Hadler J, Harrison LH, Bennett NM, et al.

Sustained Reductions in Invasive Pneumococcal Disease in the Era of Conjugate

Vaccine. J Infect Dis 2010;201:32–41. doi:10.1086/648593.

[7]  Hicks LA, Harrison LH, Flannery B, Hadler JL, Schaffner W, Craig AS, et al.

Incidence of Pneumococcal Disease Due to Non–Pneumococcal Conjugate

Vaccine (PCV7) Serotypes in the United States during the Era of Widespread

PCV7 Vaccination, 1998–2004. J Infect Dis 2007;196:1346–54.

doi:10.1086/521626.

[8]  McFetridge R, Meulen AS, Folkerth SD, Hoekstra JA, Dallas M, Hoover PA, et al.

Safety, tolerability, and immunogenicity of 15-valent pneumococcal conjugate

vaccine in healthy adults. Vaccine 2015;33:2793–9.

doi:10.1016/J.VACCINE.2015.04.025.

[9]  Satzke C, Turner P, Virolainen-Julkunen A, Adrian P V., Antonio M, Hare KM, et

al. Standard method for detecting upper respiratory carriage of *Streptococcus*

320   *pneumoniae*: Updated recommendations from the World Health Organization

321   Pneumococcal Carriage Working Group. Vaccine 2013;32:165–79.

322   doi:10.1016/j.vaccine.2013.08.062.

323 [10] O'Brien KL, Bronsdon MA, Dagan R, Yagupsky P, Janco J, Elliott J, et al.

324   Evaluation of a medium (STGG) for transport and optimal recovery of

325   *Streptococcus pneumoniae* from nasopharyngeal secretions collected during field

326   studies. J Clin Microbiol 2001;39:1021–4. doi:10.1128/JCM.39.3.1021-1024.2001.

327 [11] Carvalho M da GS, Tondella ML, McCaustland K, Weidlich L, McGee L, Mayer

328   LW, et al. Evaluation and improvement of real-time PCR assays targeting lytA,

329   ply, and psaA genes for detection of pneumococcal DNA. J Clin Microbiol

330   2007;45:2460–6. doi:10.1128/JCM.02498-06.

331 [12] Satzke C, Dunne EM, Porter BD, Klugman KP, Mulholland EK, Vidal JE, et al.

332   The PneuCarriage Project: A Multi-Centre Comparative Study to Identify the Best

333   Serotyping Methods for Examining Pneumococcal Carriage in Vaccine Evaluation

334   Studies. PLoS Med 2015;12. doi:10.1371/journal.pmed.1001903.

335 [13] Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al.

336   Geneious Basic: An integrated and extendable desktop software platform for the

337   organization and analysis of sequence data. Bioinformatics 2012;28:1647–9.

338   doi:10.1093/bioinformatics/bts199.

339 [14] Bentley SD, Aanensen DM, Mavroidi A, Saunders D, Rabbinowitsch E, Collins

340   M, et al. Genetic analysis of the capsular biosynthetic locus from all 90

341   pneumococcal serotypes. PLoS Genet 2006;2:0262–9.

342   doi:10.1371/journal.pgen.0020031.

[15]   Kapatai G, Sheppard CL, Al-Shahib A, Litt DJ, Underwood AP, Harrison TG, et

al. Whole genome sequencing of *Streptococcus pneumoniae*: development,

evaluation and verification of targets for serogroup and serotype prediction using

an automated pipeline. PeerJ 2016;4:e2477. doi:10.7717/peerj.2477.

[16]   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics

Analysis Version 7.0 for Bigger Datasets. Mol Biol Evol 2016;33:1870–4.

doi:10.1093/molbev/msw054.

[17]   Habib M, Porter BD, Satzke C. Capsular Serotyping of *Streptococcus pneumoniae*

Using the Quellung Reaction. J Vis Exp 2014. doi:10.3791/51208.

[18]   Ortika BD, Habib M, Dunne EM, Porter BD, Satzke C. Production of latex

agglutination reagents for pneumococcal serotyping. BMC Res Notes 2013;6:49.

doi:10.1186/1756-0500-6-49.

[19]   Porter BD, Ortika BD, Satzke C. Capsular Serotyping of *Streptococcus*

*pneumoniae* by Latex Agglutination. J Vis Exp 2014:51747. doi:10.3791/51747.

[20]   Geno KA, Gilbert GL, Song JY, Skovsted IC, Klugman KP, Jones C, et al.

Pneumococcal capsules and their types: Past, present, and future. Clin Microbiol

Rev 2015;28:871–99. doi:10.1128/CMR.00024-15.

[21]   Bush CA, Yang J, Yu B, Cisar JO. Chemical Structures of Streptococcus

pneumoniae Capsular Polysaccharide Type 39 (CPS39), CPS47F, and CPS34

Characterized by Nuclear Magnetic Resonance Spectroscopy and Their Relation to

CPS10A n.d. doi:10.1128/JB.01731-14.

[22]   Mavroidi A, Aanensen DM, Godoy D, Skovsted IC, Kaltoft MS, Reeves PR, et al.

Genetic relatedness of the *Streptococcus pneumoniae* capsular biosynthetic loci. J

Bacteriol 2007;189:7841–55. doi:10.1128/JB.00836-07.

[23]    Levinson G, Gutman GA. Slipped-strand mispairing: a major mechanism for DNA

sequence evolution. Mol Biol Evol 1987;4:203–21.

doi:10.1093/oxfordjournals.molbev.a040442.

[24]    Geno KA, Bush CA, Wang M, Jin C, Nahm MH, Yang J. WciG O-

Acetyltransferase Functionality Differentiates Pneumococcal Serotypes 35C and

42. J Clin Microbiol 2017;55:2775–84. doi:10.1128/JCM.00822-17.

[25]    Brady AM, Calix JJ, Yu J, Geno KA, Cutter GR, Nahm MH. Low invasiveness of

pneumococcal serotype 11A is linked to ficolin-2 recognition of O-acetylated

capsule epitopes and lectin complement pathway activation. J Infect Dis

2014;210:1155–65. doi:10.1093/infdis/jiu195.

[26]    Calix JJ, Brady AM, Du VY, Saad JS, Nahm MH. Spectrum of pneumococcal

serotype 11A variants results from incomplete loss of capsule O-acetylation. J Clin

Microbiol 2014;52:758–65. doi:10.1128/JCM.02695-13.

[27]    Spencer BL, Saad JS, Shenoy AT, Orihuela CJ, Nahm MH. Position of O-

acetylation within the capsular repeat unit impacts the biological properties of

pneumococcal serotypes 33A and 33F. Infect Immun 2017;85:e00132-17.

doi:10.1128/IAI.00132-17.

[28]    Salter SJ, Hinds J, Gould KA, Lambertsen L, Hanage W, Antonio M, et al.

Variation at the capsule locus, cps, of mistyped and non-typable*Streptococcus*

*pneumoniae* isolates. Microbiol (United Kingdom) 2012;158:1560–9.

doi:10.1099/mic.0.056580-0.

[29]    van Tonder AJ, Bray JE, Quirk SJ, Haraldsson G, Jolley KA, Maiden MCJ, et al.

389  Putatively novel serotypes and the potential for reduced vaccine effectiveness:

390  capsular locus diversity revealed among 5405 pneumococcal genomes. Microb

391  Genomics 2016;2:90. doi:10.1099/mgen.0.000090.

392 [30] Skov Sørensen UB, Yao K, Yang Y, Tettelin H, Kilian M. Capsular

393  Polysaccharide Expression in Commensal Streptococcus Species: Genetic and

394  Antigenic Similarities to *Streptococcus pneumoniae*. MBio 2016;7:e01844-16.

395  doi:10.1128/mBio.01844-16.

426 **Tables and figures**

427 Table 1. Pneumococcal 33F-1 isolates used in this study.

| Isolate | Source | Country of isolation |
|---|---|---|
| PMP1348 | Nasopharynx of healthy child (2-7 years old) | Fiji |
| PMP1349 | Nasopharynx of healthy child (5-8 weeks old) | Fiji |
| PMP1351 | Nasopharynx of healthy child (12-23 months old) | Fiji |
| PMP1352 | Nasopharynx of healthy child (12-23 months old) | Fiji |
| PMP1353 | Nasopharynx of healthy child (5-8 weeks old) | Fiji |
| PMP1379 | Nasopharynx of healthy child (12-23 months old) | Fiji |
| PMP1380 | Nasopharynx of healthy child (12-23 months old) | Fiji |
| PMP1383 | Nasopharynx of healthy adult | Fiji |
| PMP1386 | Nasopharynx of child with pneumonia | Mongolia |
| PMP1387 | Nasopharynx of child with pneumonia | Mongolia |

Figure 1. Comparison of the *wcyO* open reading frames of 33F-1 sequences to a

representative serotype 34 sequence. Only a selected portion of the DNA sequence is

shown. Numbers refer to the position number in the serotype 34 sequence with an in-

frame *wcyO* gene.

Figure 2. Maximum likelihood phylogenetic trees of 33F-1 *cps* genes with homologues

from all other serotypes. As all genes except *wchA* and *wcyO* were identical in all 33F-1

isolates only one sequence is included as a representative. Un-collapsed trees are

provided in Supplementary Figure S2. Tree for *wciC* is not shown as this gene is only

present in serotypes 33F, 33A and 37, which all have over 98% DNA sequence identity

to the 33F-1 sequence. DNA sequences were aligned using MUSCLE and trees were

constructed using the Tamura-Nei model in MEGA 7. Only bootstrap values above 50%

are shown.

```
wcyO(34)            CTA TAT AAG ATA CTT GGT AAA TTT ATG GAA ATT ATC TAT TAT TTT CAT   192
                     L   Y   K   I   L   G   K   F   M   E   I   I   Y   Y   F   H
wcyO(33F-1_Fiji)    CTA TAT GAG ATA CTT GGT AAA TTT ATG GAA ATT ATC TAT TAT TTT CAT
                     L   Y   E   I   L   G   K   F   M   E   I   I   Y   Y   F   H
wcyO(33F-1_Mongolia) CTA TAT GAG ATA CTT GGT AAA TTT ATG GAA ATT ATC TAT TAT TTT CAT
                     L   Y   E   I   L   G   K   F   M   E   I   I   Y   Y   F   H
                            150           160           170           180           190

wcyO(34)            ATG CCA TTA TTT ATG GCT ATA TCG GGT GTA TTT TTC TCT ATT CAA ATA   240
                     M   P   L   F   M   A   I   S   G   V   F   F   S   I   Q   I
wcyO(33F-1_Fiji)    ATA CCA TTA TTT ATG GCT ATA TCG GGT GTA TTT TTC TCT ATT CAA ATA
                     I   P   L   F   M   A   I   S   G   V   F   F   S   I   Q   I
wcyO(33F-1_Mongolia) ATA CCA TTA TTT ATG GCT ATA TCG GGT GTA TTT TTC TCT ATT CAA ATA
                     I   P   L   F   M   A   I   S   G   V   F   F   S   I   Q   I
                            200           210           220           230

wcyO(34)            AAA AAA GAT CGA TGG AAT AAG ATT GAG AAA TTA TTA ACT AGT AAG TTT   288
                     K   K   D   R   W   N   K   I   E   K   L   L   T   S   K   F
wcyO(33F-1_Fiji)    AAA AAA GAT CGA TGG AAT AAG ATT GAG AAA TTA TTA ACT AGT AAG TTT
                     K   K   D   R   W   N   K   I   E   K   L   L   T   S   K   F
wcyO(33F-1_Mongolia) AAA AAG ATC GAT GGA ATA AGA TTG AGA AAT TAT TAA CTA GTA AGT TTA
                     K   K   I   D   G   I   R   L   R   N   Y   *
                            250           260           270           280

wcyO(34)            AAA AGA TTA ATA TTG CCA TTT TTT GTT TTT ACT TTA TTA TAT AGT TTG   336
                     K   R   L   I   L   P   F   F   V   F   T   L   L   Y   S   L
wcyO(33F-1_Fiji)    AAA AGA TTA ATA TTG CCA TTT TTT GTT TTT TAC TTT ATT ATA TAG TTT
                     K   R   L   I   L   P   F   F   C   F   Y   F   I   I   *   F
wcyO(33F-1_Mongolia) AAA GAT TAA TAT TGC CAT TTT TTG TTT TTA CTT TAT TAT ATA GTT TGC
                            290           300           310           320           330

wcyO(34)            CCA TTA AAA TAT ATA TCA AAC TAC TAC AAT GGT GTT TCG TTT TGG AGA   384
                     P   L   K   Y   I   S   N   Y   Y   N   G   V   S   F   W   R
wcyO(33F-1_Fiji)    GCC ATT AAA ATA TAT ATC AAA CTA CTA CAA TGG TGT TTC ATT TTG GAG
wcyO(33F-1_Mongolia) CAT TAA AAT ATA TAT CAA ACT ACT ACA ATG GTG TTT CAT TTT GGA GAG
                            340           350           360           370           380
```