1 Map segmentation, automated model-building and their application to the Cryo-EM Model
2 Challenge
3
4

5 Thomas C. Terwilliger[1,2], Paul D. Adams[3,4], Pavel V. Afonine[3,5], Oleg V. Sobolev[3]
6
[1]*Los Alamos National Laboratory, Los Alamos NM 87545 USA*
[2]*New Mexico Consortium, Los Alamos NM 87544 USA*
[3]*Molecular Biophysics & Integrated Bioimaging Division, Lawrence Berkeley National*
Laboratory, Berkeley, CA 94720-8235, USA
[4]*Department of Bioengineering, University of California Berkeley, Berkeley, CA, USA*
[5]*Department of Physics and International Centre for Quantum and Molecular Structures,*
Shanghai University, Shanghai, 200444, People's Republic of China
14
15
16 E-mail: tterwilliger@newmexicoconsortium.org

17
18
19 Abstract
20

A recently-developed method for identifying a compact, contiguous region representing the
unique part of a density map was applied to 218 cryo-EM maps with resolutions of 4.5 Å or
better. The key elements of the segmentation procedure are (1) identification of all regions of
density above a threshold and (2) choice of a unique set of these regions, taking symmetry into
consideration, that maximize connectivity and compactness. This segmentation approach was
then combined with tools for automated map sharpening and model-building to generate
models for the 12 maps in the 2016 cryo-EM model challenge in a fully automated manner. The
resulting models have completeness from 24% to 82% and RMS distances from reference
interpretations of 0.6 Å to 2.1 Å.

30
31 Introduction
32
In the 2016 Cryo-EM Modeling Challenge (see
http://challenges.emdatabank.org/?q=model_challenge; accessed 2017-11-19), a total of 12
maps were supplied to contestants along with reconstruction symmetry and the sequences of
the molecules present. One of the goals of the Challenge was to fully interpret such a map
given only the map, the symmetry and the sequence information. There are a number of tools
being developed by several groups for automated interpretation of cryo-EM maps (DiMaio and
Chiu, 2016). These include methods for identification of secondary structure (Jiang et al., 2001;
Kong and Ma, 2003; Kong et al., 2004; Baker, Ju and Chiu, 2007), methods for combination of
structure-modeling tools such as Rosetta with cryo-EM model-building (Lindert et al., 2012;
Wang et al., 2015; Frenz et al., 2017), semi-automated tools for full map interpretation (Baker
et al., 2011), and automated tools based on chain-tracing (Chen et al., 2016; Collins and Si,
2017) and template-matching approaches (Zhou, Wang and Wang, 2017).

45
Prior to the 2016 cryo-EM Model Challenge, we had begun development of software for
automatic map sharpening (*phenix.auto_sharpen;* Terwilliger et al., 2018a) and interpretation
of density in cryo-EM maps (*phenix.map_to_model;* Terwilliger et al., 2018b*)* as part of the
Phenix software package (Adams et al, 2010). It was possible in principle to apply these tools
directly to the 2016 Challenge, interpreting an entire map and ignoring the symmetry of the
map. It seemed however that it would be more efficient to work with just the unique part of a
map. We reasoned that this could be done by identifying a unique part of map that contained a
complete molecule, interpreting that part of the map, and then expanding the result using the
symmetry in the map to represent the entire map. Powerful tools existed for map
segmentation (e.g., Volkmann, 2002; Baker, Chiu and Bajaj, 2006; Yu and Bajaj, 2008; Pintilie et
al., 2010), but we wanted to be able to integrate the segmentation and symmetry analysis with
automated model-building so that information from model-building could be used to make the
final choice of the regions of density representing a single molecular unit. We therefore
developed a new *Phenix* tool, *phenix.segment_and_split_map* (Terwilliger et al., 2018b) which
could be used for this purpose. Here we describe the application of
phenix.segment_and_split_map to a set of 218 cryo-EM maps selected to generally represent
the unique currently-available cryo-EM maps with resolution of 4.5 Å or better. We then
describe map segmentation, sharpening, and model-building (Terwilliger et al., 2018b) applied
to the 12 cryo-EM maps in the 2016 Cryo-EM Model Challenge.
65
66
67   Methods
68
69   Summary of map segmentation
70
The main goal of our segmentation procedure is to identify the density in a map that
corresponds to the unique part of that map. (Note that we use "density" to refer to map values.
They can be electron density, electric potential, or any other quantity that is being used to
describe the locations of atoms in the map).   A secondary goal is to choose this density in such
a way that it corresponds as closely as possible to the unique biological unit in the map. Our
overall approach to map segmentation is (1) to identify all regions of density above an
automatically-determined threshold, and (2) to choose a unique set of density regions that
maximizes connectivity and compactness, taking into account the symmetry that is present. By
default, the process is repeated with a new threshold after removing the density that has been
used in the first iteration. The density threshold for consideration of a region of density is
chosen to yield a specific volume fraction (typically 20%) of the region of the macromolecule
above the threshold.   The map is divided into regions of density above the threshold density,
where each region is composed of points above the threshold and that have at least one
neighbor above the threshold. A unique set of regions is chosen using the symmetry (if any)
supplied by the user and the criteria that the unique set should be as compact and connected
as feasible. The details of this segmentation procedure have recently been described
(Terwilliger et al., 2018b).
88

Symmetry present in a map

We identified symmetry relationships that were applied during map reconstruction using a
simplified version of approaches described by Zhang et al., (2012). In many cases the symmetry
applied during reconstruction is specified in the EM Data Bank (EMDB, Electron Microscopy
Data Bank; Lawson et al., 2016; as for example "I" for icosahedral reconstructions, "C6" for a 6-
fold symmetry axis). In others, the symmetry is specified in meta-data associated with the
deposited model in the Protein Data Bank (PDB; Bernstein et al., 1977; Berman et al., 2000). In
still others, the model deposited in the PDB contains symmetry-related copies which we
extracted with the *Phenix* tool *phenix.simple_ncs_from_pdb*. If present, we used symmetry
from the deposited models and their meta-data, and if not, we used the information from the
EMDB or literature specified in the deposition and the assumption that principal symmetry axes
(i.e., screw axes, rotational axes) are generally along the principal axes of the reconstruction to
find reconstruction symmetry in the density maps.

Map-model correlations

We calculated map-model correlations using the *Phenix* tool *phenix.map_model_cc*. This tool
identifies the region occupied by the model as all grid points in the target map within a
specified distance (typically 3 Å) of an atom in the model. Then it generates a model-based map
on the same grid and calculates the correlation of density values between the target map and
the model-based map inside the region occupied by the model.

Model-based maps were calculated in reciprocal space using elastic atomic scattering factors of
electrons for neutral atoms as described (Colliex et al., 2006, Afonine et al., 2018b). These
scattering factors are framed as the sum of Gaussian terms, represent electric potential, and
assume that all atoms are independent. These scattering factors do not include the effects of
charged residues and therefore they may be substantially incorrect for certain atoms, including
phosphates in RNA or DNA and side chains such as aspartate and glutamate. As improved
representations of electron scattering expressed as sums of Gaussian terms these become
available these can readily be incorporated in the Phenix framework.

Data used for map segmentation

We selected a group of 218 cryo-EM maps to test our segmentation algorithms. We started
with 492 maps we could extract from the EMDB in August of 2017 with simple *Phenix* tools and
that were reconstructed at resolutions of 4.5 Å or better. We excluded 91 maps where the
resolution in the EMDB and PDB differed by 0.2 Å or more or was not reported, 24 maps where
the map-model correlation was less than 0.3, and 16 maps for which the signal-to-noise in map
sharpening (Terwilliger et al., 2018a) was less than 3. We then removed map-model pairs that
were largely duplications by clustering based on sequence identity using a cutoff of 95%
identity and choosing the highest-resolution representative of each group. The sequence

identity of two structures was calculated after alignment of each chain in the first structure
with the closest-matching chain in the second structure. If either sequence was contained
within the other, the identity was considered to be 100%. Otherwise if the lengths of the
sequences differed by more than 5% , or the percentage of residues in all chains of the first
structure matching a corresponding residue in the second structure was less than 95%, the
sequences were considered to be different. Four of the maps in this set were associated with
two models, so one map-model pair was set aside for each of these, yielding 218 map-model
pairs that were analyzed in this work.
141
142   Evaluating the results of map segmentation by calculation of fraction of molecular unit within
143   the segmented region
144
We estimated the fraction of the molecular unit within the segmented region of a map from a
comparison of map-model correlations. Our method is related to the cross-correlation
variation metric described by Zhang et al. (2012) but it is extended to make an estimate of the
fraction of the molecular unit that matches the segmented map.   The segmented map has
values of zero everywhere outside the segmented region of the map. The overall idea is that if
the segmented region contains a complete molecular unit, then the map-model correlation
between one complete molecular unit and the segmented map will be the same as the map-
model correlation with the original map.   On the other hand, if the segmented region contains
part of one molecular unit and parts of symmetry-related ones, then the map-model
correlation between one intact molecular unit and the segmented map will be lower than the
correlation to the original map. We use this difference in map correlation to estimate the
fraction of a complete molecule that is within the segmented region.
157
We first calculated the map-model correlation between the original map and a map calculated
from single molecular unit extracted from the deposited model of the structure. Then we
calculated the map-model correlation between the segmented map and a single molecular unit.
The square of the ratio of these correlations is (see below) approximately equal to the fraction
of the molecular volume that is within the segmented map. The single molecular unit to
compare with the map was chosen to be a set of chains representing the unique part of the
deposited model. In cases with symmetry, each symmetry-related choice of molecular unit was
considered and the one with the highest map-model correlation was chosen.
166
The relationship between the map-model correlation for a single molecular unit and the
original map compared to the correlation for a molecular unit and a segmented map can be
calculated in a straightforward fashion with one assumption. This assumption is that the local
map-model correlation for the original map and this single molecular unit is approximately the
same everywhere in the region of the model.   With this assumption, we can readily calculate
the effect of setting all but a fraction f of the map density in the region of the model to zero.
This corresponds to calculating the map-model correlation of the segmented map to one full
molecular unit, where a fraction f of the molecular unit is present in the segmented map.
175

The correlation coefficient CC between two maps with density values represented by D_1 and D_2
can be written (after adjusting each map to set the mean density for each to zero so that $< D_1>$
$= < D_2> =0$) as,

$\quad\quad$ CC $= < D_1D_2>/\text{sqrt}(<D_1^2> <D_2^2>)$, $\quad\quad\quad\quad\quad$ (1)

where the calculation in this case is carried out over all the grid points near the model. Now
suppose we create a new map D_1' in which we set D_1 to zero at a fraction $(1-f)$ of these grid
points. Referencing Eq. (1), this means that the values of $D_1'D_2$ and $D_1'^2$ will be zero at all these
grid points, but D_2^2 will be the same. Assuming then that the values of $< D_1D_2>$, $<D_1^2>$, and
$<D_2^2>$ are approximately the same everywhere near the model, we can write that the map-
model correlation for the segmented map (CC') with all but $(1-f)$ of the map set to zero is
related to the map-model correlation for the original map (CC) by,

$\quad\quad$ CC' = CC sqrt(f) , $\quad\quad\quad\quad$ (2)

so that f, the fraction inside the mask, is approximately given by $f = CC'^2/CC^2$.

Automated model-building

The *Phenix* tool *phenix.map_to_model* has recently been described in detail (Terwilliger et al.,
2018c). The inputs required are a map file (CCP4/MRC format, Cheng et al., 2015), a sequence
file with the sequences of residues or nucleotides in each unique chain in the structure, and the
nominal resolution of the map. If symmetry was used in the reconstruction process, then the
symmetry operators can be supplied as well. All other parameters are fully optional and it is
normally not necessary for a user to adjust them.

For the model-building described here, the maps, sequence files, symmetry operators, and
resolution were all obtained from the 2016 Model Challenge web site at
http://challenges.emdatabank.org/?q=model_challenge.

The first step carried out by the map_to_model tool is to automatically sharpen the map with
the *phenix.auto_sharpen* tool (Terwilliger et al., 2018a). In this approach the map is sharpened
(or blurred) to attempt to simultaneously maximize the level of detail in the map and the
connectivity of the map.

The second step is to carry out automatic map segmentation as described above, yielding one
map that represents the unique part of the sharpened map along with a set of small maps each
representing one small region of connected density (all above a contour level determined
automatically during the segmentation process).

The third step is to carry out automatic model-building for each chain type that is represented
in the sequence file. This is done for the map representing the unique part of the sharpened

map and for each small map. Model-building is done using tools available in *Phenix* that include
placement of helices and strands in density of corresponding shapes (Terwilliger, 2010a;
Terwilliger, 2010b), tracing density along a chain and replacement with main-chain atoms
(Terwilliger, 2010c), placement of short fragments by convolution-based searches followed by
extension with 3-residue fragments from structures in the PDB (Terwilliger, 2003; Terwilliger et
al., 2018c), and recently-described methods for model-building of RNA that are extensions of
these procedures for protein (Terwilliger et al., 2018c).
227
228   The fourth step is to combine all the models. The principal method for combining models is to
229   rank all segments (fragments of a model that have no chain breaks) based on map-model
230   correlation, segment length, and secondary structure, then to go through this ranked list and
231   place whatever part of each model does not overlap with a higher-scoring model (Terwilliger et
232   al., 2018c).
233
234   After each model is built, after models are combined, and after application of reconstruction
235   symmetry to the final model, each working model is refined with real-space refinement
236   (Afonine et al., 2018a).
237
238
239
240   Data used from the Cryo-EM Model Challenge
241
The maps and reconstruction symmetry used for the 12 cryo-EM maps in the 2016 Cryo-EM
Model Challenge were taken from the Model Challenge site at
http://challenges.emdatabank.org/?q=model_challenge (accessed 2017-11-19). The Challenge
consisted of 8 unique molecules, four of which were associated with two maps at different
resolutions, leading to 12 different maps (Table I). Of these maps, most were associated with
previously-deposited models that were likely to be more accurate than the ones we built
automatically and that were therefore suitable for use as references for the accuracy of our
models. For one additional map (groEL, EMDB entry 6422) there was no deposited model,
however there is a model for a related structure in the PDB (1ss8) which we offset
superimposed on this map and used as a reference. One final structure was recently
interpreted (the proteasome structure; Veesler, D., unpublished) and we used that structure as
a reference model. We checked the map-model agreement with *phenix.map_model_cc* and
these map-model correlations ranged from 0.34 (rather low, supporting only low confidence in
the model), to 0.85, suggesting that the model is in good agreement with the map.
256
257   Results and Discussion
258
Fig. 1 illustrates the application of our map segmentation procedure (Terwilliger et al., 2018b)
to the 2.9 Å cryo-EM reconstruction of the anthrax protective antigen pore (Jiang et al., 2015).
The map has C7 symmetry (a 7-fold symmetry axis). Fig. 1A shows the 7-fold symmetry of the
pore and illustrates one of the 7 chains in purple. Fig. 1B shows the density map with 7-fold
symmetry. It can be seen that the density is much stronger for the extracellular region of the

molecule than for the transmembrane part below. The 7-fold symmetry was used along with
the map to identify symmetry-related regions of density in the map. Then a compact and
connected unique set of density regions was chosen to represent the molecule. Fig. 1C shows
the individual segmented regions of the map, and Fig. 1D shows the segmented region,
augmented by neighboring regions of density.
269
We then applied our segmentation procedure to a large set of cryo-EM maps from the EMDB
(Fig. 2). As expected, using the reconstruction symmetry of the maps in segmentation often
resulted in a very large reduction in the volume that needed to be considered to include the
unique part of each map (Fig. 2A). The average volume after segmentation and placing the
unique segmented region in a new box was 8% of the starting volume of the maps. In most
(206 of 218) of the cases illustrated in Fig. 2 we used the *add_neighbors* keyword to add a layer
of regions around the unique molecular volume in order to increase the chance of finding a
complete molecule. The 12 cases (EMD_2807, EMD_3137, EMD_5185, EMD_5600, EMD_6346,
EMD_6630, EMD_6637, EMD_6688, EMD_8598, EMD_8605, EMD_8644, EMD_9518) where
this was not done are those where the map was large (maps with 16M to 134M elements) and
we attempted to keep the size of the region to be worked on to the minimum possible.
281
Fig. 2B illustrates the fraction of the unique molecular unit that is within the unique segmented
region used for each map in Fig. 2A. This fraction of the molecule contained within the
segmented region is estimated from the map-model correlation between the molecule and a
map which is set to zero everywhere outside the segmented region, normalized to the map-
model correlation without setting any of the map to zero. If the molecule is within the
segmented region this normalized correlation will be unity, while if the molecule is split
between different segmented regions it will be smaller. As shown in Fig 2B, the fraction within
a single segmented region varies considerably among the 218 maps analyzed here, but the
mean fraction was 0.72, indicating that typically a large fraction, but not all, of the molecular
unit was contained within the segmented region.
292
We examined whether the fraction of the molecular unit contained within the segmented
region (Fig. 2B) depended on the number of symmetry copies or the resolution of the map. The
number of symmetry copies had only a small effect: maps with a single copy had an average
fraction of 0.73 and maps with 60 copies had an average of 0.71. On the other hand, resolution
had quite a substantial impact on the fraction within the segmented region: maps with
resolution of 3.5 Å or better had a mean fraction of 0.82; maps with resolution of 4 Å or worse
had a mean of 0.63.
300
We applied the combination of map sharpening, segmentation, and model-building as
implemented in the *Phenix* tool *phenix.map_to_model* (Terwilliger et al., 2018b) to the 12 maps
in the 2016 cryo-EM Challenge. The maps and corresponding reference models are listed in
Table I along with the CPU hours required for the analysis, which ranged from 7 to 422 hours.
Table II lists the number of residues that were built with C_α or P atoms within 3 Å of the
corresponding atoms in the reference model by the *phenix.map_to_model* procedure, along
with the fraction of the reference model represented by the model that was built and the

fraction of residues that were assigned the correct residue identity. The number of residues
built more than 3 Å from any residue in the reference model is also listed.
310
Overall, from 35% to 82% of the protein portions of the 12 structures were built within 3 Å of
the corresponding reference models. For the two RNA structures, 24% and 54% of the RNA
portions were built within 3 Å of the corresponding reference models. From 8% to 75% of the
protein and RNA sequences were correctly assigned.   For the non-ribosome structures, only a
small proportion of the models built did not correspond at all to the deposited models. On the
other hand, for the ribosome structures, a large fraction (over half for the 3.6 Å map) of the
protein residues built did not correspond to the deposited models. Most of these incorrectly-
built residues are located in regions that are RNA in the deposited models (recently we have
developed a tool, *phenix.remove_poor_fragments* that can remove some of these incorrectly-
built residues, but it was not available at the time of this work, TT, OS, PDA and PVA,
unpublished). The models built by *phenix.map_to_model* have RMS distances for Cα/P atoms
from reference interpretations of 0.6 Å to 2.1 Å.
323
324
The procedures developed here for map segmentation could be applied automatically to all of
the 218 maps that we examined in the tests shown in Fig. 2. Further, all 12 of the maps in the
2016 Cryo-EM Model Challenge could be automatically sharpened, segmented and partially
interpreted by the *phenix.map_to_model* procedure. It seems likely that combining the
techniques developed here with other approaches for automatic model-building might lead to
procedures that can automatically interpret an even larger part of cryo-EM maps.
331
332

333   Acknowledgements
334
This work was supported by the NIH (grant GM063210 to PDA and TT) and the *Phenix* Industrial
Consortium. This work was supported in part by the US Department of Energy under Contract
No. DE-AC02-05CH11231 at Lawrence Berkeley National Laboratory. This research used
resources provided by the Los Alamos National Laboratory Institutional Computing Program,
which is supported by the U.S. Department of Energy National Nuclear Security Administration
under Contract No. DE-AC52-06NA25396.Molecular graphics and analyses were performed with
the UCSF *Chimera* package (Petterson et al., 2005) and with *Coot* (Emsley et al., 2010).

Figure Legends
344

Figure 1. Segmentation of density for the anthrax protective antigen pore. A. Deposited
structure of anthrax protective antigen pore with one of the 7 chains in purple. B. Density map
illustrating the 7-fold symmetry used in the reconstruction. C. Individual segmented regions of
the map superimposed on a single chain from the deposited structure. Note that the deposited
structure was not used in the segmentation process. D. Illustration of the segmented region,
augmented by neighboring regions of density.
351

Figure 2. Histograms showing the results of application of the segmentation procedure to cryo-
EM maps from the EMDB. Datasets are grouped according to (panel A), the fraction of original
map required to represent the segmented region of each map, or (panel B), the fraction of each
molecular unit contained within the segmented region of each map. In each panel, the label
corresponds to the lower bound of each grouping. The values are grouped in increments of
0.05, so for example the number of datasets with values from 0.00 to 0.05 is shown over the
ordinate of "0".
359
360
361
362 References
363

364 Adams, P. D., Afonine, P.V., Bunkóczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.-
365 W. Kapral, G.J., Grosse-Kunstleve, R.W., McCoy, A.J., Moriarty, N.W., Oeffner, R., Read, R.J.,
366 Richardson, D.C., Richardson, J.S., Terwilliger, T.C. & Zwart, P.H., 2010. PHENIX: a
367 comprehensive Python-based system for macromolecular structure solution. Acta Cryst., D66,
368 213-221.
369
370 Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A., Adams, P.D.
371 (2018a). Real-space refinement in *Phenix* for cryo-EM and crystallography. Acta Cryst D., in
372 press.
373
374 Afonine, P.V., Klaholz, B.P., Moriarty, N.W., Poon, B.K., Sobolev, O.V., Terwilliger, T.C., Adams,
375 P.D., Urzhumtsev, A. (2018b). New tools for the analysis and validation of Cryo-EM maps and
376 atomic models. BioRxiv doi.org/10.1101/279844.
377
378 Baker, M.L., Yu, Z., Chiu, W., Bajaj, C., 2006. Automated segmentation of molecular subunits in
379 electron cryomicroscopy density maps. J. Structural Biology 156, 432-441.
380
381 Baker, M.L., Ju, T., Chiu, W., 2007. Identification of secondary structure elements in
382 intermediate-resolution density maps. Structure 15, 7–19.
383
384 Baker, M.L., Abeysinghe, S.S., Schuh, S., Coleman, R.A., Abrams, A., March, M.P., Hryc, C.F.,
385 Ruths, T., Chiu, W., Ju, T. (2011). Modeling protein structure at near atomic resolutions with
386 Gorgon. J. Struct. Biol. 174, 360-373.
387
388 Bai, X.C., Yan, C.Y., Yang, G.H., Lu, P.L., Ma, D., Sun, L.F., Zhou, R., Scheres, S.H.W., Shi, Y.G.
389 (2015). An atomic structure of human γ-secretase. Nature 525, 212-217.
390
391 Bartesaghi, A., Merk, A., Banerjee, S., Mathies, D., Wu, X, Milne, J.L., Subramaniam, S. (2015).
392 2.2 Å resolution cryo-EM structure of β-galactosidase in complex with a cell-permeant inhibitor.
393 Science 348, 1147-1151.
394

395 Bartesaghi, A., Matthies, D., Banerjee, S., Merk, A., Subramaniam, S. Structure of β-
396 galactosidase at 3.2- Å resolution obtained by cryo-electron microscopy. Proc. Natl. Acad. Sci.
397 USA 111, 11709-11714.
398
399 Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N. and
400 Bourne, P.E., 2000. The Protein Data Bank. Nucleic Acids Research, 28, 235-242.
401
402 Bernstein, F.C., Koetzle, T.F., Williams, G.J., Meyer, E.F., Jr., Brice, M.D., Rodgers, J.R., Kennard,
403 O., Shimanouchi, T. and Tasumi, M., 1977. The Protein Data Bank: a computer-based archival
404 file for macromolecular structures. J. Mol. Biol. 112, 535-542.
405
406 Campbell, M.G., Veesler, D., Cheng, A., Potter, C.S., Carragher, B. (2015). 2.8 Angstrom
407 resolution reconstruction of the Thermoplasma acidophilum 20S proteasome using cryo-
408 electron microscopy. eLife 4, e06380-e06380.
409
410 Cheng, A., Henderson R., Mastronarde, D., Ludtke, S., Schoenmakers, R.H.M., Short, J.,
411 Marabini, R., Dallakyan, S., Agard, D., Winn, M. (2015). J. Struct. Biol. 192, 146-150.
412
413 C. Colliex, C., Cowley, J. M., Dudarev, S. L., Fink, M., Gjønnes, J., Hilderbrandt, R., Howie, A.,
414 Lynch, D. F. , Peng, L. M. , Ren, G., Ross, A. W. , Smith, V. H., Jr, Spence, J. C. H. , Steeds, J. W.,
415 Wang, J., Whelan, M. J., Zvyagin, B. B. International Tables for Crystallography (2006). Vol. C, ch.
416 4.3, pp. 259-429.
417
418 Collins, P., Si, D. (2017). A graph based method for the prediction of backbone trace from cryo-
419 EM density maps. ACM-BCB '17 Proceedings of the 8th ACM International Conference on
420 Bioinformatics, Computational Biology, and Health Informatics.
421
422 Chen, M., Baldwin, P.R., Ludtke, S.J., Baker, M.L., 2016. De Novo modeling in cryo-EM density
423 maps with Pathwalking. J. Structural Biol. 196, 289-298.
424
425 DiMaio, F., Chiu, W. (2016). Tools for model building and optimization into near-atomic
426 resolution electron cryo-microscopy density maps. Methods. Enzymol. 679, 255-276.
427
428 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. (2010). Features and development of Coot.
429 Acta Cryst. D66, 486-501.
430
431 Fischer, N., Neumann, P., Konevega, A.L., Bock, L.V., Ficner, R., Rodnina, M.V., Stark, H. (2015).
432 Structure of the E. coli ribosome-EF-Tu complex at <3 A resolution by Cs-corrected cryo-EM.
433 Nature 520, 567-570.
434
435 Frenz, B., Walls, A.C., Egelman, E.H., Veesler, D., DiMaio, F., 2017. RosettaES: a sampling
436 strategy enabling automated interpretation of difficult cryo-EM maps. Naure Methods 14, 797-
437 803.
438

Fromm, S.A., Bharat, T.A., Jakobi, A.J., Hagen, W.J., Sachse, C. (2015). Seeing tobacco mosaic
virus through direct electron detectors. J. Struct. Biol. 189, 87-97.

Jiang, W., Baker, M.L., Ludtke, S.J., Chiu, W., 2001. Bridging the information gap: computational
tools for intermediate resolution structure interpretation. J. Mol. Biol. 308, 1033–1044.

Jiang J., Pentelute B.L., Collier R.J., Zhou, Z.H., 2015. Atomic structure of anthrax protective
antigen pore elucidates toxin translocation. Nature 521, 545-549.

Kong, Y., Ma, J., 2003. A structural-informatics approach for mining beta-sheets: locating sheets
in intermediate-resolution density maps. J. Mol. Biol. 332, 399–413.

Kong, Y., Zhang, X., Baker, T.S., Ma, J., 2004. A Structural-informatics approach for tracing beta-
sheets: building pseudo-C(alpha) traces for beta-strands in intermediate-resolution density
maps. J. Mol. Biol. 339, 117–130.

Lawson C.L., Patwardhan, A., Baker, M.L., Hryc, C., Garcia, E.S., Hudson, B.P., Lagerstedt, I.,
Ludtke, S.J., Pintilie, G., Sala, R., Westbrook, J.D., Berman, H.M., Kleywegt, G.J., Chiu, W. 2016.
EMDataBank unified data resource for 3DEM. Nucleic Acids Res. 44, D396–D403.

Li, X., Mooney P, Zheng, S., Booth, C., Braunfeld ,M.B., Gubbens, S., Agard, DA., Cheng, Y.
(2013). Electron counting and beam-induced motion correction enable near-atomic-resolution
single-particle cryoEM. Nat. Methods 10, 584-590.

Li, W., Liu, Z., Koripella, R.K., Langlois, R., Sanyal, S., Frank, J. (2015). Activation of GTP hydrolysis
in mRNA-tRNA translocation by elongation factor G. Sci. Adv. 1, e1500169.

Liao, M., Cao, E., Julius, D., Cheng, Y. (2013). Structure of the TRPV1 ion channel determined by
electron cryo-microscopy Nature 504, 107-112.

Lindert, S., Alenxander, N., Wotzel, N., Karakas, M., Stewart, P.L., Meiler, J., 2012. Em-Fold: De
novo atomic-detail protein structure determination from medium-resolution density maps.
Structure 20, 464-478.

Lu ,P.L., Bai, X.C., Ma, D., Xie, T., Yan, C.Y., Sun, L.F., Yang, G.H., Zhao, Y.Y., Zhou, R., Scheres,
S.H.W., Shi, Y.G. (2014). Three-dimensional structure of human gamma-secretase. Nature 512,
166-170.

Pettersen E.F., Goddard T.D., Huang C.C., Couch G.S., Greenblatt D.M., Meng E.C., Ferrin, T.E.
2004. UCSF Chimera--a visualization system for exploratory research and analysis. J Comput.
Chem. 25, 1605-1612.

Pintilie, G. D., Zhang, J., Goddard, T. D., Chiu, W., & Gossard, D. C., 2010. Quantitative analysis
of cryo-EM density map segmentation by watershed and scale-space filtering, and fitting of
structures by alignment to regions. Journal of Structural Biology 170, 427–438.
486
487 Roh, S.H., Hryc, C.F., Jeong, H.H., Fei, X., Jakana, J., Lorimer, G.H., Chiu, W. (2017). Subunit
488 conformational variation within individual GroEL oligomers resolved by Cryo-EM. Proc. Natl.
489 Acad. Sci. U.S.A. 114, 8259-8264.
490
491 Terwilliger, T. C. (2003). Automated main-chain model-building by template-matching and
492 iterative fragment extension. *Acta Cryst*. D59, 38-44.
493
494 Terwilliger, T. C. (2010a). Rapid model-building of β-sheets in electron density maps. Acta Cryst.
495 D66, 276-284.
496 Terwilliger, T. C. (2010b). Rapid model-building of α-helices in electron density maps. Acta
497 Cryst. D66, 268-275.
498
499 Terwilliger, T. C. (2010c). Rapid chain-tracing of polypeptide backbones in electron density
500 maps. Acta Cryst. D66, 285-294.
501
502 Terwilliger, T.C., Sobolev, O., Afonine, P.V., Adams, P.D. (2018a). Automated map sharpening by
503 maximization of detail and connectivity. bioRxiv doi: https://doi.org/10.1101/247049.
504
505 Terwilliger, T.C., Adams, P.D., Afonine, P.V., Sobolev, O.V. (2018b). A fully automatic method
506 yielding initial models from high-resolution electron cryo-microscopy maps. bioRxiv doi:
507 https://doi.org/10.1101/267138
508
509 Volkmann, N. (2002). A novel three-dimensional variant of the watershed transform for
510 segmentation of electron density maps. J. Struct. Biol. 138, 123–129.
511
512 Wang, Z., Hryc, C.F., Bammes, B., Afonine, P.V., Jakana, J., Chen, D.H., Liu, X, Baker, M.L., Kao,
513 C., Ludtke, S.J., Schmid, M.F., Adams, P.D., Chiu W. (2014). An atomic model of brome mosaic
514 virus using direct electron detection and real-space optimization. Nature Commun. 5 4808.
515
516 Wang, R. Y-R., Kudryashev, M., Li, X., Egelman, E.H., Basler, M., Cheng, Y., Baker, D., DiMaio, F.,
517 2015. De novo protein structure determination from near-atomic-resolution cryo-em maps.
518 Nature Methods 12, 335-341.
519
520
521 Yu, Z., Bajaj, C. (2008). Computational approaches for automatic structural analysis of large
522 biomolecular complexes. IEEE/AC Transactions on computational bology and bioinformatics 5
523 568-582.
524
525 Zhang, Q., Bettadapura, R., Bajaj, C. (2012). Macromolecular structure modeling from 3D EM
526 using VolRover 2.0. Biopolymers 97, 709-731.

528   Zhou N., Wang, H., Wang, J., 2017. EMBuilder: A Template Matching- based automatic model-

529   building program for high-resolution cryo-electron microscopy maps. Scientific Rep. 7, 2664.

532 Table I. Cryo-EM Model Challenge structures analyzed with *phenix.map_to_model*

Structure	Map	Reference model	Reso-lution	Reference map-model correlation	Cpu (h)	Reference
	(EMDB entry)	(PDB entry[1])	(Å)			
Beta galactosidase	2984	5A1A	2.2	0.73	32	Bartesaghi et al. (2015)
Proteasome	6287	undeposited[2]	2.8	0.81	16	Campbell et al. (2015)
E. coli 70S ribosome	2847	5AFi	2.9	0.85	422	Fischer et al. (2015)
Beta galactosidase	5995	3J7H	3.2	0.76	39	Bartesaghi et al. (2014)
Proteasome	5623	3J9i	3.3	0.77	14	Li et al. (2013)
trpV1	5778	3J5P	3.3	0.56	14	Liao et al. (2013)
TMV	2842	4UDV	3.4	0.73	7	Fromm et al. (2015)
Gamma secretase	3061	5A63	3.4	0.41	34	Bai et al. (2015)
E. coli 70S ribosome	6316	3JA1	3.6	0.38	322	Li et al. (2015)
Brome mosaic virus	6000	3J7L	3.8	0.76	16	Wang et al. (2014)
GroEL	6422	1SS8	4.1	0.83	153	Unpublished data[3]
Gamma secretase	2677	5A63	4.5	0.34	23	Lu et al. (2014)

---

[1] The PDB codes are written following the convention outlined in the editor's notes in the Computation Crystallography Newsletter (Comput. Cryst. Newsl. 2015:6; https://www.phenix-online.org/newsletter/CCN_2015_07.pdf).

[2] The recently-determined proteasome structure (Veesler, D., unpublished) was used as a reference model.

[3] The PDB entry 1ss8 was used as a model for entry EMD_6422, as used in the related entry EMD_8750 (Roh et al., 2017)

533 Table II. Results of Cryo-EM Challenge analysis with *phenix.map_to_model*

Structure	Resolution (Å)		Residues in reference model	Built within 3 Å (Residues)	RMS distance (Å)[1]	Built (%)	Matched to sequence[2] (%)	Built further than 3 Å (Residues)
Beta galactosidase	2.2		1022	842	0.6	82	75	10
Proteasome	2.8		422	327	0.6	78	59	3
E. coli 70S ribosome	2.9	Protein	6322	3212	0.7	51	23	2112
		RNA	4748	2566	1.0	54	49	280
Beta galactosidase	3.2		1022	676	1.2	66	23	26
Proteasome	3.3		427	246	1.2	58	48	4
trpV1	3.3		592	330	0.8	56	30	18
TMV	3.4		153	76	1.3	50	34	8
Gamma secretase	3.4		1223	832	1.0	68	24	80
E. coli 70S ribosome	3.6	Protein	7125	2479	2.0	35	8	3979
		RNA	4685	1140	1.9	24	26	440
Brome mosaic virus	3.8		479	198	1.5	41	11	14
GroEL	4.1		524	309	1.3	59	16	10
Gamma secretase	4.5		1223	619	2.1	51	8	185

---

[1] The RMS distances for $C\alpha$/P atoms from the reference interpretations, only including the residues built within 3 Å of the reference model.

[2] The percentage of matched to sequence is the total number of residues in the automatically-built model correctly matched to sequence divided by the total number of residues in the reference model.

Fig 1A

Fig 1B

Fig 1C

Fig 1D