1 **Joint single-cell DNA accessibility and protein epitope profiling**
2 **reveals environmental regulation of epigenomic heterogeneity**
3

4 Xingqi Chen[1,7], Ulrike M. Litzenburger[1,7]*, Yuning Wei[1,7], Alicia N. Schep[1,2, 3],
5 Edward L. LaGory[4], Hani Choudhry[5], Amato J. Giaccia[4], William J. Greenleaf[1,2,3],
6 Howard Y. Chang[1,6,]*
7

8 **Affiliations**
9 [1]Center for Personal Dynamic Regulomes, Stanford University, Stanford, CA
10 94305.
11 [2]Dept of Genetics, Stanford University, Stanford, CA 94305.
12 [3]Department of Applied Physics, Stanford University, Stanford, CA 94305.
13 [4] Division of Radiation and Cancer Biology, Department of Radiation Oncology,
14 Stanford University, Stanford, CA 94305
15 [5] Department of Biochemistry, Cancer Metabolism and Epigenetic Unit, Faculty of
16 Science, Cancer and Mutagenesis Unit, King Fahd Center for Medical
17 Research, King Abdulaziz University, Jeddah, Saudi Arabia.
18 [6] Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305.
19 [7] authors contributed equally
20 *Correspondence to: U.L. (litzenbu@stanford.edu) and H.Y.C.
21 (howchang@stanford.edu)

22
23

24 **Abstract:**
25 Here we introduce Protein-indexed Assay of Transposase Accessible Chromatin
26 with sequencing (Pi-ATAC) that combines single-cell chromatin and proteomic
27 profiling. In conjunction with DNA transposition, the levels of multiple cell surface
28 or intracellular protein epitopes are recorded by index flow cytometry and
29 positions in arrayed microwells, and then subject to molecular barcoding for
30 subsequent pooled analysis. Pi-ATAC simultaneously identifies the epigenomic
31 and proteomic heterogeneity in individual cells. Pi-ATAC reveals a casual link
32 between transcription factor abundance and DNA motif access, and deconvolute
33 cell types and states in the tumor microenvironment *in vivo*. We identify a
34 dominant role for hypoxia, marked by HIF1α protein, in the tumor
35 microvenvironment for shaping the regulome in a subset of epithelial tumor cells.

36
37
38

Introduction

Cell-to-cell variation is a universal feature that impacts normal development and human disease[1]. While recent advances in single cell research have improved our ability to document cellular phenotypic variation[1], the fundamental mechanisms that generate variability from identical DNA sequences remain elusive. Uncovering the molecular mechanism behind cellular heterogeneity would be helpful in clinical diagnosis, understanding the basic mechanism of developmental disorders, molecular basis of drug resistance in cancer and therapy of human diseases in the long term. In the last decades, studies revealed that chromatin structure is a main player regulating gene expression, and that it is tightly linked to heterogeneity in transcription and phenotype[2]. To fully understand the molecular mechanism determining cell-to-cell heterogeneity, it is essential to define the chromatin landscape in each individual cell.

Recent advances in single cell chromatin technologies, such as single cell ATAC-seq and single cell DNase-seq revealed the variation of chromatin organization across individual cells[3-5]. These technologies demonstrate that accessibility variance is associated with specific transcription factors and provide new insight into cellular variation of the "regulome"[3]. In these approaches, cells are randomly selected for next generation sequencing and the cellular variation is decoded using computational de-convolution. Thus, using available technologies, we only interpret the cellular variation and define subtypes indirect by clustering, dimensionality reduction such as principle component analysis method or projection onto a bulk scaffold. Therefore, until now, the cell-to-cell epigenetic variation cannot unambiguously be linked to the cellular phenotype or cell state. Staining of proteins for specific cell types and cell stages is helpful to indicate the cellular phenotype, for example phosphorylated Focal Adhesion Kinase for a migratory cell state[6] or HIF1α staining for cells in a hypoxic environment. While an extensive effort was put on increasing throughput of these single cell technologies[2,4], the direct linkage of cellular phenotype to the chromatin variation of individual cells remains largely ignored.

Here, we describe a novel single-cell approach, Protein-index single cell ATAC-seq (Pi-ATAC), in which we index and quantify protein expression using index Fluorescence Activated Cell Sorting (FACS) and enumerate the accessible DNA elements of the same individual cell. The combination of protein and epigenetic profile allows us to directly link the cellular phenotype and environment to the chromatin variation of individual cells. We applied Pi-ATAC to primary, heterogeneous mouse breast tumors and characterized cell states of

79 tumor-infiltrating immune cells as well as tumor cells simultaneously. In addition,
80 we link epigenetic variability of tumor cells to the hypoxic tumor
81 microenvironment. The described method allows to unbiasedly combine single-
82 cell ATAC-seq with traditional FACS and therefore would be relevant to wide
83 range of biology groups.
84
85 **Results**
86
Development of Pi-ATAC method
We were motivated to develop Protein-index single cell ATAC-seq (Pi-
ATAC) to provide two innovative advances for multiomics. First, PiATAC
enables intracellular protein analysis and DNA accessibility from the same
individual cell. Previously, ATAC-seq required live cells, and investigators could
only perform live cell sorting using cell surface markers. This limits the protein
analysis to the ~15% of cell surface proteins. We and others used conventional
flow cytometry with cell surface markers to isolate different cell types[7,8]. In Pi-
ATAC, we have developed a new method to crosslink cells and
perform intracellular protein analysis (including in the nucleus) jointly with single
cell ATAC-seq. Thus, Pi-ATAC opens the door for > 85%[9] of the proteome for
single cell multiomics.
99
Second, Pi-ATAC introduces the multimodel Indexing. In single cell
analysis, indexing means keeping track of molecules from one cell vs. all the
molecules from other cells. In Pi-ATAC, we accomplish the indexing of both
protein epitope levels and DNA regulatory landscape, and register these
orthogonal modalities at the same time, using tools widely accessible to many
investigators. Prior application of flow cytometry to ATAC-seq involved gates,
where many cells within a wide range of protein levels are lumped together. This
is a far cry from Pi-ATAC, where the level of individual protein epitopes in each
cell is precisely enumerated.
109
Pi-ATAC works on fixed cells or tissue, which then can be stored prior to
tagmentation, allowing collection of rare cells and pooling across multiple
experiments. As a result, investigators can prospectively focus their sequencing
power on rare but interesting cells. In more detail, in Pi-ATAC cells or tissue are
first fixed using paraformaldehyde (PFA), then gently dissociated and
permeabilized using the standard ATAC-seq lysis buffer (see **Methods**), followed
by antibody staining against protein epitopes of interest. As the cells are already
fixed and permeabilized, intracellular as well as intra-nuclear staining are
possible. Then, cells are transposed in bulk. The reaction is stopped by addition

of EDTA, without any purification step. As single cells are sorted into individual
wells containing the reverse crosslinking buffer (see **Methods**), fluorescence
intensities of antibodies against protein epitopes of interest are recorded and
assigned to the position of sorted cells. After reverse crosslinking, libraries are
prepared by barcoding PCR (**Figure 1a**).
The reverse crosslinking buffer used in Pi-ATAC was specially developed to be
compatible with the barcoding PCR step. In ATAC-see (a method for imaging
and sequencing DNA accessibility), tagmentation is also performed on fixed
cells, together with either intra-nuclear or cytoplasmic protein staining[10], and thus
the ATAC-see protocol also contains a reverse crosslinking step prior to a DNA
purification step. However, for single cell ATAC-seq, a DNA purification step is
not practical because the limited material from one single cell library is likely lost
during purification. Therefore, library preparation with single cell barcoding via
PCR is performed directly in the reverse crosslinking buffer. However, DNA Taq
polymerase is not compatible with sodium dodecyl sulfate (SDS) included in the
traditional reverse crosslinking buffer[11]. Hence, we developed a reverse
crosslinking buffer without SDS (see **Materials** and **Methods**), in which the DNA
Taq polymerase is not inhibited.
137
First we confirmed the functionality of the new reverse crosslinking buffer
followed directly by PCR without purification in bulk ATAC-seq libraries. Quality
of the bulk ATAC-seq libraries prepared using the new reverse crosslinking either
with or without DNA purification is comparable to the previous protocol of reverse
crosslinking buffer with purification, including (i) enrichment of fragments at
transcription start sites, (ii) fragment amount observed in open chromatin peaks
identified, and (iii) coverage on both gene promoters and distal elements
(**Supplementary Fig. 1a-e**). Together, we conclude that the new reverse
crosslinking buffer is compatible with the PCR reaction for library preparation in
Pi-ATAC.
148
Validation of Pi-ATAC workflow in single cells
We next validated the workflow and accuracy of FACS based scATAC-
seq. In order to evaluate the precision of FACS sorting, we used a mixture of two
cell types of different sizes and species ("barnyard experiment"): GM12878
human lymphoblastoid cell line (LCL) and V6.5 mouse embryonic stem cells
(ESC). Of note, the size difference between these two cell types is not a concern
in the FACS based scATAC-seq method, whereas the C1 microfluidics system
requires separation into two size-restricted integrated microfluidic chips (IFC).
First, both mouse ESC (n=144) and human LCL (n=144) were fixed with 1%
formaldehyde and mixed in a 1:1 ratio. Then ATAC was performed in bulk. Next,

we sorted single cells from the mixture of cells by FACS without regard to cell
type, and collected individual cells in 96-well plates containing the new reverse
crosslinking buffer. After sorting, plates were incubated at 65 degrees over 8hs,
followed by a Proteinase K inactivation step at 80 degrees. For library
preparation, PCR master mix and barcoding primers were added directly to the
individual wells. Finally, PCR products were pooled together; purified, and
sequenced. All fragments were then aligned to mouse and human genomes.
Indeed, we found that each individual well contained predominantly either human
or mouse DNA sequences, using a 500 fragment cutoff and 96% species
specificity (0 hybrid out of 288 cells, **Figure 1b,** see **Methods**), confirming the
precision of our index FACS sorting.

To assess the single cell Pi-ATAC library quality, we prepared libraries
from 192 GM17878 cells and comprehensively compared those to bulk GM12878
ATAC-seq datasets (n=4). Single cells that produced low quality data were
excluded from downstream analysis using nuclear fragment amount and fraction
of reads in accessible chromatin peaks, as previously described [12]. Of the 192
GM12878 cells, 168 (87.5 %) passed filter (**Supplementary Fig. 2a,** see Method
for details). Aggregation of these 168 single cells shows comparable accessibility
patterns as bulk ATAC-seq (**Supplementary Fig. 2b, Supplementary Fig. 2c**
R(77,855)=0.81, *p*<0.00001). We then measured transcription factor (TF)
variation scores in GM12878 Pi-ATAC data, using the computational tool
ChromVar [12] (**Methods**). The most significant variable TF motif was NF-κB family
(**Supplementary Fig. 2d**), which agrees with the previous report from the same
cell line by scATAC-seq approach[3].

Next, we compared Pi-ATAC data to publically available scATAC-seq
data[3]. GM12878 Pi-ATAC and GM12878 scATAC-seq data yielded similar
profiles of genome wide accessibility (*R*(77,855)=0.8, *p*<0.00001.
Supplementary Fig. 2e) as well as overall transcription factor motif variability **(***R*
(384) = 0.7, *p*<0.00001. **Supplementary Fig. 2f**). To ensure that antibody
staining of extra- or intracellular proteins does not interfere with the quality of Pi-
ATAC, 384 GM12878 cells were stained against the B-cell surface marker CD19
and intracellular phosphorylated NF-κB (**Supplementary Fig. 2g**). The antibody
staining did not substantially affect the quality of Pi-ATAC, with 298 (77.6%)
passing filter (**Supplementary Fig. 2h**). The stained GM12878 Pi-ATAC data
share concordance with previously published GM12878 scATAC-seq data in
accessibility of peaks (*R(77,855)* = 0.72, *p*<0.00001. **Supplementary Fig. 2i**), in
the genomic annotation of peak distribution (**Supplementary Fig. 2j**) and in
transcription factor motif accessibility at NF-κB and Jun motifs (**Supplementary**

Fig. 2k). Aggregation of these 298 single cells shows comparable accessibility
patterns as bulk ATAC-seq (**Figure 1c**). We next compared our GM12878 Pi-
ATAC data to available scATAC-seq data across multiple cell types[3]. Two-
dimensional t-distributed stochastic neighbor embedding (t-SNE) projection
based on TF motif accessibility revealed clustering of single cells largely
according to cell type, demonstrating Pi-ATAC GM12878 clusters together with
the published scATAC-seq GM12878 data and distinct from other cell types
(**Supplementary Fig. 2l**).
207
We next quantified the information content provided by single cells in Pi-
ATAC. Sparse DNA accessibility data from an individual cell suffice to infer TF
motif activity, based on the summed accessibility of DNA elements containing the
TF motif vs. background elements, as previously reported for scATAC-seq[12]
(**Supplementary Fig. 2d, 2l**). For individual DNA elements, we find that data
aggregation from 50 or more single cells by Pi-ATAC correlated significantly with
bulk ATAC-seq on peak accessibility quantifications (Pearson correlation
R=0.615 for 77,857 peaks in 50 cells, R=0.694 in 100 cells, $p<0.0001$,
FDR<0.01, **Figure 1d,** see **Methods**). This observation is controlled for cell-to-
cell variance and sequencing depth artifacts, as determined by 1,000 simulation
runs of subsampling equalized contribution of 500 fragments in peaks per single
cell from a randomly selected 50 cells subset. The same analysis produced
similar results in protein epitope-stained Pi-ATAC data vs. published GM12878
scATAC-seq data[3]. Similar results are obtained by down-sampling to 500
fragments in peaks from bulk ATAC-seq data to mimic single cell data ("mimic
from bulk" in **Figure 1d, see Methods**).
224
Together, these results indicate that Pi-ATAC-seq captures cell-cell
variation in active regulatory DNA as well as previous published scATAC-seq
protocols while providing protein epitope measurements. With appropriate cell
lineage and cell state characterization, Pi-ATAC data from appropriate single
cells can be aggregated to reproduce a "pseudo-bulk" genomic DNA accessibility
profile.
231
Pi-ATAC probes transcription factor protein abundance versus DNA motif
access
The K562 chronic myeloid leukemia (CML) cell line is a heterogeneous
cell mixture containing both precursor and differentiated cells[13]. Several key
transcription factors such as GATA1, GATA2 or TAL1 regulate hematopoietic
precursor status[14,15]. We, and others demonstrated that the self-renewal and
multipotency of K562 cells is correlated with high expression level of GATA2[16,17].

It is unclear whether GATA2 is a pioneer factor in this context, i.e. creating the
first DNA access at its cognate motif, or follows other TFs by accessing DNA
elements occupied by other TFs. Therefore, we applied Pi-ATAC to K562 cells
using GATA2 staining to measure the epigenetic status and heterogeneity of the
precursor cells within the K562 population (see **Methods**). To focus only on the
nuclear presence of GATA2, we isolated the K562 nuclei to exclude cytoplasmic
GATA2 as well as mitochondria contamination.
246
As expected, GATA2 staining intensity in K562 nuclei is variable, ranging
from 1 to 8400 RFU, clearly clustering into three groups, which we categorized
as high, medium and low (**Supplementary Fig. 3a**). Of the 288 nuclei sorted,
223 (77.4%) passed the filter (**Supplementary Fig. 3b**). In line with previous
scATAC-seq results[3], our Pi-ATAC results display highest variability in the motif
accessibility of the GATA and Jun/FOS family (**Supplementary Fig. 3c**). Within
the significant variable TF motifs, we observed two main sub-groups by TF motif
accessibility, one group enriched with Jun/Fos, and the other dominated by
GATA family motifs (**Supplementary Fig. 3d,** see **Methods**). We then asked
whether the GATA2 protein staining intensity correlates with the accessibility of
individual TF motifs, particularly the GATA2 motif itself. Surprisingly, we did not
find a significant correlation between accessibility of GATA2 motif itself with
GATA2 staining (**Supplementary Fig. 3e**), but we observed significant positive
correlations of GATA2 protein level with accessibility of other transcription factor
motifs, including TAL1::TCF3 (R=0.21), MTF1 (R=0.19), CEBPA (R=0.12), GSC
(R=0.15), SRF(R=0.16), and DUX(R=0.26, all with p < 0.05 and FDR<5%), which
are involved in maintaining hematopoietic cells in a de-differentiated cell
state[14,15,18-20] and determining hematopoietic cell fate in multipotential progenitor
cells[21] (**Supplementary Fig. 3e**). TAL1, TCF3 and MTF1 have been reported to
be potential binding partners in larger protein complexes with GATA2[22,23]. At the
same time, we also observed that the accessibility of TFAP motif is significantly
anti-correlated with GATA2 stain (R= -0.2, p < 0.05) (**Supplementary Fig. 3e)**.
TFAP was previously reported to drive hematopoietic differentiation[24], the
opposite role of GATA2 in CML cells. Based on direct quantification of nuclear
GATA2 protein and chromatin access from the same cell, our results suggest
that GATA2 cooperates with other TFs such as TAL1 and MTF1 to promote CML
cell self renewal in K562 cells.
274
275   Our findings suggest the feasibility to use Pi-ATAC to directly probe TF
276 mechanisms linking protein abundance, and DNA accessibility.
277
Simultaneous Pi-ATAC of tumor cells and immune cells

We next aimed to apply Pi-ATAC to primary tumor samples. Tumor tissue
is comprised of different cell types, including tumor, stromal and tumor-infiltrating
immune cells, each with heterogeneous cell sizes and cell surface markers.
Therefore we first tested whether Pi-ATAC will work on an artificial mixture of
mouse breast tumor cell line 4T1 and mouse splenocytes. This mixture allows us
to test (i) the sort efficiency of cells with comparable size differences as occurring
in the primary tumors, (ii) lineage-specific antibodies, such as the mutually
exclusive expression of EpCAM in the breast cancer cells and CD45 in
splenocytes and tumor-infiltrating immune cells. As expected, the mixture of
breast cancer cells and splenocytes are clearly distinguished and sorted using
EpCAM and CD45 antibodies (**Figure 2a, Supplementary Fig. 4a**). We sorted
192 cells and performed Pi-ATAC, of which 190 (99.0%) cells passed the filter
(**Supplementary Fig. 4b**). Importantly, aggregated single cells from EpCAM+
4T1 and CD45+ splenocytes show cell type-specific open chromatin at *Epcam*
and *Cd45* loci respectively (**Figure 2b**), confirming the specificity of the
antibodies and sorting. Within the top 30 out of 278 significant variable TF motifs,
we observed two clusters of chromatin accessibility profiles demonstrating 4T1
and mouse splenocytes respectively. Addition of the protein staining information
confirms the chromatin accessibility clustering and also reveals that three
spenocytes (1.5%) cluster with 4T1 cells (**Figure 2c, d**). At the same time, we
detect unique features of the chromatin landscape of both cell types: previously
reported Spic and Irf motifs in immune cells and Nfi family motifs in 4T1 tumor
cells (**Figure 2d**). Together, these results demonstrate the specificity and
precision of Pi-ATAC approach to deconvolute cell communities.
303
Pi-ATAC of breast tumor ecosystem
Cancer heterogeneity is a major driver of tumor evolution, progression,
and emergence of drug resistance. Although cancer heterogeneity can be
assessed with recent advances in single cell technologies, the focus lies mainly
on cell-to-cell transcriptional or mutational differences. The heterogeneity of
epigenetic information within a tumor, particularly in solid tumors, is largely
unexplored.
311
As a case study, we applied Pi-ATAC to the fast growing and highly
invasive *MMTV-PyMT* genetically engineered mouse breast tumor model[25] to
simultaneously dissect EpCAM+ tumor cells and tumor-infiltrating immune cells
(CD45+) from the same mouse breast tumor (**Figure 3a**, **Methods**). We stained
a single cell suspension of the dissected tumor with anti-EpCAM and anti-CD45
as above and performed Pi-ATAC on 384 cells (**Supplementary Fig. 5a**). We did
not exclude any tumor regions or cells, obtaining an unbiased view of the

composition of the tumor. Of all stained cells 4% were immune cells (CD45+),
28% were epithelial tumor cells (EpCAM+) (**Supplementary Fig. 5a**). The quality
of Pi-ATAC in these primary tumor cells isolated from a solid tumor was
equivalent to Pi-ATAC performed of any cell line with 369 (96.1%) cells passing
the filter (**Supplementary Fig. 5b**). Aggregated single cells from EpCAM+ and
CD45+ cells show cell type-specific open chromatin at *Epcam* and *Cd45* loci
respectively (**Supplementary Fig. 5c**).
326
Surprisingly, although the distinction of the protein staining is pronounced,
the t-SNE projection of EpCAM+ cells and CD45+ cells of bias-corrected
deviations for motifs did not form two distinct clusters (**Figure 3b**). However, the
protein information aids visualization of a polarity of the distribution of the
EpCAM+ cells and tumor-infiltrating immune cells and reveals mixing of the two
cell types. The previous observed clear separation of 4T1 and mouse splenocyte
epigenetic profiles excludes the possibility of antibody cross-contamination.
334
We next calculated the variability of TF motifs across all cells, resulting in 84
significant variable motifs (P < 0.05 after Benjamini-Hochberg correction on
multiple tests, **Supplementary Table 1**). Unsupervised hierarchical clustering of
the chromatin accessibility of the 84 TF motifs across 369 cells revealed three
modules of TF motif accessibilities (28 motifs in m1, 14 in m2 and 42 in m3) and
seven subgroups of single cells (s1-s7 with cluster size range from 27 to 68 cells)
(**Figure 3c, d(I), Supplementary Fig. 5d, e**). TF motifs of m1 consisting of
immune cell specific motif families (Spic, Runx, Ets, et. al) show uniform specific
accessibility among cells in s1 and s2 (**Figure 3c, d(II)**). M2 motifs, including
chromatin remodeling relevant TF motifs such as Smarcc1, Jun and Fos family
are consistently accessible in cells of s1 and s4 (**Figure 3c, d(III)**). M3 comprises
TF motif families Hmga, Hif, Nfi, Tead, p53, Grhl1 and Fox. We observed
heterogeneity across all cellular subgroups for TF motif accessibility in m3,
however highest mean motif accessibility in s3, s4 and s6 (**Figure 3c, d(IV),**
Supplementary Figure 6). Collectively, the seven different cellular subgroups
might reflect the heterogeneity of the composition of EpCAM+ tumor cells and
CD45+ tumor infiltrating immune cells.
352
Next, we aligned the FACS protein staining information to the seven
cellular subgroups (**Figure 3c, d(V)**). As expected, s1 and s2, the cellular
clusters exhibiting high chromatin accessibility of motifs associated with the
immune cell lineage, are significantly enriched with CD45+ cells (92.4% of 66
cells in s1 and 94.1% of 68 cells in s2, Chi-square test p<0.0001 for others, after
Benjamini-Hochberg correction of multiple tests, **Supplementary Table 2**). The
majority (102 out of 177 (57.6%)) of EpCAM+ cells fall into s3, s4 and s7 (76.7%

of 30 cells in s3, 88.2% of 68 cells in s4 and 90.7% of 43 cells in s7, enriched
with EpCAM+ cells, Chi-square test p<0.0001 for others, after Benjamini-
Hochberg correction of multiple tests, **Supplementary Table 2**). The remaining
75 of 177 EpCAM+ cells are in s5 and s6 mixed together with CD45+ cells.
EpCAM+ subgroups are characterized by accessibility of TF motifs in m3, a
module characterized by heterogeneous accessibility in various TF motifs such
as p53-, Tead-, Nfi- and Hif- families. This heterogeneous TF motif accessibility
in m3 prevents clear categorization of some cells, especially cells in s3, s5, s6
and s7; only the protein information guides the classification into immune and
tumor cells. Interestingly, m2 motifs associated with chromatin remodeling TFs
such as Smarcc and Fos show specific accessible in subgroups of both EPCAM+
(s1) and CD45+ (s4) cells (**Figure 3d(III)**), suggesting variable regulation on
different subgroups within each cell type. Our result demonstrates the Pi-ATAC
dissects chromatin profile of different cell types simultaneously from the primary
tumor, and the staining information is necessary to unambiguously define the cell
type from such complex system.
376
377   We also performed an unbiased analysis of sequence features
associated with variation in chromatin accessibility across tumor and immune
cells. We searched for enrichment of short nucleotide sequences of a specific
length k in accessible regions using ChromVAR. This k-mer analysis allows
discovery of novel motifs and is not restricted by any database entry. Because
most TF have core motifs between 5 and 8 bp, we use 6-, 7- and 8 k-mer
analysis to find enrichment of motifs of 8 or less base pairs (**Supplementary
Table 3-5**). The top enriched (p-value < 0.05 after Benjamini-Hochberg
correction of multiple tests) k-mer represents RUNX motif family, followed by
TEAD, Jun-Fos and NFI families (**Figure 3c, see Method**). As all of these k-
mers are already found using annotated TF motifs in CisBP database, we
decided to proceed using the CisPB results for downstream TF motif analysis.
389
390   Assuming the previous analyzed artificial mixture of immune and tumor
cells defines TF motif accessibility characteristic for these cell types, we
compared TF motif variability of 4T1 to splenocytes, and of CD45+ tumor
infiltrating immune cells to EpCAM+ primary tumor cells (**Figure 3e**). Of the 84
significant variable TF motifs across the primary cells 49 are shared between the
two datasets. Of note, while the top ten most variable motifs from the tumor-
immune cell mixture experiment are all m2 motifs shared with the primary cells,
most of the heterogeneous TF motif accessibility in m1 and m3 was not observed
as significantly variable in the cell line mixture. Thus, we next asked which TF
motifs are the main contributors to the heterogeneity in m3 and m1 assuming that

contributing TFs are unique features of *in vivo* tumors. Motifs along the y-axes
stand out as having additional variability of accessibility in tumor cells and
infiltrating immune cells compared to the artificial mixture (**Figure 3e**). Of these
motifs, the most variable ones are defining m1 of the hierarchical clustering
(**Figure 3e, lavender dots**). These results demonstrate that motifs accessible in
m1 are unique to the tumor infiltrating immune cells, because of their absence in
the splenocytes. While m1 motifs are accessible specifically in CD45+ cells, m3
accessibility is biased towards EpCAM+ cells (**Supplementary Figure6**).
Interestingly, the predominant TF motif specific to primary tumor cells enriched in
m3 is Hif (**Figure 3e pink dots**). We found the Hif motif to be the most significant
unique variable motif of the heterogeneous m3 module (**Figure 3d(IV)**). HIF
protein stability is tightly linked to the hypoxic environment[26], thus we
hypothesize that the hypoxic microenvironment in a solid tumor may modulate
the chromatin structure and contribute to the heterogeneity of chromatin
accessibility observed in the primary tumor.
415
416

Hypoxia drives single cell epigenetic variability in tumor microenvironment
To directly demonstrate that the hypoxic microenvironment influences the
tumor epigenetic landscape *in vivo*, we applied Pi-ATAC to EpCAM and HIF1α
stained cells in the *MMTV-PyMT* breast tumor model. We isolated 956 EpCAM+
cells and observed 762 negative, 139 low and 55 high HIF1α protein-positive
cells from the tumor (**Figure 4a**). Among these 956 cells analyzed by Pi-ATAC,
839 passed the filter (**Supplementary Fig. 7a**).
424

Focusing first on the accessibility profiles, unsupervised hierarchical
clustering revealed 2 motif clusters and 20 cell state subgroups (**Supplementary**
Figure 8a-c). The first motif cluster is dominated by high accessibility of motifs of
the ETS family, whereas the m2 is not showing clusters of accessible motifs. The
subdivision of all EpCAM+ cells into 20 subgroups suggests high epigenetic
heterogeneity within the tumor cells. Interestingly, HIF1α staining does not
correlate with any specific cell cluster, demonstrating that the protein information
provides an orthogonal measurement to the epigenetic profile.
433

We next focused on the epigenetic difference between the HIF1α positive
groups. The different groups of HIF1α level might reflect the distance of the cell
from the hypoxic center in the tumor or the time a cell was exposed to hypoxia.
The variability of transcription factor motif accessibility profiles across the three
HIF staining groups is substantially different (**Figure 4b, Supplementary Table**
6). This difference is not due to the variable sample size of the HIF staining

groups as demonstrated by downsampling simulations (**Supplementary Fig. 7b,**
Supplementary Table 6)**.** The variability of transcription factor motif accessibility
is the highest among cells not experiencing hypoxia, which are also the majority
of cells in the population (**Figure 4b, Supplementary Fig. 7b**). Among the
strongest variable motifs in the negative HIF1α group are Fos, Smarcc1 and Batf,
which were observed to be accessible in a subgroup of EpCAM+ cells all
previous experiments. The variability of low and high HIF1α stained cells is
similar; they are dominated by p53 TF family motifs. Hypoxia is known to
increase p53-protein levels via several mechanisms, which may be reflected in
TF motif activity[27]. Additional TF motifs that are highly variable in hypoxic cells
include Grainyhead and Snail, TFs involved in epithelial-mesenchymal transition
and known mediator of breast cancer metastasis[28,29] and might reflect the
evading mechanism observed in hypoxic cells[27]. These TF motifs as well as the
more uniform distribution of accessibility across all cells are in line with the
previous observed heterogeneous TF motif accessibility.
455
Fox and E2f3 were previously reported to be regulating transcription of
HIF target genes[30,31]. Indeed, the tumor Pi-ATAC data demonstrate a significant
increase of DNA accessibility of motifs for Fox family and E2f3 in the cells with
high HIF1α protein (Wilcoxon test $P < 0.05$, **Figure 4c**) in a more detailed
comparison of transcription factor motif accessibility across the three groups. At
the same time, we observed TF motifs of Pit1 and Atoh8 lost significant DNA
accessibility (Wilcoxon test $P < 0.05$) in cells with high HIF1α protein
(**Supplementary Fig. 9**). Atoh8 expression was reported to be reduced in
hypoxic conditions[32]. Pit1 is an important developmental factor[33] associated with
cancer but not previously associated to be regulated by the HIF network.
Surprisingly, we found on average decreased accessibility at HIF motifs in HIF1α
high cells compared to negative and low (**Figure 4c,** Wilcoxon test $p<0.05$).
However, we also observe a bimodal distribution of HIF motif accessibility within
this HIF1α high group (**Figure 4c, p<0.05 for Hartigans' dip statistic**). This
feature is not due to variable sample size across the three staining groups, as
shown by down-sampling simulation in negative and low HIF1α clusters to the
same number of cells as found in the HIF1α high group (*n*=42) (**Supplementary**
Fig. 10, Supplementary Table7). This bimodality shows heterogeneity in HIF
motif accessibility, which might reflect distance to the hypoxic center or time of
hypoxia exposure.
476
By organizing single cells based on HIF1α protein levels, we now can
interpret a subset of DNA accessibility dynamics that likely reflect the hypoxic
microenvironment modulating cooperating transcription factors acting on target

gene promoters and enhancers.

To confirm if the DNA accessibilities of tumor cells, particularly HIF, Fox,
E2f3, Atoh, Pit1 motifs, are indeed modulated by hypoxia and no other tumor
microenvironmental stimuli, we used an *in vitro* cell line culture system to
simulate the *in vivo* hypoxia microenvironment. We incubated the mouse breast
tumor cell line 4T1 in 1% O_2 for various time points (6hs, 12hs, 24hs, 36hs, 48hs
and 72hs). To avoid the degradation of HIF proteins under normoxia, we fixed
and collected cells in 1% O_2 followed by HIF1α protein staining and standard
bulk ATAC-seq at the indicated time points. Increased HIF protein stability was
then measured by flow cytometric analysis. We observe an immediate increase
in HIF1α protein abundance followed by a stronger increase and saturation at
about 48hs – 72hs in low O_2 (**Figure 4d**). That observation confirms that the 4T1
cells experienced bona fide hypoxia.

All bulk ATAC-seq libraries have good reproducibility at all the time points
(**Supplementary Fig. 11a-c**). We hypothesized that we might observe dynamic
chromatin changes due to low oxygen and stabilized HIF, some of which might
involve direct targets of HIF or co-factors or factors inducing HIF. We indeed
observe changes in the chromatin accessibility across time in low oxygen, with
635 at 6h to over 1000 sites opening at 72hs compared to normoxic cells.
Interestingly, only a small number of DNA elements (~20-70) lose DNA access
during this treatment (**Figure 4e**). These differential accessible chromatin sites
were annotated to 755 significantly differential accessible transcription factor
motifs, using the Cis-BP database (**Supplementary Fig. 11d, Supplementary
Table8,** see **Methods**).

This reference time course allowed us to next focus on the dynamic
changes of transcription factor motifs, which we assume to be involved in hypoxic
responses of tumor cells, because they are enriched in HIF1α high group in the
Pi-ATAC tumor experiment (**Figure 4c**). Indeed, we found that the chromatin
accessibilities are affected at HIF, Fox, E2f3, Ptx1 and Atoh8 motifs during the
different time points of hypoxia incubation (**Figure 4f, Supplementary Fig. 12**).
Also *in vitro* HIF motif accessibility undergoes dynamic changes, showing both
decrease and then increase at different time points of hypoxia treatment
confirming the slight decrease and variability observed *in vivo*. Foxp1 and E2f3
motif gain DNA accessibility with increasing duration of hypoxia incubation
(**Figure 4f**). At the same time, accessibility of Pit1 and Atoh8 motifs decrease
with longer hypoxia incubation, both results confirming the accessibility
differences observed in HIF1α high vs. negative group *in vivo* (**Supplementary**

Fig. 12).

In summary, Pi-ATAC is the first technique that allows directly linking the

microenvironment to epigenetic heterogeneity in a tumor.

Discussion

Here, we develop a novel single-cell technology, Pi-ATAC, that

simultaneously measures protein epitopes and active DNA regulatory elements

of the same individual cell. Pi-ATAC enables investigation of the basic molecular

mechanism of cellular heterogeneity from multiple angles.

Pi-ATAC directly links the cellular phenotype or microenvironment to the

epigenetic profile, because it uses antibody staining and index FACS sorting to

combine protein abundance information with the epigenetic profile of a cell.

Powerful applications are exemplified by our experiments quantifying the protein

levels of transcription factors NF-kB and HIF1α, both regulated by intricate post-

translational modification, and measuring their DNA occupancy at the same

time. Furthermore, we demonstrate that Pi-ATAC is applicable to both fixed cells

from cultured cells and solid tumors *in vivo*, without any cell size restrictions.

In Pi-ATAC, choice of antibodies allows to combine phenotypic

characterization of cells with their epigenetic profile. Pi-ATAC can directly link

protein abundance or modifications with chromatin landscape to address

mechanisms of gene regulation. We demonstrate that the epigenetic profile does

not necessarily have a one-to-one relationship with the "phenotype" indicated by

protein stain, demonstrating that both together give a more complete picture of

cell state than either one alone. The impact of this precision analysis is

highlighted in our analysis of tumor hypoxia, where the intermediate level of HIF1

protein induction is linked to tumor suppression pathway.

The tumor microenvironment has been recognized as an integral

component of malignancies in mammary and other carcinomas, contributing in

confounding ways to tumor progression, metastasis, therapy resistance, and

disease recurrence[34]. In the tumor microenvironment immune cells change from

an immune-active to an immune-suppressive state, stromal cells turn into tumor-

promoting tissue. The altered phenotype of these cells is proposed to be mainly

due to epigenetic deregulation of gene expression[35]; however until now, a direct

measurement linking the microenvironment with epigenetic changes does not

exist. These epigenetic changes are highly dynamic, causing heterogeneity of

cell states within a tumor, which contributes to drug resistance, and tumor

recurrence[36]. By directly linking the environment with HIF1α protein accumulation

and the chromatin profiles of single cells, we show that tumor hypoxia dominantly
shapes the regulome profiles of both parenchymal tumor cells and infiltrating
immune cells, becoming the dominant epigenomic signature for a minority of
cells in the tumor. Moreover, different levels or durations of hypoxia can induce
distinct TF activity states that expand tumor heterogeneity. In the future, Pi-ATAC
may be combined with single cell RNA-seq from the same cell to understand
gene regulation and cell identity of the exact same cell. The biological insight of
this new technique will be of great value for single cell research in future.
568
In summary, Pi-ATAC is easily scalable, fast and economical (~$0.15/cell
– $1.50/cell, cost calculation in **Methods**). Pi-ATAC is high throughput and easily
scalable; with sorting into 384 well plates and use of 96x90 adapter combinations
to index 8640 single cells in one experiment. Importantly, Pi-ATAC can be
performed without any specialized instrument or reagents, allowing ready access
to the broad community. In our experiments we sort and ATAC up to 1000 cells
for a tumor experiment, demonstrating scalability and throughput. Pi-ATAC may
be useful to address potential prognostic and therapeutic opportunities revealed
by these insights in future studies.

Methods:

Cell culture: GM12878 were purchased from Coriell Institute and grown in RPMI
1640 (11875-093, Gibco), 2 mM L-glutamine (25030-081, Gibco), 15% fetal
bovine serum (07905, Gibco) and 1% Pen/Strep (15140-122, Gibco). 4T1 mouse
breast cancer cells were purchased by ATCC (CRL-2539) and cultured in RPMI
1640 supplemented with 10% fetal bovine serum (07905, Gibco) and 1%
Pen/Strep (15140-122, Gibco). K562 human chronic myeloid leukemia cells were
purchased from ATCC (CCL-243) and cultured in IMDM supplemented with 10%
fetal bovine serum (07905, Gibco) and 1% Pen/Strep. V6.5 Mouse embryonic
stem cells (mESCs, Novus Biologicals) were cultured in
DMEM/F12 supplemented with N2, 10 ng/ml bFGF (R&D Systems), and 1,000
U/ml of LIF. For the 4T1 in vitro hypoxia experiments, cell lines were cultured in a
Ruskinn Invivo2 workstation maintained at 1% O_2 and 5% CO_2, and cells were
collected with trypsin digestion in the Ruskinn Invivo2 workstation and fixed
immediately after 6hs, 12hs, 24hs, 36hs, 48hs and 72hs incubation.

Mice: MMTV-PyMT mice were ordered from The Jackson Laboratory and bred
and housed in the Stanford University Research Animal Facility in accordance
with the guidelines (APLAC #14046).
596
Splenocyte isolation: A spleen was dissected from a MMTV-PyMT mouse
which did not have a tumor at the time of euthanization. Then the spleen was
sliced into small pieces. Single cells were isolated from these pieces by placing
them onto a 70 μm cell strainer and mechanically pushing the cells with a plunger
through the strainer. Red blood cells were permeabilized using ACK buffer
(Thermo Fisher, A1049201). After washing cells twice with RPMI 1640 medium,
cells were counted and used in the 4T1–splenocyte mixture Pi-ATAC experiment.

Mouse breast tumor dissection and dissociation: Tumor growth has been
monitored and mice were sacrificed at 3 month of age, when tumors reached a 1
cm^3 size. The mouse was immediately perfused using 4% Paraformaldehyde,
which was subsequently quenched with 0.125 M glycine. The tumor was
dissected and first cut into small pieces on ice, followed by enzymatic digestion
using 200 U CollagenaseV in HBSS + 0.1% FBS for 1 h at 37 degrees rotating.
After centrifugation cells were resuspended in PBS and this suspension was
filtered through a 70 μm cell strainer to remove any remaining clumps. Cells can
be stored in PBS for at least 1 week at 4 degrees.

Bulk ATAC-Seq: Fixed ATAC-seq was performed as previous described[10].
GM12878 cells were fixed with 1% formaldehyde (Sigma, USA) for 10 min and

quenched with 0.125 M glycine for 5 min at room temperature. After the fixation,
cells were counted and 50,000 cells were used per ATAC-seq reaction. The
transposition reaction follows the normal ATAC-seq protocol. After the
transposition, a reverse crosslink solution (final concentration of 50 mM Tris-Cl,
1mM EDTA, 1% SDS, 0.2M NaCl, 5 ng/ml proteinase K) was added up to 200 μl.
The mixture was incubated at 65 °C with 1200 rpm shaking in a heat block
overnight, then purified with Qiagen Mini-purification kit and eluted in 10 μl
Qiagen EB elution buffer. Sequencing libraries were prepared following the
original ATAC-seq protocol[37]. The sequencing was performed on Illumina Hi-Seq
at the Stanford Functional Genomics Facility.

Bulk Pi-ATAC-seq without elution: GM12878 cells were fixed with 1%
formaldehyde (Sigma, USA) for 10 min and quenched with 0.125 M glycine for 5
min at room temperature. After the fixation, cells were counted and 50,000 cells
were used per ATAC-seq reaction. For bulk Pi-ATAC of 4T1 cells, cells were first
fixed in hypoxia, then counted and permeabilized, followed by transposition. The
transposition reaction is the same as for standard ATAC-seq except with 0.05%
Igepal CA-630 in the lysis buffer. After 30 min transposition, at 37 degrees, the
reaction was quenched using 40 mM EDTA, then cells were centrifuged and
supernatant was discarded. 20 μl of reverse crosslinking solution was added to
the cell pellet (with final concentration of 50 mM Tris-HCl pH 8.0, 0.5% Tween20,
0.5% Igepal CA-630, 5 ng/ml proteinase K). Reverse crosslinking was performed
65 degree overnight and inactivated by a 10 min incubation at 80 degrees the
next day. The 25 μl PCR master mix (NEB, M0541S) and 5 μl of two unique
primer combinations were directly added to the reverse crosslinking mixture and
PCR was performed as previous described[38]. The PCR product was purified with
Qiagen MinElute purification kit and eluted in 20 μl Qiagen EB elution buffer. The
75x2 paired end sequencing was performed on Illumina HiSeq4000 at the
Stanford Functional Genomics Facility.

Pi-ATAC of mixture of GM12878 and mES cells:
50,000 GM12878 and 50,000 mouse ESCs were mixed 1:1. The mixture was
fixed with 1 % formaldehyde (Sigma, USA) for 10 min and quenched with 0.125
M glycine for 5 min at room temperature. Then, cells were permeabilized using
standard ATAC lysis buffer (10 mM Tris ph7.5, 10 mM NaCl, 3 mM MgCl2, fresh
0.1% NP40) and immediately spun down. Cells were then transposed in bulk; for
100,000 cells 2x ATAC reactions were calculated. After 30 min tagmentation,
reaction was quenched by addition of 40 mM EDTA. Cells were centrifuged,
supernatant discarded and pellet resuspended in PBS for single cell index
sorting. In the analysis, all alignments below 500 fragments were disregarded.
653

Pi-ATAC of mixture of 4T1 cells and splenocytes:
50,000 4T1 and 50,000 splenocytes were mixed 1:1. The mixture was fixed with 1 % formaldehyde (Sigma, USA) for 10 min and quenched with 0.125 M glycine for 5 min at room temperature. Then, cells were permeabilized using standard ATAC lysis buffer (10 mM Tris ph7.5, 10 mM NaCl, 3 mM MgCl2, fresh 0.1% NP40) and immediately spun down. Cells were then transposed in bulk; for 100,000 cells 2x ATAC reactions were calculated. After 30 min tagmentation, reaction was quenched by addition of 40 mM EDTA. After that, cells were stained at room temperature with anti mouse EpCAM-BV421 (BD Horizon, #563214) CD45-Alexa488 (BioLegend #103122, clone 30-F11) for 30 min. Cells were then washed twice, the final pellet resuspended in PBS for single cell index sorting.

Pi-ATAC of GM12878 cells: A detailed step-by-step Pi-ATAC protocol can be found at https://www.protocols.io/private/F59D7D2F8FD5E57A20E039E9CF7A9785 and will be publicly available upon publication.
GM12878 cells were counted and 100 000 cells were fixed with 1 % formaldehyde (Sigma, USA) for 10 min and quenched with 0.125 M glycine for 5 min at room temperature. Then, cells were permeabilized using standard ATAC lysis buffer (10 mM Tris ph7.5, 10 mM NaCl, 3 mM MgCl2, fresh 0.1% NP40) and immediately spun down. For indicated experiments 100,000 cells were then stained at room temperature with anti human CD19-PE Clone H1B19 (BioLegend 302207) and rabbit anti human Phospho-NF-kB p65 (Ser536, CellSignaling) followed by donkey anti rabbit 488 secondary antibody. After staining (30 min each antibody), cells were transposed in bulk; for 100,000 cells 2x ATAC reactions were calculated. After 30 min tagmentation, reaction was quenched by addition of 40 mM EDTA. Cells were centrifuged, supernatant discarded and pellet resuspended in PBS for single cell index sorting.

Index sorting and library preparation for Pi-ATAC
Single cells were index-sorted into 96-well plates without any specific staining selection using the FACS AriaII (BD Biosciences). The plates were pre-filled with 20 µl reverse crosslinking buffer (see above) per well. For each experiment, index sort files (fcs) were exported using BD FACS Software and then further analyzed by FlowJo and R.
For reverse crosslinking all 96-well plates were incubated over night at 65 degrees, then proteinase K was inactivated by 10 min incubation at 80 degrees.

Single cell libraries were prepared by adding 25 µl 2 x PCR Master Mix (NEBNext High fidelity, NEB) and 2.5 µl 25 mM barcoding primer to each well3. PCR Cycling conditions: 72°C for 5min; 98°C for 30 s; 20 cycles at 98°C for 10 s,

63°C for 30 s, and 72°C for 1 min. After that, all wells were pooled, purified using
the MinElute kit from Qiagen and eluted in 20 µl Qiagen EB elution buffer.   The
75x2 paired end sequencing was performed on Illumina HiSeq4000 at the
Stanford Functional Genomics Facility. On average we sequenced ~1000 single
cells on one HiSeq 4000 lane, resulting in an average of ~0.3 millions reads per
cell.

Pi-ATAC of K562 cells: First, we isolated the K562 nuclei by following the
protocol as reported[39]. For the Pi-ATAC, 200,000 Nuclei were transposed; for
100,000 nuclei 2x ATAC reactions were calculated. After 30 min tagmentation,
reaction was quenched by addition of 40 mM EDTA. After that, cells were stained
at room temperature with mouse anti human GATA2 (Abnova, 1-102) 30 min at
room temperature, followed by donkey anti mouse 488 secondary antibody. Cells
were then washed twice, the final pellet resuspended in PBS for single cell index
sorting.
708
Pi-ATAC of tumor cells: 1*10^6 Cells isolated from a PyMT mouse breast tumor
(see above) were stained for 30 min at room temperature for rat anti mouse
EpCAM-BV421 (BD Horizon, #563214), HIF1□-PE (R&D, #IC1935P), CD45-
Alexa488 (BioLegend #103122, clone 30-F11), followed by donkey anti rabbit
488 secondary antibody. After staining cells were transposed in bulk. Here, for
4000 final cells we tagmented 400,000 cells, meaning 8x ATAC reactions were
calculated. After 30 min tagmentation, reaction was quenched by addition of 40
mM EDTA. Cells were centrifuged, supernatant discarded and pellet
resuspended in PBS for single cell index sorting.
718
719 Single cells were index-sorted into 96-well plates without any specific staining
720 selection using the FACS AriaII (BD Biosciences). The plates were pre-filled with
721 20 µl reverse crosslinking buffer (see above) per well. For each experiment,
722 index sort files (fcs) were exported using BD FACS Software and then further
723 analyzed by FlowJo and R.
724
For reverse crosslinking all 96-well plates were incubated over night at 65
degrees, then proteinase K was inactivated by 10 min incubation at 80 degrees.
Single cell libraries were prepared by adding 25 µl 2 x PCR Master Mix
(NEBNext High fidelity, NEB) and 5 µl of two unique primer combinations, the
PCR Cycling conditions: 72°C for 5min; 98°C for 30 s; 20 cycles at 98°C for 10 s,
63°C for 30 s, and 72°C for 1 min. After that, all wells were pooled, purified using
the MinElute kit from Qiagen and eluted in 20 µl Qiagen EB elution buffer.

Pi-ATAC library purification, quantification and sequencing: The pooled
single cell Pi-ATAC library were loaded on 6% PAGE, run at 160 V for
approximately 10 minutes purification, stain the gel with SYBR Gold. One 0.5ml
tube was prepared for one sample, and a hole was made in the bottom of 0.5ml
tube with 20 gauge needle. The gel slice with range of above 150bp DNA were
cut and put in the 0.5ml (with a hole in the bottom). Next, the 0.5ml tube is put in
the 2ml tube and centrifuged with max speed for 3 minutes at room temp. After
removing the 0.5ml tubes, and 300ul Buffer (500 mM NaCl, 1 mM EDTA, 0.5%
SDS) was added in the gel. The gel is incubated at 55 degree overnight with
1400 rpm shaking. Next day, the gel is centrifuged with max speed for 3 minutes
at room temp, and the supernatant is recovered for DNA purification with ChIP
DNA Clean & Concentrator Kits (ZYMO RESEARCH, D5205). The recovered
DNA is quantified with Agilent High Sensitivity DNA ChIP and sequenced on
Illumina Hi-Seq at Stanford Functional Genomics Facility.

Pi-ATAC cost per cell: For Pi-ATAC, reagent costs is largely driven by the PCR
master mix. Cost estimate per cell: Tn5 enzyme (Illumina, $0.00049 per cell,
assuming 2.5 µl Tn5 per 50,000 cells), NEBNext® High-Fidelity 2X PCR Master
Mix,(NEB, M0541) and primers (~$1.30 per cell) and negligible costs for
antibodies per cell (about $3 per protein per 10^6 cells). In summary, it costs about
$1.50 per cell with 50 microliter PCR reaction system. In addition, we had
succeeded with low scale volume PCR reaction system at 5 microliter per well of
a 96-well plate, where the cost is ~$0.15 per cell.

Bulk Pi-ATAC data preprocessing: Paired-end reads were trimmed for Illumina
adapter sequences and transposase sequences using custom-written script and
mapped to hg19 or mm9 respectively using Bowtie2[40] v2.1.0 with parameters –
"very sensitive". Reads were subsequently filtered for alignment quality of > Q30
and were required to be properly paired. Duplicate reads were removed with
Picard (http://picard.sourceforge.net) v1.79. Reads mapping to the mitochondria
were removed and not considered. The uniquely mapped reads are merged
before peak calling if bulk samples origin from the same cell/tissue/condition.
Peak calling was performed by MACS2[41] narrow peak mode with parameters –q
0.01 –nomodel –shift 0. Peaks were filtered for following categories: (1)
mitochondrial fragments inserted in nuclear genomic sequences, by Nuclear
MiTochondrial   Sequences   (NumtS)
(http://hgdownload.cse.ucsc.edu/goldenPath/hg19/database/   or
http://hgdownload.cse.ucsc.edu/goldenPath/mm9/database/).
and (2) the consensus excludable ENCODE blacklist genomic regions
(http://mitra.stanford.edu/kundaje/akundaje/release/blacklists/hg19-human/hg19-
blacklist.xls or http://mitra.stanford.edu/kundaje/akundaje/release/blacklists/mm9-

mouse/mm9-blacklist.bed.gz).

To process comparison among bulk samples from multiple cell/tissue/conditions,
peak sets from all conditions were merged by bedtools merge to a consensus
peak list. Number of raw reads mapped to each peak in each sample was
quantified using multicov in Beltools. Peak raw counts were normalized using
DESeq2.

Pi-ATAC data processing: All single cell Pi-ATAC data were preprocessed with
custom-written script as previous described[3] and bulk preprocessed. Briefly, pair-
ended reads were trimmed with adapter sequences, were mapped to hg19 or
mm9 with Bowtie2 v2.1.0 with parameter-X2000. Uniquely mapped reads were
filtered for low-quality reads, duplicates, and reads mapped to mitochondrial
genome. Peak summit calling was processed with aggregated single cells of one
cell/tissue condition, by MACS2 narrow peak mode with parameters –q 0.05 –
nomodel –shift 0 –call summits. We annotated peaks as the summit-centered
500 base pair long genomic region. The top 50,000 peaks with least q value
reported by MACS2 were included for downstream analysis in most datasets;
whereas top 30,000 peaks were included for K562 dataset. Single cell libraries
were further filtered with requirement of at least 500 fragments mapped to
genome and at least 0.5 times the median proportion of fragments in peaks
reached by aggregated single-cells of the same cell/tissue condition
(chromVAR)[12].

Pi-ATAC data information content analysis:
The quantification of fragments mapped to each peak for the genomic DNA
accessibility was compared between bulk GM12878 Pi-ATAC samples and single
cell Pi-ATAC. Peak quantification of bulk data was calculated as mean value of
the four samples. Peak quantification of Pi-ATAC was calculated for single cell as
well as for single cells aggregation in group-size five, ten, 20, 50, 100 and 150.
To adjust cell-cell variance, 1,000 times of simulation was processed to randomly
select cells from the 192 GM cells into the group. To adjust sequencing depth
artifact, 500 of fragments passing the final filter were randomly subsampled from
the bam file for each cell in each time of simulation. The same process was
conducted for public scATAC-seq GM12878 data[3]. We also generated 50 'mimic'
single cell data through down-sampling 500 fragments passing the final filter
directly from each of the four bulk samples. Pearson correlation coefficient with
significance was calculated between peak quantification of bulk and single cell
data from each approach.

TF deviation and variability analysis: Single-cell Pi-ATAC data processing and
calculation of TF deviation was performed using the R package chromVAR as
reported[12]. TF motifs were derived from the JASPAR dataset for human data,
while Homer and Cis-BP databases were used for mouse data, with R package
motifmatchr. Briefly, for each TF, the accessibility in each cell was calculated by
subtracting the number of Pi-ATAC reads in peaks covering the corresponding
motif in the cell, and was normalized by the total TF accessibility in the cell. The
accessibility deviation value for the TF in each cell is subtracted by the mean
accessibility calculated for sets of background peaks with similar accessibility
and GC content to obtain a bias-correction, and further divided by standard
deviation of the deviations calculated for the background peak sets.
TF deviation for the hypoxic time course bulk ATAC-seq was calculated using the
R package chromVAR.
The Z-score of deviations for each TF is used for visualized in both t-SNE
projection plots and heatmaps of unsupervised hierarchical clustering based on
correlation metric, by R packages Rtsne and pheatmap.
The variability of the TF motif across single cells in the sample set was
determined by the standard deviation of the TF deviations across the cells. The
metrics is close to 1 if the motif is not significantly more variable compared to the
background peak list of that motif. To remove artifact of cell amount difference
among the three groups of mouse breast tumor assembled by HIF1α staining,
100 times of simulation was processed through downsampling 42 cells from
HIF1α low and HIF1α median group to re-calculate TF variabilities.
834
k-mer analysis: We performed an unbiased analysis of nucleotide sequence
features necessary for chromatin-accessibility variation by searching for an
enrichment of short nucleotide sequences of a specific length k in accessible
regions using ChromVAR. This k-mer analysis allows discovery of novel motifs
and is not restricted by any database entry. Because most TF have core motifs
between 5 and 8 bp, we use 6-, 7- and 8 k-mer analysis to find enrichment of
motifs of 8 or less base pairs. K-mers with significant variability were
assembled and searched against multiple motif databases by Tomtom[42] for
either similarity to known motifs or definition as a *de novo* motif.
844
845

846 **Supplementary information:** Supplementary Figure 1-12, Supplementary Table
847 1-7.
848
Acknowledgements

Supported by NIH P50-HG007735 (to H.Y.C. and W.J.G.), R35-CA209919
(H.Y.C.), Parker Institute for Cancer Immunotherapy (to H.Y.C), Swedish
Research Council International Postdoctoral Fellowship (VR-2016-06794 to
X.C.), and King Abdulaziz University (Ha.C, H.Y.C). We thank S. Kim (Stanford)
for FACS access. The sequencing data was generated on an Illumina HiSeq
4000 that was supported by NIH award S10OD018220. H.Y.C. is an Investigator
of the Howard Hughes Medical Institute.

857
Author contributions
X.C, U.M.L, W.J.G. and H.Y.C conceived and designed the study. X.C, U.M.L
and E.L.L performed experiments. Y.W designed statistical analysis and wrote
the scripts. X.C, U.M.L and Y.W performed data preprocess. Y.W, X.C and U.M.L
performed data analysis with the scripts. A.N.S performed data analysis for
Figure1d, 1e. E.L.L. H.C., and A.J.G. assisted with hypoxia experiments. X.C,
U.M.L, Y.W and H.Y.C wrote the manuscript with input from all authors. H.Y.C.
supervised all aspects of this work.
866
Competing Financial Interests statement:
H.Y.C. is affiliated with Accent Therapeutics (Founder, SAB), Epinomics
(Founder, SAB), Spring Discovery (SAB), 10X Genomics (consultant). W.J.G. is
affiliated with Epinomics (Founder, SAB), 10X Genomics (consultant), and
Guardant Health (consultant). Stanford University holds a patent on ATAC-seq
on which H.Y.C. and W.J.G. are inventors.
873
874 **Data availability:** raw and processed data available at NCBI Gene Expression
875 Omnius, accession number: GSE112091.
876
Correspondence and requests for materials should be addressed to U.M.L.
(litzenbu@stanford.edu) and H.Y.C. (howchang@stanford.edu)

Reference:

1.   Patel, A.P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in
primary glioblastoma. *Science* **344**, 1396-401 (2014).
2.   Lake, B.B. *et al.* Integrative single-cell analysis of transcriptional and
epigenetic states in the human adult brain. *Nat Biotechnol* **36**, 70-80 (2018).
3.   Buenostro, J.D. *et al.* Single-cell chromatin accessibility reveals principles of
regulatory variation. *Nature* **523**, 486-U264 (2015).
4.   Cusanovich, D.A. *et al.* Multiplex single cell profiling of chromatin accessibility
by combinatorial cellular indexing. *Science* **348**, 910-4 (2015).
5.   Jin, W. *et al.* Genome-wide detection of DNase I hypersensitive sites in single
cells and FFPE tissue samples. *Nature* **528**, 142-6 (2015).
6.   Long, W. *et al.* SRC-3Delta4 mediates the interaction of EGFR with FAK to
promote cell migration. *Mol Cell* **37**, 321-32 (2010).
7.   Buenrostro, J.D. *et al.* Integrated Single-Cell Analysis Maps the Continuous
Regulatory Landscape of Human Hematopoietic Differentiation. *Cell* **173**,
1535-1548 e16 (2018).
8.   Corces, M.R. *et al.* Lineage-specific and single-cell chromatin accessibility
charts human hematopoiesis and leukemia evolution. *Nat Genet* **48**, 1193-
203 (2016).
9.   Fagerberg, L. *et al.* Mapping the subcellular protein distribution in three
human cell lines. *J Proteome Res* **10**, 3766-77 (2011).
10. Chen, X. *et al.* ATAC-see reveals the accessible genome by transposase-
mediated imaging and sequencing. *Nat Methods* **13**, 1013-1020 (2016).
11. Wilson, I.G. Inhibition and facilitation of nucleic acid amplification. *Appl
Environ Microbiol* **63**, 3741-51 (1997).
12. Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data.
Nat Methods **14**, 975-978 (2017).
13. Lozzio, B.B., Lozzio, C.B., Bamberger, E.G. & Feliu, A.S. A multipotential
leukemia cell line (K-562) of human origin. *Proc Soc Exp Biol Med* **166**, 546-
50 (1981).
14. Goode, D.K. *et al.* Dynamic Gene Regulatory Networks Drive Hematopoietic
Specification and Differentiation. *Dev Cell* **36**, 572-87 (2016).
15. Wang, Y. *et al.* Metal-responsive transcription factor-1 (MTF-1) is essential
for embryonic liver development and heavy metal detoxification in the adult
liver. *FASEB J* **18**, 1071-9 (2004).
16. Litzenburger, U.M. *et al.* Single-cell epigenomic variability reveals functional
cancer heterogeneity. *Genome Biol* **18**, 15 (2017).
17. Kosan, C. & Godmann, M. Genetic and Epigenetic Mechanisms That Maintain
Hematopoietic Stem Cell Function. *Stem Cells Int* **2016**, 5178965 (2016).
18. Konishi, Y. *et al.* GOOSECOID inhibits erythrocyte differentiation by
competing with Rb for PU.1 binding in murine cells. *Oncogene* **18**, 6795-805
(1999).
19. Zhang, J. *et al.* Deregulation of DUX4 and ERG in acute lymphoblastic
leukemia. *Nat Genet* **48**, 1481-1489 (2016).

20. Hasemann, M.S. *et al.* C/EBPalpha is required for long-term self-renewal and
lineage priming of hematopoietic stem cells and for the maintenance of
epigenetic configurations in multipotent progenitors. *PLoS Genet* **10**,
e1004079 (2014).
21. Suh, H.C. *et al.* C/EBPalpha determines hematopoietic cell fate in
multipotential progenitor cells by inhibiting erythroid differentiation and
inducing myeloid differentiation. *Blood* **107**, 4308-16 (2006).
22. Wu, W. *et al.* Dynamic shifts in occupancy by TAL1 are guided by GATA
factors and drive large-scale reprogramming of gene expression during
hematopoiesis. *Genome Res* **24**, 1945-62 (2014).
23. Peterson, K.R. *et al.* Induction of Fetal Hemoglobin by Transcriptional Co-
Activators MTF-1 and TSPYL1. *Blood* **118**, 353 (2011).
24. Huang, J. *et al.* Dynamic Control of Enhancer Repertoires Drives Lineage and
Stage-Specific Transcription during Hematopoiesis. *Dev Cell* **36**, 9-23 (2016).
25. Guy, C.T., Cardiff, R.D. & Muller, W.J. Induction of mammary tumors by
expression of polyomavirus middle T oncogene: a transgenic mouse model
for metastatic disease. *Mol Cell Biol* **12**, 954-61 (1992).
26. Lee, J.W., Bae, S.H., Jeong, J.W., Kim, S.H. & Kim, K.W. Hypoxia-inducible factor
(HIF-1)alpha: its protein stability and biological functions. *Exp Mol Med* **36**,
1-12 (2004).
27. Sermeus, A. & Michiels, C. Reciprocal influence of the p53 and the hypoxic
pathways. *Cell Death Dis* **2**, e164 (2011).
28. Brabletz, T., Kalluri, R., Nieto, M.A. & Weinberg, R.A. EMT in cancer. *Nat Rev*
Cancer **18**, 128-134 (2018).
29. Werner, S. *et al.* Dual roles of the transcription factor grainyhead-like 2
(GRHL2) in breast cancer. *J Biol Chem* **288**, 22993-3008 (2013).
30. Aprelikova, O., Wood, M., Tackett, S., Chandramouli, G.V. & Barrett, J.C. Role of
ETS transcription factors in the hypoxia-inducible factor-2 target gene
selection. *Cancer Res* **66**, 5641-7 (2006).
31. Gao, Y. *et al.* E2F3 upregulation promotes tumor malignancy through the
transcriptional activation of HIF-2alpha in clear cell renal cell carcinoma.
Oncotarget **8**, 54021-54036 (2017).
32. Patel, N. *et al.* The transcription factor ATOH8 is regulated by erythropoietic
activity and regulates HAMP transcription and cellular pSMAD1,5,8 levels. *Br*
J Haematol **164**, 586-96 (2014).
33. Ben-Batalla, I. *et al.* Deregulation of the Pit-1 transcription factor in human
breast cancer cells promotes tumor growth and metastasis. *J Clin Invest* **120**,
4289-302 (2010).
34. Runa, F. *et al.* Tumor microenvironment heterogeneity: challenges and
opportunities. *Curr Mol Biol Rep* **3**, 218-229 (2017).
35. Rodriguez-Ubreva, J., Garcia-Gomez, A. & Ballestar, E. Epigenetic mechanisms
of myeloid differentiation in the tumor microenvironment. *Curr Opin*
Pharmacol **35**, 20-29 (2017).
36. Easwaran, H., Tsai, H.C. & Baylin, S.B. Cancer epigenetics: tumor
heterogeneity, plasticity of stem-like states, and drug resistance. *Mol Cell* **54**,
716-27 (2014).

37. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. & Greenleaf, W.J.
Transposition of native chromatin for fast and sensitive epigenomic profiling
of open chromatin, DNA-binding proteins and nucleosome position. *Nat*
Methods **10**, 1213-8 (2013).
38. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. & Greenleaf, W.J.
Transposition of native chromatin for fast and sensitive epigenomic profiling
of open chromatin, DNA-binding proteins and nucleosome position. *Nature*
Methods **10**, 1213-+ (2013).
39. Gagnon, K.T., Li, L., Janowski, B.A. & Corey, D.R. Analysis of nuclear RNA
interference in human cells by subcellular fractionation and Argonaute
loading. *Nat Protoc* **9**, 2045-60 (2014).
40. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S.L. Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. *Genome*
Biol **10**, R25 (2009).
41. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**,
R137 (2008).
42. Gupta, S., Stamatoyannopoulos, J.A., Bailey, T.L. & Noble, W.S. Quantifying
similarity between motifs. *Genome Biol* **8**, R24 (2007).
988
989

990   **Figure Legends**
991
Figure 1: **Principle of Protein-index single cell ATAC-seq (Pi-ATAC)**
a, The workflow of Pi-ATAC.
b, The precision of single cell FACS-sorting demonstrated by aligned fragment
comparison of mouse (mESC, blue) and human (GM12878, red) 1:1 cell mixture,
using a 96% species and 500 fragment cutoff.
c, UCSC genome browser track comparison of aggregated 298 single cell Pi-
ATAC to bulk ATAC-seq (both GM12878).
d, Pearson correlation of fragment counts in ATAC-seq peaks in bulk GM12878
ATAC-seq data compared to aggregated single cells from indicated single cell
ATAC-seq approaches and mimic single cells from bulk (shown are distribution of
1,000 times simulation on down sampling 500 fragments from each randomly
selected single cell or bulk sample).
1004
1005
1006
Figure 2: **Pi-ATAC precisely dissects the cell types based on epigenetic**
profile.
a, FACS sorting gating strategy and data of EpCAM+ and CD45+ cells from
mixture of 4T1 cells and mouse splenocytes; the histogram of protein staining is
presented in Supplementary Fig. 4a.
b, Genome browser tracks of aggregated Pi-ATAC 4T1 (EpCAM+) (n=95) and
splenocytes (CD45+) (n=95) at the *Epcam* and *Cd45* loci.
c, t-SNE projection generated from TF deviation z-scores showing the projection
of chromatin accessibility from Pi-ATAC 4T1 (EpCAM+) (n=95) and splenocytes
(CD45+) (n=95).
d, Heatmap of unsupervised hierarchical clustering of the 30 top variable TF
deviations from 4T1 (EpCAM+) (n=95) and splenocytes (CD45+) (n=95) Pi-
ATAC. Each column represents a cell and each row a transcription motif. The
staining cluster information from FACS was assigned to each individual cell (top
color bar).
1022
1023
Figure 3: **Pi-ATAC dissects EpCAM positive tumor cells and Tumor-**
infiltrating immune cells from the same mouse breast tumor
a, schematic illustrating the different cell types in the mouse breast tumor.
b, t-SNE projection of TF deviation z-scores of Pi-ATAC EpCAM+ (n=177) and
CD45+ (n=192) cells from the same mouse breast tumor.

c, Unsupervised hierarchical clustering of the TF deviation z-scores of all 84
significant variable TFs (p < 0.05 after Benjamini-Hochberg correction on multiple
tests) across EpCAM+ (n=177) and CD45+ (n=192) cells from the same mouse
breast tumor. Each column represents one cell and each row a transcription
factor motif. Motif modules (m1-3) and cell subgroups (s1-7) are marked with
distinguished colors. In addition, the staining cluster information from FACS was
assigned to each individual cell (top color bar).
d, t-SNE projection of TF deviations of 84 significant variable TF motifs of
EpCAM+ and CD45+ cells isolated from a tumor. **I,** color coded by cellular
subgroup information; **II-IV**, color coded by the accessibility of the TF motif with
most significant variability in each module (see Figure 3d): Ets2 in m3 (**II**),
Smarcc1 in m2 (**III**) and Hif in m1 (**IV**). Motifs are based on PWM from CisBP
database. Red is highly accessible, blue is low accessible; **V**, color coded by
immune-phenotype.
e, Scatter plot of TF motif variability calculated by ChromVAR across 4T1-
Splenocyte mixture to TF variability calculated by ChromVAR across EpCAM+ -
CD45+ primary tumor cells. Colors indicate TF motif modules as in (c). Arrow
points to Hif.
1047
1048
1049
Figure 4: **Epigenetic variability is modulated by the hypoxic**
microenvironment
a, FACS sorted EpCAM+ and HIF1α+ double positive cells from the same mouse
breast tumor; three groups of HIF1 α staining were assigned, HIF1α negative
(n=762), HIF1α low (n=139), HIF1α high (n=55) (right plot).
b, Ranking of transcription factor motif variability of different HIF1α staining
groups, HIF1α negative (n=702), HIF1α low (n=95), HIF1α high (n=42).
c, Examples of TF motif deviations which significantly change in HIF1α high
group compared to the others. In each comparison, HIF1α negative (n=702),
HIF1α low (n=95), HIF1α high (n=42). Significance of Wilcoxon test shown with
n.s. (not significant), * (p<0.05) and ** (p<0.01).
d, Flow cytometry analysis of HIF1α protein abundance in 4T1 breast cancer
cells cultured in 5% oxygen (Ctrl, gray) or 1% oxygen for indicated time points.
e, Chromatin accessibility changes of hypoxic conditions compared to control.
Red indicates higher, blue lower accessibility.
f, TF deviation examples of motifs whose accessibility increases in hypoxia over
time in 4T1 breast cancer cells (TF deviation was calculated using ChromVAR)**.**

Chang_Figure 1

Chang_Figure 2

Chang_Figure 3