Title: Precision Identification of Diverse Bloodstream Pathogens from the Gut Microbiome

Authors: Fiona B. Tamburini, BS[1]*; Tessa M. Andermann, MD, MPH[2]*; Ekaterina Tkatchenko, BS[3];

Fiona Senchyna[4], BS; Niaz Banaei[2,4], MD; Ami S. Bhatt, MD, PhD[1,3]**

1 Department of Genetics, Stanford University

2 Department of Medicine, Division of Infectious Diseases, Stanford University

3 Department of Medicine, Division of Hematology, Stanford University

4 Clinical Microbiology Laboratory, Stanford University Medical Center

* These authors contributed equally

**To whom correspondence should be addressed: asbhatt@stanford.edu

Abstract

Bloodstream infection is the most common infectious complication in hematopoietic cell transplantation

recipients. To evaluate the genomic concordance of bloodstream pathogens and bacterial strains within

the intestinal microbiome using whole genome sequencing, we developed StrainSifter, a bioinformatic

pipeline to compare nucleotide variation between bacterial isolate strains and stool metagenomes. We

applied StrainSifter to bloodstream isolates and stool metagenome samples from hematopoietic stem cell

transplant recipients with bloodstream infections. StrainSifter is designed to identify single nucleotide

variants between isolate and metagenomic short reads using stringent alignment, coverage, and variant

frequency criteria for strain comparison. We identified enteric BSI isolates that were highly concordant

with those in the gut microbiota, as well as highly concordant strains of typically non-enteric bacteria.

These findings demonstrate the utility of StrainSifter in strain matching and provide a more precise

investigation of the intestine as a reservoir of diverse pathogens capable of causing bloodstream

infections.

Introduction

Microbial pathogenicity and transmission depend in part on strain-level variability in both gene content and at the single-nucleotide variant (SNV) level, as different strains of the same organism can vary widely in their ability to cause disease (Snitkin et al. 2011; Lieberman et al. 2013). Whole genome sequencing (WGS) has facilitated the exploration of strain-level determinants of virulence and has enabled the precise tracking of pathogen transmission (Snitkin et al. 2011; Snitkin et al. 2012). Methods for strain tracking and comparison most commonly rely on either sequencing a defined number of genes or identifying single nucleotide variants (SNVs) across the whole genome. For example, whole genome sequencing of cultured *Klebsiella pneumoniae* stool isolates in ICU patients demonstrated the role of *K. pneumoniae* colonization in systemic infections (Gorrie et al. 2017; Martin et al. 2016; Nielsen et al. 2017) and genome-wide SNV comparison is routinely employed for epidemiologic strain tracking (Salipante et al. 2015; Snitkin et al. 2012; Harris et al. 2013; Chen et al. 2017; Kovanen et al. 2014). While strain comparisons have primarily been performed on bacterial isolates, newer computational tools (MIDAS, StrainPhlAn, and metaSNV) profile strain variation between metagenomic samples over time without the bias inherent in culturing isolates (Costea et al. 2017; Nayfach et al. 2016; Truong et al. 2017).

Despite the increasing use of WGS for strain comparison, bioinformatic tools have not been developed for the specific purpose of comparing disease-causing bacterial isolates to metagenomic samples at the nucleotide level within the same patient. It is desirable to achieve the highest resolution of comparison across the whole genome of an organism in order to precisely compare clinical isolates to strains within a sampled metagenome. Additionally, most tools depend on the availability of reference genomes and are thus often restricted to studying well characterized bacterial pathogens, limiting the types of organisms that are available for strain comparisons. To date, no reference-free tool exists with the stringent filtering steps necessary to identify genomic variants specific to a single strain between genomes and metagenomes. We sought to compare strain-level identity between bloodstream infection isolates and stool metagenomes using samples from immunocompromised patients at elevated risk of systemic

infections, with the goal of better understanding the origins of bloodstream infection in this patient

population.

Hematopoietic stem cell transplant (HCT) recipients are at high risk for bloodstream infections, which

occur in 20-60% of patients primarily during periods of neutropenia and mucositis (Blennow et al. 2014;

Gudiol et al. 2014; Mikulska et al. 2009). Prior to engraftment of donor stem cells, HCT recipients are

particularly susceptible to translocation of enteric organisms from the intestinal microbial reservoir into

the bloodstream. During periods of mucositis, primary bloodstream infections with enteric Gram-negative

and Gram-positive organisms are thought to arise from mucosal barrier injury (See et al. 2017). In

contrast, non-enteric commensal and environmental bacteria access the bloodstream through other routes

including intravenous lines and sites where skin epithelial integrity has become compromised. Existing

methods for strain comparison have been helpful in identifying the origins of bloodstream infections in

HCT patients using culture-based pulsed-field gel electrophoresis (PFGE), or PCR-based multi-locus

sequence typing (MLST) (Tancrede and Andremont 1985; Samet et al. 2013). Although these methods

are rapid, affordable, and standardized across many organisms, they have several limitations: 1) they

require isolation and culture of candidate pathogens from the stool or other suspected pathogen reservoirs,

a step that can introduce selection bias, and 2) they often rely on detailed characterization of a small

number of genes and therefore lack the discriminatory power required for robust, genome-wide strain

comparison (W. Li, Raoult, and Fournier 2009). No studies to date have investigated the origins of

bloodstream infections in HCT or other immunocompromised patients at the level of single nucleotide

variation across entire genomes.

In this work, we present StrainSifter, a straightforward bioinformatics pipeline based on whole genome

sequencing to comprehensively and precisely compare single nucleotide variants between isolate

genomes and metagenomes without bias introduced by culturing. Additionally, unlike most previous

strain comparison bioinformatic tools (MIDAS and StrainPhlAn), variant calling within our pipeline does

not depend on the availability of a reference genome database, a particularly relevant consideration in

immunocompromised patients who develop infections with lesser known and poorly studied opportunistic

pathogens. We have applied this tool to track bacterial strain translocation from the gut to the bloodstream

in a unique cohort of immunocompromised patients at high risk for translocation and bloodstream

infection, specifically patients with hematologic malignancies who have undergone hematopoietic stem

cell transplantation. We find that StrainSifter is able to identify strains of both intestinal mucosa-

associated (*Enterococcus faecium*, *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter cloacae, and*

Streptococcus mitis) and typically non-mucosal bacteria (*Staphylococcus epidermidis* and *Pseudomonas*

aeruginosa) within the gut microbiome as being concordant with the same strain cultured from the blood

of patients with bloodstream infections. Intriguingly, this suggests that both enteric and classically non-

enteric pathogens can colonize and even thrive within the intestinal reservoir of immunocompromised

hosts.

Methods

Cohort selection

A retrospective cohort study, approved by the institutional review board under the IRB protocol #8903

(PI: Dr. David Miklos, co-PI: Dr. Ami Bhatt), was performed at Stanford Hospital. Informed consent was

obtained from all subjects prior to specimen collection. At the time of cohort identification (July 2017), a

stool biospecimen collection containing 964 stool samples from 402 patients was available for

investigation. This collection consisted of convenience samples collected from autologous and allogeneic

stem cell transplantation patients at Stanford University Hospital between October 5 2015 and June 9

2017. Patients were included in this study if a stool sample had been collected within 30 days prior to an

episode of bloodstream infection (BSI) for which a blood isolate was also available. From this final

cohort, we sequenced all stool samples in our collection within 60 days prior to and 30 days after BSI.

Bloodstream isolate identification

Bloodstream isolates from HCT patients who received medical care at Stanford University Hospital were

obtained from the Stanford Hospital Clinical Microbiology Laboratory. All isolates considered typical

bloodstream pathogens by National Healthcare Safety Network (NHSN) guidelines are stored in a

glycerol suspension at -80°C for 12 months ("Centers for Disease Control and Prevention" 2018). Blood

culture isolates considered to be skin-associated bacteria (including viridans group *Streptococcus spp.* and

coagulase-negative *Staphylococcus spp.*) were saved if they were recovered in two or more blood culture

sets as per NHSN criteria ("Centers for Disease Control and Prevention" 2018). Isolates were identified

by standard biochemical testing and matrix-assisted laser desorption and ionization time-of-flight mass

spectrometry (MALDI-TOF) (Bruker Daltonics, Bremen, Germany).

Sample processing

Bloodstream bacterial isolates were plated on brain heart infusion agar with 10% horse blood. DNA was

extracted from isolates using the Gentra Puregene Yeast/Bact. Kit per manufacturer's instructions.   Stool

samples were collected and stored at 4°C for up to 24 hours prior to homogenization, aliquoting and

storage at 80°C. DNA was extracted from stool using the QIAamp DNA Stool Mini Kit (QIAGEN®) per

manufacturer's instructions, with an initial bead-beating step prior to extraction using the Mini-

Beadbeater-16 (BioSpec Products) and 1 mm diameter Zirconia/Silica beads (BioSpec Products). Bead-

beating consisted of 7 rounds of alternating 30 second bead-beating bursts followed by 30 seconds

cooling on ice. DNA concentration for all samples was measured using Qubit® Fluorometric Quantitation

(Life Technologies). DNA sequencing libraries from both isolates and stool were prepared using the

Nextera XT DNA Library Prep Kit (Illumina®) with isolates and stool microbiota libraries prepared at

separate times following DNAse treatment of all lab surfaces to avoid cross-contamination. Library

concentration was measured using Qubit® Fluorometric Quantitation (Life Technologies) and library

quality and size distributions were analyzed with the Bioanalyzer 2100 (Agilent). Prepared libraries were

multiplexed and subjected to 100 base pair, paired-end sequencing on the HiSeq 4000 platform

(Illumina®).

Shotgun sequencing analysis

Preprocessing

Sequencing data were demultiplexed by unique barcodes (bcl2fastq v2.20.0.422, Illumina®). Read quality was assessed using FastQC v0.11.4. Reads were deduplicated to remove PCR and optical duplicates using SuperDeduper v1.4 with the start location in the read at 5 bp (-s 5) and minimum length of 50 bp (-l 50). Deduplicated reads were trimmed using TrimGalore v0.4.4, a wrapper for CutAdapt v1.16, with a minimum quality score of 30 for trimming (-q 30), minimum read length of 50 (--length 50) and the "--nextera" flag to remove Illumina Nextera adapter sequences. Draft genomes of bacterial isolates were assembled using SPAdes v3.8.0 (Bankevich et al. 2012) with default parameters. Summary statistics for each BSI assembly were generated using 'basic_assembly_stats.py' from GAEMR v1.0.1 ("GAEMR" n.d.).

Taxonomic classification

Bloodstream isolates were taxonomically classified by aligning draft genome contigs to the BLAST nt database and identifying the most common top hit for each contig. Gut metagenomic reads were taxonomically classified via the One Codex platform, a web-based tool for assigning read-level classifications based on unique *k*-mer signatures relative to a curated reference database [2017 database] (Minot, Krumm, and Greenfield 2015).

Phylogenetic tree building and variant identification with the StrainSifter pipeline

StrainSifter is a pipeline deployed via Snakemake with modules for variant calling and phylogenetic tree building, available at GitHub (https://github.com/bhattlab/strainsifter). StrainSifter accepts as input an assembled bacterial draft genome and two or more short read data sets (isolate or metagenomic), and can report a phylogenetic tree of input samples as well as pairwise SNV counts. To build the phylogenetic trees reported in this manuscript, an NCBI RefSeq reference genome for each infectious species (based on

clinical laboratory classifications) was provided to StrainSifter. For the variant counting reported herein,

BSI draft genomes were supplied to StrainSifter. For both analyses, all stool and BSI short read datasets

were provided as input. For both phylogeny and SNV-counting modules, preprocessed short reads are

first aligned to the reference genome using the Burrows-Wheeler aligner (BWA) v0.7.10 (Li and Durbin

2009). Alignments are filtered to include only high-confidence alignments with mapping quality of at

least 60 using the "view" tool from the SAMtools suite (v1.7)(Li et al. 2009) (samtools view -b -q 60),

and further filtered using BamTools "filter" (v2.4.0) to include only reads with the desired number or

fewer mismatches (i.e. for five or fewer mismatches: bamtools filter -tag 'NM:<6') (Barnett et al. 2011)

For phylogenetic tree construction, reads with 5 or fewer mismatches were included; for determining

strain single-nucleotide variants, reads were limited to 1 or fewer mismatches. Per-base coverage is

calculated from each resulting BAM file using bedtools genomecov (v2.26.0) and processed with custom

python scripts to identify samples meeting a minimum average coverage of 5X across at least 50% of the

genome. Only samples meeting the coverage requirement are continued through the pipeline. Pileup files

are created from BAM files using SAMtools "mpileup", and are analyzed using custom python scripts to

identify bases occurring with at least 0.8 frequency at positions covered 5X or greater. Only bases with a

minimum phred score of >=20 are considered. Consensus sequences for each sample are created, wherein

bases that cannot be confidently determined given the described parameters are called as "N".

To create a phylogenetic tree, core positions are identified on a per-species basis, where core positions are

defined as positions in the reference genome where a base could be confidently called for all samples

analyzed. To generate phylogenetic trees, core positions with variants in at least one sample are identified

and concatenated into fasta files. Fasta files are aligned using MUSCLE v3.8.31 (Edgar 2004) and a

maximum-likelihood phylogenetic tree is computed using FastTree v2.1.7 (Price, Dehal, and Arkin 2010).

Phylogenetic trees are visualized in R using the ape, phangorn, and ggtree packages. Pairwise SNVs are

determined from the consensus sequences using a custom python script.

Synthetic multilocus sequence typing

Metagenomic short reads were assembled using metaSPAdes v3.8.0 (Nurk et al. 2017). Multilocus

sequence typing (MLST) schemes and sequences were downloaded from the PubMLST database

("PubMLST" n.d.). MLST gene sequences were aligned to metagenome assemblies, and the top hit for

each alignment was chosen based on E-value, percent identity, and alignment length. Only MLST

sequences that were present in the metagenomic assembly with 100% identity across the entire length of

the sequence were reported. MLST types generated by our in-house analysis were confirmed with the

SRST2 synthetic multilocus sequence typing tool (Inouye et al. 2014).

Plots

Plots were generated using the R programming language using the ggplot2, reshape2, and dplyr packages.

<u>Results</u>

Sample collection and cohort

During the study period, we identified 74 patients within our stool biospecimen collection with a

bloodstream infection (77 BSI episodes). Of these, we identified 30 patients (32 bloodstream isolates)

with at least 1 collected stool sample within 30 days prior to their bacterial BSI. From our final cohort of

30, we sequenced all stool samples collected between 60 days prior to and 30 days after the date of BSI,

83 samples in total. Patients had a median of 2 stool samples (range 1-8), with the sampling distribution

presented in Figure 1. Stools were collected at a median of 9 days prior to BSI (range -58 to +31).

Clinical characteristics of the 30 patients are listed in Table 1 (see Table S1 for individual patient data).

All patients in the cohort were on at least one antibiotic within 30 days prior to BSI, with

fluoroquinolones being the most commonly administered antibiotic.

Table 1.

Baseline characteristics	N (%)
Age (years)	
<30	4 (13%)
≤30-39	4 (13%)
≤40-49	6 (20%)
≤50-59	3 (10%)
≤60	11 (37%)
Sex (%male)	17 (57%)
Underlying diagnosis	
AML	7 (23%)
ALL	6 (20%)
CML	2 (7%)
Lymphoma	7 (23%)
MDS/Myelofibrosis	6 (20%)
Other	2 (7%)
Conditioning regimen	
Myeloablative	16 (53%)
Reduced intensity	9 (30%)
Non-myeloablative	5 (17%)
Transplant source	
Peripheral blood	20 (67%)
Bone marrow	8 (27%)
Double umbilical cord blood	2 (7%)
Type of donor	
Autologous	3 (10%)
Allogeneic	27 (90%)
Matched related donor	10 (37%)

	Matched unrelated donor	14 (52%)
	Mismatched unrelated donor	3 (11%)
TPN within 30 days		15 (50%)
Antibiotics within 30 days*		30 (100%)
	Fluoroquinolones	26 (87%)
	Beta-lactam	14 (47%)
	Carbapenem	6 (20%)
	Vancomycin (IV)	12 (40%)
Bacteremia species		
Gram-positive	*Staphylococcus aureus*	8 (25%)
	MSSA	5 (16%)
	MRSA	3 (9%)
	Coagulase-negative Staphylococcus spp.	4 (13%)
	Streptococcus mitis	7 (22%)
	Enterococcus faecium	2 (6%)
	Rothia mucilaginosa	2 (6%)
Gram-negative	*Escherichia coli*	4 (13%)
	Klebsiella spp.	3 (9%)
	Klebsiella pneumoniae	2 (6%)
	Klebsiella oxytoca	1 (3%)
	Enterobacter cloacae	1 (3%)
	Pseudomonas aeruginosa	1 (3%)

List of abbreviations: AML (acute myeloid leukemia); ALL (acute lymphoblastic leukemia); CML (chronic

myeloid leukemia).

*Antibiotics are not exclusive and will not add up to 100%

Figure 1. Stool sampling

Timeline of stool sampling relative to BSI and hematopoietic cell transplant for patients included in the

study.

Date of stool sampling relative to bacteremia

Infectious taxa are present in the gut microbiome near the time of infection

We sequenced an average of 68,704,214 reads per stool sample (20,298,792-148,162,892) and 27,999,060

reads per isolate (10,652,310-61,058,906) (Table S2). Isolate reads were assembled into draft genomes

(Table S3).

MALDI-TOF is unable to discriminate between some closely related taxa (e.g. *Streptococcus* spp.), so we

sought to confirm the clinical laboratory BSI classifications by aligning draft genome contigs to the

BLAST nt database. Of 32 BSIs, we identified 3 for which WGS and MALDI-TOF results disagreed.

Two *Streptococcus* isolates that had been classified as *Streptococcus mitis* by MALDI-TOF were

identified as *Streptococcus oralis,* and one as *Streptococcus pseudopneumoniae* by WGS (Table S4).

To determine community composition of pre- and post-BSI stool samples, we taxonomically classified

gut metagenomic reads via the One Codex platform. Considering species level read classifications, we

observe the target species in the gut at a threshold of 0.1% or greater relative abundance for 19 of 83

stools (23%) or 20 of 95 stool-BSI pairs (21%) taking into account episodes of polymicrobial BSI; one

patient developed a BSI with two bacterial species, both of which were present in the stool above

threshold level (Table 2; full taxonomic classifications in table S5). Read-level taxonomic classifiers are

often unable to discriminate between closely related species, so we also compared BSIs and stool read

classifications at the genus level. We observe the target genus in the gut community for 27% of stools, or

23% of stool-BSI pairs (Table S5, S6).

Table 2.

Patient	Day relative to BSI	BSI organism	Relative abundance (%)
1	-4	Staphylococcus aureus	0
1	3	Staphylococcus aureus	0
2	-7	Enterobacter cloacae	0
2	0	Enterobacter cloacae	0.74
2	7	Enterobacter cloacae	0
2	-7	Klebsiella pneumoniae	0
2	0	Klebsiella pneumoniae	3.47
2	7	Klebsiella pneumoniae	0.01
3	-27	Escherichia coli	87.15
3	-32	Escherichia coli	97.25
3	-33	Escherichia coli	96.4
3	-40	Escherichia coli	0.05

3	-41	Escherichia coli	0.43
3	8	Escherichia coli	0.01
3	-27	Staphylococcus aureus	0
3	-32	Staphylococcus aureus	0
3	-33	Staphylococcus aureus	0
3	-40	Staphylococcus aureus	0
3	-41	Staphylococcus aureus	0
3	8	Staphylococcus aureus	0
4	-24	Staphylococcus aureus	0
4	-30	Staphylococcus aureus	0
4	-38	Staphylococcus aureus	0
4	-43	Staphylococcus aureus	0
4	-52	Staphylococcus aureus	0
4	-58	Staphylococcus aureus	0
4	25	Staphylococcus aureus	0
4	4	Staphylococcus aureus	0
5	-7	Staphylococcus aureus	0
5	7	Staphylococcus aureus	0
6	-28	Klebsiella oxytoca	0
7	-16	Escherichia coli	0.79
7	-30	Escherichia coli	0.22
7	-51	Escherichia coli	2.05
7	-9	Escherichia coli	0.38
8	-7	Rothia mucilaginosa	0
8	0	Rothia mucilaginosa	0
9	-11	Escherichia coli	0.31
9	-18	Escherichia coli	0
9	-32	Escherichia coli	0
9	-4	Escherichia coli	0.03
9	4	Escherichia coli	0
10	-3	Staphylococcus aureus	0

11	-35	Escherichia coli	0.04
12	-14	Staphylococcus aureus	0
12	-21	Staphylococcus aureus	0
12	-35	Staphylococcus aureus	0.65
12	-42	Staphylococcus aureus	0.09
13	-1	Staphylococcus epidermidis	2.51
14	-9	Klebsiella pneumoniae	0
14	19	Klebsiella pneumoniae	0
14	26	Klebsiella pneumoniae	0
15	-12	Staphylococcus epidermidis	0
15	-12	Staphylococcus epidermidis	0
15	-33	Staphylococcus epidermidis	0
15	-33	Staphylococcus epidermidis	0
16	-10	Streptococcus mitis	0
16	-3	Streptococcus mitis	0
17	-11	Streptococcus mitis	0
17	-12	Streptococcus mitis	0
17	-5	Streptococcus mitis	0
17	4	Streptococcus mitis	0.03
18	-16	Staphylococcus epidermidis	0
18	-16	Staphylococcus epidermidis	0
19	-7	Pseudomonas aeruginosa	0
19	0	Pseudomonas aeruginosa	0.83
20	-16	Staphylococcus epidermidis	0
21	-7	Staphylococcus aureus	0.01
21	0	Staphylococcus aureus	0.25
22	-15	Streptococcus mitis	0
22	-8	Streptococcus mitis	2.33
23	-10	Enterococcus faecium	11.61
23	6	Enterococcus faecium	0
24	-1	Rothia mucilaginosa	0.04

24	-11	Rothia mucilaginosa	0
24	-13	Rothia mucilaginosa	0
24	-4	Rothia mucilaginosa	0
24	31	Rothia mucilaginosa	0
25	-11	Enterococcus faecium	0.36
25	3	Enterococcus faecium	94.09
26	-11	Streptococcus mitis	0
26	-4	Streptococcus mitis	0
27	0	Staphylococcus aureus	0.51
28	-10	Streptococcus mitis	0
28	-12	Streptococcus mitis	0.04
28	-3	Streptococcus mitis	0
28	-9	Streptococcus mitis	0.02
28	5	Streptococcus mitis	0
29	-12	Streptococcus mitis	0
29	-3	Streptococcus mitis	0
29	-8	Streptococcus mitis	0.01
29	0	Streptococcus mitis	0.03
30	-10	Streptococcus mitis	0
30	-3	Streptococcus mitis	0
30	4	Streptococcus mitis	0

Gut domination with an infectious pathogen does not always precede bloodstream infection

We sought to determine whether bloodstream infection is preceded by gut microbiota domination by a pathogen as has been previously reported (Ubeda et al. 2010; Taur et al. 2012). We defined domination as 30% species relative abundance or greater of an organism in the gut, as previously described (Taur et al. 2012). Of the 19 stools in which the BSI organism was detected, only 4 stool samples (21%) were dominated by the BSI pathogen. Stool samples from patient 3 were dominated by *Escherichia coli* (>85% relative abundance) 27, 32, and 33 days prior to the development of an *E. coli* BSI (Figure 2A; table 2);

Enterococcus faecium dominated the intestinal microbiome (94% relative abundance) in patient 25 three

days after *E. faecium* BSI (Figure 2B; table 2).

Several patients experienced BSI with organisms present at relatively lower abundance in the gut prior to

bloodstream infection: *Staphylococcus epidermidis* at approximately 2.5% relative abundance in the gut

of patient 13 one day prior to their *S. epidermidis* infection, *Pseudomonas aeruginosa* at low abundance

(0.83%) in the gut of patient 19 on the day of infection, and *Klebsiella pneumoniae* at 3.5% relative

abundance in the gut of patient 2, 9 days prior to BSI (Figure 2C,D; table 2).

We also observe gut domination by non-BSI-causing pathogens. Patient 14 experienced a *K. pneumoniae*

BSI, yet stool samples at two timepoints were dominated by other pathogens: *E. coli* at 67% relative

abundance 9 days prior to BSI and *E. faecium* at 85% relative abundance 19 days after BSI (Table 2).

Figure 2. Gut domination by a pathogen does not always precede bloodstream infection. Domination

by *E. coli* (A) and Enterococcus faecium (B) occurs prior to bacteremia, while *K. pneumoniae* (C) and *S.*

epidermidis (D) are present in the gut microbiota of their respective patients prior to BSI but not at high

abundance. Plots show species present at 2% relative abundance or greater (host reads excluded).

Phylogenetic analysis indicates close relatedness of gut and BSI strains

We developed the StrainSifter pipeline to perform nucleotide-level comparisons between genomes and

metagenomes. StrainSifter is deployed as a Snakemake workflow, and the pipeline steps are described in

Figure 3. Briefly, the inputs to StrainSifter are a draft genome (the "target" genome) and one or more

other short read datasets (isolate genomes or metagenomes). StrainSifter detects whether the target

organism is present with sufficient abundance in the short read datasets and outputs phylogenetic trees

and SNV counts between samples. Despite an abundance of research studies focused on strain-level

comparisons of microbes, there is still no clear definition of what constitutes a "strain," and it is unclear

whether genomes that differ by a few nucleotides should be considered distinct strains. To investigate the

overall relatedness of strains of each BSI species in our metagenomes and isolates, we first took a

comparative phylogenomics approach by applying StrainSifter to assess inter-patient strain relatedness.

Rather than compare strains from the same patient to publicly-available reference genomes, we instead

compared the phylogenetic relatedness of a patient's gut and BSI strains to strains from the other patient

samples in our collection. We reasoned that strains from the same geographic location are more likely to

be closely phylogenetically related than strains that are both spatially and temporally distant, so

comparing inter-patient strain relatedness in the context of the other strains in our sample collection

should more precisely indicate strains or lineages unique to a patient.

Figure 3. StrainSifter pipeline

Flowchart showing input, analysis steps, and output of the StrainSifter pipeline.

For each BSI species, we used StrainSifter to identify metagenomic samples with at least 5X coverage of

the NCBI representative genome for each target organism, found core SNVs present in each sample, and

inferred maximum-likelihood phylogenetic trees from multiple sequence alignments of concatenated core

SNVs. Reference genomes were selected based on the clinical laboratory MALDI-TOF classification of

BSI isolates. Phylogenetic trees demonstrate a higher level of intra-patient versus inter-patient genetic

relatedness among colonizing and infecting organisms (Figure 4). BSI and intestinal strains of typically

enteric species such as *Enterococcus faecium* (patient 25), *Klebsiella pneumoniae* (patient 2), *E. coli*

(patient 3, patient 7), and *S. mitis* (patient 22) are closely phylogenetically related. Unexpectedly, we

observe close intra-patient strain relatedness between gut and BSI for typically non-enteric taxa including

S. epidermidis strains (patient 13) and *Pseudomonas aeruginosa* (patient 19, not pictured). Phylogenetic

trees for *P. aeruginosa* and *Enterobacter cloacae* are not shown as these species were not observed in any

other samples.

Single-nucleotide variants reveal strain identity

We next used StrainSifter to compare single-nucleotide variants (SNVs) between strains in our sample

collection. We executed the StrainSifter pipeline for each draft genome, with all stool short read datasets

as input. StrainSifter identified samples with sufficient coverage (at least 5X coverage across 50% or

more of the genome) of the target organism in the gut metagenome and determined the number of variants

shared between each pair of input samples that contained the target strain.

We observe 1 SNV between the BSI and gut content of Patient 13 who developed a *S. epidermidis* BSI

(Table 3). This indicates that the bloodstream strain is concordant to the strain found in the gut 1 day

prior. Patient 3 developed an *E. coli* bloodstream infection and we observe zero SNVs between the BSI

strain and all samples 33, 32, and 27 days prior to the BSI, indicating that the identical E. coli BSI strain

colonized the gut over a month before the onset of infection. Further, we observe zero discriminating

SNVs between identical strains of *Pseudomonas aeruginosa* in both the blood and stool specimens.

Notably, all stool samples containing the BSI-causing organism at sufficient coverage to profile with

StrainSifter were more closely related to their respective BSI isolate based on SNV distance than to any

sample from other patients. None of the 30 patients included in our study had sufficient *Staphylococcus*

aureus in their stool samples to profile with StrainSifter, indicating that this organism infrequently

colonizes the gut of HCT patients, even at low abundance.

Table 3.

BSI organism	Sample 1	Sample 2	SNVs	Bases compared	SNVs per base compared	Total bases	Percentage of bases compared
Enterobacter cloacae	Patient 2 BSI	Patient 2 stool day 0	5	4499921	1.11E-06	4552154	98.85%
Enterococcus faecium	Patient 25 BSI	Patient 25 stool day 3	2	2931442	6.82E-07	2935264	99.87%
Escherichia coli	Patient 3 BSI	Patient 3 stool day -33	0	5032224	0.00E+00	5056158	99.53%
Escherichia coli	Patient 3 BSI	Patient 3 stool day -32	0	5032230	0	5056158	99.53%

Escherichia coli	Patient 3 BSI	Patient 3 stool day -27	0	5030454	0	5056158	99.49%
Escherichia coli	Patient 7 BSI	Patient 7 stool day -51	259	4388254	5.90E-05	5267890	83.30%
Klebsiella pneumoniae	Patient 2 BSI	Patient 2 stool day 0	4	5379270	7.44E-07	5390737	99.79%
Klebsiella pneumoniae	Patient 14 BSI	Patient 14 stool day -9	72	3924126	1.83E-05	5407718	72.57%
Pseudomonas aeruginosa	Patient 19 BSI	Patient 19 stool day 0	0	6830248	0	6882743	99.24%
Staphylococcus epidermidis	Patient 13 BSI	Patient 13 stool day -1	1	2592411	3.86E-07	2604609	99.53%
Streptococcus mitis	Patient 22 BSI	Patient 22 stool day -8	28	1345797	2.08E-05	2084062	64.58%

325 **Figure 4. Intra-patient samples are more closely related than inter-patient samples.** Branch tip

326 colors indicate stool and BSI samples from the same patient. Samples from the same patient are more

327 closely phylogenetically related than other samples.

Enterococcus faecium

Klebsiella pneumoniae

Staphylococcus epidermidis

Escherichia coli

Staphylococcus aureus

Streptococcus mitis

In silico multilocus sequence typing is concordant with StrainSifter results

To compare whole-genome sequencing-based approaches to traditional strain typing methods, we

performed *in silico* multilocus sequence typing (MLST) on BSI genomes and assembled stool

metagenomes (Table 3). In many instances, MLST types for gut metagenome strains could not be

resolved or could only be partially detected, likely due to low abundance or absence of the target

organism in the gut community. MLST data is concordant with StrainSifter SNV profiling in only 4

infections for which MLST schemes exist for the bloodstream isolate of interest (Table S7). MLST types

could not be determined for either of patient 2's BSI isolates (*K. pneumoniae* and *E. cloacae*) because the

sequences at several loci in each genome were not present in the respective MLST databases. We were

unable to type *Rothia mucilaginosa* and *Streptococcus mitis*, as standard MLST schemes do not exist for

these organisms.

<u>Discussion</u>

StrainSifter is a useful pipeline for strain detection within metagenomes

High resolution strain comparisons are required to identify the origins of bloodstream infections using

metagenomic samples. While many bioinformatic tools exist for strain comparison between clinical

isolates or SNV profiling between metagenomes, no pipeline exists for the specific purpose of detecting

the presence of a single strain in a metagenome. In this study, we have developed StrainSifter, a

straightforward and facile pipeline for identifying phylogenetic relationships between isolate and

metagenomic WGS samples.

In our study, StrainSifter was able to precisely and comprehensively compare gut metagenomic content to

the genomes of diverse bloodstream isolates using high resolution WGS. Although techniques such as *in*

silico MLST are able to compare genomes to metagenomes, MLST loci are only 450 to 500 base pairs on

average, and standard 7-gene MLST schemes therefore analyze less than 4 kilobases of DNA - on the

order of 0.1% of the average bacterial genome. By examining genome-wide variants, StrainSifter greatly

increases the confidence of strain comparisons.

Using StrainSifter, we found that the enteric organisms *E. faecium, E.coli*, *K. pneumoniae*, *E. cloacae*,

and *S. mitis* were concordant between blood and stool microbiota samples. This confirms what has

previously been reported in the literature (Tancrede and Andremont 1985; Samet et al. 2013). Previously

published studies have indicated that gut domination with a pathogen is a risk factor for subsequent

bloodstream infection (Taur et al. 2012; Ubeda et al. 2010; "Centers for Disease Control and Prevention"

2018). While *E. coli* and *E. faecium* domination events were observed preceding BSI, we find *K.*

pneumoniae, *E. cloacae,* and *S. epidermidis* in relatively low abundance in stool samples on the day of

BSI and one day prior.

Interestingly, we also find that typically non-enteric organisms *S. epidermidis* and *P. aeruginosa* were not

only present in the gut, but strains within the gut microbiota and bloodstream were concordant. Gut

colonization by *S. epidermidis* and *P. aeruginosa* is unexpected given current National Health and Safety

guidelines delimiting enteric versus non-enteric organisms (Taur et al. 2012; Ubeda et al. 2010; "Centers

for Disease Control and Prevention" 2018; Advani et al. 2017). Yet, colonization by atypical organisms

may not be surprising given the degree of disruption of the intestinal microbiota by antibiotics,

chemotherapy, and radiation. While *P. aeruginosa* intestinal colonization has been previously

documented (Nesher et al. 2014), intestinal colonization with coagulase-negative *Staphylococcus* is not

well described. In our own study, multiple stool samples were determined to have more than 80% relative

abundance of *S. epidermidis*. In addition, we observe the same colonizing strain of coagulase-negative

Staphylococcus over time in the same individual (patient 7), arguing against only transient membership

within the intestinal microbiome.

Previous research using existing strain typing and culture-based strategies has only tentatively

demonstrated evidence for gut translocation of coagulase-negative *Staphylococcus spp.* and *P. aeruginosa*

in immunocompromised patients (Wade et al. 1982; Rotstein et al. 1988; Nesher et al. 2014; MacFie et al.

1999; Costa et al. 2006). Here our straightforward approach allows for greater precision and confidence in

strain comparisons between the gut microbiota and infecting organism in the bloodstream. Given that our

analysis focused entirely on the stool microbiota, we cannot rule out the possibility of the same strain of

coagulase-negative *Staphylococcus* colonizing the entire body surface--including the gut, nares, skin, or

central line--from which the infection may have originated instead. Previous research has suggested,

however, that the majority of patients with coagulase-negative *Staphylococcus* bloodstream infection are

colonized with a multitude of strains that differ between the intestine and the skin (Herwaldt et al. 1992).

Determining whether two strains are related based on the number of SNVs that differ between their

genomes is not straightforward. While approximate bacterial mutation rates due to genetic drift have been

established, most estimates are derived from experiments in culture media. As gut and bloodstream

environments impose many selective pressures beyond genetic drift, such estimates are likely not relevant

in the clinical context. For instance, it is possible that bacterial pathogens that originate from the intestine

and translocate into the bloodstream may acquire single nucleotide variants over time due to selective

pressures. For this reason, it is difficult to define a threshold number of SNVs beyond which two strains

are not considered to be the same. Regardless of the absolute number of SNVs between strains or across

all strains analyzed with StrainSifter, strains from the same patient consistently share fewer SNVs

compared to those from other patients.

Although StrainSifter can precisely identify shared variants between genomes and metagenomes, it is

limited to profiling only the dominant strain of a given organism in a community. However, it has been

shown that gut metagenomes frequently contain only one predominant strain of each species, so

StrainSifter is likely to function optimally in most situations (Truong et al. 2017). As is the case with

existing tools for metagenomic strain comparison, StrainSifter is unable to analyze low-abundance strains

since adequate coverage (5X) is required for confident variant calling.

Future directions

StrainSifter can be used to precisely identify strains within relatively biomass-rich metagenomic samples without relying on external reference genome databases in which representation of opportunistic pathogens may be lacking. Compared to stool, samples from the nares, skin, or the environment typically have far less microbial biomass, limiting the number of sequencing reads that can be obtained. In order to provide the most precise epidemiologic information about the origins of bloodstream and other infections, StrainSifter will need to be tested in low microbial biomass samples in a larger prospective study of immunocompromised patients. Given the high resolution of WGS-based strain comparisons, we may indeed discover instances of organisms thought to originate from non-enteric sites instead colonizing and translocating from the intestine in severely immunocompromised patients, as was potentially demonstrated by *S. epidermidis* and *P. aeruginosa* in our study. Given the growing body of research on therapies to improve gut microbiota diversity and bolster colonization resistance against pathogens, more precisely identifying the origins of bloodstream infections in the stem cell transplant population may influence how hospitals and healthcare providers can most effectively work to prevent infections (Galloway-Peña, Jenq, and Shelburne 2017). Finally, once StrainSifter is able to establish the origins of an infection, the wealth of information offered by WGS can be interrogated to identify potential functional differences, including in gene content and metabolic pathways, between strains colonizing the gut, nares, or skin that result in disseminated infections compared to those that do not.

In conclusion, we have shown that StrainSifter offers a straightforward bioinformatic pipeline to precisely compare bacterial isolates to metagenomic samples, circumventing the need for external reference genome databases or culture-based methods of strain comparison. We successfully applied StrainSifter to blood culture isolates and stool samples from hematopoietic stem cell transplant patients and found classically non-enteric bacteria in the gut microbiome that were concordant with bloodstream isolates. We anticipate that StrainSifter will be an easy-to-use tool in the context of hospital or infectious outbreak

epidemiologic studies, and that characterization of the microbiome dynamics that occur prior to infection

may help us precisely identify potential reservoirs of pathogens.

Acknowledgements

We thank Joyce Kang for her assistance with stool sample processing, as well the other members of the

Bhatt Lab for providing feedback on the study design, bioinformatics pipeline, and manuscript revisions.

We would like to thank Nick Greenfield and the One Codex team for help with using their platform. We

appreciate Drs. Matthew Kelly, Chris Severyn, and Doyle Ward for their feedback and ideas on the

manuscript. We would especially like to thank the patients and nurses on the Blood and Marrow

Transplant service for their enthusiastic participation in this project. This work was supported in part by

National Science Foundation Graduate Research Fellowship (FBT), the National Institutes of Health,

National Center for Advancing Translational Science, Clinical and Translational Science Award KL2

TR000083 and UL1 TR001085 (TMA). ASB was funded in part by the National Cancer Institute NIH

K08 award, #CA184420, Damon Runyon Clinical Investigator Award, and Amy Strelzer Manasevit

Award. The content is solely the responsibility of the authors and does not necessarily represent the

official views of the NIH.

References

Advani, Sonali, Rachael Lee, Mariann Schmitz, Martha Long, and Bernard Camins. 2017. "2015 NHSN

CAUTI Definition Change and Its Impact on CLABSI Rates at an Academic Medical Center." *Open*

Forum Infectious Diseases 4 (suppl_1): S180–S180.

Bankevich, Anton, Sergey Nurk, Dmitry Antipov, Alexey A. Gurevich, Mikhail Dvorkin, Alexander S.

Kulikov, Valery M. Lesin, et al. 2012. "SPAdes: A New Genome Assembly Algorithm and Its

Applications to Single-Cell Sequencing." *Journal of Computational Biology: A Journal of*

Computational Molecular Cell Biology 19 (5): 455–77.

Barnett, Derek W., Erik K. Garrison, Aaron R. Quinlan, Michael P. Strömberg, and Gabor T. Marth.

2011. "BamTools: A C++ API and Toolkit for Analyzing and Managing BAM Files."

Bioinformatics 27 (12): 1691–92.

Blennow, O., P. Ljungman, E. Sparrelid, J. Mattsson, and M. Remberger. 2014. "Incidence, Risk Factors,

and Outcome of Bloodstream Infections during the Pre-Engraftment Phase in 521 Allogeneic

Hematopoietic Stem Cell Transplantations." *Transplant Infectious Disease: An Official Journal of*

the Transplantation Society 16 (1): 106–14.

"Centers for Disease Control and Prevention." 2018. Bloodstream Infection Event (Central Line-

Associated Bloodstream Infection and Non-Central Line-Associated Bloodstream Infection). The

Centers for Disease Control and Prevention Web Site. February 2018.

http://www.cdc.gov/nhsn/PDFs/pscManual/4PSC_CLABS current.pdf.

Chen, Yi, Yan Luo, Heather Carleton, Ruth Timme, David Melka, Tim Muruvanda, Charles Wang, et al.

2017. "Whole Genome and Core Genome Multilocus Sequence Typing and Single Nucleotide

Polymorphism Analyses of Listeria Monocytogenes Associated with an Outbreak Linked to Cheese,

United States, 2013." *Applied and Environmental Microbiology* 83 (15): 1–15.

Costa, Silvia F., Antonio A. Barone, Marisa H. Miceli, Inneke M. van der Heijden, Robson E. Soares,

Anna S. Levin, and Elias J. Anaissie. 2006. "Colonization and Molecular Epidemiology of

Coagulase-Negative Staphylococcal Bacteremia in Cancer Patients: A Pilot Study." *American Journal of Infection Control* 34 (1): 36–40.

Costea, Paul Igor, Robin Munch, Luis Pedro Coelho, Lucas Paoli, Shinichi Sunagawa, and Peer Bork. 2017. "metaSNV: A Tool for Metagenomic Strain Level Analysis." *PloS One* 12 (7): 1–9.

Edgar, Robert C. 2004. "MUSCLE: Multiple Sequence Alignment with High Accuracy and High Throughput." *Nucleic Acids Research* 32 (5): 1792–97.

"GAEMR." n.d. Accessed April 19, 2018. https://www.broadinstitute.org/software/gaemr/.

Galloway-Peña, Jessica R., Robert R. Jenq, and Samuel A. Shelburne. 2017. "Can Consideration of the Microbiome Improve Antimicrobial Utilization and Treatment Outcomes in the Oncology Patient?" *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research* 23 (13): 3263–68.

Gorrie, Claire L., Mirjana Mirčeta, Ryan R. Wick, David J. Edwards, Nicholas R. Thomson, Richard A. Strugnell, Nigel F. Pratt, et al. 2017. "Gastrointestinal Carriage Is a Major Reservoir of Klebsiella Pneumoniae Infection in Intensive Care Patients." *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* 65 (2): 1–8.

Gudiol, C., C. Garcia-Vidal, M. Arnan, I. Sánchez-Ortega, B. Patiño, R. Duarte, and J. Carratalà. 2014. "Etiology, Clinical Features and Outcomes of Pre-Engraftment and Post-Engraftment Bloodstream Infection in Hematopoietic SCT Recipients." *Bone Marrow Transplantation* 49 (6): 824–30.

Harris, Simon R., Edward J. P. Cartwright, M. Estée Török, Matthew T. G. Holden, Nicholas M. Brown, Amanda L. Ogilvy-Stuart, Matthew J. Ellington, et al. 2013. "Whole-Genome Sequencing for Analysis of an Outbreak of Meticillin-Resistant Staphylococcus Aureus: A Descriptive Study." *The Lancet Infectious Diseases* 13 (2): 130–36.

Herwaldt, L. A., R. J. Hollis, L. D. Boyken, and M. A. Pfaller. 1992. "Molecular Epidemiology of Coagulase-Negative Staphylococci Isolated from Immunocompromised Patients." *Infection Control and Hospital Epidemiology: The Official Journal of the Society of Hospital Epidemiologists of America* 13 (2): 86–92.

Inouye, Michael, Harriet Dashnow, Lesley-Ann Raven, Mark B. Schultz, Bernard J. Pope, Takehiro
Tomita, Justin Zobel, and Kathryn E. Holt. 2014. "SRST2: Rapid Genomic Surveillance for Public
Health and Hospital Microbiology Labs." *Genome Medicine* 6 (11): 90.

Kovanen, Sara M., Rauni I. Kivistö, Mirko Rossi, Thomas Schott, Ulla-Maija Kärkkäinen, Tamara
Tuuminen, Jaakko Uksila, Hilpi Rautelin, and Marja-Liisa Hänninen. 2014. "Multilocus Sequence
Typing (MLST) and Whole-Genome MLST of Campylobacter Jejuni Isolates from Human
Infections in Three Districts during a Seasonal Peak in Finland." *Journal of Clinical Microbiology*
52 (12): 4147–54.

Lieberman, Tami D., Kelly B. Flett, Idan Yelin, Thomas R. Martin, Alexander J. McAdam, Gregory P.
Priebe, and Roy Kishony. 2013. "Genetic Variation of a Bacterial Pathogen within Individuals with
Cystic Fibrosis Provides a Record of Selective Pressures." *Nature Genetics* 46 (1): 82–87.

Li, H., and R. Durbin. 2009. "Fast and Accurate Short Read Alignment with Burrows–Wheeler
Transform." *Bioinformatics* 25 (14): 1754–60.

Li, Heng, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth, Goncalo
Abecasis, Richard Durbin, and 1000 Genome Project Data Processing Subgroup. 2009. "The
Sequence Alignment/Map Format and SAMtools." *Bioinformatics* 25 (16): 2078–79.

Li, Wenjun, Didier Raoult, and Pierre-Edouard Fournier. 2009. "Bacterial Strain Typing in the Genomic
Era." *FEMS Microbiology Reviews* 33 (5): 892–916.

MacFie, J., C. O'Boyle, C. J. Mitchell, P. M. Buckley, D. Johnstone, and P. Sudworth. 1999. "Gut Origin
of Sepsis: A Prospective Study Investigating Associations between Bacterial Translocation, Gastric
Microflora, and Septic Morbidity." *Gut* 45 (2): 223–28.

Martin, Rebekah M., Jie Cao, Sylvain Brisse, Virginie Passet, Weisheng Wu, Lili Zhao, Preeti N. Malani,
Krishna Rao, and Michael A. Bachman. 2016. "Molecular Epidemiology of Colonizing and Infecting
Isolates of Klebsiella Pneumoniae." *mSphere* 1 (5): 1–12.

Mikulska, Malgorzata, Valerio Del Bono, Anna Maria Raiola, Barbara Bruno, Francesca Gualandi,
Domenico Occhini, Carmen di Grazia, Francesco Frassoni, Andrea Bacigalupo, and Claudio Viscoli.

2009. "Blood Stream Infections in Allogeneic Hematopoietic Stem Cell Transplant Recipients:

Reemergence of Gram-Negative Rods and Increasing Antibiotic Resistance." *Biology of Blood and*

Marrow Transplantation: Journal of the American Society for Blood and Marrow Transplantation

15 (1): 47–53.

Minot, Samuel S., Niklas Krumm, and Nicholas B. Greenfield. 2015. "One Codex: A Sensitive and

Accurate Data Platform for Genomic Microbial Identification." *bioRxiv*, 027607.

Nayfach, Stephen, Beltran Rodriguez-Mueller, Nandita Garud, and Katherine S. Pollard. 2016. "An

Integrated Metagenomics Pipeline for Strain Profiling Reveals Novel Patterns of Bacterial

Transmission and Biogeography." *Genome Research* 26 (11): 1612–25.

Nesher, L., K. V. I. Rolston, D. P. Shah, J. T. Tarrand, V. Mulanovich, E. J. Ariza-Heredia, and R. F.

Chemaly. 2014. "Fecal Colonization and Infection with Pseudomonas aeruginosa in Recipients of

Allogeneic Hematopoietic Stem Cell Transplantation." *Transplant Infectious Disease: An Official*

Journal of the Transplantation Society 17 (1): 33–38.

Nielsen, Karen Leth, Marc Stegger, Kristoffer Kiil, Paul A. Godfrey, Michael Feldgarden, Berit Lilje,

Paal S. Andersen, and Niels Frimodt-Møller. 2017. "Whole-Genome Comparison of Urinary

Pathogenic Escherichia Coli and Faecal Isolates of UTI Patients and Healthy Controls."

International Journal of Medical Microbiology: IJMM 307 (8): 497–507.

Nurk, Sergey, Dmitry Meleshko, Anton Korobeynikov, and Pavel A. Pevzner. 2017. "metaSPAdes: A

New Versatile Metagenomic Assembler." *Genome Research* 27 (5): 824–34.

Price, Morgan N., Paramvir S. Dehal, and Adam P. Arkin. 2010. "FastTree 2--Approximately Maximum-

Likelihood Trees for Large Alignments." *PloS One* 5 (3). Public Library of Science: e9490.

"PubMLST." n.d. Accessed April 20, 2018. https://pubmlst.org/.

Rotstein, C., D. Higby, K. Killion, and E. Powell. 1988. "Relationship of Surveillance Cultures to

Bacteremia and Fungemia in Bone Marrow Transplant Recipients with Hickman or Broviac

Catheters." *Journal of Surgical Oncology* 39 (3): 154–58.

Salipante, Stephen J., Dhruba J. SenGupta, Lisa A. Cummings, Tyler A. Land, Daniel R. Hoogestraat,

and Brad T. Cookson. 2015. "Application of Whole-Genome Sequencing for Bacterial Strain Typing

in Molecular Epidemiology." *Journal of Clinical Microbiology* 53 (4): 1072–79.

Samet, A., A. Śledzińska, B. Krawczyk, A. Hellmann, S. Nowicki, J. Kur, and B. Nowicki. 2013.

"Leukemia and Risk of Recurrent Escherichia Coli Bacteremia: Genotyping Implicates E. Coli

Translocation from the Colon to the Bloodstream." *European Journal of Clinical Microbiology &*

Infectious Diseases: Official Publication of the European Society of Clinical Microbiology 32 (11):

1393–1400.

See, Isaac, Minn M. Soe, Lauren Epstein, Jonathan R. Edwards, Shelley S. Magill, and Nicola D.

Thompson. 2017. "Impact of Removing Mucosal Barrier Injury Laboratory-Confirmed Bloodstream

Infections from Central Line-Associated Bloodstream Infection Rates in the National Healthcare

Safety Network, 2014." *American Journal of Infection Control* 45 (3): 321–23.

Snitkin, E. S., A. M. Zelazny, P. J. Thomas, F. Stock, D. K. Henderson, T. N. Palmore, J. A. Segre, and

Nisc Comparative Sequencing Program. 2012. "Tracking a Hospital Outbreak of Carbapenem-

Resistant Klebsiella Pneumoniae with Whole-Genome Sequencing." *Science Translational Medicine*

4 (148): 148ra116–48ra116.

Snitkin, Evan S., Adrian M. Zelazny, Clemente I. Montero, Frida Stock, Lilia Mijares, NISC Comparative

Sequence Program, Patrick R. Murray, and Julie A. Segre. 2011. "Genome-Wide Recombination

Drives Diversification of Epidemic Strains of Acinetobacter Baumannii." *Proceedings of the*

National Academy of Sciences of the United States of America 108 (33): 13758–63.

Tancrede, C. H., and A. O. Andremont. 1985. "Bacterial Translocation and Gram-Negative Bacteremia in

Patients with Hematological Malignancies." *The Journal of Infectious Diseases* 152 (1): 99–103.

Taur, Ying, Joao B. Xavier, Lauren Lipuma, Carles Ubeda, Jenna Goldberg, Asia Gobourne, Yeon Joo

Lee, et al. 2012. "Intestinal Domination and the Risk of Bacteremia in Patients Undergoing

Allogeneic Hematopoietic Stem Cell Transplantation." *Clinical Infectious Diseases: An Official*

Publication of the Infectious Diseases Society of America 55 (7): 905–14.

Truong, Duy Tin, Adrian Tett, Edoardo Pasolli, Curtis Huttenhower, and Nicola Segata. 2017. "Microbial

Strain-Level Population Structure & Genetic Diversity from Metagenomes." *Genome Research* 27

(4): 626–38.

Ubeda, Carles, Ying Taur, Robert R. Jenq, Michele J. Equinda, Tammy Son, Miriam Samstein, Agnes

Viale, et al. 2010. "Vancomycin-Resistant Enterococcus Domination of Intestinal Microbiota Is

Enabled by Antibiotic Treatment in Mice and Precedes Bloodstream Invasion in Humans." *The*

Journal of Clinical Investigation 120 (12): 4332–41.

Wade, J. C., S. C. Schimpff, K. A. Newman, and P. H. Wiernik. 1982. "Staphylococcus Epidermidis: An

Increasing Cause of Infection in Patients with Granulocytopenia." *Annals of Internal Medicine* 97

(4): 503–8.

591