High impact journals in ecology cover proportionally more statistically

significant findings

Silvia Ceauşu[1,2,3,4*], Luís Borda-de-Água[5], Thomas Merckx[6], Esther Sossai[3,4], Manuel

Sapage[7], Murilo Miranda[3,4], Henrique M. Pereira[3,4,5]

1. Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Aarhus

University, Aarhus C, Denmark

2. Section for Ecoinformatics and Biodiversity, Department of Bioscience, Aarhus University,

Aarhus C, Denmark

3. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig,

Germany.

4. Institute of Biology, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

5. Infraestruturas de Portugal Biodiversity Chair, CIBIO/InBIO, Campus Agrário de Vairão,

Vairão, Portugal

6. Behavioural Ecology and Conservation Group, Biodiversity Research Centre, Earth and

Life Institute, Université catholique de Louvain (UCL), Louvain-la-Neuve, Belgium

7. cE3c—Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências,

Universidade de Lisboa, Lisboa, Portugal

*corresponding author: S. Ceauşu: silvia.ceausu@mespom.eu; +49(0)341-97-33136.

Abstract

Unbiased scientific reporting is crucial for data and research synthesis. Previous studies suggest that statistically significant results are more likely to be published and more likely to be submitted to high impact journals. However, the most recent research on statistical significance in relation to journal impact factors in ecological research was published more than two decades ago or addressed a small subset of the literature. Here, we extract p-values from all articles published in 11 journals in 2012 and 2014 across a wide range of impact factors with six journals sampled in both years. Our results indicate that the proportion of statistically significant results increases with rising impact factor. Such a trend can have important consequences for syntheses of ecological data and it highlights the importance of covering a wide range of impact factors when identifying published studies for data syntheses. This trend can also lead to a biased understanding of the probability of true effects in ecology and conservation. We caution against the possible downplaying of non-significant results by either journals or authors.

Introduction

Research synthetizing published data in ecology and biological sciences is growing (1) but the validity of its results depends on unbiased reporting of research, including of statistically non-significant results. Incentives in academia that emphasize citations indices and publications in high impact factor (IF) journals may undermine this requirement. For instance, research suggests that statistically non-significant results are less likely to be published in the case of clinical trial studies (2) and are submitted to lower IF journals in ecology (3). Such a bias in publication can result in overestimating statistical significance and effect size in research synthetizing data to estimate a particular effect or phenomenon. Moreover, statistically non-significant results can be highly significant scientifically. Research on the effect of these

46 behaviours on the overall pattern of reported statistical significance in relation to IF has only

47 been conducted on a small subset of the literature (3), on a small range of IF (4) or has been

48 published more than two decades ago (5). Here, we update the research on the relationship

49 between significance levels and IF in the ecology and conservation literature within a wide

50 range of IF.

Methods

We examined how the proportion of reported significant results, expressed as p-values,

changed with increasing IF. We divided the range of IF into three intervals: low (IF<4),

medium (4≤IF<8), and high (IF≥8) to ensure that we cover a broad range of IF. We then

randomly selected at least two journals for each IF interval for both 2012 and 2014 from

journals listed in the Science Citation Index Expanded (http://mjl.clarivate.com/cgi-

bin/jrnlst/jloptions.cgi?PC=D) under four subject categories: biodiversity conservation,

biology, ecology, and evolutionary biology. We examined 11 journals (table in S1 Table)

(lowest IF 0.36 to highest IF 17.95), six journals for both years, whilst we examined three for

2014 and two for 2012 alone. We collected all p-values reported in all articles using

Examine32 Text Search from Aquila Software. We extracted exact p-values and inexact p-

values (e.g. p<0.05). All the inexact p-values were reassigned to six intervals in order to

harmonize all the reported values and calculate the proportion of significant results (Table A

in S1 Text). Due to the ubiquitous use of 0.05 as the alpha level in ecology, we considered p-

values below this level as significant. We calculated the proportion of significant results for

exact p-values and all (i.e. exact and inexact) p-values reported in each journal, for each year.

We fitted generalized linear mixed effects models (binomial distribution) to the proportion of

significant results with IF as fixed effect and journal as random effect. We used R (version

3.4.0) and the "lme4" and "arm" packages for model fitting and back-transformation of model

coefficients. We tested through a paired t-test the effect of year on the six journals that we

examined for both years. All data collected for this study are available at:

https://figshare.com/projects/High_impact_journals_cover_proportionally_more_statistically_

significant_findings/28284. The R code of the data analysis is available at:

https://github.com/SilviaCeausu/ImpactFactorsAndPvalues/blob/master/ImpactFactorPvaluec

odeGitHub2.R.

Figure 1. Proportion of a) exact p-values and b) all p-values below 0.05 reported across

impact factors (IF). The grey areas delimitate the 95% confidence intervals. The journals

considered and their impact factors are presented in S1 Table.

Results and discussion

The percentage of significant results–out of all published results within a journal– increased

with increasing IF (Figure 1). For exact p-values, the percentage of significant results

increased on average by 0.7% for each additional IF unit [95% CI: 0.3 - 1.2%]. For all p-

values (i.e. exact and inexact), the percentage increased by 1.2% for each additional IF unit

[95% CI: 0.9 - 1.5%] (Table S2). In 2012, the percentage of significant results was higher

than in 2014 (+3.3%, 95% CI: 0.2 – 6.4%) for the journals examined in both years.

Our result concurs with trends noticed in medical research (6) but they contradict results

reported for behavioural ecology by (4). The latter study concluded that neither p-values nor

statistical power varied significantly with IF but the analysis was conducted on a much

narrower range of IF (ca. 1 - 5) than our study.

Our result can arise if statistical significance influences submission or editorial decisions.

Analysing the output of doctoral dissertations in ecology, Koricheva (2) found that the

proportion of non-significant results in a study was negatively associated with IF, although

the rejection rates for non-significant results were not higher for higher IF journals. In an

article that examined clinical trials, Suñé et al (3) found that non-significant studies are less

likely to be published and, if published, more time passes between conducting the research

and publication. However, the IF of the publishing journal was not different for significant

versus non-significant results (3). These studies suggest that authors invest less effort into the

publication of their non-significant results and submit them to lower IF journals.

Our outcome can also be an effect of higher impact journals selecting studies with higher

sample or effect sizes, or requesting stricter statistical reporting and shorter articles. In studies

of the relationships between IF, and sample and effects sizes results are mixed. For meta-

analyses in ecology Lortie et al (7) did not find a relationship between IF and effect size.

Analyzing studies collected for four meta-analyses, Murtaugh found a positive correlation

between effect strength and IF in two of the four datasets (8), and Barto and Rilig (9) found

that high IF journal published the strongest effects, although in the absence of correlations

with data quality. Regarding statistical reporting, Tressoldi et al (10) suggest that higher IF

journal do not necessarily display better standards. Our data also show no indication that

higher IF journals publish more precise p-values then lower IF journals (figure in S1 Figure).

The heterogeneity of article length requirements across journals did not allow us to test

whether article length requirements play a role in our result (table in S1 Table). Moreover, we

did not analyse supplementary materials, which might include additional non-significant

results considered secondary by authors.

The large confidence intervals in our results suggest that other factors also have an influence

on publication. For example, the difference in percentage of significant results between years

suggests changes in the prominence of different research topics. However, we cannot exclude

an undervaluation of non-significant results, either by authors or by journals. This pattern may

make significant results more visible if they are published in higher IF journals than non-

significant findings, and may create an inaccurate perception of the probability of true effects

in ecology. This could lead to wasted efforts on approaches or interventions that could in

reality be far less effective than we assume (11). Publication biases could also negatively

impact our understanding of biodiversity change and its drivers if a higher proportion of non-

significant results remain unpublished compared with significant ones, especially at a time

when growing synthesis efforts are trying to shed light on important biodiversity and ecology

questions (1,12,13). Moreover, statistically non-significant results can give rise to new

theories or amendments to old ones, as it is the case of the emerging debate on the importance

of isolation in fragmented landscapes (14,15). Therefore, we advise careful consideration of

submission and publication decisions to ensure solid foundations for our scientific

understanding.

Acknowledgments

133   MS is funded by a PhD grant from Fundacão para a Ciência e a Tecnologia (FCT), Portugal

134   (ref. PD/BD/128349/2017).

References

1. Newbold T, Hudson LN, Hill SL, Contu S, Lysenko I, Senior RA, et al. Global effects of
land use on local terrestrial biodiversity. Nature. 2015;520(7545):45–50.

2. Suñé P, Suñé JM, Montoro JB. Positive outcomes influence the rate and time to
publication, but not the impact factor of publications of clinical trial results. PloS One.
2013;8(1):e54583.

3. Koricheva J. Non-significant results in ecology: a burden or a blessing in disguise? Oikos.
2003;397–401.

4. Jennions MD, Møller AP. A survey of the statistical power of research in behavioral
ecology and animal behavior. Behav Ecol. 2003;14(3):438–445.

5. Csada RD, James PC, Espie RH. The" file drawer problem" of non-significant results: does
it apply to biological research? Oikos. 1996;591–593.

6. Jannot A-S, Agoritsas T, Gayet-Ageron A, Perneger TV. Citation bias favoring statistically
significant studies was present in medical research. J Clin Epidemiol. 2013;66(3):296–
301.

7. Lortie CJ, Aarssen LW, Budden AE, Leimu R. Do citations and impact factors relate to the
real numbers in publications? A case study of citation rates, impact, and effect sizes in
ecology and evolutionary biology. Scientometrics. 2013;94(2):675–682.

8. Murtaugh PA. Journal Quality, Effect Size, and Publication Bias in Meta-Analysis.
Ecology. 2002 Apr 1;83(4):1162–6.

9. Barto EK, Rillig MC. Dissemination biases in ecology: effect sizes matter more than
quality. Oikos. 2012 Feb 1;121(2):228–35.

10. Tressoldi PE, Giofré D, Sella F, Cumming G. High impact= high statistical standards?
Not necessarily so. PloS One. 2013;8(2):e56180.

11. Meli P, Holl KD, Benayas JMR, Jones HP, Jones PC, Montoya D, et al. A global review
of past land use, climate, and active vs. passive restoration effects on forest recovery.
PloS One. 2017;12(2):e0171368.

12. Dornelas M, Gotelli NJ, McGill B, Shimadzu H, Moyes F, Sievers C, et al. Assemblage
Time Series Reveal Biodiversity Change but Not Systematic Loss. Science.
2014;344(6181):296–299.

13. Vellend M, Baeten L, Myers-Smith IH, Elmendorf SC, Beauséjour R, Brown CD, et al.
Global meta-analysis reveals no net change in local-scale plant biodiversity over time.
Proc Natl Acad Sci. 2013;110(48):19456–19459.

14. Fahrig L. Rethinking patch size and isolation effects: the habitat amount hypothesis. J
Biogeogr. 2013 Sep 1;40(9):1649–63.

15. Collinge SK. Effects of Grassland Fragmentation on Insect Species Loss, Colonization,
and Movement Patterns. Ecology. 2000;81(8):2211–26.

Supporting information captions

S1 File. Data collection protocol (separate document)

S1 Table. Information regarding the journals included in the analysis. Journal title,

publication year, the impact factor in the respective year, length limits for the main article

type of the journal, total number of articles published by each journal during the respective

year, total number of exact and inexact p-values considered. NL - no length limit specified in

the author guidelines.

Journal Name	Year	Impact factor	Length limits	Number of total published articles	Number of exact p-values	Number of inexact p-values
Agriculture Ecosystems & Environment	2012	2.859	NL	268	1980	4050
	2014	3.402		269	2260	4050
Biochemical Systematics And Ecology	2014	0.967	NL	261	393	1132
Ecology Letters	2012	17.949	5000 words	167	1263	1177
	2014	10.689		157	1041	787
Global Change Biology	2014	8.044	8000	317	2589	4853
Global Ecology and Biogeography	2012	7.223	10 printed pages, ~ 5000 words	111	418	1599
	2014	6.531		133	660	1193
Journal of Animal Ecology	2012	4.841	8500 words	150	1308	892
	2014	4.504		137	1726	1304
Journal of Ecology	2014	5.521	12 typeset pages; ~ 9500 words	155	2435	2829
Journal of Evolutionary Biology	2012	3.479	6-10 printed pages	237	4184	2418
	2014	3.232		256	4412	2828
PLOS Biology	2012	12.69	NL	153	689	520
	2014	9.343		223	560	702
Northeastern Naturalist	2012	0.362	NL	51	347	216
Wildlife Biology	2012	1.102	40000 characters	43	232	147

S2 Table. Model coefficients and selection. We tested the model used for Figure1 (main

text) against alternative random effects structures and null models. A model with both random

slope and intercept was not possible due to the low number of data points. We compared the

models according to theoretic information criteria: Akaikes' Information Criterion (AIC) and

Bayesian Information Criterion (BIC). We used the same models for both exact p-values and

all p-values. We rounded up all values to three decimal places.

Model	Coefficients (95% CI)		AIC	BIC
Exact p-values	Intercept	IF		
Final model %significant ~IF + (1|journal)	-0.364 (-0.599, -0.129)	0.032 (0.013, 0.051)	206.30	208.80
Null model - journal %significant ~ 1 + (1 | journal)	-0.187 (-0.412, 0.039)	-	214.75	216.41
Random intercept - journal %significant ~ IF + (IF - 1 | journal)	-0.38 (-0.581, -0.188)	0.069 (-0.118, 0.261)	231.75	234.25
All p-values				
Final model %significant ~ IF + (1 | journal)	0.076 (-0.11, 0.245)	0.055 (0.04, 0.071)	286.27	288.77
Random slope - journal %significant ~IF + (IF - 1 | journal)	0.027 (-0.139, 0.18)	0.121 (0.012, 0.263)	317.52	320.02
Null model - journal %significant ~1 + (1 | journal)	0.383 (0.242, 0.523)	-	339.72	341.38

206   **S1 Figure. Proportion of exact p-values below 0.05 reported in a) 2012 and b) 2014**

207   **across the range of impact factors (IF).**

211 **S2 Figure. Proportion of inexact p-values out of the total reported values across the**

212 **range of impact factors (IF).**

216