Inactivation of the monofunctional peptidoglycan glycosyltransferase SgtB allows *Staphylococcus aureus* to survive in the absence of lipoteichoic acid

Eleni Karinou[1]^, Christopher F. Schuster[1&], Manuel Pazos[2&], Waldemar Vollmer[2] and Angelika Gründling[1]*

[1]Section of Microbiology and MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London SW7 2AZ, UK

[2]The Centre for Bacterial Cell Biology, Newcastle University, Newcastle upon Tyne NE2 4AX, UK

^ Present address: The Centre for Bacterial Cell Biology, Newcastle University, Newcastle upon Tyne NE2 4AX, UK

& these authors contributed equally

* For correspondence:

Angelika Gründling

Section of Microbiology and MRC Centre of Molecular Bacteriology and Infection

Flowers Building RM 3.21

Imperial College London; London, SW7 2AZ, UK

Tel. +44 (0)2075945256

Fax +44 (0)2075943095

E-mail: a.grundling@imperial.ac.uk

Running title: Characterization of LTA negative *S. aureus* strains

Abstract

The cell wall of *Staphylococcus aureus* is composed of peptidoglycan and the anionic polymers lipoteichoic acid (LTA) and wall teichoic acid. LTA is required for growth and normal cell morphology in *S. aureus.* Strains lacking LTA are usually only viable when grown under osmotically stabilizing conditions or after the acquisition of compensatory mutations. LTA negative suppressor strains with inactivating mutations in *gdpP*, resulting in an increase in intracellular c-di-AMP levels, have been described previously. Here, we sought to identify factors other than c-di-AMP that allow *S. aureus* to survive without LTA. LTA-negative strains able to grow in un-supplemented medium were obtained and found to contain mutations in *sgtB, mazE, clpX* or *vraT*. The growth improvement through mutations in *mazE* and *sgtB* was confirmed by complementation analysis. We also show that an *S. aureus sgtB* transposon mutant, inactivated for the monofunctional peptidoglycan glycosyltransferase SgtB, displays a 4-fold increase in the MIC towards a number of cell wall-targeting antibiotics, suggesting that alteration in the peptidoglycan structure could help bacteria compensate for the lack of LTA. Muropeptide analysis of peptidoglycan isolated from a WT and *sgtB* mutant strains did not reveal any sizable alternations in the peptidoglycan structure. In contrast, the peptidoglycan isolated from an LTA-negative *ltaS* mutant strain showed a significant reduction in the fraction of highly crosslinked peptidoglycan, which was partially rescued in the *sgtB*/*ltaS* double mutant suppressor strain. Taken together, these data point towards an important function of LTA in cell wall integrity through its requirement for proper peptidoglycan assembly.

Importance

The bacterial cell wall acts as primary defence against environmental insults such as changes in osmolarity. It is also a vulnerable structure as defects in its synthesis can lead to growth arrest or cell death. The important human pathogen *Staphylococcus aureus* has a typical Gram-

positive cell wall, which consists of peptidoglycan and the anionic polymers lipoteichoic acid

(LTA) and wall teichoic acid. Several clinically relevant antibiotics inhibit the synthesis of

peptidoglycan; hence it and teichoic acids are considered attractive targets for the development

of new antimicrobials. We show that LTA is required for efficient peptidoglycan crosslinking

in *S. aureus* and inactivation of a peptidoglycan glycosyltransferase can partially rescue this

defect, altogether revealing an intimate link between peptidoglycan and LTA synthesis.

Introduction

Staphylococcus aureus is a Gram-positive bacterium found as commensal on the skin and nasal

passages of healthy individuals. However, this bacterium is also an important human pathogen

causing hospital- as well as community-acquired infections, such as serious skin infections,

osteomyelitis and endocarditis (1-3). Of major concern is the increasing resistance of this

organism to a large number of clinically relevant antibiotics (4). A number of virulence factors

contribute to the successful host colonization, immune evasion and acquisition of nutrients

within the host (5). Many of these factors are either secreted proteins or other extracellular

proteins closely associated with the bacterial cell envelope (6).

The cell envelope is essential for bacterial survival and pathogenesis but also a target

of a number of important antimicrobials. It functions as barrier and protects bacteria from

environmental insults but at the same time needs to allow the passage of solutes and nutrients

as well as sense changes in the external environment (7). *S. aureus* has a typical Gram-positive

cell envelop, which consists of a cytoplasmic membrane that is surrounded by a thick

peptidoglycan layer (8). The peptidoglycan layer is a dynamic macromolecular structure that

undergoes constant cycles of polymerisation and hydrolysis to allow bacteria to grow and

divide (7). It is composed of glycan chains made of alternating N-acetylglucosamine and N-

acetylmuramic acid residues connected by peptide bridges (9). This mesh-like sacculus is able

to protect the cell from environmental threats and at the same time withstand the high internal

osmotic pressure (10). The final steps of peptidoglycan synthesis are catalyzed by enzymes

named penicillin binding proteins (PBPs) and a coordinated action of these enzymes is crucial

for the survival of the cell (11). PBPs with glycosyltransferase and transpeptidase activity

polymerize the glycan chains and form peptide cross-bridges while the monofunctioal

transpeptidases only have the former activity (11).

S. aureus encodes four PBPs, of which PBP1, which has transpeptidase activity, and

the bifunctional PBP2, which has transpeptidase and glycosltransferase activity, are the

minimal requirement for cell survival (12). In methicillin-resistant *S. aureus* (MRSA) strains,

the alternative PBP2A, which has transpeptidase activity, is needed for β-lactam resistance in

addition to the glycosyltransferase activity of PBP2 (13, 14). Moreover, additional non-

essential proteins have been identified that are involved in peptidoglycan synthesis such as the

monofunctional glycosyltransferases SgtA and SgtB (also named Mgt) (12). Previous studies

have shown that although SgtA and SgtB have glycosyltransferase activity *in vitro*, only SgtB

can support the growth of *S. aureus* in the absence of the main glycosyltransferase PBP2.

However, this is not possible in the presence of β-lactam antibiotics when an interaction

between PBP2 and PBP2A is needed (15, 16).

In Gram-positive bacteria, the peptidoglycan layer is interspersed by a plethora of

proteins and cell wall polymers named teichoic acids (17). Teichoic acids are further

categorised into lipoteichoic acids (LTA), anchored to the outer leaflet of the cytoplasmic

membrane via a lipid moiety, and wall teichoic acids (WTA) covalently attached to the

peptidoglycan (18). Teichoic acids form an important part of the cell wall and contribute to the

physical and chemical properties of the cell wall, and to the binding of divalent cations (19,

20). While both WTA and LTA are polyanioinic cell wall polymers, they are synthesized

through separate and independent pathways in *S. aureus* and many other Gram-positive

bacteria (21, 22). Consistent with this, our work on the Gram-positive pathogen *Listeria monocytogenes* has revealed that LTA synthesis is not abrogated in the absence of WTA and *vice versa* (23). Recent work using pathway specific inhibitors and a gene interaction screen provided further evidence that the polymers are not only synthesized through separate pathways but also have distinct functions in *S. aureus* (24).

LTA is an anionic polymer made up in *S. aureus* of glycerolphosphate repeating units that are further decorated with D-Alanine residues and, as shown recently, under high salt conditions also with N-acetylglucosamine residues (25, 26). Most proteins required for its synthesis have now been identified and extensively studied over the years (22, 27). One of the key enzymes required for LTA synthesis is the lipoteichoic acid synthase enzyme LtaS (28, 29). This enzyme polymerizes the LTA backbone chain on the outside of the cell using the glycerolphosphate head group of the membrane lipid phosphatidylglycerol as substrate (29, 30). LTA is indispensable for the growth of *S. aureus* under standard laboratory growth conditions, highlighting its important physiological role (29, 31).

Previous studies indicated a function for LTA in helping to direct the cell-division machinery (32), in controlling autolysin activity (31), for biofilm formation (33), in mediating interactions with host cell receptors (17), in controlling susceptibility and/or resistance to antimicrobial peptides and in maintaining cation homeostasis (20). *S. aureus ltaS* mutants, which lack the complete LTA polymer, can in some strain backgrounds be constructed at low growth temperature (31). It has also been shown that LTA-deficient *S. aureus* strains are viable when grown under osmotically stabilizing conditions in medium containing 7.5% NaCl or 40% sucrose (31, 34). However, the bacteria display severe morphological defects including an enlarged cell size, clustering and cell division defects, even under those conditions permissive for growth (34). Bacteria can also readily acquire compensatory mutations allowing them to grow in un-supplemented medium and improving their cell morphology defects (34). The

majority of compensatory mutations previously observed were in *gdpP* leading to inactivation

of the c-di-AMP phosphodiesterase GdpP (34). The resulting increase in cellular c-di-AMP

levels allowed the bacteria to survive the cell wall stress caused by the absence of LTA, which

is now believed to be due to changes in the osmotic balance in the cell (34). However,

compensatory mutations that rescue the growth of an LTA negative *S. aureus* strain were not

only found in *gdpP* but also in other genes (34).

As part of the current study, we sought to identify these genes and further characterize

the encoded proteins to gain additional insight into why LTA is essential for the growth of *S.*

aureus and potentially uncover novel cellular functions for proteins involved in cell wall

assembly or maintenance. Using a suppressor screen approach, we found that inactivation of

the monofunctional peptidoglycan glycosyltransferase SgtB allows *S. aureus* to grow in the

absence of LTA. We further show that peptidoglycan crosslinking is significantly reduced in

the absence of LTA, which can be partially restored upon inactivation of SgtB. This in turn

might strengthen the peptidoglycan layer and, in this way, contribute to the observed growth

rescue.

Results

Identification of *S. aureus* suppressor strains able to grow in the absence of LTA in a c-

di-AMP-independent way. LTA-deficient *S. aureus* strains are viable when grown under

osmotically stabilizing conditions in medium containing 7.5% NaCl or 40% sucrose or in un-

supplemented medium after the acquisition of compensatory mutations (31, 34). The majority

of compensatory mutations previously observed were in *gdpP* leading to inactivation of the c-

di-AMP phosphodiesterase GdpP (34). In the same study, suppressor strains with mutations

outside the *gdpP* gene were noted (34). In order to characterize suppressor strains with

mutations in genes other than *gdpP* and to gain further insight into the cellular function of LTA,

a larger suppressor screen was performed and the *ltaS* mutant strains LAC*Δ*ltaS::erm*

constructed in medium supplemented either with sucrose or NaCl were plated on un-

supplemented TSA plates. A number of independently obtained suppressor colonies were

subsequently passed 4 times in fresh TSB to further improve their growth. Next, the

chromosomal DNA was isolated from 80 suppressor strains and those lacking mutations in

gdpP (coding for the c-di-AMP hydrolase) and *dacA* (coding for the c-di-AMP cyclase

enzyme) were identified by determining the sequences of these two genes. Of 80 colonies

screened, none had mutations in *dacA* and 17 strains had no mutation in *gdpP*, 7 of which were

selected for further analysis. The absence of LTA but presence of WTA in the suppressor

strains was confirmed by western blot and Alcian blue silver staining analysis, respectively

(Figs. 1A and 1B). Next, the relative cellular c-di-AMP levels in the different strains were

determined using a previously described competitive ELISA assay (35, 36) (Fig. 1C). In

contrast to the *gdpP* mutant or LTA negative suppressor strain with a mutation in *gdpP*, which

showed the expected increase in c-di-AMP levels, the 7 suppressor strains chosen for further

analysis did not show an increase in the cellular c-di-AMP concentration (Fig. 1C).

Identification of genomic alterations in the *S. aureus* suppressor strains able to grow in

the absence of LTA. Next, the genomic sequences of the seven suppressor strains were

determined and compared to that of the original *ltaS* mutant strains using a whole genome

sequencing approach. Mutations were found in *sgtB* (SAUSA300_1855), coding for the

monofunctional glycosyltransferase SgtB and mutations in this gene arose independently in

three suppressor strains (Table 1). Another strain had a mutation in SAUSA300_1254, coding

for a hypothetical membrane protein as well as SAUSA300_RS11150, encoding MazE, the

antitoxin component of a type II toxin-antitoxin module (Table 1). In the original study by

Corrigan *et al*. (34), mutations found in SAUSA300_1254 were proposed to be accessory and

required to further improve the growth of the *gdpP* mutant suppressor strains. Consistent with

a previous report (2), a large deletion in *clpX* (SAUSA300_1621), encoding a protein forming

part of an ATP-dependent protease, was observed in one strain (Table 1). Finally, a mutation

in *vraT* (SAUSA300_1867), coding for the membrane protein VraT and forming part of the

VraRST three-component system, was identified. Using our standard genome sequence

analysis workflow, an unusually large number (>300) of zero coverage regions were obtained

for suppressor strain S6, preventing us from matching a single nucleotide polymorphism with

high confidence to the observed growth rescue. Hence, suppressor stain S6 is not listed in Table

1 and was also not further analyzed. The mutations identified in the other suppressor strains

were subsequently confirmed by fluorescent automated resequencing of the respective genomic

region. Some of the mutations observed in *sgtB*, as well as the mutations in *vraT*, *clpX* and

mazE result in frameshift mutations and introduction of premature stop codons, suggesting that

the absence of the encoded proteins compensates for the lack of LTA.

Phenotypic characterization of the LTA negative *S. aureus* suppressor strains. In a

previous study, it was shown that *S. aureus* cells can grow without LTA in the absence of ClpX

(2), hence we did not further characterize the *clpX* mutant suppressor strain obtained as part of

this study. Instead our further analysis focused on suppressor strains with mutations in novel

genes, which are strains with mutations in *mazE*, *sgtB* or *vraT*. MazE, the antitoxin component

of a type II TA module, is part of the *sigB* regulon and has been shown to be essential for full

activity of the alternative sigma factor SigB (37). SigB and members of its regulon enable

bacteria to respond rapidly to environmental and antibiotic stresses and also play a role in cell

envelope homeostasis (38, 39). Various studies have investigated the role of the VraTSR three-

component regulatory system and this system has been reported to be involved in the induction

of the cell wall stressosome, mainly in the presence of cell wall targeting antibiotics (40-42).

Interestingly, the monofunctional glycosyltransferase SgtB, also identified as part of our

screen, is one of the proteins belonging to the cell wall stressosome whose expression is

regulated by VraTSR independent of the presence of cell wall targeting antibiotics (40). To

further characterize the suppressor strains, we first confirmed the growth improvement of

strains S2-*mazE*, S4-*sgtB* and S7-*vraT* when propagated in TSB. All three suppressor strains

grew better compared to the original *ltaS* mutant strain and their growth rate was only slightly

reduced compared to the WT LAC* stain (Fig. 2A). Next, the cell morphology of the WT,

original *ltaS* mutant and the three suppressor strains was assessed by microscopy following

staining of the peptidoglycan with fluorescently labelled vancomycin. The cell morphology of

the suppressor strains was considerably improved and in particular for suppressor strain S2-

mazE the division site was correctly placed in most cells (Fig. 2B). Next, the susceptibility of

the suppressor strains to a number of cell wall targeting antibiotics was determined. Although

the growth and cell morphology of the suppressor strains were improved, these strains

remained hyper-sensitive to the β-lactam antibiotic oxacillin and the MIC was reduced by 32-

fold or more for the different suppressor strains as compared to the WT LAC* strain (Fig. 3A).

The susceptibility to the cell wall or membrane targeting antibiotics lysostaphin, nisin,

vancomycin or daptomycin was also tested but no drastic differences were observed (Table 2).

Finally, the growth of the three different suppressor strains was assessed on plates containing

Congo red. Congo red is an anionic azo dye traditionally used for the detection of biofilms in

Staphylococcus. However, at higher concentrations it inhibits the growth of *S. aureus* and has

been used in the past to indicate differences in the cell wall integrity of different *S. aureus*

strains (43). Recently, the target of Congo red has been established as the LTA synthase

enzyme LtaS (44). Hence our suppressor strains, which are deleted for *ltaS* and are able to grow

in the absence of LTA, should be resistant to this dye. To test the susceptibility of the

suppressor strains, serial dilutions of overnight cultures were spotted on TSA plates containing

0.1% Congo red (Fig. 2C). The suppressor strains carrying mutations in *mazE* and *sgtB* were

significantly more resistant to Congo red than the WT strain, indicating that inactivation of

either one of these genes is indeed sufficient to bypass the LTA essentiality. However, the

suppressor strain with the mutation in *vraT* grew poorly on the Congo red plates, suggesting

that this strain might not be a bona fide suppressor strain.

Introduction of SgtB or MazE in the respective suppressor strain results in the expected

growth arrest. The result of the whole genome sequencing analysis suggested that inactivation

of MazE or SgtB is sufficient to allow *S. aureus* to grow in the absence of LTA. Introduction

of a wild-type copy of *mazE* or *sgtB* into the respective suppressor strain should restore this

phenotype and be lethal for the suppressor strains when grown in un-supplemented medium

but should not have an effect when the bacteria are propagated in medium supplemented with

40% sucrose. In order to test this, plasmids piTET-*mazE* or piTET-*sgtB,* allowing for

anhydrotetracycline (Atet) inducible gene expression, were introduced into the respective

suppressor strains. As control, these plasmids were also introduced in the WT LAC* strain.

Serial dilutions of these different strains were spotted onto TSA plates containing 100 ng/ml

or 200 ng/ml Atet for the expression of *mazE* or *sgtB* expression, respectively. As expected for

a successful complementation, the expression of *mazE* prevented the growth of the suppressor

strain S2-*mazE* (Fig. 4A) and the expression of *sgtB* prevented the growth of the suppressor

strain S4-*sgtB* (Fig. 4B) on TSA but had no effect when the bacteria were spotted on medium

supplemented with 40% sucrose (Fig. 5). Of note, the *mazE* suppressor strain S2-*mazE*

contained an additional mutation in a gene coding for a membrane protein of unknown function

with locus tag SAUSA300_125. Introduction of a WT copy of this gene in the suppressor strain

did not prevent the growth of this suppressor strain (data not shown). Taken together, the results

of this complementation analysis support the notion that inactivation of SgtB or MazE is

sufficient to allow *S. aureus* to grow in the absence of LTA.

Growth characterization and antibiotic resistance of an *S. aureus* LAC* *sgtB* mutant.

Inactivation of MazE likely has pleotropic effects due to its involvement in the activity of the

alternative sigma factor SigB and the reason as to why its inactivation allows *S. aureus* to grow

in the absence of LTA could be indirect. Hence, we next focused in trying to gain a better

understanding of the cellular function of the monofunctional peptidoglycan glycosyltransferase

SgtB and how its inactivation allows *S. aureus* to survive in the absence of LTA. To this end,

the *sgtB* mutant strain LAC**sgtB::tn* was constructed by moving the genomic region with a

transposon insertion in *sgtB* from the Nebraska transposon mutant library strain NE596 (45)

into the LAC* background. SgtB is one of two monofunctional peptidoglycan

glycosyltransferases encoded in the *S. aureus* genome and the protein can polymerize

peptidoglycan glycan chains *in vitro* (46). Although it is dispensable for the growth of *S.*

aureus, SgtB becomes necessary for bacterial survival in the absence of the main

glycosyltransferase PBP2 (12, 15, 47). Consistent with these previous observations, no

difference was observed in the cell growth and morphology of the *sgtB* mutant *S. aureus* strain

LAC**sgtB::tn* as compared to the WT LAC* strain (Fig. 6A and 6B). Our results indicate that

inactivation of SgtB allows *S. aureus* to survive in the absence of LTA, therefore the *sgtB*

mutant strain should no longer be sensitive to the Congo red dye. To test this, serial dilutions

of the WT, the *sgtB* mutant and a complementation strain were spotted on TSA plates

containing 0.1% Congo red, which inhibits the LtaS enzyme. Indeed, the *sgtB* mutant strain

was considerably more resistant to Congo red than the WT strain and this phenotype could be

complemented by introducing a functional copy of *sgtB* into the mutant strain (Fig. 6C). Next,

we also tested the susceptibility of the *sgtB* mutant and complementation strain towards the

cell wall active antibiotics oxacillin, lysostaphin, nisin, vancomycin and daptomycin. While no

differences were observed in the susceptibility of the *sgtB* mutant compared to a wild-type

strain towards most antibiotics, a slight (4-fold) and statistically significant increase in

resistance towards oxacillin was observed (Fig. 3B, Table 2). Taken together, while no growth,

or clearly visible morphological differences were observed between the WT and the *sgtB*

mutant strain, deletion of *sgtB* leads to increased Congo red dye and 4-fold increased oxacillin

antibiotic resistance in our strain background.

Inactivation of SgtB leads to an increase in peptidoglycan cross-linking in an LTA-

negative *S. aureus* strain. Since SgtB is involved in peptidoglycan synthesis and the *sgtB*

mutant displayed increased slightly increased oxacillin resistance, we hypothesised that its

deletion could somehow "strengthen" the cell wall through alterations in the peptidoglycan

structure. In order to investigate this, the muropeptide profiles of peptidoglycan isolated from

the WT LAC*, the *sgtB* mutant strain LAC*sgtB::tn*, and the *ltaS/sgtB* mutant suppressor strain

S4-*sgtB* were determined following the growth of these strains in TSB. In addition, the

muropeptide profiles were also determined for peptidoglycan isolated from these three strains

as well as the original *ltaS* mutant strain following growth in TSB supplemented with 40%

sucrose (Fig. 7). The WT and *sgtB* mutant strains showed very similar and typical *S. aureus*

muropeptide profiles (Fig. 7A). The chemical structure of several muropeptide fragments for a

number of these peaks has been previously determined in the seminal paper by de Jonge *et al.*

(48). We numbered the different peaks where possible as described in de Jonge *et al.* (48) with

peaks 1-5 being monomeric, 9 -14 dimeric, 15 trimeric and 16 and above higher oligomeric

muropeptide fragments. Quantification of the monomeric, dimeric, trimeric and higher

oligomeric peptidoglycan fragments showed that the peptidoglycan is highly crosslinked in

both strains with approximately 70% of the UV absorbing material found in the higher

oligomeric fraction (Fig. 7B and 7C). No clear differences were found between the

muropeptide profiles of the WT and *sgtB* mutant strain, either grown in TSB or TSB

supplemented with 40% sucrose (Fig. 7). In contrast to the WT and *sgtB* mutant strain, a

decrease in the higher-oligomeric peptidoglycan material was observed for the *ltaS* suppressor

strain S4-*sgtB* both in TSB and TSB sucrose medium (Fig 7). A visual inspection of the

chromatograms also indicated that peak 12, corresponding to a currently unknown

muropeptide, was reduced in *S. aureus* strains unable to produce LTA, which are the original

ltaS mutant as well as the *S4-ltaS* suppressor strain (Fig. 7A). But perhaps most notably, a

comparison of the muropeptide profiles of the peptidoglycan isolated from the original *ltaS*

mutant and the *ltaS/sgtB* suppressor strain S4-*sgtB* after growth in TSB sucrose medium

showed that, while the amount of crosslinked peptidoglycan was reduced in both strains

compared to the WT, the peptidoglycan in the suppressor strain was more crosslinked as

compared to the original *ltaS* mutant strain (Fig 7A and 7B). Taken together, our data highlight

that deletion of *ltaS* leads to a significant reduction in the amount of crosslinked peptidoglycan

in *S. aureus*. Inactivation of SgtB in a WT strain does not significantly affect the amount of

crosslinked peptidoglycan, as one might have expected based on the observed increase in

resistance oxacillin. However, in an LTA negative background strain, inactivation of SgtB

leads to an increase in peptidoglycan crosslinking, which might explain why an LTA negative

strain can grow in the absence of this monofunctional peptidoglycan glycosyltransferase.

Discussion

The anionic LTA polymer is a core component of the cell wall, essential for survival

and a number of studies have shown its importance in various cell processes (see review (22)).

Phenotypes caused by the depletion of LTA in *S. aureus* are misplacement and incomplete

formation of cell division septa, enlargement of the cells, together ultimately leading to cell

lysis (29, 34). However, how LTA mediates these roles is still unknown. In previous work, it

has been shown that LtaS and other core LTA synthesis proteins physically interact with early

and late stage cell division proteins as well as with a number of peptidoglycan synthesis

proteins (49). This indicates that LTA synthesis enzymes might at least transiently be part of

multi-enzyme complexes which might help to coordinate LTA synthesis with peptidoglycan

synthesis and cell division (49).

We have previously shown that *S. aureus* mutants producing increased intracellular c-

di-AMP levels can survive without LTA (34). It is now believed that at high c-di-AMP levels

and through the c-di-AMP-dependent regulation of potassium and osmolyte transporters, the

internal turgor pressure in the cell might be reduced so that the compromised LTA-depleted

cell wall can sustain the internal pressure (34, 50-52). Indeed, as part of the current study we

show that the absence of LTA leads to a sizable reduction in the amount of cross-linked

peptidoglycan in *S. aureus* (Fig. 7). This is consistent with the idea that in the absence of LTA

the cell wall is likely less able to sustain the high internal turgor pressure, which might also

contribute to the increased β-lactam sensitivity of LTA-negative strains observed in this and

previous studies (Table 2) (2, 34). The reason for the reduced amount of peptidoglycan

crosslinking is currently not clear but could potentially be due to mislocalization of PBPs in

the absence of LTA or PBPs having reduced enzyme activity since the proper ion homeostasis

cannot be maintained within the cell wall in the absence of LTA.

The aim of this study was to further elucidate the role of LTA in cell wall assembly and

potentially uncover additional proteins involved in the maintenance of cell wall integrity. A

suppressor screen followed by whole genome sequencing revealed mutations in genes coding

for ClpX, SgtB, MazE and VraT (Table 1) that can bypass the essentiality of LTA. Further

experimentation indicated that the strain with the mutation in *vraT* might not be a bona fide

suppressor strain. On the other hand, complementation analysis and growth assays on agar

plates containing the azo dye Congo red that prevents the growth of *S. aureus* by inhibiting the

LtaS enzyme (44) confirmed that inactivating mutations in *sgtB* and *mazE* could bypass the

essentiality of LTA. In a previous study it was found that an *S. aureus clpX* mutant readily

acquires inactivating mutations in *ltaS* resulting in the generation of LTA negative strains (2).

Conversely and consistent with this previous work, in this study we found in one of our LTA-

negative suppressor strains a large deletion in *clpX* (Table 1). ClpX is a protein folding

chaperon, which recognizes and targets proteins for degradation to the ClpP protease

component. In the absence of ClpX, *S. aureus* cells become smaller, show increased production

of autolysins and bacteria have a severe growth defect at temperatures of 30°C or lower (2).

The introduction of loss of function mutations in the gene coding for the LTA synthase LtaS

in a *clpX* mutant alleviates some of these effects. This is perhaps due to LTA depletion having

the opposite effect that leads to an increase in cell size and decreased in autolysis as reported

in some previous publications (2, 31).

Mutations in the monofunctional glycosyltransferase SgtB were the most prevalent

mutations that arose in our suppressor screen. Previous studies have shown that SgtB is not

essential for the survival of the cell and as reported in this and a previous study, an *sgtB* mutant

strain does not show any obvious growth or morphological differences as compared to a WT

strain under standard growth conditions (Fig. 6) (15). Interestingly, we found that the *sgtB*

mutant strain LAC*sgtB::tn* was 4-fold more resistant to the cell-wall targeting antibiotic

oxacillin and this phenotype could be complemented by introduction of a wild-type copy of

sgtB (Table 2). These results are in accordance with previous reports, where strains with

mutations in genes that compensate for the lack of LTA, such as *gdpP* and *clpX* mutant strains,

also show increased oxacillin resistance (2, 34).

The increased resistance to cell wall-targeting antibiotics of the *sgtB* mutant prompted

us to investigate the peptidoglycan structure of an *sgtB* mutant in more detail, as we

hypothesized that in its absence changes such as an increased crosslinking might be observed

that could potentially explain the increased resistance. However, we could not detect any

obvious changes in the muropeptide profile of an *sgtB* mutant strain as compared to a WT strain

(Fig. 7). On the other hand, the original *ltaS* mutant strain showed a significant reduction of

around 30% in the higher oligomeric crosslinked peptidoglycan (Fig. 7). But perhaps most

importantly, the peptidoglycan isolated from the *sgtB/ltaS* double mutant S4-*sgtB* suppressor

strain showed an increase in peptidoglycan crosslinking as compared to the original *ltaS* mutant

strain (Fig. 7). We speculate that this increase in peptidoglycan cross-linking could potentially

strengthen the cell wall to better sustain the high internal turgor pressure and be at least in part

responsible for the observed growth improvement.

Bacterial two-hybrid studies have indicated that the *S. aureus* SgtB protein interacts

with SgtA, PBP1, PBP2 and PBP2A (15). Therefore, SgtB and the main glycosyltransferase

PBP2 might compete for substrate during peptidoglycan biosynthesis and inactivation of the

former might increase the substrate availability and activity of the latter. This might in turn aid

in the delivery of substrate to the transpeptidase domain of the bifunctional PBP2 enzyme

resulting in increased crosslinking. Alternatively, SgtB could also affect the function of PBP4,

which has been shown to be responsible for hyper-crosslinking of the staphylococcal

peptidoglycan (53). Clearly, more experiments are needed to clarify the complex nature of

peptidoglycan biosynthesis in *S. aureus* and changes observed in the absence of LTA.

Mutations in MazE could potentially also be tied in with changes in the cell wall

structure of *S. aureus*. The MazEF type II toxin-anitoxin system is part of the *sigB* regulon and

has been shown to be required for full activity of the alternative sigma factor SigB (37). It has

also been reported that overexpression of σ^B causes cell wall thickening in *S. aureus* and

increased resistance to cell wall-targeting antibiotics (38). Therefore, inactivation of MazE, as

observed in one of our suppressor strains, could potentially also affect cell wall homeostasis

via its effect on SigB.

In summary, our results suggest that in the absence of LTA, peptidoglycan in *S. aureus*

becomes less cross-linked, which might weaken the cell wall and become less able to sustain

the high internal turgor pressure, ultimately leading to cell lysis. The suppressor mutations

obtained in *sgtB* (and perhaps also in some of the other genes observed in our screen) help the

cell survive this detrimental effect by altering and strengthening the cell wall, likely allowing

the cell wall to again better withstand the high internal turgor pressure.

Materials and Methods

Bacterial strains and culture conditions. All bacterial strains used in this study are listed in

Table 3. *Escherichia coli* strains were cultured in Luria Bertani (LB) medium and

Staphylococcus aureus strains in tryptic soya broth (TSB) at 37°C with aeration, unless

otherwise stated. When appropriate the growth medium was supplemented with antibiotics and

inducers as follow: for *E. coli* cultures: ampicillin (Amp) 100 μg/ml, chloramphenicol (Cam)

10 μg/ml, and kanamycin (Kan) 30 μg/ml; for *S. aureus* cultures: chloramphenicol (Cam) 10

μg/ml for plasmid selection and 7.5 μg/ml for chromosomally integrated plasmid selection,

erythromycin (Erm) 10 μg/ml and kanamycin (Kan) 90 μg/ml. The inducer anhydrotetracycline

(Atet) was used at a concentration of 100 or 200 ng/ml in agar plates and 50 ng/ml in broth.

Plasmid and strain construction. Strains and primers used in this study are listed in Tables 3

and 4, respectively. Strain LAC**sgtB::tn* was generated by transduction using phage Φ85 and

transducing the *sgtB* region containing a transposon insertion in *sgtB* from the Nebraska

transposon mutant library strain NE596 (45) into strain LAC*. Plasmids piTET-*sgtB* and

piTET-*mazE* for anhydrotetracycline (Atet) inducible expression of *sgtB* and *mazE* in *S. aureus*

were generated by amplifying *sgtB* or *mazE* from LAC* chromosomal DNA using primers pair

ANG2268/ANG2269 or ANG2743/ANG2744, respectively. The PCR products and plasmid

piTET were digested with AvrII and SacII, ligated and then transformed into *E. coli* XL1-Blue

yielding strain XL1-Blue piTET-*sgtB* (ANG3926) and XL1-Blue piTET-*mazE* (ANG4727).

The plasmids were shuttled through *E. coli* strain IM08B (ANG3927 and ANG4728) and

subsequently electroporated into LAC* yielding strains LAC* piTET-*sgtB* (ANG4056) and

LAC* piTET-*mazE* (ANG4729) or into the respective suppressor strains yielding strains S4-

sgtB piTET-*sgtB* (ANG4058) and S2-*mazE* piTET-*mazE* (ANG4730). The plasmid pCL55-

sgtB for complementation and expression of *sgtB* from its native promoter was generated by

amplifying the *sgtB* gene including its native promoter region from LAC* chromosomal DNA

using primers ANG2270 and ANG2271. The PCR product and plasmid pCL55 were digested

with EcoRI and BamHI, ligated and then transformed into *E. coli* XL1-Blue yielding strain

XL1-Blue pCL55-*sgtB* (ANG4708), shuttled through IM08B (ANG3923) and subsequently

electroporated into the *sgtB* mutant strain yielding strain LAC**sgtB::tn* pCL55-*sgtB*

(ANG4711). WT LAC* or mutant *S. aureus* strains with the empty vectors piTET or pCL55

(see Table 3) were used as control strains in several experiments. The sequences of all plasmid

inserts were verified by automated fluorescence sequencing at GATC Biotechnology.

Bacterial growth curves. *S. aureus* LAC* and the indicated *ltaS* suppressor strains were

grown overnight in TSB medium containing the relevant antibiotic. The original *ltaS* mutant

strains were grown in TSB containing either 7.5% NaCl or 40% sucrose. Overnight cultures

were washed three times in TSB and diluted to a starting OD_{600} of 0.05. Cultures were

incubated at 37°C with aeration and OD_{600} values determined at 2 h intervals. WT LAC* and

the *sgtB* mutant strain LAC**sgtB::tn* were grown overnight in TSB or TSB supplemented with

10 µg/ml Erm. The next day, bacteria were washed three times in TSB and diluted to a starting

OD$_{600}$ of 0.05 and 200 μl placed into a 96-well microtiter plate. Growth was monitored for 10h

using a SPECTROSTARnano plate reader (BMG Labtech). All growth curves were performed

in triplicates with average values and standard deviations plotted.

Fluorescence microscopy analysis. Cells from 1 ml overnight culture were collected by

centrifugation, washed 3 times in PBS pH 7.4 and 150 μl of these cell suspensions applied to

polylysine (0.1%, w/v) treated coverslips. Bacteria-coated coverslips were incubated with 100

μl of a 1 μg/ml BODIPY-vancomycin (Molecular Probes) solution in H$_2$O for 20 min, washed

and mounted on glass slides containing 20 μl Vectashield (Vector Laboratories). Slides were

viewed under a Zeiss Axio Imager A2, using a GFP filter set and images were captured with

an AxioCam MRc Rev.3 camera and analyzed using the Zen Pro 2012 SP2 software. The

experiment was performed in triplicate.

Minimum Inhibitory Concentrations (MIC) determination. Overnight cultures of WT, the

sgtB mutant and complementation stain as well as the indicated suppressor strains were grown

over night in TSB medium. The next day, cultures were adjusted to an OD$_{600}$ of 0.05 in TSB

and 100 μl of these suspensions were incubated in 96-well plates with 2-fold dilutions of

various antimicrobials at the following starting concentrations: oxacillin 500 μg/ml or 1 μg/ml

as appropriate, daptomycin 32 μg/ml, lysostaphin 2 μg/ml, vancomycin 32 μg/ml and nisin 64

μg/ml. Oxacillin and daptomycin containing wells were supplemented with 2% (w/v) NaCl and

0.23 mM CaCl$_2$, respectively. Plates were incubated at 37°C for 24 hours with shaking at 500

rpm. MICs were determined as the concentration of antibiotic at which growth was inhibited

by >90% compared to growth without the antibiotic.

LTA detection by western blot. LTA extraction and detection by western blot was performed

as previously described (54). Briefly, samples were prepared from 1 ml overnight cultures

normalized based on OD_{600} readings, that is cells from 1 ml culture with an OD_{600} of 6 were

suspended in 90 µl 2 × SDS protein sample buffer. 10 µl of these samples were separated on

15% SDS-polyacrylamide gels and the material subsequently transferred to PVDF membranes.

LTA was detected using the monoclonal polyglycerolphosphate-specific LTA antibody

(Hycult Biotechnology) at a 1:4,000 dilution and the HRP-conjugated anti-mouse IgG antibody

(Cell Signaling Technologies, USA) at a 1:10,000 dilution. The blots were developed by

enhanced chemiluminescence using ClarityTM Western ECL Blotting Substrate (Bio-Rad) and

imaged using the ChemiDoc Touch Imaging System (Bio-Rad). Western blots were performed

in triplicates and a representative result is shown.

Detection of WTA by Alcian blue silver staining. Flasks with 60 ml TSB medium

supplemented with the appropriate antibiotics were inoculated with single colonies of WT

LAC* and different suppressor strains and the cultures incubated overnight (18 h) with shaking

at 37°C. The original *ltaS* mutant was grown in TSB supplemented with 40% sucrose and

incubated 6-8 hours longer. Cells from an OD_{600} equivalent of 120 were harvested by

centrifuging and the bacterial pellet stored at -20°C for further processing. Wall teichoic acid

extraction and detection by Alcian blue silver staining following SDS PAGE analysis was

performed as described in Covas *et al.* (55). Briefly, cells were washed in 20 ml buffer 1 (50

mM 2-(N-morpholino)ethanesulfonic acid (MES), pH 6.5), resuspended in 20 ml buffer 2

(buffer 1 with 4% (w/v) SDS) and boiled for 60 min. Next, the cells were washed with 20 ml

buffer 2 and after transfer to 2.0 ml reaction tubes washed once more with 1.6 ml buffer 2, 1.6

ml buffer 3 (buffer 1 with 2% (w/v) NaCl) and finally with 1.6 ml buffer 1. The samples were

suspended in 1.6 ml buffer 4 (20 mM Tris–HCl, pH 8.0, 0.5 % (w/v) SDS) and incubated for

4 h at 50°C with shaking following the addition of 2 µl of Proteinase K solution (20 mg /ml)

from *Tritirachium album*. Next, the cells were collected by centrifugation, washed once with

1.6 ml buffer 3 and three times with 1.6 ml of H_2O. To release the WTA, the pellets were

suspended in 1 ml of 0.1 mM NaOH and the samples incubated for 16 h at 25°C. Next, the

samples were centrifuged and 30 µl of the supernatant containing WTA separated on native

Tris/Tricine polyacrylamide gels and the WTA visualized by Alcian blue silver staining (55).

Whole genome sequencing. WT LAC*, LAC*Δ*ltaS*$_N$, LAC*Δ*ltaS*$_S$, and the indicated

suppressor strains were cultured overnight at 37°C, cells harvested and genomic DNA

extracted. Genome sequencing was performed by MicrobesNG (University of Birmingham)

using an Illumina platform and a 250 bp paired end read kit. Sequence analysis was performed

using the CLC Genomics work bench software package. First the LAC* reads were aligned

against the published USA300 FPR3757 genome sequence (RefSeq accession number

NC_007793.1), assembled into a reference contig and the USA300 FPR3757 annotation

transferred onto the LAC* sequence. Next, the Illumina reads for the original *ltaS* mutant

strains LAC*Δ*ltaS*$_S$ and LAC*Δ*ltaS*$_N$ as well as the seven different suppressor strains were

mapped onto the assembled LAC* sequence and high frequency (> 65%) and good quality base

changes identified using the CLC Genomics Workbench software package. Genomic

alternations found in the suppressor strains but not present in the original *ltaS* mutant strain are

summarized in Table 1. The Illumina short read for the WT LAC* has been previously

published (35) and has been deposited in the European Nucleotide Archive under the study

accession number PRJEB14759. The Illumina reads for the *ltaS* mutants and *ltaS* suppressor

strains were deposited in the European Nucleotide Archive under the study accession number

PRJEB14723.

Peptidoglycan isolation and analysis. Overnight cultures of *S. aureus* strains LAC*, S4-*sgtB* and LAC*sgtB::tn* were prepared in TSB or TSB supplemented with 40% sucrose and of the original *ltaS* mutant strain LAC*$\Delta ltaS_N$ (ANG2134) in TSB supplemented with 40% sucrose. The next day, cells were back-diluted in 2 litres of the same growth medium to an OD_{600} of 0.05. The cultures were grown at 37°C until they reached an OD_{600} of approximately 1.5, cooled on ice and cells were collected by centrifugation. Peptidoglycan was purified and digested with mutanolysin as previously described (34, 48). HPLC analysis of the digested peptidoglycan material was performed as described previously (48) and the muropeptide profiles were determined for each strain and growth condition in triplicates. For the quantification of monomeric, dimeric, trimeric and higher oligomeric peptidoglycan material, the peaks were integrated. The total peak area for each muropetide profile was determined and set to 100% and the % monomeric, dimeric, trimeric, and higher oligomeric peaks calculated and the average values and standard deviation from the three profiles determined and plotted.

c-di-AMP quantification by competitive ELISA. Five ml TSB were inoculated with a single colony of *S. aureus* LAC* (WT), strain LAC* *gdpP::kan,* US3 (LAC*$\Delta ltaS_S$::erm* suppressor with mutation in *gdpP*) as well as the different LTA-negative suppressor and the tubes were incubated for 8 hours at 37°C. Next, the cultures were back-diluted to an OD_{600} of 0.05 in 10 ml TSB and grown for 15 hours at 37°C. Bacterial cells from these cultures were collected by centrifugation, cell lysates prepared and the cellular c-di-AMP levels determined using a previously described competitive ELISA method (35, 36). However, in place of determining c-di-AMP levels based on a standard curve the A_{490} values were directly compared. To this end, the A_{490} reading obtained for samples derived for the WT strain was set to 100% and the % values calculated for the samples derived from the other strains. Thee independent experiments were performed with three technical replicates and the average values and

standard deviation of the % A_{490} values determined for each strain as compared to the WT

strain. Of note, since this is a competitive ELISA, and a decrease in A_{490} readings represents

an increase in cellular c-di-AMP levels.

Acknowledgments

This research was supported by the European Research Council grant 260371 and the

Wellcome Trust grant 100289 to AG, the UK Medical Research Council (MRC) within the

AMR Cross-council initiative Collaborative Grant MR/N002679/1 to WV and the German

Research Foundation [Deutsche Forschungsgemeinschaft (DFG)] grant SCHU 3159/1-1 to

CFS. The Illumina sequencing was performed by MicrobesNG, which was supported by the

BBSRC grant BB/L024209/1.

Table 1: Genomic variations detected in the suppressor strains as compared the original
ltaS **mutant strains**
560

Strain[1]	Ref. pos.[2]	Type[3]	Ref.[4]	Allele[5]	Freq.[6]	Av. Qual.[7]	Annotations	Amino acid change[8]
ANG3711 S1-*sgtB*	2018191	SVN	T	A	100	38.5	SAUSA300_1855, monofunctional glycosyltransferase SgtB	Asn11Tyr
	2017952-2018187	DEL					SAUSA300_1855, monofunctional glycosyltransferase SgtB	
ANG3712 S2-*mazE*	1380637	INS	-	G	100	38.5	SAUSA300_1254, hypothetical membrane protein	Ile291FS
	1380639	SNV	C	A	100	38.7		Ser292Tyr
	1380641	REP	C	AG	100	38.4		Pro293fs
	2188437-2188446	DEL	TATCGGAAAA	----------			SAUSA300_RS11150, Antitoxin component of a type II TA module mazE	
ANG3713 S3-*clpX*	1775491	SNV	T	A	100	38.8	SAUSA300_1621, ATP-dependent protease ATP-binding subunit ClpX	Glu148Val
	1775495	SNV	T	C	100	38.9		Thr147Ala
	1775668	SNV	A	C	100	38.2		Val89Gly
	1775500-1775663	DEL						
ANG3714 S4-*sgtlB*	2017704	SNV	A	C	100	38	SAUSA300_1855, monofunctional glycosyltransferase SgtB	Leu173*
ANG3716 S6		N/A	N/A	N/A	N/A	N/A	N/A	N/A
ANG3717 S7-*vraT*	2027736	SNV	A	C	100	38.8	SAUSA300_1867, membrane protein Yvfq/VraT	Leu71*
ANG3718 S8-*sgtB*	2018107	INS	-	T	92.6	38.6	SAUSA300_1855, monofunctional glycosyltransferase SgtB	Ser39fs

[1]Suppressors S1, S2, S7 and S8 were derived from strain LAC*$\Delta ltaS_S$ (ANG2135) and

suppressor strains S3, S4 and S6 were derived from strain LAC*$\Delta ltaS_N$ (ANG2134).

[2]Reference position indicates base number in the LAC* reference genome

(USA300_FPR3757, NCBI CP000255.1). [3]Type of mutation with DEL indicating deletion,

SNV indicating single nucleotide variation, INS indicating insertion and REP indicating

replacement. [4]Ref. indicates base in the LAC* reference genome. [5]Allele indicates base in the

sequenced strain. [6]Freq. indicates frequency of the specific variant found in the sample reads.

[7]Av. Qual. denotes the average base quality score at the indicated position. [8]Amino acid change

in the encoded protein in LAC* versus the sequenced strain with fs indicating a frame shift and

* a premature stop codon.

Table 2: MIC values for different antibiotics given in µg/ml

	lysostaphin	nisin	vancomycin	daptomycin
LAC*	**0.125** (0.125, 0.125, 0.125)	**16** (16, 16, 16)	**4** (4, 4, 4)	**16** (16, 16, 16)
S2-*mazE*	**0.125** (0.125, 0.125, 0.125)	**2-4** (2, 2, 4)	**2** (2, 2, 2)	**4-8** (8, 8, 4)
S4-*sgtB*	**0.125-0.250** (0.125, 0.125, 0.25)	**4** (4, 4, 4)	**1** (1, 1, 1)	**8-16** (16, 8, 8)
S7-*vraT*	**0.500** (0.5, 0.5, 0.5)	**2-4** (2, 2, 4)	**4-8** (8, 4, 4)	**8** (8, 8, 8)
sgtB::tn	**0.125-0.250** (0.25, 0.125, 0.125)	**16-32** (32, 16, 32)	**4** (4, 4, 4)	**16** (16, 16, 16)
sgtB::tn **pCL55**	**0.125-0.250** (0.125, 0.125, 0.25)	**8-16** (16, 8, 16)	**4** (4, 4, 4)	**16** (16, 16, 16)
sgtB::tn **pCL55-*sgtB***	**0.125** (0.125, 0.125, 0.125)	**8-16** (8, 16, 16)	**4-8** (4, 8, 4)	**16** (16, 16, 16)

574 The MIC was defined as antibiotic concentration that leads to >90% growth inhibition as

575 compared to growth without antibiotic. MICs were determined in triplicate. The obtained MIC

576 range is shown in bold and individual MIC values of the replicate samples are shown below in

577 parenthesis.

579   **Table 3: Bacterial strains used in this study**

Strain name	Description	Reference
	***Escherichia coli* strains**	
XL1-Blue	Cloning strain; TetR - ANG127	Stratagene
IM08B	*E. coli – S. aureus* sutttle strain - ANG3723	(56)
ANG3732	IM08B pCL55; AmpR	(52)
ANG3926	XL1-Blue piTET-*sgtB*; AmpR	This study
ANG3927	IM08B piTET-*sgtB*; AmpR	This study
ANG3928	IM08B piTET; AmpR	(57)
ANG4708	XL1-Blue pCL55-*sgtB*; AmpR	This study
ANG4707	IM08B pCL55-*sgtB*; AmpR	This study
ANG4727	XL1-Blue piTET-*mazE*; AmpR	This study
ANG4728	IM08B piTET-*mazE*; AmpR	This study
	***Staphylococcus aureus* strains**	
LAC*	LAC*; Erm sensitive CA-MRSA LAC strain; ANG1575	(58)
NE596	JE2*sgtB::tn*; NARSA library transposon mutant; ErmR – ANG3930	(45)
ANG1961	LAC*gdpP::kan,* KanR	(34)
ANG2134	LAC*Δ*ltaS_N*::erm pCN38; Isolated on 7.5% NaCl; CamR ErmR	(34)
ANG2135	LAC*Δ*ltaS_S*::erm pCN38; Isolated on 40% Sucrose; CamR ErmR	(34)
ANG2137	LAC*Δ*ltaS_N*::erm suppressor S6 original colony; ErmR	(34)
ANG2140	LAC*Δ*ltaS_S*::erm suppressor S7 original colony; ErmR	(34)
ANG2143	LAC*Δ*ltaS_S*::erm suppressor S8 original colony; ErmR	(34)
ANG2434	LAC*Δ*ltaS_S*::erm suppressor with *gdpP* mutation (pass 4) (short: US3); ErmR	(34)
ANG3694	LAC*Δ*ltaSs*::erm suppressor S1 original colony; ErmR	This study
ANG3698	LAC*Δ*ltaSs*::erm suppressor S2 original colony; ErmR	This study
ANG3703	LAC*Δ*ltaSs*::erm suppressor S3, original colony; ErmR	This study
ANG3707	LAC*Δ*ltaSs*::erm suppressor S4, original colony; ErmR	This study
ANG3711	LAC*Δ*ltaS_S*::erm suppressor S1 pass 4 (short: S1-*sgtB*); ErmR	This study
ANG3712	LAC*Δ*ltaS_S*::erm suppressor S2 pass 4 (short: S2-*mazE*); ErmR	This study
ANG3713	LAC*Δ*ltaS_N*::erm suppressor S3 pass 4 (short: S3-*clpX*); ErmR	This study
ANG3714	LAC*Δ*ltaS_N*::erm suppressor S4 pass 4 (short: S4-*sgtB*); ErmR	This study
ANG3716	LAC*Δ*ltaS_N*::erm suppressor S6 pass 4 (short: S6); ErmR	This study
ANG3717	LAC*Δ*ltaS_S*::erm suppressor S7 pass 4 (short: S7-*vraT*); ErmR	This study
ANG3718	LAC*Δ*ltaS_S*::erm suppressor S8 pass 4 (short: S8-*sgtB*); ErmR	This study
ANG4054	LAC* piTET, CamR	(57)
ANG4056	LAC* piTET-*sgtB*; CamR	This study

ANG4057	S4-*sgtB* piTET; ErmR CamR	This study
ANG4058	S4-*sgtB* piTET-*sgtB*; ErmR CamR	This study
ANG4059	LAC**sgtB::tn*; ErmR	This study
ANG4710	LAC**sgtB::tn* pCL55; ErmR CamR	This study
ANG4711	LAC**sgtB::tn* pCL55-*sgtB*; ErmR CamR	This study
ANG4725	S2-*mazE* piTET; ErmR CamR	This study
ANG4729	LAC* piTET-*mazE*; CamR	This study
ANG4730	S2-*mazE* piTET-*mazE*; ErmR CamR	This study

581 **Table 4: Primers used in this study**

Primer ID	Name	Sequence
ANG2229	sausa300_1867 Fw seq	GAAGCATTTCAAATTCAATGTGCAAG
ANG2230	sausa300_1867 Rv seq	GAAGCATTTCAAATTCAATGTGCAAG
ANG2231	sgtB Fw seq	GTCGTCTCTATTGGCATTTAATAGGG
ANG2232	sgtB Rv seq	AATAATGAAGTACTAATCAAGTGGC
ANG2233	sausa300_1254 Fw seq	CATTTTGAAGAAAGGATTAATCAA
ANG2234	sausa300_1254 Rv seq	AATAATGAAGTACTAATCAAGTGGC
ANG2235	clpX Fw seq	GAAGGAACTAAAGAAGATTAATCTTC
ANG2236	clpX Rv seq	CTGGATATTGTTCTTCTTTTACTGCAC
ANG2268	5-AvrII-sgtB-pCL55iTet	CATGCCTAGGTTAAAAGAAGGAGCAAACGCATG
ANG2269	3-SacII-sgtB-pCL55iTet	CATGCCGCGGTTAACGATTTAATTGTGACATAGCC
ANG2409	5- EcoRI-promSgtB-pCL55	CATGGAATTCAAGTATTGTGGTTATCGATTG
ANG2410	3- BamH1-promSgtB-pCL55	CATGGGATCCTTAACGATTTAATTGTGACATAGCC
ANG2740	5-mazE SNP seq	AGATAATCATAGAGAAAGTCCACAGTCG
ANG2741	3-mazE SNP seq	TAAGTACGTCAGTTTTTCTTTCAATCGT
ANG2743	5-AvrII-mazE-pCL55iTet	CATGCCTAGGGGAATCAATTGGAGGTTCTCATATG
ANG2744	3-SacII-mazE-pCl55iTet	CATGCCGCGGTCATTCATTCGTTGAATTAGAAGATAAATATG

582 Restriction site sequences are underlined.

  Figures and Legends

587
588
589
Figure 1: Detection of LTA by western blot, WTA by Alcian blue silver staining and c-

di-AMP using a competitive ELISA method. (A) Detection of LTA by western-blot. Cell

extracts were prepared from LAC* (WT), an *ltaS* mutant (strain ANG2135) and the 7 *ltaS*

suppressor strains (S1-S4 and S6-S8) and separated on a 15% PAA gel. LTA was subsequently

detected by western blot using a monoclonal polyglycerolphosphate specific antibody. A

representative result from 3 independent experiments is shown. (B) Visualization of wall

teichoic acid (WTA) by native PAGE. WTAs were isolated from the same strains as shown in

(A) and separated on a 20% native gel. Bands were visualized by Alcian blue and subsequently

silver stain. One representative result from 4 independent experiments is shown. (C) c-di-AMP

detection by competitive ELISA. Cytoplasmic extracts were prepared from LAC* (WT), the

high c-di-AMP level control strains LAC* *gdpP::kan* (*gdpP*) and US3 (*ltaS/gdpP*) as well as

from the 7 *ltaS* suppressor strains (S1-S4 and S6-S8). c-di-AMP amounts were determined and

compared to the WT strain by ELISA. Of note, as this is a competitive ELISA assay, lower

A_{490} readings are obtained for samples with higher c-di-AMP levels. The A_{490} reading obtained

for the sample derived from the WT strain was set to 100% and % A_{490} reading calculated for

the test strains. The average % values and standard deviations from three independent

experiments (with three technical replicates) are plotted.

607
608

Figure 2: Growth and cell morphology of WT and mutant *S. aureus* strains. (A) Bacterial growth curves. LAC* (WT) and the suppressor strains S2-*mazE,* S4-*sgtB* and S7-*vraT* were grown overnight in TSB and the original *ltaS* mutant strain (ANG2135) in TSB with 40% sucrose. Next day, bacterial cells were washed and diluted in TSB to an OD_{600} of 0.05 and the bacterial growth was subsequently monitored over a period of 10 hours. The average OD_{600} readings from three experiments were plotted. (B) Microscopic analysis. Bacterial cells from overnight cultures of the same strains as used in panel A were washed, back-diluted and grown to mid-log phase in TSB or TSB with 40% sucrose as indicated. The bacterial cells were subsequently stained with BODIPY-vancomycin and viewed by fluorescent microscopy. (C) Analysi*s* of bacterial growth on Congo red-containing TSA plates. Overnight cultures of WT

619   and the indicated suppressor strains were serially diluted and spotted on TSA plates

620   supplemented with 0.1% Congo red and the plates incubated for 48h at 37˚C. For B and C, a

621   representative result of three independent experiments is shown.

Figure 3: Oxacillin MICs for WT and mutant _S. aureus_ strains. Oxacillin minimal inhibitory concentrations (MICs) of (A) LAC* (WT) and the suppressor strains S2-*mazE,* S4-*sgtB* and S7-*vraT* and (B) LAC* (WT), the LAC* *sgtB::tn* mutant either without integrative plasmid (-), with integrative plasmid (pCL55) or integrative complementation plasmid (pCL55-*sgtB*). Experiments were conducted with six biological replicates. Medians and 95% confidence intervals for all strains are plotted. A Kruskal-Wallis one-way analysis of variance tests was performed, which indicated significant differences. Subsequently, Dunn's tests were performed and the p-values corrected for the multiple comparisons against the WT. p-values < 0.01 (**) and < 0.001 (***).

634
635
Figure 4: Growth complementation analysis. TSA plates with spot dilutions. (A) *S. aureus*

strains LAC* piTET, LAC* piTET-*mazE*, S2-*mazE* piTET or S2-*mazE* piTET-*mazE* were

grown overnight in TSB supplemented with chloramphenicol, washed twice in PBS and serial

dilutions spotted on TSA plates supplemented with 7.5 ng/ml chloramphenicol without (left

panel) or with 100 ng/ml Atet (right panel). (B) *S. aureus* strains LAC* (WT), LAC* piTET,

LAC* piTET-sgtB, S4-*sgtB*, S4-*sgtB* piTET or S4-*sgtB* piTET-*sgtB* were grown and samples

prepared as described in A, however without chloramphenicol selection, and dilutions were

spotted on TSA plates (left panel) or TSA plates supplemented with 200 ng/ml Atet (right

panel). Representative plate images from three independent experiments are shown.

Figure 5: Growth complementation analysis. TSA sucrose plates with spot dilutions. (A) *S.*

aureus strains LAC* piTET, LAC* piTET-*mazE,* S2-*mazE* piTET or S2-*mazE* piTET-*mazE*

were grown overnight in TSB supplemented with chloramphenicol, washed twice in PBS and

serial dilutions spotted on TSA 40% sucrose plates supplemented with 7.5 ng/ml

chloramphenicol without (left panel) or with 100 ng/ml Atet (right panel). (B) *S. aureus* strains

LAC* (WT), LAC* piTET, LAC* piTET-*sgtB,* S4-*sgtB,* S4-*sgtB* piTET or S4-*sgtB* piTET-

sgtB were grown and samples prepared as described in A, however without chloramphenicol

selection, and dilutions were spotted on TSA 40% sucrose plates (left panel) or TSA 40%

sucrose plates supplemented with 200 ng/ml Atet (right panel). Representative plate images

from three independent experiments are shown.

661
662

Figure 6: Growth and cell morphology of *S. aureus* strain LAC**sgtB::tn*. (A) Bacterial
growth curves. Overnight cultures of LAC* (WT) and strain LAC**sgtB::tn* (*sgtB*) were diluted
in TSB to an OD_{600} of 0.05 and the growth monitored over a period of 10 hours using a plate
reader. The average OD_{600} values and standard deviation from three independent experiments
are plotted. (B) Microscopy analysis. *S. aureus* strains LAC* (WT) and LAC**sgtB::tn* (*sgtB*)
were grown overnight in TSB or TSB supplemented with 40% sucrose, stained with BODIPY-
vancomycin and viewed under a fluorescence microscope. (C) Susceptibility to Congo red.
Overnight cultures of strains LAC* (WT) and the *sgtB* mutant strains LAC**sgtB::tn*,
LAC**sgtB::tn* pCL55 as well as the complementation strain LAC**sgtB::tn* pCL55-*sgtB* were
serially diluted and aliquots spotted on TSA plates supplemented with 0.1% Congo red. For
panels B and C, representative result from three independent experiments are shown.

674
675

Figure 7: Peptidoglycan analysis of WT and mutant *S. aureus* strains. (A) HPLC profile of mutanolysin digested peptidoglycan. Peptidoglycan was isolated and digested with mutanolysin as described in the materials and methods section from *S. aureus* strains LAC* (WT), LAC**sgtB::tn* (*sgtB*) and the LTA negative suppressor strain S4-*sgtB* following growth in TSB medium or from *S. aureus* strains LAC* (WT), LAC**sgtB::tn* (*sgtB*), the LTA negative suppressor strain S4-*sgtB* as well as the original *ltaS* mutant strain (ANG2135) following growth in TSB 40% sucrose medium. Monomeric, dimeric, trimeric and > trimeric peptidoglycan fragments are indicated above the graphs and a representative profile from three independent samples is shown. The individual muropeptide peaks were labelled where possible with the numbers as described in de Jonge *et al*. (48). (B-C) Quantification of the different peptidoglycan peaks. The peaks corresponding to monomeric, dimeric, trimeric and > trimeric

peptidoglycan fragments were integrated and quantified for the strains used in panel A. The

combined peak area was set to 100% for each strain and the average values and standard

deviations from the three independent peptidoglycan isolations were plotted in panel B, for the

strains grown in TSB sucrose medium and in panel C for the strains grown in TSB medium.

Unpaired two-tailed student's T-tests were used to determine statistically significant

differences in monomer, dimer, trimer or > trimer fractions between WT and mutant strains.

The obtained p-values were multiplied by the Bonferroni correction factors 3 (panel B) and 2

(panel C) to consider that the WT peaks were compared to three or two different strains,

respectively. Statistically significant differences are indicated by black asterisks. For panel B,

unpaired two-tailed student's T-tests were also used to determine statistically significant

differences in monomer, dimer, trimer or > trimer fractions between the *ltaS* mutant and the

other three strains and the statistically significant differences between the *ltaS* mutant and S4-

ltaS suppressor are shown with grey asterisks. p-values < 0.05 (*), < 0.01 (**) and < 0.001

(***).

REFERENCES:

1. **Kluytmans J, Van Belkum A, Verbrugh H.** 1997. Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. Clinical microbiology reviews **10:**505-520.

2. **Baek KT, Bowman L, Millership C, Dupont Sogaard M, Kaever V, Siljamaki P, Savijoki K, Varmanen P, Nyman TA, Gründling A, Frees D.** 2016. The Cell Wall Polymer Lipoteichoic Acid Becomes Nonessential in *Staphylococcus aureus* Cells Lacking the ClpX Chaperone. MBio **7**.

3. **Francis JS, Doherty MC, Lopatin U, Johnston CP, Sinha G, Ross T, Cai M, Hansel NN, Perl T, Ticehurst JR.** 2005. Severe community-onset pneumonia in healthy adults caused by methicillin-resistant *Staphylococcus aureus* carrying the Panton-Valentine leukocidin genes. Clinical Infectious Diseases **40:**100-107.

4. **Hiramatsu K, Katayama Y, Matsuo M, Sasaki T, Morimoto Y, Sekiguchi A, Baba T.** 2014. Multi-drug-resistant *Staphylococcus aureus* and future chemotherapy. Journal of Infection and Chemotherapy **20:**593-601.

5. **Horn J, Stelzner K, Rudel T, Fraunholz M.** 2017. Inside job: *Staphylococcus aureus* host-pathogen interactions. International Journal of Medical Microbiology.

6. **Foster TJ, Geoghegan JA, Ganesh VK, Höök M.** 2014. Adhesion, invasion and evasion: the many functions of the surface proteins of *Staphylococcus aureus*. Nat Rev Microbiol **12:**49-62.

7. **Silhavy TJ, Kahne D, Walker S.** 2010. The bacterial cell envelope. Cold Spring Harbor perspectives in biology **2:**a000414.

8. **Rajagopal M, Walker S.** 2015. Envelope structures of gram-positive bacteria, p 1-44, Protein and Sugar Export and Assembly in Gram-positive Bacteria. Springer.

9.   **Vollmer W, Blanot D, De Pedro MA.** 2008. Peptidoglycan structure and architecture.

FEMS microbiology reviews **32:**149-167.

10.   **Archibald A, Hancock I, Harwood C.** 1993. Cell wall structure, synthesis, and

turnover, p 381-410, *Bacillus subtilis* and other Gram-positive bacteria. American

Society of Microbiology.

11.   **Holtje JV.** 1998. Growth of the stress-bearing and shape-maintaining murein sacculus

of *Escherichia coli*. Microbiol Mol Biol Rev **62:**181-203.

12.   **Reed P, Atilano ML, Alves R, Hoiczyk E, Sher X, Reichmann NT, Pereira PM,**

Roemer T, Filipe SR, Pereira-Leal JB, Ligoxygakis P, Pinho MG. 2015.

Staphylococcus aureus Survives with a Minimal Peptidoglycan Synthesis Machine but

Sacrifices Virulence and Antibiotic Resistance. PLoS Pathog **11:**e1004891.

13.   **Hartman BJ, Tomasz A.** 1984. Low-affinity penicillin-binding protein associated

with beta-lactam resistance in *Staphylococcus aureus*. J Bacteriol **158:**513-516.

14.   **Pinho MG, de Lencastre H, Tomasz A.** 2001. An acquired and a native penicillin-

binding protein cooperate in building the cell wall of drug-resistant staphylococci. Proc

Natl Acad Sci U S A **98:**10886-10891.

15.   **Reed P, Veiga H, Jorge AM, Terrak M, Pinho MG.** 2011. Monofunctional

transglycosylases are not essential for *Staphylococcus aureus* cell wall synthesis. J

Bacteriol **193:**2549-2556.

16.   **Wang QM, Peery RB, Johnson RB, Alborn WE, Yeh WK, Skatrud PL.** 2001.

Identification and characterization of a monofunctional glycosyltransferase from

Staphylococcus aureus. J Bacteriol **183:**4779-4785.

17.   **Weidenmaier C, Peschel A.** 2008. Teichoic acids and related cell-wall glycopolymers

in Gram-positive physiology and host interactions. Nature Reviews Microbiology

6:276.

18. **Kawai Y, Marles-Wright J, Cleverley RM, Emmins R, Ishikawa S, Kuwano M,**

Heinz N, Bui NK, Hoyland CN, Ogasawara N, Lewis RJ, Vollmer W, Daniel RA,

Errington J. 2011. A widespread family of bacterial cell wall assembly proteins. Embo

j **30:**4931-4941.

19. **Kern T, Giffard M, Hediger S, Amoroso A, Giustini C, Bui NK, Joris B, Bougault**

C, Vollmer W, Simorre JP. 2010. Dynamics characterization of fully hydrated

bacterial cell walls by solid-state NMR: evidence for cooperative binding of metal ions.

J Am Chem Soc **132:**10911-10919.

20. **Neuhaus FC, Baddiley J.** 2003. A continuum of anionic charge: structures and

functions of D-alanyl-teichoic acids in gram-positive bacteria. Microbiology and

Molecular Biology Reviews **67:**686-723.

21. **Brown S, Santa Maria JP, Jr., Walker S.** 2013. Wall teichoic acids of gram-positive

bacteria. Annu Rev Microbiol **67:**313-336.

22. **Percy MG, Gründling A.** 2014. Lipoteichoic acid synthesis and function in gram-

positive bacteria. Annu Rev Microbiol **68:**81-100.

23. **Percy MG, Karinou E, Webb AJ, Gründling A.** 2016. Identification of a

Lipoteichoic Acid Glycosyltransferase Enzyme Reveals that GW-Domain-Containing

Proteins Can Be Retained in the Cell Wall of Listeria monocytogenes in the Absence

of Lipoteichoic Acid or Its Modifications. J Bacteriol **198:**2029-2042.

24. **Santa Maria JP, Jr., Sadaka A, Moussa SH, Brown S, Zhang YJ, Rubin EJ,**

Gilmore MS, Walker S. 2014. Compound-gene interaction mapping reveals distinct

roles for *Staphylococcus aureus* teichoic acids. Proc Natl Acad Sci U S A **111:**12510-

12515.

25.   **Peschel A, Otto M, Jack RW, Kalbacher H, Jung G, Götz F.** 1999. Inactivation of

the dlt Operon in *Staphylococcus aureus* Confers Sensitivity to Defensins, Protegrins,

and Other Antimicrobial Peptides. Journal of Biological Chemistry **274:**8405-8410.

26.   **Kho K, Meredith TC.** 2018. Salt-Induced Stress Stimulates a Lipoteichoic Acid-

Specific Three Component Glycosylation System in *Staphylococcus aureus.* J Bacteriol

doi:10.1128/JB.00017-18.

27.   **Reichmann NT, Gründling A.** 2011. Location, synthesis and function of glycolipids

and polyglycerolphosphate lipoteichoic acid in Gram-positive bacteria of the phylum

Firmicutes. FEMS Microbiol Lett **319:**97-105.

28.   **Lu D, Wörmann ME, Zhang X, Schneewind O, Gründling A, Freemont PS.** 2009.

Structure-based mechanism of lipoteichoic acid synthesis by *Staphylococcus aureus*

LtaS. Proceedings of the National Academy of Sciences **106:**1584-1589.

29.   **Gründling A, Schneewind O.** 2007. Synthesis of glycerol phosphate lipoteichoic acid

in *Staphylococcus aureus*. Proc Natl Acad Sci U S A **104:**8478-8483.

30.   **Karatsa-Dodgson M, Wormann ME, Gründling A.** 2010. In vitro analysis of the

Staphylococcus aureus lipoteichoic acid synthase enzyme using fluorescently labeled

lipids. J Bacteriol **192:**5341-5349.

31.   **Oku Y, Kurokawa K, Matsuo M, Yamada S, Lee BL, Sekimizu K.** 2009. Pleiotropic

roles of polyglycerolphosphate synthase of lipoteichoic acid in growth of

Staphylococcus aureus cells. J Bacteriol **191:**141-151.

32.   **Schirner K, Marles-Wright J, Lewis RJ, Errington J.** 2009. Distinct and essential

morphogenic functions for wall-and lipo-teichoic acids in *Bacillus subtilis*. The EMBO

journal **28:**830-842.

33.   **Fedtke I, Mader D, Kohler T, Moll H, Nicholson G, Biswas R, Henseler K, Götz**

F, Zähringer U, Peschel A. 2007. A *Staphylococcus aureus ypfP* mutant with strongly

reduced lipoteichoic acid (LTA) content: LTA governs bacterial surface properties and

autolysin activity. Molecular microbiology **65:**1078-1091.

34.   **Corrigan RM, Abbott JC, Burhenne H, Kaever V, Gründling A.** 2011. c-di-AMP

is a new second messenger in *Staphylococcus aureus* with a role in controlling cell size

and envelope stress. PLoS Pathog **7:**e1002217.

35.   **Bowman L, Zeden MS, Schuster CF, Kaever V, Gründling A.** 2016. New Insights

into the Cyclic Di-adenosine Monophosphate (c-di-AMP) Degradation Pathway and

the Requirement of the Cyclic Dinucleotide for Acid Stress Resistance in

Staphylococcus aureus. J Biol Chem **291:**26970-26986.

36.   **Underwood AJ, Zhang Y, Metzger DW, Bai G.** 2014. Detection of cyclic di-AMP

using a competitive ELISA with a unique pneumococcal cyclic di-AMP binding

protein. Journal of microbiological methods **107:**58-62.

37.   **Donegan NP, Cheung AL.** 2009. Regulation of the mazEF toxin-antitoxin module in

Staphylococcus aureus and its impact on sigB expression. Journal of bacteriology

191:2795-2805.

38.   **Morikawa K, Maruyama A, Inose Y, Higashide M, Hayashi H, Ohta T.** 2001.

Overexpression of sigma factor, sigma(B), urges *Staphylococcus aureus* to thicken the

cell wall and to resist beta-lactams. Biochem Biophys Res Commun **288:**385-389.

39.   **Guldimann C, Boor KJ, Wiedmann M, Guariglia-Oropeza V.** 2016. Resilience in

the face of uncertainty: sigma factor B fine-tunes gene expression to support

homeostasis in gram-positive bacteria. Applied and environmental microbiology

82:4456-4469.

40.   **Kuroda M, Kuroda H, Oshima T, Takeuchi F, Mori H, Hiramatsu K.** 2003. Two-

component system VraSR positively modulates the regulation of cell-wall biosynthesis

pathway in *Staphylococcus aureus*. Mol Microbiol **49:**807-821.

41.   **Kuroda M, Kuwahara-Arai K, Hiramatsu K.** 2000. Identification of the up- and
826   down-regulated genes in vancomycin-resistant *Staphylococcus aureus* strains Mu3 and
827   Mu50 by cDNA differential hybridization method. Biochem Biophys Res Commun
828   **269:**485-490.

829   42.   **Boyle-Vavra S, Yin S, Jo DS, Montgomery CP, Daum RS.** 2013. VraT/YvqF is
830   required for methicillin resistance and activation of the VraSR regulon in
831   *Staphylococcus aureus*. Antimicrob Agents Chemother **57:**83-95.

832   43.   **DeFrancesco AS, Masloboeva N, Syed AK, DeLoughery A, Bradshaw N, Li GW,**
833   **Gilmore MS, Walker S, Losick R.** 2017. Genome-wide screen for genes involved in
834   eDNA release during biofilm formation by *Staphylococcus aureus*. Proc Natl Acad Sci
835   U S A **114:**E5969-E5978.

836   44.   **Vickery CR, Wood BM, Morris HG, Losick R, Walker S.** 2018. Reconstitution of
837   *Staphylococcus aureus* Lipoteichoic Acid Synthase Activity Identifies Congo Red as a
838   Selective Inhibitor. J Am Chem Soc **140:**876-879.

839   45.   **Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW.**
840   2013. A genetic resource for rapid and comprehensive phenotype screening of
841   nonessential *Staphylococcus aureus* genes. MBio **4:**e00537-00512.

842   46.   **Terrak M, Nguyen-Distèche M.** 2006. Kinetic characterization of the monofunctional
843   glycosyltransferase from *Staphylococcus aureus*. Journal of bacteriology **188:**2528-
844   2532.

845   47.   **Rebets Y, Lupoli T, Qiao Y, Schirner K, Villet R, Hooper D, Kahne D, Walker S.**
846   2014. Moenomycin resistance mutations in *Staphylococcus aureus* reduce
847   peptidoglycan chain length and cause aberrant cell division. ACS Chem Biol **9:**459-
848   467.

48.   **De Jonge B, Chang Y-S, Gage D, Tomasz A.** 1992. Peptidoglycan composition of a

highly methicillin-resistant Staphylococcus aureus strain. The role of penicillin binding

protein 2A. Journal of Biological Chemistry **267:**11248-11254.

49.   **Reichmann NT, Picarra Cassona C, Monteiro JM, Bottomley AL, Corrigan RM,**

Foster SJ, Pinho MG, Gründling A. 2014. Differential localization of LTA synthesis

proteins and their interaction with the cell division machinery in *Staphylococcus*

aureus. Mol Microbiol **92:**273-286.

50.   **Commichau FM, Gibhardt J, Halbedel S, Gundlach J, Stülke J.** 2017. A Delicate

Connection: c-di-AMP Affects Cell Integrity by Controlling Osmolyte Transport.

Trends in microbiology **26:**175-185.

51.   **Corrigan RM, Campeotto I, Jeganathan T, Roelofs KG, Lee VT, Gründling A.**

2013. Systematic identification of conserved bacterial c-di-AMP receptor proteins.

Proceedings of the National Academy of Sciences **110:**9084-9089.

52.   **Schuster CF, Bellows LE, Tosi T, Campeotto I, Corrigan RM, Freemont P,**

Gründling A. 2016. The second messenger c-di-AMP inhibits the osmolyte uptake

system OpuC in *Staphylococcus aureus*. Sci Signal **9:**ra81.

53.   **Memmi G, Filipe SR, Pinho MG, Fu Z, Cheung A.** 2008. *Staphylococcus aureus*

PBP4 is essential for beta-lactam resistance in community-acquired methicillin-

resistant strains. Antimicrob Agents Chemother **52:**3955-3966.

54.   **Gründling A, Schneewind O.** 2007. Genes required for glycolipid synthesis and

lipoteichoic acid anchoring in *Staphylococcus aureus*. J Bacteriol **189:**2521-2530.

55.   **Covas G, Vaz F, Henriques G, Pinho M, Filipe S.** 2016. Analysis of Cell Wall

Teichoic Acids in *Staphylococcus aureus*. , p 201-213. *In* Hong H (ed), Bacterial Cell

Wall Homeostasis, vol 1440. Humana Press, New York, NY.

56.   **Monk IR, Tree JJ, Howden BP, Stinear TP, Foster TJ.** 2015. Complete Bypass of

874   Restriction Systems for Major *Staphylococcus aureus* Lineages. MBio **6:**e00308-

875   00315.

876   57.   **Zeden MS, Schuster CF, Bowman L, Zhong Q, Williams HD, Gründling A.** 2018.

877   Cyclic-di-adenosine monophosphate (c-di-AMP) is required for osmotic regulation in

878   *Staphylococcus aureus* but dispensable for viability in anaerobic conditions. J Biol

879   Chem doi:10.1074/jbc.M117.818716.

880   58.   **Boles BR, Thoendel M, Roth AJ, Horswill AR.** 2010. Identification of genes involved

881   in polysaccharide-independent *Staphylococcus aureus* biofilm formation. PLoS One

882   **5:**e10146.