Title: ColE1 plasmids potentiate the fitness of *Salmonella* Heidelberg in poultry litter

[¶]Adelumola Oladeinde[1*], [¶]Kimberly Cook[1], [¶]Alex Orlek[2], [&]Greg Zock[1] [&]Kyler Herrington[3],

[&]Nelson Cox[4], [&]Jodie Plumblee Lawrence[1], [&]Carolina Hall[1]

Authors' Affiliation

[1]Bacterial Epidemiology and Antimicrobial Resistance Research Unit, U.S. National Poultry

Research Center, USDA-ARS, Athens, GA 30605, USA. [2]Nuffield Department of Medicine,

John Radcliffe Hospital, University of Oxford, Oxford, UK. [3] Department of Microbiology,

University of Georgia, Athens, GA 30602, USA. [4]Poultry Microbiological Safety and Processing

Research Unit, U.S. National Poultry Research Center, USDA-ARS, Athens, GA 30605, USA

Corresponding author* ade.oladeinde@ars.usda.gov

[¶] These authors contributed equally to this work.

[&] These authors also contributed equally to this work.

ABSTRACT

Salmonella enterica subsp. *enterica* serovar Heidelberg (*S*. Heidelberg) is a clinically-important serovar linked to food-borne illness, and commonly isolated from poultry. Investigations of a large, multistate outbreak in the USA in 2013 identified poultry litter as an important extra-intestinal environment that may have selected for specific *S*. Heidelberg strains. Poultry litter (PL) is a mixture of bedding materials and chicken excreta that contains chicken gastrointestinal (GI) bacteria, undigested feed, feathers, and other materials of host origin. In this study, we performed a series of controlled laboratory experiments which assessed the evolution of two *S*. Heidelberg strains (SH-2813 and SH-116) in PL previously used to raise 3 flocks of broiler chickens. The strains are closely related at the chromosome level, differing by only 69 single-nucleotide variants. Whole genome sequencing was performed on 86 isolates recovered after 0, 1, 7 and 14 days of evolution in PL. Only strains carrying an IncX1 (37kb), 2 ColE1 (4 and 6kb) and 1 ColpVC (2kb) plasmids survived more than 7 days in PL. Competition experiments showed that carriage of these plasmids was associated with increased overall fitness. This increased fitness was associated with an increased copy number of IncX1 and ColE1 plasmids. Further, all Col plasmid-bearing strains had similar mutations in 102 genes encoded on the chromosome and in 3 genes encoded on the IncX1 plasmid. Additionally, we observed a decrease in susceptibility to tobramycin, kanamycin, gentamicin, neomycin and fosfomycin for Col plasmid-bearing strains. Our study demonstrates how acquisition of multiple Col-like plasmids can change the evolutionary path of *S*. Heidelberg in poultry litter.

INTRODUCTION

The United States (U.S.) is the world's largest producer of poultry with over 8 billion broilers produced yearly (1). Maintaining a safe supply of poultry is increasingly important as demand

increases, production becomes more intensive and facilities become antibiotic free. Broilers

(around 22,000 broilers per house) are typically raised in open plan barns on a deep floor of

bedding materials (typically wood shavings, rice hulls or saw dust (1, 2). Poultry litter (PL) is a

mixture of bedding materials and chicken excreta that contains chicken gastrointestinal (GI)

bacteria, undigested feed, uric acid, feathers and other materials of host origin (3-6). As the cost

of bedding materials increase, land nutrient management requirements become more stringent,

and broiler production and turnover times decrease, farmers have chosen to increase the number

of flocks raised on litter before cleanout. This practice is accepted in production environments,

but the effect of litter reuse on microbial populations has not been well studied.

Available data suggest that litter re-use can result in significant changes in the

microbiological, chemical and physical composition of litter (3, 7, 8). In addition, reused litter

has been found to harbor fewer yeasts, molds, coliform bacteria and *Salmonella* compared to

fresh litter (9, 10). However, fecal shedding of food borne pathogens such as *Salmonella* into

poultry by colonized hosts can also be a source of new infection or re-inoculation. For instance,

Salmonella in 131-day-old litter was shown to colonize 82% of day-old chicks after placement

for 32 days (11) and *Salmonella enterica* subsp. *enterica* serovar Heidelberg (*S*. Heidelberg) has

been shown to persist in recycled litter, even after litter treatment (12). *Salmonella* Heidelberg is

a clinically-important serovar, linked to food-borne illness and commonly isolated from poultry.

After *S*. Typhimurium, *S*. Heidelberg is the serovar of *Salmonella* most often associated with

Salmonella-related deaths in the United States (13). In 2013, *S*. Heidelberg was responsible for a

large multistate foodborne outbreak (634 cases) linked to consumption of chicken (14).

Epidemiological and environmental investigations identified PL as an important extra-intestinal

environment that may have selected for specific *S*. Heidelberg strains (14).

Salmonella Heidelberg is highly clonal and cannot be subtyped by conventional pulsed-

field gel electrophoresis (PFGE) which poses an inherent difficulty when differentiating strains

responsible for foodborne outbreaks from those associated with background sporadic cases (15-

17). High quality core genome single nucleotide variant (hqSNV) analysis on whole genome

DNA sequences (WGS) of *S*. Heidelberg has been recommended as a better approach to

distinguish outbreak strains (15). However, core genome SNV is non-discriminatory between

isolates harboring horizontally acquired antibiotic resistance genes (ARG) versus susceptible

ones. Multidrug resistant *S*. Heidelberg isolates tend to cluster with susceptible isolates during

outbreaks and differ by only 0 - 4 SNVs (16, 17). In a study by Edirmanasinghe et al. (17), the

authors reported 65 out of 113 sequenced *S*. Heidelberg encoded the AmpC beta-lactamase

(bla$_{CMY-2}$) gene on an incompatibility group I1 plasmid (IncI1) and were resistant to important 3rd

generation cephalosporins. On the other hand, susceptible isolates (n=38) carried multiple

variants of Col plasmids and encoded no known ARG. The function of these Col plasmids is not

clear but they seem to be mobilized by co-resident conjugal plasmids, such as IncI1 and IncX,

two plasmids frequently found among *S*. Heidelberg isolates (16-19).

The majority of *S*. Heidelberg isolates sequenced and available in the National Center for

Biotechnology Information (NCBI) database were recovered from either food or clinical samples

(18, 20-22), leaving a knowledge gap in our understanding of strains of environmental origin.

The poultry litter microbiome may carry mobile genetic elements (MGE) including plasmids that

can be horizontally transferred to *S*. Heidelberg upon introduction into PL. These newly acquired

MGE may present a fitness advantage to permissive recipients, further enabling the persistence

of *S*. Heidelberg in poultry environments.

In this study, we investigated the role of PL on the fitness of two *S*. Heidelberg strains with

indistinguishable PFGE patterns, both recovered from chicken carcass in two states in USA.

Both strains harbor an IncX1 plasmid (37 kb) but differ in the occurrence of Col plasmids. We

performed a series of controlled laboratory experiments to assess the evolution of these strains in

PL that was previously used to raise three flocks of broiler chickens. We demonstrate that the

acquisition of multiple ColE1 plasmids in tandem with mutations in host chromosome and the

IncX1 plasmid contributed to the persistence of *S*. Heidelberg in PL. Additionally, this improved

fitness was associated with a decreased susceptibility to tobramycin, kanamycin, neomycin and

fosfomycin. This study elucidates for the first time how acquisition of multiple Col plasmids

from PL alters the evolutionary trajectory of *Salmonella* Heidelberg.

MATERIALS AND METHODS

Poultry Litter for microcosm evolution studies. Poultry litter (PL) for microcosm studies was

provided by the University of Georgia poultry research center in Athens, GA, USA. The PL was

previously used to raise 3 flocks of broiler chickens under simulated commercial poultry

production conditions. The litter was collected ca. 2 weeks after the third flock of birds were

removed. The PL bedding material was composed of pine shavings.

Bacterial strains and preconditioning. *S*. Heidelberg strains (SH-2813 and SH-116) used in

this study were originally recovered from broiler chicken carcass rinsates. The

genotypic/phenotypic characteristics of these strains are provided in Table 1. The strains were

preconditioned in poultry litter extract (PLE) media prepared as described in supplementary

material or in Brain Heart Infusion broth (BHIB; BD Difco, Sparks, MD) before inoculation into

PL. For each strain, 5 single colonies from Sheep Blood Agar (SBA) (Remel Inc, San Diego,

CA) were transferred to 10 ml of PLE and BHIB, and incubated at 37 ^{o}C for 24 h. Serial batch

culturing was carried out for ca. 50 generations in PLE and BHIB by transferring 100 µl of

culture to 10 ml of fresh PLE and BHIB (100-fold dilution) to yield ca. 10 gen. d^{-1}. After 5 days,

the population was used as inoculum for evolution experiments in PL microcosms.

Experimental evolution in PL. Poultry litter was collected in a 5-gallon bucket and thoroughly

mixed before 30 g aliquots were placed into 48 250 ml Nalgene wide jars and 1X PBS (3-4 ml)

was added to achieve 30 % moisture. Microcosms were incubated for 72 h at 37°C before being

individually inoculated with ca. 10^{6} colonies forming units (CFU) of either SH-2813 or SH-116

preconditioned previously in PLE or BHIB. These 4 groups/treatments were as follows: SH-2813

+ BHIB, SH-2813 + PLE, SH-116 +BHIB and SH-116 +PLE. Inoculated microcosms were

thoroughly mixed and then incubated for 14 days at 37 $^{\circ}$C (Supplementary Fig. 1). The control

microcosm remained un-inoculated. Three microcosms per group were destructively sampled at

days 0, 1, 7 and 14.

Poultry litter microcosm sampling. Poultry litter samples (25 g) were mixed with 150 ml 1X

PBS and 5-10 ml glass beads (Fisher Sci: S800242) then homogenized for 10 min at 450 rpm in

a hand wrist shaker (Boekel Scientific, Feasterville, PA). The PL mixtures were allowed to settle

for 5 min to allow settling of large debris/pine shavings and then passed through cheesecloth and

hand pressed to recover eluates which were used for downstream microbiological and chemical

analysis. The moisture content of each PL sample was determined gravimetrically by drying ca.

1 g of PL at 107 $^{\circ}$C for 24 h.

Microbiological methods. Recovered PL eluate was serially diluted and plated onto XLT-4 and

Brilliant Green Sulfa (BGS). For enrichment and recovery of injured/viable *Salmonella* spp., one

ml eluate was transferred to 20 ml of 1 % Buffered Peptone Water (BPW) and incubated 24 h

before being plated onto XLT-4 agar. All agar plates and broth tubes were incubated at 37 $^{\circ}$C for

24 – 48 h before colony forming units were enumerated. Reported counts were log-normalized

and reported per gram PL dry weight. At least three single colonies representing three unique

clones of *Salmonella* from each microcosm were randomly chosen from XLT-4 plate dilutions

and archived in 30 % Luria Bertani (LB) glycerol at -80 ^{o}C.

DNA sequencing and Bioinformatics. We sequenced 86 evolved *S*. Heidelberg isolates

encompassing clones recovered from the four treatments. Isolates were selected randomly from

the highest and lowest dilution of XLT-4 plates; and when populations decreased below LOQ,

isolates were selected after 24 h enrichment in BPW.

Genomic DNA (gDNA) was extracted from pure isolate cultures using FastDNA Spin Kit

(Mp Biomedicals, Solon, Ohio) and quantified with a Qubit Fluorometer (ThermoFisher

Scientific). Whole genome sequencing (WGS) libraries were prepared using MiSeq Nextera XT

library preparation kit (Illumina, Inc., San Diego, CA). Sequencing was performed on the

Illumina MiSeq platform with 250-bp paired end reads using the MiSeq reagent V2 (500 cycles).

All isolates were sequenced for an average coverage of 100X. Sequence reads were assembled

de novo into contigs using SPAdes (23). The quality of assembled genomes was accessed using

Quality Assessment Tool for Genome Assemblies (QUAST). (For detailed information on

assembly, coverage and quality of sequenced genomes see supplementary information).

The original strains of SH-2813 and SH-116 are referred to as ancestral strains, while

isolates from PL evolution experiments are referred to as evolved strains. Assembled contigs

were submitted to the Center for Genomic Epidemiology's multilocus sequence type (MLST)

and PlasmidFinder modules to determine the whole-genome sequence types (ST) and existing

plasmid replicon types. Prophages were identified using PHAST (24) and sequences for intact

phages were extracted, and annotated with Prokka (25). The Comprehensive Antibiotic

Resistance Database (CARD) was used for the identification of resistance genes present on

plasmid and chromosome (26).

Plasmids were classified into MOB families using methods described previously (27). To

investigate the topology of contigs identified with plasmid replicon, we used a custom script to

inspect the path in the SPAdes FASTG assembly graph file that corresponded to the plasmid

contig identified in the FASTA file; the contigs.paths file was used to determine correspondence

between FASTA contigs and FASTG edges. Plasmid contigs were assigned as 'complete

circular'; 'complete non-circular', i.e. linear but not connected to other contigs; and 'non-

complete', i.e. linear and connected to other contigs. (see Supplementary Methods for detailed

information). Complete circular plasmids were annotated using Prokka (25) and mapped with

Snapgene v 4.1.2. When coding DNA sequence (CDS) was annotated as

hypothetical/uncharacterized, we performed a PSI-BLAST search with predicted protein

sequences run to ≤3 iterations. Plasmids were analyzed by comparative analysis using

ProgressiveMAUVE (28) and CDS predicted in regions with marked differences were aligned

with MUSCLE (29). Phylogenetic tree was produced from aligned conserved relaxase or

mobilization protein sequences using the maximum likelihood method implemented in RAxML

v 8 (30) - with a rapid bootstrapping run to generate 1000 replicates.

Single Nucleotide Variants (SNVs) were determined by mapping raw sequence reads to the

chromosome and plasmid genome of *S*. Heidelberg strain AMR588-04-00435 available in NCBI

under accession numbers CP016573 and CP016575, respectively, using Burrows-Wheeler

Aligner (BWA) (31). SAM file sorting, and removal of PCR duplicates was done using

SAMtools (32). Genome Analysis ToolKit (33) with a minimum mapping quality of 30 and a

minimum base quality of 30 was used for SNV calling. A SNV based phylogenetic tree was

generated using the maximum likelihood method implemented in RAxML v 8.

Growth Assay, plasmid copy number determination and gene expression. Plasmid copy

number (PCN) and differential expression of selected transcripts was determined by performing

a growth assay on ancestral and evolved strains. Three populations were initiated from three

single colonies from strains SH-2813$_{anc}$, SH-2813$_{evol}$, SH-116$_{anc}$, and SH-116$_{evol}$. An individual

colony was transferred to 20 ml of Mueller Hinton Broth (MHB) in 50 ml centrifuge tube and

incubated overnight at 37 ^{o}C. Following overnight growth, 6 technical replicate populations (18

total per strain) were established from each culture by transferring 3 ml to 50 ml centrifuge tubes

containing 17 ml MHB. Cultures were incubated without shaking at 37 ^{o}C for 24 h. Optical

density (OD$_{600nm}$) was periodically measured in the 3 populations with a Nanodrop 2000

Spectrophotometer (ThermoFisher Scientific). Bacteria concentration was determined for 2

populations at 0.5, 4 and 24 h via serial dilution plating on R2A agar. Additionally, bacteria

culture (ca. 20 ml) was filtered onto 0.2 µm pore size polycarbonate filters (MilliporeSigma,

Burlington, MA) which were transferred to Lysing Matrix E and B tubes (MP Biomedicals,

Solon, Ohio) for genomic DNA and RNA isolation respectively; and saved at 80 ^{o}C until use.

For a complete description of PCN and gene expression determination, see Supplementary

Methods.

Competitive fitness and conjugation assay. The fitness of three strains carrying the 3 Col

plasmids (SH-2813$_{evol}$, SH-116$_{anc}$, and SH-116$_{evol}$) was determined relative to SH-2813$_{nal}$

nalidixic resistance with no Col plasmid. The SH-2813$_{nal}$ strain is isogenic to SH-2813$_{anc}$ but was

made resistant to 200 ppm nalidixic acid by spontaneous mutation in *gyrA*. Five colonies from

SBA for each strain were incubated in 9 ml of fresh MHB at 37 ^{o}C for 20 h. Thereafter, cultures

of the strains were mixed at a ratio of 1:1 (~500 µl of strain under study to 500 µl of SH-2813$_{nal)}$.

We assumed that each strain had the same initial concentration because 24 h population were not

significantly different (Table 2). Mixtures were diluted 10,000-fold in 50 ml centrifuge tubes (4

technical replicates per mixture) containing 20 ml of MHB and competed for 24 h at 37 ^{o}C

without shaking. The final population was measured by serial plating onto R2A agar

supplemented with and without 32 ppm of nal. The fitness of the strains carrying Col plasmids

relative to the SH-2813-nal strain was determined as described by ref (34). For details see

Supplementary Methods.

To pool the information from the competition assay with conjugation assay we employed

the same data set to determine the conjugation rate constant for the three Col plasmid types

present during the competition assay. Conjugation rate constants were determined as described

by Loftie-Eaton et al (35). For details see Supplementary Methods.

Antibiotic Susceptibility Testing. We performed antimicrobial susceptibility testing (AST) on

selected evolved strains following National Antimicrobial Resistance Monitoring System

(NARMS) protocols for Gram-Negative bacteria (36) and Kirby-Bauer disc diffusion assay (37).

For more details on the disc diffusion protocol used, see Supplementary Methods.

For categorizing susceptibility, breakpoints were adopted from NARMs. There is no

standardization by Clinical and Laboratory Standards Institute (38) for tobramycin, kanamycin,

neomycin, netilmicin, and fosfomycin by disc diffusion for *Salmonella* Heidelberg. Therefore,

results were interpreted with diameter difference in zone of inhibition between isolates.

Plasmid-specific genes prevalence in poultry litter. To assess the prevalence of Col plasmids

in PL, we determined the concentration of plasmid-specific genes in various litter collected for

another on-going experiment in our laboratory. These litter were sourced from 7 poultry farms in

4 states in USA, practicing either conventional or backyard method of raising chicken flocks.

DNA was extracted in triplicates from each PL as described previously for PL DNA extraction.

We performed qPCR on 2 µl of purified DNA extract as described previously in qPCR assays,

with primer sets specific to each plasmid of interest (Supplementary Table 9).

Statistical analysis. Analyses were performed using R (version 3.4.1). Survival difference

between strains preconditioned in PLE versus BHIB was done using a profile analysis followed

by a multivariate analysis of variance (MANOVA). Comparisons among strains or treatment

were performed using Wilcoxon signed-rank Test. A multiple linear regression was used to

model the relationship between IncX1 and Col plasmid copy numbers derived from whole

genome sequencing coverage.

Ethics statement. The authors did not physically interact with chickens raised on the poultry

litter used in this study, therefore, we were exempt from university guidelines and USDA/NIH

regulations regarding animal use. Institutional Animal Care and Use Committee (IACUC) -

UGA Office of Animal Care and Use.

Data availability. All sequencing data are available under NCBI Sequence Read Archive (SRA)

BioProject: PRJNA448609

RESULTS

Experimental evolution system. The survival of two *S*. Heidelberg strains (SH-2813 and SH-

116) in previously used PL was evaluated over a period of 14 days. The strains which were

recovered from chicken carcass from two states in the USA were closely related at the

chromosomal level, differing by only 69 single-nucleotide variants (SNVs). In addition, both

strains carried the IncX1 plasmid and SH-116 carried 2 ColE1-like plasmids (4 and 6 kb) and a

cryptic ColpVC plasmid (2 kb) (Table 1).

Table 1. Characteristics of the strains used in this study

Strain ID	Serotype	Plasmid Profile (size)	*Resistance profile	#Antimicrobial resistance genes	Unique Single Nucleotide Variants (SNV)
SH-2813	ST15	IncX1 (37kb)	Ery, Tyl,	AAC (6')-Iy, FosA7	69
SH-116	ST15	IncX1 (37 kb), ColE1(6 kb) ColE1 (4 kb) ColpVC (2 kb)	Ery, Tyl, Pen	AAC (6')-Iy, FosA7	49

Note. - Ery = Erythromycin; Pen = Penicillin; Tyl= Tylosin

* Susceptible to all 14 antibiotics on the NARMs Gram-negative AST panel

Present on chromosome

The concentration of the two *S*. Heidelberg strains preconditioned in either PLE or BHIB decreased by several orders of magnitude in all microcosms and their survival dynamics differed by preconditioning media and strain-type (Fig. 1a). **Figure 1. Survival curve and mutation based phylogeny of evolved *S*. Heidelberg strains**. (**a**) *S*. Heidelberg concentration from poultry litter microcosms reported in colony forming units (CFU) and normalized by litter dry weight. (**b** and **c**) Top: Single nucleotide variants (SNV) based phylogenetic tree for SH-2813 and SH-116 evolved strains, Bottom: Corresponding Venn diagram comparing number of SNV shared across time points. Clades highlighted in blue shadows denotes clones carrying 3 Col plasmids. Node tips highlighted in green rectangular boxes represents isolates recovered following overnight enrichment in buffered peptone water. Trees were rooted with SH-2813 and SH-116 ancestors. (d) Venn diagram comparing SNV shared between Col plasmid-carrying SH-2813 and SH-116 evolved strains. The concentration of SH-2813-PLE, decreased by 4-log within 1 day of inoculation into PL but rebounded 1.5-log after 7 days. In contrast, the concentration of

SH-2813-BHIB was below the limit of quantification (LOQ) within 1 day of inoculation into PL

and increased ca. 4-log by day 7. The final concentration of SH-2813-PLE and SH-2813-BHIB in

PL was log 2.5 ± 0.03 CFU g^{-1}. On the other hand, the concentration of SH-116-PLE decreased

gradually for 14 days while SH-116-BHIB was below LOQ within 1 day of inoculation into PL

and increased by ca. 3-log at day 7. The final concentration for SH-116 was below our LOQ for

direct culture counts but the strain was isolated at varying levels following enrichment in

buffered peptone water (BPW) (Supplementary Table 1).

A profile analysis followed by MANOVA on the observed bacteria concentration revealed

that preconditioning had a significant effect on SH-2813 concentrations at day 0 and day 1 (p-

value< 0.004). For SH-116, this effect was only significant at day 1 (p-value< 0.0001). We also

monitored the pH, moisture, nutrients, cations and resident aerobic bacteria present in PL during

the 14-day experiment, however we could not draw significant conclusions from analyzed results

(Supplementary Table 1 and 2)

IncX1 and Col plasmids are associated with *S*. Heidelberg persistence in PL. To investigate

the genetics of *S*. Heidelberg persistence in PL, we isolated and performed WGS on 87 isolates

obtained after 0, 1, 7 and 14 days in PL (Supplementary Table 3). All SH-2813$_{evol}$ strains (n= 27)

obtained from day 7 and 14 microcosms carried 3 Col plasmids in addition to an IncX1 plasmid,

and differed from day 0 and 1 isolates by 76 -79 unique SNV on the chromosome (Fig. 1b). It is

noteworthy that SH-2813 isolates recovered following enrichment in BPW (n = 6) were the only

day 1 strains carrying Col plasmids. For SH-116, IncX1 and Col plasmids were present in all

sequenced strains (n = 38) regardless of time of incubation in PL and isolates differed by only 0 -

4 unique SNV on the chromosome (Fig. 1c). The majority (276/306) of the plasmid contigs

represented complete circular plasmids (Supplementary Table 4). The IncX1 belonged to the

MOB_Q family of relaxases and was present in all sequenced genomes. The ColE1 plasmids taken

up by SH-2813$_{evol}$ belonged to the MOBp family and showed a conserved size of 6189 (ColE1-

6kb) and 4784 (ColE1-4kb) bp respectively. The cryptic ColpVC was 2223 bp (ColpVC-2kb).

For all complete IncX1 plasmids, there was a 21-base pair (bp) difference in strains carrying the

Col plasmids (37, 824 bp) compared to strains with no Col plasmids (37, 845 bp).

Direct-repeat deletion is associated with IncX1 copy number increase. To determine the

reason for the 21-bp size difference in the IncX1 plasmids present in *S*. Heidelberg populations

carrying Col plasmids, all complete circular IncX1 plasmid sequences were subjected to whole

genome sequence alignment and SNV identification. The 22-bp direct-repeat deletion was

located at the origin of replication (*ori*) adjacent to the replication protein (RepB) and an

additional 1-bp (G-> GA) insertion occurred in the type IV secretion protein (T4SS) VirB8 (Fig.

2a and Supplementary Table 5). **Figure 2. Deletion of 22-bp iteron of IncX1 reduces *rep***

expression. (**a**) Annotated IncX1 plasmid map of SH-2813$_{anc}$ and SH-116$_{anc}$. Regions that

acquired mutations in evolved strains carrying Col plasmids are highlighted in rectangular boxes

(**b**) Multiple alignment of the IncX1 iteron-based direct repeat (DR) origin of selected strains

from this study, and 22 *S*. Heidelberg genomes available on NCBI. (**c**) Differences in expression

of IncX1 genes in evolved SH-2813 and SH-116 compared to SH-2813$_{anc}$ and SH-116$_{anc}$ ($\log_2$

scale, n = 3, **p*.value =0.086 ;Wilcoxon signed-rank Test). The error bars represent the standard

error of the mean.

The 22-bp deletion reduced the number of direct repeats from 3 to 2, making it a truncated

ori (Fig. 2b). Direct repeat iterons of theta plasmids are known to bind Rep proteins and have

been shown to participate in negative control of their initiation (39). Deleting or adding more

copies of the iteron have been shown to increase or decrease plasmid copy number (PCN) (40).

Therefore, we hypothesized that the direct repeat deletion in IncX1 of SH-2813$_{evol}$ would result

in higher copies compared to SH-2813$_{anc}$ since the 22-bp is absent in the ancestral strain while no

change would be expected in IncX1 copies for SH-116$_{evol}$. To test this hypothesis, we determined

the copy numbers of IncX1 and Col plasmids present in ancestor (SH-2813$_{anc}$ and SH-116$_{anc}$) and

2 evolved strains (SH-2813$_{evol}$ and SH-116$_{evol}$ respectively). We assumed that these evolved

strains represent the "fittest" SH-2813 and SH-116 strains given their persistence in PL

microcosms for 14 days.

As expected, the copy number of IncX1 for SH-2813$_{evol}$ (~7 copies/cell; s.d.=1.8) was

higher than that of SH-2813$_{anc}$ (~4 copies/cell; s.d.=0.3) at 0.5 h (p. value = 0.086; Wilcoxon

rank-signed test). In contrast, SH-116 did not exhibit any significant difference in IncX1 copies

between SH-116$_{anc}$ (~5 copies/cell; s.d.=1.7) and SH-116$_{evol}$ (~6 copies/cell; s.d.=0.8) (p. value =

0.773) (Table 2).

Table 2. Plasmid copy numbers of ancestral and evolved strains.

				Plasmid Copy Number/Cell (S.D)			
Strain	Time (h)	Optical Density (S.D)	Bacteria concentration (Log CFU/ml)	IncX1	ColE1-6kb	ColE1-4kb	ColpVC
SH-2813$_{anc}$	0.5	0.015±0.00	7.98±0.0	4.17±0.3	N/A	N/A	N/A
	4	0.040±0.00	8.60±0.2	0.47±0.5	N/A	N/A	N/A
	24	0.070±0.00	8.90±0.1	0.97±0.9	N/A	N/A	N/A
SH-2813$_{evol}$	0.5	0.016±0.00	7.94±0.0	6.92±1.8	10.71±4	12.4±5.5	110.77±45
	4	0.056±0.00	11.16±2.1	0.59±0.3	3.28±0.6	2.77±1.0	86.21±25
	24	0.080±0.00	8.57±0.0	0.64±0.1	5.25±0.7	4.2±0.6	115.32±70
SH-116$_{anc}$	0.5	0.014±0.00	7.83±0.1	4.55±1.7	0.27±0.3	2.54±1.9	82.39±8
	4	0.037±0.00	8.56±0.1	0.80±1.3	0.14±0.0	0.54±0.7	54.01±26

	24	0.093±0.00	9.07±0.0	2.33±9.3	2.23±0.0	0.92±1.3	39.94±28
SH-116$_{evol}$	0.5	0.017±0.00	7.91±0.0	5.61±0.8	9.44±3.5	10±4.4	90.47±45
	4	0.065±0.00	8.73±0.2	0.60±0.8	3.83±1.7	2.9±1.3	93.83±11
	24	0.069±0.00	8.69±0.1	0.35±0.4	4.39±1.0	3.5±0.3	72.56±20

Note. – N/A not applicable

Gene expression of IncX1 replication protein and stabilization genes. We monitored the

expression of the *repB* replication gene and *relBE* genes which encode a type II toxin-antitoxin

(TA) system associated with the IncX1 plasmid. The *relBE* genes were selected since they are

adjacent to the *ori* (Fig. 2a) suggesting they may share the same promoter as *repB*; and due to

their implicated role in plasmid stabilization (41, 42). Gene expression results in SH-2813$_{evol}$

revealed a marked 4.5-fold reduction in *repB* expression and 2.2-fold reduction in *relB*

expression at 4 h relative to SH-2813$_{anc}$; while the expression levels of toxin *relE* remained

unchanged in both ancestral and evolved SH-2813 (Fig. 2c).

SH-116 strains lacks the extra 22-bp direct repeat present on IncX1, thus no change in Rep

expression was expected. However, the same pattern of expression was observed for these

transcripts for SH-116$_{evol}$ — reductions in *repB* and *relB* and no change in *relE* when compared

to SH-116$_{anc}$ (Fig. 2c). We hypothesize that the reduction in expression of Rep for evolved

strains was a result of selecting for high copy ColE1 plasmids in PL. In fact, SH-116$_{evol}$ carried

ColE1 plasmids with higher copies (ColE1-6kb =9±3.5, ColE1-4kb = 10±4.4, ColpVC-2kb =

90±45) than SH-116$_{anc}$ (ColE1-6kb =0.3±0.3 ColE1-4kb = 3±1.9, ColpVC-2kb = 82±7.9) (Table

2). This difference was further corroborated by comparing PCN determined from WGS assembly

coverage, showing that all SH-116$_{evol}$ isolates from day 7 and 14 on average had higher ColE1

plasmid copies than SH-116$_{anc}$ (Supplementary Table 6).

Further, multiple regression analysis of IncX1 and Col plasmid copies showed that IncX1

was a statistically significant predictor of Col plasmid copies in a cell (F (3, 68) =13.82, p.value

< 0.000, R^2 =0.39). SH-116 strains were predicted to have ca. six ColE1-6kb plasmid copies for

every IncX1, whereas, SH-2813 was predicted to have 3 copies per IncX1 (Fig. 3). **Figure 3.**

Correlation between IncX1 and ColE1-6kb plasmid copies. Plasmid copy number (PCN) was

determined from whole genome sequencing de-novo assembly coverage. The presence of this

high copy ColE1 plasmids correlated with faster growth rates in MHB. The optical density of

SH-2813$_{evol}$ and SH-116$_{evol}$ was 1.4 and 1.7-fold higher than that of SH-2813$_{anc}$ and SH-116$_{anc}$,

respectively, at 4 h (Table 2).

ColE1 plasmids differed in mobilization genes and non-coding RNA. ProgressiveMauve (28)

alignments showed that ColE1 plasmids present in sequenced strains were highly similar and

only differed in nucleotide regions coding for mob genes (Fig. 4; Supplementary Fig. 2b and c).

Figure 4. Col-like plasmid maps and conserved mobilization genes. (**a**, **b** and **c**) Annotated

Col plasmid maps of (**a**) ColE1-6kb (**b**) ColE1-4kb and (**c**) ColpVC from SH-116$_{anc}$. (**d and e**)

Maximum likelihood based tree of aligned conserved protein sequences of (**d**) relaxase (MbeA)

and (**e**) MbeC genes of Col plasmid-carrying strains. Clades highlighted in blue and red shadow

represent ColE1-4kb and ColE1-6kb plasmids, respectively. Trees were rooted with MbeA and

MbeC sequences of *E. coli* (Accession number: NC_001371).

We identified 5-6 open reading frames (ORF) encoding 2-3 mobilization (mob) genes, a

putative type IV TA system (Abi), a putative aminoglycoside 6-N-acetyltransferase (aac(6')-Ib)

and a hypothetical protein for ColE1-6kb (Fig. 4a). The ColE1-4kb plasmids had 4-6 ORFs

encoding a protein named macrophage stimulating factor (MSF), a hypothetical protein and 1-4

mob genes (Fig. 3b). Both ColE1 plasmids harbored a non-coding RNAI and RNAII, while

ColE1-6kb carried an additional non-coding repressor of primer (Rop) protein known to regulate

replication and ColE1 plasmid copy numbers. The ColpVC-2kb plasmids had 2-3 ORFs,

encoding 1-2 replication proteins (Rep) and a hypothetical protein (Fig. 4c). The RNAI present

in the ColE1 plasmids had ca. 62 % DNA sequence homology (data not shown).

These mob CDS were identified as MbeA and MbeC and were present in different sizes

and arrangement. In some cases, the *mbeA* gene overlapped ORF annotated as *mbeB and mbeD*

(Fig. 4; Supplementary Fig. 3), suggesting MbeB and MbeD may be part of the mob region in

some but not all ColE1 plasmids. Multiple alignments of MbeA relaxase protein sequences

showed high conservation in the N-terminal region and high variation in their C-terminal (data

not shown). According to the phylogenetic tree obtained for these conserved amino acids, MbeA

proteins belonging to ColE1-4kb plasmids clustered in a unique clade, whereas the ColE1-6kb

clustered next to the signature *E. coli* ColE1 plasmid (NC_001371) (Fig. 4d). Accordingly, *mbeC*

genes from ColE1-6kb and ColE1-4kb were represented by separate clades on phylogenetic tree

(Fig. 4e).

Col plasmids did not impose a fitness cost on host cell. Although acquisition of Col plasmids

could result in faster growth in host cells it may also translate to a fitness cost if their regulation

is not equally constrained. To establish if acquisition of Col plasmids imposed a fitness cost on *S*.

Heidelberg, we competed 2 evolved (SH-2813$_{evol}$ and SH-116$_{evol}$) and 1 ancestral (SH- 116$_{anc}$)

strain carrying three Col plasmids against a Col plasmid-free isogenic SH-2813 harboring a *gyrA*

mutation for nalidixic acid resistance (SH-2813$_{nal}$). The fitness of each strain was determined

relative to SH-2813$_{nal}$ as described by Millan et al (34). Results show that Col plasmid carrying

strains had a higher relative fitness of 1.27 (s.d. =0.12, n =3) compared to SH-2813$_{nal}$, and

differences were observed between strains (Fig. 5a). **Figure 5. Evolution of fitness and**

Conjugation rate (a) Fitness of Col plasmid-carrying strains of SH-2813 and SH-116 relative to

the Col plasmid-free ancestor of SH-2813$_{nal.}$ Bars represent means of four technical replicates for

each population. The horizontal dashed red line represents the fitness of SH-2813$_{nal}$. (b)

Conjugation rates of Col plasmids from Col plasmid-carrying strains into SH-2813$_{nal}$ ancestor.

Conjugation rates was estimated using data from fitness experiment. Bars represent means of

four technical replicates for each population.

The transfer rate for ColE1-4kb plasmid from SH-116$_{evol}$ to SH-2813$_{nal}$ was below the

detection limit of our assay. This difference in fitness between strains can be may be attributed to

a higher conjugation rate observed for ColE1-6kb and ColpVC for SH-116$_{evol}$ (Fig. 5b).

Antimicrobial resistance associated with the acquisition of Col plasmids. The ColE1-6kb

plasmid in the current study carries a putative aac (6')-Ib gene that is 56 % (*e.value* = 1<e-13)

identical to the aac (6')-Ib gene present in *Klebsiella pneumoniae* Sul1-type integron (GenBank

accession number: AAR18814.1) and 34 % (*e..value* = 6<e-28) identical to the one present in

Thiomona sp. X19 (GenBank accession number SCC93141.1); suggesting it may play a role in

aminoglycoside resistance (Fig. 6a and b). **Figure 6. Acquisition of ColE1 plasmids was**

associated with decreased susceptibility to aminoglycosides and fosfomycin. (**a**) Multiple

alignment of conserved N-terminal residues for aminoglycoside acetyltransferase (AAC-6')

genes encoded in the *S*. Heidelberg (SH-2813) chromosome and ColE1-6kb plasmid from this

study, and AAC-6' genes with known resistance phenotype. (**b**) Corresponding neighbor-joining

tree of aligned AAC-6' protein sequences. (**c**) Inhibition zone diameter distributions for evolved

strains and selected antimicrobial drugs.

Further, *S*. Heidelberg ancestral strains each carried chromosomally encoded

aminoglycoside (aac (6')-Iy) and fosfomycin (fosA7) genes (43) that have been demonstrated to

confer resistance to tobramycin, netilmicin, neomycin, amikacin, kanamycin and fosfomycin

(44-46). To assess the correlation between AMR and the acquisition of Col plasmids, we

performed antibiotic susceptibility testing (AST) on evolved *S*. Heidelberg strains (n=42) on the

NARMs antimicrobials (36). In addition, we used disc-diffusion assay (37) to evaluate

susceptibility to tobramycin, netilmicin, gentamicin, neomycin, kanamycin and fosfomycin.

The majority of the isolates tested (97.6 %) were susceptible to the 14 drugs on NARMs

AST panel (Supplementary Table 7) However, 12 of 16 (75 %) SH-2813$_{evol}$ isolates recovered

from PL on day 7 and day 14 displayed an increase in MIC for gentamicin. The average zone of

inhibition (mm) for kanamycin, tobramycin and fosfomycin for SH-2813$_{anc}$ strains was larger

than the zones measured for SH-2813$_{evol}$ strains (Wilcoxon signed-rank test; *p*-value < 0.0001)

(Fig. 6c). In contrast, SH-116 showed no significant difference in zone of inhibition between SH-

116$_{anc}$ and SH-116$_{evol}$ strains for any of the antibiotics tested (Wilcoxon signed-rank test; *p*-value

> 0.01).

Col plasmid-encoded genes are prevalent in PL. To extrapolate beyond our microcosm

evolution experiment, bioinformatics and quantitative PCR were used to assess the pattern of

distribution of these Col plasmids across genomes and in PL. DNA sequences for complete Col

plasmids from this study were subjected to BLAST searches against genomes that have been

deposited in GenBank. No BLAST hit matched the complete sequence of ColE1-6kb plasmids,

but the majority of hits covered ca. 55 % of the plasmid genome. In most cases, these hits

corresponded to mobilization genes.

For ColE1-4kb, the closest hit was 99 % identical to the colicigenic plasmid of *Salmonella*

enterica serovar Berta (pBERT) and covered the entire genome. All other plasmid hits covered

<82 % of the ColE1-4kb genome and were represented in various species of Gram-negative

bacteria. The ColpVC-2kb was >99 % identical to plasmids from 5 *Salmonella* sp, including

three *S*. Heidelberg strains isolated from Chicken in Canada (data not shown). These results

suggested that Col plasmids of poultry origin are not well represented in GenBank.

To address this possibility, we quantified Col plasmids in litter from conventional and

backyard poultry farms located in 4 states in USA. The Col plasmid genes were detected in all

litter types except 1 backyard litter, whereas the Rep protein for IncX1 was detected in 4 of 7 of

the litter types, suggesting that these genes are present in PL (Fig. 7). **Figure 7. Concentration**

of plasmid-specific genes in poultry litter of different types. Poultry litter (PL) was collected

as grab samples from multiple locations on the farm and composited in 1-gallon Whirl-Pak bags.

qPCR was performed on DNA extracted from 250 mg of PL in triplicates. The error bars

represent the standard error of the mean. Con =Conventional farming, Back = Backyard farming,

TSC =Target sequence copies

DISCUSSION

Col plasmids and IncX1 contributed to the fitness of *S*. Heidelberg

We show for the first time the evolutionary trajectory of *Salmonella enterica* serovar

Heidelberg in PL. We chose *S*. Heidelberg for its increasing role in food-borne outbreaks and its

carriage of Col-like plasmids (14, 16, 17, 22, 47). In our study, only *S*. Heidelberg strains

carrying 2 ColE1 and a cryptic ColpVC plasmids survived more than 7 days in used PL. The

ColE1-6kb relaxases (MbeA) shared >77 % homology with the relaxase of ColE1-4kb but was

more closely related to the prototype ColE1 (>96 %) plasmid of *E. coli*. When the ColE1

plasmids sequences from this study were compared to those available in NCBI, we found that

these plasmids were underrepresented in this database. The observed low frequency may be a

consequence of the bias toward sequencing strains of clinical origin, where plasmids carrying

known antibiotic resistance genes are of primary interest.

After screening multiple PL types with qPCR, we detected Col plasmid genes at high levels

in 6 of 7 litter types. Our data further suggest that the PL environment may impose selective

pressure that promotes the acquisition of these Col plasmids.

The ColE1-6kb plasmids were present in higher copy numbers than ColE1-4kb and carried 2

putative genes encoding a type-IV TA system and an aminoglycoside-acetyltransferase. The TA

system was linked by STRINGDB (48) to the widespread abortive bacterial infection (Abi)

system belonging to the cluster of orthologous group COG1106 (domain of ATPases) and the

closest homologs were those present in Pseudomonadaceae and Lactobacillaceae. In some

bacteria, this protein is annotated as ATP-binding protein or abortive infection protein. Abi

systems induce phage resistance to prophages of type 936 in *Lactococcus* and enhance plasmid

maintenance/stabilization via bacteriostatic toxicity (49). Consequently, cells carrying plasmids

encoding the Abi gene will be immune to phage attack or replication, thereby providing

protection for their clonal population. Their role in *Salmonella* remains questionable; however,

their high expression under laboratory growth conditions in our study (Supplementary Fig. 4)

suggests that they are functional and may have contributed to the persistence of *S*. Heidelberg in

PL.

We do not know the potential functions of the accessory genes on ColE1-4kb, but their

compatibility with ColE1-6kb may stem from differences in their RNAI sequences (50) and their

mob region. The ColE1 plasmid family represents plasmids that are only transmissible in the

presence of an additional plasmid with conjugative functions and usually carry a mob region

encoding specific relaxase and accessory mob genes (*mbeB*, *mbeD mbeC*). The relaxase

component is crucial for both initiation and transfer of DNA and is highly conserved (51, 52). In

addition, it has been predicted that the C-terminal region of MbeA binds ssDNA, while the N-

terminal fragment contacts MbeC (53, 54). The observed differences in C and N- terminal of

MbeA and MbeC in this study suggests that such modifications may be critical for successful co-

existence of multiple ColE1 plasmids in one host cell.

The Incompatibility factor X1 plasmid harbored by the *S*. Heidelberg strains used in our

study more than likely served a conjugative role. IncX1 is a low copy plasmid and carries

required mating pair formation system for horizontal gene transfer (HGT) including type IV

secretion systems (T4SS), coupling proteins and a relaxase (55, 56) (Fig. 2a). This plasmid was

present in all strains recovered which infers that they are stably maintained and critical to the

survival *S*. Heidelberg strains carrying them.

Stability of conjugative plasmids has been attributed to high conjugation rates and plasmid

stabilization proteins (57, 58) . The Indel present in VirB8 resulted in a frame-shift mutation in

this protein and subsequent changes in DNA regions coding for T4SS proteins. Indeed, this

mutation resulted in the fusion of a smaller VirB8 (4.4 kDa) encoding gene with a bigger one

(23.4 kDa) (Fig. 2a). In addition, differences were observed in sequences coding for the

conjugal transfer protein (TraG), an important protein for bacterial conjugation (59).

Interestingly, one strain without TraG (SH-33A0.5-116-Day7-PLE) was isolated but we do not

know the conjugal implication of this deletion (Supplementary Fig. 5)

Following the acquisition of Col plasmids, we observed a significant reduction in the

expression of replication protein (Rep) protein and a type II toxin-antitoxin (TA) gene. This

reduction in *rep* expression suggests that *rep* expression may be linked to the cost of taking up

new MGE. Reducing *rep* expression will subsequently decrease the copies of IncX1 available for

ColE1 plasmids mobilization — thus reducing the burden associated with maintaining multiple

plasmids. Also, regression analysis of IncX1 and Col copies showed that IncX1 was a good

predictor of ColE1-6kb copies per cell — evolved strains carrying high IncX1 copies also carried

high ColE1-6kb copies (Fig. 3; Supplementary Table 6). This result suggests there might be an

equilibrium between the number of "helper plasmids" to mobilizable plasmids residing in a cell.

Bacteria (TA) modules are genetic elements composed of a toxin protein component that

disrupts bacterial growth by interfering with vital cell processes and an antitoxin that impairs the

functionality of the toxin until this inhibition is abolished in response to cellular signaling. Type

II TA system carried on plasmids are important players in post-segregational killing — plasmid-

less cells are killed by the toxin, thereby establishing the dominance of plasmid-bearing cells in a

population (41, 42, 60).

At late logarithm and stationary phase growth in Mueller Hinton Broth, the *relB* gene

(antitoxin) of IncX1 was significantly downregulated in evolved strains compared to ancestral

strains, and no change was observed in *relE* (toxin) expression (Fig. 2c). This continuous

expression of *relE* will ensure that only IncX1 carrying cells are present in the population.

Therefore, disrupting the conjugative and stability genes of conjugative plasmids may represent a

critical point for limiting their persistence in the environment. For instance, a combination

treatment with linoleic and phenothiazine reduced the conjugation efficiency of six conjugative

plasmids by 3 -fold, including the IncX plasmid of RP4 (61).

Furthermore, the IncX1 present in Col plasmid-bearing strains of *S*. Heidelberg carried a 22-

bp direct repeat deletion at the origin of replication (*ori*), an Indel in VirB8 (G-> GA) and a non-

synonymous substitution in a hypothetical protein (I111T). The Rep protein of IncX1 belongs to

the family of plasmids that use the theta mode of replication similar to the F, P1, R6K and

ColE1-like plasmids. The Rep protein interacts with iteron sequences in order to control the copy

number of plasmids in a cell (39, 62).

Briefly, the Rep-iteron complexes stimulate the initiation of replication at low copy until the

regular copy number is achieved. However, when the copy number is increased beyond the

normal levels, the Rep-iteron concentration is high enough to bind almost all protein bound

iterons. This direct interaction between the plasmid molecules produces the coupling or

handcuffing of the origins, making them unavailable for the initiation of replication. Moreover,

these repeated sequences are important incompatibility determinants for iteron-based plasmids.

Strains with this deletion had higher copies of IncX1 than ancestral strain carrying the extra

22-bp direct repeat. This result supports other studies showing that the deletion of iterons

increased plasmid copy number of theta plasmids (reviewed in (63)). For example, Millan et al

(64) showed that mutations acquired in the OriV of a synthetic ColE1-like plasmid (pBGT)

carrying bla$_{TEM-1}$ β-lactamase gene increased plasmid copy number and subsequently increased

ceftazidime resistance several folds.

To expand beyond the *S*. Heidelberg strains used in our study, we performed an alignment of

the *ori* of closed "circular" IncX1 plasmids of ca. 37kb present in *S*. Heidelberg strains available

in NCBI. We observed that all the strains analyzed carried only two iterons (Fig. 2b), which

implies that this deletion is overrepresented in sequenced *S*. Heidelberg carrying IncX1 plasmids.

The majority of these strains were from clinical settings (Supplementary Table 8) which suggests

they are being recovered more often and may be associated with an increase in fitness of those

strains as was also observed in our study. Together, our results support the idea that iterons

regulate IncX1 plasmid replication and their compatibility with other plasmids.

Mutations acquired on plasmids and chromosomes after an HGT event have been shown to

compensate for the fitness cost associated with the acquisition of a new (MGE) (35, 64-66).

Here, strains carrying Col plasmids had a total of 4 plasmids and persisted for up to 14 days,

which is a "plasmid paradox," where the acquisition and maintenance of multiple plasmids over

the long term is supposed to impose a burden on their host (57, 58).

This principle has been elucidated upon, where positive epistasis minimized the cost

associated with carrying small plasmids of *Pseudomonas aeruginosa* over a long-time scale (34).

Recently, a study showed that the acquisition of three ColE1 plasmids in naïve *H. influenzae*

hosts did not impose a fitness cost as long as the bacterium compensated for the first plasmid

(67). The presence of one plasmid favored the presence of a second plasmid and the presence of

this second plasmid or third plasmid did not impose an additional significant biological cost,

even though these plasmids produced a cost when they are alone in the cell .

In this study, Col plasmid-bearing strains had increased fitness compared to a strain

without a Col plasmid and shared 102 single nucleotide variants (SNV) in common (Fig. 1d). It

is unlikely that all observed SNV are to compensate for Col plasmid acquisition, as some will be

compensatory for other MGE such as bacteriophages (68-70). For example, the prophage Gifsy-2

was found intact in 97% of *S*. Heidelberg isolates sequenced in our study, suggesting the PL

environment may have selected them for increased fitness (Supplementary Fig. 6). Further, the

ubiquity of Gifsy-2 in *S*. Heidelberg strains has been reported, which infers they are important

for *S*. Heidelberg survival (13). We observed a mutation in the replication protein (RepO) of

Gifsy-2 denoting these events are linked (Supplementary Table 5). Taken together, it is plausible

that mutations observed in this study are associated with HGT events involving both ColE1

plasmids and prophages; in addition to mutations selected for persistence in PL.

Plasmid and chromosome encoded genes mediated antibiotic resistance in *S*. Heidelberg

Another significant observation from this study is the decrease in susceptibility of Col

plasmid-bearing strains towards kanamycin, tobramycin and fosfomycin. We do not know what

these reductions in inhibition zones mean in terms of MIC as there are no current CLSI standards

(38) for *S*. Heidelberg. In addition, it has been shown that simple back- calculations from

inhibition zones to MIC can be confounded by several covariates such as serotype, host species,

and year of isolation (71).

Kanamycin and tobramycin are not used for treating *Salmonella* infections due to their

resistance in-vivo, although tobramycin in combination with fosfomycin is used to treat people

with cystic fibrosis caused by *Pseudomonas aeruginosa* (72). Since no drugs were added to the

litter used in this study (Personal Communication- Dr. Casey Ritz, University of Georgia),

experimental selection for these resistance phenotypes is unlikely. Although alternative

explanations including Col plasmid selection or population factors like competitive exclusion

and the presence of "cheaters" (73-75) may account for this result, more research is needed to

test these possibilities.

Despite the fact that the acquisition of Col plasmids coincided with the observed change in

susceptibility, we cannot say for sure if these plasmids played any role. The putative

aminoglycoside acetyltransferase (aac-(6')) gene carried on ColE1-6kb represents a fragmented

aac-6' gene and shares significant homology with the N-terminal of aac(6')-Ib of *K. pneumoniae*.

Its small size (81 amino acid residues) compared to other known aac genes (Fig. 5a), and the

absence of homologs in all antibiotic resistance genes (ARG) databases queried suggests it may

be an uncharacterized ARG. Importantly, its presence on ColE1 will result in more copies of

these gene compared to the aac-(6') gene carried as a single copy on the chromosome. More

studies will be required to determine if these two aac-(6') genes confer decreased susceptibility

towards the aforementioned aminoglycosides.

Regarding fosfomycin, Rehmann et al (43) determined that *S*. Heidelberg strains with an

inhibition zone of 26 mm were resistant to >32 µg/ml of fosfomycin, which suggests that

evolved *S*. Heidelberg strains carrying Col plasmids in our study may be resistant to fosfomycin.

The fosA7 gene is present in many *S*. Heidelberg genomes and resistance to fosfomycin has been

demonstrated in a few (43). In addition, *S*. Heidelberg harbors other chromosomal genes

associated with fosfomycin resistance including *glpT* and *uhpT* (76).

In this study, we did not observe any mutation in these genes, thus cannot correlate

fosfomycin resistance to a specific ARG. This result shows the intrinsic bias that could be

associated with estimating resistance phenotype from only known ARG that are available in

curated databases. More concerning, is the spontaneous mutant (SH-17BBPW-2813) that

emerged with reduced susceptibility to 3 classes of antibiotics following overnight enrichment in

buffered peptone water (BPW) (Supplementary Table 7). One explanation is that BPW enriched

for the limited strains present in PL or it created an environment where only the fittest strain

emerged. Therefore, we argue that multiple chromosomal mutations in addition to Col plasmids

are responsible for the reduced susceptibility displayed by evolved *S*. Heidelberg isolates,

including a non-synonymous substitution (T153A) in efflux mediated multi-drug resistance gene

acrA (77, 78).

Results from these PL evolution experiments suggest that IncX1 and ColE1 plasmids

contributed to the fitness of *S*. Heidelberg in this PL. However, if Col plasmids are a true

horizontally acquired fitness factor for *S*. Heidelberg survival in PL, one would expect SH-116 to

survive longer than SH-2813 since the ancestral populations carry Col plasmids. This was not the

case in our study (Fig. 1a). SH-116 strains were only detected in 1 of 6 microcosms by direct

count and in 3 of 6 after enrichment in buffered peptone water (BPW) after 14 days incubation in

PL. In contrast, SH-2813 was recovered by direct count in all 6 microcosms (Supplementary

Table 1).

The dynamics observed in our experiment can be modeled according to Watve et al (79)

Rock-Paper-Scissors (RPS) model of sociobiological control of plasmid copy number. In our

scenario, SH-116 cells with lower copy ColE1 plasmids were displaced by those with higher

copy ColE1 plasmids mobilized from PL microbiota until the point at which the intrinsic survival

rate of *S*. Heidelberg was greatly reduced (Fig. 1a). In contrast, a stable co-existence of SH-2813

of low, intermediate and high copy ColE1 plasmids was observed in the SH-2813 population,

since they are invading strains lacking Col plasmids to begin with.

This observation was corroborated by qPCR on extracted DNA from PL with Col plasmid

gene primer sets (Supplementary Table 9) . At day 14, the concentrations of targeted genes were

several orders of magnitude higher in SH-116 microcosms than SH-2813 (Supplementary Fig.

6), which supports the notion that the original SH-116 strains may have been displaced by those

carrying higher copy ColE1 plasmids.

In this work, we demonstrate that ColE1 plasmids acquired from PL contributed to the

persistence of *Salmonella enterica* serovar Heidelberg, however the identity of the bacterial

donors of these plasmids remains unknown. Future studies from our laboratory will attempt to

provide answers.

ACKNOWLEDGEMENTS

We are grateful to Dr. Casey Ritz for providing us with the poultry litter used in this study

and for Caroline Plymel for her help with sample analysis. We also appreciate Steven Lankin and

Mary Katherine Crews for manuscript review. Any opinions expressed in this paper are those of

the authors and do not necessarily reflect the official positions and policies of the USDA and any

mention of products or trade names does not constitute recommendation for use. The authors

declare no competing commercial interests in relation to the submitted work.

Supplementary Figures

Figure S1. Experimental design and microcosm sampling.

Figure S2. ProgressiveMauve alignment of (**a**) IncX1 (**b**) ColE1-6kb and (**c**) ColE1-4kb

plasmids from selected strains.

Figure S3. Circular maps of (**a and b**) ColE1-6kb (**c and d**) ColE1-4kb and (**d and e**) ColpVC

plasmids carried by evolved strains used in competition and conjugation experiments.

Figure S4. Gene expression of selected Col plasmid genes determined by qRT-PCR for (**a**) SH-

116_{anc} (**b**) SH-116_{evol} and (**c**) SH-2813_{evol} strains.

Figure S5. Circular map of IncX1 plasmid of strain SH-33A0.5-116-Day7-PLE.

Figure S6. Intact prophages in sequenced genomes of *S*. Heidelberg.

Figure S7. Concentration of selected Col plasmid genes determined using qPCR for (**a**) SH-

2813-PLE and (**b**) SH-116-PLE poultry litter microcosms.

Supplementary Tables

Table S1. Bacteria concentration in poultry litter microcosms.

Table S2. Physicochemical properties of poultry litter microcosms.

Table S3. Metadata on isolates sequenced.

Table S4. Identified plasmid topology.

Table S5. Single nucleotide variants shared among Col-plasmid bearing strains.

Table S6. Plasmid copy number determined by whole genome sequencing coverage.

Table S7. Antibiotic susceptibility test results.

Table S8. Metadata on NCBI *S*. Heidelberg strains carrying IncX1 plasmid.

Table S9. Primers used in this study.

Supplementary Files

Supplementary File 1. Quality assessment and coverage of sequenced SH-2813 strains.

Supplementary File 2. Quality assessment and coverage of sequenced SH-116 strains.

Supplementary File 3. Disc diffusion inhibition zone measurements.

References:

1.   USDA, NASS. Poultry production and value - 2013 summary. ISSN:1949-1573 ed2014.
p. 1-14.
2.   MacDonald JM. The economic organization of U.S. broiler production. Washington,
D.C.: Economic Research Service, USDA; 2008 June 2008. Report No.: EIB- 38 Contract No.:
EIB- 38.
3.   Wang L, Lilburn M, Yu Z. Intestinal Microbiota of Broiler Chickens As Affected by
Litter Management Regimens. Front Microbiol. 2016;7:593.
4.   Dunlop MW, McAuley J, Blackall PJ, Stuetz RM. Water activity of poultry litter:
Relationship to moisture content during a grow-out. J Environ Manage. 2016;172:201-6.
5.   Cressman MD, Yu ZT, Nelson MC, Moeller SJ, Lilburn MS, Zerby HN. Interrelations
between the Microbiotas in the Litter and in the Intestines of Commercial Broiler Chickens.
Applied and Environmental Microbiology. 2010;76(19):6572-82.
6.   Casey R, Fairchild B, Lacy M. Litter quality and broiler performance. In: Bulletin CeS,
editor. 1267 ed2005.
7.   Lovanh N, Cook KL, Rothrock MJ, Miles DM, Sistani K. Spatial shifts in microbial
population structure within poultry litter associated with physicochemical properties. Poultry
Science. 2007;86(9):1840-9.

8.   Schefferle HE. The microbiology of built up poultry litter. J Appl Bacteriol.
1965;28(3):403-11.

9.   Roll VF, Dai Pra MA, Roll AP. Research on Salmonella in broiler litter reused for up to
14 consecutive flocks. Poult Sci. 2011;90(10):2257-62.

10.   Opara OO, Carr LE, Russek-Cohen E, Tate CR, Mallinson ET, Miller RG, et al.
Correlation of water activity and other environmental conditions with repeated detection of
Salmonella contamination on poultry farms. Avian Dis. 1992;36(3):664-71.

11.   Rigby CE, Pettit JR. Some Factors Affecting Salmonella-Typhimurium Infection and
Shedding in Chickens Raised on Litter. Avian Dis. 1979;23(2):442-55.

12.   Voss-Rech D, Trevisol IM, Brentano L, Silva VS, Rebelatto R, Jaenisch FRF, et al.
Impact of treatments for recycled broiler litter on the viability and infectivity of microorganisms.
Vet Microbiol. 2017;203:308-14.

13.   Hoffmann M, Zhao S, Pettengill J, Luo Y, Monday SR, Abbott J, et al. Comparative
genomic analysis and virulence differences in closely related salmonella enterica serotype
heidelberg isolates from humans, retail meats, and animals. Genome Biol Evol. 2014;6(5):1046-
68.

14.   Gieraltowski L, Higa J, Peralta V, Green A, Schwensohn C, Rosen H, et al. National
Outbreak of Multidrug Resistant Salmonella Heidelberg Infections Linked to a Single Poultry
Company. PLoS One. 2016;11(9):e0162369.

15.   Bekal S, Berry C, Reimer AR, Van Domselaar G, Beaudry G, Fournier E, et al.
Usefulness of High-Quality Core Genome Single-Nucleotide Variant Analysis for Subtyping the
Highly Clonal and the Most Prevalent Salmonella enterica Serovar Heidelberg Clone in the
Context of Outbreak Investigations. J Clin Microbiol. 2016;54(2):289-95.

16.   Hoffmann M, Zhao SH, Pettengill J, Luo Y, Monday SR, Abbott J, et al. Comparative
Genomic Analysis and Virulence Differences in Closely Related Salmonella enterica Serotype
Heidelberg Isolates from Humans, Retail Meats, and Animals. Genome Biol Evol.
2014;6(5):1046-68.

17.   Edirmanasinghe R, Finley R, Parmley EJ, Avery BP, Carson C, Bekal S, et al. A Whole-
Genome Sequencing Approach To Study Cefoxitin-Resistant Salmonella enterica Serovar
Heidelberg Isolates from Various Sources. Antimicrob Agents Chemother. 2017;61(4).

18.   Orlek A, Phan H, Sheppard AE, Doumith M, Ellington M, Peto T, et al. A curated dataset
of complete Enterobacteriaceae plasmids compiled from the NCBI nucleotide database. Data
Brief. 2017;12:423-6.

19.   Folster JP, Pecic G, Rickert R, Taylor J, Zhao S, Fedorka-Cray PJ, et al. Characterization
of multidrug-resistant Salmonella enterica serovar heidelberg from a ground turkey-associated
outbreak in the United States in 2011. Antimicrob Agents Chemother. 2012;56(6):3465-6.

20.   Labbe G, Ziebell K, Bekal S, Macdonald KA, Parmley EJ, Agunos A, et al. Complete
Genome Sequences of 17 Canadian Isolates of Salmonella enterica subsp. enterica Serovar
Heidelberg from Human, Animal, and Food Sources. Genome Announc. 2016;4(5).

21.   Labbe G, Edirmanasinghe R, Ziebell K, Nash JH, Bekal S, Parmley EJ, et al. Complete
Genome and Plasmid Sequences of Three Canadian Isolates of Salmonella enterica subsp.
enterica Serovar Heidelberg from Human and Food Sources. Genome Announc. 2016;4(1).

22.   Evans PS, Luo Y, Muruvanda T, Ayers S, Hiatt B, Hoffman M, et al. Complete Genome
Sequences of Salmonella enterica Serovar Heidelberg Strains Associated with a Multistate Food-
Borne Illness Investigation. Genome Announc. 2014;2(3).

23.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes:
a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol.
2012;19(5):455-77.

24.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: a fast phage search tool.
Nucleic Acids Res. 2011;39(Web Server issue):W347-52.

25.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.
2014;30(14):2068-9.

26.   Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017:
expansion and model-centric curation of the comprehensive antibiotic resistance database.
Nucleic Acids Res. 2017;45(D1):D566-D73.

27.   Orlek A, Phan H, Sheppard AE, Doumith M, Ellington M, Peto T, et al. Ordering the
mob: Insights into replicon and MOB typing schemes from analysis of a curated dataset of
publicly available plasmids. Plasmid. 2017;91:42-52.

28.   Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene
gain, loss and rearrangement. PLoS One. 2010;5(6):e11147.

29.   Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. BMC Bioinformatics. 2004;5:113.

30.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics. 2014;30(9):1312-3.

31.   Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler
transform. Bioinformatics. 2009;25(14):1754-60.

32.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence
alignment/map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.

33.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010;20(9):1297-303.

34.   San Millan A, Heilbron K, MacLean RC. Positive epistasis between co-infecting
plasmids promotes plasmid survival in bacterial populations. ISME J. 2014;8(3):601-12.

35.   Loftie-Eaton W, Bashford K, Quinn H, Dong K, Millstein J, Hunter S, et al.
Compensatory mutations improve general permissiveness to antibiotic resistance plasmids. Nat
Ecol Evol. 2017;1(9):1354-63.

36.   Administration FaD. The National Antimicrobial Resistance Monitoring System Manual
of Laboratory Methods. Second ed2015.

37.   Hudzicki J. Kirby-bauer disk diffusion susceptibility test protocol
2009.

38.   Institute CLS. Performance standards for antimicrobial susceptibility testing. 2009.

39.   Chattoraj DK. Control of plasmid DNA replication by iterons: no longer paradoxical.
Mol Microbiol. 2000;37(3):467-76.

40. Nordstrom K. Control of plasmid replication--how do DNA iterons set the replication
frequency? Cell. 1990;63(6):1121-4.

41. Bustamante P, Iredell JR. Carriage of type II toxin-antitoxin systems by the growing
group of IncX plasmids. Plasmid. 2017;91:19-27.

42. Page R, Peti W. Toxin-antitoxin systems in bacterial growth arrest and persistence. Nat
Chem Biol. 2016;12(4):208-14.

43. Rehman MA, Yin X, Persaud-Lachhman MG, Diarra MS. First Detection of a
Fosfomycin Resistance Gene, fosA7, in Salmonella enterica Serovar Heidelberg Isolated from
Broiler Chickens. Antimicrob Agents Chemother. 2017;61(8).

44. Tada T, Miyoshi-Akiyama T, Dahal RK, Mishra SK, Shimada K, Ohara H, et al.
Identification of a novel 6'-N-aminoglycoside acetyltransferase, AAC(6')-Iak, from a multidrug-
resistant clinical isolate of Stenotrophomonas maltophilia. Antimicrob Agents Chemother.
2014;58(10):6324-7.

45. Tada T, Miyoshi-Akiyama T, Shimada K, Shimojima M, Kirikae T. novel 6'-n-
aminoglycoside acetyltransferase AAC(6')-Iaj from a clinical isolate of Pseudomonas
aeruginosa. Antimicrob Agents Chemother. 2013;57(1):96-100.

46. Tada T, Miyoshi-Akiyama T, Shimada K, Dahal RK, Mishra SK, Ohara H, et al. A Novel
6'-N-Aminoglycoside Acetyltransferase, AAC(6')-Ial, from a Clinical Isolate of Serratia
marcescens. Microb Drug Resist. 2016;22(2):103-8.

47. Liakopoulos A, Geurts Y, Dierikx CM, Brouwer MS, Kant A, Wit B, et al. Extended-
Spectrum Cephalosporin-Resistant Salmonella enterica serovar Heidelberg Strains, the
Netherlands(1). Emerg Infect Dis. 2016;22(7):1257-61.

48. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al.
STRING v10: protein–protein interaction networks, integrated over the tree of life. Nucleic acids
research. 2014:gku1003.

49. Dy RL, Przybilski R, Semeijn K, Salmond GP, Fineran PC. A widespread bacteriophage
abortive infection system functions through a Type IV toxin-antitoxin mechanism. Nucleic Acids
Res. 2014;42(7):4590-605.

50. Actis LA, Tolmasky ME, Crosa JH. Bacterial plasmids: replication of extrachromosomal
genetic elements encoding resistance to antimicrobial compounds. Front Biosci. 1999;4:D43-62.

51. Francia MV, Varsaki A, Garcillan-Barcia MP, Latorre A, Drainas C, de la Cruz F. A
classification scheme for mobilization regions of bacterial plasmids. FEMS Microbiol Rev.
2004;28(1):79-100.

52. Smillie C, Garcillan-Barcia MP, Francia MV, Rocha EP, de la Cruz F. Mobility of
plasmids. Microbiol Mol Biol Rev. 2010;74(3):434-52.

53. Varsaki A, Moncalian G, Garcillan-Barcia Mdel P, Drainas C, de la Cruz F. Analysis of
ColE1 MbeC unveils an extended ribbon-helix-helix family of nicking accessory proteins. J
Bacteriol. 2009;191(5):1446-55.

54. Varsaki A, Lamb HK, Eleftheriadou O, Vandera E, Thompson P, Moncalian G, et al.
Interaction between relaxase MbeA and accessory protein MbeC of the conjugally mobilizable
plasmid ColE1. FEBS Lett. 2012;586(6):675-9.

55.   Hansen LH, Bentzon-Tilia M, Bentzon-Tilia S, Norman A, Rafty L, Sorensen SJ. Design
and synthesis of a quintessential self-transmissible IncX1 plasmid, pX1.0. PLoS One.
2011;6(5):e19912.

56.   Burmolle M, Norman A, Sorensen SJ, Hansen LH. Sequencing of IncX-plasmids
suggests ubiquity of mobile forms of a biofilm-promoting gene cassette recruited from Klebsiella
pneumoniae. PLoS One. 2012;7(7):e41259.

57.   Hall JPJ, Brockhurst MA, Dytham C, Harrison E. The evolution of plasmid stability: Are
infectious transmission and compensatory evolution competing evolutionary trajectories?
Plasmid. 2017;91:90-5.

58.   MacLean RC, San Millan A. Microbial Evolution: Towards Resolving the Plasmid
Paradox. Curr Biol. 2015;25(17):R764-7.

59.   Schroder G, Lanka E. TraG-like proteins of type IV secretion systems: functional
dissection of the multiple activities of TraG (RP4) and TrwB (R388). J Bacteriol.
2003;185(15):4371-81.

60.   Hayes F. Toxins-antitoxins: plasmid maintenance, programmed cell death, and cell cycle
arrest. Science. 2003;301(5639):1496-9.

61.   Lopatkin AJ, Meredith HR, Srimani JK, Pfeiffer C, Durrett R, You L. Persistence and
reversal of plasmid-mediated antibiotic resistance. Nat Commun. 2017;8(1):1689.

62.   Giraldo R, Fernandez-Tresguerres ME. Twenty years of the pPS10 replicon: insights on
the molecular mechanism for the activation of DNA replication in iteron-containing bacterial
plasmids. Plasmid. 2004;52(2):69-83.

63.   Chattoraj DK, Schneider TD. Replication control of plasmid P1 and its host chromosome:
the common ground. Prog Nucleic Acid Res Mol Biol. 1997;57:145-86.

64.   San Millan A, Escudero JA, Gifford DR, Mazel D, MacLean RC. Multicopy plasmids
potentiate the evolution of antibiotic resistance in bacteria. Nat Ecol Evol. 2016;1(1):10.

65.   Harrison E, Guymer D, Spiers AJ, Paterson S, Brockhurst MA. Parallel compensatory
evolution stabilizes plasmids across the parasitism-mutualism continuum. Curr Biol.
2015;25(15):2034-9.

66.   San Millan A, Pena-Miller R, Toll-Riera M, Halbert ZV, McLean AR, Cooper BS, et al.
Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids.
Nat Commun. 2014;5:5208.

67.   Santos-Lopez A, Bernabe-Balas C, San Millan A, Ortega-Huedo R, Hoefer A, Ares-
Arroyo M, et al. Compensatory evolution facilitates the acquisition of multiple plasmids in
bacteria. bioRxiv. 2017.

68.   Koskella B, Brockhurst MA. Bacteria-phage coevolution as a driver of ecological and
evolutionary processes in microbial communities. FEMS Microbiol Rev. 2014;38(5):916-31.

69.   Harrison E, Hall JPJ, Paterson S, Spiers AJ, Brockhurst MA. Conflicting selection alters
the trajectory of molecular evolution in a tripartite bacteria-plasmid-phage interaction. Mol Ecol.
2017;26(10):2757-64.

70.   Harrison E, Truman J, Wright R, Spiers AJ, Paterson S, Brockhurst MA. Plasmid
carriage can limit bacteria-phage coevolution. Biol Lett. 2015;11(8).

71.   Hoelzer K, Cummings KJ, Warnick LD, Schukken YH, Siler JD, Grohn YT, et al. Agar
disk diffusion and automated microbroth dilution produce similar antimicrobial susceptibility
testing results for Salmonella serotypes Newport, Typhimurium, and 4,5,12:i-, but differ in
economic cost. Foodborne Pathog Dis. 2011;8(12):1281-8.

72.   Diez-Aguilar M, Morosini MI, Tedim AP, Rodriguez I, Aktas Z, Canton R.
Antimicrobial Activity of Fosfomycin-Tobramycin Combination against Pseudomonas
aeruginosa Isolates Assessed by Time-Kill Assays and Mutant Prevention Concentrations.
Antimicrob Agents Chemother. 2015;59(10):6039-45.

73.   Vasse M, Noble RJ, Akhmetzhanov AR, Torres-Barcelo C, Gurney J, Benateau S, et al.
Antibiotic stress selects against cooperation in the pathogenic bacterium Pseudomonas
aeruginosa. Proc Natl Acad Sci U S A. 2017;114(3):546-51.

74.   Bruger EL, Waters CM. Bacterial Quorum Sensing Stabilizes Cooperation by Optimizing
Growth Strategies. Appl Environ Microbiol. 2016;82(22):6498-506.

75.   Leinweber A, Fredrik Inglis R, Kummerli R. Cheating fosters species co-existence in
well-mixed bacterial communities. ISME J. 2017;11(5):1179-88.

76.   Castaneda-Garcia A, Rodriguez-Rojas A, Guelfo JR, Blazquez J. The glycerol-3-
phosphate permease GlpT is the only fosfomycin transporter in Pseudomonas aeruginosa. J
Bacteriol. 2009;191(22):6968-74.

77.   Webber MA, Bailey AM, Blair JM, Morgan E, Stevens MP, Hinton JC, et al. The global
consequence of disruption of the AcrAB-TolC efflux pump in Salmonella enterica includes
reduced expression of SPI-1 and other attributes required to infect the host. J Bacteriol.
2009;191(13):4276-85.

78.   Baucheron S, Tyler S, Boyd D, Mulvey MR, Chaslus-Dancla E, Cloeckaert A. AcrAB-
TolC directs efflux-mediated multidrug resistance in Salmonella enterica serovar typhimurium
DT104. Antimicrob Agents Chemother. 2004;48(10):3729-35.

79.   Watve MM, Dahanukar N, Watve MG. Sociobiological control of plasmid copy number
in bacteria. PLoS One. 2010;5(2):e9328.

A

```
                               10         20         30         40         50         60         70         80         90        100
                    |          |          |          |          |          |          |          |          |          |          |
ColE1-6kb/1-79      M A F Y V D S T A K F L T K T T E P E G S V W R Y Y Y K - - - - P G - - - - - - - - - - - N T C P A - - - - S G I Y R C T N C G D E V T C N K D D L P P Q N H R Q H N V - P G D I R - - - - W E L V V M T K
SH-Ancestral-2813/1-8 - - - - - M D I R Q M N K T N L E H W R G L R K Q L W P G - - - - D A H L A D G E E I L Q A - D Y L A S F I A M A D - C V A I G F A D - - - A S I R H D Y V N G C D S S P V A F L E G H I F V L P S F R
K.pneumoniae/1-75   - - - - - - M A Q Y K Y G N Y L N Q S Q N E A Y D T L H S - - - - P G - - - - - - - - - - - D T C A N - - - - S G I Y R C E S C G D E I A S N K G N P F P Q N H Q H A P G S G P I R - - - - W K L I V W A Q
Thiomonas/1-75      - - - - - - M A S F K Y A I H L R Q D S N N A F D Q K H P - - - - P G - - - - - - - - - - - I V A P Y - - - - A G I Y R C V V C E E I G I A Q G H T L P P Q N H R Q H E V - P G R I E - - - - W Q L L V Q A H
Stenotrophomonas/1-92 - - - - - - M T G S A A T I R P A K A A D A V A A Q L R L G L W P D - - A D D P L E T L V A A L E - D A G A V F L A C A A G Q A I G F A E - - - V R L R H D Y V N G T D S S F V G F L E G H Y V Q P Q W Q
P.aeruginosa/10-108 A P S M A N T P V G N V V P C K T P D H Q V E G W L E L R L Q L W L D G S T E E P L P E M A A A C A E P D R F G Q F L F L S P G L A E G L V E - - - V A L R T D Y V N G T E S S P V A F L E G V F V V P A S R
```

B

C

Figure showing six bar charts titled kanamycin, Tobramycin, Gentamicin, Neomycin, Netilmicin, and Fosfomycin with axes labeled "No. of Isolates" and "Zone of Inhibition (mm)". Legend: SH-116-Day0, SH-116-Day14, SH-2813-Day0, SH-2813-Day14.