Title: Hotspot mutations and ColE1 plasmids contribute to the fitness of *Salmonella* Heidelberg

in poultry litter

¶Adelumola Oladeinde[1*], ¶Kimberly Cook[1], ¶Alex Orlek[2], &Greg Zock[1] &Kyler Herrington[3],

&Nelson Cox[4], &Jodie Plumblee Lawrence[1], &Carolina Hall[1]

Authors' Affiliation

[1]Bacterial Epidemiology and Antimicrobial Resistance Research Unit, U.S. National Poultry

Research Center, USDA-ARS, Athens, GA 30605, USA. [2]Nuffield Department of Medicine,

John Radcliffe Hospital, University of Oxford, Oxford, UK. [3] Department of Microbiology,

University of Georgia, Athens, GA 30602, USA. [4]Poultry Microbiological Safety and Processing

Research Unit, U.S. National Poultry Research Center, USDA-ARS, Athens, GA 30605, USA

Corresponding author* ade.oladeinde@ars.usda.gov

¶ These authors contributed equally to this work.

& These authors also contributed equally to this work.

ABSTRACT

Salmonella enterica subsp. *enterica* serovar Heidelberg (*S*. Heidelberg) is a clinically-important

serovar linked to food-borne illness, and commonly isolated from poultry. Investigations of a

large, multistate outbreak in the USA in 2013 identified poultry litter (PL) as an important extra-

intestinal environment that may have selected for specific *S*. Heidelberg strains. Poultry litter is a

mixture of bedding materials and chicken excreta that contains chicken gastrointestinal (GI)

bacteria, undigested feed, feathers, and other materials of chicken origin. In this study, we

performed a series of controlled laboratory experiments which assessed the microevolution of

two *S*. Heidelberg strains (SH-2813 and SH-116) in PL previously used to raise 3 flocks of

broiler chickens. The strains are closely related at the chromosome level, differing from the

reference genome by 109 and 89 single nucleotide polymorphisms/InDels, respectively. Whole

genome sequencing was performed on 86 isolates recovered after 0, 1, 7 and 14 days of

microevolution in PL. Only strains carrying an IncX1 (37kb), 2 ColE1 (4 and 6kb) and 1

ColpVC (2kb) plasmids survived more than 7 days in PL. Competition experiments showed that

carriage of these plasmids was associated with increased fitness. This increased fitness was

associated with an increased copy number of IncX1 and ColE1 plasmids. Further, all Col

plasmid-bearing strains had hotspot mutations in 37 loci on the chromosome and in 3 loci on the

IncX1 plasmid. Additionally, we observed a decrease in susceptibility to tobramycin, kanamycin,

gentamicin, neomycin and fosfomycin for Col plasmid-bearing strains. Our study demonstrates

how positive selection from poultry litter can change the evolutionary path of *S*. Heidelberg.

INTRODUCTION

The United States (U.S.) is the world's largest producer of poultry with over 8 billion broilers produced yearly (1). Maintaining a safe supply of poultry is increasingly important as demand increases, production becomes more intensive and facilities become antibiotic free. Broilers (around 22,000 broilers per house) are typically raised in open plan barns on a deep floor of bedding materials (typically wood shavings, rice hulls or saw dust (1, 2). Poultry litter (PL) is a mixture of bedding materials and chicken excreta that contains chicken gastrointestinal (GI) bacteria, undigested feed, uric acid, feathers and other materials of chicken origin (3-6). As the cost of bedding materials increase, land nutrient management requirements become more stringent, and broiler production and turnover times decrease, farmers have chosen to increase the number of flocks raised on litter before cleanout. This practice is accepted in production environments, but the effect of litter reuse on microbial populations has not been well studied.

Available data suggest that litter re-use can result in significant changes in the microbiological, chemical and physical composition of litter (3, 7, 8). In addition, reused litter has been found to harbor fewer yeasts, molds, coliform bacteria and *Salmonella* compared to fresh litter (9, 10). However, fecal shedding of food borne pathogens such as *Salmonella* into poultry by colonized birds can also be a source of new infection or re-inoculation. For instance, *Salmonella* in 131-day-old litter was shown to colonize 82% of day-old chicks after placement for 32 days (11) and *Salmonella enterica* subsp. *enterica* serovar Heidelberg (*S.* Heidelberg) has been shown to persist in recycled litter, even after litter treatment (12). *Salmonella* Heidelberg is a clinically-important serovar, linked to food-borne illness and commonly isolated from poultry. After *S.* Typhimurium, *S.* Heidelberg is the serovar of *Salmonella* most often associated with *Salmonella*-related deaths in the United States (13). In

2013, *S*. Heidelberg was responsible for a large multistate foodborne outbreak (634 cases) linked

to consumption of chicken (14). Epidemiological and environmental investigations identified PL

as an important extra-intestinal environment that may have selected for specific *S*. Heidelberg

strains (14).

Salmonella Heidelberg is highly clonal and cannot be subtyped by conventional pulsed-

field gel electrophoresis (PFGE) which poses an inherent difficulty when differentiating strains

responsible for foodborne outbreaks from those associated with background sporadic cases (15-

17). High quality core genome single nucleotide variant (hqSNV) analysis on whole genome

DNA sequences (WGS) of *S*. Heidelberg has been recommended as a better approach to

distinguish outbreak strains (15). However, core genome SNV is non-discriminatory between

isolates harboring horizontally acquired antibiotic resistance genes (ARG) versus susceptible

ones. Multidrug resistant *S*. Heidelberg isolates tend to cluster with susceptible isolates during

outbreaks and differ by only 0 - 4 SNVs (16, 17). In a study by Edirmanasinghe et al. (17), the

authors reported 65 out of 113 sequenced *S*. Heidelberg encoded the AmpC beta-lactamase

(bla$_{CMY-2}$) gene on an incompatibility group I1 plasmid (IncI1) and were resistant to important 3rd

generation cephalosporins. On the other hand, susceptible isolates (n=38) carried multiple

variants of Col plasmids and encoded no known ARG. The function of these Col plasmids is not

clear but they seem to be mobilized by co-resident conjugal plasmids, such as IncI1 and IncX,

two plasmids frequently found among *S*. Heidelberg isolates (16-19).

The majority of *S*. Heidelberg isolates sequenced and available in the National Center for

Biotechnology Information (NCBI) database were recovered from either food or clinical samples

(18, 20-22), leaving a knowledge gap in our understanding of strains of environmental origin.

The poultry litter microbiome may carry mobile genetic elements (MGE) including plasmids that

can be horizontally transferred to *S*. Heidelberg upon introduction into PL. These newly acquired

MGE may present a fitness advantage to permissive recipients, further enabling the persistence

of *S*. Heidelberg in poultry environments.

In this study, we investigated the role of PL on the fitness of two *S*. Heidelberg strains with

indistinguishable PFGE patterns, both recovered from chicken carcass in two states in USA.

Both strains harbor an IncX1 plasmid (37 kb) but differ in the occurrence of Col plasmids. We

performed a series of controlled laboratory experiments to assess the microevolution of these

strains in PL that was previously used to raise three flocks of broiler chickens. Our data suggests

that acquisition of multiple ColE1 plasmids in tandem with mutations in *S*. Heidelberg

chromosome and the IncX1 plasmid may have contributed to the persistence of *S*. Heidelberg in

the PL used in this study. Additionally, the presence of Col plasmids was associated with a

decreased susceptibility to tobramycin, kanamycin, neomycin and fosfomycin.

MATERIALS AND METHODS

Poultry Litter for microcosm evolution studies. Poultry litter (PL) for microcosm studies was

provided by the University of Georgia poultry research center in Athens, GA, USA. The PL was

previously used to raise 3 flocks of broiler chickens under simulated commercial poultry

production conditions. The litter was collected approximately 2 weeks after the third flocks of

birds were removed. The PL bedding material was composed of pine shavings.

Bacterial strains and preconditioning. *S*. Heidelberg strains (SH-2813 and SH-116) used in

this study were originally recovered from broiler chicken carcass rinsates. The

genotypic/phenotypic characteristics of these strains are provided in Table 1. The strains were

preconditioned in poultry litter extract (PLE) media prepared as described in supplementary

material or in Brain Heart Infusion broth (BHIB; BD Difco, Sparks, MD) before inoculation into

PL. For each strain, 5 single colonies from Sheep Blood Agar (SBA) (Remel Inc, San Diego,

CA) were transferred to 10 ml of PLE and BHIB, and incubated at 37 °C for 24 h. Serial batch

culturing was carried out for approximately 50 generations in PLE and BHIB by transferring 100

μl of culture to 10 ml of fresh PLE and BHIB (100-fold dilution) to yield approximately 10 gen.

d^{-1}. After 5 days, the population was used as inoculum for evolution experiments in PL

microcosms.

Experimental microevolution in PL. Poultry litter was collected in a 5-gallon bucket and

thoroughly mixed before 30 g aliquots were placed into 48 250 ml Nalgene wide jars and 1X

PBS (3-4 ml) was added to achieve 30 % moisture. Microcosms were incubated for 72 h at 37°C

before being individually inoculated with approximately 10^6 colonies forming units (CFU) of

either SH-2813 or SH-116 preconditioned previously in PLE or BHIB. These 4

groups/treatments were as follows: SH-2813 + BHIB, SH-2813 + PLE, SH-116 +BHIB and SH-

116 +PLE. Inoculated microcosms were thoroughly mixed and then incubated for 14 days at 37

°C (Supplementary Fig. 1). A control microcosm remained un-inoculated. Three microcosms per

group were destructively sampled at days 0, 1, 7 and 14.

Poultry litter microcosm sampling. Poultry litter samples (25 g) were mixed with 150 ml 1X

PBS and 5-10 ml glass beads (Fisher Sci: S800242) then homogenized for 10 min at 450 rpm in

a hand wrist shaker (Boekel Scientific, Feasterville, PA). The PL mixtures were allowed to settle

for 5 min to allow settling of large debris/pine shavings and then passed through cheesecloth and

hand pressed to recover eluates which were used for downstream microbiological and chemical

analysis. The moisture content of each PL sample was determined gravimetrically by drying

approximately 1 g of PL at 107 °C for 24 h.

Microbiological methods. Recovered PL eluate was serially diluted in 1X PBS and plated on

selective or general-purpose growth media. We used Brilliant Green Sulfa (BGS) and XLT-4

agar for *Salmonella* spp., Chromagar ECC for coliforms, Enterococcosel for *Enterococcus* spp.

and Plate Count Agar (PCA) for total bacteria. For enrichment and recovery of injured/viable

Salmonella spp., 1 ml of eluate was transferred to 20 ml of 1 % Buffered Peptone Water (BPW)

and incubated 24 h before being plated onto XLT-4 agar. All agar plates and broth tubes were

incubated at 37 °C for 24 – 48 h before colony forming units were enumerated. Reported counts

were log-normalized and reported per gram PL dry weight. At least three single colonies

representing three unique clones of *Salmonella* from each microcosm were randomly chosen

from XLT-4 plate dilutions and archived in 30 % Luria Bertani (LB) glycerol at -80 °C.

DNA sequencing and bioinformatics. We sequenced 86 evolved *S*. Heidelberg isolates

encompassing clones recovered from the four treatments. Isolates were selected randomly from

the highest and lowest dilution of XLT-4 plates; and when populations decreased below limit of

quantification (LOQ = 10 CFU/g dry weight of PL), isolates were selected after 24 h enrichment

in BPW. Genomic DNA (gDNA) was extracted from pure isolate cultures using FastDNA Spin

Kit (Mp Biomedicals, Solon, Ohio) and quantified with a Qubit Fluorometer (ThermoFisher

Scientific). Whole genome sequencing (WGS) libraries were prepared using MiSeq Nextera XT

library preparation kit (Illumina, Inc., San Diego, CA). Sequencing was performed on the

Illumina MiSeq platform with 250-bp paired end reads using the MiSeq reagent V2 (500 cycles).

All isolates were sequenced for an average coverage of 100X. Sequence reads were assembled

de novo into contigs using SPAdes (23). The quality of assembled genomes was accessed using

Quality Assessment Tool for Genome Assemblies (QUAST). (For detailed information on

assembly, coverage and quality of sequenced genomes see Supplementary Files 1 and 2).

The original strains of SH-2813 and SH-116 are referred to as ancestral strains, while

isolates from PL evolution experiments are referred to as evolved strains. Assembled contigs

were submitted to the Center for Genomic Epidemiology's multilocus sequence type (MLST)

and PlasmidFinder modules to determine the whole-genome sequence types (ST) and existing

plasmid replicon types. Prophages were identified using PHAST (24) and sequences for intact

phages were extracted, and annotated with Prokka (25). The Comprehensive Antibiotic

Resistance Database (CARD) was used for the identification of resistance genes present on

plasmid and chromosome (26).

Plasmids were classified into MOB families using methods described previously (27). To

investigate the topology of contigs identified with plasmid replicon, we used a custom script to

inspect the path in the SPAdes FASTG assembly graph file that corresponded to the plasmid

contig identified in the FASTA file; the contigs.paths file was used to determine correspondence

between FASTA contigs and FASTG edges. Plasmid contigs were assigned as 'complete

circular'; 'complete non-circular', i.e. linear but not connected to other contigs; and 'non-

complete', i.e. linear and connected to other contigs. (see Supplementary Methods for detailed

information). Complete circular plasmids were annotated using Prokka (25) and mapped with

Snapgene   v   4.1.2.   When   coding   DNA   sequence   (CDS)   was   annotated   as

hypothetical/uncharacterized,   we   performed   a   PSI-BLAST   search   with   predicted   protein

sequences run to ≤3 iterations. Plasmids were analyzed by comparative analysis using

ProgressiveMAUVE (28) and CDS predicted in regions with marked differences were aligned

with MUSCLE and MAFFT (29, 30). Phylogenetic tree was produced from aligned conserved

relaxase or mobilization protein sequences using the maximum likelihood method implemented

in   Randomized   Axelerated   Maximum   Likelihood   (RAxML)   v   8   (31)   using   the

PROTGAMMA/BLOSUM62 model and a rapid bootstrapping run to generate 1000 replicates.

Samples with identical sequences were removed before phylogenetic tree was constructed. Trees

generated were rooted with MbeA and MbeC sequences of *E. coli* (Accession number:

NC001371).

Single Nucleotide Polymorphism (SNPs) and InDel analysis. SNPs/InDels were determined

by mapping raw sequence reads to the chromosome and plasmid genome of *S*. Heidelberg strain

AMR588-04-00435 available in NCBI under accession numbers CP016573 and CP016575,

respectively, using Burrows-Wheeler Aligner (BWA) (32). SAM file sorting, and removal of

PCR duplicates was done using SAMtools (33). Genome Analysis ToolKit (34) with a minimum

mapping quality of 30 and a minimum base quality of 30 was used for SNP identification.

Manipulation of variant call format files (VCF) generated was done using vcftools v 0.1.12b

(35). Shell script used   and VCF files   are   available   from   the   Dryad   Digital

Repository: https://doi.org/10.5061/dryad.pc6tp4q.

To obtain phylogenetic trees, SNPs and InDels mapping to plasmid sequences were removed

and VCF files for SH-2813 and SH-116 evolved isolates were merged. The merged files were

converted to a PHYLIP multiple alignment file using PGDSpider (36) and isolates with identical

SNPs were removed before phylogenetic tree was constructed. The PHYLIP file was analyzed

using maximum likelihood method implemented in RAxML v. 8 with the number of bootstrap

replicates criteria set to extended majority rule consensus tree (`-N autoMRE`). We used the

Jukes-Cantor (37) model of nucleotide substitution and GAMMA model of rate heterogeneity

because they were the simplest model of sequence evolution predicted using jModelTest (38).

Trees generated were rooted with either ancestral strain of SH-2813 or SH-116.

Growth assay and plasmid copy number determination. Plasmid copy number (PCN) and

differential expression of selected transcripts was determined by performing a growth assay on

ancestral and evolved strains. Three populations were initiated from three single colonies from

strains SH-2813$_{anc}$, SH-2813$_{evol}$, SH-116$_{anc}$, and SH-116$_{evol}$. An individual colony was

transferred to 20 ml of Mueller Hinton Broth (MHB) in 50 ml centrifuge tube and incubated

overnight at 37 °C. Following overnight growth, 6 technical replicate populations (18 per strain)

were established from each culture by transferring 3 ml to 50 ml centrifuge tubes containing 17

ml MHB. Cultures were incubated without shaking at 37 °C for 24 h. Optical density (OD$_{600nm}$)

was periodically measured in the 3 populations with a Nanodrop 2000 Spectrophotometer

(ThermoFisher Scientific). Bacteria concentration was determined for 2 populations at 0.5, 4 and

24 h via serial dilution plating on R2A agar. Additionally, bacteria culture (~ 20 ml) was filtered

onto 0.2 µm pore size polycarbonate filters (MilliporeSigma, Burlington, MA) which were

transferred to Lysing Matrix E and B tubes (MP Biomedicals, Solon, Ohio) for genomic DNA

and RNA isolation respectively; and saved at 80 °C until use.

Five primer sets specific to the glyceraldehyde-3-phosphate dehydrogenase, replication

initiation protein, putative toxin-antitoxin, macrophage stimulating factor, a replication protein

was designed and denoted *gapA*, repB_IncX1 Abi_ColE1-6kb, MSF_ColE1-4kb and rep

respectively. The *gapA* is a single-copy gene of *S*. Heidelberg chromosomal DNA, while repB,

Abi, MSF and rep are single-copy genes of IncX1, ColE1_6kb, ColE1_4kb and ColpVC_2kb,

respectively. Real-time qPCR amplification and analysis were performed as previously described

with a CFX96 Touch Real-Time PCR Detection System (Bio-Rad Inc., Hercules, CA).

Specificity of primer pairs was verified by melting curve analysis and amplification factor for all

assays was between 1.75 – 2.03 (Supplementary Table 9). PCN was determined as the copy ratio

of plasmid encoded genes to *gapA* (39).

Gene expression. For qRT-PCR, total RNA was isolated from membrane filters as described by

ref (40) using FastRNA Pro Blue Kit (MP Biomedicals, Solon, Ohio). Purified RNA was treated

with 4U DNA-free DNase (ThermoFisher Scientific, MA, USA) to remove genomic DNA

contamination and quantified with a Qubit Flourometer (Thermo Fisher Scientific, MA, USA).

DNase treated RNA was reversed transcribed with 100 U of high capacity cDNA reverse

transcription kit (ThermoFisher Scientific, MA, USA) per manufacturer's instructions.

Three primer sets targeting plasmid genes were designed in addition to the ones used for

PCN determination. The primers target the toxin-antitoxin proteins (*relBE*) present on IncX1 and

a putative aminoglycoside 6-N-acetyltransferase (aac- (6')) present on ColE1_6kb. qRT-PCR

was performed on generated cDNA as described for qPCR assay. Reaction mixtures (20 µl) for

all assays contained 1X SsoAdvanced Universal SYBR Green Supermix (Bio-Rad Inc.,

Hercules, CA), 250 nM (each) primers, and 2 µl (20 -100 ng) of cDNA. Data were analyzed

using the $2^{-\Delta\Delta CT}$ method described by Livak and Schmittgen (41). For each plasmid target gene,

the ratio of expression between ancestral and evolved strain was normalized to the expression of

chromosome-encoded *gapA* and *gyrB* reference genes. The statistical significance of differences

in gene expression relative to ancestor was determined using Wilcoxon signed-rank Test.

Competitive fitness and conjugation assay. The fitness of three strains carrying the 3 Col

plasmids (SH-2813$_{evol}$, SH-116$_{anc}$, and SH-116$_{evol}$) was determined relative to SH-2813$_{nal}$

nalidixic resistance with no Col plasmid. The SH-2813$_{nal}$ strain is isogenic to SH-2813$_{anc}$ but was

made resistant to 200 ppm nalidixic acid by spontaneous mutation in *gyrA*. Five colonies from

SBA for each strain were incubated in 9 ml of fresh MHB at 37 ºC for 20 h. Thereafter, cultures

of the strains were mixed at a ratio of 1:1 (~500 µl of strain under study to 500 µl of SH-2813$_{nal}$).

We assumed that each strain had the same initial concentration because 24 h population was not

significantly different (Table 3). Mixtures were diluted 10,000-fold in 50 ml centrifuge tubes (4

technical replicates per mixture) containing 20 ml of MHB and competed for 24 h at 37 °C

without shaking. The final population was measured by serial plating onto R2A agar

supplemented with and without 32 ppm of nal. The fitness of the strains carrying Col plasmids

relative to the SH-2813-nal strain was determined as described by ref (42):

$$W_{strain} = \frac{\log_e \left(N_{final,strain} | N_{initial,strain} \right)}{\log_e \left(N_{final,nal} | N_{initial,nal} \right)} \qquad (1)$$

Where W_{strain} is the fitness of the strain carrying Col plasmids, $N_{initial,strain}$ and $N_{final,strain}$ are

the numbers of cells (in colonies forming units (CFU)) of the strain carrying the Col plasmids

before and after competition, and $N_{initial,nal}$ and $N_{final,nal}$ are the numbers of cells of SH-2813-

nal before and after competition. To determine $N_{final,strain}$ we randomly screened 28 colonies

from R2A without nal on R2A and Brilliant Green Sulfur (BGS) supplemented with 32 and 200

ppm of nalidixic acid, respectively. From here, the fractions of Col plasmids donor within each

population was estimated by scoring the absence of growth on both R2A and BGS supplemented

with nal. To assess possible fitness cost associated with *gyrA* mutation, SH-2813$_{nal}$ was grown in

single culture and plated on R2A with and without nal. The ratio of cells that grew on R2A

without nal versus R2A with nal was 1.00 (s.e. =0.12, n =3).

To pool the information from the competition assay with conjugation assay we employed the

same data set to determine the conjugation rate constant for the three Col plasmid types present

during the competition assay. Conjugation rate constants were determined as described by

Loftie-Eaton et al (43). Briefly, conjugation between plasmid-containing donors (D) and

plasmid-free recipients (R) was measured under the same growth conditions used for the

competition assay. To distinguish D and R, we randomly screened 56 colonies from R2A

supplemented with 32 ppm nal for plasmid-specific gene. To do this, colonies were selected from

the lowest and highest dilution plate (10^{-5} and $10^{-7)}$ for each strain of interest, and resuspended in

96-well PCR plates containing 100 µl DEPC-treated water. Thereafter a quick-boil DNA

extraction at 95 °C for 10 mins was done. Following the extraction, qPCR was used to estimate

the proportion of positive "transconjugants" to Col plasmid-free recipients. The conjugation rate

constant γ (in ml (c.f.u x t)$^{-1}$) for the 3 Col plasmids in each competition assay was then

determined using the following equation and as described by ref (43):

$$\gamma = \psi \ln\left(1 + \frac{T}{R}\frac{N}{D}\right)\frac{1}{N-N_0} \qquad\qquad (2)$$

Where ψ is the maximum growth rate for the combined donor and recipient cultures, which is

calculated from two bacteria concentration measurements taken during exponential phase and

quantified on R2A without Nal, and calculated as follows:

$$\psi = \frac{\ln\dfrac{N_{final}}{N_{initial}}}{t_2 - t_1}$$

Total initial cell densities (*No*) and final (*N*) are the number of cells growing on R2A without nal

at 0 and 24 h respectively. T is the number of transconjugants estimated from cells growing on

R2A with nal.

Antibiotic susceptibility testing. We performed antimicrobial susceptibility testing (AST) on

selected evolved strains following National Antimicrobial Resistance Monitoring System

(NARMS) protocols for Gram-Negative bacteria (44) and Kirby-Bauer disc diffusion assay (45).

For more details on the disc diffusion protocol used, see Supplementary Methods.

For categorizing susceptibility, breakpoints were adopted from NARMs. There is no

standardization by Clinical and Laboratory Standards Institute (46) for tobramycin, kanamycin,

neomycin, netilmicin, and fosfomycin by disc diffusion for *S*. Heidelberg. Therefore, results

were interpreted with diameter difference in zone of inhibition between isolates.

Poultry litter DNA extraction. Two hundred and fifty milligrams of PL from each microcosm

were transferred to PowerBead tubes in triplicate (Qiagen Inc, Germantown, MD) and stored at -

20°C until extraction, which occurred within 2 months of sampling. DNA was extracted using a

Qiagen DNeasy Power-Soil DNA isolation kit, with the following modifications to the

manufacturer's instructions: (i) bead beating was conducted twice at 1800 rpm for 60 s, using a

Fastprep-96 bead beater (MP Biomedicals, Solon, Ohio); and (ii) cleanup step with 800 µl of

100% ethanol was performed twice before cleanup with Solution C5. After elution of extracted

DNA, an additional cleanup with Zymogen PCR inhibitor removal kit (Zymo Research Corp.

Irvine, CA) was employed as per manufacturer's instructions. Finally, all 3 purified DNA

extracts from each microcosm was pooled and concentrated in an Eppendorf Vacufuge Plus

Concentrator (Fisher Sci) at 30 °C. Pelleted DNA was reconstituted in 100 µl 1X TE buffer (10

mM Tris-HCL, 1 mM EDTA, pH 7.5) and concentration was determined fluorometrically using

Qubit Fluorometer (ThermoFisher Scientific, MA, USA). DNA was stored at -80 °C until use.

Statistical analysis. The two *S*. Heidelberg strains are the experimental units and PL represents

the sampling unit for this study. Continuous variables did not meet the assumption of a normal

distribution; therefore comparisons between strains or treatment were performed using Wilcoxon

signed-rank Test. A principal component analysis (PCA) was performed on PL physico-chemical

parameters (log-normalized) and *S*. Heidelberg concentrations to derive uncorrelated components

representing most of the data information and analyze treatment effects on *S*. Heidelberg

survival. A multiple linear regression was used to model the relationship between IncX1 and Col

plasmid copy numbers derived from whole genome sequencing coverage. Analyses were

performed using R (version 3.4.1).

Ethics statement. The authors did not physically interact with chickens raised on the poultry

litter used in this study, thereforewe were exempt from university guidelines and USDA/NIH

regulations regarding animal use. Institutional Animal Care and Use Committee (IACUC) -

UGA Office of Animal Care and Use.

Data availability. All sequencing data are available under NCBI Sequence Read Archive (SRA)

BioProject: PRJNA448609.

RESULTS

Experimental evolution system. The survival of two *S*. Heidelberg strains (SH-2813 and SH-

116) in used PL was evaluated over a period of 14 days (Supplementary Fig. 1). We chose *S*.

Heidelberg for its increasing role in food-borne outbreaks and its carriage of Col-like plasmids

(14, 16, 17, 22, 47). The PL was previously used to raise 3 *Salmonella*-free flocks. In addition,

30 g of uninoculated PL was analyzed by culture method immediately after collection and at the

start of the experiment (Day 0) for *Salmonella* isolation. No Salmonella was detected by direct

culture on selective media or by enrichment of PL in BPW. The strains used were recovered

from chicken carcass from two states in the USA and are closely related at the chromosomal

level, differing from the *S*. Heidelberg reference genome used by only 109 and 89 SNPs/InDels.

In addition, both strains carried an IncX1 plasmid and SH-116 carried 2 ColE1-like plasmids (4

and 6 kb) and a cryptic ColpVC plasmid (2 kb) (Table 1). The strains were chosen to investigate

the role of Col-like plasmids on the fitness of *S*. Heidelberg in PL. We hypothesized that *S*.

Heidelberg strains carrying Col plasmids will persist longer in PL.

Table 1. Characteristics of the strains used in this study

Strain ID	Serotype	Plasmid Profile (size)	*Resistance profile	#Antimicrobial resistance genes	¶Unique Single Nucleotide Variants (SNV)
SH-2813	ST15	IncX1 (37kb)	Ery, Tyl,	AAC (6')-Iy, FosA7	109
SH-116	ST15	IncX1 (37 kb), ColE1(6 kb) ColE1 (4 kb) ColpVC (2 kb)	Ery, Tyl, Pen	AAC (6')-Iy, FosA7	89

Note. - Ery = Erythromycin; Pen = Penicillin; Tyl= Tylosin

* Susceptible to all 14 antibiotics on the NARMs Gram-negative AST panel. Ery, Pen and Tyl

are on the NARMs Gram-positive AST panel.

Present on chromosome

¶ SNVs unique compared to reference genome (CP0165673).

The concentration of the two *S*. Heidelberg strains preconditioned in either PLE or BHIB

decreased by several orders of magnitude in all microcosms and their survival dynamics differed

by preconditioning media and strain-type (Fig. 1a).

Figure 1. Survival curve and chromosomal mutation based phylogeny of evolved *S*.

Heidelberg strains. (**a**) *S*. Heidelberg concentration from poultry litter microcosms reported in

colony forming units (CFU) and normalized by litter dry weight. (**b**) Venn diagram showing the

total number of chromosomal SNPs/InDels shared between isolates with hotspot mutations. (**c**)

Left. Maximum-likelihood tree of SH-2813 and SH-116 isolates based on 213 SNPs/InDels.

Clades and taxa highlighted in green denote isolates without hotspot mutations or Col plasmids,

while clades highlighted with cyan boxes carry hotspot mutations and 3 Col plasmids. Taxa in

parenthesis represent isolates with identical SNPs/InDels and not used in the reconstruction of

the tree shown. Isolates with sample ID's 1- 24 are evolved isolates from strain SH-2813 and

ID's 25- 48 are evolved isolates from strain SH-116. The numbers shown next to the branches

represent the percentage of replicate trees where associated taxa cluster together based on ~200

bootstrap replicates. Numbers (1- 18) denote arbitrarily assigned clade numbers. **Right**:

Corresponding venn diagram comparing the total number of SNPs/InDels shared across time

points. Tree was rooted with ancestral SH-2813.

The concentration of SH-2813-PLE, decreased by 4-log within 1 day of inoculation into PL but

rebounded 1.5-log after 7 days. In contrast, the concentration of SH-2813-BHIB was below the

LOQ within 1 day of inoculation into PL and increased approximately 4-log by day 7. The final

concentration of SH-2813-PLE and SH-2813-BHIB in PL was log 2.5 ± 0.03 CFU g^{-1}. On the

other hand, the concentration of SH-116-PLE decreased gradually for 14 days while SH-116-

BHIB was below LOQ within 1 day of inoculation into PL and increased by approximately 3-

log at day 7 (Fig. 1a). The final concentration for SH-116 was below our LOQ for direct culture

counts but the strain was isolated at varying levels following enrichment in buffered peptone

water (BPW) (Supplementary Table 1).

We determined the pH, moisture, nutrients, and cations present in PL during the 14-day

experiment (Supplementary Table 2). The litter pH and moisture level was 8.5 ± 0.23 and $32.3 \pm$

2 % at day 0 and 8.1 ± 0.09 and 16.7 ± 1 % at day 14. We observed significant differences for

pH and moisture between PLE and BHIB microcosms (*p*. value <0.001; Wilcoxon signed-rank

Test), with BHIB having higher moisture and pH (moisture =36 ± 2.1 % ; pH= 8.75 ± 0.1) than

PLE (moisture =25.4 ± 4 %; pH=8.17 ± 0.1) from day 0 to day 7. Also, percent reduction of total

organic carbon (TOC), total carbon (TC), inorganic carbon (IC), total nitrogen (TN), ammonium

(NH_4), aluminum (Al), copper (Cu), Iron (Fe) and Zinc (Zn) in BHIB microcosms was higher

than PLE microcosms after 14 days (Supplementary Table 2). To access the relationship

between these physico-chemical parameters and *Salmonella* concentration, we performed a

principal component analysis (PCA) on all variables measured. Two components of the PCA

performed had eigenvalues higher than one and captured 68.8% of the variance in the data

(Supplementary Fig. 2). Most variables were positively correlated with component 1 including

pH ($\rho = 0.79$) and moisture ($\rho = 0.87$), with the exception of Mg, Ca and Mn that were

negatively correlated with component 2.

We also determined the concentration of culturable total aerobic bacteria, coliforms and

enterococci bacteria present in PL microcosms (Supplementary Table 1). Coliform bacteria were

not recovered at any time point in our study, which agrees with studies showing low prevalence

of Gram-negative bacteria in PL (49). SH-2813 microcosms had higher aerobic bacteria (log 7.6

± 0.37 CFU g^{-1}) than SH-116 (log 6.2 ± 0.07 CFU g^{-1}).

Whole genome sequencing. All isolates sequenced had an average coverage of 101 times with a

range of 28 to 234 times Assembly size of sequenced genomes including extrachromosomal

DNA ranged from $4.75 - 4.80$ MB. Differences in genome size may be due to the presence or

absence of mobile genetic elements including phages and plasmids. The average N50 for all

isolates was 370, 097, which represents the average contig size after assembly with SPaDes

assembler. Quality assessment and coverage report for all assembled genomes are available in

Supplementary Files 1 and 2.

Phylogenetic trees and chromosomal mutations. To investigate the genetics of *S*. Heidelberg

persistence in PL, we isolated and performed WGS on 86 isolates obtained after 0, 1, 7 and 14

days in PL (Supplementary Table 3). First, we determined the genetic relatedness of the isolates

recovered by reconstructing a maximum-likelihood tree (Fig.1c) with 213 SNPs/InDels

identified in ancestral and evolved strains of SH-2813 and SH-116. We designated *S. Heidelberg*

(CP013673) as the reference genome because it differed from the two strains by a low number of

SNPs (<110) (Table 1). In addition, it carried similar IncX1 and ColpvC plasmids. After removal

of identical SNPs/InDels, 28 isolates were not included in the reconstructed tree. All SH-2813

isolates recovered after 1 day of incubation in PL were represented by 4 clades highlighted in

green (Fig. 1a). However, SH-2813 day 1 isolates recovered from PL that was previously

enriched for 24 h in BPW (SH-2813BPW-Day1) were clustered with isolates recovered after day

7 and with all the SH-116 isolates (clades 5 -18). These isolates in clades 1-4 differed by 2 – 5

SNPs/ InDels from the ancestral SH-2813. A common mutation found between these isolates (9

of 12) was in the 23S-5S Internal Transcribed Spacer (ITS) region between positions 3503197 –

3503202 (Supplementary Table 4). Isolates not carrying this mutation (SH-5B, SH-13B, and SH-

14B) had a mutation in the 16S rRNA region (position 4125232: A -> C). Mutation was also

acquired in the 23S rRNA region (position 4129873; A -> G) for 6 isolates.

All SH-2813 isolates recovered after 7 days in PL (SH-2813-Day7) and SH-2813BPW-Day1

isolates (n =6) acquired a total of 76 new mutations that were absent in day 0 and 1 isolates. For

SH-116, the number of SNP difference between two evolved isolates or the ancestral strain was

not more than 3 SNPs at any time point. A total of 62 mutations were identified between SH-

116, SH-2813-Day7 and SH-2813BPW-Day1 isolates (Fig. 1b and c). Of these mutations, 37

were SNPs/InDels present in the same loci on the chromosome and resulted in the same

nucleotide change (Table 2).

Table 2. Mutations shared between evolved *S. Heidelberg* isolates.[a]

Loci on chromosome	Location [b]	Product	Mutation change [c]
70249	intragenic	Carbamoyl phosphate synthase small subunit	Asp--> Asp
241459	intragenic	Fimbrial protein (StfE)	Val--> Arg
295492	Intergenic	Upstream 16S ribosomal rRNA	N/A
629229	intragenic	Allantoin permease	Trp--> Stop
774273	intragenic	Amino acid ABC transporter permease	Ala --> Trp
775920	Intergenic	Upstream glutamate/aspartate ABC transporter	N/A
823726	Intergenic	Upstream peptide permease	N/A
830662	intragenic	UDP-galactopyranose mutase	Phe --> Leu
832621	intragenic	Galactosyl transferase (Pseudo)	ND
832993	intragenic	Galactosyl transferase (Pseudo)	ND
1035518	intragenic	Cysteine/glutathione ABC transporter permease (CydD)	Val --> Pro
1103788	Intergenic	Upstream FtsZ	N/A
1210309	intragenic	Proline dehydrogenase (PutA)	Arg --> Glu
1288160	intragenic	Ferric-rhodotorulic acid/ferric-coprogen receptor (FhuE)	Ala --> Arg
1423283	intragenic	Cyclic di-GMP regulator (CdgR)	Gln --> Stop
1494796	intragenic	Type III secretion system export apparatus protein (SsaV)	His --> Asn
1628331	intragenic	DUF1493	ND
1649579	intragenic	NAD-dependent malic enzyme	Thr --> Thr
1689446	intragenic	1-pyrroline dehydrogenase	Arg --> Thr
2069216	Intergenic	Upstream hypothetical protein	N/A
2115173	Intergenic	Upstream cobyrinic acid a,c-diamide synthase	N/A
2333347	intragenic	Acetylase	ND
2517571	Intergenic	Upstream outer membrane protease (OmpT)	N/A
2757879	IntraRNA	5S ribosomal rRNA	N/A
2880324	intragenic	Ribonucleotide-diphosphate reductase subunit alpha (NrdA)	Asp --> Ala
3212823	intragenic	tRNA (guanosine(46)-N7)-methyltransferase (TrmB)	Arg --> Ala
3324819	intragenic	Transcriptional regulator (PadR)	Leu --> Gln
3336383	intergenic	Upstream hypothetical protein	N/A
3357216	intragenic	Transcriptional regulator (TdcA)	Phe --> Glu
3523467	Intragenic	50S ribosomal protein L15	Ala --> Arg
3594367	intragenic	Two-component system sensor histidine	Val --> Asn

		kinase (EnvZ)	
3629811	**intragenic**	**2-ketogluconate transporter**	**Ser --> Ser**
3791102	**intragenic**	**3-dehydro-L-gulonate 2-dehydrogenase**	**Gly --> Ser**
4265903	**intragenic**	**TonB-dependent vitamin B12 receptor (BtuB)**	**Ala --> Ser**
4303417	**intragenic**	**Hypothetical protein**	**Asp --> Ala**
4607527	**intragenic**	**Transposase (IS200)**	**Glu --> Val**
4652337	**intergenic**	**Upstream hypothetical protein**	**N/A**
295539	Intergenic	Upstream rRNA-16S ribosomal RNA	N/A
295548	Intergenic	Upstream rRNA-16S ribosomal RNA	N/A
295551	Intergenic	Upstream rRNA-16S ribosomal RNA	N/A
295555	Intergenic	Upstream rRNA-16S ribosomal RNA	N/A
295561	Intergenic	Upstream rRNA-16S ribosomal RNA	N/A
295564	Intergenic	Upstream rRNA-16S ribosomal RNA	N/A
926872	Intragenic	hypothetical protein[#]	Leu --> Asp
1367477	Intragenic	aldo/keto reductase[^]	Thre --> Ile
1420609	Intragenic	vitamin B12 ABC transporter permease (BtuC)	Val --> Gly
2018078	Intergenic	Upstream methyl-accepting chemotaxis protein II[#]	N/A
2049907	Intragenic	flagellin (FliC) [^]	Leu --> Ala
2115448	Intragenic	hypothetical protein[#]	Leu --> Tyr
2163440	Intragenic	Phosphomannomutase[#]	Gly --> Thr
2307329	Intragenic	bifunctional PTS fructose transporter subunit IIA/HPr[#]	Ala --> Val
2978093	Intragenic	DNA mismatch repair protein (MutS) [#]	Met--> Asn
3503198	ITS	23S -5S rRNA ITS[^]	N/A
3503200	ITS	23S -5S rRNA ITS[#]	N/A
3503201	ITS	23S -5S rRNA ITS*	N/A
3503201	ITS	23S -5S rRNA ITS*	N/A
3503201	ITS	23S -5S rRNA ITS*	N/A
3503213	ITS	23S -5S rRNA ITS[^]	N/A
3503219	ITS	23S -5S rRNA ITS[^]	N/A
3532866	Intergenic	Upstream 30S ribosomal protein S10 and leader peptidase hopD[#]	N/A
3947635	Intragenic	Cytochrome c biogenesis protein (CcmH)[^]	ND
4035545	ITS	23S -5S rRNA ITS	N/A
4035546	ITS	23S -5S rRNA ITS	N/A
4035547	ITS	23S -5S rRNA ITS[^#]	N/A
4035547	ITS	23S -5S rRNA ITS[^#]	N/A

4035548	ITS	23S -5S rRNA ITS^	N/A
4035549	ITS	23S -5S rRNA ITS	N/A
4035551	ITS	23S -5S rRNA ITS	N/A
4035556	ITS	23S -5S rRNA ITS	N/A
4035560	ITS	23S -5S rRNA ITS*	N/A
4035560	ITS	23S -5S rRNA ITS*	N/A
4035561	ITS	23S -5S rRNA ITS	N/A
4035562	ITS	23S -5S rRNA ITS*	N/A
4035562	ITS	23S -5S rRNA ITS*	N/A
4035566	ITS	23S -5S rRNA ITS	N/A
4035567	ITS	23S -5S rRNA ITS	N/A
4035570	ITS	23S -5S rRNA ITS^	N/A
4035573	ITS	23S -5S rRNA ITS	N/A
4035575	ITS	23S -5S rRNA ITS^	N/A
4035576	ITS	23S -5S rRNA ITS*	N/A
4035576	ITS	23S -5S rRNA ITS*	N/A
4271745	IntraRNA	rRNA-23S ribosomal RNA#	N/A
4271746	IntraRNA	rRNA-23S ribosomal RNA#	N/A
4271748	IntraRNA	rRNA-23S ribosomal RNA#	N/A
4271751	IntraRNA	rRNA-23S ribosomal RNA#	N/A
4289060	Intragenic	DNA-directed RNA polymerase subunit beta (RpoC) #	Ile --> Asn
4366478	Intragenic	Maltose ABC transporter substrate-binding protein (MalE) #	Gly --> Ser
4482602	Intergenic	Upstream acid phosphatase^	N/A

Note.

[a] Mutations are present in Col-plasmid carrying isolates and absent in isolates without Col plasmids. SNPs/InDels

were compared to reference genome CP016573.

[b] Intragenic- present within a coding sequence, Intergenic – non coding region between two genes, ITS – internal

transcribed spacer, IntraRNA – present within non-coding RNA.

[C] N/A – not available (SNP/InDel present in non-coding sequence), ND- not determined.

Boldness denotes hotspot mutations present in all Col-plasmid carrying isolates (n=37).

Single nucleotide variants present in <1 % of SH-2813 isolates (n=14).

^ Single nucleotide variants present in <1 % of SH-116 isolates (n=12).

* "Heterozygous" SNPs/InDels present in multicopy gene.
438
439

These mutations were in proteins or intergenic regions associated with diverse processes such as

carbomoyl phosphate synthesis, motility (StfE), galactosyltransferease (a pseudogene), two-

component system sensor histidine kinase (EnvZ), cyclic -di-GMP synthetase (c-di-GMP), SPI-2

type III secretion system apparatus protein (SsaV) , siderophore transport (BtuB and FhuE),

cysteine/glutathione transporter (CydD), 50S ribosomal protein (L15) and a protein of unknown

function (DUF1493) etc.

Twenty-five of the 62 mutations were present in ~ 70% of SH-2813 and SH-116 isolates

(Table 2). The majority of these mutations were multiple SNPs/InDels present in the 23S-5S ITS

region between positions 4033545 – 4035576 on the chromosome. Further, 14 single nucleotide

variants (SNVs) were present only in a handful of SH-2813 isolates and absent in all SH-116

isolates. For instance, strain SH-2813-17BBPW carried 5 unique SNVs including a non-

synonymous mutation (met1asn) in the DNA mismatch protein (MutS) making it a spontaneous

mutant (Table 2; Supplementary Table 4). For SH-116, 11 unique SNVs were identified in 23S-

5S ITS region (n =8) and in proteins encoding aldo/keto reductase, phase 1 flagellin (FliC) and

cytochrome c biogenesis protein (CcmH).

IncX1 and Col plasmids are associated with *S*. Heidelberg persistence in PL. We next

determined if *S*. Heidelberg isolates recovered from PL acquired new plasmids. All SH-2813$_{evol}$

isolates (n= 27) obtained from day 7 and 14 microcosms and SH-2813BPW-Day1 isolates

carried 3 Col plasmids in addition to an IncX1 plasmid. For SH-116, IncX1 and Col plasmids

were present in all sequenced strains (n = 38) regardless of time of incubation in PL. The

majority (276/306) of the plasmid contigs represented complete circular plasmids

(Supplementary Table 5). The IncX1 belonged to the MOB$_Q$ family of relaxases and was present

in all sequenced genomes. The ColE1 plasmids belonged to the MOBp family and showed a

conserved size of 6189 (ColE1-6kb) and 4784 (ColE1-4kb) bp respectively. The cryptic ColpVC

was 2223 bp (ColpVC-2kb). For all complete IncX1 plasmids, there was a 21-base pair (bp)

difference in strains carrying the Col plasmids (37, 824 bp) compared to strains with no Col

plasmids (37, 845 bp).

Direct-repeat deletion is associated with IncX1 copy number increase. To determine the

effect of the 22-bp deletion identified in the IncX1 plasmids present in *S*. Heidelberg populations

carrying Col plasmids, all complete circular IncX1 plasmid sequences were subjected to whole

genome sequence alignment and SNV identification. The 22-bp direct-repeat deletion was

located at the origin of replication (*ori*) adjacent to the replication protein (RepB) and an

additional 1-bp (G-> GA) insertion occurred in the type IV secretion protein (T4SS) VirB8 (Fig.

2a and Supplementary Table 5).

Figure 2. Deletion of 22-bp iteron of IncX1 is associated with a reduction in *rep* expression.

(**a**) Annotated IncX1 plasmid map of SH-2813$_{anc}$ and SH-116$_{anc.}$ Regions that acquired

mutations in evolved strains carrying Col plasmids are highlighted in rectangular dashed boxes

(**b**) Multiple alignment of the IncX1 iteron-based direct repeat (DR) origin of selected strains

from this study, and 22 *S*. Heidelberg genomes available on NCBI. (**c**) Differences in expression

of IncX1 genes in evolved SH-2813 and SH-116 relative to SH-2813$_{anc}$ and SH-116$_{anc}$ (log$_2$

scale, N =3, *p.value =0.086 (Wilcoxon signed-rank Test). The error bars represent the standard

error of the mean.

The 22-bp deletion reduced the number of direct repeats from 3 to 2, making it a truncated *ori*

(Fig. 2b). Direct repeat iterons of theta plasmids are known to bind Rep proteins and have been

shown to participate in negative control of their initiation (50). Deleting or adding more copies of

the iteron have been shown to increase or decrease plasmid copy number (PCN) (51). Therefore,

we hypothesized that the direct repeat deletion in IncX1 of SH-2813$_{evol}$ would result in higher

copies compared to SH-2813$_{anc}$ since the 22-bp is present in the ancestral strain while no change

would be expected in IncX1 copies for SH-116$_{evol}$. To test this hypothesis, we determined the

copy numbers of IncX1 and Col plasmids present in ancestor (SH-2813$_{anc}$ and SH-116$_{anc}$) and 2

evolved strains (SH-2813$_{evol}$ and SH-116$_{evol}$ respectively). We assumed that these evolved strains

represent the "fittest" SH-2813 and SH-116 strains given their persistence in PL microcosms for

14 days.

As expected, the copy number of IncX1 for SH-2813$_{evol}$ (~7 copies/cell; s.d.=1.8) was

higher than that of SH-2813$_{anc}$ (~4 copies/cell; s.d.=0.3) at 0.5 h (p. value = 0.086; Wilcoxon

rank-signed test). In contrast, SH-116 did not exhibit any significant difference in IncX1 copies

between SH-116$_{anc}$ (~5 copies/cell; s.d.=1.7) and SH-116$_{evol}$ (~6 copies/cell; s.d.=0.8) (p. value =

0.773) (Table 3).

Table 3. Plasmid copy numbers of ancestral and evolved strains.

Strain	Time (h)	Optical Density (S.D)	Bacteria concentration (Log CFU/ml)	Plasmid Copy Number/Cell (S.D)			
				IncX1	ColE1-6kb	ColE1-4kb	ColpVC
SH-2813$_{anc}$	0.5	0.015±0.00	7.98±0.0	4.17±0.3	N/A	N/A	N/A
	4	0.040±0.00	8.60±0.2	0.47±0.5	N/A	N/A	N/A
	24	0.070±0.00	8.90±0.1	0.97±0.9	N/A	N/A	N/A
SH-2813$_{evol}$	0.5	0.016±0.00	7.94±0.0	6.92±1.8	10.71±4	12.4±5.5	110.77±45
	4	0.056±0.00	11.16±2.1	0.59±0.3	3.28±0.6	2.77±1.0	86.21±25
	24	0.080±0.00	8.57±0.0	0.64±0.1	5.25±0.7	4.2±0.6	115.32±70
SH-116$_{anc}$	0.5	0.014±0.00	7.83±0.1	4.55±1.7	0.27±0.3	2.54±1.9	82.39±8

	4	0.037±0.00	8.56±0.1	0.80±1.3	0.14±0.0	0.54±0.7	54.01±26
	24	0.093±0.00	9.07±0.0	2.33±9.3	2.23±0.0	0.92±1.3	39.94±28
SH-116evol	0.5	0.017±0.00	7.91±0.0	5.61±0.8	9.44±3.5	10±4.4	90.47±45
	4	0.065±0.00	8.73±0.2	0.60±0.8	3.83±1.7	2.9±1.3	93.83±11
	24	0.069±0.00	8.69±0.1	0.35±0.4	4.39±1.0	3.5±0.3	72.56±20

Note. –

SH-2813$_{anc}$ - Original SH-2813 strain before inoculation into PL, SH-2813$_{evol}$ - SH-2813-12A-day14-PLE, SH-

116$_{anc}$ – Original SH-116 strain before inoculation into PL, SH-116$_{evol}$ = SH-116-36B0.5-day14-PLE.

N/A= not applicable

Gene expression of IncX1 replication protein and stabilization genes. We monitored the

expression of the *repB* replication gene and *relBE* genes which encode a type II toxin-antitoxin

(TA) system associated with the IncX1 plasmid. The *relBE* genes were selected since they are

adjacent to the *ori* (Fig. 2a) suggesting they may share the same promoter as *repB*; and due to

their implicated role in plasmid stabilization (52, 53). Gene expression results in SH-2813$_{evol}$

revealed a marked 4.5-fold reduction in *repB* expression and 2.2-fold reduction in *relB*

expression at 4 h relative to SH-2813$_{anc}$; while the expression levels of toxin *relE* remained

unchanged in both ancestral and evolved SH-2813 (Fig. 2c).

SH-116 strains lacks the extra 22-bp direct repeat present on IncX1, thus we expected no

change in Rep expression. However, there was a change in expression for these transcripts for

SH-116$_{evol}$ — reductions in *repB* and *relB* and no change in *relE* when compared to SH-116$_{anc}$

(Fig. 2c). We speculate that the reduction in expression of Rep for SH-116 evolved strains was a

result of carrying higher copy ColE1 plasmids than SH-116$_{anc}$. In fact, SH-116$_{evol}$ carried ColE1

plasmids with higher copies (ColE1-6kb =9±3.5, ColE1-4kb = 10±4.4, ColpVC-2kb = 90±45)

than SH-116$_{anc}$ (ColE1-6kb =0.3±0.3 ColE1-4kb = 3±1.9, ColpVC-2kb = 82±7.9) (Table 2). The

presence of this high copy ColE1 plasmids correlated with faster growth rates in MHB. The

optical density of SH-2813$_{evol}$ and SH-116$_{evol}$ was 1.4 and 1.7-fold higher than that of SH-2813$_{anc}$

and SH-116$_{anc}$, respectively, at 4 h (Table 3). This difference in copy number was further

corroborated by comparing PCN determined from WGS assembly coverage, showing that all

SH-116$_{evol}$ isolates from day 1, 7 and 14 on average had higher ColE1 plasmid copies than SH-

116$_{anc}$ (Supplementary Table 6). Finally, a multiple regression analysis on IncX1 and Col PCN

showed that IncX1 was a statistically significant predictor of Col plasmid copies in a cell (F (3,

68) =13.82, p.value < 0.000, R^2 =0.39). SH-116 strains were predicted to have approximately six

ColE1-6kb plasmid copies for every IncX1, whereas, SH-2813 was predicted to have 3 copies

per IncX1 (Fig. 3).

Figure 3. Correlation between IncX1 and ColE1-6kb plasmid copies. Plasmid copy number

(PCN) was determined from whole genome sequencing de-novo assembly coverage (see Table

S6.

ColE1 plasmids differed in mobilization genes and non-coding RNA. To determine if there

are genetic differences between Col plasmids carried by SH-116 and SH-2813, we used

ProgressiveMauve (28) to perform multiple whole genome alignments on DNA sequences from

complete Col plasmids. Results showed that ColE1 plasmids present in sequenced strains were

highly similar and only differed in nucleotide regions coding for mob genes (Fig. 4;

Supplementary Fig. 3b and c).

Figure 4. Linear maps of Col plasmids and phylogeny based on conserved mobilization

genes. (a) Annotated maps of Col plasmids carried by ancestral and evolved isolates of *S*.

Heidelberg used for competition experiments. (**b and c**) Maximum-likelihood based tree of

aligned conserved protein sequences of (**b**) relaxase (MbeA) and (**c**) MbeC genes of Col

plasmid-carrying isolates. Plasmids with identical MbeA (n=106) and MbeC (n=112) sequences

were not included in the reconstructed tree. The numbers shown next to the branches represent

the percentage of replicate trees where associated taxa cluster together based on 1000 bootstrap

replicates. Taxa in parenthesis represent selected isolates used for competition experiments.

Trees were rooted with MbeA and MbeC sequences of *E. coli* (NCBI accession number:

NC_001371).

We identified 5-6 open reading frames (ORF) encoding 2-3 mobilization (mob) genes, a putative

type IV TA system (Abi), a putative aminoglycoside 6-N-acetyltransferase (aac(6')-Ib) and a

hypothetical protein for ColE1-6kb (Fig. 4). The ColE1-4kb plasmids had 4-6 ORFs encoding a

protein named macrophage stimulating factor (MSF), a hypothetical protein and 1-4 mob genes.

Both ColE1 plasmids harbored a non-coding RNAI and RNAII, while ColE1-6kb carried an

additional non-coding repressor of primer (Rop) protein known to regulate replication and ColE1

plasmid copy numbers. The ColpVC-2kb plasmids had 2-3 ORFs, encoding 1-2 replication

proteins (Rep) and a hypothetical protein (Fig. 4). The RNAI present in the ColE1 plasmids had

approximately 62 % DNA sequence homology (data not shown).

These mob CDS were identified as MbeA and MbeC and were present in different sizes

and arrangement. In some cases, the *mbeA* gene overlapped ORF annotated as *mbeB and mbeD*

(Fig. 4), suggesting MbeB and MbeD was part of the mob region in some but not all ColE1

plasmids. To determine if there are differences in these proteins between Col plasmids, we

performed a multiple alignment of conserved MbeA relaxase and MbeC proteins. We observed

high conservation in the N-terminal region and high variation in their C-terminal (data not

shown). According to the phylogenetic tree reconstructed for these conserved amino acids,

MbeA proteins belonging to ColE1-4kb plasmids clustered in a unique clade, whereas the

ColE1-6kb was placed closer to the signature *E. coli* ColE1 plasmid (NC_001371) (Fig. 4b).

After the removal of identical sequences, only MbeA sequences belonging to 3 SH-2813 isolates

were included in the final tree The MbeA tree constructed for ColE1-6kb and ColE1-4kb

plasmids were represented by 1 and 2 isolates, respectively. On the other hand, the two clades

observed for MbeC tree had 9 and 5 isolates for ColE1-6kb and ColE1-4kb plasmids (Fig. 4c).

Col plasmids did not impose a fitness cost on *S*. Heidelberg. Although acquisition of Col

plasmids could result in faster growth in host bacterium it may also translate to a fitness cost if

their regulation is not equally constrained. To establish if acquisition of Col plasmids imposed a

fitness cost on *S*. Heidelberg, we competed 2 evolved (SH-2813$_{evol}$ and SH-116$_{evol}$) and 1

ancestral (SH- 116$_{anc}$) strain carrying three Col plasmids against a Col plasmid-free isogenic SH-

2813$_{anc}$ harboring a *gyrA* mutation for nalidixic acid resistance (SH-2813$_{nal}$). The fitness of each

strain was determined relative to SH-2813$_{nal}$ as described by Millan et al (42). Results show that

Col plasmid carrying strains had a higher relative fitness of 1.27 (s.d. =0.12, n =3) compared to

SH-2813$_{nal}$, and differences were observed between strains (Fig. 5a).

Figure 5. Fitness and Conjugation rates of evolved *S*. Heidelberg isolates (a) Fitness of Col

plasmid-carrying strains of SH-2813 and SH-116 relative to a Col plasmid-free ancestor of SH-

2813$_{nal}$. Bars represent means of four technical replicates for each population. The horizontal

dashed red line represents the fitness of SH-2813$_{nal}$. **(b)** Conjugation rates of Col plasmids from

Col plasmid-carrying strains to SH-2813$_{nal}$ ancestor. Conjugation rates were estimated using data

from fitness experiment. Bars represent means of four technical replicates for each population.

The transfer rate for ColE1-4kb plasmid from SH-116$_{evol}$ to SH-2813$_{nal}$ was not determined.

Symbols: Abortive phage infection protein of ColE1-6kb (Abi-ColE1-6kb), macrophage

stimulating factor protein of ColE1-4kb (MSF-ColE1-4kb) and replication protein of ColpVC

(rep-ColpVC).

The transfer rate for ColE1-4kb plasmid from SH-116$_{evol}$ to SH-2813$_{nal}$ was not determined

because none of the transconjugants screened by qPCR was positive for the MSF gene harbored

on this plasmid. Consequently, the differences in fitness between strains may be attributed to a

higher conjugation rate observed for ColE1-6kb and ColpVC for SH-116$_{evol}$ (Fig. 5b).

Antimicrobial resistance associated with the acquisition of Col plasmids. The ColE1-6kb

plasmid in the current study carries a putative aac (6')-Ib gene that is 56 % (*e.value* = 1<e-13)

identical to the aac (6')-Ib gene present in *Klebsiella pneumoniae* Sul1-type integron (GenBank

accession number: AAR18814.1) and 34 % (*e..value* = 6<e-28) identical to the one present in

Thiomona sp. X19 (GenBank accession number SCC93141.1); suggesting it may play a role in

aminoglycoside resistance (Fig. 6a and b).

Figure 6. Acquisition of ColE1 plasmids was associated with decreased susceptibility to

aminoglycosides and fosfomycin. (a) Multiple alignment of conserved N-terminal residues for

aminoglycoside acetyltransferase (AAC-6') genes encoded in the *S*. Heidelberg (SH-2813)

chromosome and ColE1-6kb plasmid from this study, and AAC-6' genes with known resistance

phenotype. (b) Corresponding neighbor-joining tree of aligned AAC-6' protein sequences. (c)

Inhibition zone diameter (mm) distributions for evolved strains and selected antimicrobial drugs.

SH-2813 (n= 27) and SH-116 (n= 18) isolates tested were recovered after 0 and 14 days of

microevolution in PL. A decrease in inhibition zone corresponds to a decrease in susceptibility to

selected antibiotic.

Further, *S*. Heidelberg ancestral strains each carried chromosomally encoded aminoglycoside

(aac (6')-Iy) and fosfomycin (fosA7) genes (54) that have been demonstrated to confer resistance

to tobramycin, netilmicin, neomycin, amikacin, kanamycin and fosfomycin (55-57). To assess

the correlation between AMR and the acquisition of Col plasmids, we performed antibiotic

susceptibility testing (AST) on evolved *S*. Heidelberg strains (n=42) on the NARMs

antimicrobials (44). In addition, we used disc-diffusion assay (45) to evaluate susceptibility to

tobramycin, netilmicin, gentamicin, neomycin, kanamycin and fosfomycin.

The majority of the isolates tested (97.6 %) were susceptible to the 14 drugs on NARMs

AST panel (Supplementary Table 7). However, 12 of 16 (75 %) SH-2813$_{evol}$ isolates recovered

from PL on day 7 and day 14 displayed an increase in MIC for gentamicin. The average zone of

inhibition (mm) for kanamycin, tobramycin and fosfomycin for SH-2813$_{anc}$ strains was larger

than the zones measured for SH-2813$_{evol}$ strains (Wilcoxon signed-rank test; *p*-value < 0.0001)

(Fig. 6c). In contrast, SH-116 showed no significant difference in zone of inhibition between SH-

116$_{anc}$ and SH-116$_{evol}$ strains for any of the antibiotics tested (Wilcoxon signed-rank test; *p*-value

> 0.01).

Col plasmid-encoded genes are prevalent in PL. To extrapolate beyond our microcosm

evolution experiment, bioinformatics and quantitative PCR were used to assess the presence of

these Col plasmids across sequenced *S*. Heidelberg genomes and in various types of PL. DNA

sequences for complete Col plasmids from this study were subjected to BLAST searches against

the non-reduntant (nr) GenBank database. No BLAST hit matched the complete sequence of

ColE1-6kb plasmids, but the majority of hits covered approximately 55 % of the plasmid

genome. In most cases, these hits corresponded to mobilization genes. For ColE1-4kb, the

closest hit was 99 % identical to the colicigenic plasmid of *Salmonella enterica* serovar Berta

(pBERT) and covered the entire genome. All other plasmid hits covered <82 % of the ColE1-4kb

genome and were represented in various species of Gram-negative bacteria. The ColpVC-2kb

was >99 % identical to plasmids from 5 *Salmonella* sp, including three *S.* Heidelberg strains

isolated from Chicken in Canada (data not shown). These results suggested that complete

sequences of Col plasmids of PL origin are not well represented in GenBank.

To address this possibility, we performed a BLAST search against the NCBI whole genome

shotgun contigs database restricting our search to only serovar *Salmonella*. This database

contains genome assemblies of incomplete genomes or incomplete chromosomes of bacteria that

were sequenced by the shotgun strategy. *Salmonella* Heidelberg and *S.* Typhimurium strains

were the major serovars with contigs matching >99 % of the ColE1-6kb plasmids. For instance,

the top 5 contigs were 100 % identical and were 6183 bp long (NCBI Bioproject

PRJNA417775). These contigs were detected in *S.* Heidelberg strains collected from broiler

breeder farms in the USA and the isolation source was environmental booties. The ColE1-4kb

and ColpVC-2kb plasmid were detected in contigs from sequenced genomes of multiple serovars

including Heidelberg, Rissen, Hadar, Typhimurium, Albert, Kentucky, Infantis, Enteritidis etc.

The percent identity and query cover for ColE1-4kb and ColpVC-2kb across the serovars ranged

from 97 -100 % and 64 -100 %, respectively. Surprisingly, majority of the *S.* Heidelberg

recovered from the broiler breeder farms carried multiple Col plasmids (data not shown). For

example, strain C_NS-007 (Accession: SAMN08031253) harbored 3 Col plasmids of sizes ~6

kb, 4 kb and 2 kb encoding the same proteins as observed in this study (Supplementary Fig. 4).

Furthermore, we determined the concentration of Col plasmids in PL collected from

conventional and backyard poultry farms located in 4 states in USA. The Col plasmid genes were

detected in all litter types except 1 backyard litter, whereas the Rep protein for IncX1 was

detected in only 1 of 7 of the litter types, suggesting that Col plasmid genes are abundant in PL

(Fig. 7a).

Figure 7. Col plasmids present in poultry litter (PL). (a) Concentration of plasmid-specific

genes and 16S rDNA in poultry litter of different origins. PL was collected as grab samples from

multiple locations of various poultry houses/farms in USA and composited in 1-gallon Whirl-Pak

bags. qPCR was performed on DNA extracted from 250 mg of PL in triplicates and pooled. Error

bars represent the standard error of the means. Symbols: Conventional farming (Con), Backyard

farming (Back), replication protein of IncX1 (IncX1-rep), replication protein of ColpVC

(ColpVC-rep), abortive phage infection protein of ColE1-6kb (ColE1-6kb-Abi), macrophage

stimulating factor protein of ColE1-4kb (ColE1-4kb-MSF) and 16S ribosomal DNA of

Enterobacteriaceae (16S rDNA). **(b)** Concentration of plasmid-specific genes in PL used for

microevolution experiment. DNA was extracted from 48 microcosms and qPCR was performed

using primers targeting the incRNAI region of ColE1-6kb and ColE1-4kb, the replication protein

of ColpVC and the 16S ribosomal DNA of Enterobacteriaceae. Note. One uninoculated

microcosm was included at day 0. ColE1 was not detected in SH-2813-BHI and SH-116-BHI

microcosms. *p.value =0.086 (Wilcoxon signed-rank Test). The error bars represent the standard

error of the mean.

Poultry litter as a selection environment for prophages. Although plasmids are the major

MGE elements that have been shown to be rapidly transferred between bacterial populations due

to the high number of plasmid copies found per cell — the loss or uptake of prophages cannot be

overlooked. Prophages can propagate horizontally between cells (lytic phages) or vertically if

they are lysogenized, typically by integrating into bacteria chromosome as prophages. Prophages

carry several virulence and accessory genes that can increase their fitness under certain

environmental conditions, such as nutrient limitation, oxidative stress, biofilm formation and

antibiotic resistance (58). However, intact prophages are like molecular time bombs that kill their

hosts upon activation of lytic cycle (59, 60). Consequently, one should expect prophage

inactivation to be under strong selection in the environment (59).

To determine if prophage selection played any role during the our microevolution

experiments, we used PHAST (24) for predictions of prophages in sequenced genomes. The

ancestral *S*. Heidelberg strains each carried 3 intact (Gifsy-2, PaV-LD, sal-4) and 2 defective

prophages (Gifsy-1, BcepMu). Only phages found intact by PHAST were further analyzed in this

study. Gifsy-2 (NC_010393) prophage harbored a Cu-Zn superoxide dismutase (SOD) and an

attachment invasion locus (OmpX) gene with known oxidative stress (61, 62) and virulence (63)

functions respectively. PaV-LD (NC_016564) is a planktonic phage carrying Salmochelins

(siderophores of *Salmonella* enterica and uropathogenic *Escherichia coli* strains) transcribed

from *iroBCDEN* genes and sal-4 (NC_030919) carried accessory gene for polymyxin resistance

(*arnC*). We observed that the sal-4 prophage shared the same proteins as enteroP22 and sal-5,

and the number of query hits matching them was similar (data not shown). For convenience, we

have named all prophages predicted as P22 or sal-5, sal-4 in this study. The prophage Gifsy-2,

PaV-LD and sal-4 were found intact in 84, 19, and 45 of the 85 sequenced evolved strains,

respectively. Strain SH-32A-116-Day7-PLE had no intact phage (Supplementary Fig. 5).

DISCUSSION

Positive selection for hotspot mutations during microevolution in PL. Repeated,

phylogenetically unlinked nucleotide changes that results in mutation of the same amino acid

positions are called hotspot mutations (64-66). Two common types of hotspot mutations have

been described: (1) parallel mutations that lead to the same amino acid change in the same

position and (ii) coincidental mutations that lead to a different amino acid change in the same

position (64). These forms of convergent evolution are strong indicators of positive selection and

not expected to occur by chance alone (65). We identified 37 parallel mutations carried by all

SH-116 isolates recovered and in SH-2813-Day7 and SH-2813BPW-Day1 isolates. A significant

proportion of these mutations were point mutations identified in proteins with multiple functions.

For instance, a non-synonymous mutation (gln123stop) was present in the cyclic di- GMP

regulator (CdgR) — a universal bacterial second messenger that regulates a variety of processes,

including cell motility, intercellular interactions, biofilm formation and dispersal and responses

to diverse environmental changes (67). The amino acid change resulted in a premature stop

codon in position 123. A potential effect of such mutation is a truncated, incomplete or

nonfunctional CdgR. High levels of *cdgR* have been demonstrated to promote biofilm formation

and sessility, while low levels promote motility and the synthesis of virulence factors in human

and animal pathogens (68). A mutation (val196Asn) was also identified in a two-component

system sensor histidine kinase (EnvZ) that is involved in the regulation of osmoregulation and

controls the expression of outer membrane porin proteins OmpF and OmpC (69). Missense

mutation in *envZ* gene has been shown to reduce the expression of porin genes, thus reducing the

permeability of the outer membrane of *Salmonella* to cephalosporin antibiotics (70).

Non-synonymous mutations identified in ferric-coprogen receptor (FhuE) and tonB-

dependent vitamin B12 (BtuB) suggest that these mutations are related. FhuE serves as a

receptor for ferric-coprogen and ferric-rhodotorulic acid and uptake of ferric-coprogen is BtuB

dependent. Together, they are required for the uptake and utilization of cobalamins and

siderophores (71, 72). These two genes are controlled by Fur regulator and a RNA riboswitch,

and their expression is repressed when iron and cyanocobolamin reaches a certain level (71). The

effect of this mutation on their interaction or expression in *S*. Heidelberg is beyond the scope of

this study. Multiple SNPs/InDels was identified in the 23S-5S ITS region of isolates with hotspot

mutations. A large fraction of these mutations were insertions ranging in size from 2 − 70 bp and

were found in multiple loci within the ITS region (4035545 − 4035576). The high number and

frequency of these mutations may reflect a recent emergence of new lineages and/or their relative

evolutionary instability (64). The 23S-5S rRNA ITS region of *Escherichia coli* (*E. coli*) have

been shown to be the most hypervariable region found throughout the ribosomal RNA (*rrn*)

operon, and has also been reported for *Salmonella* Typhi (73). To assess this variability across *S*.

Heidelberg genomes, we reconstructed a phylogenetic tree with the 23S-5S ITS DNA sequences

from 22 complete genomes of *S*. Heidelberg available in GenBank (Supplementary Fig. 6). In

addition, we included the ITS region for 14 isolates from this study. The GenBank strains

clustered with either day 0 or day 14 isolates from this study. One explanation could be that the

strains clustering with day 14 isolates have been co-selected by a recent common environmental

pressure such as PL, whereas the strains clustering with day 0 isolates have not undergone a

common selection pressure. However, it is important to mention that the GenBank strains (19 of

22 strains) were isolated in Canada and may have biased the result of the reconstructed tree.

Another possible explanation is that this region of the *rrn* operon may be prone to indel

accumulation and is relatively unstable. For example, mutation was not identified in this region

in some of the isolates having hotspot mutations for both SH-2813 (6 isolates) and SH-116 (6

isolates).

High frequency hotspot mutations have been identified in the core genomes of *E. coli* and

Salmonella serovar *enterica* (64, 66, 74). Furthermore, hotspot mutations are more likely to be of

recent origin than long term (65). Chattopadhyay et al (64) performed an in-depth analysis on

hotspot mutations in multiple *E. coli*, *Shigella* and *Salmonella* sequence types and concluded that

the number of recent hotspot mutations were significantly higher than long term ones. In

addition, the frequency of recent hotspot mutations increased by the addition of more genomes to

their analysis. Taken together, our result can be explained by the so-called "source sink"

dynamics of pathogen microevolution. The "source" can be defined as an evolutionary stable

habitat (e.g. chicken gut), while the "sink" as an unstable transient compartment of a habitat (e.g.

poultry litter) where *S*. Heidelberg is introduced occasionally. Under this scenario, hotspot

mutations that are adaptive for survival in PL are continuously selected for on a short term, but

may be selected out upon colonization or entry into the original stable habitat. This tradeoff may

result in different hotspot mutations in *S*. Heidelberg strains isolated from PL, chicken gut or

humans.

Col plasmids and IncX1 contributed to the fitness of *S*. Heidelberg. In this study, only *S*.

Heidelberg strains carrying 2 ColE1 and cryptic ColpVC plasmids survived more than 7 days in

used PL. The ColE1-6kb relaxases (MbeA) shared >77 % homology with the relaxase of ColE1-

4kb but was more closely related to the prototype ColE1 (>96 %) plasmid of *E. coli*. When the

ColE1 plasmids sequences from this study were compared to those available in NCBI, we found

that these plasmids were underrepresented in the nr database but abundant in the genome

assemblies of incompletely assembled *Salmonella* genomes. The observed low frequency of

complete ColE1 plasmids harbored by *Salmonella* in NCBI may be a consequence of the bias

towards sequencing strains carrying plasmids with known antibiotic resistance genes or

virulence. After screening multiple PL of diverse origin with qPCR, we detected Col plasmid

genes at high levels in 6 of 7 litter types. Our data further suggest that the PL environment may

impose selective pressure that promotes the acquisition of these Col plasmids.

The ColE1-6kb plasmids were present in higher copy numbers than ColE1-4kb and carried 2

putative genes encoding a type-IV TA system and an aminoglycoside-acetyltransferase. The TA

system was linked by STRINGDB (75) to the widespread abortive bacterial infection (Abi)

system belonging to the cluster of orthologous group COG1106 (domain of ATPases) and the

closest homologs were those present in Pseudomonadaceae and Lactobacillaceae. In some

bacteria, this protein is annotated as ATP-binding protein or abortive infection protein. Abi

systems induce phage resistance to prophages of type 936 in *Lactococcus* and enhance plasmid

maintenance/stabilization via bacteriostatic toxicity (76). Consequently, cells carrying plasmids

encoding the Abi gene will be immune to phage attack or replication, thereby providing

protection for their clonal population. Their role in *Salmonella* remains questionable; however,

their high expression under laboratory growth conditions in our study (Supplementary Fig. 7)

suggests that they are functional and may have contributed to the persistence of *S*. Heidelberg in

PL.

We do not know the potential functions of the accessory genes on ColE1-4kb, but their

compatibility with ColE1-6kb may stem from differences in their RNAI sequences (77) and their

mob region. The ColE1 plasmid family represents plasmids that are only transmissible in the

presence of an additional plasmid with conjugative functions and usually carry a mob region

encoding specific relaxase and accessory mob genes (*mbeB*, *mbeD mbeC*). The relaxase

component is crucial for both initiation and transfer of DNA and is highly conserved (78, 79). In

addition, it has been predicted that the C-terminal region of MbeA binds ssDNA, while the N-

terminal fragment contacts MbeC (80, 81). The observed differences in C and N- terminal of

MbeA and MbeC in this study suggests that such modifications may be critical for successful co-

existence of multiple ColE1 plasmids in one host cell. We used these conserved mob protein

sequences for phylogenetic tree reconstruction and assessed if there is evolutionary relatedness

between the ColE1 plasmids from this study. The majority of the isolates (11 of 14) included in

the MbeC tree were from SH-116, while the 3 isolates included in the MbeA tree were from SH-

2813 strain. These results infers that the ColE1 plasmids carried by both SH-2813$_{evol}$ and SH-116

isolates have identical relaxases but differ in MbeC proteins.

The Incompatibility factor X1 plasmid harbored by the *S*. Heidelberg strains used in our

study more than likely served a conjugative role. IncX1 is a low copy plasmid and carries

required mating pair formation system for horizontal gene transfer (HGT) including type IV

secretion systems (T4SS), coupling proteins and a relaxase (82, 83) (Fig. 2a). This plasmid was

present in all strains recovered which infers that they are stably maintained and critical to the

survival *S*. Heidelberg strains carrying them. Stability of conjugative plasmids has been

attributed to high conjugation rates and plasmid stabilization proteins (84, 85) . The InDel

present in VirB8 resulted in a frame-shift mutation in this protein and subsequent changes in

DNA regions coding for T4SS proteins. Indeed, this mutation resulted in the fusion of a smaller

VirB8 (4.4 kDa) encoding gene with a bigger VirB8 gene (23.4 kDa) (Fig. 2a). In addition,

differences were observed in sequences coding for the conjugal transfer protein (TraG), an

important protein for bacterial conjugation (86). Interestingly, one strain without TraG (SH-

33A0.5-116-Day7-PLE) was isolated but we do not know the conjugal implication of this

deletion (Supplementary Fig. 8)

Following the acquisition of Col plasmids, we observed a significant reduction in the

expression of replication protein (Rep) protein and a type II toxin-antitoxin (TA) gene. This

reduction in *rep* expression suggests that *rep* expression may be linked to the cost of taking up

new MGE. Reducing *rep* expression will subsequently decrease the copies of IncX1 available for

ColE1 plasmids mobilization — thus reducing the burden associated with maintaining multiple

plasmids. Also, regression analysis of IncX1 and Col copies showed that IncX1 was a good

predictor of ColE1-6kb copies per cell — evolved strains carrying high IncX1 copies also carried

high ColE1-6kb copies (Fig. 3; Supplementary Table 6). This result suggests there might be

equilibrium between the numbers of "helper plasmids" to mobilizable plasmids residing in a cell.

Bacteria (TA) modules are genetic elements composed of a toxin protein component that

disrupts bacterial growth by interfering with vital cell processes and an antitoxin that impairs the

functionality of the toxin until this inhibition is abolished in response to cellular signaling. Type

II TA system carried on plasmids are important players in post-segregational killing — plasmid-

less cells are killed by the toxin, thereby establishing the dominance of plasmid-bearing cells in a

population (52, 53, 87). At late logarithm and stationary phase growth in Mueller Hinton Broth,

the *relB* gene (antitoxin) of IncX1 was significantly downregulated in evolved strains compared

to ancestral strains, and no change was observed in *relE* (toxin) expression (Fig. 2c). This

continuous expression of *relE* will ensure that only IncX1 carrying cells are present in the

population. Therefore, disrupting the conjugative and stability genes of conjugative plasmids

may represent a critical point for limiting their persistence in the environment. For instance, a

combination treatment with linoleic and phenothiazine reduced the conjugation efficiency of six

conjugative plasmids by 3 -fold, including the IncX plasmid of RP4 (88).

Furthermore, the IncX1 present in Col plasmid-bearing strains of *S*. Heidelberg carried a 22-

bp direct repeat deletion at the origin of replication (*ori*), an Indel in VirB8 (G-> GA) and a non-

synonymous substitution in a hypothetical protein (I97T). Isolates with this deletion had higher

copies of IncX1 than ancestral strain carrying the extra 22-bp direct repeat. This result supports

other studies showing that the deletion of iterons increased plasmid copy number of theta

plasmids (reviewed in (90)). For example, Millan et al (91) showed that mutations acquired in

the OriV of a synthetic ColE1-like plasmid (pBGT) carrying bla$_{TEM-1}$ β-lactamase gene increased

plasmid copy number and subsequently increased ceftazidime resistance several folds.

To expand beyond the *S*. Heidelberg strains used in our study, we performed an alignment of

the *ori* of closed "circular" IncX1 plasmids of approximately 37kb present in *S*. Heidelberg

strains available in NCBI. We observed that all the strains analyzed carried only two iterons (Fig.

2b), which implies that this mutation is overrepresented in sequenced *S*. Heidelberg carrying

IncX1 plasmids. The majority of these strains were from clinical settings (Supplementary Table

8) which suggests they are being recovered more often and may be associated with an increase in

fitness of those strains as was also observed in our study. Taken together, our results support the

idea that iterons regulate IncX1 plasmid replication and their compatibility with other plasmids.

Mutations acquired on plasmids and chromosomes after an HGT event have been shown to

compensate for the fitness cost associated with the acquisition of a new MGE (43, 91-93). Here,

strains carrying Col plasmids had a total of 4 plasmids and persisted for up to 14 days, which is a

"plasmid paradox," where the acquisition and maintenance of multiple plasmids over the long

term is supposed to impose a burden on their host (84, 85). This principle has been elucidated

upon, where positive epistasis minimized the cost associated with carrying small plasmids of

Pseudomonas aeruginosa over a long-time scale (42). Recently, a study showed that the

acquisition of three ColE1 plasmids in naïve *H. influenzae* hosts did not impose a fitness cost as

long as the bacterium compensated for the first plasmid (94). The presence of one plasmid

favored the presence of a second plasmid and the presence of this second plasmid or third

plasmid did not impose an additional significant biological cost, even though these plasmids

produced a cost when they are alone in the cell. Col plasmid-bearing isolates from this study had

increased fitness compared to a isolate without a Col plasmid and carried 37 hotspot mutations

(Fig. 1d). It is unlikely that all observed SNPs/InDels are to compensate for Col plasmid

acquisition, as some will be compensatory for other MGE such as bacteriophages (95-97). For

example, the prophage Gifsy-2 was found intact in 97% of *S*. Heidelberg isolates sequenced in

our study, suggesting the PL environment may have selected them for increased fitness

(Supplementary Fig. 5). Further, the ubiquity of Gifsy-2 in *S*. Heidelberg strains has been

reported, which infers they are important for *S*. Heidelberg survival (13). We identified a hotspot

mutation in the intergenic region upstream of cell division protein (FtsZ) of Gifsy-2, an

important protein for cell division and lambda bacteriophage infection in *E. coli* (98) (Table 2).

In summary, it is more than likely that the hotspot mutations observed in this study are

associated with HGT events involving plasmids and prophages, in addition to mutations selected

for persistence in PL.

Plasmid and chromosome encoded genes mediated antibiotic resistance in *S*. Heidelberg.

Another significant observation from this study is the decrease in susceptibility of Col plasmid-

bearing strains towards kanamycin, tobramycin and fosfomycin. We do not know what these

reductions in inhibition zones mean in terms of MIC as there are no current CLSI standards (46)

for *S*. Heidelberg. In addition, it has been shown that simple back- calculations from inhibition

zones to MIC can be confounded by several covariates such as serotype, host species, and year of

isolation (99). The chickens raised on the PL used for this study were not administered any drugs

(Personal Communication- Dr. Casey Ritz, University of Georgia), thus experimental selection

for these resistance phenotypes is highly unlikely. It is a possibility that Col plasmid selection or

population factors like competitive exclusion and the presence of "cheaters" (100-102) could

account for this result but more research will be needed to test these possibilities.

Despite the fact that the acquisition of Col plasmids coincided with the observed change in

susceptibility, we cannot say for sure if these plasmids played any role. The putative

aminoglycoside acetyltransferase (aac-(6')) gene carried on ColE1-6kb represents a fragmented

aac-6' gene and shares significant homology with the N-terminal of aac(6')-Ib of *K. pneumoniae*.

Its small size (81 amino acid residues) compared to other known aac genes (Fig. 5a), and the

absence of homologs in all antibiotic resistance genes (ARG) databases queried suggests it may

be an uncharacterized ARG. Importantly, its presence on ColE1 will result in more copies of

these gene compared to the aac-(6') gene carried as a single copy on the chromosome. More

studies will be required to determine if these two aac-(6') genes confer decreased susceptibility

towards the aforementioned aminoglycosides.

Regarding fosfomycin, Rehmann et al (54) determined that *S*. Heidelberg strains with an

inhibition zone of 26 mm were resistant to >32 µg/ml of fosfomycin, which suggests that

evolved *S*. Heidelberg strains carrying Col plasmids in our study may be resistant to fosfomycin.

The fosA7 gene is present in many *S*. Heidelberg genomes and resistance to fosfomycin has been

demonstrated in a few (54). In addition, *S*. Heidelberg harbors other chromosomal genes

associated with fosfomycin resistance including *glpT* and *uhpT* (103). We did not observe any

mutation in these genes, thus cannot correlate fosfomycin resistance to a specific ARG. This

result shows the intrinsic bias that could be associated with estimating resistance phenotype from

only known ARG that are available in curated databases. More concerning, is the spontaneous

mutant (SH-17BBPW-2813) that emerged with reduced susceptibility to 3 classes of antibiotics

following overnight enrichment in buffered peptone water (BPW). One explanation is that BPW

enriched for the limited strains present in PL or it created an environment where only the fittest

strain emerged. Therefore, we argue that chromosomal mutations in addition to Col plasmids are

responsible for the reduced susceptibility displayed by evolved *S*. Heidelberg isolates.

Results from this PL microevolution experiment suggest that chromosomal mutations and

Col plasmids contributed to the fitness of *S*. Heidelberg. Col plasmids were present in all *S*.

Heidelberg isolates sequenced after 7 days in PL suggesting that their presence is required for

survival. However, the population of SH-116 strains were significantly lower in PL at day 14

compared to SH-2813 populations. We expected SH-116 to be in higher abundance than SH-

2813 since the ancestral populations carried Col plasmids. This was not the case in this study

(Fig. 1a). SH-116 strains were only detected in 1 of 6 microcosms by direct count and in 3 of 6

after enrichment in buffered peptone water (BPW) after 14 days incubation in PL. In contrast,

SH-2813 was recovered by direct count in all 6 microcosms (Supplementary Table 1). The

dynamics observed in our experiment can be modeled according to Watve et al (104) Rock-

Paper-Scissors (RPS) model of sociobiological control of plasmid copy number. In this model,

when a mutant bacterium with higher copy number plasmid was introduced in the presence of a

wildtype low copy plasmid, the wildtype was driven to extinction. This high copy mutant was

then vulnerable to invasion by another mutant that had higher copy numbers, which in turn was

vulnerable to invasion by another mutant with higher plasmid copies. In our scenario, SH-116$_{anc}$

population with lower copy ColE1 plasmids were displaced by those with higher copy ColE1

plasmids mobilized from PL microbiota until the point at which the intrinsic survival rate of *S*.

Heidelberg was greatly reduced (Fig. 1a). In contrast, a stable co-existence of SH-2813 of low,

intermediate and high copy ColE1 plasmids was observed in the SH-2813 population. Based on

the RPS model, it is likely that SH-2813 isolates with higher copy plasmids will end up invading

isolates with lower copies. Therefore we should expect a system without a unique equilibrium.

The ColE1 plasmids carried by SH-116 isolates had mob protein (MbeC) sequences that

were represented by 11 evolved isolates on the reconstructed phylogenetic tree (Fig. 4c), whereas

the SH-2813 isolates had only 3 evolved isolates. This suggests that SH-116 may be acquiring

new ColE1 plasmids from bacteria present in PL. This observation was corroborated by qPCR on

extracted DNA from PL with Col plasmid gene primer sets. At day 0 and 14, the copies of

ColE1-6kb and ColpVC were significantly higher in SH-116-PLE microcosms than SH-2813-

PLE (Fig. 7b), which supports the notion that the original SH-116 strains may have been

displaced by those carrying higher copy ColE1 plasmids. For the BHI microcosms, we did not

detect any plasmid gene even though 16S rRNA result showed an abundance of bacterial DNA

(data not shown). We have attributed this discrepancy to the significant differences in moisture

and pH between BHI and PLE microcosms. To avoid contamination and mix up of strains,

evolution experiments with SH-2813 and SH-116 strains were done on different days. In

addition, incubation of microcosms was done in separate incubators (similar make and model)

and in separate laboratories. The plastic containers with BHI and microcosms were placed on the

upper and bottom layer of the incubators. To avoid drying out of PL in incubator, a 2-liter bucket

of water was placed on the upper layer next to the BHI microcosms. An unforeseen outcome of

this experimental setup was an increase in moisture for BHI microcosms. Moisture has been

shown to be a significant factor changing the microbiota of PL (7). Thus, it is plausible that this

difference in moisture reduced the population of resident microbiota carrying Col plasmids in the

BHI microcosms compared to PLE. Further research from our laboratory will be exploring this

hypothesis.

CONCLUSION

Hotspot mutations and ColE1 plasmids acquired under positive selection from PL contributed to the persistence of *S*. Heidelberg observed in this study; however the identity of the bacterial donors of these Col plasmids remains unknown. Further, due to space limitation we did not discuss in detail exactly how each mutation observed in evolved isolates of *S*. Heidelberg can confer a fitness advantage in PL. Also, we did not identify mutations that could have been due to homologous intragenic recombination events. Owing to the large amount of information obtained from the SNPs/InDels analyzed across 86 genomes, it was not practical to perform an in-depth analysis on every one of them. Therefore, we have provided a list of all the identified mutations, their corresponding VCF files and UNIX command line used (Supplementary Table 4 and Dryad Digital Repository: https://doi.org/10.5061/dryad.pc6tp4q).

ACKNOWLEDGEMENTS

We are grateful to Dr. Casey Ritz for providing us with the poultry litter used in this study and for Caroline Plymel for her help with sample analysis. We wish to thank Dr. Karamat Sistani for valuable assistance with litter nutrient analysis. We also appreciate Steven Lankin and Mary Katherine Crews for manuscript review. Any opinions expressed in this paper are those of the authors and do not necessarily reflect the official positions and policies of the USDA and any mention of products or trade names does not constitute recommendation for use. The authors declare no competing commercial interests in relation to the submitted work.

Supplementary Figures

Figure S1. Experimental design.

Figure S2. Principal component analysis (PCA) of poultry litter characterizing the differences

between microcosms. Data represent 48 microcosms. Positively correlated variables point to the

same side of the plot and negative correlated variables point to opposite sides of the graph.

Symbol: Colored by contributions to the principal component (Contrib).

Figure S3. ProgressiveMauve alignment of (**a**) IncX1 (**b**) ColE1-6kb and (**c**) ColE1-4kb whole

plasmid DNA sequence from selected isolates from this study. DNA regions that differ are

highlighted with horizontal black rectangular boxes.

Figure S4. Linear maps of Col plasmids detected in an incomplete assembly of a *S*. Heidelberg

strain. Strain NS-007 (NCBI accession: SAMN08031253) was isolated from a broiler breeder

farm in the USA under Bioproject PRJNA417775. Contig size for ColE1-6kb, ColE1-4kb and

ColpVC-2kb are 6183, 4778 and 2105 bp, respectively.

Figure S5. Intact prophages in sequenced genomes of *S*. Heidelberg from this study.

Figure S6. Maximum likelihood tree of *S*. Heidelberg strains available in GenBank based on

23S-5S internal transcribed spacer. The consensus 23S-5S ITS sequence of isolates included

from this study were extracted by mapping whole genome sequence reads to *S*. Heidelberg

reference genome (CP016573)

Figure S7. Gene expression of selected Col plasmid genes determined by qRT-PCR for (**a**) SH-

116$_{anc}$ (**b**) SH-116$_{evol}$ and (**c**) SH-2813$_{evol}$ isolates. Log2 fold-change is relative to the expression

of chromosome encoded reference genes *gapA* and *gyrB*. Symbols: replication protein of

ColpVC (ColpVC-rep), abortive phage infection protein of ColE1-6kb (ColE1-6kb-Abi),

macrophage stimulating factor protein of ColE1-4kb (ColE1-4kb-MSF) and hypothetical protein

(hypo).

Figure S8. Circular map of IncX1 plasmid of isolate SH-33A0.5-116-Day7-PLE.

Supplementary Tables

Table S1. Bacteria concentration in poultry litter microcosms.

Table S2. Physicochemical properties of poultry litter microcosms.

Table S3. Metadata on isolates sequenced.

Table S4. SNPs/InDels identified in sequenced *S*. Heidelberg genomes from this study compared

to the reference genome.

Table S5. Identified plasmids topology.

Table S6. Plasmid copy number determined by whole genome sequencing coverage.

Table S7. Antibiotic susceptibility test results.

Table S8. Metadata on NCBI *S*. Heidelberg strains carrying IncX1 plasmid.

Table S9. Primers used in this study.

Supplementary Files

Supplementary File 1. Quality assessment and coverage of sequenced SH-2813 strains.

Supplementary File 2. Quality assessment and coverage of sequenced SH-116 strains.

Supplementary File 3. Disc diffusion inhibition zone measurements.

References:

1.   USDA, NASS. Poultry production and value - 2013 summary. ISSN:1949-1573 ed2014.

p. 1-14. Retrieved from

"https://usda.mannlib.cornell.edu/usda/nass/PoulProdVa//2010s/2014/PoulProdVa-04-29-

2014.pdf".

2.   MacDonald JM. The economic organization of U.S. broiler production. Washington,

D.C.: Economic Research Service, USDA; 2008 June 2008. Report No.: EIB- 38 Contract No.:

EIB- 38.

3. Wang L, Lilburn M, Yu Z. Intestinal microbiota of broiler chickens as affected by litter

management regimens. Front Microbiol. 2016;7:593.

4. Dunlop MW, McAuley J, Blackall PJ, Stuetz RM. Water activity of poultry litter:

Relationship to moisture content during a grow-out. J Environ Manage. 2016;172:201-6.

5. Cressman MD, Yu ZT, Nelson MC, Moeller SJ, Lilburn MS, Zerby HN. Interrelations

between the microbiotas in the litter and in the intestines of commercial broiler chickens.

Applied and Environmental Microbiology. 2010;76(19):6572-82.

6. Casey R, Fairchild B, Lacy M. Litter quality and broiler performance. University of

Georgia Cooperative Extension bulletin No. 1267. 2005.

7. Lovanh N, Cook KL, Rothrock MJ, Miles DM, Sistani K. Spatial shifts in microbial

population structure within poultry litter associated with physicochemical properties. Poultry

Science. 2007;86(9):1840-9.

8. Schefferle HE. The microbiology of built up poultry litter. J Appl Bacteriol.

1965;28(3):403-11.

9. Roll VF, Dai Pra MA, Roll AP. Research on *Salmonella* in broiler litter reused for up to

14 consecutive flocks. Poult Sci. 2011;90(10):2257-62.

10. Opara OO, Carr LE, Russek-Cohen E, Tate CR, Mallinson ET, Miller RG, et al.

Correlation of water activity and other environmental conditions with repeated detection of

Salmonella contamination on poultry farms. Avian Dis. 1992;36(3):664-71.

11. Rigby CE, Pettit JR. Some factors affecting *Salmonella*-Typhimurium infection and

shedding in chickens raised on litter. Avian Dis. 1979;23(2):442-55.

12.   Voss-Rech D, Trevisol IM, Brentano L, Silva VS, Rebelatto R, Jaenisch FRF, et al.

Impact of treatments for recycled broiler litter on the viability and infectivity of microorganisms.

Vet Microbiol. 2017;203:308-14.

13.   Hoffmann M, Zhao S, Pettengill J, Luo Y, Monday SR, Abbott J, et al. Comparative

genomic analysis and virulence differences in closely related *Salmonella enterica* serotype

Heidelberg isolates from humans, retail meats, and animals. Genome Biol Evol. 2014;6(5):1046-

68.

14.   Gieraltowski L, Higa J, Peralta V, Green A, Schwensohn C, Rosen H, et al. National

outbreak of multidrug resistant *Salmonella* Heidelberg infections linked to a single poultry

company. PLoS One. 2016;11(9):e0162369.

15.   Bekal S, Berry C, Reimer AR, Van Domselaar G, Beaudry G, Fournier E, et al.

Usefulness of high-quality core genome single-nucleotide variant analysis for subtyping the

highly clonal and the most prevalent *Salmonella enterica* serovar Heidelberg clone in the context

of outbreak investigations. J Clin Microbiol. 2016;54(2):289-95.

16.   Hoffmann M, Zhao SH, Pettengill J, Luo Y, Monday SR, Abbott J, et al. Comparative

Genomic Analysis and Virulence Differences in Closely Related *Salmonella enterica* Serotype

Heidelberg Isolates from Humans, Retail Meats, and Animals. Genome Biol Evol.

2014;6(5):1046-68.

17.   Edirmanasinghe R, Finley R, Parmley EJ, Avery BP, Carson C, Bekal S, et al. A Whole-

Genome Sequencing Approach To Study Cefoxitin-Resistant *Salmonella enterica* Serovar

Heidelberg Isolates from Various Sources. Antimicrob Agents Chemother. 2017;61(4).

18.   Orlek A, Phan H, Sheppard AE, Doumith M, Ellington M, Peto T, et al. A curated dataset

of complete Enterobacteriaceae plasmids compiled from the NCBI nucleotide database. Data

Brief. 2017;12:423-6.

19.   Folster JP, Pecic G, Rickert R, Taylor J, Zhao S, Fedorka-Cray PJ, et al. Characterization

of multidrug-resistant *Salmonella enterica* serovar Heidelberg from a ground turkey-associated

outbreak in the United States in 2011. Antimicrob Agents Chemother. 2012;56(6):3465-6.

20.   Labbe G, Ziebell K, Bekal S, Macdonald KA, Parmley EJ, Agunos A, et al. Complete

Genome Sequences of 17 Canadian Isolates of *Salmonella* enterica subsp. *enterica* Serovar

Heidelberg from Human, Animal, and Food Sources. Genome Announc. 2016;4(5).

21.   Labbe G, Edirmanasinghe R, Ziebell K, Nash JH, Bekal S, Parmley EJ, et al. Complete

Genome and Plasmid Sequences of Three Canadian Isolates of *Salmonella* enterica subsp.

enterica Serovar Heidelberg from Human and Food Sources. Genome Announc. 2016;4(1).

22.   Evans PS, Luo Y, Muruvanda T, Ayers S, Hiatt B, Hoffman M, et al. Complete Genome

Sequences of *Salmonella enterica* Serovar Heidelberg Strains Associated with a Multistate Food-

Borne Illness Investigation. Genome Announc. 2014;2(3).

23.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes:

a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol.

2012;19(5):455-77.

24.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: a fast phage search tool.

Nucleic Acids Res. 2011;39(Web Server issue):W347-52.

25.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.

2014;30(14):2068-9.

26.   Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017:

expansion and model-centric curation of the comprehensive antibiotic resistance database.

Nucleic Acids Res. 2017;45(D1):D566-D73.

27.   Orlek A, Phan H, Sheppard AE, Doumith M, Ellington M, Peto T, et al. Ordering the

mob: Insights into replicon and MOB typing schemes from analysis of a curated dataset of

publicly available plasmids. Plasmid. 2017;91:42-52.

28.   Darling AE, Mau B, Perna NT. ProgressiveMauve: multiple genome alignment with gene

gain, loss and rearrangement. PLoS One. 2010;5(6):e11147.

29.   Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and

space complexity. BMC Bioinformatics. 2004;5:113.

30.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80.

31.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 2014;30(9):1312-3.

32.   Li H, Durbin R. Fast and accurate short read alignment with Burrows−Wheeler

transform. Bioinformatics. 2009;25(14):1754-60.

33.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence

alignment/map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.

34.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The

Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA

sequencing data. Genome Res. 2010;20(9):1297-303.

35.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant

call format and VCFtools. Bioinformatics. 2011;27(15):2156-8.

36.   Lischer HE, Excoffier L. PGDSpider: an automated data conversion tool for connecting

population genetics and genomics programs. Bioinformatics. 2012;28(2):298-9.

37.   Jukes TH, Cantor CR. Evolution of protein molecules. Mammalian protein metabolism.

1969;3(21):132.

38.   Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics

and parallel computing. Nat Methods. 2012;9(8):772.

39.   Lee C, Kim J, Shin SG, Hwang S. Absolute and relative QPCR quantification of plasmid

copy number in *Escherichia coli*. J Biotechnol. 2006;123(3):273-80.

40.   Oladeinde A. Interacting Effects of Sunlight, Dissolved Organic Matter and Reactive

Oxygen Species on *Escherichia coli* and *Enterococcus faecalis* Survival. University of Georgia,

2017, PhD dissertation.

41.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative PCR and the $2^{-\Delta\Delta CT}$ method. methods. 2001;25(4):402-8.

42.   San Millan A, Heilbron K, MacLean RC. Positive epistasis between co-infecting

plasmids promotes plasmid survival in bacterial populations. ISME J. 2014;8(3):601-12.

43.   Loftie-Eaton W, Bashford K, Quinn H, Dong K, Millstein J, Hunter S, et al.

Compensatory mutations improve general permissiveness to antibiotic resistance plasmids. Nat

Ecol Evol. 2017;1(9):1354-63.

44.   Food and Drug Administration (FDA). The National Antimicrobial Resistance

Monitoring System Manual of Laboratory Methods. Second ed2015. Retrieved from "

https://www.fda.gov/downloads/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/Natio

nalAntimicrobialResistanceMonitoringSystem/UCM453381.pdf"

45.   Hudzicki J. Kirby-bauer disk diffusion susceptibility test protocol. 2009.

46.   Clinical Laboratory Standard Institute (CLSI). Performance standards for antimicrobial

susceptibility testing. 2009.

47.   Liakopoulos A, Geurts Y, Dierikx CM, Brouwer MS, Kant A, Wit B, et al. Extended-

Spectrum Cephalosporin-Resistant *Salmonella enterica* serovar Heidelberg Strains, the

Netherlands(1). Emerg Infect Dis. 2016;22(7):1257-61.

48.   Wasilenko JL, Fratamico PM, Narang N, Tillman GE, Ladely S, Simmons M, et al.

Influence of primer sequences and DNA extraction method on detection of non-O157 Shiga

toxin-producing *Escherichia coli* in ground beef by real-time PCR targeting the eae, stx, and

serogroup-specific genes. J Food Prot. 2012;75(11):1939-50.

49.   Nandi S, Maurer JJ, Hofacre C, Summers AO. Gram-positive bacteria are a major

reservoir of Class 1 antibiotic resistance integrons in poultry litter. P Natl Acad Sci USA.

2004;101(18):7118-22.

50.   Chattoraj DK. Control of plasmid DNA replication by iterons: no longer paradoxical.

Mol Microbiol. 2000;37(3):467-76.

51.   Nordstrom K. Control of plasmid replication--how do DNA iterons set the replication

frequency? Cell. 1990;63(6):1121-4.

52.   Bustamante P, Iredell JR. Carriage of type II toxin-antitoxin systems by the growing

group of IncX plasmids. Plasmid. 2017;91:19-27.

53.   Page R, Peti W. Toxin-antitoxin systems in bacterial growth arrest and persistence. Nat

Chem Biol. 2016;12(4):208-14.

54.   Rehman MA, Yin X, Persaud-Lachhman MG, Diarra MS. First Detection of a

Fosfomycin Resistance Gene, fosA7, in *Salmonella enterica* Serovar Heidelberg Isolated from

Broiler Chickens. Antimicrob Agents Chemother. 2017;61(8).

55.   Tada T, Miyoshi-Akiyama T, Dahal RK, Mishra SK, Shimada K, Ohara H, et al.

Identification of a novel 6'-N-aminoglycoside acetyltransferase, AAC(6')-Iak, from a multidrug-

resistant clinical isolate of *Stenotrophomonas maltophilia*. Antimicrob Agents Chemother.

2014;58(10):6324-7.

56.   Tada T, Miyoshi-Akiyama T, Shimada K, Shimojima M, Kirikae T. novel 6'-n-

aminoglycoside acetyltransferase AAC(6')-Iaj from a clinical isolate of *Pseudomonas*

aeruginosa. Antimicrob Agents Chemother. 2013;57(1):96-100.

57.   Tada T, Miyoshi-Akiyama T, Shimada K, Dahal RK, Mishra SK, Ohara H, et al. A Novel

6'-N-Aminoglycoside Acetyltransferase, AAC(6')-Ial, from a Clinical Isolate of *Serratia*

marcescens. Microb Drug Resist. 2016;22(2):103-8.

58.   Wang X, Kim Y, Ma Q, Hong SH, Pokusaeva K, Sturino JM, et al. Cryptic prophages

help bacteria cope with adverse environments. Nat Commun. 2010;1:147.

59.   Bobay LM, Touchon M, Rocha EP. Pervasive domestication of defective prophages by

bacteria. Proc Natl Acad Sci U S A. 2014;111(33):12127-32.

60.   Feiner R, Argov T, Rabinovich L, Sigal N, Borovok I, Herskovits AA. A new perspective

on lysogeny: prophages as active regulatory switches of bacteria. Nat Rev Microbiol.

2015;13(10):641-50.

61.   Golubeva YA, Slauch JM. *Salmonella enterica* serovar Typhimurium periplasmic

superoxide dismutase SodCI is a member of the PhoPQ regulon and is induced in macrophages. J

Bacteriol. 2006;188(22):7853-61.

62.   Pacello F, Ceci P, Ammendola S, Pasquali P, Chiancone E, Battistoni A. Periplasmic

Cu,Zn superoxide dismutase and cytoplasmic Dps concur in protecting *Salmonella enterica*

serovar Typhimurium from extracellular reactive oxygen species. Biochim Biophys Acta. 2008;1780(2):226-32.

63.   Meng X, Liu X, Zhang L, Hou B, Li B, Tan C, et al. Virulence characteristics of extraintestinal pathogenic *Escherichia coli* deletion of gene encoding the outer membrane protein X. J Vet Med Sci. 2016;78(8):1261-7.

64.   Chattopadhyay S, Paul S, Kisiela DI, Linardopoulou EV, Sokurenko EV. Convergent molecular evolution of genomic cores in *Salmonella enterica* and *Escherichia coli*. J Bacteriol. 2012;194(18):5002-11.

65.   Chattopadhyay S, Weissman SJ, Minin VN, Russo TA, Dykhuizen DE, Sokurenko EV. High frequency of hotspot mutations in core genes of *Escherichia coli* due to short-term positive selection. Proc Natl Acad Sci U S A. 2009;106(30):12412-7.

66.   Tenaillon O, Rodriguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, et al. The molecular diversity of adaptive convergence. Science. 2012;335(6067):457-61.

67.   Chou SH, Galperin MY. Diversity of Cyclic Di-GMP-Binding Proteins and Mechanisms. J Bacteriol. 2016;198(1):32-46.

68.   Dow JM, Fouhy Y, Lucey J, Ryan RP. Cyclic di-GMP as an intracellular signal regulating bacterial biofilm formation. Biofilm mode of life. 2007.

69.   Siryaporn A, Goulian M. Cross-talk suppression between the CpxA–CpxR and EnvZ–OmpR two-component systems in E. coli. Molecular microbiology. 2008;70(2):494-506.

70.   Sun S, Berg OG, Roth JR, Andersson DI. Contribution of gene amplification to evolution of increased antibiotic resistance in *Salmonella* Typhimurium. Genetics. 2009;182(4):1183-95.

71.   Noinaj N, Guillier M, Barnard TJ, Buchanan SK. TonB-dependent transporters: regulation, structure, and function. Annu Rev Microbiol. 2010;64:43-60.

72.   Cadieux N, Bradbeer C, Kadner RJ. Sequence changes in the ton box region of BtuB

affect its transport activities and interaction with TonB protein. J Bacteriol. 2000;182(21):5954-

61.

73.   Anton AI, Martinez-Murcia AJ, Rodriguez-Valera F. Intraspecific diversity of the 23S

rRNA gene and the spacer region downstream in *Escherichia coli*. J Bacteriol.

1999;181(9):2703-9.

74.   Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. Tests of parallel

molecular evolution in a long-term experiment with *Escherichia coli*. Proc Natl Acad Sci U S A.

2006;103(24):9107-12.

75.   Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al.

STRING v10: protein–protein interaction networks, integrated over the tree of life. Nucleic acids

research. 2014:gku1003.

76.   Dy RL, Przybilski R, Semeijn K, Salmond GP, Fineran PC. A widespread bacteriophage

abortive infection system functions through a Type IV toxin-antitoxin mechanism. Nucleic Acids

Res. 2014;42(7):4590-605.

77.   Actis LA, Tolmasky ME, Crosa JH. Bacterial plasmids: replication of extrachromosomal

genetic elements encoding resistance to antimicrobial compounds. Front Biosci. 1999;4:D43-62.

78.   Francia MV, Varsaki A, Garcillan-Barcia MP, Latorre A, Drainas C, de la Cruz F. A

classification scheme for mobilization regions of bacterial plasmids. FEMS Microbiol Rev.

2004;28(1):79-100.

79.   Smillie C, Garcillan-Barcia MP, Francia MV, Rocha EP, de la Cruz F. Mobility of

plasmids. Microbiol Mol Biol Rev. 2010;74(3):434-52.

80.   Varsaki A, Moncalian G, Garcillan-Barcia Mdel P, Drainas C, de la Cruz F. Analysis of

ColE1 MbeC unveils an extended ribbon-helix-helix family of nicking accessory proteins. J

Bacteriol. 2009;191(5):1446-55.

81.   Varsaki A, Lamb HK, Eleftheriadou O, Vandera E, Thompson P, Moncalian G, et al.

Interaction between relaxase MbeA and accessory protein MbeC of the conjugally mobilizable

plasmid ColE1. FEBS Lett. 2012;586(6):675-9.

82.   Hansen LH, Bentzon-Tilia M, Bentzon-Tilia S, Norman A, Rafty L, Sorensen SJ. Design

and synthesis of a quintessential self-transmissible IncX1 plasmid, pX1.0. PLoS One.

2011;6(5):e19912.

83.   Burmolle M, Norman A, Sorensen SJ, Hansen LH. Sequencing of IncX-plasmids

suggests ubiquity of mobile forms of a biofilm-promoting gene cassette recruited from *Klebsiella*

pneumoniae. PLoS One. 2012;7(7):e41259.

84.   Hall JPJ, Brockhurst MA, Dytham C, Harrison E. The evolution of plasmid stability: Are

infectious transmission and compensatory evolution competing evolutionary trajectories?

Plasmid. 2017;91:90-5.

85.   MacLean RC, San Millan A. Microbial Evolution: Towards Resolving the Plasmid

Paradox. Curr Biol. 2015;25(17):R764-7.

86.   Schroder G, Lanka E. TraG-like proteins of type IV secretion systems: functional

dissection of the multiple activities of TraG (RP4) and TrwB (R388). J Bacteriol.

2003;185(15):4371-81.

87.   Hayes F. Toxins-antitoxins: plasmid maintenance, programmed cell death, and cell cycle

arrest. Science. 2003;301(5639):1496-9.

88.   Lopatkin AJ, Meredith HR, Srimani JK, Pfeiffer C, Durrett R, You L. Persistence and

reversal of plasmid-mediated antibiotic resistance. Nat Commun. 2017;8(1):1689.

89.   Giraldo R, Fernandez-Tresguerres ME. Twenty years of the pPS10 replicon: insights on

the molecular mechanism for the activation of DNA replication in iteron-containing bacterial

plasmids. Plasmid. 2004;52(2):69-83.

90.   Chattoraj DK, Schneider TD. Replication control of plasmid P1 and its host chromosome:

the common ground. Prog Nucleic Acid Res Mol Biol. 1997;57:145-86.

91.   San Millan A, Escudero JA, Gifford DR, Mazel D, MacLean RC. Multicopy plasmids

potentiate the evolution of antibiotic resistance in bacteria. Nat Ecol Evol. 2016;1(1):10.

92.   Harrison E, Guymer D, Spiers AJ, Paterson S, Brockhurst MA. Parallel compensatory

evolution stabilizes plasmids across the parasitism-mutualism continuum. Curr Biol.

2015;25(15):2034-9.

93.   San Millan A, Pena-Miller R, Toll-Riera M, Halbert ZV, McLean AR, Cooper BS, et al.

Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids.

Nat Commun. 2014;5:5208.

94.   Santos-Lopez A, Bernabe-Balas C, San Millan A, Ortega-Huedo R, Hoefer A, Ares-

Arroyo M, et al. Compensatory evolution facilitates the acquisition of multiple plasmids in

bacteria. bioRxiv. 2017.

95.   Koskella B, Brockhurst MA. Bacteria-phage coevolution as a driver of ecological and

evolutionary processes in microbial communities. FEMS Microbiol Rev. 2014;38(5):916-31.

96.   Harrison E, Hall JPJ, Paterson S, Spiers AJ, Brockhurst MA. Conflicting selection alters

the trajectory of molecular evolution in a tripartite bacteria-plasmid-phage interaction. Mol Ecol.

2017;26(10):2757-64.

97.   Harrison E, Truman J, Wright R, Spiers AJ, Paterson S, Brockhurst MA. Plasmid

carriage can limit bacteria-phage coevolution. Biol Lett. 2015;11(8).

98.   Haeusser DP, Hoashi M, Weaver A, Brown N, Pan J, Sawitzke JA, et al. The Kil peptide

of bacteriophage lambda blocks *Escherichia coli* cytokinesis via ZipA-dependent inhibition of

FtsZ assembly. PLoS Genet. 2014;10(3):e1004217.

99.   Hoelzer K, Cummings KJ, Warnick LD, Schukken YH, Siler JD, Grohn YT, et al. Agar

disk diffusion and automated microbroth dilution produce similar antimicrobial susceptibility

testing results for *Salmonella* serotypes Newport, Typhimurium, and 4,5,12:i-, but differ in

economic cost. Foodborne Pathog Dis. 2011;8(12):1281-8.

100.   Vasse M, Noble RJ, Akhmetzhanov AR, Torres-Barcelo C, Gurney J, Benateau S, et al.

Antibiotic stress selects against cooperation in the pathogenic bacterium Pseudomonas

aeruginosa. Proc Natl Acad Sci U S A. 2017;114(3):546-51.

101.   Bruger EL, Waters CM. Bacterial Quorum Sensing Stabilizes Cooperation by Optimizing

Growth Strategies. Appl Environ Microbiol. 2016;82(22):6498-506.

102.   Leinweber A, Fredrik Inglis R, Kummerli R. Cheating fosters species co-existence in

well-mixed bacterial communities. ISME J. 2017;11(5):1179-88.

103.   Castaneda-Garcia A, Rodriguez-Rojas A, Guelfo JR, Blazquez J. The glycerol-3-

phosphate permease GlpT is the only fosfomycin transporter in *Pseudomonas aeruginosa*. J

Bacteriol. 2009;191(22):6968-74.

104.   Watve MM, Dahanukar N, Watve MG. Sociobiological control of plasmid copy number

in bacteria. PLoS One. 2010;5(2):e9328.

1289

1290

A

Legend:
- IncX1-rep (red)
- ColpVC-rep (blue)
- ColE1-6kb-Abi (purple)
- ColE1-4kb-MSF (yellow)
- 16S rDNA (green)

B

ColE1-6kb | ColpVC-2kb | 16S rDNA

Legend:
- SH-2813-PLE (red)
- SH-116-PLE (blue)
- Control-No Salmonella (black)

A

```
                        10        20        30        40        50        60        70        80        90       100
                         |         |         |         |         |         |         |         |         |         |
ColE1-6kb/1-79      MAFYVDSTAKFLTKTTEPEGSVWRYYYK---PG------------NTCPA----SGIYRCTNCGDEVTCNKDDPLPPQNHRQHNV-PGDIR----WELVVMTK
SH-Ancestral-2813/1-8-----------MDIRQMNKTNLEHWRGLRKQLWPGPDDAHLADGEEILQA-DYLASFIAMAD-GVAIGFAD---ASIRHDYVNGCDSSPVAFLEGIFVLPSFR
K.pneumoniae/1-75      ----MAQYKYGNYLNQSQNEAYDTLHS--PG------------DTCAN----SGIYRCESCGDEIASNKGNPFPPQNHHQHAPGSGPIR---WKLIVWAQ
Thiomonas/1-75         ----MASFKYAIHLRQDSNNAFDQKHP--PG------------IVAPY----AGIYRCVVCGDEIGIAQGHTLPPQNHHQHPAGRGRIE---WQLLVQAH
Stenotrophomonas/1-92  ----MTGSAATIRPAKAADAVAWAQLRLGLWPD--ADDPLETLVAALAE-DAGAVFLACAAGGQAIGFAE---VRLRHDYVNGTDSSPVGFLEGWYVQPQWQ
P.aeruginosa/10-108    APSMANTPVVGVVVPCKTPDHPGWLELRLQLWPDGSTEEFLPEMAAACAEPDRFGQFLFLSPGGLAEGLVE---VALRTDYVNGTESSPVAFLEGVFVVPASR
```

B

C

Figure containing six bar charts showing No. of Isolates vs Zone of Inhibition (mm) for kanamycin, Tobramycin, Gentamicin, Neomycin, Netilmicin, and Fosfomycin.

Legend: SH-116-Day0, SH-116-Day14, SH-2813-Day0, SH-2813-Day14

Decrease in susceptibility to antibiotics