A high-throughput method for unbiased quantitation and categorisation of

nuclear morphology

Benjamin Matthew Skinner[1]

Claudia Cattoni Rathje[2]

Joanne Bacon[1]

Emma Elizabeth Philippa Johnson[1]

Erica Lee Larson[3,4]

Emily Emiko Konishi Kopania[4]

Jeffrey Martin Good[4]

Gullalaii Yousafzai[2]

Nabeel Ahmed Affara[1]

Peter James Ivor Ellis[2,5]

[1] Department of Pathology, University of Cambridge, Cambridge, CB2 1QP, UK

[2] School of Biosciences, University of Kent, Canterbury, CT2 7NJ, UK

[3] Department of Biological Sciences, University of Denver, Denver, CO, USA

[4] Division of Biological Sciences, University of Montana, MT, USA

[5] Corresponding author

Running title: Nuclear morphology analysis in mouse sperm

Keywords: spermatogenesis, morphometrics, fertility, image analysis, F1 cross

Abstract

The physical arrangement of chromatin in the nucleus is cell type and species specific. This

is particularly evident in sperm, in which most of the cytoplasm has been lost; the shape of

the nucleus reflects the shape of the cell. Mice have distinctive falciform ('hook shaped')

sperm heads and nuclei. Quantification of the differences in shape variation between mouse

species and lines often relies on manual measurement and classification that leads to

subjective results, making comparisons within and between samples difficult.

We have developed an analysis program for assessing the morphology of asymmetric

nuclei, and characterised the sperm of mice from a range of inbred, outbred and wild-derived

mouse lines. We find that laboratory lines have elevated sperm shape variability both within

and between samples in comparison to wild-derived inbred lines, and that sperm shape in

the F1 offspring of CBA and C57Bl6J lines is subtly affected by the direction of the cross.

Hierarchical clustering can distinguish distinct sperm shapes with greater efficiency and

reproducibility than even experienced manual assessors. We quantified the range of

morphological defects in the inbred BALB/c line, demonstrating we can identify different

morphological subgroups. This approach has applications for studies of sperm development,

infertility and toxicology.

1 Introduction

Nuclei are complex, dynamic structures within a cell. For many cell types, the nucleus is

generally spherical, but for other cell types the nucleus adopts a distinctive shape [1]. One of

the most profound changes to nuclear shape occurs during spermatogenesis: mammalian

sperm tend to have a spatulate, or 'paddle' shape, meaning the nucleus both condenses and

reshapes. The chromatin becomes wound ~4-6 times more tightly than in metaphase,

mediated via replacement of histones with smaller protamines [2], and various cytoskeletal

elements coordinating to shape the nucleus [3].

In rodents, this process is even more elaborate: most rodents, including mice, have a

falciform 'hook-shaped' sperm, with varying degrees of hook length and body shape

between species (e.g. [4]). The mouse sperm head shape develops through a series of

interacting mechanical forces, reshaping the nucleus via the cytoskeleton and

nucleoskeleton. The sperm head is divided into developmental 'modules', each of which is

shaped by particular cytoskeletal components [5]. When these processes go awry, distinct

morphological abnormalities can result (e.g. [6]), linking phenotype with the underlying

genetic alterations. The reshaping of the nucleus is itself a distinct process from the

chromatin condensation and repackaging [3]. Reshaping precedes transition and protamine

replacement, and chromatin condensation then follows.

Analytical methods for categorising and quantifying sperm head shape variation have

developed markedly over the years, and the advent of computational processing of images

has dramatically increased the quality of data we can capture, and the sophistication of the

analyses. To date, morphometric approaches in sperm have fallen into three main groups;

the measurement of basic parameters such as lengths, widths, and areas of objects, the use

of elliptic fourier analysis to investigate differences in the two dimensional outline of the

object, and the use of Procrustes analyses to examine differences in fixed landmarks within

the sperm head. Each of these approaches has advantages and disadvantages.

Basic measures such as area and length were the first statistics recorded describing sperm

morphology (e.g. [7–9]. These still remain useful, especially in situations such as CASA

analysis for fertility screening, in which an assessment of semen quality must be made

rapidly across many different cells [10]. However, the parameters measured by these

analyses are dominated by the size of the object, not the shape, and can make it difficult to

consistently assess the number of normal sperm across populations [11].

In contrast, elliptic Fourier descriptors [12] allow an arbitrary closed two dimensional shape

to be decomposed into harmonic amplitudes describing the curvature of the object

perimeter, thus allowing subtle variations in shape to be discovered [13]. This approach has

proved powerful for demonstrating differences between species, lines within a species, and

different treatments (e.g. [14–16]). However, the approach has the drawback that the shape

parameters and underlying mathematics are difficult for biologists to understand and relate

back to the biological structure that is affected [17]. Moreover, since Fourier analyses rely on

smooth harmonic deformations of an underlying elliptical outline, sharp points - such as

found at the tip of a mouse sperm - tend to be poorly fitted [18].

The third major method, Procrustes-based geometric morphometric analysis, uses

landmarks and semilandmarks within the object to align individual samples to consistent

size, position and orientation (e.g. [4]). Principal component analysis (PCA) can then be

used to identify the major varying landmarks distinguishing samples [5]. This approach has

the advantage of tightly relating the variation to physical structures within the object:

however, since objects are aligned by a least-squares method rotating about the centroid,

objects are susceptible to smearing of landmarks in highly variable regions, and can require time-consuming manual placement of landmarks.

In terms of the biological field of application, sperm shape analysis has proven useful in three main interrelated areas: infertility, speciation, and toxicology. In infertility, while abnormal sperm morphology is extremely common in infertile knockout lines, the role played by specific types and extents of shape defect remains to be elucidated, as does the extent to which teratozoospermia can be used as an indicator of other sperm defects (e.g. DNA damage or defective motility [19]). Deregulation of reproductive processes is a major contributor to speciation through the induction of hybrid male sterility [20]. In particular, sperm shape abnormalities are a feature of house mouse hybrid sterility, with a range of mapped quantitative trait loci known, particularly on the sex chromosomes but also on autosomes [21–24].

Sperm shape is used as an assessment of genotoxicity and/or reproductive toxicity of compounds (e.g. [25,26]. These studies often carry out a manual classification of sperm into various categories of morphological abnormality, based on previously described sperm shapes. The manual element thus makes this application both time consuming, and prone to operator bias. A further problem is that the classes of abnormality described are often arbitrarily chosen, and vary between studies. Use of a scoring chart, based on the morphological abnormalities typical for one experimental system, may therefore compromise the ability to quantitate abnormalities in a different system. It would be far more useful to have an automated and reproducible method that is able to discover categories of morphological abnormality within a sperm population, without prior training.

To address these needs for unbiased measurement, analysis and categorisation of nuclear morphologies, we have developed a new image analysis programme that generates

quantitative information on the underlying regions of the nucleus that differ within and

between samples, independent of nuclear size.

We have validated the software on different mouse lines, and can quickly analyse hundreds

of images. Here, we demonstrate the use of this software to compare a range of different

inbred, outbred and wild-derived lines (revealing the effects of inbreeding depression and

potentially hybrid dysgenesis), to unravel the morphological variation in a single sample

(revealing different classes of abnormality in an inbred line), and to trace genetic influences

on sperm morphology in a reciprocal F1 cross between CB57Bl6 and CBA lines (revealing

contrasting effects of the parental genomes on sperm size and shape).

Methods

Mouse lines

All animal procedures were in accordance with the United Kingdom Animal Scientific

Procedures Act 1986 and the University of Montana Institute for Animal Care and Use

Committee (protocol 002-13) and were subject to local ethical review. Animals were sourced

as indicated in Table 1; either from an approved supplier (Charles River Laboratories,

Manston, UK), bred at Cambridge University Central Biomedical Services (Home Office

licenses 80/2451 and 70/8925 held by PE), or bred at the University of Montana. Breeding

colonies at the University of Montana were established from mice purchased from Jackson

Laboratories (Bar Harbor, ME) or were acquired from Francois Bonhomme (University of

Montpellier). Animals were housed singly or in small groups, sacrificed via CO_2 followed by

cervical dislocation (UM) or only cervical dislocation and tissues collected *post mortem* for

analysis.

Line Name	Sample ID	Note	Samples Imaged	Source
C57Bl6/J	C57Bl6	Inbred	2 individual animals (C57 3, 4)	CRL
CBA/Ca	CBA	Inbred	3 individual animals (CBA1, 2, 3)	CRL
B6CBA	B6CBA	F1 offspring of C57Bl6 (f) and CBA (m) inbred lines	3 individual animals (B6CBA 1, 2, 4)	CRL
CBAB6	CBAB6	F1 offspring of CBA (f) and C57Bl6 (m) inbred lines	4 individual animals (CBAB6 1, 2, 3, 4)	CRL
CD1	CD1	Outbred	1 pool of 15 males	CRL
DBA/1J	DBA	Inbred	2 individual animals (DBA 1, 2)	CRL

BALB/cAnN Crl	BALB/c	Inbred	2 individual animals (Balbc 1, 2)	CRL
FVB/N	FVB	Inbred	2 individual animals (FVB 1, 2)	CRL
MF1Y^{RIII}	MF1Y^{RIII}	Outbred	2 pools (MF1YRIII 1, 2) of 8 males each	Bred at Uni. Cambridge
LEWES/EiJ	*M. m. domesticus*	*M. m. domesticus* Wild-derived inbred	2 pools (LEW 1, 2) of 2 males each	Bred at Uni. Montana
PWK/PhJ	*M. m. musculus*	*M. m. musculus* Wild-derived inbred	2 pools (PWK 2, 3) of 2 males each	Bred at Uni. Montana
STF	*M. spretus*	*M. spretus* Wild-derived inbred	2 pools (STF 1, 2) of 2 males each	Bred at Uni. Montana

Table 1: Mouse lines analysed for this study. CRL; Charles River Laboratories, Manston,
UK.

Sperm collection and fixation

The vasa deferentia and caudae epididymes were dissected from each animal, and the

contents squeezed out into 1ml PBS (scaled up accordingly if multiple animals were pooled).

The sperm were transferred to a microfuge tube, and tissue clumps were allowed to settle.

Sperm were transferred to a new tube and pelleted at 500g for 5mins. The supernatant was

removed, and the sperm fixed dropwise with either 3:1 methanol-acetic acid or 2%

paraformaldehyde (PFA) in PBS. Sperm were again pelleted at 500g for 5mins, and washed

in fixative twice more. Samples were stored at -20°C (methanol-acetic acid) or 4°C (PFA).

Imaging

Samples were diluted in fixative as required to obtain an evenly-spread preparation, and 8µl

of sample dropped onto a slide and allowed to air dry. Slides were counterstained with 16µl

VectorShield with DAPI (Vector Labs) under a 22x50mm cover slip and imaged at 100x on

an Olympus BX-61 epifluorescence microscope equipped with a Hamamatsu Orca-ER

C4742-80 cooled CCD camera and appropriate filters. Images were captured using Smart-

Capture 3 (Digital Scientific UK). To validate the reproducibility of the software, sample

images were also gathered on three other microscopes: (1) an Olympus BX61 with a

Hamamatsu C10600 orca r² camera, (2) an Olympus BX61 with a Hamamatsu Orca-03G

camera, and (3) a Nikon Microphot-SA epifluorescence microscope with a Photometrics

Metachrome II CH250 cooled CCD camera.

Nucleus detection and morphological analysis

Image analysis was performed using a custom program designed as a plugin for the freely

available image analysis program ImageJ [27]. The plugin, Nuclear Morphology Analysis.

The core software was developed using Java 8, with the user interface written using Swing.

The software is available at http://bitbucket.org/bmskinner/nuclear_morphology/wiki/Home/

together with full installation instructions, an online wiki user manual, and example images

for testing. The analyses described here were conducted using software version 1.13.6. The

program allows for (a) detection of objects within fluorescence images and (b) morphological

analysis of objects as sperm nuclei using a species-specific set of rules for identifying

biologically relevant structures. The detection strategy is outlined in the supplementary

methods.

Once nuclei were acquired from a set of images, they were consistently oriented and

aligned. We used a modification of the Zahn-Roskies (ZR) transform [28] to 'unroll' the

outline of each sperm nucleus by measuring the interior angle of the sperm at each point

around the nuclear perimeter to generate a linear trace we refer to as the angle profile

(Figure 1); further details and validation of the robustness of the method are given in the

supplementary data and Supplementary figures 1-11.

Figure 1: Shapes are detected by measuring the internal angles around the periphery of the
nucleus. Angles from key features in an example nucleus (A) are plotted in (B). The actual
profile for the entire perimeter is shown in (C).

Statistical analysis and clustering

Following segmentation, standard nuclear parameters were measured: area, perimeter and

aspect ratio, the width of the nuclear body versus the length of the hook as described in

other papers (e.g. [9], and the lengths of each perimeter segment. Data was exported for

further processing in R. Differences between datasets were tested using a pairwise Wilcoxon

rank sum test, with Bonferroni multiple testing correction. In order to quantify the variability of

the nuclear shapes, we developed a new per-nucleus measure defined as the root-mean-

square difference between the per-nucleus angle profile and the median angle profile for the

dataset, averaged across the length of the angle profile. The coefficient of variability

(standard deviation / mean) was also calculated for each of the other measured parameters.

The 'average shape' of the nuclei was calculated by averaging the x and y coordinates at

consistent semilandmarks taken as fractions of the perimeter across all nuclei, vertically

aligned and with their centres of mass at (0,0). This yielded a 'consensus nucleus'

visualising the overall shape of the population. Clustering was implemented via the WEKA

data mining software library [29].

Results

Detection and quantification of sperm shape in C57Bl6 and CBA mice

The difference between CBA and C57Bl6 sperm is distinguishable to the trained eye, and

makes a useful demonstration of the software's features. The angle profiles generated are

distinct for each genotype (Figure 2A). CBA sperm have a larger cross-sectional area, are

longer, and also have slightly shorter hooks than C57Bl6 sperm (Figure 2B/C). These

differences are reflected in the profiles; the long narrow tail in the CBAs appears as a

smooth curve at x=50 in the profile, while the shorter, wider C57Bl6s show a distinct dip

corresponding to the sharper curve of the dorsal angle before the acrosome. The shorter

hook of the CBAs is also seen as a narrow peak at x=10; the longer hook of the C57Bl6s

has a correspondingly wider peak. Automated segmentation of the nuclear profile allows

quantification and significance testing of the inter-line differences in each separate region of

the nuclear profile (see Supplementary figures 4, 5, 14).

CBA and C57Bl6 have previously been characterised by Wyrobek et al [9], who measured

160 nuclei of each genotype by manual tracing of projected microscope images of eosin-

stained sperm heads. We found our measured values to be similar (Supplementary Table 7)

but slightly smaller - as expected given that their measurements are for the entire sperm

head rather than just the nucleus. The body widths are within 0.3µm, and our bounding

heights are approximately 1.2µm smaller, consistent with our measurements lacking the

acrosomal cap (~0.15µm [30]), and the proteinaceous part of the sperm hook. We measured

the CBAs to be 12% longer than the C57Bl6s, again close to the previously published

13.5%.

Figure 2: A) Comparison of shape profiles between C57Bl6 (yellow) and CBA (blue), showing the median and interquartile range of the nuclear shape profiles. B) Consensus nuclei from each population, and the overlap showing the regions differing. C) Size and shape measurements between the lines. The prominent dorsal angle in C57Bl6 nuclei is marked with an asterisk.

Comparison of sperm morphology and variability across lines demonstrates the effects of

inbreeding depression and hybrid dysgenesis

With the software tested on CBA and C57Bl6, we wanted to investigate the extent to which

sperm shape variability within and between lines is affected by two factors: inbreeding

depression and the complex inter-subspecific mosaic origin of classical laboratory strains.

We selected a panel of inbred laboratory lines and compared them to (a) outbred laboratory

lines, and (b) wild-derived inbred lines (Table 1). Biological replicate samples from the inbred

lines represent either single animals (lab lines) or a pool of two animals (wild-derived inbred

lines). For the outbred lines, several individuals were pooled to ensure we were capturing

the diversity of the population as a whole.

Variability within each line was assessed using a new measure based on the similarity of

each cell's angle profile to the median for that line (see Methods). This was found to

correlate well with other population measures of variability such as the coefficients of

variation for area, bounding height and perimeter (Supplementary table 1). A comparison of

the overlaid average nuclear shape is shown in Figure 3. In addition to each line having a

characteristic sperm morphology, different lines showed different levels of intra-sample

variability. A breakdown by biological replicates shows that these data reflect true line

differences rather than biological differences between individual animals or technical

differences between imaging sessions or choice of fixative (Supplementary figures 9-11).

The BALB/c mice have the most variable shape profiles of all the lines we analysed, as well

as the highest coefficient of variability in area, height and width (Supplementary tables 1, 2).

The other inbred laboratory lines all showed low intra-line variability despite the fact that

there were marked differences in sperm size and shape between lines. Of the inbred

laboratory lines tested, CBA and DBA had the lowest intra-sample variability. The two

outbred lines, CD1 and MF1Y[RIII] both showed slightly higher intra-sample variability. This

may reflect the fact that these samples were pooled samples derived from multiple

genetically unique individuals. Turning to the wild-derived lines, all three lineages analysed

(*M. m. domesticus*, *M. m. musculus* and *M. spretus*) had lower variability than any of the

standard laboratory lines, despite that fact that these wild-derived lines are inbred.

Figure 3: Parameters for additional lines examined, with representative nuclei and population consensus. Samples are coloured according to their type: from left to right: inbred (yellow), outbred (white) and inbred wild-derived (blue).

Segmentation of sperm profiles allows detailed analysis of elements of sperm morphology

Once "unrolled" into an angle profile, this profile can then be segmented at local minima and maxima (see Methods) to identify specific landmarks within the sperm head shape. Some landmarks are consistently found across all lines, such as the tip of the apical hook and the point of maximum curvature at the base of the sperm head, while other landmarks such as the dorsal angle and the indentation at the tail attachment site are variable between lines. As an example of how this can be used to compare samples, since we had already found the presence of a dorsal angle to vary between CBA and C57Bl6 sperm, we examined how this varied across the full data set. Of all the lines studied, only five showed a clear dorsal angle, with the others having a smoother profile posterior to the acrosome. The distance from the rear reference point to the dorsal angle was characteristic for each of these five lines, as was the variability in this measurement, with BALB/c mice showing highest variability. Supplementary Figure 15 discusses the ubiquitous and variable landmarks discovered by the segmentation analysis and shows the detailed segmentation pattern for each line, while Supplementary Table 4 gives the numerical segment length data for each line.

C57Bl6 / CBA F1 cross males demonstrate the effects of each parental genotype on sperm shape and the relief of inbreeding depression by heterosis.

The differences we saw between inbred and outbred laboratory lines made us curious as to the impact of line background and genetic interactions thereof. We investigated one specific reciprocal F1 cross, between C57Bl6 and CBAs. The use of F1 animals is important here as it relieves the effects of inbreeding depression caused by fixation of deleterious recessive variants in each of the parental lines, but still yields a uniform population of genetically

identical males from each cross. B6CBA mice are the F1 offspring of a female B6 with a

male CBA and CBAB6 mice are the reciprocal cross. Sperm morphology for both F1 lines

matches the CBA parental line closely, indicating a dominant effect of the CBA genotype

(Figure 4A). In terms of sperm cross-sectional area, both types of F1 sperm are much more

similar to the CBA parent, while being fractionally larger than either parental line (Figure 4B).

Males from both directions of the F1 cross showed less variability in their sperm shape

compared to either parent line, suggestive of a degree of heterosis in the F1s.

The reciprocal cross data allows us to look for parent-of-origin effects on sperm shape. We

found two such differences, in sperm cross-sectional area and in bounding width. CBAB6s

have a slightly larger sperm area than the B6CBAs (19.3 square microns versus 18.6 square

microns, p<0.001) and the region around the posterior of the nucleus is widened in the

CBAB6s, intermediate to CBA and C57Bl6 (Figure 4B/C). The differences around the

posterior are largely driven by changes in the dorsal angle, which is present in C57, absent

in CBA, and virtually absent in both reciprocal F1 cross males (Figure 4D). For bounding

width, we find that this parameter is influenced by the male parent: CBAB6 and B6CBA are

significantly different to each other (p=0.0016), as are C57Bl6 and CBA (p=1.27E-12), but

there is no significant difference between C57Bl6 and CBAB6 (p=0.18) or between CBA and

B6CBA (p=0.095). This suggests that this aspect of sperm shape may be influenced either

by sex chromosome or mitochondrial background or by autosomal imprinted loci.

Figure 4: Subtle differences can be seen between a CBAB6 (CBA mother) and a B6CBA
(C57Bl6 mother). Both are intermediate to the parental shapes, but CBAB6 sperm are wider,
and their shape is closer to that of the C57Bl6. A) Consensus nuclei B) Size measurements;
C) Overlay of consensus nuclei; D) comparison of angle profiles; the tail attachment region is
expanded in the inset.

Hierarchical clustering can separate samples based on shape differences

Next, we turned our attention to the analysis of morphological variation within a given

population. In particular, we considered that cluster analysis of the sperm from a single

sample would give an unbiased breakdown of the different morphological sub-populations

contained therein. We used a hierarchical clusterer, as implemented by the WEKA data

mining tool [29] to separate sperm based on their shape profiles.

We tested the clustering algorithm by pooling images from C57Bl6 and CBA and analysing

them as a single sample. Since C57Bl6 and CBA sperm are slightly different sizes, the

simplest partitioning of the mixed set is a binary cut-off at a given threshold for nuclear area.

Passing the nuclear areas to a hierarchical clusterer and selecting the two most distinct

clusters using the Ward clustering method was 83-85**%** accurate at separating the individual

sperm by line. To determine whether shape-based hierarchical clustering could improve

upon this, we sampled values from the angle profile for each nucleus at regular intervals

(corresponding to the original window proportion) and provided these as inputs to the

clustering algorithm. This clustering was markedly more accurate than a simple size-based

cut-off, and separated the two genotypes with 91-95% success (Supplementary table 6). In

head-to-head tests using a representative subset of 50 nuclei from each genotype, the

clusterer performed at least as well (96%) as experienced assessors (97% accuracy), and

substantially better than novice assessors (75% accuracy) (Supplementary figure 12).

Hierarchical clustering can detect morphological subgroups within a sample

Having demonstrated that cluster analysis can recover different shapes from a mixed

population of known composition, we looked at its use for novel shape discovery within a

single highly variable population. Since the BALB/c line showed the highest variability in our

line survey, we chose this as our test sample. A cluster analysis based on angle profile alone

found four major groups of sperm shape, from mostly normal through to severe hyper-

condensation of the sperm (Figure 5). The final class is still highly variable compared to the

other classes; clustering these nuclei further reveals a separation of two separate types of

hypercondensation (Supplementary figure 16) as previously described [31]. While the most

normal sperm had near-normal placement of the dorsal angle, and a normal tail attachment

site, the most heavily distorted sperm showed frequent presence of additional sharp angles

in the sperm outline, effacement of the tail attachment site due to compression of the rear of

the sperm head, and an ever more prominent and misplaced dorsal angle that may reflect

altered microtubule dynamics during nuclear shaping (see Discussion).

*Figure 5: The overall population of BALB/c sperm appears distorted compared to other lines
(grey), but clustering reveals separate classes of morphology, from mostly normal (green) to
highly condensed (yellow).*

2
3
4
5

Discussion

We present here a morphological analysis tool designed to study nuclear morphology, with the ability to automatically identify key landmarks in the nuclear outline and quantitatively measure a range of nuclear and sub-nuclear parameters. Here, we demonstrate the use of this software to analyse the highly asymmetrical shape of the mouse sperm nucleus; however it is a generally applicable tool suitable for analysis of all sizes and shapes of nuclei. A companion paper ([32], submitted for publication) demonstrates its use in comparing sperm from boars judged to be suitable/unsuitable for use in artificial insemination.

Comparison of this method with other nuclear shape analysis methods

The key advantage offered by the software presented here is automation of the steps involved in object detection, shape decomposition and comparison. At the object detection stage, we use an edge detection algorithm that is markedly more effective than the fixed-threshold detection used in other packages, particularly in the presence of inhomogeneous staining of the bright chromocenter and dim apical hook. At the shape decomposition step, we introduce a modification of the Zahn-Roskies transform [28] that sensitively detects the various angular landmarks around the sperm periphery without the need for manual intervention. Together, these innovations massively increase the number of nuclei that can be quantified and compared to each other, with a total of 8,749 nuclei being measured during this study, and over 22,000 nuclei in the companion paper analysing boar sperm [32]. This for the first time permits the use of sample sizes that accurately capture not only fixed size and shape differences between samples, but also the detection and classification of intra-sample variability. Our method is robust to differences between camera and microscope setups and fixation techniques, making it amenable to analysis of large numbers of images, and potentially to automated image capturing from whole slide scanners.

While there are other features of sperm morphology that we do not yet address in this

package, the modular design of our software allows additional analysis pipelines to be added

at a subsequent date, and for features from different fluorescence channels to be associated

with specific nuclei and analysed in relation to them. We anticipate that other sperm

morphological features such as the extent and thickness of the acrosome, the proteinaceous

tip of the hook, the presence of cytoplasmic droplets, and the length and morphology of the

tail will be amenable to our approach by combining nuclear staining for orientation with

phase contrast imaging, tubulin immunostaining, MitoTracker, SpermBlue or other stains.

Since we are imaging fixed cells, the nuclei also remain available for interrogation by

chromosome painting or other molecular cytogenetic approaches, e.g. to detect aneuploid

cells and correlate their chromosomal status with their nuclear morphology.

Comparison of sperm shape within and between lines

Our observations support previous studies (e.g. [4,9]), add further information on the precise

regions of the sperm head that that differ between lines, and demonstrate the variability of

sperm morphology within each given line. In particular, we examined the presence and

placement of the dorsal angle of the sperm. This feature is created by pressure from the

manchette: a cone-shaped array of microtubules that forms behind the nucleus and slides

backwards during spermiogenesis, shaping the rear of the sperm head in the process.

Defects in katanin p80, a microtubule severing protein, lead to failure of this process and

abnormal compression of the base of the sperm head [6]. The narrowing of the tail

attachment site seen in FVB and BALB/c males, together with the prominent dorsal angle

seen in both lines (especially the latter) may indicate that manchette migration is abnormal in

these males.

Comparison of sperm variability within and between lines

The greatest variability we saw was in the BALB/c animals. This line is known to have poor

sperm morphology and high levels of sperm aneuploidy. Kishikawa et al [31] observed

different classes of abnormality, which we were able to recapitulate. In their analysis, the

authors found chromosomal abnormalities in 35% of highly abnormal sperm, but also in 15%

of sperm that were morphologically 'normal' by their criteria. Given that our new analysis

detects additional classes of more subtle shape difference that were not discriminated in the

earlier analysis, we hypothesise that these new abnormal classes may also be enriched for

chromosomal defects compared to the most normal sperm. Further differences await

characterisation: different classes and levels of sperm abnormalities have been described

depending on the particular subline and age of the animal [33].

Consistent with [34], we found that an F1 cross between C57Bl6 and CBA laboratory lines

lowered sperm shape variability (see below), suggestive of a degree of inbreeding

depression that was relieved by heterosis. However, the least variable lines we examined

were the wild-derived inbred lines PWK, LEW and STF, representing *M. m. musculus*, *M.m.*

*domesticu*s and *M. spretus* respectively. Since these three lines are also inbred, this

suggests that the wide variety of sperm shapes in laboratory lines, and the elevated level of

intra-individual variability in all the laboratory lines is not primarily a consequence of

inbreeding depression.

Instead, this is potentially linked to the status of the laboratory mouse as a hybrid between

several mouse subspecies - a factor that may have disrupted regulatory interactions

throughout the genome. Against this, PWK, despite being predominantly of *musculus* origin,

nevertheless has substantial introgression of domesticus DNA, of the order of ~6-7% of the

genome [35,36]. The degree of disruption may therefore depend on both the direction of

introgression and the specific regions involved. The recent finding of polymorphic hybrid incompatibilities within both *musculus* and *domesticus* subspecies shows that multiple regions of the genome contribute to hybrid breakdown and hybrid sterility. Consequently, the various different classical and wild-derived inbred lines may have fixed different combinations of incompatible alleles that collectively destabilise sperm development to varying extents in each line [37].

X/Y mismatch is a strong potential contributor to regulatory disruption, since most laboratory lines carry a *musculus* Y on a predominantly *domesticus* background [35]. The copy number of the ampliconic genes on the X and Y chromosomes varies markedly between *musculus* and *domesticus* subspecies, and the relative copy number of these genes is known to be important for normal sperm morphology [38–40]. However, while most of the laboratory lines we examined do indeed have mismatched X/Y chromosomes [35,41,42], the FVB X and Y are both of *domesticus* origin, indicating that the alterations in sperm shape in this line are not due to X/Y mismatch.

An alternative but not mutually exclusive explanation for the difference between classical laboratory inbred lines and wild-derived inbred lines is that the classical lines have been selected over multiple generations for their ability to breed well in captivity - indeed FVB is particularly known for its fecundity [43]. It may seem paradoxical that selection for high fecundity could adversely affect male fertility parameters: however, under laboratory conditions of non-competitive mating, co-housing a single male with one or more females, it is likely that reproductive output is driven largely by maternal factors. Thus, even though laboratory lines are fertile under lab breeding conditions, their sperm may be uncompetitive in mixed mating experiments compared to a pure species background. The morphology of the FVB zygote pronucleus is independent of the paternal genetic background, and the efficiency of FVB sperm for IVF appears unexceptional [44]. Sperm morphology and fertilisation success in laboratory mice has been shown to evolve rapidly in response to

competitive mating experiments, indicating that the baseline competitive ability of laboratory

line sperm is sub-optimal [45,46]. Intriguingly, it has even been shown that in lines

experimentally selected for high fecundity, male fertility and sperm morphology/motility

parameters are compromised, suggestive of a trade-off between the male and female factors

necessary for high fecundity in a laboratory environment [47].

Relevance for speciation, fertility, and toxicology studies

Abnormal sperm head morphology has emerged as a common form of hybrid male sterility in

mice [21–24,48]. Some sterility factors broadly impair spermatogenesis, resulting in reduced

sperm counts, lower motility, and abnormal morphology. However, several studies have now

shown that hybrid sterility QTL in mice often correspond to specific reproductive phenotypes

[24]. The challenges of manually quantifying morphology in large mapping panels has

necessitated the use of crude categorical scores [21,23,48], hampering quantitative

precision and likely limiting the ability to draw causal links between hybrid incompatibilities

and specific aspects of sperm morphological development.

Our approach assists in two ways: firstly by enabling more rigorous quantitation of sperm

shape, and secondly by enabling the large sample sizes and systematic approach needed

for mapping studies. As a proof of principle, we have compared males from a reciprocal

cross between C57Bl6 and CBA mice, and identified a dominant effect of the CBA genotype

on sperm shape. Within this, however, there are subtle differences between the CBAB6 and

B6CBA animals, suggesting an effect of either chromosome constitution or imprinting on

sperm bounding width. This demonstrates the usefulness of this approach for understanding

subtle features of mouse sperm nuclear development, and the potential to use this software

for genetic mapping of the various determinants of mouse sperm head shape.

Fertility rate and IVF efficiency has been correlated with the genetic background of sperm

among inbred mouse lines [49]. Furthermore, many studies have shown that the genetic

background of a line can influence sperm morphology. For example, deletion of the long arm

of the Y chromosome results in a more severe phenotype on B10.BR background than on

CBA [50]. Mashiko et al [16] have suggested morphology of sperm is associated with

fertilising efficiency in at least two mouse lines (B6D2F1 and C57Bl6/N). Since particular

genetic mutations in mouse sperm shape are associated with characteristic nuclear shape

abnormalities (e.g. [19]), detailed examination of sperm from natural mutant and/or targeted

knowckout animals may point to pathways of interest for understanding spermiogenesis and

male fertility more generally.

In toxicological analysis, rodent sperm are conventionally manually classified into classes of

predefined morphological abnormality (e.g. [26,51]). The hierarchical clustering implemented

within the software is able to separate nuclei based on shape as accurately as an

experienced manual sperm scorer; however it is much faster and more consistent. This may

be of use in samples where the nature and degree of abnormalities is hard for humans to

reliably quantify. It is also important to understand and quantify normal morphological

variation between lines since different lines can have different responses to toxicological

agents [52]. While many studies of toxicology using rodent models are conducted on rats,

the extra information available in the mouse sperm head still makes them a useful model

system. The fact that specific genetic lesions cause specific shape changes means that the

sperm shape might in principle give information not just about the presence/absence of

toxicity but also its mode of action. This level of analysis would complement existing studies

of sperm function, which, in clinical settings or in automated CASA platforms (e.g. [53]), is

still lacking detailed morphological data [10].

Conclusions

We present a new software package for the rapid, high-throughput, replicable analysis and comparison of nucleus shape in mouse sperm. By using a range of mouse lines, we have demonstrated the ability of the software to discriminate subtle differences between lines, and to reproducibly separate the nuclei into morphological groups. This has applications for studies of speciation, fertility and understanding the impact of genotoxic compounds. The analysis steps are generalisable and will work on many symmetric or asymmetric shapes of nuclei including, but not limited to sperm from other species.

Acknowledgements

We thank the animal handling staff at the University of Kent, University of Cambridge, University of Montana and Charles River Laboratories. We also thank the experimental test subjects who volunteered to classify C57Bl6 and CBA sperm.

Funding

BMS was supported by the Leverhulme Trust (grant RPG337) and the Biotechnology and Biological Sciences Research Council (BBSRC, grant BB/N000129/1). EEPJ was supported by BBSRC training grant BB/L502443/1. PE and CCR were supported by HEFCE (University of Kent) and by the BBSRC (grant BB/N000463/1). JMG and ELL were supported by the Eunice Kennedy Shriver National Institute of Child Health and Human Development of the National Institutes of Health (R01-HD073439 and R01-HD094787) and the National Institute of General Medical Sciences (R01-GM098536). EEKK was supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. (DGE-1313190).

Competing Interests

We have no competing interests.

Authors' contributions

Conceptualisation, BMS and PE; Methodology, BMS and PE; Software and Validation, BMS;

Investigation, CCR, JB, EEPJ and GY; Data Curation and Formal Analysis, BMS and CCR;

Visualisation, BMS; Supervision and Project Administration, PE; Writing - Original Draft,

BMS and PE; Writing - Review and Editing, BMS, CCR, JMG, ELL and PE; Resources,

JMG, ELL, EEKK, NA and PE; Funding Acquisition, NA and PE. All authors gave final

approval for publication.

References

1. Skinner BM, Johnson EEP. 2017 Nuclear morphologies: their diversity and functional
relevance. *Chromosoma* **126**, 195–212. (doi:10.1007/s00412-016-0614-5)

2. Patankar A, Parte P. 2017 Sperm Chromatin Compaction and Male Infertility. In *Male
Infertility: Understanding, Causes and Treatment*, pp. 295–315. Springer, Singapore.
(doi:10.1007/978-981-10-4017-7_17)

3. Meistrich ML. 1993 Nuclear Morphogenesis during Spermiogenesis. In *Molecular
Biology of the Male Reproductive System* (ed D de Kretser), pp. 67–97. San Diego,
USA: Academic Press.

4. Varea Sánchez M, Bastir M, Roldan ERS. 2013 Geometric Morphometrics of Rodent
Sperm Head Shape. *PLoS ONE* **8**, e80607. (doi:10.1371/journal.pone.0080607)

5. Medarde N, Muñoz-Muñoz F, López-Fuster MJ, Ventura J. 2013 Variational modularity
at the cell level: insights from the sperm head of the house mouse. *BMC Evolutionary
Biology* **13**, 179. (doi:10.1186/1471-2148-13-179)

6. O'Donnell L *et al.* 2012 An Essential Role for Katanin p80 and Microtubule Severing in
Male Gamete Production. *PLoS Genet* **8**, e1002698.
(doi:10.1371/journal.pgen.1002698)

7. Friend GF. 1936 Memoirs: The Sperms of the British Muridae. *Journal of Cell Science*

8. Cummins JM, Woodall PF. 1985 On mammalian sperm dimensions. *J. Reprod. Fertil.*
75, 153–175.

9. Wyrobek AJ, Meistrich ML, Furrer R, Bruce WR. 1976 Physical characteristics of mouse
sperm nuclei. *Biophys. J.* **16**, 811–825. (doi:10.1016/S0006-3495(76)85730-X)

10. Mortimer ST, van der Horst G, Mortimer D. 2015 The future of computer-aided sperm analysis. *Asian J Androl* **17**, 545–553. (doi:10.4103/1008-682X.154312)

11. Davis RO, Gravance CG. 1994 Consistency of Sperm Morphology Classification Methods. *Journal of Andrology* **15**, 83–91. (doi:10.1002/j.1939-4640.1994.tb01690.x)

12. Kuhl FP, Giardina CR. 1982 Elliptic Fourier features of a closed contour. *Computer graphics and image processing* **18**, 236–258.

13. Diaz G, Zuccarelli A, Pelligra I, Ghiani A. 1989 Elliptic Fourier analysis of cell and nuclear shapes. *Computers and Biomedical Research* **22**, 405–414. (doi:10.1016/0010-4809(89)90034-7)

14. Ostermeier GC, Sargeant GA, Yandell BS, Evenson DP, Parrish JJ. 2001 Relationship of Bull Fertility to Sperm Nuclear Shape. *Journal of Andrology* **22**, 595–603. (doi:10.1002/j.1939-4640.2001.tb02219.x)

15. Utsuno H, Miyamoto T, Oka K, Shiozawa T. 2014 Morphological alterations in protamine-deficient spermatozoa. *Hum. Reprod.* **29**, 2374–2381. (doi:10.1093/humrep/deu225)

16. Mashiko D, Ikawa M, Fujimoto K. 2017 Mouse spermatozoa with higher fertilization rates have thinner nuclei. *PeerJ* **5**, e3913. (doi:10.7717/peerj.3913)

17. Ostermeier GC, Sargeant GA, Yandell BS, Parrish JJ. 2001 Measurement of Bovine Sperm Nuclear Shape Using Fourier Harmonic Amplitudes. *Journal of Andrology* **22**, 584–594. (doi:10.1002/j.1939-4640.2001.tb02218.x)

18. Zelditch ML, Swiderski DL, Sheets HD. 2012 *Geometric Morphometrics for Biologists: A Primer*. Academic Press.

19. de Boer P., Vries M., Ramos L. 2014 A mutation study of sperm head shape and motility in the mouse: lessons for the clinic. *Andrology* **3**, 174–202. (doi:10.1111/andr.300)

20. Shorter JR *et al.* 2017 Male Infertility Is Responsible for Nearly Half of the Extinction Observed in the Mouse Collaborative Cross. *Genetics* **206**, 557–572. (doi:10.1534/genetics.116.199596)

21. Oka A, Mita A, Sakurai-Yamatani N, Yamamoto H, Takagi N, Takano-Shimizu T, Toshimori K, Moriwaki K, Shiroishi T. 2004 Hybrid breakdown caused by substitution of the X chromosome between two mouse subspecies. *Genetics* **166**, 913–924.

22. Campbell P, Nachman MW. 2014 X–Y Interactions Underlie Sperm Head Abnormality in Hybrid Male House Mice. *Genetics* **196**, 1231–1240. (doi:10.1534/genetics.114.161703)

23. Good JM, Dean MD, Nachman MW. 2008 A Complex Genetic Basis to X-Linked Hybrid Male Sterility Between Two Species of House Mice. *Genetics* **179**, 2213–2228. (doi:10.1534/genetics.107.085340)

24. White MA, Stubbings M, Dumont BL, Payseur BA. 2012 Genetics and Evolution of Hybrid Male Sterility in House Mice. *Genetics* , genetics.112.140251. (doi:10.1534/genetics.112.140251)

25. Bakare AA, Okunola AA, Adetunji OA, Jenmi HB. 2009 Genotoxicity assessment of a pharmaceutical effluent using four bioassays. *Genet Mol Biol* **32**, 373–381. (doi:10.1590/S1415-47572009000200026)

26. Das S, Upadhaya P, Giri S. 2016 Arsenic and smokeless tobacco induce genotoxicity, sperm abnormality as well as oxidative stress in mice in vivo. *Genes and Environment* **38**, 4. (doi:10.1186/s41021-016-0031-2)

27. Abramoff MD, Magelhaes PJ, Ram SJ. 2004 Image processing with ImageJ. *Biophotonics Int* **11**, 36–42.

28. Zahn, CT, Roskies, RZ. 1972 Fourier descriptors for plane closed curves. *IEEE Transactions, Computers* , 269–281.

29. Hall M, Frank E, Holmes G, Pfahringer B, Reutemann P, Witten IH. 2009 The WEKA Data Mining Software: An Update. *SIGKDD Explor. Newsl.* **11**, 10–18. (doi:10.1145/1656274.1656278)

30. Lin Y-W, Hsu T-H, Yen PH. 2013 Mouse sperm acquire a new structure on the apical hook during epididymal maturation. *Asian J Androl* **15**, 523–528. (doi:10.1038/aja.2013.46)

31. Kishikawa H, Tateno H, Yanagimachi R. 1999 Chromosome Analysis of BALB/c Mouse Spermatozoa with Normal and Abnormal Head Morphology. *Biol Reprod* **61**, 809–812. (doi:10.1095/biolreprod61.3.809)

32. Mandawala AA, Skinner BM, Walling GA, Fowler KE, Harvey S. 2018 Sperm morphology differences associated with pig fertility. *bioRxiv* , 314708. (doi:10.1101/314708)

33. Ohta H, Sakaide Y, Wakayama T. 2009 Age- and substrain-dependent sperm abnormalities in BALB/c mice and functional assessment of abnormal sperm by ICSI. *Hum Reprod* **24**, 775–781. (doi:10.1093/humrep/den456)

34. Wyrobek AJ, Gordon LA, Burkhart JG, Francis MW, Kapp RW, Letz G, Malling HV, Topham JC, Whorton MD. 1983 An evaluation of the mouse sperm morphology test and other sperm tests in nonhuman mammals: A report of the U.S. environmental protection agency Gene-Tox program. *Mutation Research/Reviews in Genetic Toxicology* **115**, 1–72. (doi:10.1016/0165-1110(83)90014-3)

35. Yang H *et al.* 2011 Subspecific origin and haplotype diversity in the laboratory mouse. *Nature Genetics* **43**, 648–655. (doi:10.1038/ng.847)

36. Sarver BAJ, Keeble S, Cosart T, Tucker PK, Dean MD, Good JM. 2017 Phylogenomic Insights into Mouse Evolution Using a Pseudoreference Approach. *Genome Biol Evol* **9**, 726–739. (doi:10.1093/gbe/evx034)

37. Larson EL *et al.* 2018 The Evolution of Polymorphic Hybrid Incompatibilities in House Mice. *Genetics* , genetics.300840.2018. (doi:10.1534/genetics.118.300840)

38. Conway SJ, Mahadevaiah SK, Darling SM, Capel B, Rattigan AM, Burgoyne PS. 1994 Y353/B: a candidate multiple-copy spermiogenesis gene on the mouse Y chromosome. *Mammalian Genome* **5**, 203–210. (doi:10.1007/BF00360546)

39. Cocquet J, Ellis PJI, Yamauchi Y, Mahadevaiah SK, Affara NA, Ward MA, Burgoyne PS. 2009 The Multicopy Gene Sly Represses the Sex Chromosomes in the Male Mouse Germline after Meiosis. *PLoS Biol* **7**, e1000244. (doi:10.1371/journal.pbio.1000244)

40. Cocquet J, Ellis PJI, Mahadevaiah SK, Affara NA, Vaiman D, Burgoyne PS. 2012 A Genetic Basis for a Postmeiotic X Versus Y Chromosome Intragenomic Conflict in the Mouse. *PLoS Genet* **8**, e1002900. (doi:10.1371/journal.pgen.1002900)

41. Gubbay J, Collignon J, Koopman P, Capel B, Economou A, Münsterberg A, Vivian N, Goodfellow P, Lovell-Badge R. 1990 A gene mapping to the sex-determining region of the mouse Y chromosome is a member of a novel family of embryonically expressed genes. *Nature* **346**, 245–250. (doi:10.1038/346245a0)

42. Mardon G, Mosher R, Disteche CM, Nishioka Y, McLaren A, Page DC. 1989 Duplication, deletion, and polymorphism in the sex-determining region of the mouse Y chromosome. *Science* **243**, 78–80. (doi:10.1126/science.2563173)

43. Silver LM. 1995 *Mouse Genetics: Concepts and Applications*. Oxford, New York: Oxford University Press.

44. Taketo M *et al.* 1991 FVB/N: an inbred mouse strain preferable for transgenic analyses. *PNAS* **88**, 2065–2069. (doi:10.1073/pnas.88.6.2065)

45. Firman RC, Simmons LW. 2008 Polyandry, sperm competition, and reproductive success in mice. *Behav Ecol* **19**, 695–702. (doi:10.1093/beheco/arm158)

46. Firman RC, Simmons LW. 2010 Experimental evolution of sperm quality via postcopulatory sexual selection in house mice. *Evolution* **64**, 1245–1256. (doi:10.1111/j.1558-5646.2009.00894.x)

47. Michaelis M, Sobczak A, Koczan D, Langhammer M, Reinsch N, Schoen J, Weitzel JM. 2017 Selection for female traits of high fertility affects male reproductive performance and alters the testicular transcriptional profile. *BMC Genomics* **18**. (doi:10.1186/s12864-017-4288-z)

48. White MA, Steffy B, Wiltshire T, Payseur BA. 2011 Genetic Dissection of a Key Reproductive Barrier Between Nascent Species of House Mice. *Genetics* **189**, 289–304. (doi:10.1534/genetics.111.129171)

49. Sztein JM, Farley JS, Mobraaten LE. 2000 In Vitro Fertilization with Cryopreserved Inbred Mouse Sperm. *Biology of Reproduction* **63**, 1774–1780. (doi:10.1095/biolreprod63.6.1774)

50. Styrna J, Kilarski W, Krzanowska H. 2003 Influence of the CBA genetic background on sperm morphology and fertilization efficiency in mice with a partial Y chromosome deletion. *Reproduction* **126**, 579–588. (doi:10.1530/rep.0.1260579)

51. Al-Sarar AS, Abobakr Y, Bayoumi AE, Hussein HI, Al-Ghothemi M. 2014 Reproductive toxicity and histopathological changes induced by lambda-cyhalothrin in male mice. *Environ. Toxicol* **29**, 750–762. (doi:10.1002/tox.21802)

52. Bone W, Walden CM, Fritsch M, Voigtmann U, Leifke E, Gottwald U, Boomkamp S, Platt FM, van der Spoel AC. 2007 The sensitivity of murine spermiogenesis to miglustat is a quantitative trait: a pharmacogenetic study. *Reprod Biol Endocrinol* **5**, 1. (doi:10.1186/1477-7827-5-1)

1   53. Butts IAE, Ward MAR, Litvak MK, Pitcher TE, Alavi SMH, Trippel EA, Rideout RM. 2011
2   Automated sperm head morphology analyzer for open-source software. *Theriogenology*
3   **76**, 1756-1761.e3. (doi:10.1016/j.theriogenology.2011.06.019)