Myosin-II activity generates a dynamic steady state with continuous actin turnover in a minimal actin cortex

Sonal, Kristina A. Ganzinger, Sven K. Vogel, Jonas Mücksch, Philipp Blumhardt and Petra Schwille*

Max Planck Institute of Biochemistry, 82152 Martinsried, Germany

*Correspondence: schwille@biochem.mpg.de

ABSTRACT

Dynamic reorganization of the actomyosin cytoskeleton allows a fine-tuning of cell shape

that is vital to many cellular functions. It is well established that myosin-II motors generate

the forces required for remodeling the cell surface by imparting contractility to actin

networks. An additional, less understood, role of myosin-II in cytoskeletal dynamics is

believed to be in the regulation of actin turnover; it has been proposed that myosin activity

increases actin turnover in various cellular contexts, presumably by contributing to

disassembly. *In vitro* reconstitution of actomyosin networks has confirmed the role of

myosin in actin network disassembly, but factors such as diffusional constraints and the use

of stabilized filaments have thus far limited the observation of myosin-assisted actin

turnover in these networks. Here, we present the reconstitution of a minimal dynamic actin

cortex where actin polymerization is catalyzed on the membrane in the presence of myosin-

II activity. We demonstrate that myosin activity leads to disassembly and redistribution in

this simplified cortex. Consequently, a new dynamic steady state emerges in which actin

filaments undergo constant turnover. Our findings suggest a multi-faceted role of myosin-II

in fast remodeling of the eukaryotic actin cortex.

INTRODUCTION

Rapid and controlled modulation of cell shape provides the foundation for various cellular

functions, such as division, growth, migration and adhesion. In eukaryotic cells, much of this

modulation can be attributed to the dynamics of the actomyosin cortex that lines the inner

surface of the plasma membrane. The cortex is a thin network of actin filaments associated

with myosin motors (see Salbreux et al., 2012 for a comprehensive review). It is further

reinforced by a collection of actin binding proteins that mediate filament-filament and

filament-membrane interactions. Additionally, myosin-II motors drive contractility of the

actin meshwork, generating forces at different length scales that enable fast remodeling of

the cell surface.

Another factor that imparts dynamicity to the cortex is the continuous turnover of all the

protein components of this network. Relative rates of turnover can determine the

mechanical properties of the cortex and are, therefore, subject to local regulation. For

instance, actin turnover has significant consequences on cortical tension (Tinevez et al.,

2009) and various network elements are implicated in imparting different turnover rates to

subpopulations of actin (Fritzsche et al., 2013). The rate of actin turnover even within the

same cell has been shown to vary between functionally distinct regions of the cortex

(Murthy and Wadsworth, 2005). Turnover rates of actin can be tuned by manipulation of

either assembly or disassembly processes. Lipids in the bilayer can orchestrate local

polymerization via activation of nucleators such as the Arp2/3 complex (Higgs and Pollard,

2000). Disassembly, on the other hand, is believed to be enhanced by myosin activity itself,

although dedicated severing proteins also exist (Blanchoin et al., 2000). Suppression of

myosin activity was implicated in reduced actin turnover in a range of cellular contexts

(Guha et al., 2005, Medeiros et al., 2006, Murthy and Wadsworth, 2005, Yang et al., 2012).

Due to the complexity of cellular environments, *in vitro* experiments have proved more

amenable for direct observation of myosin-II driven severing of actin filaments (Haviv et al.,

2008, Murrell and Gardel, 2012, Vogel et al., 2013) as well as network disassembly

(Reymann et al., 2012). While various other reconstitution studies have addressed the

mesoscopic effects of myosin contractility on actin network reorganization, a direct

investigation of the role of myosin in network turnover has been lacking. The observation of

filament turnover has previously been limited either by factors like stabilized actin

filaments, diffusion constraints, restricted polymerization (Koster et al., 2016, Linsmeier et

al., 2016, Murrell and Gardel, 2012, Smith et al., 2007, Stam et al., 2017, Vogel et al., 2017,

Vogel et al., 2013); or by a focus on network-level changes rather than individual network

components (Carvalho et al., 2013, Soares e Silva et al., 2011, Kohler et al., 2011, Bussonnier

et al., 2014).

By using a biomimetic actin cortex with minimal restraints on the mobility of components

and coexistence of assembly and disassembly, we demonstrate that myosin-II activity can

generate turnover in membrane-associated dendritic actin networks. This observation was

enabled by the following features incorporated during reconstitution: 1) polymerization

driven by catalysts localized to the membrane and 2) no additional anchoring of either the

myosin or the actin. While sharing qualitative attributes of the contraction process with

previous studies, our minimal cortex additionally showed breakdown of actin networks on

the membrane, redistribution of network components and finally, recruitment of the

released components into new assembly-- altogether, the essential aspects of filament

turnover.

RESULTS

Assembly of a dynamic actomyosin cortex on supported lipid bilayers

For the reconstitution of a biomimetic minimal cortex with the potential for actin network

turnover, we required a strong catalyst of actin polymerization to oppose myosin-driven

disassembly. The constitutively active version of the murine N-Wasp, also known as "VCA",

served as the catalyst. This protein was anchored to a supported lipid bilayer (SLB) with the

stable association of a 10x Histidine tag to nickelated lipids (Nye and Groves, 2008). Not only

does the membrane localization of the catalyst emulate the cellular scenario, but it also

allows us to preferentially image the polymerization process, by using Total Internal

Reflection Fluorescence (TIRF) microscopy. VCA has been extensively studied *in vitro* and

has been shown to promote the autocatalytic growth of a branched actin network, when

combined with the Arp2/3 complex (Pantaloni et al., 2000, Rohatgi et al., 1999). Such

"dendritic" network assembly is fast and, therefore, a suitable choice for counterbalancing

disassembly by myosin activity. Muscle myosin-II motors were preassembled into filaments

and introduced into the system after actin assembly (Fig. 1A).

We started with the characterization of the actin assembly process in the absence of myosin

motors. As expected, after a short time lag, surface actin intensity increased fast on the

VCA-functionalized SLBs in the presence of Arp2/3 and actin monomers (Fig. 1B, Movie 1).

The observed time profiles are characteristic of the autocatalytic nature of dendritic actin

network assembly (Pantaloni et al., 2000). Actin network growth was dense and resulted in

bundle-like organization of filaments with time (Fig. 1C). In comparison, actin assembly was

significantly weaker when Arp2/3 was excluded from the reaction and nearly non-existent in

the absence of VCA (Fig. 1B). While VCA alone was sufficient to induce some actin assembly,

the resulting meshwork was coarser than in the presence of Arp2/3 (Fig. S1). This Arp2/3-

independent assembly can be attributed to the ability of VCA to enhance actin filament

elongation on surfaces by increasing the local availability of monomers (Bieling et al., 2018).

In contrast, Arp2/3 alone failed to initiate any actin polymerization on the membrane (Fig.

1B, Fig. S1), since VCA not only activates the Arp2/3 complex, but is also necessary to

mediate actin-membrane interactions in this minimal cortex. Consequently, adding VCA to

this sample immediately leads to increased actin intensity on the surface and initiation of

filament assembly (Fig. S1). Altogether, these trends confirm that the actin assembly we

observe on the membrane is VCA-dependent, with considerable contribution from Arp2/3-

induced branching.

Interestingly, the diffusional mobility of the VCA on the membrane appeared to be essential

for the observed actin network growth, as holes in the lipid bilayer were largely devoid of

actin, even though VCA accumulated there (Fig. S2). We also confirmed that VCA intensity

on the membrane remained stable during actin assembly (Fig. S3, Movie 1). Actin was grown

for longer periods to bring the growth process close to saturation (Fig. S4) before adding

myofilaments.

Introducing myosin-II to the dendritic actin network resulted in condensation of actin into

contraction foci. Interestingly, two distinct length scales of actomyosin contraction were

observed under our experimental conditions (Fig. 1D, Movie 2). In some samples, the

contraction foci were distributed homogeneously with shorter distances between them

(9.1±2.2 μm) and actin enrichment in these foci was less pronounced. In other cases, we

observed a distinct coarsening process with actin condensing into dense foci, separated by

larger length scales (20.3±7.4 μm). The characteristic length scales of the two contraction

modes were calculated as nearest neighbor distance of the contraction foci (Fig. 1E). The

observation of multistage coarsening is in line with previous studies on disordered actin

networks, both membrane-associated and in solution (Backouche et al., 2006, Kohler et al.,

2011, Linsmeier et al., 2016, Murrell and Gardel, 2012, Smith et al., 2007, Soares e Silva et

al., 2011, Stam et al., 2017, Vogel et al., 2013).

The bimodal outcome of contraction led us to investigate the factors that impact length

scale of contraction in membrane-associated dendritic actin networks. We found that this

length scale responded to manipulations in surface density of VCA, Arp2/3 concentration,

myosin concentration and branch capping (Fig. S5). A five-fold increase in VCA surface

density, leading to a net increase in membrane attachment of the actin network, reduced

the length scale by one-third (Fig. S5A). Further, enhanced branching resulting from

increasing Arp2/3 concentrations trended towards lower length scale (Fig. S5B). Lower

myosin concentrations and the introduction of branch capping, presumably resulting in

shorter filament lengths, also favored lower length scales (Fig. S5C-D). These dependencies

corroborate a previous *in vitro* study, where membrane adhesion and filament cross-linking

were shown to affect contraction length scale in a disordered actin network (Murrell and

Gardel, 2012).

Myosin-II breaks down membrane-associated actin networks

Irrespective of the length scale of contraction, we found that myosin activity led to

disassembly of the actin cortex (Fig. 2). The initial stages of myosin contraction showed

bundling of actin filaments, followed by a compaction of the actin network (Fig. 2A a- white

arrows, Movie 3). As the actin condensed into contraction foci, rips in the actin network

could be observed. Finally, this process resulted in a decrease in actin intensity on the

surface, indicating network breakdown. Myofilaments were associated with the bundles as

well as the condensed actin foci (Fig. 2A b, yellow arrows). They, however, accumulated on

the membrane even as the actin intensity dropped with disassembly (Fig. 2B).

The length scale of the rips in the actin network determined whether the mode of

contraction was homogeneous or coarse, with larger rips observed in the latter case. In both

cases, actin intensity on the membrane reduced by 50% within a time range of 5±1.7

minutes after addition of myosin (Fig. S6). The contraction events were myosin-specific and

required a continued supply of ATP, as the process can be stalled at the bundling stage in

the absence of ATP regeneration (Movie 4). Since myosin continues to bind actin in the

absence of ATP, a catastrophic disintegration of the bundles was observed once new ATP

was made available by the addition of an ATP regeneration mix.

The dense actin condensates formed during coarsening allowed a distinct observation of

actin filament breakdown upon contraction. Representative images and kymograph of one

of these foci illustrate the disassembly of actin filaments in the core after condensation (Fig.

2C a, b). Following the individual radial profiles of actin and myosin intensities in these foci

at different time points, we observed that the condensation process was followed by a

hollowing of the foci (Fig. 2D). The actin intensity at the core decreased drastically, while the

myosin intensity showed a comparatively minor reduction (Fig. 2C b, 2D). This indicates that

as contraction continues in the condensed actin foci, the actin filaments disintegrate rapidly

under the stresses generated by the enriched myosin motors. A similar hollowing of actin

condensates with myosin activity was described in solution experiments (Soares e Silva et

al., 2011).

Individual events of filament or bundle fragmentation were difficult to observe in the dense

network. However, as actin condensed into coarser foci, bundles of actin could be observed

distinctly linking the foci. As contraction proceeded these bundles ruptured abruptly and

recoiled, revealing the tensile stresses acting on them (Fig. S7- left panel, Kymograph).

Myofilaments could be seen as distinct puncta along the bundles, presumably cross-linking

the actin filaments (Fig. S7- lower right panel). This explicitly shows one mode of filament

breakage occurring during contraction in the dendritic network.

169   Redistribution in active networks results in a dynamic steady state of actin
170   distribution

The stark contrast in actin distribution generated during coarsening also enabled us to

observe phenomena which were not discernable during homogeneous contraction.

Although coarsening initially generated a very heterogeneous actin distribution on the

membrane, eventually the actin network resumed a seemingly uniform organization (Movie

5). In Figure 3A, we use the Coefficient of Variation (CV) of the intensity distribution of actin

as an indicator of the heterogeneity in its physical distribution on the membrane. CV was

calculated as the ratio of standard deviation to mean of the actin intensity distribution at

each time point. The increase in CV coincided with the contraction process upon addition of

myofilaments, peaking at the height of coarsening (Fig. 3A a-c). The CV then dropped as the

actin was redistributed and finally results in an actin distribution nearly as homogeneous as

before the addition of myosin (Fig. 3A d-f, graph). This observation suggests that large-scale

heterogeneity in actin distribution is unstable in this dynamic minimal cortex. Under the

combined influence of myosin-driven disassembly and actin polymerization, reorganization

continues till a more homogeneous actin distribution is achieved. This distribution was

similar to that observed when contraction proceeded homogeneously (Fig. 3A- graph).

The restored uniformity in actin distribution can be attributed to two phenomena: the loss

of actin from contraction foci and the recovery of actin filaments on the remaining regions

of the membrane. These two processes occurred simultaneously, as can be illustrated by

plotting the opposing trends in actin intensity in these two regions of the membrane (Fig.

S8). The loss of actin resulted from breakdown of condensed actin in the foci as described in

the previous result (Fig. 2D) and also observable in Figure 3A (c-d). The concomitant

reappearance of actin in other regions was accompanied by formation of new myosin foci in

these parts (Fig. 3B a, white arrows, Movie 6). A kymograph of a stretch between two

contraction foci revealed that both actin and myosin contribute to the newly emerging node

(Fig. 3B b-c, data representative of 32 such nodes). Myosin intensity, however, increased

with a slight delay after actin intensity in the new foci (Fig. 3B c, inset), suggesting that actin

drives the reorganization. This delay was variable between nodes, ranging from 5 to 40

seconds. The source of the material for the new foci cannot be distinctly discerned, but both

actin and myosin were exchanged with the neighboring foci (Movie 6, Fig. 3B b, white

arrows). Myosin accumulated in the new foci with time, whereas actin intensity remained

constant (Fig. 3B b, c). These observations indicate that myosin follows actin filaments and

the redistribution of the former is limited as compared to the latter.

As can be expected, this reorganization resulted in a reduction in the distance between foci

we previously measured as contraction length scale during coarsening (Fig. 3C a, b). The

final values approached the length scale of homogeneous contraction (Fig. 3C), indicating

that this actomyosin network has a favored length scale that is restored in spite of initial

disturbances. The resultant actin distribution resembles a dynamic steady state with

filament exchange between the foci observed even after an hour since addition of myosin

(Movie 7, Fig. S9).

Unlike myosin, VCA localization on the membrane was stable during contraction (Fig. S10A)

and unaffected by coarsening (Fig. S10B), suggesting that VCA-driven actin polymerization

could contribute to reappearance of actin during redistribution. Interestingly, VCA also

enriched in the late-stage actomyosin foci formed on the membrane, possibly resulting in

enhanced polymerization at these sites (Fig. S10B). However, while redistribution of actin

resulted in local increase in actin intensity, mean actin intensity on the entire membrane did

not increase after the initial reduction during contraction (Fig. S11). Thus, a considerable

proportion of the actin network is permanently lost from the membrane, coinciding with

appearance of condensates in solution (Fig. S12A), reminiscent of previous solution studies

(Soares e Silva et al., 2011). The shape dynamics of these condensates indicate that myosin

is active in them (Fig. S12B), possibly trapping the actin filaments by cross-linking and

recapture.

New assembly of actin filaments reveals myosin-induced actin turnover in the minimal cortex

New assembly of actin filaments reveals myosin-induced actin turnover in

the minimal cortex

To investigate the role of myosin-driven fragmentation in actin filament turnover, we asked

whether the network components released during myosin contraction can be reused for

new polymerization. We reasoned that if excess monomers were washed off from the actin

network before addition of myofilaments, any polymerization observed after myosin activity

would largely be derived from the filament fragments and monomers generated by myosin-

mediated breakdown of actin. To isolate the actin growth process, myosin-II activity was

blocked using the inhibitor blebbistatin after the initial fragmentation and redistribution of

actin had taken place. We found that the surface intensity of actin gradually increased after

the inactivation of myosin (Fig. S13), suggesting that myosin breaks down the actin network

to the level of monomers, which are then competent for new polymerization. In accordance

with our hypothesis, novel polymerization was only observed if myosin had broken down

the actin network, since no increase in actin intensity was observed if blebbistatin was

added before actin network breakdown (Fig. S13). This indicates that breakdown by myosin

facilitates new actin assembly by releasing monomers after network growth has saturated.

To follow the formation of new filaments in presence of myosin activity in greater detail, we

designed a pulse-chase type experiment with two differently labeled actin monomers (Fig.

4A). The actin network was assembled as usual with 10% Alexa568-labeled actin (Label1).

Excess monomers in solution were then removed by extensive washing. Along with

myofilaments, additional actin monomers (1 µM) were added to the system, 10% of which

were labeled with Alexa488 (Label2). Additional Arp2/3 complex was also provided so that

the process of reassembly is not limited by its availability. We then observed whether the

newly assembled actin on the membrane contained both labels.

We found that the proportion of Label2 in the membrane-associated actin increased over

time after the initial breakdown of the Label1-network (Fig. 4B). This continuous increase

confirmed that the actin filaments observed on the membrane were undergoing constant

turnover in the presence of myosin activity. The steady increase in Label2 intensity on the

membrane was accompanied by a very slight increase in Label1 intensity. The difference in

the rates of change was probably due to the increase in total amount of actin on the

membrane in this period and the total amount of Label2-actin in the system being higher.

In the intermediate stages of label mixing, filaments with uniform distribution of Label1

could be observed with speckled occurrence of Label2 (Fig. 4C). Since the actin on the

membrane appeared bundled in the presence of myosin, individual filaments could not be

distinguished. However, single pixel line profiles along two distinct bundles showed that the

intensities of the two labels appear to be anti-correlated (Fig. 4D). Such filaments of mixed,

interspersed labeling would only be observed if monomers of both Label1 and Label2 were

available for polymerization, i.e. only if Label1 monomers were released from the pre-

assembled network upon myosin contraction. Altogether, these observations suggest that

the dynamic steady state actin distribution that emerges in this minimal cortex is comprised

of actin filaments undergoing continuous turnover under the combined influence of new

assembly of actin and myosin activity.

DISCUSSION

In this study, we have reconstituted a minimal dynamic actin cortex to elucidate myosin-

assisted turnover of actin filaments. We employed a TIRF microscopy approach to study the

effects of myosin-II contraction on the reorganization of actively polymerizing branched

actin networks on membranes. The key distinguishing feature of the minimal actin cortex

described in this study is the coexistence of catalytic actin polymerization with the

contractile activity of myosin. This enabled us to demonstrate myosin-II-stimulated actin

turnover by observing the three essential processes of: 1) disassembly of pre-existing actin

network, 2) redistribution of released network components, and 3) re-use of these

components for new network assembly (Fig. 5A).

Altogether, our observations of myosin-driven reorganization of branched actin networks on

membrane are well in line with previous studies, which also describe a distinct network

coarsening upon myosin addition both in solution (Backouche et al., 2006, Kohler et al.,

2011, Soares e Silva et al., 2011) or on membranes (Koster et al., 2016, Linsmeier et al.,

2016, Murrell and Gardel, 2012, Smith et al., 2007, Stam et al., 2017, Vogel et al., 2013).

However, beyond this condensation process, we further observed the subsequent de novo

polymerization and redistribution of actin on the membrane (Fig. 5B), with only one of the

solution studies previously reporting a similar phenomenon of an emergent dynamic steady

state (Kohler et al., 2011). This finding highlights the dynamic nature of the minimal

actomyosin cortex that we have reconstituted. In particular, the significance of including

polymerization catalyzed by VCA is reinforced by the observation of less dynamic

condensates in solution, as VCA is membrane-bound and not present in any significant

amount in solution.

On a lipid bilayer, fragmentation of individual actin filaments by myofilaments, in isolation

or within a dense network, has been demonstrated (Murrell and Gardel, 2012, Vogel et al.,

2013). However, the breakdown and removal of membrane-bound actin after initial

compaction was likely impaired by strong membrane adhesion of the biotinylated actin

filaments (Vogel et al., 2013) or the use of crowding agents in solution (Murrell and Gardel,

2012). In contrast, in the minimal cortex described here, only the nucleation promoting

factor (VCA) is firmly bound to the nickelated lipids in the membrane. The membrane

association of the actin, resulting from the cumulative effect of multivalent interactions

between the actin filaments and the VCA, is persistent but dynamic. This set-up is a

reasonable approximation of cellular environments, where proteins linking actin networks

to membrane undergo fast turnover (Fritzsche et al., 2014), effectively enabling a highly

dynamic association by an alternate mechanism. Such reversible filament anchoring may be

crucial for the myosin-driven disassembly of dendritic actin networks that we observe, as

similar processes were reported when actin networks were grown from a related nucleation

promoter, adhered to a glass surface (Reymann et al., 2012). Notably, we observed actin

disassembly in the form of a distinct hollowing of actin condensates on the membrane,

previously reported only in solution actomyosin aggregates (Soares e Silva et al., 2011). The

solution study speculated that myosin coalesces in the centre of contraction foci when

further translocation of myofilaments along the actin filaments is restricted by collision with

other myofilaments. The polarized motion of myosin motors coupled with the capacity of

actin filaments to reorient would normally result in the formation of a myosin-rich core with

actin filaments extending outwards from it. The fragmentation of actin filaments under

myosin-generated stress further allows a considerable fraction of the actin to disperse even

if actin fragments remain associated with the myosin in the core, leading to an "actin shell"

organization. However, in contrast with those observed in solution, these shell-like

structures are transient in our minimal cortex, presumably because the polymerization

machinery enables further redistribution of actin.

When coarsening was observed during contraction in our minimal cortex, some regions of

the membrane were left largely devoid of actin. The resulting contrast in actin distribution

enabled direct observation of the redistribution of network components on the membrane.

Synergistic relocation of actin and myosin, facilitated by actin polymerization, led to a more

homogeneous distribution of contraction foci. The active network then continued to show

long-term flows of actin filaments over this preferred length scale, resulting in the

emergence of a dynamic steady state of actin distribution, previously described only in

solution clusters (Kohler et al., 2011). Such fluidization of membrane-associated actin

networks by motor activity may aid the fast shape dynamics of the eukaryotic actin cortex.

The transition of length scale of actomyosin dynamics that results from redistribution

provides a compelling avenue for further investigation. While we offer some preliminary

analyses of the factors that affect the length scale of actomyosin contraction, further study

is needed to determine why our cortical active network shows bimodal length scales. We

observed a dependence on net membrane association of the actin network, branching

densities and possibly filament length and active motor concentration, consistent with

previous observations during contraction of disordered actin networks on membrane

(Murrell and Gardel, 2012). These are all factors that are subject to extensive biochemical

regulation in cell cortices and could, therefore, contribute to the coordination of actomyosin

activity over different length scales in a cell.

Finally, we elucidate the contribution of myosin activity in actin filament turnover in this

reconstituted cortex with the direct observation of the phenomenon. The notion of myosin

enhancing actin turnover rates has been suggested in various cellular contexts (Guha et al.,

2005, Medeiros et al., 2006, Murthy and Wadsworth, 2005, Yang et al., 2012, Wilson et al.,

2010), but the inhibition of myosin activity is a fairly complex perturbation due to the many

roles of myosin-II motors in the cytoskeleton. By disassembling actin networks, myosin can

amplify local cortical remodeling by providing fast access to a pool of network components

for new assembly. Since actin organization is to be coordinated over micron-scale distances

in the cell from a limited pool of components, this suggests an interesting mechanism by

which myosin activity could support faster dynamics on a shorter scale. Furthermore, the

negative feedback on contractility due to disassembly could form the basis of the pulsatory

dynamics observed during certain developmental processes (Nishikawa et al., 2017, Martin

et al., 2009).

In this simplified cortex, we selected only the minimal components required for dendritic

network assembly and contraction to gain a clearer understanding of the role of each

participating component. Altogether, this study demonstrates that myosin activity, when

counterbalanced by actin polymerization, can induce actin turnover in a minimal cortex and

result in the formation of an active network in dynamic steady state on the membrane. An

interesting avenue for more detailed investigation is to further dissect the actomyosin

organization in the emergent distribution and determine the influence of other structural

components of cortical networks. A detailed comparison with non-muscle myosin-II in a

similar set-up will also enhance our understanding of a cellular setting. Importantly, our

experiments suggest that the reconstituted dynamic actin cortex described in this study is

readily amenable to increased complexity and controlled manipulations. It, therefore, holds

great potential for providing further insights into the functioning of its biological inspiration.

MATERIALS AND METHODS

Proteins and other reagents

VCA domain (amino acids 400-501) of murine N-Wasp (Uniprot ID Q91YD9) was cloned into a modified pCoofy vector with 10x Histidine tag. The protein was then purified by a combination of affinity and anion exchange chromatography. In brief, the protein was expressed in a 1 L culture of Rosetta T1 strain of BL21 cells by inducing with 0.5 mM IPTG and growing the cells for 16 hours at 16°C. For purification, cells were lysed by sonication. The cell lysate was purified with the His tag on a HisTrap HP column (GE Healthcare, Chicago, IL, USA) via a chromatography system. The protein containing fractions were combined and diluted 10 times for ion exchange chromatography on a MonoQ column (GE Healthcare, Chicago, IL, USA). The selected protein fractions were buffer exchanged into phosphate buffer saline supplemented with 10% Glycerol. All buffers included 1 mM TCEP to prevent protein dimerization.

VCA was labeled with Atto488-maleimide (28562) from Sigma-Aldrich Corporation, St. Louis, MI, USA. The dye was used in 20-fold excess of the protein (by mass) and the mix was incubated overnight in a cold room with gentle rotation. Buffer exchange was done on a small volume Sephadex G-25 column. This was followed by dialysis for complete removal of the dye. The final storage buffer of the protein was 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10% Glycerol and 1 mM TCEP. Labeling efficiency was quantified by spectrophotometric analysis.

Purification of myosin-II purification from rabbit muscle, as well as labeling with Alexa Fluor 488, was carried out as described previously (Vogel et al., 2013).

Unlabeled actin monomers from rabbit skeletal muscle (AKL99) and Arp2/3 complex from porcine brain (RP01P) were purchased from Cytoskeleton, Inc., Denver, CO, USA. Actin monomers conjugated with Alexa Fluor 568 (A12374) and Alexa Fluor 488 (A12373) were acquired from ThermoFischer Scientific, Waltham, MA, USA. Capping protein CapZ (non-muscle, human recombinant) was purchased from HYPERMOL EK, Germany.

Blebbistatin (B0560), Phosphocreatinine (P7936), Creatine Phosphokinase (C3755), Pyranose

oxidase (P4234) and Catalase (C40) were purchased from Sigma-Aldrich Corporation, St.

Louis, MI, USA.

Preparation of lipid bilayer

DOPC and DGS-NTA lipids were purchased from Avanti Polar Lipids, Inc., Alabaster, AL, USA.

Small unilamellar vesicles with desired proportion of DOPC:DGS-NTA lipids (with 0.005%

Atto655-DOPE) were prepared by rehydrating a lipid film in SLB buffer (50 mM Tris-HCl pH

7.5, 150 mM KCl) followed by sonication. The composition was 99 mol% DOPC and 1 mol%

DGS-NTA in all experiments, unless otherwise mentioned. High precision coverglass (no. 1.5,

Marienfeld-Superior, Lauda-Königshofen, Germany) was treated with Piranha solution (3:1

H_2SO_4:H_2O_2, overnight), rinsed with copious amounts of milliQ water, dried and

subsequently briefly treated with O_2 plasma (Femto plasma cleaner, Diener-plasma,

Ebhausen, Germany). Reaction chambers were generated on these coverglasses by press-to-

seal silicone isolators (GBL664206; Grace Bio-labs, Bend, OR, USA). 0.1 mg/ml of vesicles

were deposited in the silicone chambers with frequent pipetting, followed by 15 minutes of

incubation. The bilayers were gently washed first with SLB buffer and then with VCA buffer

(10 mM Tris-HCl pH 7.5, 50 mM NaCl, 1 mM DTT) and stored in this buffer till sample

preparation.

Sample preparation

A lipid bilayer was flushed again with VCA buffer and then incubated with 200 nM 10xHis-

VCA in the same buffer. For experiments where labeled VCA was required, Atto488-labeled

VCA was included in a 1 in 10 ratio with unlabeled VCA. A premix containing actin

monomers (10 % Alexa568-labeled), Arp2/3 complex and ATP regeneration system

(Phosphocreatine and Creatine Phosphokinase) was made at the same time in G-buffer (10

mM Tris-HCl pH 8.0, 0.2 mM $CaCl_2$, 0.2 mM ATP, 1 mM DTT). After 20 minutes, the excess

VCA was washed off and actin pre-mix was added. An image was taken for the reference

intensity value for normalization before triggering polymerization by adding the 5x G-to-F

buffer (50 mM Tris-HCl pH 7.5, 250 mM KCl, 10 mM $MgCl_2$, 1 mM DTT with variable ATP). A

typical sample finally contained 1 µM actin monomers and 10 nM Arp2/3, unless otherwise

mentioned in the figures. Additionally, the sample also included the ATP regeneration

system (20 mM Phosphocreatinine, 53 U/ml Creatine Phosphokinase), and an oxygen

scavenger system (37 U/ml Pyranose Oxidase, 90 U/ml Catalase and 0.8 % Glucose). The

final buffer conditions were 10 mM Tris-HCl pH 7.5, 50 mM KCl, 1mM $MgCl_2$, 1mM DTT with

ATP adjusted to 0.1 mM. The sample volumes were 40 ml.

Myofilaments were pre-assembled by making a 4x stock of the myosin in a buffer of 10 mM

Tris pH 7.5, 50 mM KCl, 2 mM $MgCl_2$, 1 mM DTT and incubating for 10 minutes before

adding to the actin network by replacing ¼ of the sample volume. Where washing was

required before addition of myosin, a 4-fold replacement of sample volume was carried out

with gentle mixing.

In all cases, evaporation was minimized by storing the samples in closed boxes with wet

tissue till they were used. For putting on the microscope, a hydrophobic pen was used to

draw a "moat" around the silicone well and this was filled with water. A small plastic lid was

used to cover the sample, including the moat, during imaging.

Variations from this basic protocol have been mentioned in the text, where relevant.

Blebbistatin was dissolved in DMSO and used at a final concentration of 20 µM.

For experiments in which a second label of actin was needed, Alexa488-labeled actin

monomers were used. For pulse-chase experiments in Figure 4, Alexa488-labeled

monomers were mixed with unlabeled actin monomers at a 1 in 10 ratio. 1 µM of this 10 %

labeled actin was added to the sample, along with 10 nM Arp2/3, as described in the text.

Image acquisition

Fluorescence images were recorded on a home-built objective-type TIRF microscope

(Mucksch et al., 2018), constructed around a Nikon Ti-S microscope body with oil immersion

objective (Nikon SR Apo TIRF, 100x, NA 1.49, Nikon GmbH, Düsseldorf, Germany). In brief,

the excitation laser lines (490 nm (Cobolt Calypso, 50 mW nominal), 561 nm (Cobolt Jive, 50

mW nominal) and 640 nm (Cobolt 06-MLD, 140 mW nominal)) were attenuated with an

acusto-optical tunable filter (Gooch & Housego TF-525-250) and spatially filtered by a

polarization-maintaining single-mode fiber (kineFLEX-P-3-S-405.640-0.7-FCS-P0 and

kineMATIX, Qioptiq, Hamble, UK). After the fiber, the excitation beam was collimated,

extended 3-fold by a telescope of achromatic doublets and moved off the optical axis by a

mirror and a focusing lens, both on a piezo-controlled translation stage. This way, the

excitation beam was focused off-axis on the back focal plane of the objective. Excitation and

emission light were separated using a four-color notch beam splitter (zt405/488/561/640rpc

flat, AHF Analysentechnik, Tübingen, Germany). For single-color detection, the fluorescence

light was going through a 4f lens system, including spectral filtering (525/50, 593/46,

705/100) and detected on an electron-multiplying charge-coupled device (EMCCD) camera

(iXon Ultra 897, Andor Technologies, Belfast, UK). For dual-color schemes, the image was

laterally clipped in a conjugated focal plane. The clipped images were spectrally separated

(T555LPXR, Chroma), bandpass filtered (525/50, 593/46) and focused side-by-side on the

same EMCCD. To minimize phototoxicity and –bleaching, the AOTF transmission was

synchronized with the EMCCD acquisition using a controller card (PCle-6323, National

Instruments) with a customized LabView software (National Instruments, Austin, USA).

Imaging interval was either 2 or 5 seconds, with exposure times of 50 or 100 msec. A

custom-built focus stabilization eliminated drift of the focus position, based on a feedback

loop, which monitored the position of an infrared laser beam, back-reflected from the

sample interface.

For imaging in solution, as shown in Figure S12, a Yokogawa scan head CSU10-X1 spinning

disk system set up on a Nikon Eclipse Ti inverted microscope body was used. Detection was

done with an EMCCD camera (iXon Ultra 897, Andor Technologies, Belfast, UK) and a 3i solid

state diode laser stack with 488 nm, 561 nm and 640 nm laser lines (3i, Denver, Colorado,

USA) was used for excitation. Imaging was done with a UPLanSApo 60x/1.20 Water UIS2

objective (Olympus, Japan).

Image processing and analysis

ImageJ (Rueden et al., 2017) was used for image processing and analysis. Rolling ball

background subtraction was used to highlight features in Figure 2A. Rolling ball radii were

100 pixel for actin and 20 pixel for myosin.

Customized macros were made for most intensity and length scale analyses. For intensity

measurement, the macro included a module to specify a square region (edge length 200

pixel) of the sample and then measure mean intensity and standard deviation for each

frame. For calculation of Coefficient of Variance in Figure 3, the ratio between the mean

intensity and the standard deviation was calculated for each time frame. The macro for

length scale analysis combined the use of multipoint selection and the mathematical

transformation of the pixel positions to lengths in microns. Additional plugins used were

Radial_Profile_Angle_Ext and Kymo_wide_reslice (available for download from

https://imagej.nih.gov). The radial intensity profiles in Figure 2D were made by using the

Radial_Profile_Angle_Ext plugin, which integrates the intensities at each point along the

radius of a circular selection. The profile was mirrored to plot the average intensity along

the diameter for aiding visualization of the node organization. For Kymo_wide_reslice, the

pixel thickness of the selected line is specified with corresponding data in the figure legend.

The software FFmpeg was used for making movies from time series data (available for

download from http://ffmpeg.org/).

Statistical analysis and plotting of graphs was done in Origin. Fitting of curves was also done

in Origin using the in-built model "ExpGrow1". Fitting excluded the initial 3 minutes for

achieving a better fit. For significance testing in length scale measurements, values were

pooled across samples for each condition. The nonparametric Mann-Whitney test was used

to compare two populations at a 0.05 significance level.

ACKNOWLEDGMENTS

The authors would like to thank Prof. Klemens Rottner for sharing plasmids and Sigrid Bauer, Dr. Katharina Nakel, Michaela Schaper and Kerstin Andersson, for technical assistance. This work is partially supported by MaxSynBio Consortium, which is jointly funded by the Federal Ministry of Education and Research of Germany and the Max Planck Society. KAG has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 703132. JM is grateful for financial support from the excellence cluster Nanosystems Initiative Munich. S, JM and PB acknowledge support from the International Max Planck Research School for Molecular and Cellular Life Sciences (IMPRS-LS). JM and PB acknowledge support from the Center for NanoScience (CeNS).

REFERENCES

BACKOUCHE, F., HAVIV, L., GROSWASSER, D. & BERNHEIM-GROSWASSER, A. 2006. Active gels: dynamics of patterning and self-organization. *Phys Biol,* 3**,** 264-73.

BIELING, P., HANSEN, S. D., AKIN, O., LI, T. D., HAYDEN, C. C., FLETCHER, D. A. & MULLINS, R. D. 2018. WH2 and proline-rich domains of WASP-family proteins collaborate to accelerate actin filament elongation. *EMBO J,* 37**,** 102-121.

BLANCHOIN, L., POLLARD, T. D. & MULLINS, R. D. 2000. Interactions of ADF/cofilin, Arp2/3 complex, capping protein and profilin in remodeling of branched actin filament networks. *Curr Biol,* 10**,** 1273-82.

BUSSONNIER, M., CARVALHO, K., LEMIERE, J., JOANNY, J. F., SYKES, C. & BETZ, T. 2014. Mechanical detection of a long-range actin network emanating from a biomimetic cortex. *Biophys J,* 107**,** 854-62.

CARVALHO, K., LEMIERE, J., FAQIR, F., MANZI, J., BLANCHOIN, L., PLASTINO, J., BETZ, T. & SYKES, C. 2013. Actin polymerization or myosin contraction: two ways to build up cortical tension for symmetry breaking. *Philos Trans R Soc Lond B Biol Sci,* 368**,** 20130005.

FRITZSCHE, M., LEWALLE, A., DUKE, T., KRUSE, K. & CHARRAS, G. 2013. Analysis of turnover dynamics of the submembranous actin cortex. *Mol Biol Cell,* 24**,** 757-67.

FRITZSCHE, M., THOROGATE, R. & CHARRAS, G. 2014. Quantitative analysis of ezrin turnover dynamics in the actin cortex. *Biophys J,* 106**,** 343-53.

GUHA, M., ZHOU, M. & WANG, Y. L. 2005. Cortical actin turnover during cytokinesis requires myosin II. *Curr Biol,* 15**,** 732-6.

HAVIV, L., GILLO, D., BACKOUCHE, F. & BERNHEIM-GROSWASSER, A. 2008. A cytoskeletal demolition worker: myosin II acts as an actin depolymerization agent. *J Mol Biol,* 375**,** 325-30.

HIGGS, H. N. & POLLARD, T. D. 2000. Activation by Cdc42 and PIP(2) of Wiskott-Aldrich syndrome protein (WASp) stimulates actin nucleation by Arp2/3 complex. *J Cell Biol,* 150**,** 1311-20.

KOHLER, S., SCHALLER, V. & BAUSCH, A. R. 2011. Structure formation in active networks. *Nat Mater,* 10**,** 462-8.

KOSTER, D. V., HUSAIN, K., ILJAZI, E., BHAT, A., BIELING, P., MULLINS, R. D., RAO, M. & MAYOR, S. 2016. Actomyosin dynamics drive local membrane component organization in an in vitro active composite layer. *Proc Natl Acad Sci U S A,* 113**,** E1645-54.

LINSMEIER, I., BANERJEE, S., OAKES, P. W., JUNG, W., KIM, T. & MURRELL, M. P. 2016. Disordered actomyosin networks are sufficient to produce cooperative and telescopic contractility. *Nat Commun,* 7**,** 12615.

MARTIN, A. C., KASCHUBE, M. & WIESCHAUS, E. F. 2009. Pulsed contractions of an actin-myosin network drive apical constriction. *Nature,* 457**,** 495-9.

MEDEIROS, N. A., BURNETTE, D. T. & FORSCHER, P. 2006. Myosin II functions in actin-bundle turnover in neuronal growth cones. *Nat Cell Biol,* 8**,** 215-26.

MUCKSCH, J., BLUMHARDT, P., STRAUSS, M. T., PETROV, E. P., JUNGMANN, R. & SCHWILLE, P. 2018. Quantifying Reversible Surface Binding via Surface-Integrated Fluorescence Correlation Spectroscopy. *Nano Lett*.

MURRELL, M. P. & GARDEL, M. L. 2012. F-actin buckling coordinates contractility and severing in a biomimetic actomyosin cortex. *Proc Natl Acad Sci U S A,* 109**,** 20820-5.

MURTHY, K. & WADSWORTH, P. 2005. Myosin-II-dependent localization and dynamics of F-actin during cytokinesis. *Curr Biol,* 15**,** 724-31.

NISHIKAWA, M., NAGANATHAN, S. R., JULICHER, F. & GRILL, S. W. 2017. Controlling contractile instabilities in the actomyosin cortex. *Elife,* 6.

NYE, J. A. & GROVES, J. T. 2008. Kinetic control of histidine-tagged protein surface density on supported lipid bilayers. *Langmuir,* 24**,** 4145-9.

PANTALONI, D., BOUJEMAA, R., DIDRY, D., GOUNON, P. & CARLIER, M. F. 2000. The Arp2/3 complex branches filament barbed ends: functional antagonism with capping proteins. *Nat Cell Biol,* 2**,** 385-91.

REYMANN, A. C., BOUJEMAA-PATERSKI, R., MARTIEL, J. L., GUERIN, C., CAO, W., CHIN, H. F., DE LA CRUZ, E. M., THERY, M. & BLANCHOIN, L. 2012. Actin network architecture can determine myosin motor activity. *Science,* 336**,** 1310-4.

ROHATGI, R., MA, L., MIKI, H., LOPEZ, M., KIRCHHAUSEN, T., TAKENAWA, T. & KIRSCHNER, M. W. 1999. The interaction between N-WASP and the Arp2/3 complex links Cdc42-dependent signals to actin assembly. *Cell,* 97**,** 221-31.

RUEDEN, C. T., SCHINDELIN, J., HINER, M. C., DEZONIA, B. E., WALTER, A. E., ARENA, E. T. & ELICEIRI, K. W. 2017. ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics,* 18**,** 529.

SALBREUX, G., CHARRAS, G. & PALUCH, E. 2012. Actin cortex mechanics and cellular morphogenesis. *Trends Cell Biol,* 22**,** 536-45.

SMITH, D., ZIEBERT, F., HUMPHREY, D., DUGGAN, C., STEINBECK, M., ZIMMERMANN, W. & KAS, J. 2007. Molecular motor-induced instabilities and cross linkers determine biopolymer organization. *Biophys J,* 93**,** 4445-52.

SOARES E SILVA, M., DEPKEN, M., STUHRMANN, B., KORSTEN, M., MACKINTOSH, F. C. & KOENDERINK, G. H. 2011. Active multistage coarsening of actin networks driven by myosin motors. *Proc Natl Acad Sci U S A,* 108**,** 9408-13.

STAM, S., FREEDMAN, S. L., BANERJEE, S., WEIRICH, K. L., DINNER, A. R. & GARDEL, M. L. 2017. Filament rigidity and connectivity tune the deformation modes of active biopolymer networks. *Proc Natl Acad Sci U S A,* 114**,** E10037-E10045.

TINEVEZ, J. Y., SCHULZE, U., SALBREUX, G., ROENSCH, J., JOANNY, J. F. & PALUCH, E. 2009. Role of cortical tension in bleb growth. *Proc Natl Acad Sci U S A,* 106**,** 18581-6.

VOGEL, S. K., GREISS, F., KHMELINSKAIA, A. & SCHWILLE, P. 2017. Control of lipid domain organization by a biomimetic contractile actomyosin cortex. *Elife,* 6.

VOGEL, S. K., PETRASEK, Z., HEINEMANN, F. & SCHWILLE, P. 2013. Myosin motors fragment and compact membrane-bound actin filaments. *Elife,* 2.

WILSON, C. A., TSUCHIDA, M. A., ALLEN, G. M., BARNHART, E. L., APPLEGATE, K. T., YAM, P. T., JI, L., KEREN, K., DANUSER, G. & THERIOT, J. A. 2010. Myosin II contributes to cell-scale actin network treadmilling through network disassembly. *Nature,* 465**,** 373-7.

YANG, Q., ZHANG, X. F., POLLARD, T. D. & FORSCHER, P. 2012. Arp2/3 complex-dependent actin networks constrain myosin II function in driving retrograde actin flow. *J Cell Biol,* 197**,** 939-56.

LIST OF ABBREVIATIONS

Arp2/3	Actin-related protein 2/3
ATP	Adenosine Triphosphate
CV	Coefficient of Variation
DGS-NTA	1,2-Dioleoyl-sn-glycero-3-phosphoethanolamine
DOPC	1,2-Dioleoyl-sn-glycero-3-phosphocholine
IPTG	Isopropyl β-D-1-thiogalactopyranoside
NPF	Nucleation Promoting Factor
s.d.	Standard deviation
SLB	Supported Lipid Bilayer
TCEP	tris(2-carboxyethyl)phosphine
TIRF	Total Internal Reflection Fluorescence
VCA	Verpolin homology, cofilin, and acidic domain

FIGURES

Figure 1. Assembly of a dynamic actomyosin cortex on supported lipid bilayers.

A) Schematic depiction of the design of a minimal dynamic actomyosin cortex. A dendritic actin network is assembled on a supported lipid bilayer by Arp2/3-induced branch

nucleation, catalyzed by VCA anchored to nickelated lipids (DGS-NTA). Myofilaments are introduced after actin assembly and the system is observed by TIRF microscopy.

B) Time profiles of actin assembly on the membrane and its dependence on different reaction components. The growth is slower in the absence of Arp2/3 and undetectable in the absence of VCA. Addition of VCA (arrow) to the latter shows immediate increase in actin intensity. Concentrations used were 1 μM actin, 10 nM Arp2/3 complex and 100 μM ATP on an SLB with 1 mol% DGS-NTA.

C) Time frames from the profile of the VCA + Arp2/3 reaction depicted in B. Bundling can be observed at 30 minutes. Intensity scale is provided as reference for comparison with images of control conditions in Figure S1.

D) Representative images of the two modes of actomyosin contraction observed in the presence of 0.8 μM myosin-II. The modes differ in density and distribution of condensates.

E) Characteristic length scale of contraction calculated as nearest neighbor distance between foci. Plot shows 25-75 percentile as box, range within 1.5 interquartile region as whiskers, median as line and mean as square. Individual data points are depicted as closed circles. Length scales are significantly different between the two modes ($p < 0.05$; Mann-Whitney test).

Figure 2. Myosin-II breaks down membrane-associated actin networks.

A) After addition of myofilaments, dendritic actin networks show bundling, compaction, network rips and finally removal from the membrane (a). White arrows highlight the mentioned feature in each frame. Panel b shows the location of myofilaments (green) in the corresponding frames of panel a. Yellow arrows highlight myosin in actin bundles and foci.

B) Time profiles of actin and myosin intensities from the sample described in A. The actin intensity is normalized to the value before addition of myosin and decreases as the myosin intensity on the surface increases.

C) Selected time frames from TIRF imaging (a) and kymograph (b) depicting the release of actin from a single condensate formed during coarsening. The kymograph shows intensities averaged over a 10-pixel thick vertical line through the center of the condensate. Data is representative of 10 condensates observed with labeled myosin.

D) Radially integrated intensities of actin and myosin at the different time points depicted in C showing hollowing of the actin core and reduction in actin in the foci. The decrease in myosin intensity is less pronounced in the same duration.

Figure 3. Redistribution in active networks restores homogeneous length scales.

A) Heterogeneity of actin distribution during contraction is represented by the Coefficient of Variation (CV) of actin intensity in the image plotted over time. CV is calculated as the standard deviation divided by the mean for actin intensity distribution at each time frame. Images corresponding to the time points highlighted by cyan circles in the profile of coarsening are shown on the right (a-f).

B) (a) The reappearance of actomyosin foci in regions where actin was removed during coarsening is highlighted by white arrows in selected time frames. (b) The kymograph

plotted from a 10 pixel-thick line connecting two foci shows the emergence of new foci in between. White arrows indicate transfer of material between the foci. Actin and myosin are individually shown in grayscale in the lower frames. (c) The graph shows actin and myosin intensity changes in the newly emerging node measured in a circular region of 10-pixel diameter. The graph inset shows normalized intensities in the first 5 minutes as the new foci emerges.

C) In samples showing coarsening, the distance between neighboring foci decreases over time, coming closer to the value of length scale of homogeneous contraction (gray line with the shaded area depicting mean±s.d.). Asterisk indicates samples were differences are significant after transition ($p < 0.05$; Mann Whitney test). Representative images of the transition of actin distribution in a sample are shown.

Figure 4. Actin undergoes turnover in the minimal dynamic cortex.

A) Schematic of experiment design to follow the turnover of actin filaments observed on the membrane after myosin contraction.

B) Left panel shows kymograph from an experiment of the type described in A. Label1 intensity shows a gradual reduction while the increase in Label2 shows a sharper increase. The right panel shows the corresponding graph showing intensity changes of the two labels in the actin network as well as the ratio of the two labels (dark red).

C) Label2 appears as speckles on the actin structures observed at an intermediate time-point, indicating that the Label2 has an interspersed distribution on filaments with homogenous distribution of Label1.

D) A 1-pixel wide line profile along two of the bundles in C showing normalized intensities of the two labels. The graphs appear to be anti-correlated, indicating that filaments of mixed labeling exist in these bundles.

Figure 5. Myosin activity generates a dynamic steady state actin distribution with continuous turnover in a minimal actin cortex.

A) Schematic summarizing the three key aspects of actin turnover in the minimal dynamic actomyosin cortex reconstituted in this study: 1) breakdown of membrane-bound branched actin networks by myosin, 2) redistribution of network components on the membrane, and 3) incorporation of these network components in the new actin assembly on the membrane.

B) Schematic comparing the results of our study with a previous reconstitution of a minimal actin cortex, where pre-formed actin filaments were attached to an SLB via neutravidin-mediated bridging of biotinylated actin monomers and biotinylated lipids (Vogel et al. 2013).

Our experimental design overcomes constraints on actin assembly and diffusion by using the VCA-Arp2/3 actin filament nucleation module that enables both fast actin polymerization as well as dynamic membrane attachment. Consequently, while myosin-II-driven reorganization of actin was limited to coarsening in that study, we observe a subsequent redistribution resulting in the emergence of a dynamic steady state actin distribution in our minimal dynamic actin cortex.