S100A8/A9 mediate the reprograming of normal mammary epithelial cells

induced by dynamic cell-cell interactions with adjacent breast cancer cells.

Seol Hwa Jo (cho5761@hanmail.net)[1], Woo Hang Heo (woohang92@gmail.com)[1],

Bok Sil Hong (boksil427@gmail.com)[2], Ju Hee Kim (jhsubzero@naver.com)[2], Jiwoo

Lee (leehy69@snu.ac.kr)[2], Eun-Shin Lee (silvershoe99@gmail.com)[3], Han-Byoel Lee

(hblee80@gmail.com)[3], Wonshik Han (hanw@snu.ac.kr)[2,3,4], Yeonju Park

(pyj11@snu.ac.kr)[5], Dong-Sup Lee (dlee5522@snu.ac.kr)[5], Nam Hoon Kwon

(nanarom1@snu.ac.kr)[6], Min Chul Park (minchul.park@neomics.com)[6], Jeesoo Chae

(moverm0210@gmail.com)[4,7], Jong-Il Kim (jongil@snu.ac.kr)[4,7], Dong-Young Noh

(dynoh@snu.ac.kr)[3,4,8], Hyeong-Gon Moon (moonhg74@snu.ac.kr)[2,3,4].

[1]Interdisciplinary Graduate Program in Cancer Biology, Seoul National University

College of Medicine, Seoul, Korea,

[2]Center for Medical Innovation, Biomedical Research Institute, Seoul National

University Hospital, Seoul, Korea,

[3]Department of Surgery, Seoul National University College of Medicine, Seoul, Korea,

[4]Genomic Medicine Institute, Medical Research Center, Seoul National University

College of Medicine, Seoul, Korea,

[5]Department of Biomedical Sciences, Seoul National University College of Medicine,

Seoul, Korea,

[6]Medicinal Bioconvergence Research Center, Seoul National University, Suwon, Korea,

[7]Department of Biochemistry, Seoul National University College of Medicine, Seoul,

Korea

[8]Cancer Research Institute, Seoul National University College of Medicine, Seoul,

Korea

Address correspondence to:

Dong-Young Noh, Seoul National University College of Medicine, 103 Daehak-ro,

Jongno-gu, Seoul, 03080, South Korea. Phone: 82-2-2072-2772; Fax: 82-2-766-3975;

E-mail: dynoh@snu.ac.kr

Hyeong-Gon Moon, Seoul National University College of Medicine, 103 Daehak-ro,

Jongno-gu, Seoul, 03080, South Korea. Phone: 82-2-2072-2634; Fax: 82-2-766-3975;

E-mail: mooonhg74@snu.ac.kr

Abstract

Background: Tumor microenvironment drive the progression of solid tumors. Among the various cell types in the breast cancer microenvironments, non-tumorigenic mammary epithelial cells are often located in the vicinity of malignant cancer cells. However, the biologic characteristics of non-tumorigenic mammary epithelial cells adjacent to breast cancer cells are poorly understood.

Methods: Non-tumorigenic mammary epithelial MCF10A cells were co-cultured with various breast cancer cells and the extent of cell-cell interactions was evaluated by time-lapse imaging. The molecular and functional traits of the co-cultured MCF10A cells were investigated by using RNA sequencing, phosphor-protein arrays, pathway analysis, and various *in vitro* assays. The molecular changes observed in vitro were validated using breast cancer xenograft mouse models. The phenotype changes in the S100A8/A9-overexpressing MCF10A cells were investigated to determine the causal relationship of S100A8/A9 upregulation observed in co-cultured MCF10A cells.

Results: When the cells were co-cultured directly, there were dynamic cell-cell interactions between the MCF10A cells and breast cancer cells including lamellipodia- or nanotube-like contacts and transfer of extracellular vesicles. MCF10A cells exhibited features of epithelial-mesenchymal transition, and showed increased capacity of cell proliferation, migration, colony formation, and 3-dimensional sphere formation upon the direct co-culture with breast cancer cells. Transcriptome analysis and phosphor-protein array suggested that several cancer-related pathways are significantly dysregulated in MCF10A cells after the direct co-culture with breast cancer cells. S100A8/A9 showed distinct up-regulation in the co-cultured MCF10A cells and orthotropic xenograft of syngeneic mouse mammary tumors. When S100A8/A9 overexpression was induced in

MCF10A cells, the cells showed phenotypic features of directly co-cultured MCF10A

cells in terms of *in vitro* cell behaviors and signaling activities.

Conclusions: This study suggests the possibility of dynamic cell-cell interactions

between non-tumorigenic mammary epithelial cells and breast cancer cells that could

lead to a substantial transition of molecular profiles and functional characteristics in

mammary epithelial cells. S100A8/A9, along with other dysregulations in cancer-

related signaling pathways, may mediate this phenotype transition of non-tumorigenic

mammary epithelial cells.

Keywords: Breast cancer, Mammary epithelial cells, Microenvironment, Cell-cell

interaction, S100A8/A9, mTOR

Backgrounds

In solid tumors, the complex tumor microenvironment controls all steps of tumor progression and metastasis [1, 2]. The tumor microenvironment is comprised of various endogenous and recruited cells that undergo dynamic cell-cell interactions with malignant epithelial cells and contribute to the tumor cell's behaviors [3, 4]. For example, cancer-associated fibroblasts actively remodel extracellular matrix and immune microenvironment, and cancer-associated adipocytes provide inflammatory milieu that support tumor growth [5, 6]. Moreover, recent efforts to target the immune microenvironment have shown promising therapeutic responses in selected solid tumors [7]. Therefore, understanding the molecular mechanisms of the tumor-microenvironment interactions can provide scientific basis for developing novel therapeutic strategies that target the tumor microenvironment [3, 8, 9].

Normal epithelial cells are closest neighbors to the malignant transformed cells in human epithelial tumors arising from solid organs. During the early steps of carcinogenesis, the normal epithelial cells may exert tumor-suppressive effects by promoting protrusion of transformed epithelial cells from the epithelial layers [10-12]. However, the tumor-suppressive effects of normal epithelial cells may not last throughout the solid tumor progression. While the normal myoepithelial cells obtained from healthy human breast tissues contribute to the maintaining polarity of mammary epithelial cells and suppress aberrant growth, the myoepithelial cells derived from breast cancer tissues failed to restore physiologic polarity in mammary epithelial cells and showed increased expression of various chemokines such as CXCL12 [13, 14]. These reports suggest a potential functional transition of normal epithelial cells caused by adjacent malignant epithelial cells which may contribute the progression of solid

tumors.

In this study, we show that breast cancer cells and non-tumorigenic mammary epithelial

cells undergo dynamic cell-cell interactions that lead to a substantial reprograming of

molecular characteristics of the mammary epithelial cells. The reprograming of normal

mammary epithelial cells includes phenotypes changes as well as dysregulations of

mRNA expression and cell signaling activities. Our data suggests that S100A8/A9

upregulation in non-tumorigenic mammary epithelial cells may play a critical role in the

phenotype shifting induced by adjacent cancer cells.

139
140

Methods

Cell culture

Cells were purchased from Korean Cell Line Bank (Seoul, Korea). Non-tumorigenic

mammary epithelial MCF10A cells were maintained in a 1:1 mixture of Dulbecco's

Modified Eagle's Medium (DMEM) and Ham's F12 medium (F12) with 5% horse serum,

20 ng/mL epidermal growth factor (EGF), 100 ng/mL cholera toxin, 10 μg/mL insulin,

and 500 ng/mL hydrocortisone. MCF7 and MDA-MB-231 breast cancer cells were

cultured in DMEM with 10% FBS, 1% penicillin, and 1% streptomycin. SK-BR3 breast

cancer cells were cultured in RPMI 1640 with 10% FBS, 1% penicillin, and 1%

streptomycin. For fluorescence tagged cells, puromycin was added.

Co-culture of MCF10A cells and breast cancer cells

To optimize the culture medium for co-culture, we tested the effect of various mixture of

MDA-MB-231 culture media and MCF10A culture media on cell survival. Based on the

effect on cell proliferation, we chose 7:3 ratio mixture of MDA-MB-231 media and

MCF10A media for the direct co-culture of the cells (Additional File 1). For effective

separation of MCF10A cells and MDA-MB-231 cells after the direct co-culture, MCF10A

cells and MDA-MB-231 cells were transfected with GFP and RFP using lentiviral

vectors, respectively. After the direct co-culture, cells were isolated using a FACS Aria II

cell sorter (Becton Dickinson, NJ) as shown in the Additional File 2.

Indirect co-culture was performed by using transwell inserts (pore size 0.4um). Cancer

cells were seeded on the membrane of the insert and MCF10A cells were seeded on

the bottom six well plates. For controls, same cells were seeded in the bottom wells

and on the transwell inserts.

***In vitro* assays measuring cell phenotypes**

Proliferation assays were conducted using CellTiter-Glo Luminescent Cell Viability

Assay (Promega, Madison, USA) following the manufacturer's protocol. Cells were

seeded in triplicate into 96-well plates at a density of 2,000 cells per well. For migration

assay, $2x10^4$ cells were seeded in an insert (8μm pore size) with serum free media and

media with 10% FBS was added in lower chambers. Cells were incubated for 20hrs

and fixed with 4% paraformaldehyde and stained with 0.1% crystal violet. For colony

formation assay, $2x10^3$ cells were seeded into 6-well plates. After 2 weeks, colonies

were fixed in 4% paraformaldehyde and stained with 0.1% crystal violet.

RNA Sequencing and qPCR

RNA sequencing libraries were prepared using TruSeq RNA Access library kit (Illumina,

Inc., San Diego, CA, USA) according to the manufacturer`s protocol. After validation of

the libraries, using Agilent DNA screentape D1000 kit on a TapeStation (Agilent

Technologies, Santa Clara, CA, USA), the hybridization steps were performed using

exome capture probes and streptavidin coated beads. RNA sequencing was performed

by HiSeq 2000 (Illumina,San Diego, USA by the Macrogen Incorporated). We

processed reads from the sequencer and aligned them to the Homo sapiens (hg19).

After aligning reads to genome, Cufflinks v2.2.1 was used to assemble aligned reads

into transcripts and to estimate their abundance. The transcript counts in isoform and

gene level were calculated, and the relative transcript abundances were measured in

FPKM (Fragments Per Kilobase of exon per Million fragments mapped) from Cufflinks.

RNA sequencing data of breast cancer was generated through a separate project to

characterize the genomic profiles of the primary breast tumor and patient-derived

xenograft tumors which will be presented in other reports (IRB No. 1402-054-555).

For qPCR, total RNA was extracted from isolated cells with TRIzol (Favorgen, Taiwan).

PrimeScript 1st strand cDNA Synthesis Kit (Takara, Japan) were used for reverse

transcription of RNA, and resulting cDNA was amplified using Power SYBR® Green

PCR Master Mix (Applied Biosystems, CA). The information on the primer sequences

used in this study is listed in the Additional File 3.

Phosphorylation array and western blotting

For phosphorylation array, protein extraction was done with buffer following the

manufacturer's protocol. Protein concentration was measured by BCA assay kit

(Thermo scientific, Palm Springs, CA, USA). Phosphorylation array was performed by

using the human phosphor-kinase array kit (R&D systems, USA).

Proteins were harvested with RIPA buffer (Thermo scientific, Palm Springs, CA, USA),

protease & phosphatase inhibitor and 0.5M EDTA solution. Protein concentration was

measured by BCA assay kit (Thermo scientific, Palm Springs, CA, USA). Cell lysates

were loaded onto 10% gels and transferred to a PVDF membrane. The membrane was

blocked with 5% skim milk and incubated with primary antibody overnight at 4℃. The

peroxidase-conjugated secondary antibody was used for detection. Bands were

detected by LAS. The information on the antibodies used in this study is listed in the

Additional File 4.

Cell imaging

For time-lapse imaging of cells during the direct co-culture, $9x10^3$ GFP-transfected

MCF10A cells and $2x10^4$ RFP-transfected MDA-MB-231 cells were seeded in 8-well

chambers and cultured for 24 hours. After 24 hours culture, live cells images were

taken every 10 minutes with fixed position for 16 hours by using Leica confocal

microscope (Leica TCS SP8, Leica Microsystems Ltd, Korea), and the obtained images

were analyzed by Leica LAS X program.

For *in vivo* imaging, the experiments were performed in accordance with the Animal

Care and Use Committee guidelines of Woosung BSC (Suwon, Korea). Hair-removed

ear skin of C57BL/6 mouse (10 weeks old) was injected with 4.6 x 10^4 cells of the

mixture MCF10A-GFP (2.3 x 10^4) and MDA-MB-231-RFP (2.3 x 10^4) cell lines. The

mice were anesthetized with an intraperitoneal injection of Zoletil (30 mg/kg, Virbac,

Carros, France) and Rompun (10 mg/kg, Bayer-Korea, Seoul, Korea) before imaging.

The mice were placed on the heated plate of a motorized XYZ translational stage. The

in vivo movement of MCF10A and MDA-MB-231 was monitored by modified custom-

built laser-scanning confocal microscopy {Choe, 2015 #3}. GFP-expressing cells were

visualized at an excitation wavelength of 491 nm and detected through a bandpass

filter of 502 nm to 537 nm (Semrock Inc, Rochester, NY, USA). RFP-expressing cells

were imaged at an excitation wavelength of 532 nm and detected using a bandpass

filter of 562 nm to 596 nm (Semrock Inc). Cell movement was visualized at 1 min

interval for 2 hr. After acquired from the imaging system, 512 x 512 pixel images were

then compensated with Matlab (Mathworks, Natick, MA, USA) and reconstructed by

ImageJ software.

S100A8/A9 expression in xenograft tumor-bearing mouse fatpad

5-week-old BALB/c mice were purchased from KOATECH (Seoul, Korea) and housed

in Seoul National University Hospital Clinical Research Institute's Specific Pathogen

Free zone. All experiments were approved by the Institutional Animal Care and Use

Committee in Seoul National University Hospital (SNUH-IACUC, 17-0165-S1A0 (1)). To

obtain control mouse fatpads, eight-week-old non-tumor-bearing BALB/c mouse was

sacrificed and mammary fat pad was resected. To obtain tumor-bearing mouse fat pad,

2 X 10^5 4T1 mouse mammary epithelial cancer cells were injected in 6-week-old

BALB/c mouse's mammary fat pad. At two weeks after the tumor injection, the mouse

was sacrificed and tumor-bearing mammary fatpad was resected. The resected fatpads

were fixed with 4% PFA and embedded in paraffin.

S100A8/A9-overexpressing MCF10A cell line

The coding sequences of S100A8 and S100A9 were acquired by RT-PCR and cloned

into pCDH-GFP and pCDH-RFP vectors. MCF10A cells were transfected with lentivirus

S100A8-GFP and S100A9-RFP construct. Transfected cells were selected by

puromycin.

Stastatic analysis

In general, most data represent the mean ± S.D and are representative of 3

independent experiments, except for RNA sequencing and phospho-protein arrays

which are two independent experiments. Graph Pad Prism (ver. 7.01) was used for

generating graphs and heatmaps and performing statistical tests. P values were

calculated from unpaired two-tailed Student's t tests or Mann-Whitney test as

appropriate.

To identify differentially expressed genes, we filtered genes with one more than zeroed

FPKM values and the data were log2-transformed and subjected to quantile

normalization. Statistical significance of the differential expression data was determined

using fold change in which the null hypothesis was that no difference exists among

samples. Gene pathway analysis for the DEG was done based on KEGG pathway

(http://www.genome.jp/kegg/pathway.html).

Results

Breast cancer cells show dynamic cell-cell interactions with non-transformed

mammary epithelial cells *in vitro* and *in vivo*.

First, we determined the presence and the extent of cell-cell interactions *in vitro*

between the breast cancer cells and mammary epithelial cells. We co-cultured the

RFP-transfected breast cancer cells (MDA-MB-231) with GFP-transfected non-

transformed mammary epithelial cells (MCF10A) using *in vitro* direct co-culture method.

While the majority of MCF10A cells maintained the clusters of adherent cells, MDA-MB-

231 cells showed spreading patterns of cell growth and the cells infiltrated between the

MCF10A cell clusters (Additional File 5). The time-lapse imaging of the cells showed

that MDA-MB-231 cells had more frequent cell movements than the MCF10A cells and

the cells showed various dynamic cell-cell interaction patterns (Additional File 6 and 7).

MDA-MB-231 cells formed both lamellipodia-like structures for adjacent cells and

nanotube-like projections for long-range cell-cell interactions (Figure 1a and 1b,

Additional File 5) [15, 16]. The lamellipodia-like structures of MDA-MB-231 cells

actively contacted the MCF10A cells and a portion of extended lamellipodia could

remain as extracellular vesicles which were then engulfed by adjacent MCF10A cells

(Figure 1c, Additional File 7).

Among the various cell-cell interactions, exchanges of extracellular vesicles were

frequently seen between the cells. MCF10A cells engulfed extracellular vesicles

originated from MDA-MB-231 cells, and the vesicles were often transferred to the

nucleus while some stayed at cytoplasm (Figure 1d and 1e). Additionally, a small

proportion of co-cultured cells (1~2%) exhibited mixed fluorescence (Figure 1f,

Additional File 2).

To observe the cell-cell interactions *in vivo*, we inoculated a mixture of MDA-MB-231

cells and MCF10A cells in the earlobes of nude mouse and obtained the time-lapse

imaging data. After one hour of the inoculation, the movements of the MDA-MB-231

cells around the MCF10A cells were detectable (Additional File 8 and 9). After 24 hours,

the cell-cell interactions and exchange of vesicles were more frequently seen between

the cells (Figure 1g, Additional File 9). These data suggest that the cancer cells and

epithelial cells can undergo a dynamic range of physical contacts and cell-cell

interactions both *in vitro* and *in vivo*.

Direct co-culture with breast cancer cells induce phenotypic changes in MCF10A

cells

We then determined whether the cell-cell interactions with breast cancer cells induce

phenotype changes in MCF10A cells. First, we co-cultured MCF10A cells with breast

cancer cell lines using in-direct co-culture system. MCF10A cells co-cultured indirectly

with cancer cells showed no significant changes in cell morphology or the cell growth

patterns in 2D and 3D cultures (Additional File 10). However, when the MCF10A cells

were cultured in the direct co-culture system with breast cancer cells, the MCF10A cells

showed substantial phenotypic changes. Directly co-cultured MCF10A cells, sorted by

GFP-expression, showed changes in cell morphology such as transition into spindle-

shaped cells and loss of cell-cell adhesions (Figure 2a).

Additionally, the MCF10A cells showed marked decrease in E-cadherin expression

when they were directly co-cultured with MDA-MB-231 cells (Figure 2b). When the

isolated MCF10A cells were grown *in vitro*, the proliferation rate, colony formation

capacity, and the cell migration ability were significantly enhanced in cells directly co-

cultured with MDA-MB-231 cells (Figure 2c-2e). MCF10A cells also showed higher

numbers and larger sizes of cell spheres when they were grown in 3D matrigels

(Figure 2f). Collectively, our data show that direct co-culture with breast cancer cells

induce substantial phenotype changes in non-tumorigenic mammary epithelial cells.

Transcriptome changes and pathway dysregulations in MCF10A cells when

directly co-cultured with MDA-MD-231 cells

To understand the molecular mechanisms underlying the phenotype changes induced

by direct co-culture, we analyzed the transcriptome profiles of MCF10A cells when they

were directly co-cultured with MDA-MB-231 cells. A total of 151 genes were

dysregulated in directly co-cultured MCF10A cells more than two-fold in expression

(Figure 3a, Additional File 11). Again, we observed the decreased expression of

mammary epithelial differentiation markers such as CDH1 or CD24. Based on the gene

expression profiles, we identified several cancer-related pathways, including metabolic

pathway, cell adhesion, growth factor signaling, and TP53 pathway, that were

significantly dysregulated in MCF10A cells after the direct co-culture (Figure 3b).

To determine the changes in the cell signaling in MCF10A cells when the cells were

directly co-cultured with MDA-MB-231 cells, we used the phosphorylation antibody

array to profile the phosphor-protein signaling pathways. Many of the candidate

phosphor-proteins showed differential levels of expression in the directly co-cultured

MCF10A cells. PRAS40, STAT6, ERK1/2 levels were substantially increased while

HSP60, β-catenin, RSK1/2/3, STAT3, and p53 levels were downregulated (Figure 3c).

These results suggest that non-tumorigenic mammary epithelial cells undergo a

significant shift in transcriptomic and proteomic characteristics.

S100A8/A9 induction in peritumoral mammary epithelial cells and stromal

tissues

Among the differentially expressed genes, S100A8 and S100A9 genes were highly

upregulated in directly co-cultured MCF10A cells and their increased expression was

validated in mRNA and protein levels (Figure 3a and 4a). When S100A8 and S100A9

mRNA levels in normal breast and breast cancer tissues were examined using RNA

sequencing data comprised of 65 normal and 68 cancer tissues, both genes were

significantly upregulated in breast cancer tissues when compared to normal breast

tissues (Figure 4b). Furthermore, the mRNA dataset from Finak et al [17] showed that

S100A8 and S100A9 genes were significantly upregulated in the adjacent stroma

tissues in human breast cancer (Figure 4c).

We then used mouse mammary carcinoma xenograft model to determine whether the

induction of S100A8/A9 expression by breast cancer cells also occurs *in vivo*. BALB/c

mouse fatpads were injected with 4T1 murine mammary carcinoma cells and were

subsequently harvested. Compared to non-tumor bearing fatpads, mouse mammary

epithelial cells and peritumoral stromal cells showed increased expression of both

S100A8 and S100A9 (Figure 4d and 4e). These findings in addition to the above

transcriptome data indicate that breast cancer cells induce S100A8/9 expression in

tumor microenvironment especially in non-tumorigenic mammary epithelial cells.

S100A8/A9 upregulation contribute to the phenotypic and molecular changes in

the co-cultured MCF10A cells

To determine the functional importance of S100A8 and S100A9 gene expression in

mammary epithelial cells, we established a stable MCF10A cells that overexpress

S100A8 and S100A9 genes. Transduction of S100A8 overexpression vector resulted in

upregulation of both S100A8 and S100A9 proteins (Additional File 12). S100A8-

overexpressing MCF10A cells showed higher rate of cell proliferation when compared

to the control cells (Figure 5a). Furthermore, S100A8-overexpressing MCF10A cells

showed increased cell migration, invasion, colony formation, and 3-dimensional cell

growth which are characteristics of the MCF10A cells directly co-cultured with breast

cancer cells. (Figure 5b-5d).

To investigated the effect of S100A8 and S100A9 expression on the signaling pathway

activities, we repeated the phosphorylation antibody array in S100A8-overexpressed

MCF10A cells (Figure 5e-5f). More than half of the signaling proteins observed in

S100A8-overexpressed MCF10A cells showed concordant expression patterns with

directly co-cultured MCF10A cells and only one protein showed opposite expression

pattern between the two cells (Figure 5g). The dysregulation of signaling pathways in

both S100A8 overexpressing MCF10A cells and the MCF10A cells directly co-cultured

with MDA-MB-231 cells were further validated using western blotting as shown in

Figure 5h. Taken together, out data demonstrate that, at least in part, the phenotypic

and molecular transitions seen in the mammary epithelial cells when co-cultured with

breast cancer cells can be explained by the upregulation of S100A8/A9.

Discussion

Normal mammary epithelial cells are closest neighboring cells to malignant epithelial

cells during the initial process of carcinogenesis. Also, cancer cells often contact

adjacent normal epithelial cells during the invasion process. The underlying hypothesis

of our work was that normal epithelial cells adjacent to cancer cells may undergo

molecular and functional transition despite the normal microscopic appearance. Indeed,

we observed that non-tumorigenic epithelial cells undergo a substantial molecular and

phenotypic changes when the cells were directly exposed to the breast cancer cells.

Non-tumorigenic mammary epithelial MCF10A cells showed significant increase in cell

proliferation and migration when they were directly co-cultured with breast cancer cells.

Along with the changes in cell behavior, directly co-cultured MCF10A cells also

exhibited spindle-shaped morphologies and significant downregulation of epithelial

adhesion markers such as E-cadherin and Zo-1. Previous studies have shown that E-

cadherin expression in normal epithelial cells plays an important role during the

process initial carcinogenesis by regulating cell protrusion formation of neighboring

transformed cells [12, 18]. Additionally, three-dimensional direct co-culture with breast

cancer cells induced loss of epithelial differentiation features in non-tumorigenic MDCK

cells [19]. Finally, Trujillo et al [20] have shown that normal epithelial cells adjacent

breast tumors show increased expression of EMT markers such as α-SMA and

S100A4. Our data, along with these previous reports, suggests that breast cancer cells

may induce epithelial mesenchymal transition in adjacent normal mammary epithelial

cells via direct cell-cell contacts.

Using the *in vitro* direct co-culture approach, we observed a dynamic and complex cell-

cell interactions between breast cancer cells and non-tumorigenic mammary epithelial

cells that include diverse physical contacts and vesicle transfers. We were also able to

demonstrate that the tumor cells and normal epithelial cells exhibit various physical

interactions *in vivo* by showing live images taken from the mouse earlobe model. Such

dynamic interactions have also been reported in a three-dimensional co-culture

experiment [20] and between various stromal cell types in microenvironment [21].

Recently, Roh-Johnson et al [22] have shown that the cytoplasmic transfer during the

cell-cell contacts between melanoma cells and macrophages is critical in cancer cell

invasion *in vivo*. The clinical and biologic consequences of this cell-cell interactions

between malignant cancer cells and adjacent normal epithelial cells are unclear.

Normal epithelial cells exert tumor-suppressive effects during the initial phase of

carcinogenesis by cell competition and protrusion of neighboring transformed cells [18,

23, 24]. Alternatively, tumor cells can reprogram the adjacent normal epithelial cells to

form a pro-tumorigenic microenvironment at the late stages of carcinogenesis [13]. This

phenomenon of tumor-driven reprograming of microenvironment has been shown for

other cell types such as fibroblast or adipocytes in breast cancer [5, 6]. Further

research is needed to clarify the role of molecular transformation of normal epithelial

cells during the breast cancer progression.

In molecular levels, we observed that S100A8/A9 expression levels are significantly

up-regulated when non-tumorigenic MCF10A epithelial cells were co-cultured with

MDA-MB-231 cells. The up-regulation of S100A8/A9 was also observed in a syngeneic

mouse xenograft model of 4T1 murine mammary carcinoma cells. S100A8/A9 act as

alarmins that trigger damage-associated molecular patterns molecules and modulate

the immune response [25]. S100A8/A9 overexpression in MCF10A cells resulted in a

similar cell phenotype seen in directly co-cultured MCF10A cells in terms of in vitro cell

behaviors and cell signaling activities. Moon A and the colleagues have previously

shown that S100A8/A9 overexpression can induce phenotype transformation in

MCF10AHRAS cells by   . Our results show that S100A8/A9, potentially critical

regulators of cell behavior, could be induced in non-tumorigenic mammary epithelial

cells during the dynamic cell-cell interactions with adjacent breast cancer cells.

Additionally, the directly co-cultured and S100A8/A9-overexpressing cells both showed

increased phosphorylation of PRAS40-Thr^{246} and its upstream Akt-Ser^{473}.

Phosphorylation of Akt-Ser^{473} and PRAS40-Thr^{246} activate mTOR signaling that affects

diverse biologic aspects of the epithelial cells and other cells of tumor

microenvironment [26-28]. The biologic implications and therapeutic potentials of the

observed mTOR signaling dysregulation in mammary epithelial cells in breast cancer

microenvironment should be further explored.

Our study has several limitations. First, we have used a limited number of cell lines

during the present study. The molecular transitions in non-tumorigenic mammary

epithelial cells might vary along the different subtypes of breast cancers. Second, the

effect of this phenotype changes in mammary epithelial cells on the breast cancer cells

in terms of breast cancer growth and metastasis has not been investigated using *in*

vivo breast cancer models. Finally, although we've demonstrated the up-regulation of

S100A8/A9 expression in mouse fatpad xenograft tumors, we were not able to

examine the S100A8/A9 up-regulation in human breast cancer tissues. Further studies

investigating the biologic consequence and clinical implications of the molecular

changes in the non-tumorigenic mammary epithelial cells in breast cancer

microenvironment are needed.

Conclusion

Conclusion

In conclusion, our study demonstrate that breast cancer cells may induce substantial

molecular changes in non-tumorigenic mammary epithelial cells via dynamic cell-cell

interactions. As the results, mammary epithelial cells undergo a phenotype transition

which involves more active proliferation and migration. S100A8/A9 may play a pivotal

role during this phenotype transition of mammary epithelial cells. Our study provides

scientific basis for pursuing a novel therapeutic strategy that targets the non-

tumorigenic mammary epithelial cells in tumor microenvironments.

Declarations

Ethics approval and consent to participate

This study does not include data on the human subjects. For animal experiments, the

fatpad xenograft tumor experiment was approved by the Institutional Animal Care and

Use Committee in Seoul National University Hospital (SNUH-IACUC, 17-0165-S1A0

(1)), and the *in vivo* imaging experiments were performed in accordance with the

Animal Care and Use Committee guidelines of Woosung BSC (Suwon, Korea).

Consent for publication

This manuscript does not contain any individual person's data in any form (including

individual details, images or videos).

Availability of data and materials

Data supporting these results are available from the authors upon request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Basic Science Research Program through the National

Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and

Technology (2015R1D1A1A02061904), by the grant from the National R&D Program

for Cancer Control, Ministry for Health and Welfare, Republic of Korea (A1520250),

and by the grant of the Korea Health Industry Development Institute (KHIDI), funded by

the Ministry of Health & Welfare, Republic of Korea (HI13C2148).

Authors' contributions

SHJ performed the bulk of the work reported, including writing the initial draft,

generating cell lines, and phospho-protein arrays. WHH performed fatpad tumor

xenograft studies, PCR, western blotting and writing the initial draft. BSH, JHK

performed YP and DSL assisted and provide guidance to cell sorting and animal

experiments. NHK and MCP performed in vivo cell imaging studies. ESL, HBL, and WH

analyzed the transcriptome data and pathway analysis. JC and JIK provided RNA

sequencing data of breast cancer and analyzed the target gene expression levels.

DYN and HGM initiated the study, provided overall direction and oversight to the

project, and finalized the manuscript. All authors read and approved the final

manuscript.

Acknowledgements

Not applicable.

References

1. Joyce JA, Pollard JW: **Microenvironmental regulation of metastasis**. *Nat Rev Cancer* 2009, **9**(4):239-252.

2. Hanahan D, Coussens LM: **Accessories to the crime: functions of cells recruited to the tumor microenvironment**. *Cancer Cell* 2012, **21**(3):309-322.

3. Place AE, Jin Huh S, Polyak K: **The microenvironment in breast cancer progression: biology and implications for treatment**. *Breast Cancer Res* 2011, **13**(6):227.

4. Gajewski TF, Schreiber H, Fu YX: **Innate and adaptive immune cells in the tumor microenvironment**. *Nat Immunol* 2013, **14**(10):1014-1022.

5. Kalluri R: **The biology and function of fibroblasts in cancer**. *Nat Rev Cancer* 2016, **16**(9):582-598.

6. Lee J, Hong BS, Ryu HS, Lee HB, Lee M, Park IA, Kim J, Han W, Noh DY, Moon HG: **Transition into inflammatory cancer-associated adipocytes in breast cancer microenvironment requires microRNA regulatory mechanism**. *PLoS One* 2017, **12**(3):e0174126.

7. Pardoll DM: **The blockade of immune checkpoints in cancer immunotherapy**. *Nat Rev Cancer* 2012, **12**(4):252-264.

8. Klemm F, Joyce JA: **Microenvironmental regulation of therapeutic response in cancer**. *Trends Cell Biol* 2015, **25**(4):198-213.

9. Fang H, Declerck YA: **Targeting the tumor microenvironment: from understanding pathways to effective clinical trials**. *Cancer Res* 2013, **73**(16):4965-4977.

10. Porazinski S, de Navascues J, Yako Y, Hill W, Jones MR, Maddison R, Fujita Y, Hogan C: **EphA2 Drives the Segregation of Ras-Transformed Epithelial**

Cells from Normal Neighbors. *Curr Biol* 2016, **26**(23):3220-3229.

11. Saitoh S, Maruyama T, Yako Y, Kajita M, Fujioka Y, Ohba Y, Kasai N, Sugama N,

Kon S, Ishikawa S *et al*: **Rab5-regulated endocytosis plays a crucial role in**

apical extrusion of transformed cells. *Proc Natl Acad Sci U S A* 2017,

114(12):E2327-E2336.

12. Hogan C, Dupre-Crochet S, Norman M, Kajita M, Zimmermann C, Pelling AE,

Piddini E, Baena-Lopez LA, Vincent JP, Itoh Y *et al*: **Characterization of the**

interface between normal and transformed epithelial cells. *Nat Cell Biol*

2009, **11**(4):460-467.

13. Gudjonsson T, Ronnov-Jessen L, Villadsen R, Rank F, Bissell MJ, Petersen OW:

Normal and tumor-derived myoepithelial cells differ in their ability to

interact with luminal breast epithelial cells for polarity and basement

membrane deposition. *J Cell Sci* 2002, **115**(Pt 1):39-50.

14. Allinen M, Beroukhim R, Cai L, Brennan C, Lahti-Domenici J, Huang H, Porter

D, Hu M, Chin L, Richardson A *et al*: **Molecular characterization of the tumor**

microenvironment in breast cancer. *Cancer Cell* 2004, **6**(1):17-32.

15. Mattila PK, Lappalainen P: **Filopodia: molecular architecture and cellular**

functions. *Nat Rev Mol Cell Biol* 2008, **9**(6):446-454.

16. Rustom A, Saffrich R, Markovic I, Walther P, Gerdes HH: **Nanotubular**

highways for intercellular organelle transport. *Science* 2004,

303(5660):1007-1010.

17. Finak G, Bertos N, Pepin F, Sadekova S, Souleimanova M, Zhao H, Chen H,

Omeroglu G, Meterissian S, Omeroglu A *et al*: **Stromal gene expression**

predicts clinical outcome in breast cancer. *Nat Med* 2008, **14**(5):518-527.

18. Kajita M, Hogan C, Harris AR, Dupre-Crochet S, Itasaki N, Kawakami K,

Charras G, Tada M, Fujita Y: **Interaction with surrounding normal epithelial**

cells influences signalling pathways and behaviour of Src-transformed

cells. *J Cell Sci* 2010, **123**(Pt 2):171-180.

19. Ivers LP, Cummings B, Owolabi F, Welzel K, Klinger R, Saitoh S, O'Connor D,

Fujita Y, Scholz D, Itasaki N: **Dynamic and influential interaction of cancer**

cells with normal epithelial cells in 3D culture. *Cancer Cell Int* 2014,

14(1):108.

20. Trujillo KA, Heaphy CM, Mai M, Vargas KM, Jones AC, Vo P, Butler KS, Joste

NE, Bisoffi M, Griffith JK: **Markers of fibrosis and epithelial to mesenchymal**

transition demonstrate field cancerization in histologically normal tissue

adjacent to breast tumors. *Int J Cancer* 2011, **129**(6):1310-1321.

21. Fong EL, Wan X, Yang J, Morgado M, Mikos AG, Harrington DA, Navone NM,

Farach-Carson MC: **A 3D in vitro model of patient-derived prostate cancer**

xenograft for controlled interrogation of in vivo tumor-stromal

interactions. *Biomaterials* 2016, **77**:164-172.

22. Roh-Johnson M, Shah AN, Stonick JA, Poudel KR, Kargl J, Yang GH, di

Martino J, Hernandez RE, Gast CE, Zarour LR *et al*: **Macrophage-Dependent**

Cytoplasmic Transfer during Melanoma Invasion In Vivo. *Dev Cell* 2017,

43(5):549-562 e546.

23. Leung CT, Brugge JS: **Outgrowth of single oncogene-expressing cells from**

suppressive epithelial environments. *Nature* 2012, **482**(7385):410-413.

24. Kajita M, Fujita Y: **EDAC: Epithelial defence against cancer-cell competition**

between normal and transformed epithelial cells in mammals. *J Biochem*

2015, **158**(1):15-23.

25. Srikrishna G: **S100A8 and S100A9: new insights into their roles in**

malignancy. *J Innate Immun* 2012, **4**(1):31-40.

26. Wiza C, Nascimento EB, Ouwens DM: **Role of PRAS40 in Akt and mTOR**

signaling in health and disease. *Am J Physiol Endocrinol Metab* 2012,

302(12):E1453-1460.

27.   Kim LC, Cook RS, Chen J: **mTORC1 and mTORC2 in cancer and the tumor**

microenvironment. *Oncogene* 2017, **36**(16):2191-2201.

28.   Duluc C, Moatassim-Billah S, Chalabi-Dchar M, Perraud A, Samain R, Breibach

F, Gayral M, Cordelier P, Delisle MB, Bousquet-Dubouch MP *et al*:

Pharmacological targeting of the protein synthesis mTOR/4E-BP1

pathway in cancer-associated fibroblasts abrogates pancreatic tumour

chemoresistance. *EMBO Mol Med* 2015, **7**(6):735-753.

Figure legends

Figure 1. Dynamic cell-cell interactions between breast cancer cells and non-

tumorigenic mammary epithelial cells.

Representative images of cell-cell interactions between RFP-transfected MDA-MB-231

breast cancer cells and GFP-transfected MCF10A non-tumorigenic mammary epithelial

cells were captured using time-lapse confocal microscopy (a-d). **a** MDA-MB-231 cells

use lamellipodia-like structures to contacting MCF10A cells (yellow arrow) or distant

MCF10A cells (white arrow). **b** Nanotube-like structures extending from MDA-MB-231

cells are seen (white arrows). **c** RFP-expressing vesicles from MDA-MB-231 cells are

transferred to MCF10A cells (white arrows). **d-e** The area within the white rectangle in

Figure 1d is shown in 1e. Transferred vesicles from MDA-MB-231 cells are located in

both nucleus (white arrow) and cytoplasm (yellow arrow) of MCF10A cells. **f** A minority

of co-cultured cells show dual fluorescence. **g** The representative images of the *in vivo*

behavior of MDA-MB-231 cells and MCF10A cells co-injected in the earlobe of mouse

are shown with migrating extracellular vesicles (white arrows).

Figure 2. Phenotype transition of non-tumorigenic mammary epithelial cells

when directly co-cultured with breast cancer cells.

a MCF10A cells were directly co-cultured with MDA-MB-231, SK-BR3, and MCF7

breast cancer cells. After sorting out the MCF10A cells, the cellular morphologies are

shown. **b** The expression levels of epithelial-mesenchymal transition markers were

measured by PCR in MCF10A cells after co-culture with MDA-MB-231 cells (upper)

and quantified (lower). **c** Relative growth rate of co-cultured MCF10A cells was

measured based on ATP level with CellTiter-Glo reagent. **d** The representative images

and quantified results of colony-formation assay for co-cultured MCF10A cells are

shown. **e** The representative images and quantified results of transwell migration assay

for co-cultured MCF10A cells are shown. **f** The representative images and quantified

results of 3-dimensional Matrigel culture assay for co-cultured MCF10A cells are shown.

Error bars denote mean ± SD. *P < 0.05, **P < 0.01, ***P < 0.001. P values are

determined by the Mann-Whitney test.

Figure 3. Transcriptomic and proteomic profiles of non-tumorigenic mammary

epithelial cells co-cultured with breast cancer cells.

a Heatmap shows the top 60 genes that were up- or down-regulated in MCF10A cells

co-cultured with MDA-MB-231 cells. **b** Heatmap shows the significant enrichment of

KEGG pathways based on the RNA sequencing data. Pathways significantly

dysregulated in both datasets are listed on the right side. **c** Results of phosphor-protein

array experiments are shown. Spots with more than two-fold changes are marked with

numbered rectangles and listed below (upper). The overall quantification results of

phospho-protein array are shown (lower).

Figure 4. S100A8/A9 up-regulation in breast cancer tissues and

microenvironments.

a S100A8/A9 mRNA levels and protein levels in MCF10A cells co-cultured with MDA-

MB-231 cells are shown. **b** The S100A8/A9 mRNA levels extracted from human breast

cancer RNA sequencing data are shown. **c** S100A8/A9 mRNA levels in breast cancer

stromal tissues are shown (Finak et al. GSE9014). **d** Mouse mammary fatpad tissues

bearing 4T1 breast cancer cells were stained against S100A8 and S100A9. Lower

panels show magnified images of the upper insets. **e** Non-tumor-bearing mouse fatpad

were stained with S100A8 and S100A9. *P < 0.05, **P < 0.01, ***P < 0.001. P values are

determined by the unpaired Student t-test.

Figure 5. S100A8/A9-overexpression induce the phenotypes of non-tumorigenic

mammary epithelial cells co-cultured with breast cancer cells.

a Relative growth rate of S100A8/A9-overexpressing MCF10A cells was determined

based on ATP level with CellTiter-Glo reagent. **b** The representative images and

quantified results of transwell migration assay for S100A8/A9-overexpressing MCF10A

cells are shown. **c** The representative images and quantified results of Matrigel

invasion assay for S100A8/A9-overexpressing MCF10A cells are shown. **d** The

representative images and quantified results of colony-formation assay for S100A8/A9-

overexpressing MCF10A cells are shown. **e-f** Results of phosphor-protein array

experiments with S100A8/A9-overexpressing MCF10A cells (e) and quantified

expression of significantly dysregulated proteins (f) are shown. **g** The comparison of

phospho-protein array patterns for co-cultured MCF10A cells and S100A8/9-

overexpressing MCF10A cells are shown. For example, 'Down/Up' designates spots

that are down-regulated in co-cultured MCF10A cells and up-regulated in S100A8/A9-

overexpressing MCF10A cells, respectively. **h** Western blotting results showing the

protein expression levels of selected dysregulated proteins in both co-cultured

MCF10A cells and S100A8/A9-overexpressing MCF10A cells. Error bars denote mean

± SD. **P < 0.01, ***P < 0.001. P values are determined by the Mann-Whitney test.

a

b

c

¹PRAS40 (T246), ²β-catenin, ³p53 (S15), ⁴STAT3 (Y705), ⁵HSP60, ⁶RSK1/2/3 (S380/S286/S377),