Assessing distinct patterns of cognitive aging using tissue-specific brain age prediction based on diffusion tensor imaging and brain morphometry

Geneviève Richard[a,b,c,*], Knut Kolskår[a,b,c], Anne-Marthe Sanders[a,b,c], Tobias Kaufmann[a], Anders Petersen[d], Nhat Trung Doan[a], Jennifer Monereo Sánchez[a], Dag Alnæs[a], Kristine M. Ulrichsen[a,b,c], Erlend S. Dørum[a,b,c], Ole A. Andreassen[a], Jan Egil Nordvik[c], Lars T. Westlye[a,b,*]

[a] NORMENT, KG Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Norway

[b] Department of Psychology, University of Oslo, Norway

[c] Sunnaas Rehabilitation Hospital HT, Nesodden, Norway

[d] Center for Visual Cognition, Department of Psychology, University of Copenhagen, Copenhagen, Denmark

*Corresponding authors:

Geneviève Richard & Lars T. Westlye

Email: genevieve.richard@medisin.uio.no, l.t.westlye@psykologi.uio.no

Postal address: Oslo University Hospital, P.O.Box 4956 Nydalen, 0424 OSLO, Norway

Telephone: +47 23 02 73 50, Fax: +47 23 02 73 33

Short title:

Tissue-specific brain age prediction

Key words:

Brain age, machine learning, white matter, gray matter, DTI, T1

Disclosures:

The authors declare no competing financial interests.

Abstract

Multimodal imaging enables sensitive measures of the architecture and integrity of the human

brain, but the high-dimensional nature of advanced brain imaging features poses inherent

challenges for the analyses and interpretations. Multivariate age prediction reduces the

dimensionality to one biologically informative summary measure with potential for assessing

deviations from normal lifespan trajectories. A number of studies documented remarkably

accurate age prediction, but the differential age trajectories and the cognitive sensitivity of

distinct brain tissue classes have to a lesser extent been characterized.

Exploring differential brain age models driven by tissue-specific classifiers provides a

hitherto unexplored opportunity to disentangle independent sources of heterogeneity in brain

biology. We trained machine-learning models to estimate brain age using various

combinations of FreeSurfer based morphometry and diffusion tensor imaging based indices of

white matter microstructure in 612 healthy controls aged 18-87 years. To compare the tissue-

specific brain ages and their cognitive sensitivity we applied each of the 11 models in an

independent and cognitively well-characterized sample (n=265, 20-88 years). Correlations

between true and estimated age in our test sample were highest for the most comprehensive

brain morphometry (r=0.83, CI:0.78-0.86) and white matter microstructure (r=0.79, CI:0.74-

0.83) models, confirming sensitivity and generalizability. The deviance from the

chronological age were sensitive to performance on several cognitive tests for various models,

including spatial Stroop and symbol coding, indicating poorer performance in individuals

with an over-estimated age. Tissue-specific brain age models provide sensitive measures of

brain integrity, with implications for the study of a range of brain disorders.

Introduction

Increasing age is a major risk factor for cognitive decline and neurodegeneration, and

deviating lifespan trajectories in brain structure and function is a sensitive marker in several

common neurological and mental disorders (Cole & Franke 2017). The maturing and aging

brain is highly heterogeneous in term of individual trajectories and in term of brain regions

and mechanisms involved (Fjell et al. 2013; Westlye et al. 2010b). Understanding the

individual determinants and heterogeneity of the developing and aging brain is imperative for

identifying persons at risk for various brain disorders, and for developing and applying

effective and targeted treatments.

Exploring different modalities acquired by magnetic resonance imaging (MRI)

provide a powerful tool to investigate age-related differences in both gray- and white- matter

tissue classes across brain regions. However, the richness and complexity of the information

provided by advanced imaging pipelines challenges its interpretation. Together, the

multifactorial age-related variability and the richness of imaging measures have motivated the

development of biologically informative summary measures based on brain imaging data.

Using machine-learning to estimate the biological age of the brain based on neuroimaging

data is one such approach (Cole & Franke 2017; Cole et al. 2018; Kaufmann et al. 2018).

Deviation from the normative trajectory is a highly relevant biomarker for the integrity of the

brain in healthy and clinical populations (Marquand et al. 2016; Wolfers et al. in press). Brain

age gap is a heritable trait showing regionally specific genetic overlaps with major brain

disorders, including schizophrenia and multiple sclerosis (Kaufmann et al. 2018), and

accumulating evidence supports increased brain age in several clinical groups, including

patients with schizophrenia (Kaufmann et al. 2018; Schnack et al. 2016), Alzheimer's disease

(Amoroso et al. 2017; Kaufmann et al. 2018), HIV (Cole et al. 2017b; Kuhn et al. 2018)

multiple sclerosis (Kaufmann et al. 2018) and cardiovascular risk factors (Franke et al. 2013;

Habes et al. 2016). Indeed, while individuals with brains estimated as younger than their

chronological age have been shown to be more physically active (Steffener et al. 2016),

augmented brain age has been associated with poor health (Ronan et al. 2016), poor cognitive

performance (Liem et al. 2017), early neurodegenerative diseases (Gaser et al. 2013), and

increased mortality (Cole et al. 2017a). Less is known about the regional heterogeneity, i.e. to

which degree different brain regions, systems or compartments show differential aging

patterns and sensitivity to cognitive performance. Brain gray and white matter compartments,

which can be assessed and quantified using T1-weighted imaging and diffusion tensor

imaging (DTI), respectively, comprise distinct tissue classes with largely differential

biological and environmental modifiers and age trajectories (Bennett et al. 2010; Cao et al.

2017; Fjell et al. 2013; Salat et al. 2005; Storsve et al. 2014; Westlye et al. 2010a; Westlye et

al. 2010b). Therefore, allowing for differential brain age models for these distinct classes

provides an opportunity to disentangle independent sources of heterogeneity in brain aging.

Thus, to identify common and unique aging patterns with potentially differential

sensitivity to cognitive function, we aimed to test the complementary value of tissue-specific

prediction by comparing brain age estimated using different combinations of FreeSurfer based

morphometric measures (regional cortical thickness, surface area and volume) and white

matter microstructure features (DTI based fractional anisotropy and mean, radial and axial

diffusivity) across the brain. Based on previous studies on brain aging, we expected high

accuracy and generalizability of the age prediction models (Cole & Franke 2017). Since tissue

specific brain age models capture biologically distinct information, we anticipated that the

different FreeSurfer based brain morphometry and white matter microstructure models would

only partly reflect common variance, and therefore provide complementary information with

differential sensitivity to cognitive performance. Given that brain age predictions might be

sensitive to the overall integrity of the brain (Liem et al. 2017), we anticipated that individuals

with an over-estimated brain age would show lower cognitive performance, in particular

among the elderly, and that the tissue-specific brain age models would show partly

differential cognitive sensitivity.

To ensure generalizability, we trained the models in a large publicly available training

set (n=612, 18-87 years) and validated their performance using 10-fold cross-validation

before applying to an independent and well characterized test set (n=265, 20-88 years). We

assessed the cognitive sensitivity using linear and non-linear models with performance on a

range of paper-and-pencil and computerized tests comprising different large-scale cognitive

domains (processing speed, executive functioning, working memory, attention, and general

intellectual abilities) and cognitive scores based on computational models as dependent

variables and age, sex and brain age gap (BAG, estimated brain age minus chronological age)

as independent variables. For transparency, we report results both at an uncorrected level and

corrected using false discovery rate (FDR) and Bonferroni methods to control the error rate.

Materials and methods

Table 1 summarizes key demographics. We included data from healthy volunteers from two

independent cohorts: (1) the Cambridge Centre for Ageing and Neuroscience (Cam-CAN)

sample (http://www.mrc-cbu.cam.ac.uk/datasets/camcan/; (Shafto et al. 2014; Taylor et al.

2017)) and (2) StrokeMRI, which is an ongoing study on the determinants of stroke recovery,

brain health and successful aging (Dorum et al. 2016; Dorum et al. 2017). Figure 1 shows the

age distribution for each sample.

Volunteers were recruited to Cam-CAN through a large-scale collaborative research

project funded by the Biotechnology and Biological Sciences Research Council (BBSRC,

grant number BB/H008217/1), the UK Medical Research Council and University of

Cambridge. For more information, see www.cam-can.org. Among the 650 datasets made

available, 17 were excluded based on missing or poor quality DTI data and 21 due to poor

T1-weighted data quality. Data from the remaining 612 individuals (age 18-87, mean = 54.41,

SD = 18.26, 314 females) were included.

Healthy individuals were recruited to StrokeMRI through advertisement in

newspapers, social media and word-of-mouth. All participants completed a comprehensive

cognitive assessment, multimodal MRI and blood sampling for clinical biochemical analysis,

various biomarkers and genotyping. MRI and cognitive assessments were performed on two

subsequent days. Exclusion criteria included history of stroke, dementia, or other neurologic

and psychiatric diseases, alcohol- and substance abuse, medications significantly affecting the

nervous system and counter indications for MRI. In addition, individuals scoring lower than

25 on the Montreal Cognitive Assessment (MoCA; Nasreddine et al. 2005) were assessed for

inclusion based on their age, level of education and performance on other cognitive tests. No

participants were excluded based on a single low score. A neuroradiologist reviewed all scans

and 14 participants with clinically significant abnormalities were excluded. Additional

exclusion criteria included missing or incomplete MRI or cognitive data (n=2), or poor

quality images (n=20). The remaining 265 participants (age 20-88, mean = 56.95, SD = 14.84,

168 females) were included in further analyses. The study was approved by the Regional

Committee for Medical and Health Research Ethics (South-East Norway), and conducted in

accordance with the Helsinki declaration. All subjects signed an informed consent prior to

participating and received a compensation for their participation.

Fig. 1 Histogram of the age distribution for each sample.

Cognitive assessment in StrokeMRI

Cognitive performance was assessed with a set of neuropsychological and computerized tests

assumed to be sensitive to cognitive aging, including the MoCA, the vocabulary and matrix

subtests of the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler 1999), the

California Verbal Learning Test (CVLT-II; Delis et al. 2000), and the Delis-Kaplan Executive

Function System (D-KEFS) color word interference test (Stroop; Delis et al. 2001). We

included several computerized tests from the Cognitive Assessment at Bedside for iPAD

(CABPad; Willer et al. 2016), including motor speed, verbal fluency (phonological and

semantic), working memory, spatial Stroop (executive control of attention), spatial attention

span, and symbol digit coding tests. In addition, in order to assess the specificity of cognitive

associations using computation modeling, we included three mathematically independent

parameters based on the Theory of Visual Attention (TVA; Bundesen 1990; Bundesen &

Habekost 2008), including short-term memory storage (K), processing speed (C), perceptual

threshold (t_0). These parameters based on computational modeling of response patterns have

been shown to be sensitive to age, brain structure and function in healthy individuals

(Espeseth et al. 2014; Wiegand et al. 2018) and a range of brain disorders (Habekost 2015;

Habekost & Starrfelt 2009). Here, we used a TVA-based modeling of a whole report

(Sperling 1960), in which six letters were briefly presented for different exposure durations

and the participant's task was to accurately report as many letters as possible. Task error rate

was also assessed (i.e. number of incorrect letters out of reported letters).

MRI acquisition

Cam-CAN participants were scanned on a 3T Siemens TIM Trio scanner with a 32-channel

head-coil at Medical Research Council (UK) Cognition and Brain Sciences Unit (MRC-

CBSU) in Cambridge, UK. DTI data was acquired using a twice-refocused spin echo

sequence with the following parameters a repetition time (TR) of 9100 ms, echo time (TE) of

104 ms, field of view (FOV) of 192 x 192 mm, voxel size: 2 mm^3, 66 axial slices using 30

directions with b= 1000 s/mm^2, 30 directions with b= 2000 s/mm^2, and 3 b=0 images (Shafto

et al. 2014). Only the b=[0,1000] were used in the current analysis. High-resolution 3D T1-

weighted data was acquired using a magnetization prepared rapid gradient echo (MPRAGE)

sequence with the following parameters: TR: 2250 ms, TE: 2.99 ms, inversion time (TI): 900

ms, flip angle (FA): 9°, FOV of 256 x 240 x 192mm; voxel size =1 mm^3 isotropic, GRAPPA

acceleration factor of 2, scan time 4:32 minutes (Shafto et al. 2014).

StrokeMRI participants were scanned on a 3T GE 750 Discovery MRI scanner with a

32-channel head coil at Oslo University Hospital. Paddings were used to reduce head motion.

DTI data were acquired using an echo planar imaging (EPI) sequence with the following

parameters: TR/TE/flip angle: 8150 ms/83.1 ms/90°, FOV: 256 x 256 mm, slice thickness: 2

mm, in-plane resolution: 2 mm, 60 directions (b=1000 s/mm^2) and 5 b=0 volumes, scan time:

8:58 min. In addition, 7 b=0 volumes with reversed phase-encoding direction were acquired.

High-resolution T1-weighted data was acquired using a 3D IR-prepared FSPGR (BRAVO)

with the following parameters: repetition time: 8.16 ms, echo time: 3.18 ms, flip angle: 12º,

voxel size: 1 × 1 × 1 mm, field of view: 256 x 256 mm, 188 sagittal slices, scan time: 4:43

minutes.

DTI processing and analysis

Diffusion MRI data from both samples were processed locally using the Oxford Center for

Functional Magnetic Resonance Imaging of the Brain (FMRIB) Software Library (FSL)

(http://www.fmrib.ox.ac.uk/fsl). To correct for geometrical distortions, motion and eddy

currents, data were preprocessed using topup (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/topup) and

eddy (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/eddy) respectively (Andersson et al. 2003; Smith et

al. 2004). Topup uses information from the reversed phase-encoded image, resulting in pairs

of images (blip-up, blip-down) with distortions going in opposite directions. From these

image pairs the susceptibility-induced off-resonance field was estimated and the two images

were combined into a single corrected one (Andersson et al. 2003; Smith et al. 2004). This

step was performed on StrokeMRI data only. Eddy detects and replaces slices affected by

signal loss due to bulk motion during diffusion encoding, which is performed within an

integrated framework along with correction for susceptibility induced distortions, eddy

currents and motion (Andersson & Sotiropoulos 2016). Although these processing steps have

been shown to strongly increase the temporal signal-to-noise ratio (tSNR) (Doan et al. 2017),

we performed additional visual inspection to identify and remove poor quality data.

Fractional anisotropy (FA), eigenvector, and eigenvalue maps were calculated using

dtifit in FSL. Mean diffusivity (MD) was defined as the mean of all three eigenvalues, radial

diffusivity (RD) as the mean of the second and third eigenvalue, and axial diffusivity (AD) as

the principal eigenvalue.

Voxelwise analysis of FA, MD, AD and RD were carried out using Tract-Based

Spatial Statistics (TBSS; Smith et al. 2006), part of FSL (Smith et al. 2004). First, all subjects'

FA data were aligned to a common space using the nonlinear registration tool FNIRT

(Andersson et al. 2007a; Andersson et al. 2007b). Next, the mean FA image was created and

thinned to create a mean FA skeleton, which represents the centers of all tracts common to all

participants. Each subject's aligned FA data was then projected onto this skeleton and the

resulting data fed into voxelwise cross-subject statistics. The same warping and

skeletonization was repeated for MD, AD and RD. We thresholded and binarized the mean

FA skeleton at FA>0.2. For each individual, we calculated the mean skeleton FA, MD, AD

and RD, as well as mean values within 23 regions of interest (ROIs) based on two

probabilistic white matter atlases provided with FSL, i.e. the CBM-DTI-81 white-matter

labels atlas and the JHU white-matter tractography atlas (Hua et al. 2008; Mori et al. 2005;

Wakana et al. 2007), yielding a total of 96 DTI features per individual.

T1 processing

T1-weighted images were processed using FreeSurfer 5.3 (http://surfer.nmr.mgh.harvard.edu;

(Dale et al. 1999)) including brain extraction, intensity normalization, automated tissue

segmentation, generation of white and pial surfaces (Dale et al. 1999). All reconstructions

were visually assessed and corrected as appropriate. Cortical parcellation was performed

using the Desikan–Killiany atlas (Desikan et al. 2006; Fischl et al. 2004) and subcortical

segmentation was performed based on a probabilistic atlas (Fischl et al. 2002). In addition to

global features (intracranial volume, total surface area, whole-cortex mean thickness), mean

thickness, total surface area, and volume for each cortical ROI, as well as the volume of

subcortical structures were computed yielding a set of 251 FreeSurfer based features.

Age prediction

Eleven different models were trained to estimate age based on the feature sets described

above (one based on FreeSurfer T1 features, one based on WM DTI features, one including

all T1 and DTI features, in addition to eight models based on a smaller subset of features,

including models based on FA, MD, AD, RD, sub-cortical volume, volume, area and

thickness to further explore the modality specificity of the estimations).

Due to the systematic differences in the brain features across scanners as well as non-

linear effects of age, we fit a generalized additive model (GAM; Hastie 2017) for each feature

and regressed out the effects of scanning site and age², accounting all models for age and sex.

In addition, we regressed out the estimated total intracranial volume from the area and volume

features. Next, for each model, we created a training data matrix by concatenating all the

features for all participants in the training sample (Cam-CAN), which were used as input to

estimate age. We used the *xgboost* framework in R (http://xgboost.readthedocs.io/en/latest/R-

package/xgboostPresentation.html), an efficient and scalable implementation of gradient

boosting machine learning techniques, to build the prediction models. The following

parameters were used: learning rate (eta) = 0.1, nround = 5000, gamma = 1, max_depth = 6,

subsample=0.5. To estimate the performance of our models, we used a 10-fold cross-

validation procedure within the training sample and repeated the cross-validation step 1000

times to provide a robust estimate of model predictive accuracy. Next, we tested the

performance of our trained models by predicting age in unseen healthy subjects in the test

sample (StrokeMRI).

For each feature set, we calculated the correlation between the predicted and the

chronological age as a measure of the model performance, in addition to the mean absolute

error (MAE, in years). For each individual, we calculated the discrepancy between the

estimated and the chronological age, i.e. the BAG, for each model. The MAE was calculated

from the BAG for each model. Since we were interested in the effect of BAG independently

of age, the effect of age was regressed out for each BAG using linear models.

Statistical analysis

Statistical analysis was performed using R (http://www.r-project.org). For cognitive data, we

used *outlierTest* from the car package (Fox & Weisberg 2011) to identify the most extreme

observations based on a linear model, including age and sex. Twenty-five observations were

identified as outliers and treated as missing values based on a Bonferroni corrected p < 0.05.

To visualize the associations between the cognitive tests and to form cognitive domain scores

based on the correlation patterns, we performed hierarchical clustering using the default

setting of the heatmap.2 package in gplots (Warnes et al. 2016), which uses hclust (Müllner

2013) to form clusters based on the complete linkage method. Briefly, this is a step-wise

clustering process that merges the two nearest clusters until only one single cluster remains,

maximizing distance between individuals components between two clusters.

For each cognitive measure and summary score based on the clusters formed form the

clustering step above, we used linear models to test for the effect of age and sex. Since

cognitive performance may show non-linear associations with age, we performed an

additional analysis including both age and age^2 in the models. Then, for each test showing a

significant association with age, we tested whether adding BAG to the models lead to an

improved model fit. More specifically, we tested for differential associations with cognitive

function by comparing the parameter estimates for the different BAG models using Fisher z-

transformation. To test the assumption that increased BAG is more relevant for cognitive

function among the elderly, we tested for age by BAG interactions on cognitive performance.

For transparency, we report both uncorrected p-values and p-values adjusted using FDR

(Benjamini & Hochberg 1995; Wright 1992) and Bonferroni correction using a factor of 495

(11 brain gaps and 45 cognitive features).

Results

Brain age prediction

10-fold cross-validation on the training sample revealed high correlations between

chronological and predicted age for the DTI based white matter microstructure (r=0.87) and

FreeSurfer based morphometric (r=0.88) models. Likewise, the correlations for FA (r=.76),

MD (r=.80), AD (r=.83), RD (r=78), sub-volume (r=.84), volume (r=.80), area (r=.70) and

thickness (r=.79) based models also confirmed reasonable model performance.

Most models accurately predicted age in the independent test set. Figure 2A shows a

correlation matrix for the 11 BAGs. Figure 2B shows the correlations between the

chronological age and the predicted age in the test sample for each model with their

confidence intervals, ranging from (r=.86, CI:.82-.89, MAE= 6.14) for the combined model to

r=.58 (CI:.49-.65, MAE=10.24) for the model based on area. Figure 2 (D to F) show the

estimated age from the three models that performed best among the 11 feature sets, i.e. the

combined DTI and T1 feature models (r=.86, CI:.82-.89, MAE= 6.14), the 251 FS T1 features

(r=.83, CI:.78-.86, MAE= 6.76), and the 96 WM DTI features (r=.79, CI:.74-.83, MAE=7.28).

Fig. 2 Comparison between the 11 BAG models. (a) Heatmap of the correlation between
different BAGs. (b) Correlations between the chronological age and the predicted age in the
test sample for each model with their confidence intervals. (c) Mean and standard error of the
45 p-values ($-\log_{10}(p)$) for the cognitive scores and composite scores for each row (i.e.
BAGs), with a higher mean representing a stronger global association across tests. (d)
Correlation between the chronological age of each subjects and the combined age, (e) the
brain morphometry age, and (f) the white matter microstructure age.

Cognitive assessments and associations with BAGs

Table 1 summarizes descriptive statistics and associations with age and sex for each of the 49

cognitive scores, derived features and domain scores. Linear models revealed 45 significant

associations with age after correcting for multiple comparisons, with the strongest effect sizes

for the symbol coding test, motor speed, spatial Stroop and spatial attention span. Since non-

linear models revealed significant associations with age² only with the color word Stroop 3

(inhibition) and its derived scores (See supplementary Table S1), the main models presented

here are linear in order to keep the model to its simplest form. Figure 3 shows a correlation

matrix across all normalized cognitive scores with the variables sorted according to the

hierarchical clustering. Several variables were highly correlated, and the clustering solution

generally suggested seven broad cognitive domains including (Cluster 1) memory and

learning (CVLT, attention span, MoCA), (Cluster 2) visual processing speed (TVA

processing speed and perceptual threshold), (Cluster 3) verbal skills (phonological and

semantic flow), (Cluster 4) attentional control and speed (spatial Stroop), (Cluster 5)

executive control and speed (color-word Stroop), (Cluster 6) reasoning and psychomotor

speed (matrix, symbol coding and motor speed, short-term memory storage (TVA-parameter

K)), and (Cluster 7) working memory.

Fig. 3 Hierarchical clustering of the cognitive features. Each cognitive score was normalized

and when required the scores were multiplied by -1 to ensure that positive scores represent

good performance. The higher panel shows the dendrogram resulting from the hierarchical

clustering of the scores in 7 cognitive domains.

Table 2 shows summary statistics for the associations between cognitive performance and

BAG using linear models, including age and sex as covariates. Figure 4 shows a heatmap of

the association between cognitive scores and brain age gaps for which the significant

associations have been marked with an asterisk. Supplementary Table S1 and Fig. S1 shows

the summary statistics and the heatmap of the associations between cognitive performance and BAG using non-linear models. Figure 2C shows the mean and standard error of the 45 p-values ($-\log_{10}(p)$) for the cognitive scores and composite scores for each row (i.e. BAGs), with a higher mean representing a stronger cumulative association across tests.

Figure 5 shows a scatter plot of the 2 strongest associations, which were found between the most comprehensive model (all features combined) and spatial Stroop congruent trials and number of responses, respectively, indicating poorer performance with higher BAG. Fisher z-transformation revealed no statistically significant differences in the cognitive associations between linear models using tissue-specific BAG. No significant interactions were found between BAG and age on cognitive performance.

Fig. 4 Heatmap of the association between cognitive scores and brain age gaps. The color scale depicts the minus log of the p-values ($-\log_{10}(p)$) for each association. The association marked with a small star represents significant associations after FDR correction, and the one marked with a big star shows significant associations after Bonferroni correction.

Fig. 5 Scatter plots of the 2 strongest associations between cognitive measures and BAG. The color gradient represents the age where lighter color is assigned to older individuals, and darker color to younger individuals. All associations indicate worse performance with higher brain age gap.

Discussion

Brain aging is highly heterogeneous, and expanding our understanding of the biological determinants of human aging is imperative for reducing the burden of age-related cognitive decline and neurodegenerative disorders. An estimate of an individual's deviation from the expected lifespan trajectory in brain structure and function may provide a sensitive measure of individual brain integrity and health, both in presumably healthy individuals and in patients suffering from various brain disorders.

The biological heterogeneity of the brain strongly suggests that the concept of a single brain age is too simple, and that tissue-specific brain age models may provide increased sensitivity and specificity in relation to cognitive and mental functions. In line with this view, our main findings demonstrate that different combinations of FreeSurfer based brain morphometry and DTI based white matter microstructural indices can be used to accurately predict the age of individuals, but that the shared variance from the different models suggest that they reflect partly non-overlapping processes of brain aging. Further, the results revealed partly differential sensitivity to cognitive performance; with the strongest cumulative associations across cognitive tests for brain age gaps estimated using RD. Even though our data provide no strong evidence of independent associations with cognitive performance in the current sample of healthy individuals, tissue specific age prediction models might better

inform us about the individual determinants and heterogeneity of the aging brain compared to

models collapsing several brain compartments by potentially capturing distinct measures of

brain aging.

Brain age prediction

For the age prediction models, our results demonstrated that the 11 different combinations of

FreeSurfer based morphometric measures (regional cortical thickness, surface area and

volume) and white matter microstructure features (diffusion tensor imaging (DTI) based

fractional anisotropy and mean, radial and axial diffusivity) across the brain age models

accurately predicted the age of an individual with a mean absolute error between 6.14 and

10.23 years. Brain morphometry and white matter microstructure models had a MAE of 6.76

and 7.28 respectively, which correspond with previous publications (Cole et al. 2016; Han et

al. 2014; Valizadeh et al. 2017). In general, combining features and modalities increased the

performance, and the highest performing model included a combination of both brain

morphometry and white matter microstructure (mean absolute error of 6.14 years). Moreover,

the correlations between the different brain age gaps suggested a relatively low level of shared

variance (mean correlation = 0.51, SD=0.13). Together these findings support the notion that

tissue specific brain age models capture biologically distinct information. This is in line with

the characteristic lifespan patterns of global linear decreases in gray matter volume and the

nonlinear trajectories of total white matter volume and DTI based metrics of white matter

microstructure (Cox et al. 2016; Fjell et al. 2013; Ge et al. 2002; Liu et al. 2017; Raz et al.

2010; Westlye et al. 2010b), highlighting that the different compartments carry unique

biological information and that combining different modalities lead to a better estimation the

age of individuals (Cherubini et al. 2016; Liem et al. 2017; Madan & Kensinger 2018).

Cognitive associations

We performed a comprehensive cognitive assessment of the test sample, confirming previous

evidence of substantial age-related differences in cognitive performance across a range of

tests and domains. Hierarchical clustering of the cognitive features indicated a characteristic

pattern of covariance, largely reflecting broad cognitive domains, including memory and

learning, visual processing speed, verbal skills, attentional and executive control, reasoning

and psychomotor speed, and working memory. Ninety percent of the included cognitive

features showed age-differences, with the largest effect sizes observed for speed-based

measures, such as symbol coding test, which measures mental and visuo-motor speed (Willer

et al. 2016). This is in line with the well-established literature on age-related decline in

information processing speed in healthy aging (Bennett et al. 2010; Craik & Salthouse 2008;

Harada et al. 2013). Importantly, not only tasks measuring reaction time, but also various

TVA measures based on computational modeling, such as short-term memory storage (K),

processing speed (C), and perceptual threshold (t_0) showed strong associations with age, in

line with previous studies (Espeseth et al. 2014; Habekost 2015; Habekost et al. 2013;

McAvinue et al. 2012; Wiegand et al. 2018).

Based on the assumption that brain age captures variance related to the integrity of the

brain, we anticipated that individuals with an over-estimated age would show lower cognitive

performance, and that the tissue-specific brain age models would show partly differential

sensitivity. To test these hypotheses, we used linear models to explore the associations

between cognitive performance and BAG, with age and sex as covariates, and directly

compared the parameter estimates from the different brain age models. We found a significant

association between performance on several tests and BAG beyond the age associations,

indicating lower performance in individuals with higher BAG. Briefly, one significant

association was found for WM DTI, five for combined BAG, two for the sub-volume, one for

the RD and one for the MD BAG. The strongest associations were found with the spatial

Stroop congruent trials, and number of responses. These findings support that the deviance

between the estimated age and the chronological age captures relevant biological information

regarding the cognitive performance of an individual. Whereas we found no significantly

different associations between brain age models, the association with symbol digit coding test

was only seen for WM DTI BAG, while associations with Stroop 3 and 4 were observed only

for sub-volume BAG, suggesting some specificity that should be investigated in future studies

including larger samples and a broader spectrum of mental health, cognitive and brain

phenotypes, both across healthy and clinical samples. We speculate that the contributions of

the different modalities in predicting age and the associations with both cognitive

performance, but also age-related illnesses vary across the age-span, as it does during

maturational age (Brown et al. 2012). Thus, future studies might benefit from investigating

modality specific brain-age estimation using specific age range.

Limitations

The present findings do not come without limitations. First, although reducing the

dimensionality of complex brain imaging data to a biologically informative brain age is a

powerful method to assess deviations from normal lifespan trajectories in brain health,

findings from this data reduction method are limited in specificity. Here, we attempted to both

reduce the complexity of the information while keeping some modality specificity measured

by different MRI parameters. Finding a balance between specificity and precision represents

an interesting challenge for future studies. Moreover, causality and individual level

trajectories cannot be established based on cross-sectional data. Therefore, future longitudinal

studies are needed to inform us about the relevance of the differential trajectories of the

tissue-specific brain age prediction with implications for the study of a range of brain

disorders. Next, although the age distribution of the test sample is irrelevant to the individual

prediction accuracy, the relative overrepresentation of older individuals in the test sample is a

limitation when investigating interactions between BAG and age. Thus, although the lack of

brain by BAG interactions on cognitive function did not support our hypothesis that increased

BAG is more relevant for cognitive function among the elderly, future studies including

individuals across a broader range of function are needed to characterize the lifespan

dynamics in the associations between brain and behavior. Although we covered a relatively

broad spectrum of structural brain features, the link between imaging based indices of brain

structure and brain function is elusive, and brain age models including other brain imaging

features, including functional measures, might provide a sensitive supplement to the current

models. Lastly, whereas the results showed some numerical differences in the cognitive

sensitivity of the different combinations of FreeSurfer based morphometry and white matter

microstructure models, these differences were not statistically significant, and the hypothesis

that tissue specific models provide increased specificity in terms of cognitive associations

remains to be further explored in future studies.

In conclusion, we have demonstrated that models based on different combinations of

brain morphometry and white matter microstructural indices provide partly differential

information about the aging brain, emphasizing the relevance of tissue-specific brain age

models in the study of brain and mental function in health and disease.

Acknowledgements

This study was supported by the Norwegian ExtraFoundation for Health and Rehabilitation

(2015/FO5146), the Research Council of Norway (249795, 248238), the South-Eastern

Norway Regional Health Authority (2014097, 2015044, 2015073), Sunnaas Rehabilitation

Hospital, and the Department of Psychology, University of Oslo, and approved by the

Regional Committee for Medical and Health Research Ethics (South-East). Volunteers were

recruited to StrokeMRI through advertisement in newspapers, social media and word-of-

mouth. In addition, volunteers were recruited to Cam-CAN through a large-scale

collaborative research project funded by the Biotechnology and Biological Sciences Research

Council (BBSRC, grant number BB/H008217/1), the UK Medical Research Council and

University of Cambridge.

References

Amoroso N, Diacono D, Fanizzi A, La Rocca M, Monaco A, Lombardi A, Guaragnella

C, Bellotti R, Tangaro S, and Alzheimer's Disease Neuroimaging I. 2017. Deep learning

reveals Alzheimer's disease onset in MCI subjects: results from an international challenge. *J*

Neurosci Methods. 10.1016/j.jneumeth.2017.12.011

Andersson JLR, Jenkinson M, and Smith S. 2007a. TR07BP1: Non-linear optimisation.

FMRIB Analysis Group Technical Reports: FMRIB Analysis Group.

Andersson JLR, Jenkinson M, and Smith S. 2007b. TR07JA2: Non-linear registration,

aka spatial normalization. FMRIB Analysis Group Technical Reports: FMRIB Analysis

Group.

Andersson JLR, Skare S, and Ashburner J. 2003. How to correct susceptibility

distortions in spin-echo echo-planar images: application to diffusion tensor imaging.

Neuroimage 20:870-888. 10.1016/S1053-8119(03)00336-7

Andersson JLR, and Sotiropoulos SN. 2016. An integrated approach to correction for

off-resonance effects and subject movement in diffusion MR imaging. *Neuroimage* 125:1063-

1078. 10.1016/j.neuroimage.2015.10.019

Benjamini Y, and Hochberg Y. 1995. Controlling the False Discovery Rate - a Practical

and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-*

Methodological 57:289-300.

Bennett IJ, Madden DJ, Vaidya CJ, Howard DV, and Howard JH, Jr. 2010. Age-related

differences in multiple measures of white matter integrity: A diffusion tensor imaging study

of healthy aging. *Hum Brain Mapp* 31:378-390. 10.1002/hbm.20872

Brown TT, Kuperman JM, Chung Y, Erhart M, McCabe C, Hagler DJ, Jr.,

Venkatraman VK, Akshoomoff N, Amaral DG, Bloss CS, Casey BJ, Chang L, Ernst TM,

Frazier JA, Gruen JR, Kaufmann WE, Kenet T, Kennedy DN, Murray SS, Sowell ER,

Jernigan TL, and Dale AM. 2012. Neuroanatomical assessment of biological maturity. *Curr*

Biol 22:1693-1698. 10.1016/j.cub.2012.07.002

Bundesen C. 1990. A theory of visual attention. *Psychol Rev* 97:523-547.

Bundesen C, and Habekost T. 2008. Principles of Visual Attention: Linking Mind

and Brain. *Oxford University Press*. 10.1093/acprof:oso/9780198570707.001.0001

Cao B, Mwangi B, Passos IC, Wu MJ, Keser Z, Zunta-Soares GB, Xu DP, Hasan KM,

and Soares JC. 2017. Lifespan Gyrification Trajectories of Human Brain in Healthy

Individuals and Patients with Major Psychiatric Disorders. *Scientific Reports* 7. ARTN 511

10.1038/s41598-017-00582-1

Cherubini A, Caligiuri ME, Peran P, Sabatini U, Cosentino C, and Amato F. 2016.

Importance of Multimodal MRI in Characterizing Brain Tissue and Its Potential Application

for Individual Age Prediction. *Ieee Journal of Biomedical and Health Informatics* 20.

10.1109/Jbhi.2016.2559938

Cole JH, and Franke K. 2017. Predicting Age Using Neuroimaging: Innovative Brain

Ageing Biomarkers. *Trends Neurosci*. 10.1016/j.tins.2017.10.001

Cole JH, Marioni RE, Harris SE, and Deary IJ. 2018. Brain age and other bodily 'ages':

implications for neuropsychiatry. *Mol Psychiatry*. 10.1038/s41380-018-0098-1

Cole JH, Poudel RPK, Tsagkrasoulis D, Caan MWA, Steves C, Spector TD, and

Montana G. 2016. Predicting brain age with deep learning from raw imaging data results in a

reliable and heritable biomarker. *ARXIV*. 2016arXiv161202572C

Cole JH, Ritchie SJ, Bastin ME, Valdes Hernandez MC, Munoz Maniega S, Royle N,

Corley J, Pattie A, Harris SE, Zhang Q, Wray NR, Redmond P, Marioni RE, Starr JM, Cox

SR, Wardlaw JM, Sharp DJ, and Deary IJ. 2017a. Brain age predicts mortality. *Mol*

Psychiatry. 10.1038/mp.2017.62

Cole JH, Underwood J, Caan MWA, De Francesco D, van Zoest RA, Leech R, Wit

FWNM, Portegies P, Geurtsen GJ, Schmand BA, van der Loeff MFS, Franceschi C, Sabin

CA, Majoie CBLM, Winston A, Reiss P, Sharp DJ, and Collaboration C. 2017b. Increased

brain-predicted aging in treated HIV disease. *Neurology* 88:1349-1357.

10.1212/Wnl.0000000000003790

Cox SR, Ritchie SJ, Tucker-Drob EM, Liewald DC, Hagenaars SP, Davies G, Wardlaw

JM, Gale CR, Bastin ME, and Deary IJ. 2016. Ageing and brain white matter structure in

3,513 UK Biobank participants. *Nat Commun* 7:13629. 10.1038/ncomms13629

Craik FIM, and Salthouse TA. 2008. *The handbook of aging and cognition*. New York:

Psychology Press.

Dale AM, Fischl B, and Sereno MI. 1999. Cortical surface-based analysis - I.

Segmentation and surface reconstruction. *Neuroimage* 9:179-194. DOI

10.1006/nimg.1998.0395

Delis DC, Kaplan E, and Kramer JH. 2001. *Delis-Kaplan Executive Function*

System:Technical Manual. San Antonio, TX: Harcourt Assessment Company.

Delis DC, Kramer JH, Kaplan E, and Ober BA. 2000. *California Verbal Learning Test-*

Second Edition (CVLT-II). San Antonio, TX: Psychological Corporation.

Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL,

Dale AM, Maguire RP, Hyman BT, Albert MS, and Killiany RJ. 2006. An automated labeling

system for subdividing the human cerebral cortex on MRI scans into gyral based regions of

interest. *Neuroimage* 31:968-980. 10.1016/j.neuroimage.2006.01.021

Doan NT, Engvig A, Persson K, Alnaes D, Kaufmann T, Rokicki J, Cordova-Palomera

A, Moberget T, Braekhus A, Barca ML, Engedal K, Andreassen OA, Selbaek G, and Westlye

LT. 2017. Dissociable diffusion MRI patterns of white matter microstructure and connectivity

in Alzheimer's disease spectrum. *Scientific Reports* 7. ARTN 45131

10.1038/srep45131

Dorum ES, Alnaes D, Kaufmann T, Richard G, Lund MJ, Tonnesen S, Sneve MH,

Mathiesen NC, Rustan OG, Gjertsen O, Vatn S, Fure B, Andreassen OA, Nordvik JE, and

Westlye LT. 2016. Age-related differences in brain network activation and co-activation

during multiple object tracking. *Brain and Behavior* 6. 10.1002/brb3.533

Dorum ES, Kaufmann T, Alnaes D, Andreassen OA, Richard G, Kolskar KK, Nordvik

JE, and Westlye LT. 2017. Increased sensitivity to age-related differences in brain functional

connectivity during continuous multiple object tracking compared to resting-state.

Neuroimage 148:364-372. 10.1016/j.neuroimage.2017.01.048

Espeseth T, Vangkilde SA, Petersen A, Dyrholm M, and Westlye LT. 2014. TVA-based

assessment of attentional capacities-associations with age and indices of brain white matter

microstructure. *Front Psychol* 5:1177. 10.3389/fpsyg.2014.01177

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A,

Killiany R, Kennedy D, Klaveness S, Montillo A, Makris N, Rosen B, and Dale AM. 2002.

Whole brain segmentation: Automated labeling of neuroanatomical structures in the human

brain. *Neuron* 33:341-355. Doi 10.1016/S0896-6273(02)00569-X

Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH, Busa E,

Seidman LJ, Goldstein J, Kennedy D, Caviness V, Makris N, Rosen B, and Dale AM. 2004.

Automatically parcellating the human cerebral cortex. *Cerebral Cortex* 14:11-22.

10.1093/cercor/bhg087

Fjell AM, Westlye LT, Grydeland H, Amlien I, Espeseth T, Reinvang I, Raz N, Holland

D, Dale AM, Walhovd KB, and Alzheimer Disease Neuroimaging I. 2013. Critical ages in the

life course of the adult brain: nonlinear subcortical aging. *Neurobiol Aging* 34:2239-2247.

10.1016/j.neurobiolaging.2013.04.006

Fox J, and Weisberg S. 2011. An {R} Companion to Applied Regression. In: Second,

editor. Thousand Oaks, CA: Sage.

Franke K, Gaser C, Manor B, and Novak V. 2013. Advanced BrainAGE in older adults

with type 2 diabetes mellitus. *Frontiers in Aging Neuroscience* 5. ARTN 90

10.3389/fnagi.2013.00090

Gaser C, Franke K, Kloppel S, Koutsouleris N, Sauer H, and Initi AsDN. 2013.

BrainAGE in Mild Cognitive Impaired Patients: Predicting the Conversion to Alzheimer's

Disease. *PLoS One* 8. ARTN e67346

10.1371/journal.pone.0067346

Ge Y, Grossman RI, Babb JS, Rabin ML, Mannon LJ, and Kolson DL. 2002. Age-

related total gray matter and white matter changes in normal adult brain. Part I: volumetric

MR imaging analysis. *AJNR Am J Neuroradiol* 23:1327-1333.

Habekost T. 2015. Clinical TVA-based studies: a general review. *Front Psychol* 6:290.

10.3389/fpsyg.2015.00290

Habekost T, and Starrfelt R. 2009. Visual attention capacity: a review of TVA-based

patient studies. *Scand J Psychol* 50:23-32. 10.1111/j.1467-9450.2008.00681.x

Habekost T, Vogel A, Rostrup E, Bundesen C, Kyllingsbaek S, Garde E, Ryberg C, and

Waldemar G. 2013. Visual processing speed in old age. *Scand J Psychol* 54:89-94.

10.1111/sjop.12008

Habes M, Guray E, Toledo JB, Zhang T, Bryan RN, Janowitz D, Doshi J, von

Sarnowski B, Hegenscheid K, Voelzke H, Schminke U, Hoffmann W, Grabe HJ, and

Davatzikos C. 2016. Non-Resilient Brain Aging in Association with Cardiovascular Risk and

White Matter Hyperintensities: The Ship Study. *Alzheimer's & Dementia* 12:P226-P227.

10.1016/j.jalz.2016.06.407

Han CE, Peraza LR, Taylor JP, and Kaiser M. 2014. Predicting Age across Human

Lifespan Based on Structural Connectivity from Diffusion Tensor Imaging. *2014 Ieee*

Biomedical Circuits and Systems Conference (Biocas):137-140.

Harada CN, Natelson Love MC, and Triebel KL. 2013. Normal cognitive aging. *Clin*

Geriatr Med 29:737-752. 10.1016/j.cger.2013.07.002

Hastie T. 2017. gam: Generalized Additive Models.

Hua K, Zhang JY, Wakana S, Jiang HY, Li X, Reich DS, Calabresi PA, Pekar JJ, van

Zijl PCM, and Mori S. 2008. Tract probability maps in stereotaxic spaces: Analyses of white

matter anatomy and tract-specific quantification. *Neuroimage* 39:336-347.

10.1016/j.neuroimage.2007.07.053

Kaufmann T, van der Meer D, Doan NT, Schwarz E, Lund MJ, Agartz I, Alnæs D,

Barch DM, Baur-Streubel R, Bertolino A, Bettella F, Beyer MK, Bøen E, Borgwardt S,

Brandt CL, Buitelaar J, Celius EG, Cervenka S, Conzelmann A, Córdova-Palomera A, Dale

AM, de Quervain DJF, Di Carlo P, Djurovic S, Dørum ES, Eisenacher S, Elvsashagen T,

Espeseth T, Fatouros-Bergman H, Flyckt L, Franke B, Frei O, Haatveit B, Haberg AK, Harbo

HF, Hartman CA, Heslenfeld D, Hoekstra PJ, Høgestøl EA, Jernigan T, Jonassen R, Jönsson

EG, Kirsch P, Kloszewska I, Kolskar K-K, Landrø NI, Le Hellard S, Lesch K-P, Lovestone S,

Lundervold A, Lundervold AJ, Maglanoc LA, Malt UF, Mecocci P, Melle I, Meyer-

Lindenberg A, Moberget T, Norbom LB, Nordvik JE, Nyberg L, Oosterlaan J, Papalino M,

Papassotiropoulos A, Pauli P, Pergola G, Persson K, Richard G, Rokicki J, Sanders A-M,

Selbæk G, Shadrin AA, Smeland OB, Soininen H, Sowa P, Steen VM, Tsolaki M, Ulrichsen

KM, Vellas B, Wang L, Westman E, Ziegler GC, Zink M, Andreassen OA, and Westlye LT.

2018. Genetics of brain age suggest an overlap with common brain disorders. *bioRxiv*.

Kuhn T, Kaufmann T, Doan NT, Westlye LT, Jones J, Nunez RA, Bookheimer SY,

Singer EJ, Hinkin CH, and Thames AD. 2018. An augmented aging process in brain white

matter in HIV. *Hum Brain Mapp*. 10.1002/hbm.24019

Liem F, Varoquaux G, Kynast J, Beyer F, Masouleh SK, Huntenburg JM, Lampe L,

Rahim M, Abraham A, Craddock RC, Riedel-Heller S, Luck T, Loeffler M, Schroeter ML,

Witte AV, Villringer A, and Margulies DS. 2017. Predicting brain-age from multimodal

imaging data captures cognitive impairment. *Neuroimage* 148:179-188.

10.1016/j.neuroimage.2016.11.005

Liu K, Yao S, Chen K, Zhang J, Yao L, Li K, Jin Z, and Guo X. 2017. Structural Brain

Network Changes across the Adult Lifespan. *Front Aging Neurosci* 9:275.

10.3389/fnagi.2017.00275

Madan CR, and Kensinger EA. 2018. Predicting age from cortical structure across the

lifespan. *Eur J Neurosci* 47:399-416. 10.1111/ejn.13835

Marquand AF, Rezek I, Buitelaar J, and Beckmann CF. 2016. Understanding

Heterogeneity in Clinical Cohorts Using Normative Models: Beyond Case-Control Studies.

Biol Psychiatry 80:552-561. 10.1016/j.biopsych.2015.12.023

McAvinue LP, Habekost T, Johnson KA, Kyllingsbaek S, Vangkilde S, Bundesen C,

and Robertson IH. 2012. Sustained attention, attentional selectivity, and attentional capacity

across the lifespan. *Atten Percept Psychophys* 74:1570-1582. 10.3758/s13414-012-0352-6

Mori S, Wakana S, van Zijl PCM, and Nagae-Poetscher LM. 2005. MRI Atlas of

Human White Matter. In: Science E, editor. 1st Edition ed: Elsevier Science. p 276.

Müllner D. 2013. {fastcluster}: Fast Hierarchical, Agglomerative Clustering Routines

for {R} and {Python}. *Journal of Statistical Software* 53:1-18.

Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I,

Cummings JL, and Chertkow H. 2005. The montreal cognitive assessment, MoCA: A brief

screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*

53:695-699. DOI 10.1111/j.1532-5415.2005.53221.x

Raz N, Ghisletta P, Rodrigue KM, Kennedy KM, and Lindenberger U. 2010.

Trajectories of brain aging in middle-aged and older adults: regional and individual

differences. *Neuroimage* 51:501-511. 10.1016/j.neuroimage.2010.03.020

Ronan L, Alexander-Bloch AF, Wagstyl K, Farooqi S, Brayne C, Tyler LK, Fletcher

PC, and Cam-CAN. 2016. Obesity associated with increased brain age from midlife.

Neurobiology of Aging 47:63-70. 10.1016/j.neurobiolaging.2016.07.010

Salat DH, Tuch DS, Greve DN, van der Kouwe AJW, Hevelone ND, Zaleta AK, Rosen

BR, Fischl B, Corkin S, Rosas HD, and Dale AM. 2005. Age-related alterations in white

matter microstructure measured by diffusion tensor imaging. *Neurobiology of Aging* 26:1215-

1227. 10.1016/j.neurobiolaging.2004.09.017

Schnack HG, van Haren NE, Nieuwenhuis M, Hulshoff Pol HE, Cahn W, and Kahn RS.

2016. Accelerated Brain Aging in Schizophrenia: A Longitudinal Pattern Recognition Study.

Am J Psychiatry 173:607-616. 10.1176/appi.ajp.2015.15070922

Shafto MA, Tyler LK, Dixon M, Taylor JR, Rowe JB, Cusack R, Calder AJ, Marslen-

Wilson WD, Duncan J, Dalgleish T, Henson RN, Brayne C, Matthews FE, and Cam-CAN.

2014. The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) study protocol: a

cross-sectional, lifespan, multidisciplinary examination of healthy cognitive ageing. *Bmc*

Neurology 14. ARTN 204

10.1186/s12883-014-0204-1

Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE,

Watkins KE, Ciccarelli O, Cader MZ, Matthews PM, and Behrens TEJ. 2006. Tract-based

spatial statistics: Voxelwise analysis of multi-subject diffusion data. *Neuroimage* 31:1487-

1505. 10.1016/j.neuroimage.2006.02.024

Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg

H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J,

Zhang YY, De Stefano N, Brady JM, and Matthews PM. 2004. Advances in functional and

structural MR image analysis and implementation as FSL. *Neuroimage* 23:S208-S219.

10.1016/j.neuroimage.2004.07.051

Sperling G. 1960. The information available in brief visual presentations. *Psychological*

Monographs: General and Applied 74:1-29. 10.1037/h0093759

Steffener J, Habeck C, O'Shea D, Razlighi Q, Bherer L, and Stern Y. 2016. Differences

between chronological and brain age are related to education and self-reported physical

activity. *Neurobiology of Aging* 40:138-144. 10.1016/j.neurobiolaging.2016.01.014

Storsve AB, Fjell AM, Tamnes CK, Westlye LT, Overbye K, Aasland HW, and

Walhovd KB. 2014. Differential Longitudinal Changes in Cortical Thickness, Surface Area

and Volume across the Adult Life Span: Regions of Accelerating and Decelerating Change.

Journal of Neuroscience 34:8488-8498. 10.1523/Jneurosci.0391-14.2014

Taylor JR, Williams N, Cusack R, Auer T, Shafto MA, Dixon M, Tyler LK, Cam C,

and Henson RN. 2017. The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) data

repository: Structural and functional MRI, MEG, and cognitive data from a cross-sectional

adult lifespan sample. *Neuroimage* 144:262-269. 10.1016/j.neuroimage.2015.09.018

Valizadeh SA, Hanggi J, Merillat S, and Jancke L. 2017. Age prediction on the basis of

brain anatomical measures. *Hum Brain Mapp* 38:997-1008. 10.1002/hbm.23434

Wakana S, Caprihan A, Panzenboeck MM, Fallon JH, Perry M, Gollub RL, Hua KG,

Zhang JY, Jiang HY, Dubey P, Blitz A, van Zijl P, and Mori S. 2007. Reproducibility of

quantitative tractography methods applied to cerebral white matter. *Neuroimage* 36:630-644.

10.1016/j.neuroimage.2007.02.049

Warnes GR, Bolker B, Bonebakker L, Gentleman R, Liaw WHA, Lumley T, Maechler

M, Magnusson A, Moeller S, Schwartz M, and Venables B. 2016. gplots: Various R

Programming Tools for Plotting Data.

Wechsler D. 1999. *Wechsler Abbreviated Scale of Intelligence (WASI)*: Psychological

Corporation.

Westlye LT, Walhovd KB, Dale AM, Bjornerud A, Due-Tonnessen P, Engvig A,

Grydeland H, Tamnes CK, Ostby Y, and Fjell AM. 2010a. Differentiating maturational and

aging-related changes of the cerebral cortex by use of thickness and signal intensity.

Neuroimage 52:172-185. 10.1016/j.neuroimage.2010.03.056

739   Westlye LT, Walhovd KB, Dale AM, Bjornerud A, Due-Tonnessen P, Engvig A,

740   Grydeland H, Tamnes CK, Ostby Y, and Fjell AM. 2010b. Life-Span Changes of the Human

741   Brain White Matter: Diffusion Tensor Imaging (DTI) and Volumetry. *Cerebral Cortex*

742   20:2055-2068. 10.1093/cercor/bhp280

743   Wiegand I, Lauritzen MJ, Osler M, Mortensen EL, Rostrup E, Rask L, Richard N,

744   Horwitz A, Benedek K, Vangkilde S, and Petersen A. 2018. EEG correlates of visual short-

745   term memory in older age vary with adult lifespan cognitive development. *Neurobiol Aging*

746   62:210-220. 10.1016/j.neurobiolaging.2017.10.018

747   Willer L, Pedersen PM, Forchhammer HB, and Christensen HK. 2016. Cognitive

748   assessment at bedside for iPad: A preliminary validation of a novel cognitive test for stroke

749   patients. *European Stroke Journal* 1:294-301. 10.1177/2396987316665233

750   Wolfers T, Doan NT, Kaufmann T, Alnæs D, Moberget T, Agartz I, Buitelaar J, Ueland

751   T, Melle I, Beckmann CF, Franke B, Andreassen OA, Westlye LT, and Marquand A. in press.

752   Extensive interindividual differences in schizophrenia and bipolar disorder: mapping

753   biological heterogeneity in reference to normative brain ageing *JAMA Psychiatry*.

754   Wright SP. 1992. Adjusted P-Values for Simultaneous Inference. *Biometrics* 48:1005-

755   1013. Doi 10.2307/2532694

Figure legends

Fig. 1 Histogram of the age distribution for each sample.

Fig. 2 Comparison between the 11 BAG models. (a) Heatmap of the correlation between

different BAGs. (b) Correlations between the chronological age and the predicted age in the

test sample for each model with their confidence intervals. (c) Mean and standard error of the

45 p-values ($-\log_{10}(p)$) for the cognitive scores and composite scores for each row (i.e.

BAGs), with a higher mean representing a stronger global association across tests. (d)

Correlation between the chronological age of each subjects and the combined age, (e) the

brain morphometry age, and (f) the white matter microstructure age.

Fig. 3 Hierarchical clustering of the cognitive features. Each cognitive score was normalized

and when required the scores were multiplied by -1 to ensure that positive scores represent

good performance. The higher panel shows the dendrogram resulting from the hierarchical

clustering of the scores in 7 cognitive domains.

Fig. 4 Heatmap of the association between cognitive scores and brain age gaps. The color

scale depicts the minus log of the p-values ($-\log_{10}(p)$) for each association. The association

marked with a small star represents significant associations after FDR correction, and the one

marked with a big star shows significant associations after Bonferroni correction.

Fig. 5 Scatter plots of the 2 strongest associations between cognitive measures and BAG. The

color gradient represents the age where lighter color is assigned to older individuals, and

darker color to younger individuals. All associations indicate worse performance with higher

brain age gap.

Table legends

Table 1. Demographics and cognitive information. * significant associations between

cognitive measures with age after FDR correction, ** significant associations between

cognitive measures with age after Bonferroni correction

Table 2. Cognitive associations with Brain Age Gap (BAG) – statistics. * FDR significant **

Bonferroni significant

Supplementary Materials

Fig. S1. Heatmap of the association between cognitive scores and brain age gaps using non-

linear models, including age, age^2 and sex as covariates. The color scale depicts the minus log

of the p-values ($-\log_{10}(p)$) for each association. The association marked with a small star

represents significant associations after FDR correction, and the one marked with a big star

shows significant associations after Bonferroni correction.

Table S1. Cognitive associations with Brain Age Gap (BAG) using non-linear models,

including age, age^2 and sex as covariates – statistics. * FDR significant ** Bonferroni

significant.

835 **Tables**

836 Demographics and cognitive information

	Cam-CAN	StrokeMRI Mean (SD)	Range (IQR)	Main effect Age *t (p)*	Main effect Sex *t (p)*
Total N (% females)	612 (51.3%)	265 (63.4%)			
Mean age (SD)	54.41 (18.26)	56.95 (14.84)			
Age range	18-87	20-88			
MoCA	-	27.60 (1.72)	21 – 30 (2)	-4.57 (<0.001)**	-2.32 (0.021)
WASI words	-	65.27 (6.60)	44 – 79 (10)	4.72 (<0.001)**	0.10 (0.920)
WASI matrix	-	25.39 (5.64)	7 – 35 (6)	-7.60 (<0.001)**	-0.28 (0.776)
CVLT learning 1-5	-	48.92 (11.37)	17 – 73 (15.5)	-5.05 (<0.001)**	-5.26 (<0.001)
CVLT interference	-	5.53 (2.15)	0 – 13 (3)	-4.33 (<0.001)**	-0.41 (0.681)
CVLT recall	-	10.83 (3.42)	0 – 16 (5)	-6.50 (<0.001)**	5.94 (<0.001)
CVLT delayed recall	-	11.39 (3.44)	0 – 16 (5)	-4.97 (<0.001)**	-5.51 (<0.001)
CVLT recognition hit	-	14.70 (1.50)	8 – 16 (2)	-2.62 (0.0093)*	-2.68 (0.008)
CVLT recognition errors	-	3.79 (3.92)	0 – 18 (4)	5.22 (<0.001)**	4.18 (<0.001)
CVLT recog misses	-	1.30 (1.49)	0 – 8 (2)	2.62 (0.0093)*	2.68 (0.008)
CVLT recog false alarm	-	2.46 (3.48)	0 – 18 (3)	4.45 (<0.001)**	3.59 (0.0004)
CVLT recog correct rejection	-	44.20 (3.92)	30 – 48 (4)	-5.22 (<0.001)**	-4.18 (<0.001)
CVLT d'	-	2.97 (0.72)	0.97 – 3.90 (1.11)	-5.01 (<0.001)**	-4.50 (<0.001)
STROOP 1	-	31.14 (5.66)	21 – 50 (7)	5.05 (<0.001)**	2.44 (0.015)
STROOP 2	-	22.12 (3.49)	14 – 35 (4)	2.89 (0.004)*	2.27 (0.024)
STROOP 3	-	55.86 (14.13)	10 – 108 (15)	7.55 (<0.001)**	2.97 (0.003)
STROOP 4	-	61.74 (14.85)	33 – 117 (19)	7.51 (<0.001)**	1.77 (0.078)
STROOP mean 1 and 2	-	26.54 (4.16)	18.5 – 42 (5)	4.47 (<0.001)**	2.47 (0.014)
STROOP 3 minus mean 1 and 2	-	81.94 (16.51)	34.5 – 145 (18.5)	7.31 (<0.001)**	3.02 (0.003)
STROOP 4 minus mean 1 and 2	-	87.64 (16.73)	53.5 – 142 (24)	7.52 (<0.001)**	1.85 (0.066)
CP – Right motor speed	-	79.56 (23.34)	34 – 153 (32)	-12.25 (<0.001)**	-0.36 (0.716)
CP – Left motor speed	-	81.36 (17.80)	39 – 131 (26)	-12.07 (<0.001)**	0.20 (0.842)
CP – FAS Phonological flow	-	54.70 (14.53)	14 – 95 (19.75)	-0.61 (0.541)	-2.58 (0.011)
CP – FAS Semantic flow	-	51.00 (10.14)	27 – 81 (13)	-2.93 (0.004)*	-3.93 (<0.001)

CP – Visual WM forward ls	-	4.23 (1.01)	2 – 7 (2)	-5.31 (<0.001)**	0.29 (0.774)
CP – Visual WM forward ss	-	5.45 (1.87)	1 – 10 (3)	-6.59 (<0.001)**	-0.25 (0.803)
CP – Visual WM backward ls	-	3.80 (1.28)	0 – 8 (1)	-4.60 (<0.001)**	-1.85 (0.065)
CP – Visual WM backward ss	-	4.56 (2.08)	0 – 12 (3)	-5.48 (<0.001)**	-1.02 (0.309)
CP – Visual WM ss	-	9.96 (3.57)	1 – 21 (4)	-7.04 (<0.001)**	-0.95 (0.342)
CP – Spatial stroop congruent (ms)	-	674.42 (132.77)	410 – 1159 (181)	8.52 (<0.001)**	-1.03 (0.304)
CP – Spatial stroop incongruent (ms)	-	929.52 (198.01)	462 – 1827 (269)	9.41 (<0.001)**	-0.75 (0.451)
CP – Spatial stroop Errors	-	2.17 (2.41)	0 – 11 (3)	0.73 (0.463)	1.59 (0.113)
CP – Spatial stroop numb of reps	-	119.63 (16.64)	55 – 166 (22)	-9.67 (<0.001)**	1.23 (0.219)
CP – Spatial stroop incong – cong (ms)	-	252 (110)	20 – 678 (134.5)	5.73 (<0.001)**	-0.68 (0.498)
CP – Spatspan ls	-	5.37 (1.78)	1 – 10 (2)	-9.12 (<0.001)**	-4.88 (<0.001)
CP – Spatspan tot	-	29.87 (12.43)	3 – 55 (18)	-9.28 (<0.001)**	-4.66 (<0.001)
CP – Coding corr	-	54.50 (12.11)	24 – 88 (16)	-16.69 (<0.001)**	-2.46 (0.015)
CP – Coding error	-	0.67 (0.99)	0 – 5 (1)	-1.10 (0.271)	1.56 (0.121)
TVA – Short-term memory storage (K)	-	3.38 (0.77)	1.46 – 5.53 (1.09)	-7.75 (<0.001)**	-1.52 (0.129)
TVA – Processing speed (C)	-	31.55 (14.07)	5.99 – 89.67 (14.75)	-4.69 (<0.001)**	0.41 (0.6847)
TVA – Perceptual threshold (t_0)	-	23.01 (14.05)	0 – 79.75 (17.59)	5.72 (<0.001)**	-1.94 (0.053)
TVA – Error rate	-	0.10 (0.06)	0.0035 – 0.3316 (0.0983)	-1.35 (0.177)	0.67 (0.502)
Cluster 1	-	-	-	-7.19 (<0.001)**	-5.16 (<0.001)
Cluster 2	-	-	-	-7.28 (<0.001)**	1.61 (0.110)
Cluster 3	-	-	-	-2.01 (0.045)*	-3.99 (<0.001)
Cluster 4	-	-	-	-9.98 (<0.001)**	1.25 (0.212)
Cluster 5	-	-	-	-6.86 (<0.001)**	-2.56 (0.011)
Cluster 6	-	-	-	-15.79 (<0.001)**	-1.08 (0.282)
Cluster 7	-	-	-	-6.50 (<0.001)**	-0.77 (0.440)

837 Table 1. Demographics and cognitive information. * significant associations between

838 cognitive measures with age after FDR correction, ** significant associations between

839 cognitive measures with age after Bonferroni correction

843   Cognitive associations with Brain Age Gap (BAG) – statistics

Test	Adj R^2 no-BAG	BAG	Main effect Age t (p)	Main effect Sex t (p)	Main effect BAG t (p)	Adj R^2
MoCA	0.0907	T1	-4.5596 (<0.001)	-2.3145 (0.021)	-0.124 (0.901)	0.0878
		DTI	-4.5599 (<0.001)	-2.3155 (0.021)	1.5914 (0.113)	0.0966
		Combined	-4.5653 (<0.001)	-2.3176 (0.021)	-0.4626 (0.644)	0.0885
WASI words	0.0731	T1	4.7118 (<0.001)	0.1020 (0.919)	-0.2169 (0.828)	0.0704
		DTI	4.7056 (<0.001)	0.1121 (0.911)	-0.8126 (0.417)	0.0727
		Combined	4.7091 (<0.001)	0.1041 (0.917)	-0.4827 (0.630)	0.0711
WASI matrix	0.1791	T1	-7.6061 (<0.001)	-0.2785 (0.781)	-0.9158 (0.361)	0.1793
		DTI	-7.6610 (<0.001)	-0.2624 (0.793)	-1.6546 (0.099)	0.1854
		Combined	-7.6128 (<0.001)	-0.2726 (0.785)	-1.1102 (0.268)	0.1806
CVLT learning 1-5	0.1810	T1	-5.0373 (<0.001)	-5.2514 (<0.001)	-0.2505 (0.802)	0.1750
		DTI	-5.0418 (<0.001)	-5.2533 (<0.001)	-0.3608 (0.719)	0.1753
		Combined	-5.0387 (<0.001)	-5.2522 (<0.001)	-0.2492 (0.803)	0.1750
CVLT interference	0.0664	T1	-4.3256 (<0.001)	-0.4062 (0.685)	-0.9588 (0.339)	0.0626
		DTI	-4.3218 (<0.001)	-0.4104 (0.682)	-0.2391 (0.811)	0.0594
		Combined	-4.3202 (<0.001)	-0.4101 (0.682)	-0.1875 (0.851)	0.0594
CVLT recall	0.2438	T1	-6.4897 (<0.001)	-5.9257 (<0.001)	-0.4868 (0.627)	0.2397
		DTI	-6.4885 (<0.001)	-5.9257 (<0.001)	-0.1245 (0.901)	0.2391
		Combined	-6.5080 (<0.001)	-5.9373 (<0.001)	-1.1114 (0.268)	0.2427
CVLT delayed recall	0.1850	T1	-4.9636 (<0.001)	-5.4973 (<0.001)	0.1421 (0.887)	0.1808
		DTI	-4.9611 (<0.001)	-5.4969 (<0.001)	0.224 (0.823)	0.1809
		Combined	-4.9655 (<0.001)	-5.4954 (<0.001)	-0.3038 (0.762)	0.1810
CVLT recognition hits	0.0494	T1	-2.6125 (0.010)	-2.6822 (0.008)	-0.8586 (0.391)	0.0486
		DTI	-2.6144 (0.010)	-2.6786 (0.008)	0.0946 (0.925)	0.0459
		Combined	-2.6212 (0.009)	-2.6854 (0.008)	-1.0724 (0.285)	0.0501
CVLT recognition errors	0.1526	T1	5.2227 (<0.001)	4.1850 (<0.001)	-0.8471 (0.398)	0.1528
		DTI	5.2115 (<0.001)	4.1755 (<0.001)	-0.5651 (0.573)	0.1514
		Combined	5.2139 (<0.001)	4.1740 (<0.001)	-0.2537 (0.800)	0.1506
CVLT recog misses	0.0494	T1	2.6125 (0.010)	2.6822 (0.008)	0.8586 (0.391)	0.0486
		DTI	2.6144 (0.010)	2.6786 (0.008)	-0.0946 (0.925)	0.0459
		Combined	2.6212 (0.009)	2.6854 (0.008)	1.0724 (0.285)	0.0501
CVLT recog false alarm	0.1150	T1	4.4519 (<0.001)	3.5827 (<0.001)	-0.776 (0.439)	0.1146
		DTI	4.4378 (<0.001)	3.5803 (<0.001)	-0.5207 (0.603)	0.1134
		Combined	4.4418 (<0.001)	3.5788 (<0.001)	-0.3488 (0.728)	0.1129
CVLT recog correct rejection	0.1526	T1	-5.2227 (<0.001)	-4.1850 (<0.001)	0.8471 (0.398)	0.1528
		DTI	-5.2115 (<0.001)	-4.1755 (<0.001)	0.5651 (0.573)	0.1514
		Combined	-5.2139 (<0.001)	-4.1740 (<0.001)	0.2537 (0.800)	0.1506
CVLT d'	0.1566	T1	-5.0074 (<0.001)	-4.4914 (<0.001)	0.3628 (0.717)	0.1536
		DTI	-5.0021 (<0.001)	-4.4969 (<0.001)	0.8538 (0.394)	0.1556
		Combined	-5.0038 (<0.001)	-4.4902 (<0.001)	0.1699 (0.865)	0.1533
STROOP 1	0.1118	T1	5.1466 (<0.001)	2.4999 (0.013)	2.6939 (0.008)	0.1299

		DTI	5.0968 (<0.001)	2.4769 (0.014)	1.6664 (0.097)	0.1147
		Combined	5.2111 (<0.001)	2.5317 (0.012)	3.3767 (<0.001)*	0.1434
STROOP 2	0.0477	T1	2.8868 (0.004)	2.2619 (0.025)	0.1557 (0.876)	0.0433
		DTI	2.8768 (0.004)	2.2489 (0.025)	-0.4639 (0.643)	0.0440
		Combined	2.8949 (0.004)	2.2713 (0.024)	0.4976 (0.619)	0.0442
STROOP 3	0.2104	T1	7.5930 (<0.001)	2.9898 (0.003)	1.5092 (0.133)	0.2109
		DTI	7.6511 (<0.001)	3.0224 (0.003)	2.231 (0.027)	0.2190
		Combined	7.6793 (<0.001)	3.0233 (0.003)	2.5768 (0.011)	0.2240
STROOP 4	0.1887	T1	7.5403 (<0.001)	1.7884 (0.075)	1.2397 (0.216)	0.1906
		DTI	7.5847 (<0.001)	1.8121 (0.071)	1.7368 (0.084)	0.1953
		Combined	7.6387 (<0.001)	1.8247 (0.069)	2.3662 (0.019)	0.2033
STROOP mean 1 and 2	0.0949	T1	4.5089 (<0.001)	2.5033 (0.013)	1.5875 (0.114)	0.0978
		DTI	4.4750 (<0.001)	2.4760 (0.014)	0.3927 (0.695)	0.0894
		Combined	4.5432 (<0.001)	2.5399 (0.012)	2.0254 (0.044)	0.1034
STROOP 3 minus mean 1 and 2	0.2051	T1	7.3383 (<0.001)	3.0427 (0.003)	1.1397 (0.256)	0.2021
		DTI	7.3613 (<0.001)	3.0703 (0.002)	1.3546 (0.177)	0.2038
		Combined	7.4197 (<0.001)	3.1063 (0.002)	2.1881 (0.030)	0.2130
STROOP 4 minus mean 1 and 2	0.1936	T1	7.5360 (<0.001)	1.8671 (0.063)	0.8763 (0.382)	0.1919
		DTI	7.5297 (<0.001)	1.8697 (0.063)	0.6331 (0.527)	0.1907
		Combined	7.6081 (<0.001)	1.9215 (0.056)	1.7531 (0.081)	0.1993
CP – Right motor speed	0.3695	T1	-12.2893 (<0.001)	-0.3592 (0.720)	-1.5504 (0.122)	0.3676
		DTI	-12.2318 (<0.001)	-0.3612 (0.718)	-0.3435 (0.732)	0.3620
		Combined	-12.3125 (<0.001)	-0.3587 (0.720)	-1.8139 (0.071)	0.3697
CP – Left motor speed	0.3630	T1	-12.1437 (<0.001)	0.2100 (0.834)	-1.9945 (0.047)	0.3634
		DTI	-12.0669 (<0.001)	0.2081 (0.835)	-0.8704 (0.385)	0.3555
		Combined	-12.2516 (<0.001)	0.2149 (0.830)	-2.9047 (0.004)	0.3740
CP – FAS Semantic flow	0.0840	T1	-2.9562 (0.003)	-3.9454 (<0.001)	-2.0826 (0.038)	0.0960
		DTI	-2.9607 (0.003)	-3.9388 (<0.001)	-2.0997 (0.037)	0.0963
		Combined	-2.9513 (0.004)	-3.9389 (<0.001)	-1.8308 (0.068)	0.0926
CP – Visual WM forward ls	0.0936	T1	-5.3071 (<0.001)	0.2850 (0.776)	-0.5838 (0.560)	0.0906
		DTI	-5.3392 (<0.001)	0.2963 (0.767)	-1.7204 (0.087)	0.0999
		Combined	-5.3059 (<0.001)	0.2853 (0.776)	-0.3127 (0.755)	0.0897
CP – Visual WM forward ss	0.1416	T1	-6.5795 (<0.001)	-0.2502 (0.803)	-0.2158 (0.829)	0.1375
		DTI	-6.6000 (<0.001)	-0.2448 (0.807)	-1.1695 (0.243)	0.1420
		Combined	-6.5786 (<0.001)	-0.2496 (0.803)	-0.02 (0.984)	0.1373
CP – Visual WM backward ls	0.0852	T1	-4.5941 (<0.001)	-1.8511 (0.065)	-0.1047 (0.917)	0.0820
		DTI	-4.6170 (<0.001)	-1.8545 (0.065)	-1.3334 (0.184)	0.0884
		Combined	-4.6051 (<0.001)	-1.8550 (0.065)	-0.8013 (0.424)	0.0843
CP – Visual WM backward ss	0.1022	T1	-5.4741 (<0.001)	-1.0181 (0.310)	-0.2721 (0.786)	0.1015
		DTI	-5.4971 (<0.001)	-1.0179 (0.310)	-1.3043 (0.193)	0.1072
		Combined	-5.4898 (<0.001)	-1.0215 (0.308)	-1.0074 (0.315)	0.1048
CP – Visual WM ss	0.1607	T1	-7.0322 (<0.001)	-0.9515 (0.342)	-0.3013 (0.763)	0.1591
		DTI	-7.0622 (<0.001)	-0.9511 (0.342)	-1.3634 (0.174)	0.1649

		Combined	-7.0399 (<0.001)	-0.9528 (0.342)	-0.6665 (0.506)	0.1603
CP – Spatial stroop congruent	0.2288	T1	8.6156 (<0.001)	-1.0080 (0.314)	2.1921 (0.029)	0.2288
		DTI	8.6687 (<0.001)	-1.0021 (0.317)	2.6995 (0.007)	0.2362
		Combined	8.8278 (<0.001)	-0.9828 (0.327)	3.9007 (<0.001)**	0.2588
CP – Spatial stroop incongruent	0.2548	T1	9.5489 (<0.001)	-0.7429 (0.458)	2.6569 (0.008)	0.2700
		DTI	9.5931 (<0.001)	-0.7587 (0.449)	2.8817 (0.004)	0.2735
		Combined	9.7197 (<0.001)	-0.7378 (0.461)	3.8071 (<0.001)**	0.2903
CP – Spatial stroop numb of reps	0.2731	T1	-9.7755 (<0.001)	1.2211 (0.223)	-2.2212 (0.027)	0.2753
		DTI	-9.8507 (<0.001)	1.2328 (0.219)	-2.9614 (0.003)	0.2859
		Combined	-9.9891 (<0.001)	1.2198 (0.224)	-3.8816 (<0.001)**	0.3027
CP – Spatial stroop incong – cong	0.1012	T1	5.7663 (<0.001)	-0.6595 (0.510)	1.5611 (0.120)	0.1134
		DTI	5.7466 (<0.001)	-0.6678 (0.505)	0.9705 (0.333)	0.1081
		Combined	5.7568 (<0.001)	-0.6584 (0.511)	1.2056 (0.229)	0.1099
CP – Spatspan ls	0.3055	T1	-9.1038 (<0.001)	-4.8656 (<0.001)	-0.032 (0.975)	0.3009
		DTI	-9.1746 (<0.001)	-4.9104 (<0.001)	-1.5749 (0.117)	0.3077
		Combined	-9.1043 (<0.001)	-4.8663 (<0.001)	-0.075 (0.940)	0.3009
CP – Spatspan total	0.3057	T1	-9.2664 (<0.001)	-4.6439 (<0.001)	0.1074 (0.915)	0.3024
		DTI	-9.3260 (<0.001)	-4.6815 (<0.001)	-1.3773 (0.170)	0.3076
		Combined	-9.2686 (<0.001)	-4.6461 (<0.001)	-0.0612 (0.951)	0.3024
CP – Coding corr	0.5387	T1	-16.7647 (<0.001)	-2.5004 (0.013)	-1.6149 (0.108)	0.5352
		DTI	-17.0893 (<0.001)	-2.5467 (0.012)	-3.3998 (<0.001)*	0.5510
		Combined	-17.0071 (<0.001)	-2.5604 (0.011)	-3.0056 (0.003)	0.5467
TVA - Short-term memory storage (K)	0.2013	T1	-7.7691 (<0.001)	-1.5196 (0.130)	-1.1179 (0.265)	0.1981
		DTI	-7.8117 (<0.001)	-1.5383 (0.125)	-2.0302 (0.043)	0.2070
		Combined	-7.7525 (<0.001)	-1.5195 (0.130)	-0.9537 (0.341)	0.1970
TVA - Perceptual threshold (t_0)	0.0764	T1	5.7303 (<0.001)	-1.9470 (0.053)	0.9617 (0.337)	0.1141
		DTI	5.7333 (<0.001)	-1.9444 (0.053)	1.1066 (0.270)	0.1152
		Combined	5.7523 (<0.001)	-1.9587 (0.051)	1.8346 (0.068)	0.1226
TVA - Processing speed (C)	0.1304	T1	-4.6692 (<0.001)	0.3969 (0.692)	0.8093 (0.419)	0.0723
		DTI	-4.6800 (<0.001)	0.4053 (0.686)	0.1402 (0.889)	0.0699
		Combined	-4.6827 (<0.001)	0.3944 (0.694)	0.8916 (0.374)	0.0728
Cluster 1	0.2470	T1	-7.1741 (<0.001)	-5.1567 (<0.001)	-0.1927 (0.847)	0.2440
		DTI	-7.1623 (<0.001)	-5.1410 (<0.001)	0.3683 (0.713)	0.2443
		Combined	-7.1805 (<0.001)	-5.1641 (<0.001)	-0.3879 (0.699)	0.2443
Cluster 2	0.1720	T1	-7.2680 (<0.001)	1.6030 (0.110)	-0.1013 (0.919)	0.1687
		DTI	-7.2785 (<0.001)	1.6062 (0.110)	-0.6549 (0.513)	0.1701
		Combined	-7.2740 (<0.001)	1.6104 (0.109)	-0.6382 (0.524)	0.1700
Cluster 3	0.0698	T1	-2.0177 (0.045)	-3.9824 (<0.001)	-0.8103 (0.419)	0.0686
		DTI	-2.0337 (0.043)	-3.9969 (<0.001)	-1.84 (0.067)	0.0783
		Combined	-2.0185 (0.045)	-3.9877 (<0.001)	-0.9765 (0.330)	0.0697
Cluster 4	0.2783	T1	-10.1319 (<0.001)	1.2314 (0.219)	-2.5436 (0.012)	0.2937
		DTI	-10.1479 (<0.001)	1.2377 (0.217)	-2.5207 (0.012)	0.2933
		Combined	-10.3013 (<0.001)	1.2196 (0.224)	-3.6163 (<0.001)*	0.3113

Cluster 5	0.1772	T1	-6.8872 (<0.001)	-2.5902 (0.010)	-1.1084 (0.269)	0.1779
		DTI	-6.8667 (<0.001)	-2.5805 (0.010)	-0.5825 (0.561)	0.1750
		Combined	-6.9577 (<0.001)	-2.6481 (0.009)	-1.9103 (0.057)	0.1858
Cluster 6	0.5092	T1	-15.9345 (<0.001)	-1.1148 (0.266)	-1.8971 (0.059)	0.5145
		DTI	-15.9719 (<0.001)	-1.1080 (0.269)	-2.0875 (0.038)	0.5160
		Combined	-16.0156 (<0.001)	-1.1196 (0.264)	-2.459 (0.015)	0.5193
Cluster 7	0.1399	T1	-6.4852 (<0.001)	-0.7736 (0.440)	-0.3433 (0.732)	0.1369
		DTI	-6.5210 (<0.001)	-0.7689 (0.443)	-1.6007 (0.111)	0.1452
		Combined	-6.4926 (<0.001)	-0.7759 (0.439)	-0.63 (0.529)	0.1379

844 Table 2. Cognitive associations with BAG – statistics. * FDR significant ** Bonferroni

845 significant.

Supplementary Figures

Fig. S1. Heatmap of the association between cognitive scores and brain age gaps using non-

linear models, including age, age^2 and sex as covariates. The color scale depicts the minus log

of the p-values (-$\log_{10}(p)$) for each association. The association marked with a small star

represents significant associations after FDR correction, and the one marked with a big star

shows significant associations after Bonferroni correction.

889 **Supplementary Tables**

890 Cognitive associations with Brain Age Gap (BAG) using non-linear models, including age,

891 age^2 and sex as covariates – statistics.

Test	Adj R^2 no-BAG	BAG	Main effect Age t(p)	Main effect Age^2 t(p)	Main effect Sex t(p)	Main effect BAG t(p)	Adj R^2
MoCA	0.088	T1	-4.5563 (<0.001)	-0.2752 (0.7834)	-2.1792 (0.0302)	-0.1146 (0.9088)	0.0845
		DTI	-4.5575 (<0.001)	-0.3078 (0.7585)	-2.1727 (0.0307)	1.5937 (0.1122)	0.0934
		Combined	-4.561 (<0.001)	-0.2413 (0.8095)	-2.1894 (0.0295)	-0.4399 (0.6604)	0.0851
WASI words	0.0792	T1	4.7079 (<0.001)	-1.5614 (0.1197)	0.4698 (0.6389)	-0.1719 (0.8637)	0.0756
		DTI	4.7017 (<0.001)	-1.5532 (0.1216)	0.4775 (0.6334)	-0.7843 (0.4336)	0.0778
		Combined	4.7058 (<0.001)	-1.5325 (0.1266)	0.4655 (0.642)	-0.3554 (0.7226)	0.076
WASI matrix	0.1825	T1	-7.6085 (<0.001)	-1.3223 (0.1873)	0.048 (0.9618)	-0.8787 (0.3804)	0.1817
		DTI	-7.6627 (<0.001)	-1.311 (0.1911)	0.0606 (0.9517)	-1.6227 (0.1059)	0.1876
		Combined	-7.6132 (<0.001)	-1.2678 (0.206)	0.0407 (0.9676)	-1.0117 (0.3126)	0.1825
CVLT learning 1-5	0.1757	T1	-5.0181 (<0.001)	0.5198 (0.6036)	-5.2162 (<0.001)	-0.2662 (0.7903)	0.1727
		DTI	-5.0228 (<0.001)	0.5204 (0.6032)	-5.2181 (<0.001)	-0.3723 (0.71)	0.1729
		Combined	-5.0196 (<0.001)	0.5363 (0.5922)	-5.2205 (<0.001)	-0.2959 (0.7675)	0.1727
CVLT interference	0.0593	T1	-4.3175 (<0.001)	-0.062 (0.9506)	-0.3788 (0.7051)	-0.9545 (0.3407)	0.0589
		DTI	-4.3143 (<0.001)	-0.086 (0.9316)	-0.3772 (0.7064)	-0.2366 (0.8132)	0.0558
		Combined	-4.3124 (<0.001)	-0.075 (0.9403)	-0.3794 (0.7047)	-0.1796 (0.8576)	0.0557
CVLT recall	0.2401	T1	-6.4676 (<0.001)	0.6091 (0.543)	-5.8925 (<0.001)	-0.5048 (0.6141)	0.2379
		DTI	-6.4667 (<0.001)	0.5965 (0.5514)	-5.8892 (<0.001)	-0.1381 (0.8902)	0.2371
		Combined	-6.4859 (<0.001)	0.6973 (0.4863)	-5.9255 (<0.001)	-1.1682 (0.2438)	0.2412
CVLT delayed recall	0.1822	T1	-4.9433 (<0.001)	0.6624 (0.5083)	-5.4919 (<0.001)	0.1193 (0.9052)	0.179
		DTI	-4.9411 (<0.001)	0.6618 (0.5087)	-5.4912 (<0.001)	0.2079 (0.8354)	0.1791
		Combined	-4.9454 (<0.001)	0.6981 (0.4857)	-5.4986 (<0.001)	-0.3668 (0.7141)	0.1794
CVLT recognition hits	0.0464	T1	-2.615 (0.0095)	-0.3694 (0.7121)	-2.5144 (0.0125)	-0.8444 (0.3992)	0.0453
		DTI	-2.6175 (0.0094)	-0.3994 (0.6899)	-2.504 (0.0129)	0.1039 (0.9173)	0.0427
		Combined	-2.622 (0.0093)	-0.3004 (0.7641)	-2.5328 (0.0119)	-1.0382 (0.3002)	0.0467
CVLT recognition errors	0.1567	T1	5.2609 (<0.001)	1.4118 (0.1592)	3.7395 (<0.001)	-0.8964 (0.3709)	0.1561
		DTI	5.2485 (<0.001)	1.3947 (0.1643)	3.7336 (<0.001)	-0.5993 (0.5495)	0.1546
		Combined	5.2514 (<0.001)	1.4099 (0.1598)	3.7277 (<0.001)	-0.3837 (0.7015)	0.1539
CVLT recog misses	0.0464	T1	2.615 (0.0095)	0.3694 (0.7121)	2.5144 (0.0125)	0.8444 (0.3992)	0.0453
		DTI	2.6175 (0.0094)	0.3994 (0.6899)	2.504 (0.0129)	-0.1039 (0.9173)	0.0427
		Combined	2.622 (0.0093)	0.3004 (0.7641)	2.5328 (0.0119)	1.0382 (0.3002)	0.0467
CVLT recog false alarm	0.1191	T1	4.4883 (<0.001)	1.4053 (0.1612)	3.1557 (0.0018)	-0.8236 (0.411)	0.1179
		DTI	4.4728 (<0.001)	1.3906 (0.1656)	3.1567 (0.0018)	-0.5544 (0.5798)	0.1167
		Combined	4.4784 (<0.001)	1.4159 (0.158)	3.1475 (0.0018)	-0.4786 (0.6326)	0.1164
CVLT recog correct rejection	0.1567	T1	-5.2609 (<0.001)	-1.4118 (0.1592)	-3.7395 (<0.001)	0.8964 (0.3709)	0.1561
		DTI	-5.2485 (<0.001)	-1.3947 (0.1643)	-3.7336 (<0.001)	0.5993 (0.5495)	0.1546
		Combined	-5.2514 (<0.001)	-1.4099 (0.1598)	-3.7277 (<0.001)	0.3837 (0.7015)	0.1539
CVLT d′	0.1548	T1	-5.0157 (<0.001)	-0.7018 (0.4835)	-4.1938 (<0.001)	0.3855 (0.7002)	0.1519

		DTI	-5.0104 (<0.001)	-0.7106 (0.478)	-4.1973 (<0.001)	0.8695 (0.3854)	0.1539
		Combined	-5.0122 (<0.001)	-0.7079 (0.4796)	-4.1897 (<0.001)	0.2343 (0.8149)	0.1516
STROOP 1	0.1181	T1	5.2178 (<0.001)	1.8817 (0.061)	1.9786 (0.0489)	2.6477 (0.0086)	0.1385
		DTI	5.1698 (<0.001)	1.915 (0.0566)	1.9488 (0.0524)	1.6361 (0.103)	0.1239
		Combined	5.2707 (<0.001)	1.7146 (0.0876)	2.0444 (0.0419)	3.243 (0.0013)	0.1499
STROOP 2	0.0565	T1	2.955 (0.0034)	1.8769 (0.0617)	1.7273 (0.0853)	0.0898 (0.9286)	0.0528
		DTI	2.9455 (0.0035)	1.897 (0.059)	1.7093 (0.0886)	-0.5228 (0.6016)	0.0538
		Combined	2.9595 (0.0034)	1.8439 (0.0664)	1.7389 (0.0833)	0.3307 (0.7411)	0.0531
STROOP 3	0.2519	T1	7.9387 (<0.001)	4.0215 (<0.001)**	2.0391 (0.0425)	1.4584 (0.146)	0.2552
		DTI	8.0033 (<0.001)	4.0506 (<0.001)**	2.0666 (0.0398)	2.245 (0.0256)	0.2635
		Combined	8.0064 (<0.001)	3.9001 (<0.001)**	2.0915 (0.0375)	2.3542 (0.0193)	0.265
STROOP 4	0.2092	T1	7.727 (<0.001)	2.7144 (0.0071)	1.1078 (0.269)	1.1959 (0.2329)	0.2105
		DTI	7.7745 (<0.001)	2.7322 (0.0067)	1.1277 (0.2605)	1.7316 (0.0846)	0.2154
		Combined	7.8094 (<0.001)	2.5945 (0.01)	1.1659 (0.2448)	2.2006 (0.0287)	0.221
STROOP mean 1 and 2	0.1065	T1	4.6014 (<0.001)	2.1809 (0.0301)	1.8851 (0.0606)	1.5137 (0.1314)	0.1111
		DTI	4.5693 (<0.001)	2.2194 (0.0274)	1.8487 (0.0657)	0.3205 (0.7489)	0.1033
		Combined	4.6255 (<0.001)	2.0629 (0.0402)	1.9411 (0.0534)	1.8349 (0.0677)	0.1148
STROOP 3 minus mean 1 and 2	0.2486	T1	7.6928 (<0.001)	4.0734 (<0.001)**	2.0223 (0.0442)	1.0347 (0.3018)	0.2488
		DTI	7.7161 (<0.001)	4.0759 (<0.001)**	2.0485 (0.0416)	1.2741 (0.2038)	0.2504
		Combined	7.7517 (<0.001)	3.9419 (<0.001)**	2.0994 (0.0368)	1.8804 (0.0612)	0.2561
STROOP 4 minus mean 1 and 2	0.2225	T1	7.7912 (<0.001)	3.224 (0.0014)*	1.0274 (0.3052)	0.7875 (0.4317)	0.2213
		DTI	7.7853 (<0.001)	3.233 (0.0014)*	1.0273 (0.3053)	0.5573 (0.5778)	0.2203
		Combined	7.8437 (<0.001)	3.1135 (0.0021)*	1.0945 (0.2748)	1.4951 (0.1362)	0.2263
CP - Right motor speed	0.3621	T1	-12.2583 (<0.001)	0.4241 (0.6718)	-0.4499 (0.6532)	-1.56 (0.12)	0.3656
		DTI	-12.201 (<0.001)	0.3813 (0.7033)	-0.4415 (0.6592)	-0.3491 (0.7273)	0.3599
		Combined	-12.2818 (<0.001)	0.5259 (0.5994)	-0.4741 (0.6358)	-1.8479 (0.0658)	0.368
CP - Left motor speed	0.3586	T1	-12.1901 (<0.001)	-1.3758 (0.1701)	0.5329 (0.5946)	-1.9551 (0.0517)	0.3656
		DTI	-12.1163 (<0.001)	-1.4124 (0.159)	0.5396 (0.5899)	-0.8474 (0.3976)	0.3579
		Combined	-12.2849 (<0.001)	-1.2158 (0.2252)	0.4997 (0.6177)	-2.7995 (0.0055)	0.3752
CP - FAS semantic flow	0.1047	T1	-3.0413 (0.0026)	-2.5748 (0.0106)	-3.2544 (0.0013)	-2.0265 (0.0437)	0.1153
		DTI	-3.0473 (0.0025)	-2.6035 (0.0098)	-3.2424 (0.0013)	-2.0796 (0.0385)	0.116
		Combined	-3.0316 (0.0027)	-2.4889 (0.0134)	-3.2639 (0.0012)	-1.642 (0.1018)	0.1105
CP - Visual WM forward ls	0.0897	T1	-5.3048 (<0.001)	-0.2803 (0.7795)	0.3424 (0.7323)	-0.5645 (0.5729)	0.0873
		DTI	-5.3364 (<0.001)	-0.2747 (0.7838)	0.352 (0.7252)	-1.71 (0.0885)	0.0966
		Combined	-5.3039 (<0.001)	-0.2873 (0.7742)	0.3445 (0.7307)	-0.2836 (0.7769)	0.0864
CP - Visual WM forward ss	0.1388	T1	-6.5947 (<0.001)	-0.6439 (0.5202)	-0.0916 (0.9271)	-0.1756 (0.8608)	0.1355
		DTI	-6.6142 (<0.001)	-0.6289 (0.53)	-0.0903 (0.9281)	-1.1522 (0.2503)	0.1399
		Combined	-6.5945 (<0.001)	-0.657 (0.5118)	-0.0873 (0.9305)	0.0421 (0.9664)	0.1354
CP - Visual WM backward ls	0.0852	T1	-4.6076 (<0.001)	-0.9343 (0.351)	-1.567 (0.1184)	-0.068 (0.9458)	0.0816
		DTI	-4.6299 (<0.001)	-0.9132 (0.362)	-1.5757 (0.1163)	-1.3135 (0.1902)	0.0878
		Combined	-4.6163 (<0.001)	-0.8736 (0.3831)	-1.5842 (0.1144)	-0.7237 (0.4699)	0.0835
CP - Visual WM backward ss	0.1026	T1	-5.4768 (<0.001)	-0.619 (0.5365)	-0.8352 (0.4044)	-0.2473 (0.8049)	0.0993

		DTI	-5.4994 (<0.001)	-0.604 (0.5464)	-0.8389 (0.4023)	-1.2897 (0.1983)	0.105
		Combined	-5.4903 (<0.001)	-0.5466 (0.5852)	-0.8555 (0.3931)	-0.9558 (0.3401)	0.1023
CP - Visual WM ss	0.1618	T1	-7.0453 (<0.001)	-0.9494 (0.3433)	-0.6911 (0.4902)	-0.2639 (0.7921)	0.1588
		DTI	-7.0749 (<0.001)	-0.9354 (0.3504)	-0.6944 (0.4881)	-1.3432 (0.1804)	0.1645
		Combined	-7.051 (<0.001)	-0.9076 (0.365)	-0.7016 (0.4836)	-0.5866 (0.558)	0.1597
CP - Spatial stroop congruent	0.2152	T1	8.6103 (<0.001)	0.5061 (0.6132)	-1.099 (0.2728)	2.1677 (0.0311)	0.2265
		DTI	8.6646 (<0.001)	0.539 (0.5904)	-1.1012 (0.2719)	2.6844 (0.0078)	0.234
		Combined	8.8148 (<0.001)	0.27 (0.7874)	-1.0172 (0.3101)	3.8562 (<0.001)**	0.2561
CP - Spatial stroop incongruent	0.2498	T1	9.5215 (<0.001)	-0.3129 (0.7546)	-0.6399 (0.5228)	2.6625 (0.0083)	0.2674
		DTI	9.5663 (<0.001)	-0.2497 (0.803)	-0.6711 (0.5028)	2.8797 (0.0043)	0.2708
		Combined	9.6908 (<0.001)	-0.5285 (0.5976)	-0.5807 (0.5619)	3.8325 (<0.001)*	0.2883
CP - Spatial stroop numb of reps	0.2613	T1	-9.7578 (<0.001)	-0.1542 (0.8776)	1.2196 (0.2237)	-2.2091 (0.0281)	0.2725
		DTI	-9.8334 (<0.001)	-0.1782 (0.8587)	1.2369 (0.2173)	-2.951 (0.0035)	0.2831
		Combined	-9.967 (<0.001)	0.0905 (0.928)	1.1582 (0.2479)	-3.8672 (<0.001)**	0.2999
CP - Spatial stroop incong - cong	0.1076	T1	5.7387 (<0.001)	-0.9619 (0.337)	-0.4011 (0.6887)	1.5968 (0.1116)	0.1131
		DTI	5.7189 (<0.001)	-0.9105 (0.3635)	-0.4223 (0.6732)	0.9815 (0.3273)	0.1075
		Combined	5.7304 (<0.001)	-1.0014 (0.3176)	-0.3888 (0.6978)	1.2835 (0.2005)	0.1099
CP - Spatspan ls	0.3027	T1	-9.0803 (<0.001)	0.8225 (0.4116)	-4.9183 (<0.001)	-0.0628 (0.95)	0.3
		DTI	-9.1511 (<0.001)	0.8555 (0.3931)	-4.9703 (<0.001)	-1.5904 (0.113)	0.307
		Combined	-9.0818 (<0.001)	0.8301 (0.4073)	-4.9202 (<0.001)	-0.1448 (0.885)	0.3
CP - Spatspan total	0.3033	T1	-9.2413 (<0.001)	0.5919 (0.5545)	-4.6451 (<0.001)	0.085 (0.9324)	0.3006
		DTI	-9.3007 (<0.001)	0.6245 (0.5329)	-4.6899 (<0.001)	-1.3875 (0.1665)	0.3059
		Combined	-9.2443 (<0.001)	0.6028 (0.5472)	-4.6491 (<0.001)	-0.1118 (0.9111)	0.3006
CP - Coding corr	0.5307	T1	-16.7381 (<0.001)	-0.3279 (0.7433)	-2.3396 (0.0201)	-1.596 (0.1117)	0.5335
		DTI	-17.0629 (<0.001)	-0.3481 (0.7281)	-2.3804 (0.018)	-3.3875 (<0.001)*	0.5495
		Combined	-16.9736 (<0.001)	-0.1442 (0.8855)	-2.4411 (0.0153)	-2.976 (0.0032)	0.5449
TVA - Short-term memory storage (K)	0.1941	T1	-7.7533 (<0.001)	-0.0891 (0.9291)	-1.4529 (0.1475)	-1.1129 (0.2668)	0.1949
		DTI	-7.7958 (<0.001)	-0.0894 (0.9288)	-1.471 (0.1426)	-2.0247 (0.044)	0.2039
		Combined	-7.7352 (<0.001)	-0.0343 (0.9726)	-1.4655 (0.144)	-0.9449 (0.3456)	0.1938
TVA - Perceptual threshold (t_0)	0.1146	T1	5.7716 (<0.001)	0.9866 (0.3248)	-2.1312 (0.034)	0.9264 (0.3551)	0.1141
		DTI	5.7764 (<0.001)	1.0081 (0.3144)	-2.1342 (0.0338)	1.0951 (0.2745)	0.1153
		Combined	5.784 (<0.001)	0.8478 (0.3973)	-2.1074 (0.0361)	1.7412 (0.0829)	0.1217
TVA - Processing speed (C)	0.0699	T1	-4.66 (<0.001)	0.1122 (0.9107)	0.3556 (0.7225)	0.804 (0.4222)	0.0686
		DTI	-4.6708 (<0.001)	0.135 (0.8927)	0.3579 (0.7207)	0.1387 (0.8898)	0.0662
		Combined	-4.6734 (<0.001)	0.0619 (0.9507)	0.3657 (0.7149)	0.8815 (0.3789)	0.0691
Cluster 1	0.2446	T1	-7.1703 (<0.001)	-0.4829 (0.6296)	-4.8899 (<0.001)	-0.1702 (0.865)	0.2416
		DTI	-7.1589 (<0.001)	-0.5024 (0.6158)	-4.8709 (<0.001)	0.3825 (0.7024)	0.242
		Combined	-7.1752 (<0.001)	-0.4542 (0.6501)	-4.8989 (<0.001)	-0.3391 (0.7349)	0.2419
Cluster 2	0.1695	T1	-7.2665 (<0.001)	-0.5147 (0.6072)	1.6797 (0.0943)	-0.0814 (0.9352)	0.1662
		DTI	-7.2772 (<0.001)	-0.5154 (0.6068)	1.6829 (0.0937)	-0.6514 (0.5154)	0.1676
		Combined	-7.2707 (<0.001)	-0.4611 (0.6451)	1.6725 (0.0957)	-0.5914 (0.5548)	0.1674
Cluster 3	0.0767	T1	-2.065 (0.0399)	-1.6934 (0.0916)	-3.4589 (<0.001)	-0.7735 (0.4399)	0.0753

		DTI	-2.0812 (0.0384)	-1.693 (0.0917)	-3.4728 (<0.001)	-1.8207 (0.0698)	0.0849
		Combined	-2.0636 (0.0401)	-1.6405 (0.1021)	-3.4746 (<0.001)	-0.848 (0.3972)	0.0757
Cluster 4	0.2757	T1	-10.1014 (<0.001)	0.3766 (0.7068)	1.098 (0.2733)	-2.5518 (0.0113)	0.2912
		DTI	-10.118 (<0.001)	0.31 (0.7568)	1.1208 (0.2635)	-2.5199 (0.0124)	0.2908
		Combined	-10.2703 (<0.001)	0.5794 (0.5629)	1.0357 (0.3014)	-3.6468 (<0.001)*	0.3095
Cluster 5	0.2124	T1	-7.1604 (<0.001)	-3.4563 (<0.001)*	-1.6891 (0.0924)	-1.0197 (0.3089)	0.2125
		DTI	-7.1405 (<0.001)	-3.4709 (<0.001)*	-1.6759 (0.095)	-0.5011 (0.6167)	0.21
		Combined	-7.2104 (<0.001)	-3.3405 (0.001)*	-1.7586 (0.0799)	-1.6393 (0.1024)	0.2177
Cluster 6	0.51	T1	-15.9382 (<0.001)	-1.0984 (0.2732)	-0.8171 (0.4147)	-1.8518 (0.0653)	0.5149
		DTI	-15.9778 (<0.001)	-1.1226 (0.2627)	-0.8054 (0.4214)	-2.0589 (0.0406)	0.5165
		Combined	-16.0092 (<0.001)	-0.9591 (0.3385)	-0.8532 (0.3944)	-2.3608 (0.019)	0.5191
Cluster 7	0.1381	T1	-6.5016 (<0.001)	-0.6594 (0.5102)	-0.5961 (0.5516)	-0.3018 (0.7631)	0.135
		DTI	-6.5363 (<0.001)	-0.643 (0.5208)	-0.5958 (0.5518)	-1.5824 (0.1148)	0.1432
		Combined	-6.5065 (<0.001)	-0.6229 (0.5339)	-0.6062 (0.5449)	-0.5676 (0.5708)	0.1358

892 **Table S1.** Cognitive associations with Brain Age Gap (BAG) using non-linear models,

893 including age, age^2 and sex as covariates – statistics. * FDR significant ** Bonferroni

894 significant.

895