**A Genome-Wide Association Analysis Reveals a Role for Recombination in the Evolution**

**of Antimicrobial Resistance in *Burkholderia multivorans***

Julio Diaz Caballero[1], Shawn T. Clark[2,3], Pauline W. Wang[4], Sylva L. Donaldson[4], Bryan

Coburn[5], D. Elizabeth Tullis[6], Yvonne C.W. Yau[3,7], Valerie J. Waters[8], David M. Hwang[2,3,9],

David S. Guttman[1,4,*]

[1] Department of Cell and Systems Biology, University of Toronto, Toronto, Ontario, Canada

[2] Latner Thoracic Surgery Laboratories, University Health Network, University of Toronto,

Toronto, Ontario, Canada

[3] Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario,

Canada

[4] Centre for the Analysis of Genome Evolution and Function, University of Toronto, Toronto,

Ontario, Canada

[5] Division of Infectious Diseases, Department of Medicine, University Health Network, University

of Toronto, Toronto, Ontario, Canada

[6] Adult Cystic Fibrosis Clinic, St. Michael's Hospital, Toronto, Ontario, Canada

[7] Department of Pediatric Laboratory Medicine, Division of Microbiology, The Hospital for Sick

Children, Toronto, Ontario, Canada

[8] Department of Pediatrics, Division of Infectious Diseases, The Hospital for Sick Children,

University of Toronto, Toronto, Ontario, Canada

[9] Department of Pathology, University Health Network, Toronto, Ontario, Canada.

[*] To whom correspondence may be addressed. Email: david.guttman@utoronto.ca

Running title: *B. multivorans* GWAS reveals a role for recombination in the evolution of

antimicrobial resistance

Keywords: *Burkholderia multivorans*, cystic fibrosis, evolution, genome wide association

analyses, bacterial population dynamics, recombination

## **Abstract** (300 words)

Cystic fibrosis (CF) lung infections caused by members of the *Burkholderia cepacia* complex, such as *Burkholderia multivorans*, are associated with high rates of mortality and morbidity. We performed a population genomic study of 111 *B. multivorans* sputum isolates from a single CF patient through three stages of infection including the initial incident infection, deep sampling of a one-year period of chronic infection, and deep sampling of a post-transplant recolonization. We reconstructed the evolutionary history of the population and used a lineage-controlled genome-wide association study (GWAS) approach to identify genetic variants associated with antibiotic resistance. We found that the incident isolate was more susceptible to agents from three antimicrobial classes (β-lactams, aminoglycosides, quinolones), while the chronic isolates diversified into distinct genetic lineages with reduced antimicrobial susceptibility to the same agents. The post-transplant reinfection isolates displayed genetic and phenotypic signatures that were distinct from sputum isolates from all CF lung specimens. There were numerous examples of parallel pathoadaptation, in which individual loci, or even the same codon, were independently mutated multiple times. This set of loci was enriched for functions associated with virulence and resistance. Our GWAS approach identified one variant in the *ampD* locus (which was independently mutated four times in our dataset) associated with resistance to β-lactams, and two non-synonymous polymorphisms associated with resistance to both aminoglycosides and quinolones, affecting an *araC* family transcriptional regulator, which was independently mutated three times, and an outer member porin, which was independently mutated twice. We also performed recombination analysis and identified a minimum of 14 recombination events. Parallel pathoadaptive loci and polymorphisms associated with β-lactam resistance were over-represented in these recombinogenic regions. This study illustrates the power of deep, longitudinal sampling coupled with evolutionary and lineage-corrected GWAS analyses to reveal how pathogens adapt to their hosts.

## Author Summary

Cystic fibrosis (CF) is a common lethal genetic disorder that affects individuals of European descent and predisposes them to chronic lung infections. Among the organisms involved in these infections, bacteria from the *Burkholderia cepacia* complex (BCC) are often associated with poor clinical prognosis. This study examines how the most prevalent BCC species among CF patients, *B. multivorans*, evolves within a CF patient and identifies mutations underlying antibiotic resistance and adaptation to both the native CF lung and a non-CF lung allograft. We demonstrate that *B. multivorans* can diversify phenotypically and genetically within the CF lung, with a complex population structure underlying a chronic infection We noted that isolates collected after the patient was re-infected post-transplant were more closely related to descendants of the incident clone than to those recovered in the weeks prior to transplant. We used a genome-wide association method to identify genes associated with resistance to the β-lactam antibiotics: aztreonam and ceftazidime. Many of these variants were found in regions that show patterns of recombination (genetic exchange) between strains. We also found that genes which were mutated multiple times during overall infection were more likely to be found in regions showing signals consistent with recombination. The presence of multiple independent mutations in a gene is a very strong signal that the gene helps bacteria adapt to their environment. Overall, this study provides insight into how pathogens adapt to the host during long-term infections, specific genes associated with antibiotic resistance, and the origin of new and recurrent infections.

## Introduction

The *Burkholderia cepacia* complex (BCC) describes a highly diverse group of at least 20 closely related species within the genus *Burkholderia* that can cause serious opportunistic infections in humans [1, 2]. Individuals with the fatal genetic disease cystic fibrosis (CF) are particularly susceptible to chronic BCC infections, which are commonly associated with rapid decline in lung function, high rates of mortality and poor post-transplant outcome [3, 4]. Of the BCC species, *Burkholderia multivorans* and *Burkholderia cenocepacia* account for 85-97% of all BCC found in CF patients [5]; however, *B. multivorans* infections have surpassed *B. cenocepacia* in prevalence over the past decade [6]. Many BCC that are CF-associated are intrinsically virulent and antibiotic resistant and require strict infection control practices, as they can be transmitted between patients [7-10]. Despite a wealth of knowledge describing the molecular basis of these pathogenic properties and their evolution in strains of the well-studied *B. cenocepacia*, little is known about the factors that govern these attributes in *B. multivorans* [9].

Dissecting the molecular basis of complex adaptive traits in bacterial pathogens, such as antimicrobial resistance, can be difficult as a single phenotype may be influenced by a large number of loci that interact with each other as well as their environment. Resistance in the BCC is associated with alterations to outer membrane permeability, the expression of multidrug efflux pumps and β-lactamases, and diversification of antimicrobial targets [11]. Consequently, methods that focus on identifying polymorphisms in single genes with large effects may miss the majority of loci that modulate phenotypes in more subtle ways. The development of genome-wide association studies (GWAS) has expanded our ability to identify loci of small effect size that have been associated with numerous diseases and other related phenotypes of interest in humans [12, 13].   In contrast, the application of GWAS to analyze bacterial behaviors has been slower to gain traction for a number of inter-related reasons: 1) clonal reproduction of microbes leads to confounding associations due to common ancestry, often referred to as population structure; 2) recombination in bacteria, which is more analogous to gene conversion than eukaryotic recombination, occurs at variable rates among different species and is not linked to reproduction; 3) the unpredictable nature of recombination results in the erratic breakdown of linkage disequilibrium between selected sites and distal neutral sites; and 4) selection can be extremely strong, resulting in the relatively rapid fixation of not only a selected allele, but entire genomes due to the linkage disequilibrium [14, 15].

Nevertheless, several recent studies have proposed novel approaches to overcome these

challenges. These methods include using cluster membership [16-18], phylogenetic history [15,

19, 20], or lineage effects [21] to differentiate mutations leading to a phenotypic outcome from

mutations related to the genetic background of the bacterial population. While these methods

hold tremendous promise for identifying genetic variation underlying bacterial phenotypes of

interest, they generally focus on cross sectional sampling of diverse isolates and populations.

Their power has not been established for the fine-scale analysis of individual bacterial

populations evolving over short time scales, with strong positive selection and restricted

recombination [14, 22]. The application of fine-scale evolutionary analysis to bacterial

populations is especially important in the context of clinically significant pathogen infections,

where evolution is associated with adaptation to the host environment and antimicrobial

treatment [23].

In this study, we take a fine-scale approach to microbial GWAS to examine the genetic basis of

antimicrobial resistance within a *B. multivorans* population that had been sampled longitudinally

from a single patient over a ten-year period. We characterized the genomic diversity in this

population and assessed associations between all genetic variants and multiple antibiotic

resistance phenotypes. Using a clustering-based approach to control for population structure

and linkage disequilibrium, our analysis identified single nucleotide polymorphisms (SNPs) that

were associated with resistance to β-lactams, aminoglycosides, and quinolones. In addition, we

found that both multiply-mutated loci (those that are targets of parallel pathoadaptation) and β-

lactam resistance-associated variants were overrepresented in recombinogenic regions of the

*B. multivorans* genome

## Results

For our evolutionary analysis and GWAS, we used a series of *B. multivorans* isolates that were

cultured from respiratory specimens obtained from an adult male with CF (CF170, being

followed by the CF Clinic at St. Michael's Hospital, Toronto, Canada). In a ten-year period,

patient CF170 acquired an incident (i.e. initial) lung *B. multivorans* infection, developed a

chronic *B. multivorans* lung infection, received a double lung transplant, and finally experienced

a *B. multivorans* re-colonization of the allograft three years post-transplant.  Isolates from each

of these three phases of his *B. multivorans* infection are represented in this study (Fig 1). We

defined these isolates as 1) the single isolate recovered from the patient's first infection – the

'incident infection' isolate; 2) 100 isolates collected six to seven years post-incident infection

from ten sputum specimens (ten isolates per specimen) over approximately a one-year period –

the 'chronic infection' isolates; and 3) ten isolates collected from a single expectorated sputum

sample ten years after the incident infection, and three years after the patient underwent a

double lung transplant – the 'post-transplant' isolates.  Patient CF170 was being treated with

alternating cycles of antibiotic therapy while chronically infected, with 13 antibiotics being

administered at different intervals and durations over the course of the chronic infection

sampling period (Fig 1). The genomes of all 111 isolates were whole-genome sequenced on the

Illumina platform, yielding a median coverage depth of 117X (S1 Fig). Multi-locus sequence

typing was performed *in silico* by extracting seven loci from the whole genome sequence data

(*atpD, gltB, gyrB, recA, lepA, phaC, trpB*) and comparing them to the *Burkholderia cepacia*

complex MLST Databases in pubMLST. This analysis revealed that all isolates were clonally

related and of the sequence type ST-783 [24].

**Genomic diversity and phylogenetic analysis suggest underlying population structure.**

The *de novo* genome assembly of a single isolate recovered from the third chronic infection

sputum sample was used as the reference for the mapping assembly of all other isolates. This

particular isolate was chosen as the reference since it had the best overall *de novo* assembly

metrics. The reference assembly consisted of 6,444,123 bases across 26 contigs, which were

pseudo-scaffolded against the complete genome of *B. multivorans* ATCC 17616 (Fig 2A).

Through a conservative variant calling pipeline [25], a total of 1,878 SNPs and 327 indels

segregating among the 111 isolates were identified, with 1,034, 666, and 177 SNPs being found

on chromosomes, 1, 2, and 3 respectively. Only a single SNP was found in a contig which did

not map to the ATCC 17616 genome. Overall, 740 (39.4%) SNPs and 168 (51.4%) indels were

parsimonious informative (PI, i.e. non-singleton), and 226 (12.0%) SNPs and 99 (30.3%) indels

segregated in at least two sampling time points. From the 1,878 SNPs, 70.6%, 15.7%, and

13.7% were non-synonymous, synonymous, and intragenic substitutions respectively (Fig 2C).

47.7% of the intergenic SNPs were found in putative regulatory regions (defined as the

intergenic region within 150 bases from the start codon of any gene). The population showed a

genetic diversity average of 123.39 ± 120.55 (number of SNP differences, mean ± standard

deviation) pairwise differences. The distribution of these difference suggested an underlying

population structure since genetic diversity was not uniform even among isolates from the same

specimen (S2 Fig).

We reconstructed the core genome phylogenetic relationships among all isolates using an

alignment of the 1,878 SNPs and a Bayesian approach (Fig 3A). The root of this tree was

identified by adding *B. multivorans* ATCC17616 to the analysis. The tree topology indicates that

the incident infection isolate diverged from the other 110 isolates at the base of the tree. The ten

isolates from the post-transplant sample are again highly divergent (relative to the total diversity)

and form a basally branching, monophyletic clade, while the chronic sample isolates form a less

divergent, monophyletic clade. Moreover, there seem to be subgroups among the chronic

infection isolates suggesting population structure. This structure is also observed in a network-

based phylogenetic approach (S3 Fig), where two groups of isolates from the chronic infection

sampling cluster in a star-like phylogeny. Star phylogenies are characterized by roughly equal

divergence from the common ancestor, and are associated with recent purges in genetic

variation [26].

**Population structure analysis clusters the isolates into five groups.**  We used the Monte

Carlo Markov Chain analysis of SNPs and indels implemented in STRUCTURE to infer

population structure among the 111 isolates. We identified the lowest number of subpopulations

that maximized the likelihood of data; hence determining the underlying population structure in

the data without overestimating the number of subpopulations [27]. There were three

subpopulations that arose from single common ancestors, which we labelled groups R, B, and

G, comprising 54, 26, and 10 isolates, respectively (Fig 3C-D).  The ancestral composition of

the incident isolate and seven of the chronic infection isolates, recovered at collection points T1,

T2 and T10, resembled a combination of the three identified subpopulations. This group of

isolates was labeled RBG. Another group labeled RB (13 isolates) has an admixed ancestry

from the ancestral subpopulations of R and B.

Isolates from groups RBG and RB were found in low frequencies through different samples from

the chronic infection period (Fig 3B). In contrast, isolates from group R or B were more

dominant in this same period. The isolates from group R were first observed at the third time

point of the chronic infection samples, and they remained the most abundant group in

subsequent chronic samples (Fig 4). In contrast, the abundance of group B isolates decreased

over time. The genetic diversity, measured as number of SNPs, significantly differed between

these groups (one-way ANOVA: $F(4,1902) = 1,371.452$, p-value $< 0.0001$), with group G (those

recovered exclusively post-transplant) being the most diverse, followed by groups RBG and RB,

then groups R and B (S4a Fig).

The time to the most recent common ancestor (tMRCA) calculated as days before the last

sample for all isolates and the various STRUCTURE-defined groups is shown in Supplementary

Figure 4c.  This analysis shows that the RGB group, which includes all of the chronic infection

isolates as well as the post-transplant isolates, coalesced to a common ancestor at roughly the

same time as the full isolate collection, including the incident infection (S4c Fig). This result

supports the hypothesis that the infection of the transplanted lung came from the same source

as the original incident isolate, despite being separated by approximately ten years, as opposed

to a clone that persisted and diversified in the lung of the patient during chronic colonization.

Additionally, it appears that groups R and B diverged at approximately the same time (S4c Fig).

Unfortunately, we are unable to determine if these were allopatric populations that colonized

distinct regions in the lung, or sympatric populations that coexisted within the same

compartment due to our sampling of expectorated sputum.

**Selection analysis supports positive selection in the population.** We determined the ratio

of non-synonymous to synonymous substitutions ($d_N/d_S$) as an estimate of selection.  Since we

expect that time has allowed natural selection or genetic drift to have acted on the multi-time

segregating mutations more so than on variants that segregate in a single sample, we

determined the $d_N/d_S$ both for all SNPs in each group, as well as for only those that segregate in

at least two time-points – 'multi-time' SNPs (S4b Fig). The $d_N/d_S$ for the overall population was

1.35 (95% confidence interval, CI = 1.19-1.53) and 1.34 for multi-time SNPs (CI = 0.94-1.96),

which may indicate weak positive selection, or simply the segregation of mildly deleterious

variants.  Only groups R and RB multi-time SNPs showed $d_N/d_S$ above the neutral expectation

of 1.0 (group R $d_N/d_S$ = 2.05, CI = 0.57-11.15, group RB $d_N/d_S$ = 2.38, CI = 1.08-6.18), although

the confidence intervals for the group R are quite large. All other groups had $d_N/d_S$ ratios only

slightly elevated (ranging from 1.19-1.63), although the differences between groups were not

statistically significant.

Further support for positive selection comes from a significantly negative Tajima's D test (D = -

2.21, P < 0.01) and Fu and Li's tests (D* = -6.11, P < 0.02; F* = -5.20, P < 0.02). While all three

of these results can be explained by both positive selection and recent population expansion,

the combination of these results with the high nucleotide diversity and $d_N/d_S$ > 1.0 is most

consistent with positive selection.

**GWAS identification of variants associated with antibiotic resistance.** We assumed that the intensive antibiotic exposure during the chronic infection sampling period would result in strong selection for resistance-associated genotypes in *B. multivorans.* Minimum inhibitory concentrations (MICs) for two β-lactams (aztreonam, ceftazidime), two aminoglycosides (tobramycin and amikacin), and the fluoroquinolone ciprofloxacin were determined for all isolates. Isolates from the three phases of infection had distinct susceptibility profiles. The incident isolate had MICs of 8 µg/mL or less for all agents tested, while all chronic infection and post-transplant isolates had higher MICs for both aminoglycosides (t-test $p < 0.0001$, Fig 3E), but variable MICs for β-lactams and fluoroquinolone tested (range: ≤8 to >512 µg/mL).

The 1,878 SNP positions segregating among the 111 isolates were grouped in 149 distinct mutational profiles (i.e. one or more SNP positions that share the same pattern of reference vs. alternative base among the strain collection, S5 Fig). Prior to population control, each of these mutational profiles was examined for a statistical association to the five tested antibiotics at six different levels of resistance and these associations were corrected for multiple testing by taking into consideration the number of tests. Five mutational profiles (comprising 17 SNPs) associated with resistance to both β-lactam antibiotics, and one mutational profile (comprising 2 SNPs) associated specifically with ceftazidime (S6 and S7 Fig). Ten mutational profiles (comprising 250 SNPs) were associated with resistance to amikacin, tobramycin, and ciprofloxacin. Additionally, two mutational profiles (comprising 31 SNPs) associated with resistance to both aminoglycosides, and four mutational profiles (comprising 33 SNPs) associated specifically with ceftazidime.

Next, we tested these variants against population structure controls, counting only those associated variants that were observed in multiple subpopulation groups as determined by the population structure analysis. This criterion could be satisfied by one of two mechanisms: 1) the mutations arose in the subpopulations through multiple independent mutational events, or 2) they arose in a common ancestor of multiple subpopulations and have been maintained in multiple lineages while being lost in others. Out of all mutational profiles associated with elevated MICs for both β-lactams, one (comprising a single SNP) passed the population structure control (S6b Fig). This SNP was found in 20.4% of isolates in group R, and 50% of isolates in group RBG. This variant leads to a non-synonymous amino acid substitution in the sequence of the *ampD* gene (BMUL_2790), a locus extensively studied for its role in resistance to β-lactams [28, 29]. This mutation was predicted to have a deleterious effect on AmpD by

PROVEAN analysis (score = -8.0, S8a Fig). In fact, the *ampD* locus was independently mutated

four other times within our dataset. A second SNP in *ampD* was found in a mutational profile

that was similarly associated with β-lactam MICs; nevertheless, it failed to pass the population

structure control. Additionally, two mutational profiles associated to the aminoglycosides and

ceftazidime showed evidence of multiple independent polymorphic events (S6e Fig). One of

these mutational profiles, which comprises a single SNP, is represented by a non-synonymous

substitution in an *araC* family transcriptional regulator locus (BMUL_3951). PROVEAN analysis

indicates that this mutation is unlikely to have a deleterious effect on the locus (score = 6.906).

The second mutational profile, again including only a single SNP, gave rise to a non-

synonymous substitution in locus BMUL_3342, which is annotated as an outer member protein

(porin). While this mutation is not expected to end in a deleterious effect (PROVEAN score =

3.273), it occurs in a locus that is independently mutated two other times.

**Additional variants associated with pathoadaptation can be detected by identifying multi-**

**mutated loci.** Loci that are independently mutated multiple times provide strong evidence of

selection by parallel pathoadaptation [30]. We observed 328 loci that were independently

mutated multiple times in our collection (Table 1). Given the genome size and the total number

of polymorphisms (both SNPs and indels), we only consider the 62 loci with three or more

independent mutations to be statistically significant (p-value < 0.05/[1,878 SNPs + 327 indels =

2205 polymorphisms]). 184 SNPs (9.8%) and 26 indels (8.0 %) were found in these 62 loci. We

excluded the possibility that multiply mutated loci showed excess polymorphism simply due to

an increased mutational rate by examining the mutational class spectrum for the multiply

mutated loci relative to the genome-wide average. While the rate of non-synonymous,

synonymous and intergenic mutations among all 1,878 SNPs is 70.6%, 15.7%, and 13.7%

respectively, the mutational class spectrum of the SNPs found among multiply mutated loci is

83.1% non-synonymous, 11.7% synonymous, and 3.2% intergenic substitutions. Therefore, the

mutational class distribution of SNPs found in multiply mutated loci is significantly skewed

toward an excess of non-synonymous mutations (P < 0.0001, chi-square test).

Some of these multi-mutated loci are known to play significant roles in antibiotic resistance. For

example, a gene encoding a probable transcriptional regulator protein of MDR efflux pump

cluster (BMUL_0641), which has been associated with drug resistance in multiple pathogens

[31-33], has seven independently acquired mutations, and the probability of any gene being

mutated seven times is $1.65 \times 10^{-23}$. A locus with five multiple mutations (P = $6.48 \times 10^{-16}$) encodes

N-acetylmuramoyl-L-alanine amidase (AmpD, BMUL_2790), which is associated with resistance

to β-lactam antibiotics [28]. Moreover, a functional enrichment analysis revealed the

phosphorelay signal transduction system GO function overrepresented in multiply mutated

genes compared to the functional annotation of the whole genome (P = 0.050). The

phosphorelay signal transduction system has been previously described as a therapeutic target,

given that it controls the expression of genes encoding virulence factors [34].

We also found ten genes that had two independent mutations located in the same or adjacent

codon (Table 2). The mutational class spectrum of the SNPs associated with this observation is

of 90%, 10% and 0% of non-synonymous, synonymous, and intergenic substitutions,

respectively. In this case, the fraction of non-synonymous mutations is significantly higher than

the fraction found for both all SNPs, as well as all the SNPs in the multiply mutated loci (P <

0.00001, chi-square test). One of the genes with multiple independent mutations in the same

codon encodes for RNA polymerase sigma factor (RpoD), which is associated with the

expression of housekeeping genes [35]. One of the mutations in this locus is fixed between the

post-transplant isolates and the rest of the isolates, and the other mutation is fixed between the

isolates in group RBG collected in the tenth sample time and the rest of the isolates.

**Parallel pathoadaptive variants are overrepresented in recombinogenic regions.** We

identified a minimum of 14 recombination events in our full dataset based on the four-gamete

tests of Hudson and Kaplin [36] (Fig 2D). Three of these events were identified between sites in

different genome assembly contigs; therefore, they were not considered in downstream

recombination analysis. The nucleotide length of this recombinogenic regions ranged from

4,783 bases to 192,532 bases, and these regions account for 15.1% of the assembled genome.

298 (15.9%) out of the total 1,878 SNPs and 47 indels (14.4%) occur in these regions, which is

not significantly different than expected given the recombinogenic proportion of the genome.

We next looked to see if there was an association between recombination and the evolution of

antibiotic resistance. 51 (18.3%) of the 279 SNPs associated with both aminoglycosides tested

(amikacin & tobramycin), and 42 (14.9%) of the 281 SNPs linked to ciprofloxacin are found in

recombinogenic regions (Fig 5A). These ratios fail to reject the null hypothesis that these

mutations are randomly distributed around the genome. On the other hand, 52.9% (9 of 17

SNPs) and 47.4% (9 of 19 SNPs) of the SNPs associated with aztreonam and ceftazidime,

respectively, are found in recombinogenic regions, which are significantly different than

expected by chance (p < 0.0001, chi square test).

Finally, 49 (26.6%) of the 184 SNPs and 4 (8.5%) of the 47 indels found in loci independently

mutated three or more times occur in the identified recombinogenic regions (Fig 5B). Thus,

while SNPs involved in multi-mutated loci are overrepresented in recombinogenic regions more

than expected (P < 0.0001, chi square test), indels in multi-mutated genes are not significantly

underrepresented.

**Discussion**

Our study investigated how *B. multivorans* evolves within the lungs of an individual afflicted with

CF using a deep longitudinal sampling design (i.e. multiple isolates obtained per sputum

sample) to capture both the overall population diversity and the temporal shifts that occurred at

different phases of the infection, including the colonization of a new allograft. To identify the

source of genetic diversity in this *B. multivorans* population, we needed to understand: 1) the

genetic relationships between the incident isolate that was recovered from the first BCC-positive

sputum culture, the chronic strains that persisted in the population, and the population of strains

that re-established an infection post-transplant; 2) whether there were multiple colonization

events of the patient by divergent clones; 3) how genetic diversity was generated and dispersed

in the population; and 4) how the pathogen adapts and responds to clinical treatment. While we

were unable to address all of these questions, we have concluded that the chronic population

originated from either the incident isolate, or a clone that shared a recent common ancestor with

the incident isolate. Furthermore, all of the chronic isolates descended from a single common

ancestor, ruling out multiple independent colonization events.

One clear signal is that the *B. multivorans* isolates recovered from the post-transplant lung did

not originate from the chronic population. In fact, it appears that the post-transplant isolates

came from a new infection that originated from the same source as the incident infection.

Unfortunately, the source of these infections cannot be determined, and could be either the

environment or the patient's upper respiratory tract. In the former case, it is likely that the patient

lived in the same home or locale over the course of the study, and that the ancestral *B.*

*multivorans* clone is endemic in that environment. Alternatively, in the latter case, the upper

respiratory tract is known to act as a reservoir for a number of CF pathogens [37].

Consequently, it is possible that clonal descendants of the ancestral or incident strains resided
in the patient's upper airways since the incident infection. Some transplant procedures attempt
to clean the nasal reservoir prior to transplant via nasal washing / scraping, but we do not know
if this was done for this patient. If this hypothesis is true, it would explain why the post-transplant
isolates have an antibiotic susceptibility pattern much more similar to the chronic isolates than
the incident isolate. We also note that the post-transplant population is much more genetically
diverse than any of the chronic populations. This could suggest that this population was rapidly
adapting to an environmental change, such as the shift from CF to non-CF conditions, which
would include, differences in immune response, the composition of the allograft microbiome,
and treatment regimens. Alternatively, it could reflect colonization by a population of related
strains. It is possible that given sufficient time this population would eventually be winnowed
down to a single surviving clone (as is seen with the incident infection) due to selection and / or
genetic drift

A major motivator for this study was to better understand how pathogens adapt to their hosts
over the course of disease progression and treatment; an issue that can be addressed using
statistical association tests. Correcting for the genetic structure of the bacterial population poses
a challenge to the implementation of these tests. Population structure in this context refers
relationships among strains due to descent form a common ancestor and limited recombination.
This structure results in the linkage of segregating genetic variation around the genome, which
makes it very difficult to distinguish a causal mutation that is responsible for a phenotype of
interest from a neutral variant that occurred in the same genetic background. In the absence of
recombination, the neutral mutation will have the same population distribution as the causal
mutation due to genetic hitchhiking. This issue is particularly prevalent when studying largely
isolated and recently evolved populations, such as the case of pathogens evolving within a host.

To overcome these two issues, we imposed a lineage control filter on our GWAS approach, in
which we focused only on mutations that occurred in multiple, distinct, genetic lineages. This
pattern can best be explained by recombination of polymorphisms between lineages, but
formally, could also be due to extensive gene loss. Our analysis showed that linkage
disequilibrium was only disrupted in a relatively small number of polymorphism (those
polymorphisms shown as orange circles; S7b-e Fig). This reinforces the need for deep sampling
since the infrequent recombination signals may have been missed if isolates were only collected
from a single sample, or if only single isolates were recovered from each sample. Consequently,

the tractability of GWAS in this *B. multivorans* population was greatly enhanced by our sampling

schema.

Using the established lineage structure of the *B. multivorans* population as control for our

association study, we identified two non-synonymous SNPs associated with resistance to the

aminoglycosides amikacin and tobramycin, and to the quinolone ciprofloxacin. One of these

SNPs occurs in a locus encoding the transcription factor AraC, which is involved in the global

regulation of efflux pumps, while the other SNP was found in a locus annotated as a porin.

Although not specific to aminoglycosides or quinolones, overexpression of efflux pumps and

repression of porin proteins has been reported as important mechanisms of antibiotic resistance

for bacteria [38]. Neither mutation is projected to significantly vary the function of the encoding

protein.

Additionally, we identified a single SNP associated with resistance to the β-lactams aztreonam

and ceftazidime. This SNP occurs in the *ampD* gene, which is a negative regulator of the β-

lactamase AmpC, and it is expected to have a deleterious effect in the encoding protein. This

observation is not unexpected as bacteria treated with β-lactams would benefit from the

constitutive overproduction of β-lactamase. Overall, AmpD seems to play an important role in

the pathoadaptation of this *B. multivorans* population since four other independent non-

synonymous mutations, all of which are expected to have deleterious effects on the protein,

occur at this locus (S8a Fig).

Our use of the population control criterion of only considering mutations present in multiple

lineages meant that we excluded some variants associated to virulence, such as one of the four

mutations in *ampD*, which was statistically associated with β-lactam resistance. Without our

population control it would be impossible to identify causative mutations from hitchhiking

variants that are in linkage disequilibrium with the causative mutation. Filtering in this manner

reduces the number of false positives; nevertheless, variants underlying phenotypes of interest

could be segregating in linkage disequilibrium blocks, and therefore, may not be identified in our

GWAS approach (false negatives).

We observed that mutations associated with resistance to β-lactams (prior to lineage controls)

occur disproportionately in recombinogenic regions (Fig 2F), while variants associated with both

aminoglycosides or ciprofloxacin are more randomly distributed with respect to recombinogenic

regions. The study patient received both long-term maintenance β-lactam and aminoglycoside

treatments in addition to multiple short-term β-lactam treatments that included cycles of

ceftazidime, piperacillin/tazobactam, meropenem, and cefepime. This more aggressive and

varied course of treatment with β-lactams could potentially explain the increased role of

recombination in the dissemination of putatively beneficial polymorphisms, similar to what has

been observed in other pathogens [39, 40].

Our analysis identified genes under strong selection by focusing on loci with a statistical excess

of independent mutations (i.e. parallel pathoadaptation) [25, 41, 42]. Examining multi-mutated

loci can reveal the heterogeneous selective pressures that bacteria must adapt to in order to

reside within the lung. For instance, a gene encoding a transcription regulator of multidrug

resistance efflux pumps independently accumulated seven different mutations leading to eight

unique alleles in our population of 111 *B. multivorans* isolates. We also found seven different

alleles of a locus encoding cyclic β-1,2-glucan synthase, which is linked to bacteria's ability to

elude host cell defenses [43]. A number of loci underlying virulence-associated traits, such as

quorum sensing and biofilm production, also carry multiple independent mutations. Particularly

interesting are multiply mutated loci with no characterized function, or with no prior linkage to

resistance or virulence. These loci include a NAD-glutamate dehydrogenase locus BMUL_4010,

which was mutated five independent times over the course of the study, and a glycosyl

transferase protein (BCEN2424_5592), not previously seen in *B. multivorans* that was mutated

six times (4 SNPs and 2 indels) during the course of the study. Examples such as these provide

excellent candidates for characterizing the cryptic resistome – loci previously not known to be

involved in antimicrobial resistance. In addition, the strongest signals of parallel pathoadaptation

involve those cases where mutations occur independently in the same or adjacent codon.

These observations point to a very specific form of selective pathoadaptation, which identifies

the specific residue or region of the locus that potentially plays a role in selective advantage and

may affect a conserved function.

Finally, our study highlighted an intriguing role for recombination in the development of

antimicrobial resistance in *B. multivorans*. We observed that multi-mutated loci were over-

represented within recombinogenic regions, along with an excess of mutations associated with

β-lactam resistance. This suggests that while recombination plays an important role in the

pathoadaptation of this *B. multivorans* population, its selective benefit may be environment

dependent.

Our study illustrates the relevance of deep, longitudinal sampling to the implementation of
GWAS approaches in a population under positive selection. We identified the potential genetic
basis behind the antibiotic resistance of a *B. multivorans* population in a single host. Moreover,
this approach allowed us to study variants associated to antibiotic resistance and revealed that
resistance to β-lactams may be passed within the population via recombination. This study is
limited to *in silico* predictions of the impact mutations on protein function, and future efforts
should include functional validation of these mutants; nevertheless, many of the identified genes
are already well-established targets for antibiotic resistance. Additionally, our findings are
restricted to a single patient and a single bacterial species; extending this approach in other
systems under positive selection will be required to establish the generalizability of the findings.
Nevertheless, this study is one of the first examining in depth the fine-scale evolution of *B.*
*multivorans* in the lungs of a CF patient as it transitions from an early infection to chronic
infections and the eventual reinfection of a transplanted allograft.

## Materials and Methods

**Ethics statement.** All protocols involving the collection, handling and laboratory use of respiratory specimens were approved by the Research Ethics Boards of St. Michael's Hospital (Protocol #09-289) (Toronto, Canada) and the University Health Network (Protocol #09-0420-T) (Toronto, Canada). We obtained informed consent from the study subject prior to specimen collection and sputa were produced voluntarily. All experiments involving clinical specimens were performed in accordance with the *Tri-Council Policy Statement: Ethical Conduct for Research Involving Humans*, of the Canadian Institutes of Health Research, the Natural Sciences and Engineering Research Council of Canada, and the Social Sciences and Humanities Research Council of Canada.

**Specimen collection and isolation of *B. multivorans*.** Sputum specimens were collected by expectoration from a 29-year-old male (CF170), with a homozygous ΔF508 CFTR genotype being followed at the Adult CF Clinic at St. Michael's Hospital (Toronto, Canada). Ten sputum specimens were collected over a 10-month period while the patient was in the advanced stages of CF lung disease (assessed by the forced expiratory volume in 1 second ($FEV_1$), $FEV_1$ which was 27-39 % predicted throughout the course of the study), and an additional sputum specimen obtained after the patient had undergone double lung transplantation. All specimens were processed for bacterial culture as previously described [44]. After 48h of incubation, cultures were visually inspected, and each distinct colony morphotype was described using eight characteristics of physical appearance (pigmentation, size, surface texture, surface sheen, opacity, mucoidy, autolysis and margin shape). Ten colonies were selected from each sputum culture in relation to the diversity of colony types present. The incident isolate was obtained from the *Burkholderia cepacia* complex repository at St. Michael's Hospital and was recovered from the first BCC positive sputum culture produced by the study patient (Toronto, Canada). Isolates were stored at −80°C in 20% (v/v) glycerol after a 20h subculture in LB broth (Wisent Inc., QC, CA) and confirmed as *Burkholderia* spp. by a secondary subculture onto both *Burkholderia cepacia* selective (BCSA) (HiMedia Laboratories, Mumbai, IN) and MacConkey (Becton Dickinson, MD, USA) agars, as well as being tested for growth at 42°C. The *recA* gene was sequenced from each isolate as described by Spilker *et al.* for preliminary speciation [45].

**Antimicrobial susceptibility testing.** Each isolate confirmed as *B. multivorans* was screened for antimicrobial susceptibility by agar dilution using Clinical and Laboratory Standards Institute

procedures [46]. We tested susceptibility to representatives of the β-lactam (aztreonam [ATM],

ceftazidime [CAZ]), fluoroquinolone (ciprofloxacin [CIP]) and aminoglycoside (amikacin [AMK],

tobramycin [TOB]) (Sigma-Aldrich, ON, Canada) classes. Minimum inhibitory concentrations

(MIC), defined as the lowest concentration of each antibiotic to inhibit growth, were reported as

the median MIC of three independent experiments. Growth was assessed following 24 to 48 h

of incubation on Mueller-Hinton agar (Becton, Dickinson, MD, USA). The *B. multivorans* ATCC

17616 strain was included as a positive control, while *P. aeruginosa* ATCC 27853 and *E. coli*

ATCC 25922 were used as quality controls.

**Sequencing and Quality Control.** *B. multivorans* isolates were whole-genome sequenced on

the MiSeq and NextSeq Illumina platforms. The number of bases sequenced per isolate ranged

from 213 to 2,262 million bases, and the median was 1,026 million bases. Trimmomatic v. 0.33

was used to remove adapters and quality trim the sequencing reads from each isolate

(parameter settings: PE -phred33 ILLUMINACLIP:adapters.fa:2:30:10 LEADING:5 TRAILING:5

SLIDINGWINDOW:4:25) [47]. Sequencing reads with guanine homopolymers longer than ten

bases were trimmed with cutadapt v. 1.9.1 (parameter settings: -a "G{10}") [48]. Reads bellow

100 bases were removed using Trimmomatic v. 0.33 (parameter settings: PE –phred33

MINLENGTH:100). The resulting quality-controlled sequencing reads yielded a median read

depth per position of 117X (range 32-276X).

*De novo* **and Reference Mapping Assembly.** Each of the isolates was *de novo* assembled

using the CLC Genomics Workbench v. 8.0.1 (Aarhus, Denmark). Contigs with a scaffolding

depth lower than 10X and/or with a size smaller than 1 Kb were removed from further analyses.

Isolate CF170-3b, which was sequenced with 250 bp-long paired-end reads, yielded the best

assembly metrics in 26 contigs with lengths ranging from 1,010 to 1,243,078 bases and an N50

of 654,231. The final assembly length of the CF170-3b isolate was of 6,444,123 bp. These

contigs were annotated at the RAST server using the native gene caller and Classic RAST as

the annotation scheme [47]. Further, this genome was functionally annotated with blast2go v

4.1.9 [49] including blastx v. 2.6.0+ [50]. Statistical results from the functional enrichment

analysis were Bonferroni corrected for multiple testing using the number of multiply-mutated

genes (P-value/62). The contigs of the CF170-3b genome were used as the reference for

mapping assembly of each remaining isolate. We performed three different reference-mapping

assemblies including BWA v 0.7.12 [51], LAST v 284v [52] and novoalign v 2.08.03 (Novocraft

Technologies).

**Single Nucleotide Polymorphism (SNP) and indel Calling.** SAMtools and BCFtools v 0.1.19
were used to produce the initial set of variants [53]. We implemented a method previously
described to detect SNPs among the 111 isolates [25, 54]. First, 1,878 high-confidence
polymorphic positions were identified using the following criteria: 1) variant Phred quality score
of ≥ 30 and 2) variants must be found at least 150 bp away from either the edge of the reference
contig or an indel. Second, we reviewed each high-confidence polymorphic position in each
isolate with a relaxed Phred score threshold of 25. Support for either the reference or the SNP
call was verified with a multi-hypothesis correction which required that at least 80% of the
sequencing reads endorsed the SNP or the reference. If the data did not support either base,
then the position was called as an ambiguous base ('N'). The ambiguous call rate was lower
than 0.1%.

Candidate indels detected by BWA and SAMtools were examined by realigning mapped and
unmapped sequencing reads to the indel regions using Dindel v. 1.01 [55]. High-confidence
indel positions were defined as sites with: 1) variant Phred quality score of ≥ 35; 2) at least two
forward and two reverse reads; and 3) sequencing coverage ≥ 10. These indel positions were
reviewed in each isolate. The final indel call required a Phred quality score ≥ 25 and an allele
frequency ≥ 80%. Ambiguous indel calls were defined as those where the allele frequency was
≤ 20%.

**Population and Single Genome Sequencing Evaluation.** We performed bulk population
sequencing on the post-transplant specimen to confirm that our isolate sampling depth
appropriately represented the real *B. multivorans* population diversity (S9 Fig). The sequencing
reads from each of the ten isolates from the post-transplant sample were rarified to 1/10$^{th}$ of the
number of sequencing reads produced by the population sequencing experiment. These reads
were combined in corresponding paired-end fasta files. Next, population and single isolate
sequencing reads were mapped to the *de novo* assembled genome of the CF170-3b isolate
using BWA. Mutation allele frequencies for each experiment were estimated was previously
described by Lieberman *et al.* [54].

**Phylogenetic, Population Structure, Coalescent and Recombination Analyses.** Using the
1,878 SNPs, we created a genome-wide alignment to reconstruct the phylogenetic relationships
among the 111 isolates. The phylogeny was calculated using MrBayes v. 3.2.6 [56]. The

nucleotide substitution model that best fit our data was the General Time Reversible (GTR) with

gamma-distributed rate variation across sites (LnL=-12,858.8103, AIC= 26,175.6205) as

calculated with jModelTest v. 2.1.10 [57]. The Bayesian analysis was run through two different

chains of 1 million Markov Chain Monte Carlo (MCMC) generations sampled every 100 MCMC

generations and the burn-in period was of 250,000 MCMC generations. The final average

standard deviation of split frequencies was of $7.7 \times 10^{-3}$, and the potential scale reduction factor

(PSRF) of the substitution model parameters ranged from 1 - $4.74 \times 10^{-4}$ to 1 + $6.84 \times 10^{-4}$. The

phylogeny was rooted with *B. multivorans* ATCC 17616 [58]. The network-based phylogenetic

analysis was performed using SplitsTree v 4.14.4 [59]. We employed the Jukes-Cantor distance

matrix to implement the neighbor-net Network (Fit=99.418).

The variance among the 111 isolates, including SNPs and indels, was employed to investigate

the population structure using the Structure software v 2.3.4 [60]. Structure employs a Bayesian

algorithm to detect the number of ancestral populations (K), also known as clusters, which

describe the variance and covariance observed in a test population. The number of clusters

ranging from 1-10 was tested in triplicates through 1 million MCMC generations sampled every

1,000 MCMC generations and a burn-in period of 250,000 MCMC generations. We used the

correlated allele frequencies model, and admixture was allowed in these analyses. We plotted

the estimated ln probability of data for the tested levels of K, and identified the smallest stable K

as the optimum value since it maximized the global likelihood of the data (S10 Fig) [61]. The

estimated ln probability of data plateaus at K=3, where the variance of ln likelihood ranges from

2,517.3 to 2,466.7. Assuming three ancestral populations, the isolates were classified into five

different groups according to their ancestry. Isolates whose ancestry is attributed exclusively

(>90%) to either ancestral population one, two, or three are grouped in group red (R), (B), or

(G), respectively. Group RB includes isolates with admixed ancestry from clusters one and two

(at least 10% of both cluster one and two, and less than 10% of cluster three). Isolates whose

ancestral composition is made up from a combination of all three clusters (at least 10% of each

cluster) are in group RBG.

We used BEAST v. 1.8.4 to implement a Bayesian approach to inferring the time to the most

recent common ancestor (tMRCA) for the entire population and each group individually [62].

Next, we employed the GTR nucleotide substitution model, and estimated the nucleotide

substitution frequencies with MEGA7 using the Maximum Likelihood Estimate of the Substitution

Matrix tool ([AC] = 0.0092, [AG] = 0.4278, [AT] = 0.0016, [CG] = 0.0262, [GT] = 0.0062, and

[CT] = 0.5290). Preliminary analyses consisting of duplicate 10 million generations and a 10%

burn-in were used to estimate the appropriate molecular clock and demographic models. We

tested the Bayesian skygrid, constant size and the exponential, logarithmic and expansion

growth population size models using three different molecular clock models (strict and the

lognormal and exponential uncorrelated relaxed clocks). The exponential relaxed uncorrelated

molecular clock and the Bayesian skygrid model was inferred the most appropriate given our

data ([AIC] = 26,228.421) [63]. The final analysis was run in duplicate for 1 billion MCMC

generations sampled every 1,000 MCMC generation, and the burn-in period was set at 10% of

the MCMC generations.

Population genetic tests and detection of recombination events in each contig were performed

with DnaSP v. 5.10.01 [64].

**SNP to Phenotype Association.** We tested the null hypothesis that the presence or absence

of each of the 1,878 SNPs, summarized in 149 distinct mutational profiles, is equally likely found

in antibiotic resistant isolates using Fisher's exact test. These tests were conducted for each

examined antibiotic at six different MIC resistance thresholds (≤16, 32, 64, 128, 256 and ≤512

MIC). For each test, we created a contingency table reflecting the distribution of each mutation

profile in isolates with lower and greater MIC than each resistance threshold. *P* values were

adjusted based on the total number of tests (number of mutational profiles), and only

associations with a *P* value < 3.36 X $10^{-4}$ (0.05 / 149) were considered significant to control for

multiple testing. Next, we simulated gains or losses of these mutational events following a

continuous-time Markov chain along a ClonalFrameML v. 1.0-19 phylogeny as implemented in

GLOOME v. 01.266 using the default parameters [65, 66]. We defined independent mutational

events as those with a probability greater than 0.95 and to control for population structure, we

required multiple independent mutational events in at least two STRUCTURE-defined groups.

*In silico* mutation impact prediction. To predict the potential impact of non-synonymous

SNPs on the biological function of a protein, we employed PROVEAN v. 1.1.3 [67]. These

calculations were performed on the GPC supercomputer at the SciNet HPC Consortium [68].

## Acknowledgements

This research was funded by an Emerging Team Grant awarded to D.S.G from the Canadian Institutes of Health Research (CIHR) and Cystic Fibrosis Canada (CMF108027). J.D.C. was supported by an Ontario Trillium Scholarship. S.T.C. was supported by an Ontario Graduate Scholarship.  B.C. was supported by a CIHR Fellowship.

# References

1. Vandamme P, Dawyndt P. Classification and identification of the Burkholderia cepacia complex: Past, present and future. Syst Appl Microbiol. 2011;34(2):87-95. Epub 2011/01/25. doi: 10.1016/j.syapm.2010.10.002. PubMed PMID: 21257278.

2. De Smet B, Mayo M, Peeters C, Zlosnik JE, Spilker T, Hird TJ, et al. Burkholderia stagnalis sp. nov. and Burkholderia territorii sp. nov., two novel Burkholderia cepacia complex species from environmental and human sources. Int J Syst Evol Microbiol. 2015;65(7):2265-71. Epub 2015/04/16. doi: 10.1099/ijs.0.000251. PubMed PMID: 25872960.

3. Courtney JM, Bradley J, McCaughan J, O'Connor TM, Shortt C, Bredin CP, et al. Predictors of mortality in adults with cystic fibrosis. Pediatr Pulmonol. 2007;42(6):525-32. Epub 2007/05/01. doi: 10.1002/ppul.20619. PubMed PMID: 17469153.

4. Stephenson AL, Sykes J, Berthiaume Y, Singer LG, Aaron SD, Whitmore GA, et al. Clinical and demographic factors associated with post-lung transplantation survival in individuals with cystic fibrosis. J Heart Lung Transplant. 2015;34(9):1139-45. Epub 2015/06/20. doi: 10.1016/j.healun.2015.05.003. PubMed PMID: 26087666.

5. Drevinek P, Mahenthiralingam E. Burkholderia cenocepacia in cystic fibrosis: epidemiology and molecular mechanisms of virulence. Clin Microbiol Infect. 2010;16(7):821-30. doi: 10.1111/j.1469-0691.2010.03237.x. PubMed PMID: 20880411.

6. Lipuma JJ. The changing microbial epidemiology in cystic fibrosis. Clin Microbiol Rev. 2010;23(2):299-323. Epub 2010/04/09. doi: 10.1128/CMR.00068-09. PubMed PMID: 20375354; PubMed Central PMCID: PMC2863368.

7. Lipuma JJ. Update on the Burkholderia cepacia complex. Current opinion in Pulmonary Medicine. 2005;11(6):528-33. Epub 2005/10/12. doi: 00063198-200511000-00010 [pii]. PubMed PMID: 16217180.

8. Jones AM, Dodd ME, Govan JR, Barcus V, Doherty CJ, Morris J, et al. *Burkholderia cenocepacia* and *Burkholderia multivorans*: influence on survival in cystic fibrosis. Thorax. 2004;59(11):948-51. Epub 2004/11/02. doi: 59/11/948 [pii]
10.1136/thx.2003.017210. PubMed PMID: 15516469; PubMed Central PMCID: PMC1746874.

9. Leitao JH, Sousa SA, Ferreira AS, Ramos CG, Silva IN, Moreira LM. Pathogenicity, virulence factors, and strategies to fight against *Burkholderia cepacia* complex pathogens and related species. Appl Microbiol Biotechnol. 2010;87(1):31-40. Epub 2010/04/15. doi: 10.1007/s00253-010-2528-0. PubMed PMID: 20390415.

10. Zlosnik JE, Zhou G, Brant R, Henry DA, Hird TJ, Mahenthiralingam E, et al. Burkholderia species infections in patients with cystic fibrosis in British Columbia, Canada. 30 years'

experience. Ann Am Thorac Soc. 2015;12(1):70-8. Epub 2014/12/05. doi:

10.1513/AnnalsATS.201408-395OC. PubMed PMID: 25474359.

11. Rhodes KA, Schweizer HP. Antibiotic resistance in Burkholderia species. Drug Resist

Updat. 2016;28:82-90. Epub 2016/09/14. doi: 10.1016/j.drup.2016.07.003. PubMed PMID:

27620956; PubMed Central PMCID: PMCPMC5022785.

12. Price AL, Spencer CC, Donnelly P. Progress and promise in understanding the genetic

basis of common diseases. Proc Biol Sci. 2015;282(1821):20151684. doi:

10.1098/rspb.2015.1684. PubMed PMID: 26702037; PubMed Central PMCID:

PMCPMC4707742.

13. McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, Ioannidis JP, et al. Genome-

wide association studies for complex traits: consensus, uncertainty and challenges. Nat Rev

Genet. 2008;9(5):356-69. doi: 10.1038/nrg2344. PubMed PMID: 18398418.

14. Power RA, Parkhill J, de Oliveira T. Microbial genome-wide association studies: lessons

from human GWAS. Nat Rev Genet. 2017;18(1):41-50. doi: 10.1038/nrg.2016.132. PubMed

PMID: 27840430.

15. Farhat MR, Shapiro BJ, Kieser KJ, Sultana R, Jacobson KR, Victor TC, et al. Genomic

analysis identifies targets of convergent positive selection in drug-resistant Mycobacterium

tuberculosis. Nat Genet. 2013;45(10):1183-9. doi: 10.1038/ng.2747. PubMed PMID:

23995135; PubMed Central PMCID: PMCPMC3887553.

16. Chewapreecha C, Harris SR, Croucher NJ, Turner C, Marttinen P, Cheng L, et al. Dense

genomic sampling identifies highways of pneumococcal recombination. Nat Genet.

2014;46(3):305-9. Epub 2014/02/11. doi: 10.1038/ng.2895. PubMed PMID: 24509479;

PubMed Central PMCID: PMCPMC3970364.

17. Chewapreecha C, Marttinen P, Croucher NJ, Salter SJ, Harris SR, Mather AE, et al.

Comprehensive identification of single nucleotide polymorphisms associated with beta-

lactam resistance within pneumococcal mosaic genes. PLoS Genet. 2014;10(8):e1004547.

doi: 10.1371/journal.pgen.1004547. PubMed PMID: 25101644; PubMed Central PMCID:

PMCPMC4125147.

18. Chen PE, Shapiro BJ. The advent of genome-wide association studies for bacteria. Curr

Opin Microbiol. 2015;25:17-24. doi: 10.1016/j.mib.2015.03.002. PubMed PMID: 25835153.

19. Sheppard SK, Didelot X, Meric G, Torralbo A, Jolley KA, Kelly DJ, et al. Genome-wide

association study identifies vitamin B5 biosynthesis as a host specificity factor in

Campylobacter. Proc Natl Acad Sci U S A. 2013;110(29):11923-7. doi:

10.1073/pnas.1305559110. PubMed PMID: 23818615; PubMed Central PMCID:

PMCPMC3718156.

20. Chaston JM, Newell PD, Douglas AE. Metagenome-wide association of microbial

determinants of host phenotype in Drosophila melanogaster. MBio. 2014;5(5):e01631-14.

doi: 10.1128/mBio.01631-14. PubMed PMID: 25271286; PubMed Central PMCID:

PMCPMC4196228.

21. Earle SG, Wu CH, Charlesworth J, Stoesser N, Gordon NC, Walker TM, et al. Identifying

lineage effects when controlling for population structure improves power in bacterial

association studies. Nat Microbiol. 2016;1:16041. doi: 10.1038/nmicrobiol.2016.41. PubMed

PMID: 27572646; PubMed Central PMCID: PMCPMC5049680.

22. Didelot X, Maiden MC. Impact of recombination on bacterial evolution. Trends Microbiol.

2010;18(7):315-22. doi: 10.1016/j.tim.2010.04.002. PubMed PMID: 20452218; PubMed

Central PMCID: PMCPMC3985120.

23. Hughes D, Andersson DI. Evolutionary consequences of drug resistance: shared principles

across diverse targets and organisms. Nat Rev Genet. 2015;16(8):459-71. doi:

10.1038/nrg3922. PubMed PMID: 26149714.

24. Jolley KA, Maiden MC. BIGSdb: Scalable analysis of bacterial genome variation at the

population level. BMC Bioinformatics. 2010;11:595. doi: 10.1186/1471-2105-11-595.

PubMed PMID: 21143983; PubMed Central PMCID: PMCPMC3004885.

25. Diaz Caballero J, Clark ST, Coburn B, Zhang Y, Wang PW, Donaldson SL, et al. Selective

Sweeps and Parallel Pathoadaptation Drive Pseudomonas aeruginosa Evolution in the

Cystic Fibrosis Lung. MBio. 2015;6(5):e00981-15. doi: 10.1128/mBio.00981-15. PubMed

PMID: 26330513; PubMed Central PMCID: PMCPMC4556809.

26. McVean G. The structure of linkage disequilibrium around a selective sweep. Genetics.

2007;175(3):1395-406. doi: 10.1534/genetics.106.062828. PubMed PMID: 17194788;

PubMed Central PMCID: PMCPMC1840056.

27. Kalinowski ST. The computer program STRUCTURE does not reliably identify the main

genetic clusters within species: simulations and implications for human population structure.

Heredity (Edinb). 2011;106(4):625-32. doi: 10.1038/hdy.2010.95. PubMed PMID: 20683484;

PubMed Central PMCID: PMCPMC3183908.

28. Hwang J, Kim HS. Cell Wall Recycling-Linked Coregulation of AmpC and PenB beta-

Lactamases through ampD Mutations in Burkholderia cenocepacia. Antimicrob Agents

Chemother. 2015;59(12):7602-10. doi: 10.1128/AAC.01068-15. PubMed PMID: 26416862;

PubMed Central PMCID: PMCPMC4649219.

29. Kong KF, Schneper L, Mathee K. Beta-lactam antibiotics: from antibiosis to resistance and
bacteriology. APMIS. 2010;118(1):1-36. doi: 10.1111/j.1600-0463.2009.02563.x. PubMed
PMID: 20041868; PubMed Central PMCID: PMCPMC2894812.

30. Wood TE, Burke JM, Rieseberg LH. Parallel genotypic adaptation: when evolution repeats
itself. Genetica. 2005;123(1-2):157-70. PubMed PMID: 15881688; PubMed Central PMCID:
PMCPMC2442917.

31. Westfall LW, Carty NL, Layland N, Kuan P, Colmer-Hamood JA, Hamood AN. mvaT
mutation modifies the expression of the Pseudomonas aeruginosa multidrug efflux operon
mexEF-oprN. FEMS Microbiol Lett. 2006;255(2):247-54. doi: 10.1111/j.1574-
6968.2005.00075.x. PubMed PMID: 16448502.

32. da Silva PE, Von Groll A, Martin A, Palomino JC. Efflux as a mechanism for drug resistance
in Mycobacterium tuberculosis. FEMS Immunol Med Microbiol. 2011;63(1):1-9. doi:
10.1111/j.1574-695X.2011.00831.x. PubMed PMID: 21668514.

33. Coyne S, Courvalin P, Perichon B. Efflux-mediated antibiotic resistance in Acinetobacter
spp. Antimicrob Agents Chemother. 2011;55(3):947-53. doi: 10.1128/AAC.01388-10.
PubMed PMID: 21173183; PubMed Central PMCID: PMCPMC3067115.

34. Stephenson K, Hoch JA. Two-component and phosphorelay signal-transduction systems as
therapeutic targets. Curr Opin Pharmacol. 2002;2(5):507-12. Epub 2002/09/27. PubMed
PMID: 12324251.

35. Potvin E, Sanschagrin F, Levesque RC. Sigma factors in Pseudomonas aeruginosa. FEMS
Microbiol Rev. 2008;32(1):38-55. doi: 10.1111/j.1574-6976.2007.00092.x. PubMed PMID:
18070067.

36. Barton NH. Richard Hudson and Norman Kaplan on the Coalescent Process. Genetics.
2016;202(3):865-6. doi: 10.1534/genetics.116.187542. PubMed PMID: 26953263; PubMed
Central PMCID: PMCPMC4788121.

37. Folkesson A, Jelsbak L, Yang L, Johansen HK, Ciofu O, Hoiby N, et al. Adaptation of
Pseudomonas aeruginosa to the cystic fibrosis airway: an evolutionary perspective. Nat Rev
Microbiol. 2012;10(12):841-51. Epub 2012/11/14. doi: 10.1038/nrmicro2907. PubMed PMID:
23147702.

38. Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. Molecular mechanisms of
antibiotic resistance. Nat Rev Microbiol. 2015;13(1):42-51. doi: 10.1038/nrmicro3380.
PubMed PMID: 25435309.

39. Garcia-Solache M, Lebreton F, McLaughlin RE, Whiteaker JD, Gilmore MS, Rice LB.
Homologous Recombination within Large Chromosomal Regions Facilitates Acquisition of

beta-Lactam and Vancomycin Resistance in Enterococcus faecium. Antimicrob Agents
Chemother. 2016;60(10):5777-86. doi: 10.1128/AAC.00488-16. PubMed PMID: 27431230;
PubMed Central PMCID: PMCPMC5038250.
40. Aubert D, Naas T, Nordmann P. Integrase-mediated recombination of the veb1 gene
cassette encoding an extended-spectrum beta-lactamase. PLoS One. 2012;7(12):e51602.
doi: 10.1371/journal.pone.0051602. PubMed PMID: 23251590; PubMed Central PMCID:
PMCPMC3518468.
41. Marvig RL, Sommer LM, Molin S, Johansen HK. Convergent evolution and adaptation of
Pseudomonas aeruginosa within patients with cystic fibrosis. Nat Genet. 2015;47(1):57-64.
doi: 10.1038/ng.3148. PubMed PMID: 25401299.
42. Sokurenko EV, Hasty DL, Dykhuizen DE. Pathoadaptive mutations: gene loss and variation
in bacterial pathogens. Trends Microbiol. 1999;7(5):191-5. PubMed PMID: 10354593.
43. Arellano-Reynoso B, Lapaque N, Salcedo S, Briones G, Ciocchini AE, Ugalde R, et al.
Cyclic beta-1,2-glucan is a Brucella virulence factor required for intracellular survival. Nat
Immunol. 2005;6(6):618-25. doi: 10.1038/ni1202. PubMed PMID: 15880113.
44. Clark ST, Diaz Caballero J, Cheang M, Coburn B, Wang PW, Donaldson SL, et al.
Phenotypic diversity within a Pseudomonas aeruginosa population infecting an adult with
cystic fibrosis. Sci Rep. 2015;5:10932. doi: 10.1038/srep10932. PubMed PMID: 26047320;
PubMed Central PMCID: PMCPMC4456944.
45. Spilker T, Baldwin A, Bumford A, Dowson CG, Mahenthiralingam E, LiPuma JJ. Expanded
multilocus sequence typing for burkholderia species. J Clin Microbiol. 2009;47(8):2607-10.
doi: 10.1128/JCM.00770-09. PubMed PMID: 19494070; PubMed Central PMCID:
PMCPMC2725695.
46. CLSI. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow
Aerobically; Approved Standard M07-A9. Wayne, PA: Clinical and Laboratory Standards
Institute; 2012.
47. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20. doi: 10.1093/bioinformatics/btu170. PubMed PMID:
WOS:000340049100004.
48. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011. 2011;17(1). doi: 10.14806/ej.17.1.200
pp. 10-12.
49. Gotz S, Garcia-Gomez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, et al. High-
throughput functional annotation and data mining with the Blast2GO suite. Nucleic Acids

Res. 2008;36(10):3420-35. Epub 2008/05/01. doi: 10.1093/nar/gkn176. PubMed PMID:

18445632; PubMed Central PMCID: PMCPMC2425479.

50. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST

and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res.

1997;25(17):3389-402. Epub 1997/09/01. PubMed PMID: 9254694; PubMed Central

PMCID: PMCPMC146917.

51. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25(14):1754-60. doi: 10.1093/bioinformatics/btp324. PubMed PMID:

19451168; PubMed Central PMCID: PMCPMC2705234.

52. Shrestha AM, Frith MC. An approximate Bayesian approach for mapping paired-end DNA

reads to a reference genome. Bioinformatics. 2013;29(8):965-72. doi:

10.1093/bioinformatics/btt073. PubMed PMID: 23413433; PubMed Central PMCID:

PMCPMC3624798.

53. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. doi:

10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed Central PMCID:

PMCPMC2723002.

54. Lieberman TD, Michel JB, Aingaran M, Potter-Bynoe G, Roux D, Davis MR, Jr., et al.

Parallel bacterial evolution within multiple patients identifies candidate pathogenicity genes.

Nat Genet. 2011;43(12):1275-80. doi: 10.1038/ng.997. PubMed PMID: 22081229; PubMed

Central PMCID: PMCPMC3245322.

55. Albers CA, Lunter G, MacArthur DG, McVean G, Ouwehand WH, Durbin R. Dindel: accurate

indel calls from short-read data. Genome Res. 2011;21(6):961-73. doi:

10.1101/gr.112326.110. PubMed PMID: 20980555; PubMed Central PMCID:

PMCPMC3106329.

56. Ronquist F, Huelsenbeck JP. MrBayes 3: Bayesian phylogenetic inference under mixed

models. Bioinformatics. 2003;19(12):1572-4. PubMed PMID: 12912839.

57. Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics

and parallel computing. Nat Methods. 2012;9(8):772. doi: 10.1038/nmeth.2109. PubMed

PMID: 22847109; PubMed Central PMCID: PMCPMC4594756.

58. Stanier RY, Palleroni NJ, Doudoroff M. The aerobic pseudomonads: a taxonomic study. J

Gen Microbiol. 1966;43(2):159-271. doi: 10.1099/00221287-43-2-159. PubMed PMID:

5963505.

59. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol Biol
Evol. 2006;23(2):254-67. doi: 10.1093/molbev/msj030. PubMed PMID: 16221896.

60. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus
genotype data. Genetics. 2000;155(2):945-59. PubMed PMID: 10835412; PubMed Central
PMCID: PMCPMC1461096.

61. Porras-Hurtado L, Ruiz Y, Santos C, Phillips C, Carracedo A, Lareu MV. An overview of
STRUCTURE: applications, parameter settings, and supporting software. Front Genet.
2013;4:98. doi: 10.3389/fgene.2013.00098. PubMed PMID: 23755071; PubMed Central
PMCID: PMCPMC3665925.

62. Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and
the BEAST 1.7. Mol Biol Evol. 2012;29(8):1969-73. doi: 10.1093/molbev/mss075. PubMed
PMID: 22367748; PubMed Central PMCID: PMCPMC3408070.

63. Gill MS, Lemey P, Faria NR, Rambaut A, Shapiro B, Suchard MA. Improving Bayesian
population dynamics inference: a coalescent-based model for multiple loci. Mol Biol Evol.
2013;30(3):713-24. doi: 10.1093/molbev/mss265. PubMed PMID: 23180580; PubMed
Central PMCID: PMCPMC3563973.

64. Librado P, Rozas J. DnaSP v5: a software for comprehensive analysis of DNA
polymorphism data. Bioinformatics. 2009;25(11):1451-2. doi: 10.1093/bioinformatics/btp187.
PubMed PMID: 19346325.

65. Didelot X, Wilson DJ. ClonalFrameML: efficient inference of recombination in whole bacterial
genomes. PLoS Comput Biol. 2015;11(2):e1004041. doi: 10.1371/journal.pcbi.1004041.
PubMed PMID: 25675341; PubMed Central PMCID: PMCPMC4326465.

66. Cohen O, Ashkenazy H, Belinky F, Huchon D, Pupko T. GLOOME: gain loss mapping
engine. Bioinformatics. 2010;26(22):2914-5. doi: 10.1093/bioinformatics/btq549. PubMed
PMID: 20876605.

67. Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the functional effect of amino
acid substitutions and indels. PLoS One. 2012;7(10):e46688. doi:
10.1371/journal.pone.0046688. PubMed PMID: 23056405; PubMed Central PMCID:
PMCPMC3466303.

68. Chris L, Daniel G, Leslie G, Richard P, Neil B, Michael C, et al. SciNet: Lessons Learned
from Building a Power-efficient Top-20 System and Data Centre. Journal of Physics:
Conference Series. 2010;256(1):012026.

69. Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos: an
information aesthetic for comparative genomics. Genome Res. 2009;19(9):1639-45. doi:

10.1101/gr.092759.109. PubMed PMID: 19541911; PubMed Central PMCID:

PMCPMC2752132.

70. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and

annotation of phylogenetic and other trees. Nucleic Acids Res. 2016;44(W1):W242-5. doi:

10.1093/nar/gkw290. PubMed PMID: 27095192; PubMed Central PMCID:

PMCPMC4987883.

71. Miller CA, McMichael J, Dang HX, Maher CA, Ding L, Ley TJ, et al. Visualizing tumor

evolution with the fishplot package for R. BMC Genomics. 2016;17(1):880. doi:

10.1186/s12864-016-3195-z. PubMed PMID: 27821060; PubMed Central PMCID:

PMCPMC5100182.

72. Lieberman TD, Flett KB, Yelin I, Martin TR, McAdam AJ, Priebe GP, et al. Genetic variation

of a bacterial pathogen within individuals with cystic fibrosis provides a record of selective

pressures. Nat Genet. 2014;46(1):82-7. doi: 10.1038/ng.2848. PubMed PMID: 24316980;

PubMed Central PMCID: PMCPMC3979468.

73. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics. 2014;30(9):1312-3. doi: 10.1093/bioinformatics/btu033.

PubMed PMID: 24451623; PubMed Central PMCID: PMCPMC3998144.

74. Carrasco-Lopez C, Rojas-Altuve A, Zhang W, Hesek D, Lee M, Barbe S, et al. Crystal

structures of bacterial peptidoglycan amidase AmpD and an unprecedented activation

mechanism. J Biol Chem. 2011;286(36):31714-22. doi: 10.1074/jbc.M111.264366. PubMed

PMID: 21775432; PubMed Central PMCID: PMCPMC3173140.

**Table 1. Parallel Pathoadapted Loci with Multiple Independent Mutations**

| Locus | Encoded Protein | No. of SNPs/Indels | Probability [a] |
|---|---|---|---|
| BMUL_0641 | Probable transcription regulator protein of MDR efflux pump cluster | 7/0 | $1.65 \times 10^{-23}$ |
| BCEN2424_5592 [c] | Glycosyltransferase 36 | 4/2 | $1.03 \times 10^{-19}$ |
| BMUL_4010 | NAD-glutamate dehydrogenase | 5/0 | $6.48 \times 10^{-16}$ |
| BMUL_0487 | Hypothetical protein | 5/0 | $6.48 \times 10^{-16}$ |
| BMUL_4327 | Porin | 3/2 | $6.48 \times 10^{-16}$ |
| BMUL_2790 | N-acetyl-anhydromuranmyl-L-alanine amidase (AmpD) | 5/0 | $6.48 \times 10^{-16}$ |
| BMUL_1598 | Amino acid adenylation domain-containing protein | 4/0 | $4.06 \times 10^{-12}$ |
| BMUL_0353 | YD repeat-containing protein | 3/1 | $4.06 \times 10^{-12}$ |
| BMUL_0449 | Preprotein translocase subunit (SecB) | 4/0 | $4.06 \times 10^{-12}$ |
| BMUL_2632 | Chaperone protein (DnaJ) | 4/0 | $4.06 \times 10^{-12}$ |
| BMUL_4942 | Signal transduction histidine kinase (CheA) | 3/1 | $4.06 \times 10^{-12}$ |
| BMUL_2775 | UDP-N-acetylmuramate--L-alanyl-gamma-D-glutamyl-meso-diaminopimelate ligase | 4/0 | $4.06 \times 10^{-12}$ |
| BMUL_1444 | Transcription termination factor (Rho) | 4/0 | $4.06 \times 10^{-12}$ |
| BMUL_0954 | Glycoside hydrolase 15-like protein | 4/0 | $4.06 \times 10^{-12}$ |
| BMUL_4115 | Outer membrane autotransporter | 4/0 | $4.06 \times 10^{-12}$ |
| BMUL_0250 | 50S ribosomal protein L4 (RplD) | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_5547 | Conjugation protein (TrbI) | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_2931 | TPR repeat-containing protein | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3678 | Integral membrane sensor signal transduction histidine kinase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3503 | L-serine dehydratase 1 | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_0690 | RND efflux system outer membrane lipoprotein | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_0663 | Alpha/beta hydrolase fold protein | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_0431 | Histidine kinase | 1/2 | $2.55 \times 10^{-8}$ |
| BMUL_4510 | Signal transduction histidine kinase (CheA) | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_1970 | Major facilitator transporter | 3/0 | $2.55 \times 10^{-8}$ |

| | | | |
|---|---|---|---|
| BMUL_2008 | Major facilitator transporter | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_2621 | DNA mismatch repair protein (mutL) | 1/2 | $2.55 \times 10^{-8}$ |
| BMUL_4037 | Esterase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3977 | Metallophosphoesterase | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_4949 | Aldehyde dehydrogenase | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_3951 | Transcriptional regulator (AraC) | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_6019 | Cytosine/purines uracil thiamine allantoin permease | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_0307 | Amino acid carrier protein | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_5501 | Cytochrome c oxidase subunit I | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_5087 | Short-chain dehydrogenase/reductase SDR | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_4813 | RNA polymerase sigma factor RpoD | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3197 | Beta-galactosidase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3212 | Feruloyl-CoA synthase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3315 | PA-phosphatase like phosphoesterase | 1/2 | $2.55 \times 10^{-8}$ |
| BMUL_3752 | Peptidoglycan-binding (LysM) | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3615 | Aldehyde oxidase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_1686 | Ribonuclease R | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_4615 [b] | Amidophosphoribosyltransferase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_4605 | UTP-glucose-1-phosphate uridylyltransferase | 3/0 | $2.55 \times 10^{-8}$ |
| ABD05_14940 [d] | Isochorismatase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_1431 | GAF modulated sigma54 specific transcriptional regulator (Fis) | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_1377 | N-acetyltransferase GCN5 | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_0964 | DNA polymerase III subunit alpha (DnaE) | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_0692 | Carbohydrate kinase FGGY | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_0477 | Error-prone DNA polymerase (DnaE2) | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_0443 | Phosphoenolpyruvate-protein phosphotransferase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_3068 | Aldehyde dehydrogenase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_4835 | Hypothetical protein | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_1873 | UvrD/REP helicase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_2536 | Hypothetical protein | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_2710 | Outer membrane autotransporter | 3/0 | $2.55 \times 10^{-8}$ |

| | | | |
|---|---|---|---|
| BMUL_0123 | Heavy metal translocating P-type ATPase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_0116 | Acyl-CoA dehydrogenase domain-containing protein | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_0075 | Two component transcriptional regulator | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_4226 | 4-hydroxyphenylpyruvate dioxygenase | 3/0 | $2.55 \times 10^{-8}$ |
| BMUL_4749 | Amino acid permease | 2/1 | $2.55 \times 10^{-8}$ |
| BMUL_4798 | Integrase catalytic region | 1/2 | $2.55 \times 10^{-8}$ |

[a] Calculated based on the probability of resampling with replacement any locus n times, given a genome size of N. $P = (1/N)^{(n-1)}$. We used $(n-1)$ since we are calculating the probability for any locus, rather than a specific locus.

[b] A mutation occurred in the intergenic region flanking the start codon of this locus.

[c] This locus is not found in ATCC 17616 The homolog with highest similarity is in *B. cenocepacia* DDS 22E-1

[d] This locus is not found in ATCC 17616 The homolog with highest similarity is in *B. cenocepacia* HI2424

**Table 2. Pairs of Mutations Occurring in the Same or in Neighboring Codons.**

| Encoded Protein | Proximity |
| --- | --- |
| Regulatory protein GntR, HTH:GntR, C-terminal | Adjacent codon |
| Oligopeptide ABC transporter, periplasmic oligopeptide-binding protein (OppA) | 2 codons away |
| Citrate-proton symporter | 2 codons away |
| CDP-6-deoxy-delta-3,4-glucoseen reductase-like | 2 codons away |
| RNA polymerase sigma factor (RpoD) [a] | Same codon |
| Endo-1,4-beta-xylanase Z precursor [b] | Adjacent codon |
| Isoquinoline 1-oxidoreductase beta subunit [b] | 2 codons away |
| LSU ribosomal protein L4p (L1e) [b] | Same codon |
| Chaperone protein (DnaJ) [c] | Adjacent codon |
| Probable transcription regulator protein of MDR efflux pump cluster [d] | 2 codons away |

a Loci additionally mutated 1 more time. Additional mutation is synonymous.

b Loci additionally mutated 1 more time. Additional mutation is non-synonymous.

c Locus additionally mutated 2 more times. All non-synonymous mutations.

d Locus additionally mutated 5 more times. All non-synonymous mutations.

# Figure Legends

**Fig 1. Time course of *B. multivorans* infection in study patient CF170.** A total of 111 *B. multivorans* isolates from twelve collection times were used in this study (1 isolate from the initial infection, 10 isolates from each of 10 sputum samples collected during chronic infection, and 10 isolates from a sputum sample obtained during a post-transplant infection). Antibiotic treatment history during the chronic infection period is shown in the lower panel. Black bars indicate antibiotic administration, while hashed bars indicate intermittent exposure in that time block (only relevant prior to the start of chronic sampling). The method of antibiotic administration is shown as intravenous (iv), inhaled (inh), or oral (po).

**Fig 2. Genomic Characterization of 111 *B multivorans* isolates.** (A) Contigs (gray outer ring) of the *de novo* reference were arranged according to the three chromosomes of the complete genome of *B. multivorans* ATCC 17616. This genome was obtained from expectorated sputum collected in the third chronic infection sample. (B) Genome annotation according to RAST. (C) SNP count per 10 Kb as a function of their location in the contigs. Non-synonymous (orange), synonymous (yellow), putative regulatory (dark grey) and intergenic (light grey). (D) Indel (blue) count per 10 Kb. (E) Recombinogenic regions, as predicted by DnaSP Hudson-Kaplin four gamete test, are shown as red blocks. (F) Variants Associated with Antibiotic Resistance. From outermost to innermost ring: aztreonam and ceftazidime (β-lactam), amikacin and tobramycin (aminoglycoside), and ciprofloxacin (quinolone). This figure was prepared with circus v. 0.69 [69].

**Fig 3. Population structure and antibiotic resistance profiles.** (A) Phylogenetic relationships of the 111 *B. multivorans* isolates were estimated employing a Bayesian approach based on genome-wide single nucleotide polymorphisms (SNPs). (B) Time of collection for each isolate. (C) Population structure analysis as assessed by Structure v2.3.4 with three expected ancestral subpopulations. Ancestral subpopulations are coded as red (R), blue (B), and green (G). (D) Isolates are grouped based on their ancestral composition. Group R, B, G, RB, and RBG are shaded in red, blue, green, purple, and grey respectively. (E) Antibiotic susceptibility for each isolate, the highest black circle represents the MIC (μg/mL), to the β-lactams: aztreonam and ceftazidime, the aminoglycosides: amikacin and tobramycin, and the quinolone: ciprofloxacin are shown as filled circles at six different concentration thresholds. This figure was elaborated at the interactive tree of life (iTOL) website v. 3 [70].

**Fig 4. Population genomics of the community over time.** Groups R, B, G, RB, and RBG are coloured in red, blue, green, purple, and grey respectively. (A) Frequency of each group over time. (B) The clonal graph was created with the assumption that RGB is the group of isolates resembling the ancestor of all the isolates, and RB is the group of isolates resembling the ancestor of group R and B. The distance between sample times is relative to the actual number of days between them. This plot was created using fishplot v. 0.3 [71].

**Fig 5. Distribution of pathoadaptive variants in recombinogenic regions of the genome.** (A) Distribution of the mutations associated with the tested antibiotics in the identified recombinogenic regions and in the rest of the genome (\*\*\* $p < 0.0001$, chi square test with multiple test correction). (B) Distribution of the mutations in multi-mutated loci in the identified recombinogenic regions and in the rest of the genome (\*\*\* $p < 0.001$, chi square test with multiple test correction).

## Supporting Information

**S1 Fig. Sequencing coverage.** Whole genome sequencing of 111 isolates of *B. multivorans* in the Illumina platform. (A) Distribution of number of bases sequenced per isolate. (B) Distribution of median read depth per position.

**S2 Fig. Genetic diversity over time.** (A) Pairwise nucleotide differences between isolates collected from the same collection sample. Incident infection is not included since only one isolate was recovered from that time point. (B) Nucleotide differences between each isolate and the incident infection isolate.

**S3 Fig. Neighbor-Net phylogeny.** This network-based phylogeny was calculated in SplitsTree v. 4.14.4. Individual strain names at the tips of each branch have been replaced with pie charts indicating the distribution of dates during which the strains were sampled (indicated by the circular legend).

**S4 Fig. Genetic diversity and selection analysis per group.** (A) Pairwise nucleotide differences between isolates from the same group based on ancestry. (B) $d_N/d_S$ per group calculated including all SNPs and using only SNPs observed in multiple time points (MTP). dN/dS and the respective confidence intervals were calculated as described by Lieberman *et al*. [72]. (C). Time to Most Recent Common Ancestry (tMRCA) as estimated using the BEAST software for each group. The x axis represents the log of the years before the last sampling time. The whiskers for each data point show the 95% high probability density intervals.

**S5 Fig. SNP positions with identical distribution of reference or alternative bases across the strain collection are grouped into mutational profiles.** Here, "0"s and "1"s represent the reference or alternative base, respectively, at each SNP position for each strain. SNP1 is the only position where only Strain1 has a base alternative to the reference. Hence, mutational profile 1, 1-0-0-0, comprises only one SNP. On the other hand, Strain4 is the only strain with a variant base for positions SNP2 and SNP3. Therefore, mutational profile 2, 0-0-0-1, comprises SNP2 and SNP3.

**S6 Fig. Mutational profiles associated with antibiotic resistance.** (A) Maximum Likelihood phylogeny of 111 *B. multivorans* isolates was elaborated using RaxML v. 7.0.4 with a GTR +

1019 gamma model and 1,000 bootstraps [73]. Here, we show all mutation profiles associated with
1020 antibiotic resistance prior to lineage control in black and with lineage control in orange. (B)
1021 resistance to both β-lactams, (C) to amikacin only, (D) to both aminoglycosides, (E) to both
1022 aminoglycosides and to ciprofloxacin, (F) and to ciprofloxacin only. A filled circle represents a
1023 SNP call in the corresponding isolate compared to the reference.

1024

**S7 Fig. Resistance levels at which genetic associations are statistically significant.**
Mutational profiles were tested for association against six levels of antibiotic resistance (<16,
<32, <64, <128, <256 and <512 MIC) to five antibiotics (amikacin, tobramycin, aztreonam,
ceftazidime and ciprofloxacin). Black boxes show the levels of resistance at which the
mutational profiles were statistically significant including multi-testing correction. Associations to
ciprofloxacin antibiotic resistance are shown up to <128 MIC since no isolate had a MIC of 256
or greater in relation to that antibiotic.

1032

**S8 Fig. Mutations in *ampD* locus.** (A) Distribution of the PROVEAN scores of all identified
non-synonymous substitutions highlighting SNPs in multi-mutated loci (yellow) and in the *ampD*
gene (red or blue if associated to β -lactam resistance). Red lines represent thresholds from
most specific (highest), to most sensitive (lowest) to determine if a mutation is deleterious to the
function of the gene in which it occurs. (B) Crystal structure of protein product of AmpD (PDB
ID:2Y2B, [74]) in complex with reaction products. Mutations found in our *B. multivorans*
population are colored in red or blue (mutations associated with β-lactam resistance).

1040

**S9 Fig. Population and single isolate sequencing.** Sequencing reads from each isolate from
the post-transplant sample were rarified to 1/10th of the number of reads in the population
sequencing experiment; then they were combined so that the number of reads would be the
same for both experiments. Sequencing reads from the population and single isolate
experiments were mapped to the same reference as described above. Mutation allele
frequencies for both experiments were calculated using the quality thresholds described by
Lieberman *et al*. [54]. (A) Grey circles represent mutation allele frequencies in the deep
population sequencing experiment (y axis) versus in single isolate sequencing (x axis). The
dashed line represents the $x=y$ function and the solid line is the best fit line taking into account
all data points ($R^2$=0.9928, 95% confidence interval= 0.9918-0.9937). Red circles represent
alleles found in the single isolate sequencing experiment but not in the deep sequencing one.

Fixed mutations between the reference and all the post-transplant isolates are colored blue. (B)

Proportion of false positives in the single isolate sequencing experiment.

1054

1055    **S10 Fig.  Determining the number of ancestral populations that explain the variance and**

1056    **covariance in CF170 *B. multivorans* population.** (A) We ran three independent chains for

1057    each K between one and ten. The estimated ln probability of data plateaus at K=3 in all chains.

**Tree scale: 0.1**

**A**

**B** INCIDENT / T1 / T2 / T3 / T4 / T5 / T6 / T7 / T8 / T9 / T10 / POST-TRANSPANT — Collection Time

**C** % Ancestral Contribution

**D** group R / group B / group RB / group RBG / group G

**E**

Aztreonam — β-lactams — ≤512 / 256 / 128 / 64 / 32 / ≤16

Ceftazidime — ≤512 / 256 / 128 / 64 / 32 / ≤16

Amikacin — Aminoglycosides — ≤512 / 256 / 128 / 64 / 32 / ≤16

Tobramycin — ≤512 / 256 / 128 / 64 / 32 / ≤16

Ciprofloxacin — Quinolone — ≤512 / 256 / 128 / 64 / 32 / ≤16

**A**

Frequency

**B**

Group RBG
Group RB
Group B
Group R
Group G

Incident Infection

T1 T2 T3 T4 T5 T6 T7 T8 T9 T10

Chronic Infection

Post Transplant Infection