Computational Causal Modeling of the Dynamic Biomarker Cascade in Alzheimer's Disease

Jeffrey R. Petrella[1], Wenrui Hao[2], Adithi Rao[1], P. Murali Doraiswamy[3]

for the Alzheimer's Disease Computational Modeling Initiative

Departments of Radiology[1] and Psychiatry[3], Duke University Health System

Department of Mathematics[2], Penn State University

Send Correspondence to Jeffrey R. Petrella, MD, Department of Radiology, DUMC - Box 3808, Durham, NC 27710-3808; Telephone: 919-684-7218; Fax: 919-684-7157; Email: jeffrey.petrella@duke.edu

Abstract:

Background: Alzheimer's disease (AD) is a major public health concern and there is an urgent need to better understand its complex biology and develop effective therapies. AD progression can be tracked in patients though validated imaging and spinal fluid biomarkers of pathology and neuronal loss. We still, however, lack a coherent quantitative model that explains how these biomarkers interact and evolve over time. Such a model could potentially help identify the major drivers of disease in individual patients and simulate response to therapy prior to entry in clinical trials. A current theory of AD biomarker progression, known as the dynamic biomarker cascade model, hypothesizes AD biomarkers evolve in a sequential, but temporally overlapping manner. A computational model incorporating assumptions about the underlying biology of this theory and its variations would be useful to test and refine its accuracy with longitudinal biomarker data from clinical trials.

Methods: We implemented a causal model to simulate time-dependent biomarker data under the descriptive assumptions of the dynamic biomarker cascade theory. We modeled pathologic biomarkers (beta-amyloid and tau), neuronal loss biomarkers and cognitive impairment as non-linear first order ordinary differential equations (ODEs) to include amyloid-dependent and non-dependent neurodegenerative cascades. We tested the feasibility of the model by adjusting its parameters to simulate three specific natural history scenarios in early-onset autosomal dominant AD and late-onset AD, and determine whether computed biomarker trajectories agreed with current assumptions of AD biomarker progression. We also simulated the effects of anti-amyloid therapy in late-onset AD.

Results: The computational model of early-onset AD demonstrated the initial appearance of amyloid, followed by biomarkers of tau and neurodegeneration, followed by onset of cognitive decline based on cognitive reserve, as predicted by prior literature. Similarly, the late-onset AD computational models demonstrated the first appearance of amyloid or non-amyloid-related tauopathy, depending on the

magnitude of comorbid pathology, and also closely matched the biomarker cascades predicted by prior literature. Forward simulation of anti-amyloid therapy in symptomatic late-onset AD failed to demonstrate any slowing in progression of cognitive decline, consistent with prior failed clinical trials in symptomatic patients.

Conclusions: We have developed and computationally implemented a mathematical causal model of the dynamic biomarker cascade theory in AD. We demonstrate the feasibility of this model by simulating biomarker evolution and cognitive decline in early and late-onset natural history scenarios, as well as in a treatment scenario targeted at core AD pathology. Models resulting from this causal approach can be further developed and refined using patient data from longitudinal biomarker studies, and may in the future play a key role in personalizing approaches to treatment.

Introduction:

Alzheimer's disease (AD), one of the leading public health priorities in the U.S., is projected to affect over 15 million people by 2050. The high failure rate of clinical drug trials over the past decade is in large part rooted in an incomplete understanding of its complex causal mechanisms (3). Genetic pathway analyses implicate over 1000 different molecular species and over 30 metabolic pathways in the pathophysiology of AD, including amyloid and tau proteinopathies, inflammation, microglial activation, alterations in signaling pathways, cholesterol metabolism, and cholinergic function (4). It is therefore likely that AD is not a single disease, but a common end-stage pathway resulting from multiple interacting etiologies. Effective treatment will likely require a personalized medicine approach to track disease progression, determine the major pathophysiologic drivers and tailor an appropriate therapy.

AD progression can be tracked in patients, from pre-symptomatic to late-stage disease, through several validated biomarkers. These are biomarkers of AD core pathology (cerebrospinal fluid and PET scan markers of beta-amyloid and tau proteins), and biomarkers of neuronal loss (FDG-PET and volumetric MR imaging). Data from the Alzheimer's Disease Neuroimaging Initiative (ADNI), and other naturalistic studies, have led to a hypothetical model of disease progression known as the dynamic AD biomarker cascade theory (5), which hypothesizes that AD biomarkers evolve in a sequential, but temporally overlapping manner. According to the hypothesis, amyloid pathology is an early event, leading to tau pathology, followed by neuronal loss and cognitive decline. Additional refinements of the model have been proposed to make it more generalizable to community-based aging populations, including the addition of suspected non-amyloid pathology (SNAP) (e.g., cerebrovascular disease, age-related changes and non-AD tauopathies), as well as the concept of cognitive reserve (e.g. protective factors such as genetics or education), both of which could influence the variability in onset of biomarkers and cognitive decline (2). Although this theoretical model has been operationalized into a categorical scheme for

classifying patients, the system remains descriptive, and makes no assumptions regarding putative causal relationships among biomarkers (6). Understanding how these biomarkers interact, evolve over time and result in cognitive expression of disease will be essential to harness them in a personalized medicine approach to AD diagnosis and treatment. Given the complexity of AD, a rigorous mathematical and computational modeling approach, such as that offered by systems biology, will be a critical component.

The tools of systems biology may be used to incorporate clinical biomarkers of disease progression into a computational model to determine the major patho-etiologic drivers of disease in individual patients and help simulate the effects of potential interventions. One modeling approach, known as causal modeling, refers to an explicitly formulated mathematical description of the biological phenomena of interest, based on existing knowledge, in terms of cause and effect relationships (7). This is in contrast to correlative models which merely describe statistical associations between variables without regard to the mechanism driving the phenomena under investigation. Our goal was to construct and test the feasibility of an initial computational causal model (CCM) of AD biomarker progression, based on the updated dynamic AD biomarker cascade theory (1, 2). This would enable the theory to be tested rigorously with existing data and further refined as new data becomes available.

Methods:

For the construction of the causal model, assumptions about biomarker relationships and temporal course were drawn from prior literature (1, 2, 5). We tested whether the CCM would lead to the predicted biomarker trajectories described in the literature, and whether it would predict failed outcome of anti-amyloid therapy started late in the disease course (8). Figure 1 shows the variables and their relationships in the computational model.

Computational Model Construction:

Figure 1: Diagram depicting causal modeling implementation of the biomarker cascade model in Alzheimer's disease (1, 2). Blue circles represent biomarker quantities. A_β represents amyloid pathology. Its initial value determines when during the lifespan the amyloid cascade begins. τ_ρ represents amyloid-related tau pathology (p-tau). τ_o represents age-related and/or suspected non-Alzheimer pathology (SNAP)-related tauopathy. N represents neuronal dysfunction/loss. C represents cognitive impairment. λ values are growth rate constants, and δ a degredation/clearance rate constant. Amyloidopathy, aging, SNAP (suspected non-Alzheimer's pathology), ApoE status and cognitive reserve are constants that modify the onset of the growth cascades. Anti-amyloid therapy is a function of time. Descriptions of all variables and parameters are listed in Table 1 and Table 2, respectively.© 2017 – 2018 Duke University. All Rights Reserved.

We implemented the above causal model, using the ordinary differential equation (ODE) toolbox in

MATLAB (Mathworks®, Natwick, MA), as system of non-linear first order ODEs to include amyloid

dependent and non-dependent neurodegenerative cascades. The amyloid dependent cascade is

initiated by amyloid beta, A_β, and mediated via phosphorylated tau, τ_ρ. The non-amyloid-dependent

cascades are initiated by comorbidities, e.g., aging and/or suspected non-Alzheimer pathology (SNAP),

either directly, or indirectly via non-amyloid dependent tauopathy, $\tau_{o,}$. Initiation of cognitive decline, C,

is directly determined by neurodegeneration, comorbidities, genetic factors and cognitive reserve. The

equations are as follows (© 2017 – 2018 Duke University. All Rights Reserved):

$$\frac{dA_\beta}{dt} = \lambda_{A_\beta} A_\beta \left(K_{A_\beta} - A_\beta \right) + \lambda_{A_\beta A_0} A_o - \delta_{A_\beta} A_{Rx}(t)$$

$$\frac{d\tau_\rho}{dt} = \lambda_{\tau_\rho A_\beta} A_\beta + \lambda_{\tau_\rho} \tau_\rho \left(K_{\tau_\rho} - \tau_\rho \right)$$

$$\frac{d\tau}{dt} = \lambda_{\tau_\rho A_\beta} A_\beta + \lambda_{\tau_\rho} \tau_\rho \left(K_{\tau_\rho} - \tau_\rho \right) + \lambda_{\tau_o AS} AS$$

$$\frac{dN}{dt} = \lambda_{N\tau_o} (\tau - \tau_\rho) + \lambda_{N\tau_\rho} \tau_\rho + \lambda_N N(K_N - N) + \lambda_{NAS} AS$$

$$\frac{dC}{dt} = \lambda_{CN} NR + \lambda_C C(K_C - C) + \lambda_{CAS} AS + \lambda_{C\varepsilon} \varepsilon$$

where A_β represents amyloid pathology, τ_ρ represents amyloid-related tau pathology (p-tau), τ_o

represents age-related and/or SNAP-related tauopathy, τ represents total tau pathology, defined as the

sum of τ_ρ and τ_o, N represents neuronal dysfunction/loss, and C represents cognitive impairment. λ

defines numerous rate constants. $\lambda_{A_\beta}, \lambda_{\tau_\rho}, \lambda_N$ and λ_C reflect the logistic growth rates of the various

biomarker cascades. The remaining rate constants reflect linear growth rates of the biomarkers and

determine the influence of various factors on the time-of-onset of the subsequent biomarker cascades.

λ_{CN}, for example, is a rate constant that reflects the influence of neurodegeneration, which is modified

by cognitive reserve, for example education level. This, along with comorbid pathologies and genetic risk

alleles, determines the age of onset of the cognitive decline cascade. δ_{A_β} represents a degredation rate

constant for A_β, and in this model, mediates the effects of anti-amyloid therapy. A_o represents

amyloidopathy, A_{Rx} (t) represents a time-dependent function for anti-amyloid therapy, AS represents

aging and/or SNAP, R represents cognitive reserve, and ε represents ApoE allele status. Descriptions of all variables and parameters are listed in Table 1 and Table 2, respectively.

Additional assumptions of the model are as follows: 1) Biomarker cascade growth is implemented via a logistic growth model with carrying capacity K. K is adjusted using a least squares minimization procedure so all biomarkers achieve maximal level of 1 at age 100 years. This is done to configure biomarker curves in a sigmoidal shape with a progressively steeper slope in the right-hand tail for later changing biomarkers, as described in the hypothetical model; 2) At time t=0, A_β is set to a very small number. This is done to initiate the amyloid cascade sometime during the lifespan, even in the absence of amyloidopathy. For simplicity, amyloidopathy A_o, is set to zero in all models, and a slightly larger initial value of A_β is used for the early-onset model, whereas a slightly smaller value is used in the late-onset models. All other biomarker initial values are set to zero, except for total tau in the late onset models, which is set to the minimum biomarker level on the graphs; 3) The minimal biomarker level on the graphs is set to 0.1 to allow for different onset-delays for the sigmoidal shaped biomarker curves that depend upon both biology and biomarker sensitivity; 4) Amyloidopathy, SNAP, aging, and ApoE status are constants across the age span that add linearly to the growth rate of the biomarkers and cause earlier initiation of the amyloid, tau, neurodegenerative and/or cognitive decline cascades; 5) Cognitive reserve is a constant that modifies the effect of neuronal degeneration on the onset of cognitive decline. A lower value is used in the low risk group, and a higher value in the high-risk group. 6) Anti-amyloid therapy, once initiated, is assumed to be maintained throughout the lifespan, and A_{Rx} (t) is simulated as a Heaviside step function, H(n), using the half maximum convention,

$$H[n] = \begin{cases} 0, & n < 0 \\ 1/2, & n = 0 \\ 1, & n > 0 \end{cases}$$

, where n represents the age of initiation of therapy. In the case of anti-amyloid therapy, carrying capacity, K, was determined for all biomarkers in the no-therapy condition, in a natural history context, with δ_{A_β} set to zero. δ_{A_β} was then changed to positive number to simulate the effects of amyloid degradation, with fixed K values based on natural history. This was done to assure that the evolution of biomarkers in the pre-therapy interval was in no way influenced by the administration of therapy later in the course of the disease.

To determine the feasibility of the CCM, we parameterized and tested four versions: 1) familial autosomal dominant AD, 2) late-onset amyloid-first AD, 3) late onset tau-first AD, and 4) anti-amyloid therapy in late-onset amyloid-first AD dementia. In the first three scenarios, the goal was to determine whether manipulating the CCM parameters in a physiological meaningful manner could reproduce biomarker trajectories that closely matched those visually depicted in the literature. In the fourth scenario, we determined whether the model would predict the outcome of recently failed clinical trials of anti-amyloid therapy administered in symptomatic late-onset AD (8).

Results:

Computational model of familial autosomal dominant AD

Figure 2 illustrates that the cascade of early-onset familial AD derived from prior literature (1) (left) matches closely to the output generated by our DCM (right). Specifically, they both demonstrate the initial appearance of amyloid, followed by tau and neurodegeneration, then followed by the onset of cognitive decline. It also shows how cognitive reserve could modify the cascade. Although tau in these figures represents total-tau, in this scenario it is dominated by p-tau (amyloid-related tau). Model parameters are shown in Table 2.

Figure 2: Model of early-onset, autosomal dominant AD. The red, blue and yellow curves represent the evolution of amyloid, tau and neuronal biomarker levels, respectively, over the course of the disease. The green curves represent cognition in two hypothetical high and low risk groups based on low and high cognitive reserve. Our CCM-generated curves (right) closely match the schematic model curves on the left (adapted from (1) with permission) from prior literature.

Computational Model of beta-amyloid-first late-onset AD

The late-onset AD CCM (Figure 3) output shows that amyloid appears first, followed by total tau and neurodegeneration. In this CCM, the arrival of amyloid is delayed compared to that in early-onset AD but reaches detection threshold prior to total tau. The CCM trajectories visually matched those predicted in the literature (1).   Model parameters are shown in Table 2.

Figure 3: CCM of late-onset, amyloid-first AD (right panel) matches closely the pattern hypothesized in the literature (left panel – adapted from (1) with permission). The red, blue and yellow curves represent the evolution of amyloid, tau and neuronal biomarker levels, respectively, over the course of the disease. In the left panel, the blue and yellow lines are combined into a single purple line, per the original theory in which tau was considered a neurodegenerative marker . The green curves represent cognition in two hypothetical high and low risk groups, based on cognitive reserve

Computational model of delayed beta-amyloid late-onset AD

In this CCM (Figure 4), the arrival of total-tau precedes that of amyloid and initiates neurodegeneration, whereas the subsequent appearance of amyloid accelerates this process.   Our CCM mimics a condition described in literature as suspected non-amyloid pathology (SNAP) in some ways (absence of initial amyloid) but illustrates a mixed pathology concept, where amyloid and amyloid-related tau contribute to cognitive decline at later stages. Model parameters are shown in Table 2.

Figure 4: The red, blue and yellow curves represent the evolution of amyloid, tau and neuronal biomarker levels, respectively, over the course of the disease. In the left panel (adapted from (1) with permission), the blue and yellow lines are combined into a single purple line, per the original theory in which tau was considered a neurodegenerative marker . The green curves represent cognition in two hypothetical high and low risk groups, based on cognitive reserve. The CCM (right panel) closely matches the trajectories proposed in the literature.

Simulation of anti-amyloid therapy administered in amyloid-first late-onset AD dementia.

Figure 5 depicts the outcome of our CCM of anti-amyloid therapy when given to amyloid-first late-onset

AD dementia patients after symptom onset. This model output shows no benefit on the onset or slope

of cognitive decline, despite the amyloid level dropping substantially from its peak. Tau levels drop

marginally. The model mimics the results of recent failed anti-amyloid therapy trials in probable AD

dementia (8). In this model, anti-amyloid therapy would have to be given before a hypothetical tipping

point to show benefits on cognition. Model parameters are shown in Table 2.

Figure 5: Left panel shows the untreated condition for comparison, reproduced from Figure 3. Right panel showing the CCM simulation of the effect of anti-amyloid therapy administered in AD after symptom onset. The red curve shows marked decline in brain amyloid levels, the blue line shows a small decrease in tau levels and green lines show there is no significant effect on cognitive decline onset or rate, consistent with the many failed trials.

Discussion:

We have implemented a CCM that incorporates the three clinically available categories of biomarkers to track AD progression, amyloidopathy, tauopathy, and neurodegeneration. The model effectively simulates the temporal evolution of the biomarkers and their relation to cognitive decline as described in previous literature (1, 2, 5), taking into account late verses early onset, the influence of aging and co-occurring non-AD-related brain pathology common in the elderly, and the concept of cognitive resilience to AD pathologic changes. In addition, we simulate the effects of a disease-modifying therapy given late in the disease course, after patients become symptomatic. This CCM was developed both as a means to test existing theories and as a new resource for the field that can be refined as our knowledge advances.

The hypothetical model of the AD pathological cascade, originally published in 2010 (5), and updated in 2013 (2), is based largely on cross-sectional biomarker data, due to limited individual longitudinal biomarker data. It postulates a temporal evolution markers of amyloid pathology, tau pathology, and neurodegeneration, represented as sequential plots of biomarker abnormality over time, leading to cognitive impairment. Three different pathological and neuronal loss scenarios were considered, early onset familial AD, amyloid-first late onset AD, and non-amyloid first (SNAP) late onset AD (1). We created and parameterized a CCM, based on assumptions of underlying biology inherent in the AD pathological cascade, to successfully simulate these three natural history scenarios. Our CCM of early-onset autosomal dominant AD closely matched the temporal order and shape of the biomarker trajectories in the literature schematic (5), and is also supported by empirical data from the longitudinal Dominantly Inherited Alzheimer Network (DIAN) study (9). Our CCM of amyloid-first late onset AD and SNAP-first model of AD also closely simulated the curves postulated in the literature (2). Lastly, our simulation of anti-amyloid therapy in late-onset AD dementia mimicked the negative findings from several failed clinical trials of anti-amyloid therapies (8). Of note, most of the disease modifying

treatment trials in preclinical disease or in mild AD, both several recently completed and ongoing, target

beta-amyloid or tau pathologies.

There are some limitations to our work. First, the hypothetical model (1) on which we built our CCM

may not be entirely accurate. Although, some aspects AD pathological cascade model, for example the

ordering and shape of the biomarker curves, have been validated using data-driven approaches (10, 11),

the model continues to evolve as more natural history and clinical trial data becomes available. Second,

we did not parameterize our models using actual patient biomarker data. Rather, our goal in this work

was to construct and test the feasibility of a CCM that would best match the hypothetical biomarker

trajectories proposed in the literature (1). Subsequent goals will include adjusting parameters based on

longitudinal data and iteratively refining the model itself as new knowledge becomes available. Third,

our CCM is a simplified causal model of biomarkers interacting with each other, an abstraction that does

not model the actual underlying cellular and molecular processes. Prior CCM efforts in AD have modeled

the disease at a molecular and cellular level (12, 13) as well as at a whole-brain, systems level using MRI

and EEG data (14, 15). There have been few CCM approaches that have specifically focused on clinical

AD biomarkers (14, 16), and none have incorporated all three clinically available categories of

biomarkers, amyloidopathy, tauopathy, and neurodegeneration. Overcoming the above limitations will

require large real-world data sets of individual longitudinal biomarker trajectories across the cognitive

continuum as well as integration of genomic, cellular and biomarker knowledge. Such efforts are

underway on an international scale (17-20).

A key strength of the CCM is that it allows for testing underlying causal assumptions in an integrated

fashion, unlike other published correlative mathematical models of clinical biomarkers (10, 11, 21-27)

that treat them independently or fit the data without considering its underlying causal structure. For

example, several studies validated the temporal ordering of biomarkers, without attempting to explain

the underlying disease mechanism by which this temporal ordering arises (10, 11, 26, 27). Causal models allow for testing the effects of non-linear interactions among multiple AD biomarkers and comorbid conditions that cannot be deduced by intuition alone, as well as for predicting response to single and combination therapies. A CCM can be implemented in a "forward" manner to simulate new data or in a "backward" manner, using a Bayesian inversion procedure, to infer the causal architecture of the system based on existing data. This approach has been applied extensively to re-construct mechanistic models of brain function and disease, including AD, from electrophysiologic and imaging data (14, 16, 28). Unlike descriptive models of disease, which can be become increasingly difficult to validate, particularly as datasets of biomarker trajectories become larger and more complex, CCM's can be easily scaled up to increasing degrees of complexity. It is our hope that once a CCM resource for clinical AD biomarkers is created, it will be parameterized based on data from patient studies, expanded and iteratively refined over time. Ultimately such a model would create a global resource for the field to translate existing knowledge, personalize care and accelerate drug discovery for this devastating disorder (29).

References:

1.   Jack CR, Jr., Holtzman DM. Biomarker modeling of Alzheimer's disease. Neuron. 2013;80(6):1347-58. doi: 10.1016/j.neuron.2013.12.003. PubMed PMID: 24360540; PMCID: 3928967.

2.   Jack CR, Jr., Knopman DS, Jagust WJ, Petersen RC, Weiner MW, Aisen PS, Shaw LM, Vemuri P, Wiste HJ, Weigand SD, Lesnick TG, Pankratz VS, Donohue MC, Trojanowski JQ. Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. Lancet neurology. 2013;12(2):207-16. PubMed PMID: 23332364.

3.   Geerts H, Dacks PA, Devanarayan V, Haas M, Khachaturian ZS, Gordon MF, Maudsley S, Romero K, Stephenson D, Brain Health Modeling I. Big data to smart data in Alzheimer's disease: The brain health modeling initiative to foster actionable knowledge. Alzheimer's & dementia : the journal of the Alzheimer's Association. 2016;12(9):1014-21. doi: 10.1016/j.jalz.2016.04.008. PubMed PMID: 27238630.

4.   Ogishima S, Mizuno S, Kikuchi M, Miyashita A, Kuwano R, Tanaka H, Nakaya J. A map of Alzheimer's disease-signaling pathways: a hope for drug target discovery. Clin Pharmacol Ther. 2013;93(5):399-401. Epub 2013/03/21. doi: 10.1038/clpt.2013.37. PubMed PMID: 23511713.

5.   Jack CR, Jr., Knopman DS, Jagust WJ, Shaw LM, Aisen PS, Weiner MW, Petersen RC, Trojanowski JQ. Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. Lancet neurology. 2010;9(1):119-28. doi: 10.1016/S1474-4422(09)70299-6. PubMed PMID: 20083042; PMCID: 2819840.

6.   Jack CR, Jr., Bennett DA, Blennow K, Carrillo MC, Feldman HH, Frisoni GB, Hampel H, Jagust WJ, Johnson KA, Knopman DS, Petersen RC, Scheltens P, Sperling RA, Dubois B. A/T/N: An unbiased descriptive classification scheme for Alzheimer disease biomarkers. Neurology. 2016;87(5):539-47. doi: 10.1212/WNL.0000000000002923. PubMed PMID: 27371494; PMCID: 4970664.

7.   Voit EO. A First Course in Systems Biology. New York, NY: Garland Science,Taylor & Francis Group LLC; 2013. p. 19-50.

8.   Cummings JL, Morstorf T, Zhong K. Alzheimer's disease drug-development pipeline: few candidates, frequent failures. Alzheimers Res Ther. 2014;6(4):37. Epub 2014/07/16. doi: 10.1186/alzrt269. PubMed PMID: 25024750; PMCID: PMC4095696.

9.   Bateman RJ, Xiong C, Benzinger TL, Fagan AM, Goate A, Fox NC, Marcus DS, Cairns NJ, Xie X, Blazey TM, Holtzman DM, Santacruz A, Buckles V, Oliver A, Moulder K, Aisen PS, Ghetti B, Klunk WE, McDade E, Martins RN, Masters CL, Mayeux R, Ringman JM, Rossor MN, Schofield PR, Sperling RA, Salloway S, Morris JC, Dominantly Inherited Alzheimer N. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. The New England journal of medicine. 2012;367(9):795-804. doi: 10.1056/NEJMoa1202753. PubMed PMID: 22784036; PMCID: 3474597.

10.   Jedynak BM, Lang A, Liu B, Katz E, Zhang Y, Wyman BT, Raunig D, Jedynak CP, Caffo B, Prince JL, Alzheimer's Disease Neuroimaging I. A computational neurodegenerative disease progression score: method and results with the Alzheimer's disease Neuroimaging Initiative cohort. Neuroimage. 2012;63(3):1478-86. doi: 10.1016/j.neuroimage.2012.07.059. PubMed PMID: 22885136; PMCID: 3472161.

11.   Young AL, Oxtoby NP, Daga P, Cash DM, Fox NC, Ourselin S, Schott JM, Alexander DC, Alzheimer's Disease Neuroimaging I. A data-driven model of biomarker changes in sporadic Alzheimer's disease. Brain : a journal of neurology. 2014;137(Pt 9):2564-77. doi: 10.1093/brain/awu176. PubMed PMID: 25012224; PMCID: 4132648.

12.   Geerts H, Spiros A, Roberts P, Carr R. Towards the virtual human patient. Quantitative Systems Pharmacology in Alzheimer's disease. Eur J Pharmacol. 2017;817:38-45. Epub 2017/06/07. doi: 10.1016/j.ejphar.2017.05.062. PubMed PMID: 28583429.

13.	Hao W, Friedman A. Mathematical model on Alzheimer's disease. BMC Systems Biology. 2016;10:108. doi: 10.1186/s12918-016-0348-2. PubMed PMID: PMC5116206.

14.	Iturria-Medina Y, Carbonell FM, Sotero RC, Chouinard-Decorte F, Evans AC. Multifactorial causal model of brain (dis)organization and therapeutic intervention: Application to Alzheimer's disease. NeuroImage. 2017;152:60-77. doi: https://doi.org/10.1016/j.neuroimage.2017.02.058.

15.	Penny W, Iglesias-Fuster J, Quiroz YT, Lopera FJ, Bobes MA. Dynamic Causal Modeling of Preclinical Autosomal-Dominant Alzheimer's Disease. J Alzheimers Dis. 2018. Epub 2018/03/23. doi: 10.3233/JAD-170405. PubMed PMID: 29562504.

16.	Iturria-Medina Y, Sotero RC, Toussaint PJ, Mateos-Perez JM, Evans AC, Alzheimer's Disease Neuroimaging I. Early role of vascular dysregulation on late-onset Alzheimer's disease based on multifactorial data-driven analysis. Nature communications. 2016;7:11934. doi: 10.1038/ncomms11934. PubMed PMID: 27327500; PMCID: 4919512.

17.	Bertram L, McQueen MB, Mullin K, Blacker D, Tanzi RE. Systematic meta-analyses of Alzheimer disease genetic association studies: the AlzGene database. Nature genetics. 2007;39(1):17-23. Epub 2006/12/29. doi: 10.1038/ng1934. PubMed PMID: 17192785.

18.	Ellis KA, Bush AI, Darby D, De Fazio D, Foster J, Hudson P, Lautenschlager NT, Lenzo N, Martins RN, Maruff P, Masters C, Milner A, Pike K, Rowe C, Savage G, Szoeke C, Taddei K, Villemagne V, Woodward M, Ames D, Group AR. The Australian Imaging, Biomarkers and Lifestyle (AIBL) study of aging: methodology and baseline characteristics of 1112 individuals recruited for a longitudinal study of Alzheimer's disease. Int Psychogeriatr. 2009;21(4):672-87. Epub 2009/05/28. doi: 10.1017/S1041610209009405. PubMed PMID: 19470201.

19.	Morris JC, Aisen PS, Bateman RJ, Benzinger TL, Cairns NJ, Fagan AM, Ghetti B, Goate AM, Holtzman DM, Klunk WE, McDade E, Marcus DS, Martins RN, Masters CL, Mayeux R, Oliver A, Quaid K, Ringman JM, Rossor MN, Salloway S, Schofield PR, Selsor NJ, Sperling RA, Weiner MW, Xiong C, Moulder KL, Buckles VD. Developing an international network for Alzheimer research: The Dominantly Inherited Alzheimer Network. Clin Investig (Lond). 2012;2(10):975-84. Epub 2012/11/10. doi: 10.4155/cli.12.93. PubMed PMID: 23139856; PMCID: PMC3489185.

20.	Mueller SG, Weiner MW, Thal LJ, Petersen RC, Jack CR, Jagust W, Trojanowski JQ, Toga AW, Beckett L. Ways toward an early diagnosis in Alzheimer's disease: the Alzheimer's Disease Neuroimaging Initiative (ADNI). Alzheimer's & dementia : the journal of the Alzheimer's Association. 2005;1(1):55-66. Epub 2007/05/04. doi: 10.1016/j.jalz.2005.06.003. PubMed PMID: 17476317; PMCID: PMC1864941.

21.	Donohue MC, Jacqmin-Gadda H, Le Goff M, Thomas RG, Raman R, Gamst AC, Beckett LA, Jack CR, Jr., Weiner MW, Dartigues JF, Aisen PS, Alzheimer's Disease Neuroimaging I. Estimating long-term multivariate progression from short-term data. Alzheimer's & dementia : the journal of the Alzheimer's Association. 2014;10(5 Suppl):S400-10. doi: 10.1016/j.jalz.2013.10.003. PubMed PMID: 24656849; PMCID: 4169767.

22.	Guerrero R, Schmidt-Richberg A, Ledig C, Tong T, Wolz R, Rueckert D, Alzheimer's Disease Neuroimaging I. Instantiated mixed effects modeling of Alzheimer's disease markers. Neuroimage. 2016;142:113-25. doi: 10.1016/j.neuroimage.2016.06.049. PubMed PMID: 27381077.

23.	Jack CR, Jr., Wiste HJ, Lesnick TG, Weigand SD, Knopman DS, Vemuri P, Pankratz VS, Senjem ML, Gunter JL, Mielke MM, Lowe VJ, Boeve BF, Petersen RC. Brain beta-amyloid load approaches a plateau. Neurology. 2013;80(10):890-6. Epub 2013/03/01. doi: 10.1212/WNL.0b013e3182840bbe. PubMed PMID: 23446680; PMCID: PMC3653215.

24.	Oxtoby NP, Young AL, Fox NC, Daga P, Cash DM, Ourselin S, Schott JM, Alexander DC. Learning imaging biomarker trajectories from noisy Alzheimer's disease data using a Bayesian multilevel model. Lecture Notes in Computer Science. 2014;8677:85-94.

25.	Villemagne VL, Burnham S, Bourgeat P, Brown B, Ellis KA, Salvado O, Szoeke C, Macaulay SL, Martins R, Maruff P, Ames D, Rowe CC, Masters CL, Australian Imaging B, Lifestyle Research G. Amyloid

beta deposition, neurodegeneration, and cognitive decline in sporadic Alzheimer's disease: a prospective cohort study. Lancet neurology. 2013;12(4):357-67. Epub 2013/03/13. doi: 10.1016/S1474-4422(13)70044-9. PubMed PMID: 23477989.

26.   Fonteijn HM, Modat M, Clarkson MJ, Barnes J, Lehmann M, Hobbs NZ, Scahill RI, Tabrizi SJ, Ourselin S, Fox NC, Alexander DC. An event-based model for disease progression and its application in familial Alzheimer's disease and Huntington's disease. Neuroimage. 2012;60(3):1880-9. doi: 10.1016/j.neuroimage.2012.01.062. PubMed PMID: 22281676.

27.   Jedynak BM, Liu B, Lang A, Gel Y, Prince JL, Alzheimer's Disease Neuroimaging I. A computational method for computing an Alzheimer's disease progression score; experiments and validation with the ADNI data set. Neurobiol Aging. 2015;36 Suppl 1:S178-84. Epub 2014/12/03. doi: 10.1016/j.neurobiolaging.2014.03.043. PubMed PMID: 25444605; PMCID: PMC4267989.

28.   Kiebel SJ, Garrido MI, Moran R, Chen CC, Friston KJ. Dynamic causal modeling for EEG and MEG. Hum Brain Mapp. 2009;30(6):1866-76. Epub 2009/04/11. doi: 10.1002/hbm.20775. PubMed PMID: 19360734.

29.   Voit EO. Systems Biology in Medicine and Drug Development. In: Science G, editor. A First Course in Systems Biology: Taylor & Francis Group, LLC; 2013. p. 347-72.

Table 1: Model Variables

Variable	Description
A_β	amyloid beta pathology
τ_ρ	amyloid-related phospho tau pathology
τ_o	SNAP-related tau pathology
N	neuronal degeneration
C	cognitive impairment

SNAP = suspected non-Alzheimer's pathology

Table 2: Model parameters for early- and late-onset AD and anti-amyloid therapy scenarios

Parameter	Description	Early onset	Late onset amyloid-first	Late onset SNAP-first	Anti-amyloid Rx (Late onset amyloid-first)
$A_{\beta 0}$	amyloid beta pathology – initial value	0.05	0.01*	0.01*	0.01*
τ_{p0}	tau (amyloid-related) pathology– initial value	0	0	0	0
τ_{o0}	tau (SNAP-related) pathology – initial value	0	0.05*	0.05*	0.05*
N_0	neurodegeneration – initial value	0	0	0	0
C_0	cognitive impairment – initial value	0	0	0	0
$K_{A\beta}$	amyloid beta pathology - carrying capacity	1	1	1	1
$K_{\tau p}$	tau (amyloid-related) - carrying capacity	1	1	1	1
K_C	cognitive impairment - carrying capacity	1	1	1	1
K_N	neurodegeneration - carrying capacity	1	1	1	1
$\lambda_{A\beta}$	amyloid cascade - growth rate	0.08	0.12*	0.1*	0.12*
$\lambda_{A\beta AO}$	amyloid pathology from amyloidopathy - growth rate	0	0	0	0
$\delta_{A\beta}$	amyloid degradation/clearance rate	0	0	0	0.04*
$\lambda_{\tau p\beta}$	tau (amyloid-related) from amyloid – growth rate	0.025	0.025	0.025	0.025
$\lambda_{\tau p}$	tau (amyloid-related) pathol cascade – growth rate	0.05	0.05	0.05	0.05
$\lambda_{\tau oAS}$	tau (SNAP-rel) path from aging/SNAP - growth rate	0.002	0.002	0.002	0.002
λ_C	cognitive impairment cascade - growth rate	0.05	0.05	0.05	0.05
λ_{CN}	cognitive impair from neurodegen - growth rate	0.001	0.001	0.001	0.001
λ_{CAS}	cognitive impairment from aging/SNAP - growth rate	0	0	0	0
λ_{CE}	cognitive impairment from genetic risk - growth rate	0.001	0.001	0.001	0.001
λ_N	neurodegeneration cascade growth rate	0.05	0.05	0.05	0.05
$\lambda_{N\tau p}$	neurodegen from tau (amyloid-related) - growth rate	0.025	0.025	0.025	0.025
λ_{NTo}	neurodegen from tau (SNAP-related) - growth rate	0.0075	0.0075	0.0075	0.0075
λ_{NAS}	neurodegeneration from aging/SNAP - growth rate	0	0	0	0
AO	amyloidopathy	0	0	0	0
ARx	age of onset of anti-amyloid therapy	0	0	0	65*
AS	aging, SNAP	0	1*	2*	1*
E	apoE allele genetic risk	0	0	0	0
R_1	cognitive reserve (low risk)	1	1	1	1
R_2	cognitive reserve (high risk)	25	25	25	25

* Signifies a change in parameters from those of the early-onset AD model.

SNAP = suspected non-Alzheimer's pathology

Conflicts of Interest:

The authors disclose no relevant conflicts of interest.

Funding Statement:

The authors performed this work as part of their employment at Duke University Medical System and

Penn State University, and received no specific funding from other sources.