Studies on the mechanism of membrane mediated general anesthesia.

Mahmud Arif Pavel[1,2], E. Nicholas Petersen[1,2], Hao Wang[1,2], Richard A. Lerner[3], and Scott B.

Hansen[1,2,*]

[1]Department of Molecular Medicine, [2]Department of Neuroscience, The Scripps Research

Institute, Jupiter, Florida 33458, USA.

[3]Department of Chemistry, The Scripps Research Institute, La Jolla, California 92037, USA

*Correspondence: shansen@scripps.edu

ABSTRACT

Inhaled anesthetics are a chemically diverse collection of hydrophobic molecules that robustly

activate TWIK related K+ channels (TREK-1) and reversibly induce loss of consciousness. For a

hundred years anesthetics were speculated to target cellular membranes, yet no plausible

mechanism emerged to explain a membrane effect on ion channels. Here we show that inhaled

anesthetics (chloroform and isoflurane) activate TREK-1 through disruption of palmitate-

mediated localization of phospholipase D2 (PLD2) to lipid rafts and subsequent production of

signaling lipid phosphatidic acid (PA). Catalytically dead PLD2 robustly blocks anesthetic TREK-

1 currents in cell patch-clamp. Localization of PLD2 renders the anesthetic-insensitive TRAAK

channel sensitive. General anesthetics chloroform, isoflurane, diethyl ether, xenon, and propofol

disrupt lipid rafts and activate PLD2. In the whole brain of flies, anesthesia disrupts rafts and

PLD[null] flies resist anesthesia. Our results establish a membrane mediated target of inhaled

anesthesia and suggest PA helps set anesthetic sensitivity *in vivo*.

INTRODUCTION

In 1846 William Morton demonstrated general anesthesia with inhaled anesthetic diethyl ether (*1*). For many anesthetics (but not all), lipophilicity is the single most significant indicator of potency; this observation is known as the Meyer-Overton correlation (*2*, *3*). This correlation, named for its discoverers in the late 1800's, and the chemical diversity of anesthetics (Supplementary Fig. S1a) drove anesthetic research to focus on perturbations to membranes as a primary mediator of inhaled anesthesia (*3*). Over the last two decades, enantiomer selectivity of anesthetics suggested a chiral target and direct binding to ion channels emerged (*4*). But the possibility of a membrane mediated effect has remained (*5–7*).

In 1997 a theory emerged suggesting that disruption of ordered lipids surrounding a channel could activate the channel (*8*). Disruption of ordered lipids (sometimes referred to as lipid rafts (*9*)) allows phospholipase D2 (PLD2) to translocate out of lipid domains and experience a new chemical environment (*10*) (Supplementary Fig. S1b-c). PLD2 actives Twik related potassium channel-1 (TREK-1). If inhaled anesthetics can disrupt lipid domains to activate a channel, this would constitute a mechanism distinct from the usual receptor-ligand interaction and establish a definitive membrane mediated mechanism for an anesthetic.

TREK-1 is an anesthetic-sensitive two-pore-domain potassium (K2P) channel. Xenon, diethyl ether, halothane, and chloroform robustly activate TREK-1 at concentrations relevant to their clinical use (*11*, *12*) and genetic deletion of TREK-1 decreases anesthesia sensitivity in mice (*13*). Here we show inhaled anesthetics disrupt palmitate-mediated localization in the membranes of cultured neuronal, muscle cells, and in the whole brain of anesthetized flies. The disruption releases PLD2 from muscle and neuronal cells which signal downstream to activate TREK-1 channels. This result establishes the membrane as a pertinent target of inhaled anesthetics.

Anesthetics perturb nanoscale lipid heterogeneity (GM1 domains).

The best studied lipid domains contain saturated lipids cholesterol and sphingomyelin (e.g.

monosialotetrahexosylganglioside1 (GM1)) (see Supplementary Fig. S1b-c) and bind cholera

toxin B (CtxB) with high affinity (*9*). Anesthetics lower the melting temperature and expand GM-

1 domains in artificial membranes and membrane vesicles (*14–16*). Lipids are thought to exist in

heterogenous mixtures with features smaller than 100 nm that are best characterized with super-

resolution imaging (e.g., direct stochastical optical reconstruction microscopy (dSTORM) and

stimulated emission depletion (STED)) (*17–20*). Palmitoylation, a posttranslational modification

that covalently attaches a 16 carbon lipid, localizes proteins to GM1 lipids (*21*) (Supplementary

Fig. S1c) including many ion channels (*22*). We used dSTORM to test the hypothesis that

anesthetics harness lipid heterogeneity to gate TREK-1 (*23*).

To test an anesthetic specific perturbation to GM1 lipids, we treated neuroblastoma 2A (N2A)

cells with anesthetic chloroform at 1mM concentration and monitored fluorescent CtxB clustering

by dSTORM (Fig. 1a). Chloroform strongly increased both the diameter and area of GM1 domains

in the cell membrane (Fig. 1b-c, Supplementary Fig. S2a). Isoflurane (1 mM) behaved similar

chloroform (Fig. 1a-c). The Ripley's radius, a measure of space between domains, decreased

dramatically for both chloroform and isoflurane (Supplementary Fig. S2b) suggesting the domains

expand (*24*) and possibly divide as the number of rafts per cell increased (Fig. 1d-e,

Supplementary Fig. S2c). Methyl-β-cyclodextrin (MβCD), a chemical that disrupts GM1 domains

by removing cholesterol (*10*), reduced the total number of domains (Fig.1d). Binning the domains

into small (0-150 nm) and large (150-500nm) revealed a clear shift from small to large domains

in the presence of inhaled anesthetics and revealed the opposite effect after MβCD treatment

(Supplementary Fig. S2d). Similar results were obtained in C2C12 myoblasts (muscle cells);

chloroform strongly increased both the diameter and area of GM1 domains. (Supplementary Fig.

S2e-h)

Mechanism of anesthetic sensitivity in TREK-1 channels

To confirm the observed disruption is capable of producing a biological signal, we studied TREK-1 activation by inhaled anesthetics. Activation of TREK-1 by inhaled anesthetics, was previously shown to require a disordered loop in the channel's C-terminus (*11*) (Supplementary Fig. S3a-b). The enzyme phospholipase D2 (PLD2) also binds to and activates through the same C-terminal region in TREK-1 (*25*). PLD2 is palmitoylated at cysteines near its Pleckstrin homology (PH) domain, which is required to localize it to GM1 lipids (*26*). The PH domain also binds PIP_2 which opposes the localization by palmitoylation (Supplementary Fig. S1c). Disruption of GM1 lipids by mechanical force activates PLD2 by substrate presentation—the enzyme translocated from GM1 to PIP_2 lipids near its substrate (*10*). If anesthetics disrupt GM1 domains, then we expect PLD2 or a complex of PLD2 and TREK-1 to translocate and activate TREK-1. If correct, the PLD2 dependent TREK-1 activation would confirm lipid disruption as a component of TREK-1 anesthetic sensitivity and a membrane mediated pathway.

To test our hypothesis and the role of lipid heterogeneity in anesthetic sensitivity of TREK-1, we applied chloroform to HEK293 cells over-expressing TREK-1 and a catalytically dead K758R PLD2 mutant (xPLD2) that blocks anionic lipid production (e.g. PA and PG) (*27*). We over expressed TREK-1 in HEK293 cells to avoid the confounding effects of endogenous channels other than TREK-1 found in N2A and C2C12 cells. We found xPLD2 blocked all detectible chloroform specific current (Fig. 2a-c). Cells not transfected with TREK-1 had no observable current and were unaffected by chloroform treatment (Supplementary Fig. S3c) showing the xPLD's effect is specific to TREK-1 currents. We confirmed xPLD overexpression with western blot analysis (Supplementary Fig. S3d). This result suggests a two-step mechanism for anesthetic action on TREK-1 channels. First, anesthetics disrupt GM1 domains releasing PLD2 and second, the enzyme binds to the C-terminus of TREK-1 and they translocates to substrate and activates

the channel through increased local concentration of anionic lipid (Fig. 2d). Similarly, PLD could

be pre-bound to TREK-1 and translocation as a complex to PIP_2 domains and PLD substrate.

Transfer of anesthetic sensitivity to TRAAK.

TWIK-related arachidonic acid-stimulated K+ channel (TRAAK) is an anesthetic insensitive

homolog of TREK-1 (Supplementary Fig. S3e). Interestingly, native TRAAK is also insensitive to

PLD2 (*25*). However, concatenating PLD2 to the N-terminus maximally activates TRAAK and

introduction of the PLD2 binding domain from TREK-1 renders TRAAK PLD2 sensitive (*25*). If

PLD2 is responsible for anesthetic sensitivity in TREK-1, we reasoned we could render TRAAK

anesthetic sensitive by introducing the PLD2 binding site into the C-terminus of TRAAK (Fig. 3a).

We over expressed the previously characterized PLD2 sensitive TRAAK chimera (*25*)

(TRAAK/ctTREK) in HEK cells. As expected, in the presence of 1mM chloroform, TRAAK/ctTREK

robustly responded to chloroform (Fig. 3b-d). To confirm the response is due to PLD2 localization

and not a direct interaction of the anesthetic with a structural feature of the TREK-1 C-terminus,

we over expressed the chimera with xPLD2 and found chloroform had no effect on the channel.

This result suggests the disordered C-terminus exerts its anesthetic effect through binding to

PLD2 and not direct binding of anesthetic to the C-terminus (Fig. 3e).

The lack of TREK-1 current in the presence of anesthetic (Fig 2b-c) suggests direct binding of

anesthetic is insufficient to activate the channel absent PLD2 activity. This result opposes what

was previously thought, that the anesthetic directly activates TREK-1(*28*). To confirm our result

and distinguish the contribution of an indirect anesthetic effect from a direct one, we tested the

contribution of direct binding to TREK-1 anesthetic sensitivity in a flux assay. Ion channels are

routinely expressed, purified, and assayed in vesicles and purified TREK-1 channels robustly

recapitulates small molecules binding to channels using a flux assay in a cell free system (*29–*

31).

Functionally reconstituted TREK-1 in 16:1 phosphatidylcholine (PC) liposomes with 18:1

phosphatidylglycerol (PG) (85:15 mol% ratio) was unaffected by chloroform or isoflurane (1 mM)

(Supplementary Fig. S3f-g). Changing the lipids and ratios to 18:1 PC/18:1 PG (90/10 mol%)(*30*)

also had no effect (data not shown). To assure that the channel was properly reconstituted and

in conditions capable of increased potassium flux, we reconstituted a mutant TREK-1 with double

cysteines that covalently lock TREK-1 in the activated state (*30*, *32*). Compared to the open

TREK-1 control, inhaled anesthetics failed to activate TREK-1 (Supplementary Fig. S3f-g). We

showed in a previous study, using the same assay, a clear direct inhibition with local anesthetics

(*31*).

Anesthetics displace PLD2 out of GM1 lipids.

For raft disruption to be biologically significant it must transduce the disruption into a biological

signal. To test our hypothesis that the disruption gives rise to PLD2 activity and TREK-1 activation,

we directly imaged palmitate-mediated PLD2 localization using dSTORM imaging and anesthetic

induced translocation of PLD2 out of GM1 lipids. We measure translocation, by calculating a

pairwise correlation of PLD2 with CTxB before and after anesthetic treatment. Pairwise correlation

avoids potential artifacts from over sampling, and the limits of the user selected parameters used

when determining raft size (Supplementary Fig. S4). Both N2A and C2C12 endogenously express

TREK-1 and PLD2 (see methods) allowing us to characterize their translocation under

endogenous promoters. TREK-1 is not palmitoylated, but palmitoylation could sequester PLD2

away from TREK-1 (Supplementary Fig. S1b-c) or sequester the pre-bound PLD2/TREK-1

complex into GM1 domains away from PLD's activating lipid (PC) or both (*10*).

Treating the N2A cells with chloroform or isoflurane (1 mM), caused PLD2 to translocate away

from GM1 lipids (Fig 4a). Pair correlation analysis showed PLD2 strongly associated with GM1

domains prior to anesthetic treatment (Fig 4b, grey trace) but only weakly associated after

treatment (green traces). This highly significant anesthetic-induced change in PLD2 localization

was true across all sizes of GM1 domains ((Fig. 4b-c), pair correlation analysis) suggesting the

energetics of PLD2 partitioning into GM1 lipids changed, not merely a change in the size of the

domain. The anesthetic-specific translocation was similar in magnitude to MβCD stimulated

translocation (Fig. 2b-c) and implies that anesthetic expansion of GM-1 domains has the same

effect on palmitate-mediated localization. Expansion is therefore a form of raft disruption (Fig. 4d).

We obtained similar pair correlation results in C2C12 cells treated with chloroform

(Supplementary Fig. S5). The analysis of pairwise correlation with and without anesthetic, that

are otherwise treated identical, also largely rules out artifacts from antibody or toxin labeling or

permeabilization as the differences in PLD2 localization.

Anesthetics activate PLD2 through raft disruption.

If the membrane is a general mechanism for TREK-1 mediated anesthesia, then most known

activators of TREK-1 should also activate PLD2. We tested enzymatic activation of PLD2 by

treating live cells with a spectrum of chemically diverse inhaled anesthetics and monitoring activity

using an assay that couples PLD's choline release to a fluorescent signal (*10*) (Fig. 5a-e). Diethyl

ether, chloroform, isoflurane, and xenon all significantly activated PLD2 in N2A (Fig. 5a-e,h) and

C2C12 cells (Supplementary Fig. S6a-d,f). Isoflurane had the greatest effect (Fig. 5h) in N2A cells

and chloroform had the greatest effect among inhaled anesthetics in C2C12 cells (Supplementary

Fig. S6b,f). The dose response of chloroform and isoflurane reveals an EC50 of 1.0 ± 0.56 mM

and $0.8 \pm$ mM respectively (Fig. 5i) in N2A cells. The Hill slopes were 2.4 ± 2.8 and 1.3 ± 0.89

respectively. Ketamine, an injectable NMDA receptor specific anesthetic (*33*) and F6, a non-

immobilizer that defies the Meyer-Overton rule, had no effect on PLD activity, as expected (Fig.

5f-h). Dose responses at low concentration were highly variable making the determination of the

Hill slope difficult to determine. The PLD2 assay also lacks the ability to mimicked partitioning into

the membrane from continuous inhalation.

We also tested the injectable general anesthetics propofol (50 µM) (*4*). Propofol robustly activated

PLD2 in N2A cells (Fig. 5d, h). If our mechanism is correct, then propofol should lead to TREK-1

activation. As predicted, propofol, robustly increased TREK-1 currents (Fig 5k) in whole cell patch-

clamp. Propofol's effect was significant (Fig. 5m, $p=0.017$, two tailed Student's t test,) and co-

transfection of xPLD2 with TREK-1 completely blocked the propofol specific current (Fig. 5l).

Hence, PLD2 activity predicts channel function and this result suggests propofol works through

the same pathway as inhaled anesthetics to activate TREK-1 albeit with less potency in C2C12

(Supplementary Fig. S6d,f).

***In vivo* PLD dependent anesthetic sensitivity.**

If anesthetics disrupt membranes to activate PLD *in vivo*, then blocking PLD could block or

attenuate anesthesia in an animal. Establishing anesthetic sensitivity of PLD *in vivo* would also

establish the membrane and PA signaling as important up stream mediators of anesthetic action

independent of a channel. To monitor sedation *in vivo*, we recorded single-animal measurements

(activity and position) of *Drosophilae melanogaster* (fruit fly) in a vertically mounted chamber (Fig.

6a, Petersen unpublished data). Flies without functional PLD (PLD[null]) and wildtype (with PLD)

were subjected to chloroform vapor and monitored for sedation. Sedation was determined by 5

minutes of continuous inactivity with a vertical position at the bottom of the fly chamber (Fig. 6a).

Sedation of PLD[null] flies with 2.8 mM chloroform required almost twice the exposure as wildtype

flies (~600 vs. 350 sec, $p<0.0001$) indicating a highly significant resistance to anesthesia in PLD[null]

(Fig. 6b). Almost all flies were anesthetized before the first PLD^{null} fly passed out. The

concentration of chloroform is that of the vapor not the concentration in the animal. All flies

eventually lost consciousness suggesting PLD sets the threshold rather than dictates anesthesia.

Disruption of GM1 domains in whole brains

To confirm the presence of GM1 domains in brain tissue, we dissected whole brains from adult

flies and labeled them with CTxB and an antibody against the pan-neuronal marker Elav. Confocal

imaging (Figure 6c) of CTxB-labeled lipids (GM1, green) showed robust expression of GM1 lipids

throughout the fly brain. High concentrations were observed on the membrane of the cell bodies

(Figure 6c, zoom) as expected for GM1. GM1 was also observed in clusters, confirming that lipid

domains also exist in the CNS of flies, making them similar in organization to our cell culture.

Lastly, we asked if anesthetic disruption of lipid rafts observed in cell culture also exists in the

whole brain of an anesthetized animal. To test this, we anesthetized adult flies with chloroform,

dissected their brains and characterized GM1 domains in those brains compared with a no-

anesthesia control using dSTORM super resolution imaging. Consistent with cell culture, GM1

domains were expanded in fly brains treated with anesthetic (Fig. 6e). The number of domains

was found to decrease by ~10%.

DISCUSSION

We conclude that the membrane is a target of inhaled anesthetics and that PA and disruption of

palmitate mediated lipid localization contributes to anesthesia *in vivo*. Our proposed model for

TREK-1 in mammalian cells is consistent with most known properties of inhaled anesthetics

(Supplementary Fig. S5a) despite it utilizing an indirect mechanism through PLD2 and lipid

binding sites (*11*, *12*, *28*). The disruption of palmitoylate mediated localization from a lipid

nanodomain nicely explains how the C-terminus renders a channel anesthetic sensitive when

the domain is highly charged, devoid of structure, and has no obvious hydrophobicity expected

to bind an anesthetic (Supplementary Fig. S5b).

TREK-1 is not conserved in flies precluding direct downstream testing in the whole animal.

These experiments will need to be done in a mammalian model. Nonetheless, our model does

not rely on a single channel, rather we expect the lipid to contribute to central regulation of

excitability through direct binding to multiple ion channels. Recently PA emerged as class of

lipid-regulated ion channel modulating excitability and pain (*23, 34*). PA's importance is directly

supported by the finding that a single protein that modulates PA production, PLD, dramatically

shifted the anesthetic threshold in an animal (Fig. 6b). The conclusion is also indirectly

supported by the observation of GM1 domains throughout the brain of flies (Fig. 6c), the

observed disruption of GM1 domains in the whole brain tissue of chloroform treated flies (Fig.

6e), and the modular transfer of anesthetic sensitivity to TRAAK by localization of a PA

producing enzyme (Fig. 3b).

The importance of PA is evident by the fact that it is upstream of TREK-1. The latency (time

delay) of PLD mixing in these experiments were dictated by our relatively slow application of

anesthetic by gravity flow see methods). However theoretical estimates of latency based on

distance of PLD from its substrate suggest a PLD dependent latency of ~650 μs (*10*), much

faster than our application of anesthetic ((>10 sec). In rafts that are mechanically disrupted, we

have measured a PLD2 latency of TREK-1 activation with an upper limit of 2.1 ms (Petersen et.

al unpublished data); we expect disruption by anesthetic is similar. PA can also affect

membrane curvature and hydrophobic mismatch, and anesthetics can in theory affect these

properties to regulate TREK-1(*35*), but these potential mechanism, if they contribute, appear to

be insignificant compared to anesthetic activation of PLD (Fig. 2a-b).

Over the past decade anesthetic research has focused on direct binding of inhaled anesthetics

to allosteric sites on proteins (*2*, *4*, *36*). This was not the case for TREK-1, but our results do not

preclude direct binding for other channels. In some instances, the direct binding may be

competition of anesthetics with lipid regulatory sites (*23*) which would be a hybrid protein/lipid

mechanism. Contributing to a protein based narrative, anesthetics are also known to be

enantiomer selective (*37–39*). Interestingly ordered lipids are also chiral and the ability of an

anesthetic to partition into ordered domains will need to be considered in light of the mechanism

presented here.

Many channels are directly palmitoylated (*22*) and they could be modulated by a change in ion

channel localization. For example, the anesthetic channel γ-aminobutyric acid A receptor

(GABA$_A$R) gamma subunit is palmitoylated (*22*) and the alpha subunit was recently shown to

bind PIP$_2$ (*40*). Hence GABA$_A$R is comprised of precisely the same features that render PLD2

anesthetics sensitivity. If and how these feature function in GABA$_A$R is not known. Many

important signaling molecules are palmitoylated including tyrosine kinases, GTPases, CD4/8,

and almost all G-protein alpha subunits (*41*). Anesthetic disruption of palmitate mediated

localization of these proteins likely contributes to modulation channels in a membrane

dependent manner. The overall contribution of the membrane will need to be studies for each

channel and regulator independently.

Lastly, we considered the biophysical effect of anesthetics on the bulk membranes or 'non-raft'

membranes. We saw very little effect of clinical concentration of anesthetics on TREK-1

reconstituted into (DOPC) liposomes in our flux assay (Supplementary Fig. 3a-b), a mimic of

bulk lipids. This result agrees with previous studies that showed the effect of anesthetics on bulk

lipids is insufficient to activate a channel (*42*) at clinical concentrations despite the fact that

anesthetics fluidize and thin membranes (*43*). TREK-1 is very sensitive to membrane thickness

(*44*). It's possible we failed to test an optimal thickness that is responsive in artificial systems,

however, the fact that xPLD2 blocked all detectible anesthetic currents in whole cells suggests,

in a biological membrane, domain disruption and TREK-1 and PLD2 translocation are the

primary mechanism for anesthetic activation of TREK-1, not thinning of bulk lipids.

Our data show a pathway where anionic lipids are central mediators of anesthetic action on ion

channels and these results suggest lipid regulatory molecules and lipid binding sites in channels

may be effective targets for treating nervous system disorders and understanding the thresholds

that govern intrinsic nerve cell excitability. Thus, the system we describe here obviously did not

evolve to interact with inhaled anesthetics and a search as to what the endogenous analogue

that activates this physiological system is warranted.

REFERENCES

1. V. M. Seutin, Mechanisms of actions of inhaled anesthetics. *N. Engl. J. Med.* **349**, 909–910; author reply 909-910 (2003).

2. B. W. Urban, M. Bleckwenn, M. Barann, Interactions of anesthetics with their targets: Non-specific, specific or both? *Pharmacol. Ther.* **111**, 729–770 (2006).

3. A. Kopp Lugli, C. S. Yost, C. H. Kindler, Anaesthetic mechanisms: update on the challenge of unravelling the mystery of anaesthesia. *Eur. J. Anaesthesiol.* **26**, 807–20 (2009).

4. N. P. Franks, General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. *Nat. Rev. Neurosci.* **9**, 370–86 (2008).

5. J. M. Sonner, A hypothesis on the origin and evolution of the response to inhaled anesthetics. *Anesth. Analg.* **107**, 849–54 (2008).

6. J. M. Sonner, R. S. Cantor, Molecular Mechanisms of Drug Action: An Emerging View. *Annu. Rev. Biophys.* **42**, 143–167 (2013).

7. M. Weinrich, D. L. Worcester, The actions of volatile anesthetics: a new perspective. *Res. Pap. Acta Cryst.* **74**, 1169–1177 (2018).

8. R. A. Lerner, A hypothesis about the endogenous analogue of general anesthesia. *Proc Natl Acad Sci U S A*. **94**, 13375–13377 (1997).

9. D. Lingwood, K. Simons, Lipid rafts as a membrane-organizing principle. *Science*. **327**, 46–50 (2010).

10. E. N. Petersen, H.-W. Chung, A. Nayebosadri, S. B. Hansen, Kinetic disruption of lipid rafts is a mechanosensor for phospholipase D. *Nat. Commun.* **7**, 13873 (2016).

11. A. J. Patel *et al.*, Inhalational anesthetics activate two-pore-domain background K+ channels. *Nat. Neurosci.* **2**, 422–6 (1999).

12. M. Gruss *et al.*, Two-pore-domain K+ channels are a novel target for the anesthetic gases xenon, nitrous oxide, and cyclopropane. *Mol Pharmacol*. **65**, 443–452 (2004).

13. C. Heurteaux *et al.*, TREK-1, a K+ channel involved in neuroprotection and general anesthesia. *EMBO J.* **23**, 2684–2695 (2004).

14. E. Gray, J. Karslake, B. B. Machta, S. L. Veatch, Liquid general anesthetics lower critical temperatures in plasma membrane vesicles. *Biophys. J.* **105**, 2751–9 (2013).

15. D. Papahadjopoulos, K. Jacobson, G. Poste, G. Shepherd, Effects of local anesthetics on membrane properties. I. Changes in the fluidity of phospholipid bilayers. *Biochim. Biophys. Acta*. **394**, 504–19 (1975).

16. A. G. Lee, Model for action of local anaesthetics. *Nature*. **262**, 545–8 (1976).

17. S. a Jones, S.-H. Shim, J. He, X. Zhuang, Fast, three-dimensional super-resolution imaging of live cells. *Nat. Methods.* **8**, 499–508 (2011).

18. B. Huang, W. Wang, M. Bates, X. Zhuang, Three-dimensional super-resolution imaging by stochastic optical reconstruction microscopy. *Science*. **319**, 810–3 (2008).

19. E. Betzig *et al.*, Imaging intracellular fluorescent proteins at nanometer resolution. *Science*. **313**, 1642–1645 (2006).

20. S. T. Hess, T. P. K. K. Girirajan, M. D. Mason, Ultra-high resolution imaging by fluorescence photoactivation localization microscopy. *Biophys. J.* **91**, 4258–4272 (2006).

21. I. Leventhal, D. Lingwood, M. Grzybek, U. Coskun, K. Simons, Palmitoylation regulates raft affinity for the majority of integral raft proteins. *Proc. Natl. Acad. Sci.* **107**, 22050–22054 (2010).

22. M. J. Shipston, Ion channel regulation by protein palmitoylation. *J. Biol. Chem.* **286**, 8709–8716 (2011).

23. C. V. Robinson, T. Rohacs, S. B. Hansen, Tools for Understanding Nanoscale Lipid Regulation of Ion Channels. *Trends Biochem. Sci.* (2019), doi:10.1016/j.tibs.2019.04.001.

24. J. J. Kinnun, R. Bittman, S. R. Shaikh, S. R. Wassall, DHA Modifies the Size and Composition of Raftlike Domains: A Solid-State2H NMR Study. *Biophys. J.* **114**, 380–391 (2018).

25. Y. Comoglio *et al.*, Phospholipase D2 specifically regulates TREK potassium channels via direct interaction and local production of phosphatidic acid. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 13547–52 (2014).

26. M. McDermott, M. J. O. Wakelam, A. J. Morris, Phospholipase D. *Biochem. Cell Biol.* **82**, 225–53 (2004).

27. T. C. Sung *et al.*, Mutagenesis of phospholipase D defines a superfamily including a trans-Golgi viral protein required for poxvirus pathogenicity. *EMBO J.* **16**, 4519–30 (1997).

28. E. J. Bertaccini, R. Dickinson, J. R. Trudell, N. P. Franks, Molecular modeling of a tandem two pore domain potassium channel reveals a putative binding site for general anesthetics. *ACS Chem. Neurosci.* **5**, 1246–1252 (2014).

29. Z. Su, E. C. Brown, W. Wang, R. MacKinnon, Novel cell-free high-throughput screening method for pharmacological tools targeting K+ channels. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 1602815113- (2016).

30. C. Cabanos, M. Wang, X. Han, S. B. Hansen, A Soluble Fluorescent Binding Assay Reveals PIP2 Antagonism of TREK-1 Channels. *Cell Rep.* **20**, 1287–1294 (2017).

31. M. A. Pavel, H. Chung, E. N. Petersen, S. B. Hansen, Polymodal Mechanism for TWIK-Related K+ Channel Inhibition by Local Anesthetic. *Anesth. Analg.* **(in press)** (2019), doi:10.1213/ANE.0000000000004216.

32. S. G. Brohawn, E. B. Campbell, R. MacKinnon, Physical mechanism for gating and mechanosensitivity of the human TRAAK K+ channel. *Nature*. **516**, 126–130 (2014).

33. J. F. MacDonald *et al.*, Actions of ketamine, phencyclidine and MK-801 on NMDA receptor currents in cultured mouse hippocampal neurones. *J. Physiol.* **432**, 483–508 (1991).

34. S. B. Hansen, Lipid agonism: The PIP2 paradigm of ligand-gated ion channels. *Biochim. Biophys. Acta*. **1851**, 620–628 (2015).

35. R. S. Cantor, The lateral pressure profile in membranes: a physical mechanism of general anesthesia. *Biochemistry*. **36**, 2339–44 (1997).

36. H. C. Hemmings *et al.*, Emerging molecular mechanisms of general anesthetic action. *Trends Pharmacol. Sci.* **26**, 503–510 (2005).

37. E. I. Eger *et al.*, Minimum alveolar anesthetic concentration values for the enantiomers of isoflurane differ minimally. *Anesth. Analg.* **85**, 188–192 (1997).

38. R. Dickinson, I. White, W. R. Lieb, N. P. Franks, Stereoselective loss of righting reflex in rats by isoflurane. *Anesthesiology*. **93**, 837–843 (2000).

39. D. F. Covey *et al.*, Enantioselectivity of pregnanolone-induced gamma-aminobutyric acid(A) receptor modulation and anesthesia. *J. Pharmacol. Exp. Ther.* **293**, 1009–16 (2000).

40. D. Laverty *et al.*, Cryo-EM structure of the human α1β3γ2 GABAA receptor in a lipid bilayer. *Nature*. **565**, 516–520 (2019).

41. C. Aicart-Ramos, R. A. Valero, I. Rodriguez-Crespo, Protein palmitoylation and subcellular trafficking. *Biochim. Biophys. Acta - Biomembr.* **1808**, 2981–2994 (2011).

42. K. F. Herold, R. L. Sanford, W. Lee, O. S. Andersen, H. C. Hemmings, Clinical concentrations of chemically diverse general anesthetics minimally affect lipid bilayer properties. *Proc. Natl. Acad. Sci. U. S. A.*, 201611717 (2017).

43. R. D. Booker, A. K. Sum, Biophysical changes induced by xenon on phospholipid bilayers. *Biochim. Biophys. Acta - Biomembr.* **1828**, 1347–1356 (2013).

44. A. Nayebosadri, E. N. Petersen, C. Cabanos, S. B. Hansen, A Membrane Thickness Sensor in TREK-1 Channels Transduces Mechanical Force. *SSRN Electron. J.* (2018), doi:10.2139/ssrn.3155650.

45. M. J. Mlodzianoski *et al.*, Sample drift correction in 3D fluorescence photoactivation localization microscopy. *Opt. Express*. **19**, 15009–19 (2011).

46. M. A. Kiskowski, J. F. Hancock, A. K. Kenworthy, On the use of Ripley's K-function and

its derivatives to analyze domain size. *Biophys. J.* **97**, 1095–1103 (2009).

47.   P. Sengupta, T. Jovanovic-Talisman, J. Lippincott-Schwartz, Quantifying spatial

organization in point-localization superresolution images using pair correlation analysis.

Nat. Protoc. **8**, 345–54 (2013).

48.   J. M. Hartley *et al.*, Super-Resolution Imaging and Quantitative Analysis of Membrane

Protein/Lipid Raft Clustering Mediated by Cell-Surface Self-Assembly of Hybrid

Nanoconjugates. *Chembiochem*. **16**, 1725–9 (2015).

49.   P. Sengupta *et al.*, Probing protein heterogeneity in the plasma membrane using PALM

and pair correlation analysis. *Nat. Methods*. **8**, 969–75 (2011).

ACKNOWLEDGEMENTS

We thank Andrew S. Hansen for assisting with experimental design and discussion and
comments on the manuscript, Manasa Gudheti of Vutara for help with dSTORM data
processing, Michael Frohman for mPLD2 cDNA, and Guillaume Sandoz for chimeric TRAAK
cDNAs, and Stuart Forman for helpful discussion. This work was supported by a Director's New
Innovator Award (1DP2NS087943-01 to S.B.H.) from the NIH, a graduate fellowship from the
Joseph B. Scheller & Rita P. Scheller Charitable Foundation to E.N.P. We are grateful to the Iris
and Junming Le Foundation for funds to purchase a super-resolution microscope, making this
study possible.

AUTHOR INFORMATION

[1]Department of Molecular Medicine, [2]Department of Neuroscience, The Scripps Research
Institute, Jupiter, Florida 33458, USA.

[3]Department of Chemistry, The Scripps Research Institute, La Jolla, California 92037, USA

CONTRIBUTIONS

MAP, RAL, and SBH designed the experiments and wrote the manuscript. MAP performed all

electrophysiology, dSTORM, and PLD2 enzymes assays with help from ENP for dSTORM

imaging, and PLD2 assays.

COMPETING INTERESTS

The authors declare no competing interests.

MATERIALS & CORRESPONDENCE

shansen@scripps.edu

Figures and Figure Legends:

Figure 1 | Inhaled anesthetics disrupt GM1 domain structure. a, Representative reconstructed

super-resolution (dSTORM) images of GM1 domains (lipid rafts) before and after treatment with

chloroform (1 mM), isoflurane (1 mM), or MβCD (100 μM) (Scale bars: 1 μm). **b-d,** Bar graphs

comparing the average sizes (**b**) and areas (**c**) quantified by cluster analysis (± s.e.m., n = 2842-

7382). **d**, Quantified number of rafts per cell. (± s.e.m., n=10) (Student's t-test results:

****P<0.0001) **e**, Model representation of raft disruption by anesthetics. GM1 lipids (blue) form

ordered domains. Inhaled anesthetic (orange hexagon) intercalate and disrupt lipid order causing

the domain to expand.

Figure 2 | Activation of TREK-1 by inhaled anesthetic is PLD2 dependent. a,

Representative TREK-1 whole-cell currents activated by chloroform (1 mM) in physiological K^+

gradients. The current-voltage relationships (I-V curves) were elicited by 1-s depolarizing pulses

from -100 to 100 mV in +20 mV increments. **b**, Representative I-V curves showing that co-

expression of a catalytically inactive mutant of PLD2 (xPLD2 = PLD2_K758R) abolishes the

TREK-1 activation by chloroform. **c,** Bar graph showing the ~2-fold increase of TREK1 current

when activated by chloroform (1 mM) (n = 11) at +40 mV (± s.e.m.). **d,** Schematic

representation of TREK-1 activation by inhaled anesthetics. Anesthetic disruption of GM1

domains causes PLD2 to localize with TREK-1 and its substrate phosphatidylcholine (PC) in the

disordered region of the membrane. As PLD2 hydrolyzes PC to phosphatidic acid (PA), the

anionic lipid binds to a known gating helix (grey cylinder), with a lipid binding site (cyan)(*30*),

that activates TREK-1. Student's t-test results: *P < 0.05; **P<0.01; ***P<0.001; NS ≥ P.0.05.

Figure 3 | PLD2 localization renders TRAAK anesthetic sensitive. Native TRAAK is an anesthetic insensitive channel. **a,** Cartoon showing the experimental setup. TRAAK is fused with the C-terminus of TREK-1 (TRAAK/ctTREK). The BLD binding site is depicted in green. **b-c,** Representative I-V curve showing TRAAK/ctTREK-1 is activated by chloroform when co-expressed with mouse PLD2 (mPLD2) (**b**). The co-expression of the catalytically inactive PLD2 (xPLD2) abolishes the chloroform activation of TRAAK/ctTREK-1 chimeric channel (± s.e.m., n = 7) (**c**). **d,** Bar graph summarizing TRAAK/ctTREK-1 chimeric channel current in the presence or absence of xPLD2 and chloroform (1mM) at +40 mV (± s.e.m., n = 11) (Student's t-test results: ****P<0.0001, **P<0.01; NS ≥ P.0.05.) **e,** Model mechanism showing that anesthetics activate the TRAAKctTREK-1 chimeric channel through raft disruption and PLD2 substrate presentation; xPLD2 abolishes the activation (the color scheme is as in Fig. 2).

Figure 4 | Inhaled anesthetics displace PLD2 from GM1 domains. a, Representative super-

resolution (dSTORM) images of fluorescently labeled CTxB (lipid raft, from Fig. 1a) and PLD2

before treatment (Control) and after treatment with chloroform (1 mM), isoflurane (1mM), and

MβCD (100 μM) in N2A cells (scale bars: 1 μm). **b**, Average cross-correlation functions (C(r))

showing a decrease in PLD2 association with ordered GM1 domains after treatment with

anesthetic or MβCD. **c**, Comparison of the first data point in (**b**) (5 nm radius) (± s.e.m., n = 10-

17). **d**, Schematic representation of PLD2 in GM1 domain before (left) and after (right) anesthetic treatment. Palmitoylation drives PLD into GM1 domains (blue) away from its unsaturated PC substrate (yellow outline). Anesthetics (orange hexagon) disrupts GM-1 domains causing the enzyme to translocate where it finds its substrate PC in the disordered region of the cell.

Figure 5 | General anesthetics activates phospholipase D2 (PLD2) through raft disruption.

(**a-e**) Live cell assays showing the effect of anesthetics on PLD2 activity in N2A cells. Chloroform

(1 mM) (**a**), isoflurane (1 mM) (**b**), diethyl ether (1 mM) (**c**), propofol (50 µM) (**d**) and xenon (0.044

µM) (**e**) increased the PLD2 activity as compared with the control cells. Ketamine (50 µM) (**f**) and

the non-immobilizer F6 (**g**) had no effect on the PLD2 activity and the activity was inhibited by a

PLD2 specific inhibitor (2.5-5 µM) (mean ± s.e.m., n = 4). **h**, Summary of normalized anesthetic

induced activity of PLD2 in (a-g) at 60 min. (mean ± s.e.m., n = 4). **i-j**, Dose response of chloroform

(EC_{50} = ~1.0 mM) (**i**) and isoflurane (EC_{50} = ~0.8 mM) (**j**) on PLD2 activity through raft disruption.

k-l, Representative I-V curves showing the effects of propofol on TREK-1 in HEK293 cells using

1   whole cell patch clamp (**k**), and with xPLD2(**l**). **m,** Summary of TREK-1 currents showing an ~2-

2   fold increase when activated by propofol (25-50 µM) (n = 6) at +40 mV (± s.e.m.).

Figure 6 | Phospholipase D (PLD) regulates anesthetic sensitivity in *Drosophila*

***Melanogaster.* a**, Diagram depicting the setup of the anesthetic treatment and the positional

recordings of the flies. Fly positions were used to confirm anesthesia. **b**, Sedation curves showing

the percent of unconscious flies over time after chloroform treatment (2.8 mmol/L of air volume)

in both WT and PLD[null] flies. $p < 0.00001$. **c**, Confocal images showing robust labeling of GM1

domains (CtxB, green) in the membranes of labeled neurons (pan-neuronal Elav antibody, purple)

of the whole fly brain. **d-e**. Flies were treated with chloroform and the GM1 domains of whole brain

tissue were assayed for anesthetic induced disruption by super resolution imaging (dSTORM). **d**,

Super resolution images with (right) and without (left) sedating chloroform. **e,** Quantitation of raft

size diameter from fixed whole fly brain with and without chloroform. Similar to raft disruption in

tissue culture, the GM1 domains expand in brains of flies treated with chloroform. Grey dotted

lines indicate a hypothetical zoom compared to the low-resolution imaging of confocal in **c**.

SUPPLEMENTARY INFORMATION

Supplementary Figure 1 | GM1 domains and PLD2 activation by substrate presentation. a,

Chemical structures of select inhaled anesthetics are shown. Diversity ranges from xenon, a

single hydrophobic atom, to chloroform, chlorinated carbon, and ether compounds. **b**, the

organization of PLD2 and lipid rafts on the surface of a cell are shown (not to scale). GM-1

domains are comprised of GM1 saturated lipids and cholesterol, palmitoylation drives PLD2 into

GM1 domains (blue circle) where it is sequestered away from its substrate polyunsaturated PC

and PIP_2 domains (Red circle with yellow outline). **c**, Side view of the membrane in (b) with PLD2

localized to the inner leaflet through two palmitoylation sites (blue lines). GM1 domains also

commonly referred to as the liquid ordered phase (Lo) are thicker than the liquid disordered phase

(Ld). We previously showed mechanical force disrupts lipid rafts and causing PLD2 translocates

to the PIP_2 domains. PLD2 has a PIP_2 binding site and resides in equilibrium between GM1

domains and PIP_2 domains. PIP_2 is polyunsaturated and preferentially localized with unsaturated

PC—PLD2 generates unsaturated PA and PG products through substrate presentation(*10*).

Supplementary Figure 2 | Inhaled anesthetics disrupt GM1 domains. a, Particle size

distribution curve of the lipid rafts in N2A cells after chloroform, isoflurane, and MβCD treatments

(n=10). The number of rafts smaller than 150 nm decreased and the rafts larger than 150 nm

increased (see panel (d) for quantification). **b**, Derivatives of Ripley's K-Function (H(r))

demonstrating the separation of GM1 domains with or without treatment of the inhaled anesthetics

(1 mM) or MβCD (100 µM) (± s.e.m., n = 10). **c**, Bar graphs showing the amount of particle

localizations acquired within the fixed frames numbers (10000) (± s.e.m., n = 5). **d,** Histograms

showing the size distribution of lipid rafts from N2A cells in (a) binned from 0-150 nm and 150-

500 nm. Chloroform (1mM) and isoflurane (1mM) both shift from small to large and MβCD (100

µM) shifts from large to small (± s.e.m., n = 10). Both observations are a form of disruption. **e,**

Representative super-resolution (dSTORM) images of lipid rafts from C2C12 cells before and

after the treatment of the inhaled anesthetics or MβCD (100 µM) (Scale bars: 1 µm). **f**, Particle

1   size distribution curve of the lipid rafts after the chloroform, isoflurane, and MβCD treatments from

2   multiple C2C12 cells (n = 6-8). **g-h**, Bar graphs comparing the average lipid raft sizes (**g**) and

3   areas (**h**) quantified by cluster analysis showing the disruption of lipid raft by chloroform and

4   MβCD (± s.e.m., n = 2844-6525). Student's t-test results: ***P<0.001; ****P<0.0001.

Supplementary Figure 3 | General anesthetics activate TREK-1 through PLD2. a, Cartoon

depicting the amino acids that are known to play a role in anesthetic activation and the PLD2

dependent lipid-gating of TREK-1 channels. **b,** multiple-sequence alignment between the

swapped C-terminus of mouse TREK-1 (293-411) and TRAAK (255-398) generated with Clustal

Omega (EMBL-EBI). The colors and features correspond with panel (a). **c,** Current-voltage

relationship (I-V) for TREK-1 in wild type HEK 293T. No change in whole cell currents by 1 mM

of chloroform. **d**, Western blot showing the over expression of wtPLD2, mPLD2 and xPLD2 from

transfected HEK 293T cells. **e,** Bar graph summarizing wt. TRAAK channel currents when

activated by chloroform (1mM) at +40 mV (± s.e.m., n = 5) confirming earlier studies that

TRAAK is anesthetic insensitive. **f,** Ion flux assay of purified TREK-1 reconstituted into DOPC

(16:1) liposomes with 15 mol% DOPG anionic lipid. Anesthetics chloroform (1 mM), isoflurane

(1 mM), and propofol (50 µM) had no significant effect on channel currents compare to the

double cysteine gain of function mutation (Cys GOF) (± s.e.m., n = 3-5). **g,** Bar graph comparing

the relative activity from **f**.

Supplementary Figure 4 | Effect of search radius on observed raft size. a, Representative

images showing the raw images with particle localizations (left panel, scale bar = 5 μm),

clustered particles (DBSCAN) to determine raft sizes (middle panel, scale bar = 5 μm), and the

zoomed in (right panel, scale bar = 0.5 μm) using 100 nm search radius. Arbitrary colors are

assigned to particles of the same cluster. **b**, Bar graphs showing the changes in raft sizes after

anesthetic treatment at various search radius (ε) when at least 10 localizations were clustered.

c, Cartoon depicting the size variation at different search radius. A radius that is too small will

treat each particle as an individual cluster and give cluster that are too small. A search radius

that is too large clusters separate molecules giving an artificially large size. The search radius of

100 nm shown in (a) and used for this study shows appropriate clustering and a significant

increase in size.

Supplementary Figure 5 | Inhaled anesthetics displace PLD2 from GM1 domains in C2C12 cells. a, Representative super-resolution (dSTORM) images of fluorescently labeled lipid raft (CTxB) and PLD2 before and after the treatment with chloroform (1 mM), and MβCD (100 µM) in C2C12 cells (Scale bars: 0.5 µm). **b-c**, Average Cross-correlation functions (C(r)) (**b**) and the function normalized at C(r=5) (**c**) shows chloroform (1 mM) and MβCD (100 µM), disrupt PLD2 localization into lipid raft in C2C12 cells (± s.e.m., n = 5). Student's t-test results: *P < 0.05; ***P<0.001; ****P<0.0001; NS ≥ P.0.05.

Supplementary Figure 6 | Activation of PLD2 by general anesthetics in C2C12 cells. a-e,

Diethyl ether (1 mM) (**a**), chloroform (1 mM) (**b**), isoflurane (1 mM) (**c**), and higher concentration

of propofol (400 μM), but not 50 μM propofol (**d**) (green lines) activate PLD2 compared to control

cells. Ketamine (50 μM, blue line) had no observed effect on PLD2 activity. (**e**) PLD2 inhibitor

(2.5-5 μM) inhibited the activity (± s.e.m., n = 4). **f**, Normalized summary effect of anesthetics on

PLD2 activity at 60 min from the above experiments (± s.e.m., n = 4). Student's t-test results: *P

< 0.05; ***P<0.001; ****P<0.0001; NS ≥ P.0.05.

Methods

Sample preparation for Super Resolution Microscopy (d-STORM)

Super Resolution Microscopy was performed on N2A and C2C12 cells. Confluent cells were first differentiated overnight with serum-free DMEM in 8-well chamber slides (Nunc Lab-Tek Chamber Slide System, Thermo Scientific). Cells were then washed and treated with anesthetics or other drugs for 10 min. Chambers containing volatile anesthetic were tightly sealed with aluminum stickers. Cells were then chemically fixed with 3% paraformaldehyde and 0.1% glutaraldehyde in PBS for 10 min at room temperature with shaking, and the fixing solution was quenched by incubating with 0.1% $NaBH_4$ for 7 min followed by three times 10 min wash with PBS. Anesthetics or the drugs were also added into the fixing solution to ensure its effect on the cell. Fixed cells were then permeabilized with 0.2% Triton-X 100 in PBS for 15 min except the cells receiving the CTxB treatment. Cells were blocked using a standard blocking buffer (10% BSA, 0.05% Triton in PBS) for 90 min at room temperature. For labelling, anti-PLD2 antibody (Cell Signaling) with 1:500 dilution and CTxB (Life Technologies) with 1:1000 dilution in the blocking buffer were simultaneously added to cells and incubated for 60 min at room temperature. Cells were then extensively washed with 1% BSA, 0.05% Triton in PBS for five times 15 min each before labeling with the secondary antibody diluted into the blocking buffer and incubating for 30 min. Prior to labeling, the secondary antibody was conjugated to either Alexa 647 (to detect CTxB raft) or Cy3B (to detect PLD2). The incubation with secondary antibody was followed by above extensive wash and a single 5 min wash only with PBS. Labeled cells were then post-fixed with the previous fixing solution for 10 min without shaking followed by three times 5 min washes with PBS and two 3 min washes with deionized distilled water. To elucidate the lipid raft disruption by anesthetics or other drugs, compounds were applied to the reaction buffer at these concentrations: chloroform (1 mM) (Fisher Scientific); isoflurane (1 mM) (Sigma); mβCD

(100 µM) (Fisher); diethyl ether(1mM) (Sigma); ketamine (50 µM) (Cayman Chemicals); xenon

(Praxair).

d-STORM Image Acquisition and Analysis

Imaging was performed with A Zeiss Elyra PS1 microscope using TIRF mode equipped with an

oil-immersion 63X objective. Andor iXon 897 EMCCD camera was used along with the Zen 10D

software for image acquisition and processing. The TIRF mode in the dSTORM imaging provided

low background high-resolution images of the cell membrane. A total of 10,000 frames with an

exposure time of 18 ms were collected for each acquisition. Excitation of the Alexa Fluor 647

dye was achieved using 642 nm lasers and Cy3B was achieved using 561 nm lasers. Cells were

imaged in a photo-switching buffer suitable for dSTORM: 1% betamercaptoethanol, 0.4 mg

glucose oxidase and 23.8 µg Catalase (oxygen scavengers), 50 mM Tris, 10 mM NaCl, and

10% glucose at pH 8.0. Sample drift during the acquisition was corrected for by an

autocorrelative algorithm(*45*) or tracking several immobile, 100 nm gold fiducial markers or

TetraSpec beads using the Zen 10D software. The data were filtered to eliminate molecules

with localization precisions >50 nm.

Super-resolved Images were constructed using the default modules in the Zen Software. Each

detected event was fitted to a 2D Gaussian distribution to determine the center of each point

spread function (PSF) plus the localization precision. The Zen software also has many rendering

options including the options to remove the localization errors, outliers in brightness and size. The

super-resolved images have an arbitrary resolution of 128 pixel/μm. To determine the raft size

determination and the cross-correlations, the obtained localization coordinates were converted to

be compatible to Vutara SRX software (version 5.21.13) by an Excel macro. Cross-correlation

and raft size estimation were calculated through cluster analysis using the default analysis

package in the Vutara SRX software(*10, 46–48*). Cross-correlation function *c*(*r*) estimates the

spatial scales of co-clustering of two signals ─ the probability of localization of a probe to

distance *r* from another probe(*49*). Raft sizes are the size of clusters determined by measuring

the area of the clusters comprising of more than 10 observations.

In Vivo PLD2 activity measurements

A nonradioactive method was performed to measure in vivo PLD2 activity as described previously

(ref) (Fig. S2). Briefly, N2A or C2C12 cells were seeded into 96-well flat culture plates with

transparent-bottom to reach confluency (~ 5 x 104 per well). Then the confluent cells were

differentiated with serum-free DMEM for a day and washed with 200 µL of phosphate buffer saline

(PBS). The PLD assay reactions were promptly begun by adding 100 µL of working solution with

or without anesthetics. The working solution contained 50 µM Amplex red, 1 U per ml horseradish

peroxidase, 0.1 U per ml choline oxidase, and 30 µM dioctanoyl phosphatidylcholine (C8-PC).

Anesthetics were directly dissolved into the working buffer from freshly made stocks and

incubated overnight before assay reagents were added. In case of volatile anesthetics, 96-well

plates were tightly sealed with aluminum sticky films after adding the reaction buffer. The PLD

activity and the background (lacking cells) was determined in triplicate for each sample by

measuring fluorescence activity with a fluorescence microplate reader (Tecan Infinite 200 PRO,

reading from bottom) for 2 hours at 37°C with at excitation wavelength of 530 nm and an emission

wavelength of 585 nm. Subsequently, PLD activity was normalized by subtracting the background

and to the control activity. Data were then graphed (Mean ± SEM) and statistically analyzed

(student t-test) with GraphPad Prism software (v6.0f).

Electrophysiology

Whole cell patch clamp recordings of TREK1 currents were made from TREK-1-transfected HEK

293T cells as described previously (lab ref, Comoglio et. al. 2014). Briefly, HEK 293T were

cultured in growth media [DMEM, 10% heat-inactivated fetal bovine serum, 1% penicillin/

streptomycin] in a humidified incubator (95% air and 5% CO_2) at 37°C. When the HEK 293T cells

were ~90% confluent, they were seed at 50% confluency per 35-mm dish containing 15mm glass

coverslips coated with poly-D-lysine (1 mg/ml) to ensure good cell adhesion. The cells were then

transiently transfected using X-tremeGENE (Sigma) with a total of 1µg of DNA per dish. For co-

transfection of TREK1, TRAAK with PLD2 or PLD2-K758R cells were transfected with a ratio of

1:3. Human TREK1 pCEH and mouse PLD2 was kindly provided by Dr. Stephen B. Long, Sloan

Kettering Institute, NY. TRAAK/Ct-TREK1 (starting at Gly 293) pIRES2eGFP and TREK1/Ct-

TRAAK (starting at Gly 255) pIRES2eGFP was kindly provided by Dr. Sandoz Guillaume, iBV

CNRS, Université de Nice Sophia Antipolis, France. HEK 293T cells were obtained from ATCC

(Manassas, VA). Human TRAAK was a gift from Dr. Steve Brohawn, University of California,

Berkeley. Transfected cells were then visualized and selected for electrophysiology 24-48 hours

post transfection using green fluorescent protein. Standard whole-cell currents were recorded at

room temperature with Axopatch 200B amplifier and Digidata 1440A (Molecular Devices) and

measured with Clampex 10.3(Molecular Devices) at sample rate of 10 kHz and filtered at 2 kHz.

The recording micropipettes were made from the Borosilicate glass electrode pipettes (B150-86-

10, Sutter Instrument) by pulling with the Flaming/Brown micropipette puller (Model P-1000, Sutter

instrument). The micropipette resistances were ranged from 3-7 MΩ and filled with the internal

solution (in mM): 140 KCl, 3 $MgCl_2$, 5 EGTA, 10 HEPES, 10 TEA pH 7.4 (adjusted with KOH).

The external bath solution contained (in mM): 145 NaCl, 4 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 10 HEPES, 10

TEA pH 7.4 (adjusted with NaOH). After the voltage offset was adjusted to zero current between

the patch electrode and the bath solution, the whole cell configuration was achieved by repetitive

gentle suctions on cells sealed at 1-10 GΩ. In the whole cell configuration, cells were held at

−60 mV and currents were elicited by voltage steps command (at -100 to +100 mV from V_{hold} = -

60 mV) and voltage ramp commands (-100 mV to +50 mV in 5 ms). Volatile anesthetic,

chloroform, was applied using a gravity-driven (5 ml/min) gas-tight perfusion systems (Valves and

tubing were made of PTFE). HEK 293T cells were perfused with control solution or the test

solution that contained the volatile anesthetic. Chloroform was dissolved based on the anesthetic

saturation experiments that it has 66.6 mM solubility in water at 37°C(*11*). Subsequently, data

were replayed and analyzed using Clampfit 10 (Molecular Devices) to generate current-voltage

relationship (I-V Curve) from voltage steps protocol. Student's *t*-test was applied to assess

statistical significance using Prism6 (GraphPad software) and judged significant at $p < 0.001$. The

values represented in the graphs are Mean ± SEM.

Chanel Purification and Flux Assay

TREK-1 channel protein purification and Flux assay were done as previously described(*29*, *30*).

Briefly, Pichia yeast was used to express zebrafish TREK-1 (1-322 amino acids) containing GFP

at C-terminus. Followed by cryo milling, the extraction of the proteins was done in dodecyl-β-d-

maltoside (DDM) with protease inhibitors. The proteins were then purified on a cobalt affinity

column to homogeneity followed by size exclusion chromatography (SEC). The final SEC buffer

contained 20 mM Tris (pH 8.0), 150 mM KCl, 1 mM EDTA, and 2 mM DDM. All proteins were

collected with a predominant monodispersed peak corresponding to the expected molecular

weight (MW) of the assembled channel protein plus GFPs. This Purified TREK-1 was used to

generate Proteoliposomes by mixing 1:100 TREK-1/lipids. The ratio of the Lipids (85% DOPC

and 15% DOPG) were mixed, dried, and solubilized in rehydration buffer (150 mM KCl, 20 mM

HEPES [pH 7.4]) and calibrated with 3 mM DDM before the channel mixing. DDM was then

removed by BioBeads (Bio-Rad) and the proteoliposomes (5 μL) were sonicated and added to

195 μL of flux assay buffer (150 mM NaCl, 20 mM HEPES [pH 7.4], 2 μM 9-amino-6-chloro-2-

methoxyacridine [ACMA]) in a 96-well plate at room temperature. Flux was initiated by the addition

of the protonophore carbonyl cyanide m-chlorophenyl hydrazone (CCCP) (3.2 μM).

SDS PAGE and Western Blot

HEK 293T cells were transiently transfected using X-treme gene 9 (Sigma Aldrich). After 48 hours

of the transfection, cells were homogenized in the ice cold lysis buffer comprising 1.5% DDM

(Anatrace), 10% Glycerol (Fisher Scientific), 1 mM EDTA (Sigma), and 1: 100 protease inhibitors

(calbiochem) in 1X PBS, pH 7.5. Lysates were centrifuged at 17,000g for an hour at 4°C to collect

the supernatant. Protein sample were diluted 1:1 with 2X SDS Laemmli buffer (Biorad) and

incubated for 30 mins at 37°C. The denatured proteins were then separated on 4-20% SDS

polyacrylamide gel (Mini-PROTEAN® TGX™, Biorad) and blotted onto polyvinylidene difluoride

membranes, and probed with rabbit monoclonal antibody against PLD2 (E1Y9G, Cell Signaling).

Blot signals were visualized with a AlphaImager™ Gel Imaging System (Alpha Innotech).

Drosophila

General protocols

Flies were maintained in stocks at 25C. For all experiments, male flies were isolated from stocks

1-4 days old and allowed to recover in vials of no more than 10 flies for 24-48 hours before use

in protocols.

Anesthesia in Drosophila melanogaster

Anesthesia in flies were performed by volatiles and aerosols administered with positional

recording (VAAPR, Petersen unpublished data) in a custom-built narrow-width vertical chambers

placed in front of a camera to monitor the fly's positions with or without the chloroform treatment.

Wildtype and PLD^{null} flies were gently loaded into the designated chamber using mouth aspiration

and the hoses used for compound delivery were attached to the chamber. Flies were allowed to

habituate in the chamber while preparing treatment mixtures (~15 min). Mixtures without

chloroform contains only water with methylcyclohexanol (MCH) (1: 250), an aversive odor to

increase baseline activity of the flies. Chloroform were delivered at known concentrations by

calculating the partial pressures of the anesthetic in known dilutions of solvent. Both air control

and chloroform air were passed through flow meters to control the overall flow rate of 290

mL/min/chamber. Air was given to all flies for 2 min to record baseline activity/position after which

the chloroform was applied to the experimental flies. Data was analyzed by the Noah.py program

for position and activity(ref). Time of half maximal anesthesia (T50i) values were obtained using

a best-fit variable slope curve; sedation curves were analyzed using the Mantel-Cox test. Curve

fitting and statistical tests were determined using GraphPad Prism 6.

Brain Imaging

Flies were placed into control vial or vial containing a chloroform-soaked tissue. Flies were

removed and brains were isolated and placed into buffer containing PBS or PBS+chloroform on

ice. After dissections, brains were transferred to a buffer containing PBS with 3%

paraformaldehyde and 0.1% glutaraldehyde and allowed to fix while rocking overnight at 4C. The

following day the tissue was rinsed with PBS containing 0.5% TritonX-100 and 0.5% BSA. This

buffer was used for all additional steps unless otherwise stated. Tissue was rinsed 2X for 1hr after

which buffer with additional 30mg of BSA was added and allowed to rock at rt for 1.5 hours. Buffer

was removed and identical buffer containing primary antibodies (DSHB, Rat-Elav-7E8A10, 1:200)

was added to the tissue after which it was rocked for 3 hr at rt and then overnight at 4C. The

following day buffer was used to rinse the tissue as was done previously. Secondary antibody

(Jackson ImmunoResearch, 112-606-072, 1:500) was then applied in a similar manner as before

but rocking at 4C was extended to 5 days. On the 5[th] day, the tissue was rinsed once again 2X

and then in PBS alone. Tissue was then prepared for imaging. CTxB (1:500) was treated as a

secondary antibody. Super-res imaging was performed as described above. For whole brain

imaging confocal imaging was used after mounting the tissue onto a coverslip using standard

protocols.