Geographic variation in adult and embryonic desiccation tolerance in a terrestrial-breeding frog

Rudin-Bitterli, T.S.[1,2][*], Evans, J.P.[1,2] & Mitchell, N.J.[1]

[1] - School of Biological Sciences, The University of Western Australia, Crawley, Western Australia 6009, Australia

[2] - Centre for Evolutionary Biology, The University of Western Australia, Crawley, Western Australia 6009, Australia

Keywords: amphibians, desiccation tolerance, intra-specific variation, climate change, adaptation, phenotypic plasticity, environmental sensitivity

ABSTRACT

Intra-specific variation in the ability of individuals to tolerate environmental perturbations is often neglected when considering the impacts of climate change. Yet this information is potentially crucial for mitigating any deleterious effects of climate change on threatened species. Here we assessed patterns of intra-specific variation in desiccation tolerance in the frog *Pseudophryne guentheri*, a terrestrial-breeding species experiencing a drying climate. Adult frogs were collected from six populations across a rainfall gradient and their dehydration and rehydration rates were assessed. We also compared desiccation tolerance of embryos and hatchlings originating from within-population parental crosses from four of the six populations. Embryos were reared on soil at three soil-water potentials, ranging from wet to dry (ψ = -10, -100 & -400 kPa), and their desiccation tolerance was assessed across a range of traits including survival, time to hatch after inundation, wet mass at hatching, hatchling malformations and swimming performance. We found significant and strong patterns of intra-specific variation in almost all traits, both in adults and first generation offspring. Adult frogs exhibited clinal variation in their water balance responses, with populations from drier sites both dehydrating and rehydrating more slowly compared to frogs from more mesic sites. Similarly, desiccation tolerance of embryos and hatchlings was significantly greater in populations from xeric sites. Taken together, our findings suggest that populations within this species will respond differently to the regional reduction in rainfall predicted by climate change models. We emphasise the importance of considering geographic variation in phenotypic plasticity when predicting how species will respond to climate change.

INTRODUCTION

Understanding how organisms will respond to rapid environmental change is a major challenge for conservation and evolutionary biologists (Hoffmann and Sgro 2011). Most commonly, projections of climate change impacts on communities and species have been based on studies of the effects of environmental change on single populations (Moran et al. 2016). However, species are not uniform entities (Bolnick et al. 2003, 2011); individuals and populations vary genetically and phenotypically within species (Endler 1977), and intra-specific variation can be as great as trait variation across species (Albert et al. 2010; Des Roches et al. 2018). Therefore, models based on the assumption that all members of a species will respond similarly to ecological challenges may be invalid (see Kolbe et al. 2010; Kelly et al. 2012; Valladares et al. 2014; Llewelyn et al. 2016; Moran et al. 2016). In particular, information on the environmental sensitivity of range-edge populations will be vital for understanding future changes in species distributions, since it is at range edges where colonisations and extinctions primarily occur as the climate changes (Valladares et al. 2014; Rehm et al. 2015).

Clinal studies investigating patterns of phenotypic differentiation along environmental gradients can provide insight into how environmental stress has shaped the tolerance of populations (Keller et al. 2013). For example, in anuran amphibians, acid tolerance (Räsänen et al. 2003) and thermal tolerance (Hoppe 1978) vary between populations in a pattern consistent with the cline of each environmental stressor. By quantifying and comparing the sensitivity of populations along environmental clines, estimates of a species' capacity to adjust, either via phenotypic plasticity or local adaptation (or a combination of both), to altered environmental

conditions can be generated (Llewelyn et al. 2016; Pontes-da-Silva et al. 2018). For example, range-edge populations may hold a reservoir of pre-adapted genes that might, with sufficient gene flow, facilitate adaptation at the species level (Aitken and Whitlock 2013). Studies of variation in stress tolerance along environmental clines have demonstrated the capacity for populations to adjust their phenotypes to local conditions (Hoffmann and Harshman 1999; Hoffmann et al. 2002; Arthur et al. 2008; Gilchrist et al. 2008; Sinclair et al. 2012; Keller et al. 2013), and have shown that phenotypic divergence is often adaptive (Rajpurohit and Nedved 2013).

Water availability is a key environmental factor that is particularly important for amphibians. Amphibian skin is highly permeable to water (Young et al. 2005) and thus the ability of anurans to occupy terrestrial habitats is dependent on their ability to resist and avoid desiccation. In adult anurans, hydration state affects signalling behaviours and reproductive success (Mitchell 2001), locomotor performance (Gatten 1987; Hillman 1987), and predator avoidance and the ability to catch prey (Titon et al. 2010). Similarly, water availability has major effects on offspring fitness, particularly in terrestrial-breeding species, as the outer capsule of their eggs is almost completely permeable to water (Bradford and Seymour 1988*a*; Mitchell 2002; Andrewartha et al. 2008). Terrestrial embryos that develop on relatively dry soils are typically smaller (Mitchell 2002; Andrewartha et al. 2008; Eads et al. 2012), develop more slowly (Bradford and Seymour 1985), have reduced survival (Martin and Cooper 1972; Bradford and Seymour 1988*a*; Eads et al. 2012) and are more often malformed (Mitchell 2002; Eads et al. 2012). Thus, low environmental water availability is likely to induce strong directional selection through its negative effects on survival, reproduction and growth. This in turn can potentially lead to genetic or

plastic differences in desiccation tolerance among populations that occur across a range of hydric environments. Despite this, there are few studies of population-level variation in anuran desiccation tolerance across clines of water availability (but see Van Berkum et al. 1982), and no studies investigating whether such patterns exist at early developmental stages, where selection on desiccation tolerance should be especially strong given that embryos cannot move to escape unfavourable microclimates.

Here, we quantified intra-specific variation in traits that reflected adult and embryonic desiccation tolerance in populations of the terrestrial-breeding frog *Pseudophryne guentheri*. This species is highly suited to investigating geographic patterns in trait variation, as populations are distributed over a large area of southwestern Australia (Fig. 1) that spans a pronounced rainfall gradient (~300-1250 mm annual rainfall). Furthermore, *P. guentheri* inhabits a region that has experienced a substantial decline in winter rainfall over the past 40 years (19% reduction since the 1970s) (Smith 2004; IOCI 2012; Andrich and Imberger 2013; CSIRO and Bureau of Meteorology 2016). Rainfall in this region is predicted to decline further in the coming decades (Gallant et al. 2007; Bates et al. 2008; Smith and Power 2014; CSIRO and Bureau of Meteorology 2015) and there is concern that the range of this species will contract (Arnold 1988). We investigated whether adult *P. guentheri* show clinal variation in their water balance and whether the desiccation tolerance of their offspring differs in line with the rainfall gradient. We focused on quantifying variation in a range of traits putatively tied to fitness, including survival, time to hatch after inundation, wet mass at hatching, hatchling malformations and swimming performance.

MATERIALS AND METHODS

Ethics statement

All animal experiments were conducted in accordance with the University of Western Australia's (UWA) Animal Ethics Committee (permit number RA/3/100/1466). Fieldwork was conducted under permit SF010807 issued by the Western Australian Department of Biodiversity, Conservation and Attractions.

Study species

Pseudophryne guentheri (Anura: Myobatrachidae) is a small (26-33 mm snout-to-vent length) terrestrial-breeding frog endemic to the southwest of Western Australia (Tyler and Doughty 2009). Its range abuts that of an inland, arid-adapted member of this widespread genus (*P. occidentalis*). Breeding takes place in autumn and early winter following seasonal rainfall. Males excavate burrows in areas that are likely to be flooded and call from the burrow entrance to attract females (Anstis 2013). After a female has selected and approached a male, mating occurs inside burrows. Females deposit clutches of 60 to 300 eggs directly onto the soil and encapsulated embryos develop terrestrially (Anstis 2013). Hatching occurs when burrows flood in late winter (Eads et al. 2012) and tadpoles complete their development in ephemeral water bodies in about three months (Anstis 2013).

Animal collection and study sites

Adult *P. guentheri* were collected by hand and in pit-fall traps from six study sites located in the centre and northern limit of the species' range (Fig 1) in May and June 2016. Sites were distributed across a rainfall gradient, with the wettest site (population 1) receiving ~790 mm of rain per year and the driest site (population 6) receiving ~330 mm of rain per year (Table 1). In total, 10 to 19 (mean = 15.8, Table

1) calling males were collected from each site. Gravid females were more difficult to

locate and hence our collection of females was restricted to four sites (Table 1).

Adult frogs were temporally housed in small (4.4 L) plastic containers containing

moist sphagnum moss and transported to the University of Western Australia in

Perth within five days of collection. Frogs were then housed in a controlled-

temperature room at 16°C with an 11/13 h light/dark photoperiod to mimic winter

conditions, and were fed a diet of pinhead crickets.

Dehydration and rehydration assays in adult males

Rates of dehydration and rehydration were assessed in 90 adult males collected

across all six populations, with a minimum sample size of 10 males per population

(mean = 15). Frogs were allowed to acclimate to 16°C for two to five days, during

which time they were fed *ad libitum*. Body mass during this period remained stable (±

2.5% body mass). One day prior to commencement of the experiments, all food was

removed to minimise weight changes due to defecation during water balance

experiments, and sphagnum moss was moistened to ensure that frogs were fully

hydrated.

Dehydration rate

After the urinary bladder was emptied by cannulation, frogs were weighed (= 100%

standard mass) and then placed individually in desiccation cages. Desiccation cages

consisted of a PVC ring (diameter = 5.5 cm, height = 2 cm) wrapped in Nylon mesh

that allowed frogs to be weighed without handling. Desiccation cages holding frogs

were weighed (± 0.01g, GX-600 high precision balance, A&D Weighing, NSW,

Australia), and placed into a desiccating chamber containing Drierite (W. A.

Hammond, Xenia, OH, USA). Cages were weighed every 30 min until frogs reached

85% of their standard mass. This amount of water loss via evaporation and

respiration was reversible and did not cause adverse effects, and frogs moved very

little during trials.

Rehydration rate

Immediately after the desiccation trials, frogs were removed from the cages and

placed in lidded petri dishes containing deionised water to a level of 1 cm. This

ensured that a frog's ventral patch was exposed to the water at all times and

minimised their movements. Rehydration occurred rapidly and frogs were blotted dry

and weighed every 10 min until regaining 100% of their standard mass.

As rates of water loss and water uptake are dependent on body size (Withers et al.

1982; Wygoda M. 1984; Titon and Gomes 2015), dehydration and rehydration rates

are expressed as area specific measurements, calculated for each frog by the

equation

$$\text{Dehydration rate} = \frac{W_1 - W_2}{[SA \times (T_2 - T_1)]}$$

where W_1, and W_2 indicate initial and final body mass, and T_1 and T_2 represent start

and end time of each trial (Liu and Hou 2012). The same equation was used to

calculate area-specific rehydration rates. The surface area of the frogs was

estimated as SA (cm^2) = 9.9 (standard weight)$^{0.56}$ (McClanahan and Baldwin 1969).

In-vitro fertilisation methods

In-vitro fertilisation was used to perform controlled within-population crosses in the laboratory. The eggs of each female were divided equally into five groups and fertilised separately with sperm from one of five males. This allowed us to control for potential parental compatibility effects on offspring fitness, which have been identified in this species (Eads et al. 2012) and in other anurans (Dziminski et al. 2008). Due to the restricted number of sires, we used the same male's sperm to fertilise several females in some populations (see Statistical Analysis). Males were euthanized via ventral immersion in <0.03% benzocaine solution for 10 min, followed by double pithing. Both testes were removed, blotted dry and weighed to the nearest 0.1 mg (precision balance XS204, Mettler Toledo) and placed on ice. Testes were then macerated in 20 to 615 µL (adjusted according to the weight of the testes) of chilled 1:1 standard amphibian ringer (SAR; 113mM NaCl, 2mM KCl, 1.35 mM $CaCl_2$, and 1.2 mM $NaHCO_3$). This buffer has a similarly osmolality to the male's reproductive tract and keeps spermatozoa in an inactive state (Byrne et al. 2015), allowing sperm storage for extended periods of time (days - weeks) without considerable declines in motility (Browne et al. 2001; Kouba et al. 2003). Sperm concentrations in testes macerates were measured using an improved Neubauer haemocytometer (Hirschmann Laborgeräte, Eberstadt, Germany), and sperm suspensions were diluted with 1:1 SAR to produce stock solution of 100 sperm per µl.

Ovulation in females was induced via two subcutaneous injections of the hormone LHRHa over the course of two days (Silla 2011). On day one, a priming dose of 20% of the ovulatory dose was administered, followed by an ovulatory dose (2 µg LHRHa

per 1g female standard weight, diluted in 100 µl of SAR) 22 hours after the first injection. Approximately ten hours after administration of the ovulatory dose, eggs were gently squeezed from each female onto a clean surface. They were then moistened with SAR and divided equally among five small petri dishes and placed on ice until fertilisation. Following Eads et al. (2012), a small volume of sperm suspension was pipetted onto one edge of the petri dish, followed by a larger volume of diluted SAR solution (one part SAR to four parts de-ionised water) which activated the sperm. To promote fertilisation, each dish was manually agitated for 20 sec to mix the solutions and eggs. After 15 minutes, eggs were backlit and photographed (while submerged in water to minimize refraction) using a digital imaging camera (Leica DFC320) attached to a light microscope (Leica MZ7.5) at X 6.3 magnification. These images were used to measure the ovum diameter of a random sample of 50 eggs from each female, using ImageJ software (Abràmoff et al. 2004). One hour after combining eggs and sperm, fertilisation success was scored by counting eggs that had rotated (Gosner Stage 1; Gosner 1960).

Soil preparation and embryo incubation

Fertilised eggs were assigned to one of three water-potential (ψ) rearing environments: a wet treatment (ψ = -10 kPa), an intermediate treatment (ψ = -100 kPa) and a dry treatment (ψ = -400 kPa). These water potentials represented a range of hydric conditions in which egg clutches develop in the field, which depends largely on the time since rainfall (N. J. Mitchell, unpubl. data).

Soil water potentials were established by oven-drying homogenised soil, previously collected from a *P. guentheri* breeding site, at 80 °C for 24 hours. Soil was

distributed into small, lidded containers (dimensions: 7.5 x 11 x 4 cm, 50 g of soil per container), and re-wetted with an appropriate mass of deionised water using a soil water-retention curve previously determined for the same soil type (N. J. Mitchell, unpubl. data), and allowed to equilibrate. The water content of the soil (g/g of dry soil) was approximately 50% in the wet treatment, 33% in the intermediate treatment, and 21% in the dry treatment.

Eggs were distributed onto soils 7 - 9 hours after fertilisation, with three replicates per water potential treatment and family (sire-by-dam combination). Small plastic rings (nylon plumbing olives, 12 mm in diameter) were placed to surround egg clusters and were labelled to identify individual crosses. The containers housing the eggs and soils were then placed in incubators (model i-500, Steridium, Australia) set at 16 ± 0.5 °C and weighed every two days to ensure that soil water potentials remained stable throughout incubation. Embryos were monitored every two days and any mouldy eggs were removed. Embryonic survival was recorded for each family as the percentage of fertilised eggs that hatched (see below).

Desiccation tolerance of embryos and hatchlings

Time to hatching

At 33 days after fertilisation (when embryos were approximately at Gosner Stage 26), hatching was induced by placing embryos individually in small tubes containing 2 ml of deionised water (Bradford and Seymour 1988*b*; Eads et al. 2012). Embryos were monitored every 30 min, and once a hatchling escaped the egg capsule, the time to hatching was recorded.

 Swimming performance

Swimming performance was recorded 6 - 12 hours after hatching on a subset of

hatchlings (N = 556 across all populations, with a minimum sample size of 30 per

population and treatment). For this purpose, individual hatchlings were placed in a

petri dish (diameter = 15 cm) containing water to a level of 1 cm. After an initial

acclimation period of 1 min, a glass cannula was used to gently poke the tail of each

hatchling, which elicited a burst swimming response. A video camera (Canon

PowerShot G16, recording at 60 fps) installed 30 cm above the petri dish was used

to film three burst swimming responses for each hatchling, and their movement was

later tracked and analysed using EthoVision v8.5 software (Noldus et al. 2001).

EthoVision enabled the quantification of the following swimming parameters:

maximum velocity (cm s^{-1}), mean velocity (cm s^{-1}) and total distance moved (cm).

We also recorded mean meander (deg cm^{-1}), a measure of the straightness of the

swimming response, as dry rearing environments can lead to asymmetrically shaped

hatchlings (Eads et al. 2012) that swim in a more circular motion (N. J. Mitchell, pers.

observations). A hatchling was considered moving when it exceeded 0.45 cm s^{-1}.

Since each video recording contained three burst swimming responses with periods

of no movement in between them, EthoVision only analysed frames in which a

hatchling moved faster than 0.45 cm s^{-1} (consequently merging the three swimming

responses for each hatchling). Immediately following the swimming performance

trials, hatchlings were euthanized in <0.03% benzocaine and preserved in 10%

neutral buffered formalin.

 Hatchling wet weight and malformations

 Wet masses of preserved hatchlings were recorded to the nearest 0.001 g after

 blotting on tissue. Hatchlings were then photographed in lateral view while

 submerged in water using a digital imaging camera (Leica DFC320) attached to a

 light microscope (Leica MZ7.5) at X 6.3 magnification. These images were used to

 score any malformations for each hatchling. Malformations consisted of three forms

 of notochord malformations (scoliosis – lateral curvature of the spine; lordosis –

 concave curvature of the spine; and kyphosis – convex curvature of the spine) and

 edemas (abnormal accumulation of fluids in tissues) (Fig. 2). Collectively, these

 deformities represented 84% of the total number of malformations observed.

 Malformations were quantified for each combination of treatment and sire-by-dam

 family as the percentage of hatchlings that showed a malformation of any kind.

 Statistical analysis

 All analyses were performed in R version 3.4.3 (R Development Core Team 2017).

 Adult de- and rehydration rates

 Rehydration rate was $\log_{10}$-transformed and the rate of dehydration was transformed

 using the Box-Cox method (Box and Cox 1964) to fulfil assumptions of normality. We

 then used one-way ANOVAs, followed by Tukey HSD post-hoc tests, to test whether

 dehydration and rehydration rates significantly differed between populations. Linear

 regression analyses were used to test for relationships between dehydration or

 rehydration rates and mean annual rainfall.

Desiccation tolerance of embryos and hatchlings

For four populations where we created within-population crosses we used linear mixed-effects models (with restricted maximum-likelihood methods; REML) to investigate variation in offspring fitness traits resulting from the water potential treatment. To ensure that data complied with assumptions of normality, Q-Q plots of residuals were inspected and, where necessary, data were transformed using the Box-Cox method (Box and Cox 1964). These transformations were performed for hatchling wet weight, time to hatching, total distance moved, and mean meander. Linear mixed-effect models were run using the lme4 package in R (Bates et al. 2015), with treatment, population and the population-by-treatment interaction treated as fixed effects and dam fitted as a random effect. We also added sires as random effects to all models to control for the repeated use of sperm donors across IVF trials. Ovum size was added as a covariate in all analyses to control for some trait variation due to maternal effects (Eads et al. 2012). The significance levels of fixed effects were evaluated using Wald chi-squared tests on the full model.

Fertilization rates, survival and malformation data were binomial variables and thus a generalized linear mixed-effects model (GLMM) with a logit-link function was used for the analysis of these traits. Fixed effects were treatment, population and the population-by-treatment interaction, and sire and dam were added as random effects. The significance of the fixed effects of treatment and population and their interaction, were evaluated using Wald Z tests, and 95% Wald confidence intervals were obtained using the "confint" function from the MASS package in R (Venables and Ripley 2002).

RESULTS

Dehydration and rehydration rates

Across all six study populations, males lost 15% of their standard mass on average after 4.98 ± 1.39 SD h (range: 2.03 - 8.20 h) in the desiccation chamber, and on average rehydrated from this benchmark to full hydration in less than an hour (0.82 ± 0.39 SD h, range: 16 - 120 min). Time to dehydrate and rehydrate correlated strongly with body size ($P < 0.001$), and therefore only area-specific dehydration and rehydration rates are used to compare populations (see Materials and Methods). Rates of dehydration differed significantly among populations ($F_{5,84}=11.58$, $P < 0.001$), with the highest in males from the wettest site (population 1) and a trend towards decreasing rates with increasing aridity (Fig. 3A). Rehydration rates followed a similar pattern (Fig 3B), but the population effect was not significant ($F_{5,84} = 0.964$, $P = 0.445$). We found significant linear relationships between dehydration and rehydration rates and mean annual rainfall (dehydration rate: regression coefficient = 99.75 ± 13.66 SE, t = 2.27, $P = < 0.001$, rehydration rate: regression coefficient = 3.42 ± 1.51 SE, t = 2.27, $P = 0.026$) (Fig. 3A & B), with frogs originating from more mesic sites dehydrating and rehydrating faster than frogs from xeric populations.

Desiccation tolerance of embryos and hatchlings

Survival

Embryonic survival on wet soils (high water potential) was high (> 93%) in all four populations (Fig. 4A). Lower soil water potentials negatively affected survival rates, although the severity of the treatment effect varied among populations, as evident from significant population-by-treatment effects (Table 2). The population from the wettest site (population 1) showed the largest reduction in survival in response to dry

and intermediate soil moisture, and this difference tended to decrease in populations

originating from sites of increasing aridity (Fig. 4A). As such, low soil water potentials

had very little effect on embryonic survival in the driest population (population 5),

with over 88% of embryos surviving to hatching stage.

Time to hatching

Across all populations, the average time that the terrestrial embryos required to

hatch after submergence in water significantly increased with decreasing soil water

potentials (Table 2, Fig. 4B). That is, embryos reared on relatively wet soils hatched

more quickly than their siblings reared on drier soils. However, there were marked

differences in hatching times between populations, and the way treatment affected

time to hatching also differed significantly among populations (Table 2, Fig. 4B). As

such, hatchlings originating from the driest population (population 5) hatched

significantly more quickly than hatchlings from the other three populations,

irrespective of the rearing environment (Fig. 4B). Hatchlings reared on wet soils from

populations 1, 2 and 3 hatched after approximately 17 hours, but required

approximately 30 hours when reared at intermediate water potentials. Furthermore,

hatchlings originating from the wettest site (population 1) took > 40 h to hatch, but

this increase in hatching time was less substantial in population 2, and populations 3

and 5 exhibited similar hatching times in dry and intermediate rearing environments

(Fig. 4B).

Wet mass at hatching

Low soil water potentials significantly reduced the wet mass of hatchlings. Similar to

the pattern observed in the previous two traits, the population-by-treatment

interaction was significant, suggesting that the severity of the treatment effect varied among populations (Table 2). Hatchlings from the wettest site (population 1) showed the largest reduction in hatchling size in response to low soil water potentials, and this decline in size decreased in populations originating from sites with increasing aridity (Fig 4C). Accordingly, hatchling wet mass from the driest site (population 5) hardly varied between dry, intermediate or wet rearing environments (Fig. 4C). As expected, ovum size significantly affected the wet mass of hatchlings (Table 2).

Malformations

The total percentage of malformed hatchlings was low (~ 5%) in the wet treatment in all four populations (Fig 4D) but increased with decreasing soil water potential. Malformation rates did not differ significantly between populations, and there were no significant population-by-treatment interactions for this trait. Soil water potential treatments did not significantly affect the relative occurrence of notchord or edema type malformations, but there were significant population effects for both these traits (Table 2, Fig. 4D).

Swimming performance

Low soil water potential significantly reduced the swimming performance of hatchlings (Table 3), with maximum and mean swimming velocity decreasing (Fig. 5A & B) and the total distance moved declining in dry and intermediate rearing environments (Fig. 5D, Table 3). Furthermore, mean meander increased in response to low soil water potentials (Fig. 5C), indicating that hatchlings swam more linearly when reared in wetter conditions. The effect of the soil water potential treatment on swimming performance differed significantly among populations (Table 3), with the

wettest population (population 1) showing the largest declines in velocity and largest

increases in mean meander in response to the dry and intermediate treatment. In

contrast, swimming performance in hatchlings from the driest population (population

5) was hardly affected by the dry treatment (Fig. 5)

DISCUSSION

Our findings demonstrate significant intra-specific variation in traits related to

desiccation tolerance in *Pseudophryne guentheri* (adult males and first generation

offspring), consistent with patterns of genetic adaptation and/or phenotypic plasticity

in response to local water availability. One interpretation of this finding is that our

results may, at least in part, reflect cryptic speciation, which is becoming increasingly

apparent in this (and related) genera of Australian ground frogs (e.g. Donnellan et al.

2012; Catullo et al. 2014). Recent genetic analysis of ten *P. guentheri* populations

(including the six studied here), which was based on more than 12,000 single-

nucleotide polymorphism (SNP) markers, showed strong structure (overall F_{ST} 0.186)

but clear evidence for populations clustering within a well-supported species

(Cummins 2017). Hence the trait divergence reported here most likely reflects local

adaptation to climatic (and potentially edaphic) variables, and emphasizes why

considering population variation in fitness-related traits will often be necessary for

understanding how a species might respond to climate change.

Dehydration and rehydration rates in adult males

Our observation that populations significantly differed in their water balance

(dehydration and rehydration rates) was in accordance with annual variation in

rainfall in these populations (Fig. 3). Dehydration negatively affects survival and

fitness in anurans through its effects on locomotor performance (Claussen 1974;

Gatten 1987; Moore and Gatten 1989; Köhler et al. 2011), which can impede

predator escape, foraging behaviour, territorial defence, signalling and mating

(Mitchell 2001; Titon and Gomes 2015). Consequently, lower rates of dehydration

are likely to benefit frogs occupying dry habitats, as they allow individuals to leave

moist retreats for longer to perform ecologically important behaviours (Feder and

Londos 1984; Winters and Gifford 2013). While our experiments were not designed

to determine whether variation in dehydration rates was a consequence of genetic

adaptation or phenotypic plasticity, our results reflect interspecific patterns for

amphibians reported elsewhere. For example, in anurans, resistance to evaporative

water loss (EWL) correlates with environmental water availability (Wygoda M. 1984;

De Andrade and Abe 1997), with terrestrial specialists exhibiting lower rates of EWL

than species occupying primarily aquatic habitats (Young et al. 2005). Similarly,

Winters and Gifford (2013) found variation in EWL rates at the population level, with

populations of lungless salamander *(Plethodon montanus)* from dry, low-elevation

areas dehydrating more slowly relative to those from wetter, higher-elevation areas.

Together these results are consistent with patterns of directional selection for lower

dehydration rates in areas where water is frequently scarce.

One finding less consistent with the current understanding of amphibian water

balance was our observation that mean annual rainfall and rates of rehydration had a

significant positive relationship (Fig. 3D). That is, males from xeric sites rehydrated

more slowly than males from mesic sites. This finding is in contrast to interspecific

patterns, where rehydration rates are generally highest in species from arid

environments (Ewer 1952; Bentley et al. 1958; Dumas 1966; Van Berkum et al.

1982; Titon and Gomes 2015). Similarly, the single study we found that investigated

rehydration rates of an anuran species at the population level demonstrated higher

rates of water uptake in populations from drier habitats (Van Berkum et al. 1982).

The ability to absorb water rapidly can extend the time that a frog can be active by

decreasing the amount of time spent in a refuge for rehydration (Van Berkum et al.

1982), suggesting that fast and efficient hydration is likely beneficial for frogs,

particularly those inhabiting dry areas.

One explanation for the slower rehydration rates observed in the more xeric

populations could be that the mechanisms that reduce evaporative water loss also

reduce water uptake. The thickness of the epidermis (stratum corneum), for

example, affects both dehydration and rehydration rates in anurans (Toledo and

Jared 1993). There is also evidence that skin thickness is associated with habitat

aridity at the species level (Toledo and Jared 1993), with species from drier areas

producing a thicker epidermis which reduces dehydration rates whilst also impeding

water uptake. Further, aquaporins (water channel proteins) embedded within the

pelvic patch (Kubota et al. 2006) can influence rehydration rates in anurans, and

variation in the density and type of aquaporins among species has been interpreted

as an adaptation of frogs to their respective hydric environment (Suzuki et al. 2007;

Ogushi et al. 2010). It is currently unknown whether similar patterns of morphological

or physiological differences in the skin properties exist at the population level, and

we encourage future studies to explore the underlying mechanisms that drive intra-

specific differences in dehydration and rehydration rates across hydric gradients.

An alternative explanation for our rehydration rate data is that our method of

exposing the ventral patch of dehydrated frogs to water may not have adequately

reflected natural behaviours. Terrestrial frogs rarely rehydrate in free standing water,

and instead water uptake is most often achieved via exposure of the ventral patch to

moist soil (Wells 2007). Whilst our design minimised potential behavioural co-factors

that could influence rehydration rates (e.g. frogs rehydrating on soil could employ a

range of tactics, including burying and/or adopting different postures that aid water

uptake), future work comparing rates of water uptake in water and moist soil would

be beneficial for an evaluation of the generality of the rehydration patterns obtained

here.

Desiccation tolerance of embryos and hatchlings

We found substantial intra-specific variation in traits related to desiccation tolerance

in *P. guentheri* embryos and hatchlings. Low soil water potentials generally reduced

the expression of traits putatively linked to fitness, although the severity of this effect

varied greatly among populations, with populations from the wettest site (pop. 1)

being the most negatively affected by dry rearing environments. Embryonic survival,

for example, was reduced by ~35% in the dry treatment in the wettest population, but

only by ~5% in the population originating from the driest site (Fig. 4A). Whilst

negative effects of desiccation on the survival of terrestrial embryos are well

established (Martin and Cooper 1972; Bradford and Seymour 1988*a*; Mitchell 2002;

Eads et al. 2012), the present study is, to our knowledge, the first to report such

marked differences among populations. Nevertheless, we are only able to speculate

about the mechanisms driving such intra-specific differences in embryonic survival

across the dry rearing environments. One possible explanation is that the structural

composition of the egg jelly coat may differ among populations. For example,

females from drier areas may produce eggs with thicker capsules, which would limit

water loss during embryonic development (Martin and Cooper 1972). Alternatively,

there may be physiological processes that enhance desiccation tolerance of

embryos developing on drier soils. For example, embryos may slow their metabolism

and development rates and extend the period of subsistence on energy reserves to

reduce their water requirements (Bradford and Seymour 1985; Podrabsky et al.

2010). We eagerly anticipate future studies designed to uncover the mechanisms

underlying variation in desiccation resistance among populations.

We found marked intra-specific differences in the time an embryo was able to hatch

after submerging it in water. In particular, embryos from the driest population (pop. 5)

hatched quickly, irrespective of the water potential of the soil on which they were

reared, whereas time to hatching was prolonged in populations originating from

areas with increasing annual precipitation (Fig 4B). Faster hatching times are likely

beneficial, as they can allow hatchlings to be washed into stable pools of standing

water, where they can feed and seek refuge (Warkentin 2011*a*). This response may

be particularly vital in areas where annual precipitation is low and where ephemeral

water bodies also dissipate more quickly due to higher ambient temperatures. At the

same time, immediate hatching after submergence is not always advantageous. For

example, embryos may delay hatching if they are too premature to be fully

competent as free-swimming tadpoles (Warkentin 2011*a*).

The strong intra-specific variation in hatching time raises questions about the

mechanisms driving these differences. Hatching in amphibians is highly plastic

(Warkentin 2011*a*) and can be achieved through chemical or mechanical processes,

or a combination of both (Duellmann and Trueb 1986). The chemical hatching

mechanism is more common (Warkentin 2011*a*), and embryos of many species have

hatching glands which release proteolytic enzymes in response to hatching cues

(such as predators or low PO_2) that digest components of the egg (Carroll and

Hedrick 1974; Nokhbatolfoghahai and Downie 2007). It is possible that embryos

produced from drier sites had denser hatching glands, synthesised a larger amount

of enzyme or released enzymes from vesicles more rapidly (Nokhbatolfoghahai and

Downie 2007; Kawaguchi et al. 2009; Cohen et al. 2016). Alternatively, intra-specific

differences in hatching time could potentially arise due to variation in development

rate. Development rate is negatively affected by low soil water potentials (Bradford

and Seymour 1985; Packard 1991) and it is possible that embryos from mesic

populations were too premature to hatch after 33 days when reared on dry soils. For

example, hatching glands may not have developed, or embryos may have lacked the

sensory capacities to perceive the hatching cue (Warkentin 2011*b*). However,

hatching glands in Myobatrachids develop relatively early (from Gosner stage 17)

and are fully present in embryos by Gosner stage 19 (Anstis 2010), making it unlikely

that any slight differences in the maturity of embryos in our study affected their ability

to hatch.

A further potential driver of intra-specific differences in hatching time may be the

structural composition of the egg capsule. For example, egg capsules from

populations occupying drier habitats may swell more rapidly when they rehydrate

when flooded, resulting in mechanical rupture of the perivitelline membrane, or an

outer egg cuspule layer, and so releasing the embryo (Duellmann and Trueb 1986).

While understanding the mechanisms driving differences in hatching time was not

the focus of our study, in light of our findings we suggest that *P. guentheri* offers an

ideal experimental system to explore this further.

Low soil water potentials reduced the wet mass of hatchlings and increased the

occurrence of hatchling malformations, in line with earlier studies (Taigen et al. 1984;

Mitchell 2002; Andrewartha et al. 2008; Eads et al. 2012). Embryos were preserved

6 - 12 hours after hatching, and therefore differences in wet weight are unlikely to be

the result of variation in body water content as hatchlings had ample time to hydrate.

Retarded growth on drier soils may be a consequence of reduced metabolic rates

(Bradford and Seymour 1985; Mitchell 2002) and yolk consumption (Packard 1999).

However, wet mass at hatching was only reduced slightly in hatchlings from the

driest site when reared on dry soils, whereas wet mass decreased significantly in

hatchlings originating from more mesic sites (Fig. 4C). Malformations are associated

with insufficient swelling of the egg capsule on dry substrates, which leads to

embryos being unable to rotate freely within the perivitelline space and sometimes

adhering to the perivitelline membrane (Bradford and Seymour 1988*a*; Mitchell 2002;

Andrewartha et al. 2008). Tadpole malformations can persist after metamorphosis

(Plowman et al. 1994) and are therefore likely to have long-term consequences for

survival and fitness.

The effect that water potential treatment had on body size and shape explained

differences in swimming performance among hatchings from different populations,

with hatchlings swimming more slowly, less straight and moving a smaller distance if

they were reared on drier soils (Fig 5). Decreased swimming performance is likely to

have negative fitness consequences, as rapid swimming allows tadpoles to escape predators and generally aids foraging (Webb 1986; Watkins 1996; Wilson and Franklin 1999; Teplitsky et al. 2005; Walker et al. 2005; Langerhans 2009). As observed for other traits studied here, there was pronounced intra-specific variation in the effects of dry soils on hatchling swimming performance. Swimming performance was unaffected by the dry or intermediate soil water potential treatments in the driest population (pop. 5), but decreased steeply in more mesic populations in response to incubation on dry soils. Interestingly, hatchlings from the driest population swam comparatively poorly, even though other effects of the dry rearing environments were minimal. One explanation could be that egg capsules from mesic populations are more permeable to water, and consequently swell relatively more on moist soils while also dehydrating more rapidly on dry soils. Thus, the benefit of moist soils may be greater in mesic populations. Alternatively, there may be selection for smaller body size in hatchlings from drier sites (Fig. 4C), enhancing their survival via a reduction of deleterious malformations due to lesions with the perivitelline membrane, but at a cost to swimming performance.

In summary, we show significant intra-specific variation in traits associated with desiccation tolerance in *P. guentheri* adults, and in the early developmental stages of their offspring. While dry rearing environments generally had a negative effect on traits putatively linked to fitness, the severity of treatment effects varied greatly among populations in a pattern consistent with the cline in annual precipitation. Together our findings suggest that water availability has influenced patterns of population variation in desiccation tolerance in *P. guentheri*, mediated either by phenotypic plasticity, local adaptation, or a combination of both. Whether the species

will persist under the predicted rapid declines in annual rainfall in southwestern

Australia remains unclear. Prior work (Eads et al. 2012) revealed that at least one *P.*

guentheri population lacks sufficient additive genetic variation in desiccation

tolerance to adapt to changes in water availability. Depending on the generality of

these patterns, the species may be limited in its ability to adapt to conditions beyond

its current range (e.g. Kellermann et al. 2009). Yet, if the observed intra-specific

variation in desiccation tolerance is the result of local adaptation, dry-adapted

populations could possess a reservoir of genes that may facilitate adaptive

responses to changes in water availability. As many *P. guentheri* populations are

now isolated due to extensive habitat fragmentation (Arnold 1988; Hobbs 1993),

limiting gene flow (Cummins 2017), the dry-adapted populations identified here could

be used as source populations for assisted, targeted gene flow (Aitken and Whitlock

2013) in future efforts to conserve mesic populations.

ACKNOWLEDGEMENTS

This work would not have been possible without great support from Stewart

Macdonald who located many populations and collected breeding adults, and

Brighton Downing who provided extensive assistance in the field and laboratory. We

further thank our volunteers who assisted with collecting adults from breeding

choruses: Marcus Lee, J. P. Lawrence, Callum Donohue, Blair Bentley and

Savannah Victor. This research was supported by funding from the School of

Biological Sciences at the University of Western Australia, the ANZ Holsworth

Wildlife Research Endowment and the Australian Government's National

Environmental Science Programme through the Threatened Species Recovery Hub.

T. Rudin-Bitterli was supported by the International Postgraduate Research

Scholarship and the C.F.H. & E.A. Jenkins Postgraduate Research Scholarship.

REFERENCES

Aitken, S. N., and M. C. Whitlock. 2013. Assisted gene flow to facilitate local adaptation to

climate change. Annual Review of Ecology, Evolution, and Systematics 44:367–388.

Albert, C. H., W. Thuiller, N. G. Yoccoz, R. Douzet, S. Aubert, and S. Lavorel. 2010. A multi-

trait approach reveals the structure and the relative importance of intra- vs. interspecific

variability in plant traits. Functional Ecology 24:1192–1201.

Andrewartha, S. J., N. J. Mitchell, and P. B. Frappell. 2008. Phenotypic differences in

terrestrial frog embryos: effect of water potential and phase. The Journal of Experimental

Biology 211:3800–3807.

Andrich, M. A., and J. Imberger. 2013. The effect of land clearing on rainfall and fresh water

resources in Western Australia: a multi-functional sustainability analysis. International

Journal of Sustainable Development & World Ecology 20:549–563.

Anstis, M. 2010. A comparative study of divergent embryonic and larval development in the

Australian frog genus *Geocrinia* (Anura: Myobatrachidae). Records of the Western

Australian Museum 25:399–440.

Anstis, M. 2013. Tadpoles and frogs of Australia. New Holland Publishers, London.

Arnold, G. W. 1988. Possible effects of climatic change on wildlife in Western Australia.

Pages 375–386 *in* G. I. Pearman, ed. Greenhouse: Planning for climate change. E. J. Brill,

Leiden.

Arthur, A. L., A. R. Weeks, and C. M. Sgrò. 2008. Investigating latitudinal clines for life

history and stress resistance traits in *Drosophila simulans* from eastern Australia. Journal of

Evolutionary Biology 21:1470–1479.

Bates, B. C., P. Hope, B. Ryan, I. Smith, and S. Charles. 2008. Key findings from the Indian

Ocean Climate Initiative and their impact on policy development in Australia. Climatic Change 89:339–354.

Bates, D., M. Mächler, B. M. Bolker, and S. C. Walker. 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67:1–48.

Bentley, P. J., A. K. Lee, and A. R. Main. 1958. Comparison of dehydration and hydration of two genera of frogs (*Heleioporus* and *Neobatrachus*) that live in areas of varying aridity. Experimental Biology 35:677–684.

Bolnick, D. I., P. Amarasekare, M. S. Araújo, R. Bürger, J. M. Levine, M. Novak, V. H. W. Rudolf, et al. 2011. Why intraspecific trait variation matters in community ecology. Trends in Ecology and Evolution 26:183–192.

Bolnick, D. I., R. Svanbäck, J. A. Fordyce, L. H. Yang, J. M. Davis, C. D. Hulsey, and M. L. Forister. 2003. The ecology of individuals: incidence and implications of individual specialization. The American Naturalist 161:1–28.

Box, G. E. P., and D. R. Cox. 1964. An analysis of transformations. Journal of the Royal Statistical Society. Series B (Methodological) 26:211–252.

Bradford, D. F., and R. S. Seymour. 1985. Energy conservation during the delayed-hatching period in the frog *Pseudophryne bibroni*. Physiological Zoology 58:491–496.

———. 1988*a*. Influence of water potential on growth and survival of the embryo, and gas conductance of the egg, in a terrestrial breeding frog, *Pseudophryne bibronii*. Physiol. Zool. 61:470–474.

———. 1988*b*. Influence of environmental PO2 on embryonic oxygen consumption, rate of development, and hatching in the frog *Pseudophryne bibroni*. Physiological Zoology 61:475–482.

Browne, R. K., J. Clulow, and M. Mahony. 2001. Short-term storage of cane toad (*Bufo marinus*) gametes. Reproduction 121:167–173.

Byrne, P. G., C. Dunne, A. J. Munn, and A. J. Silla. 2015. Environmental osmolality influences sperm motility activation in an anuran amphibian. Journal of Evolutionary Biology 28:521–534.

Carroll, E. J., and J. L. Hedrick. 1974. Hatching in the toad *Xenopus laevis*: morphological

events and evidence for a hatching enzyme. Developmental biology 38:1–13.

Catullo, R. A., P. Doughty, and J. S. Keogh. 2014. A new frog species (Myobatrachidae:

Uperoleia) from the Northern Deserts region of Australia, with a redescription of *U.*

trachyderma. Zootaxa 3753:251–262.

Claussen, D. L. 1974. Water balance and jumping ability in anuran amphibians. American

Zoologist 14:1257.

Cohen, K. L., M. A. Seid, and K. M. Warkentin. 2016. How embryos escape from danger: the

mechanism of rapid, plastic hatching in red-eyed treefrogs. The Journal of Experimental

Biology 219:1875–1883.

CSIRO and Bureau of Meteorology. 2015. *Climate change in Australia. Information for*

Australia's natural resource management regions: technical report. CSIRO and Bureau of

Meteorology, Australia.

CSIRO, and Bureau of Meteorology. 2016. *State of the Climate 2016*. CSIRO Publishing,

Collingwood, VIC, Australia.

Cummins, D. 2017. A genome-wide search for genetic diversity and local adaptation in a

terrestrial breeding frog highlights vulnerability to climate change. Honours Thesis. University

of Western Australia.

De Andrade, D. V., and A. S. Abe. 1997. Evaporative water loss and oxygen uptake in two

casque-headed tree frogs, *Aparasphenodon brunoi* and *Corythomantis greeningi* (Anura,

Hylidae). Comparative Biochemistry and Physiology - A Physiology 118:685–689.

Des Roches, S., D. M. Post, N. E. Turley, J. K. Bailey, A. P. Hendry, M. T. Kinnison, J. A.

Schweitzer, et al. 2018. The ecological importance of intraspecific variation. Nature Ecology

& Evolution 2:57–64.

Donnellan, S. C., M. J. Mahony, and T. Bertozzi. 2012. A new species of *Pseudophryne*

(Anura: Myobatrachidae) from the central Australian ranges. Zootaxa 69–85.

Duellmann, W. E., and L. Trueb. 1986. Biology of amphibians. McGraw-Hill, New York, NY.

Dumas, P. C. 1966. Studies of the *Rana* species complex in the Pacific Northwest. Copeia

1966:60–74.

Dziminski, M. A., J. D. Roberts, and L. W. Simmons. 2008. Fitness consequences of

parental compatibility in the frog *Crinia georgiana*. Evolution 62:879–886.

Eads, A. R., N. J. Mitchell, and J. P. Evans. 2012. Patterns of genetic variation in desiccation

tolerance in embryos of the terrestrial-breeding frog, *Pseudophryne guentheri*. Evolution

66:2865–2877.

Endler, J. A. 1977. Geographic variation, speciation, and clines. Monographs in population

biology 10:1–246.

Ewer, R. F. 1952. The effects of posterior pituitary extracts on water balance in *Bufo carens*

and *Xenopus laevis*, together with some general considerations of anuran water economy.

Journal of Experimental Biology 29:429–439.

Feder, M. E., and P. L. Londos. 1984. Hydric constraints upon foraging in a terrestrial

salamander, *Desmognathus ochrophaeus* (Amphibia: Plethodontidae). Oecologia 64:413–

418.

Gallant, A. J. E., K. J. Hennessy, and J. S. Risbey. 2007. Trends in rainfall indices for six

Australian regions : 1910-2005. Australian Meteorolgical Magazine 56:223–239.

Gatten, R. E. 1987. Activity metabolism of anuran amphibians: Tolerance to dehydration.

Physiological Zoology 60:576–585.

Gilchrist, G. W., L. M. Jeffers, B. West, D. G. Folk, J. Suess, and R. B. Huey. 2008. Clinal

patterns of desiccation and starvation resistance in ancestral and invading populations of

Drosophila subobscura. Evolutionary Applications 1:513–523.

Gosner, K. L. 1960. A simplified table for staging anuran embryos and larvae with notes on

identification. Herpetologica 16:183–190.

Hillman, S. S. 1987. Dehydrational effects on cardiovascular and metabolic capacity in two

amphibians. Physiological Zoology 60:608–613.

Hobbs, R. J. 1993. Effects of landscape fragmentation on ecosystem processes in the

Western Australian wheatbelt. Biological Conservation 64:193–201.

Hoffmann, A. A., A. Anderson, and R. Hallas. 2002. Opposing clines for high and low

temperature resistance in *Drosophila melanogaster*. Ecology Letters 5:614–618.

Hoffmann, A. A., and L. G. Harshman. 1999. Desiccation and starvation resistance in

Drosophila: patterns of variation at the species, population and intrapopulation levels.

Heredity 83:637–643.

Hoffmann, A. A., and C. M. Sgro. 2011. Climate change and evolutionary adaptation. Nature

470:479–485.

Hoppe, D. M. 1978. Thermal tolerance in tadpoles of the chorus frog *Pseudacris triseriata*.

Herpetologica 34:318–321.

IOCI. 2012. *Western Australia's weather and climate: a synthesis of Indian Ocean Climate*

Initiative stage 3 research: Summary for policymakers. (B. Bates, C. Frederiksen, & J.

Wormworth, eds.). Indian Ocean Climate Initiative (IOCI), Commonwealth Scientific and

Industrial Research Organization (CSIRO) and Bureau of Meteorology (BoM), BoM,

Melbourne, VIC, Australia.

Kawaguchi, M., H. Fujita, N. Yoshizaki, J. Hiroi, H. Okouchi, Y. Nagakura, T. Noda, et al.

2009. Different hatching strategies in embryos of two species, pacific herring *Clupea pallasii*

and Japanese anchovy *Engraulis japonicus*, that belong to the same order Clupeiformes,

and their environmental adaptation. Journal of Experimental Zoology Part B: Molecular and

Developmental Evolution 312:95–107.

Keller, I., J. M. Alexander, R. Holderegger, and P. J. Edwards. 2013. Widespread phenotypic

and genetic divergence along altitudinal gradients in animals. Journal of Evolutionary Biology

26:2527–2543.

Kellermann, V., B. Van Heerwaarden, C. M. Sgrò, and A. A. Hoffmann. 2009. Fundamental

evolutionary limits in ecological traits drive *Drosophila* species distributions. Science

325:1244–1246.

Kelly, M. W., E. Sanford, and R. K. Grosberg. 2012. Limited potential for adaptation to

climate change in a broadly distributed marine crustacean. Proceedings of the Royal Society

B: Biological Sciences 279:349–356.

Köhler, A., J. Sadowska, J. Olszewska, P. Trzeciak, O. Berger-Tal, and C. R. Tracy. 2011.

Staying warm or moist? Operative temperature and thermal preferences of common frogs (*Rana temporaria*), and effects on locomotion. Herpetological Journal 21:17–26.

Kolbe, J. J., M. Kearney, and R. Shine. 2010. Modeling the consequences of thermal trait variation for the cane toad invasion of Australia. Ecological Applications 20:2273–2285.

Kouba, A. J., C. K. Vance, M. A. Frommeyer, and T. L. Roth. 2003. Structural and functional aspects of *Bufo americanus* spermatozoa: effects of inactivation and reactivation. Journal of Experimental Zoology. Part A, Comparative Experimental Biology 295:172–82.

Kubota, M., T. Hasegawa, T. Nakakura, H. Tanii, M. Suzuki, and S. Tanaka. 2006. Molecular and cellular characterization of a new aquaporin, AQP-x5, specifically expressed in the small granular glands of *Xenopus* skin. J Exp Biol 209:3199–3208.

Langerhans, R. B. 2009. Morphology, performance, fitness: functional insight into a post-Pleistocene radiation of mosquitofish. Biology Letters 5:488–491.

Liu, J. N., and P. C. L. Hou. 2012. Cutaneous resistance to evaporative water loss in Taiwanese arboreal rhacophorid frogs. Zoological Studies 51:988–995.

Llewelyn, J., S. L. Macdonald, A. Hatcher, C. Moritz, and B. L. Phillips. 2016. Intraspecific variation in climate-relevant traits in a tropical rainforest lizard. Diversity and Distributions 22:1000–1012.

Martin, A. A., and A. K. Cooper. 1972. The ecology of terrestrial anuran eggs, genus *Crinia* (Leptodactylidae). Copeia 1:163–168.

McClanahan, L. J., and R. Baldwin. 1969. Rate of water uptake through the integument of the desert toad, *Bufo punctatus*. Comparative Biochemistry And Physiology 28:381–389.

Mitchell, N. J. 2001. Males call more from wetter nests: effects of substrate water potential on reproductive behaviours of terrestrial toadlets. Proceedings. Biological sciences / The Royal Society 268:87–93.

———. 2002. Low tolerance of embryonic desiccation in the terrestrial nesting frog *Bryobatrachus nimbus* (Anura: Myobatrachinae). Copeia 2002:364–373.

Moore, F. R., and R. E. J. Gatten. 1989. Locomotor performance of hydrated, dehydrated, and osmotically stressed anuran amphibians. Herpetologica 45:101–110.

Moran, E. V., F. Hartig, and D. M. Bell. 2016. Intraspecific trait variation across scales:

Implications for understanding global change responses. Global Change Biology 22:137–

150.

Nokhbatolfoghahai, M., and J. R. Downie. 2007. Amphibian hatching gland cells: Pattern and

distribution in anurans. Tissue and Cell 39:225–240.

Noldus, L. P. J. J., A. J. Spink, and R. A. J. Tegelenbosch. 2001. EthoVision: A versatile

video tracking system for automation of behavioral experiments. Behavior Research

Methods, Instruments & Computers 33:398–414.

Ogushi, Y., A. Tsuzuki, M. Sato, H. Mochida, R. Okada, M. Suzuki, S. D. Hillyard, et al.

2010. The water-absorption region of ventral skin of several semiterrestrial and aquatic

anuran amphibians identified by aquaporins. American Journal of Physiology-Regulatory,

Integrative and Comparative Physiology 299:R1150–R1162.

Packard, G. C. 1991. Physiological and ecological importance of water to embryos of

oviparous reptiles. Pages 213–228 *in* D. C. Deeming and M. W. J. Ferguson, eds. Egg

incubation: Its effects on embryonic development in brids and reptiles. Cambridge University

Press.

Packard, G. C. 1999. Water relations of chelonian eggs and embryos: is wetter better?

American Zoologist 39:289–303.

Plowman, M. C., S. Grbac-Ivankovic, J. Martin, S. M. Hopfer, and F. W. Sunderman. 1994.

Malformations persist after metamorphosis of *Xenopus laevis* tadpoles exposed to Ni2+,

Co2+, or Cd2+ in FETAX assays. Teratogenesis, Carcinogenesis, and Mutagenesis 14:135–

144.

Podrabsky, J. E., A. Tingaud-Sequeira, and J. Cerdà. 2010. Metabolic dormancy and

responses to environmental desiccation in fish embryos. Topics in Current Genetics 21:203–

226.

Pontes-da-Silva, E., W. E. Magnusson, B. Sinervo, G. H. Caetano, D. B. Miles, G. R. Colli, L.

M. Diele-Viegas, et al. 2018. Extinction risks forced by climatic change and intraspecific

variation in the thermal physiology of a tropical lizard. Journal of Thermal Biology 73:50–60.

R Development Core Team. 2017. A Language and Environment for Statistical Computing,

Version 3.4.3. R Foundation for Statistical Computing, Vienna, Austria: URL http://www.R-

project.org/.

Rajpurohit, S., and O. Nedved. 2013. Clinal variation in fitness related traits in tropical

drosophilids of the Indian subcontinent. Journal of Thermal Biology 38:345–354.

Räsänen, K., A. Laurila, and J. Merilä. 2003. Geographic variation in acid stress tolerance of

the moor frog, *Rana arvalis*. I. Local adaptation. Evolution 57:352–362.

Rehm, E. M., P. Olivas, J. Stroud, and K. J. Feeley. 2015. Losing your edge: Climate change

and the conservation value of range-edge populations. Ecology and Evolution 5:4315–4326.

Silla, A. J. 2011. Effect of priming injections of luteinizing hormone-releasing hormone on

spermiation and ovulation in Günther's Toadlet, *Pseudophryne guentheri*. Reproductive

Biology and Endocrinology 9:68.

Sinclair, B. J., C. M. Williams, and J. S. Terblanche. 2012. Variation in thermal performance

among insect populations. Physiological and Biochemical Zoology 85:594–606.

Smith, I. 2004. An assessment of recent trends in Australian rainfall. Australian

Meteorological Magazine 53:163–173.

Smith, I., and S. Power. 2014. Past and future changes to inflows in Perth (Western

Australia) dams. Journal of Hydrology: Regional studies 2:84–96.

Suzuki, M., T. Hasegawa, Y. Ogushi, and S. Tanaka. 2007. Amphibian aquaporins and

adaptation to terrestrial environments: A review. Comparative Biochemistry and Physiology -

A Molecular and Integrative Physiology 148:72–81.

Taigen, T. L., F. H. Pough, and M. M. Stewart. 1984. Water balance of terrestrial anuran

(*Eleutherodactylus coqui*) eggs: importance of parental care. Ecology 65:248–255.

Teplitsky, C., S. Plenet, J. P. Léna, N. Mermet, E. Malet, and P. Joly. 2005. Escape

behaviour and ultimate causes of specific induced defences in an anuran tadpole. Journal of

Evolutionary Biology 18:180–190.

Titon, B., and F. R. Gomes. 2015. Relation between water balance and climatic variables

associated with the geographical distribution of anurans. PLoS ONE 10:e0140761.

doi:10.1371/ journal.pone.0140761.

Titon, B., C. A. Navas, J. Jim, and F. R. Gomes. 2010. Water balance and locomotor

performance in three species of neotropical toads that differ in geographical distribution.

Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology

156:129–135.

Toledo, R. C., and C. Jared. 1993. Cutaneous adaptations to water balance in amphibians.

Comparative Biochemistry and Physiology -- Part A: Physiology 105:593–608.

Tyler, M. J., and P. Doughty. 2009. Frogs of Western Australia. Western Australian Museum,

Perth.

Valladares, F., S. Matesanz, F. Guilhaumon, M. B. Araújo, L. Balaguer, M. Benito-Garzón,

W. Cornwell, et al. 2014. The effects of phenotypic plasticity and local adaptation on

forecasts of species range shifts under climate change. Ecology Letters 17:1351–1364.

Van Berkum, F., F. H. Pough, M. M. Stewart, and P. F. Brussard. 1982. Altitudinal and

interspecific differences in the rehydration abilities of Puerto Rican frogs. Physiological

Zoology 55:130–136.

Venables, W. N., and B. D. Ripley. 2002. Modern applied statistics with S (fourth edi.).

Springer, New York.

Walker, J. A., C. K. Ghalambor, O. L. Griset, D. McKenney, and D. N. Reznick. 2005. Do

faster starts increase the probability of evading predators? Functional Ecology 19:808–815.

Warkentin, K. M. 2011*a*. Plasticity of hatching in amphibians: Evolution, trade-offs, cues and

mechanisms. Integrative and Comparative Biology 51:111–127.

———. 2011*b*. Environmentally cued hatching across taxa: Embryos respond to risk and

opportunity. Integrative and Comparative Biology 51:14–25.

Watkins, T. B. 1996. Predator-mediated selection on burst swimming performance in

tadpoles of the Pacific tree frog, *Pseudacris regilla*. Physiological Zoology 69:154–167.

Webb, P. W. 1986. Effect of body form and response threshold on the vulnerability of four

species of teleost prey attacked by largemouth bass (*Micropterus salmoides*). Canadian

Journal of Fisheries and Aquatic Sciences 43:763–771.

Wells, K. D. 2007. Water relations. Pages 82–121 *in* K. D. Wells, ed. The Ecology and

Behavior of Amphibians. The University of Chicago Press, Chicago.

Wilson, R. S., and C. E. Franklin. 1999. Thermal acclimation of locomotor performance in

tadpoles of the frog *Limnodynastes peronii*. Journal of Comparative Physiology - B

Biochemical, Systemic, and Environmental Physiology 169:445–451.

Winters, A., and M. E. Gifford. 2013. Geographic variation in the water economy of a

lungless salamander. Herpetological Conservation and Biology 8:741–747.

Withers, P. C., S. S. Hillman, R. C. Drewes, and O. M. Sokol. 1982. Water loss and nitrogen

excretion in sharp-nosed reed frogs (*Hyperolius nasutus*: anura, Hyperoliidae). The Journal

of Experimental Biology 97:335–343.

Wygoda M. 1984. Low cutaneous evaporative water loss in arboreal frogs. Physiological

Zoology 57:329–337.

Young, J. E., K. A. Christian, S. Donnellan, C. R. Tracy, and D. Parry. 2005. Comparative

analysis of cutaneous evaporative water loss in frogs demonstrates correlation with

ecological habits. Physiological and Biochemical Zoology 78:847–856.

 TABLES

 Table 1. Site characteristics and sample numbers (N) for each *P. guentheri*

 population sampled, with populations numbered by increasing aridity.

Collection site	Population No	Longitude	Latitude	N ♂	♀	F_1*	Annual mean precipitation (mm)	Annual mean temperature (°C)
Chidlow	1	31°53'05.5"S,	116°18'48.0"E	17	5	647	788	17.2
Flint Plot	2	32°17'01.4"S,	116°31'24.1"E	12	4	453	654	16.7
Pingelly	3	32°28'25.9"S,	116°58'27.9"E	19	8	911	428	16.6
Dudinin	4	32°49'17.4"S,	117°53'01.0"E	18	0	0	358	16.5
Binnu	5	28°02'30.8"S,	114°39'36.0"E	19	5	870	352	19.9
Mullewa	6	28°31'07.3"S,	115°38'11.4"E	10	0	0	329	20.5

889 Note: Climate data were obtained from the Bureau of Meteorology and are

890 interpolated values for the specific coordinates of each population, averaged from

891 1980 to 2017. *First generation offspring obtained via within-population parental

892 crosses from four localities.

 Table 2. Mixed-effects model results of embryonic and larval *P. guentheri* traits

 associated with desiccation tolerance

Trait	*N*	Source	df	X^2	*P*	Sig.
Embryonic survival (%)	2844	Treatment	2	98.398	< 0.001	***
		Population	3	8.874	0.031	*
		Population x Treatment	6	19.924	0.003	**
		Ovum size	1	0.926	0.336	ns
Time to hatching (h)	2378	Treatment	2	745.596	< 0.001	***
		Population	3	104.731	< 0.001	***
		Population x Treatment	6	173.249	< 0.001	***
		Ovum size	1	0.455	0.499	ns
Wet weight at hatching (mg)	2344	Treatment	2	864.308	< 0.001	***
		Population	3	5.641	0.130	ns
		Population x Treatment	6	295.199	< 0.001	***
		Ovum size	1	3.770	0.05	*
Proportion of hatchlings malformed	2382	Treatment	2	59.152	< 0.001	***
		Population	3	5.934	0.115	ns
		Population x Treatment	6	2.940	0.816	ns
		Ovum size	1	2.545	0.111	ns
Proportion of notochord malformations	2382	Treatment	2	2.929	0.231	ns
		Population	3	18.234	< 0.001	***
		Population x Treatment	6	12.272	0.056	ns
		Ovum size	1	0.006	0.939	ns
Proportion of edema malformations	2382	Treatment	2	2.811	0.245	ns
		Population	3	13.673	0.003	**
		Population x Treatment	6	2.350	0.885	ns
		Ovum size	1	1.489	0.222	ns

895　**Table 3.** Mixed-effects model results of swimming performance traits in *P. guentheri*

896　hatchlings originating from four populations along a natural rainfall gradient.

Trait	N	Source	df	X^2	P	Sig.
Maximum velocity (cm s^{-1})	556	Treatment	2	197.270	< 0.001	***
		Population	3	12.499	0.006	**
		Population x Treatment	6	95.770	< 0.001	***
		Ovum size	1	0.4347	0.510	ns
Mean velocity (cm s^{-1})	556	Treatment	2	197.270	< 0.001	***
		Population	3	12.499	0.006	**
		Population x Treatment	6	95.770	< 0.001	***
		Ovum size	1	0.4347	0.510	ns
Mean meander (deg cm^{-1})	556	Treatment	2	138.066	< 0.001	***
		Population	3	11.747	0.008	**
		Population x Treatment	6	25.783	< 0.001	***
		Ovum size	1	4.518	0.036	*
Total distance moved (cm)	556	Treatment	2	66.881	< 0.001	***
		Population	3	5.699	0.127	ns
		Population x Treatment	6	17.385	0.008	**
		Ovum size	1	1.765	0.184	ns

 FIGURE LEGENDS

Figure 1. Map showing the distribution of *P. guentheri* in Western Australia (grey

line; based on occurrence records from the Atlas of Living Australia) and the location

of collection sites, overlaid with annual mean rainfall data (mm). Populations are

numbered by increasing aridity. Female *P. guentheri* left, male right.

Figure 2. Types of malformations in *P. guentheri* hatchlings scored in this study: (A)

normal shaped hatchling without malformations, (B) hatchling with scoliosis, (C)

hatchling with lordosis, (D) hatchling with kyphosis and (E) hatchling with edema.

Figure 3. Variation in (A) dehydration and (B) rehydration rates (mean ± SE) among

the six *P. guentheri* populations located along a natural rainfall gradient (790 - 330

mm/year). The x-axis lists populations from wettest to driest sites. Populations that

do not share the same letters are significantly different (Tukey HSD post-hoc tests, *P*

< 0.05). Relationship between annual mean rainfall (mm) and (C) dehydration rate

(mean ± SE) and (D) rehydration rate (mean ± SE) in six *P. guentheri* populations.

For statistics, see results section.

Figure 4.

Embryonic and larval *Pseudophryne guentheri* trait responses (mean with 95%

confidence intervals) for four populations, reared on soil at three water potentials; dry

(white circles), intermediate (grey circles) and wet (black circles). The x-axis shows

populations arranged from the wettest (1) to the driest (5) sites. (A) embryonic

survival (proportion of fertilised eggs that hatched), (B) time to hatching (h), (C) wet

weight at hatching (mg), (D) proportion of malformed hatchlings, (E) proportion of

malformations in the notochord category and (F) proportion of malformations in

the edema category.

Figure 5.

Swimming performance (mean with 95% confidence intervals) of *P. guentheri*

hatchlings from four populations, reared on soil at three water potentials; dry (white

circles), intermediate (grey circles) and wet (black circles). The x-axis shows

populations arranged from the wettest (1) to the driest (5) sites. (A) maximum

velocity (cm s^{-1}), (B) mean velocity (cm s^{-1}), (C) mean meander (deg cm $^{-1}$) and (D)

total distance moved (cm).

 FIGURES

 Fig.1

934 **Fig.2**

 Fig.3

Fig.4

 Fig.5