Early transcriptional responses after dengue vaccination mirror the

response to natural infection and predict neutralizing antibody titers

Running title: Early correlates of immunity to dengue

Stephen J. Popper[a], Fiona R. Strouts[a*], Janet C. Lindow[b*#], Henry K. Cheng[a], Magelda

Montoya[c], Angel Balmaseda[d], Anna P. Durbin[e], Stephen S. Whitehead[f], Eva Harris[c], Beth D.

Kirkpatrick[b], David A. Relman[a,g,h]

[a] Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA

[b] Vaccine Testing Center, University of Vermont College of Medicine, Burlington, VT 05401,

USA

[c] Division of Infectious Diseases and Vaccinology, School of Public Health, University of

California, Berkeley, Berkeley, CA 94720-3370, USA

[d] Laboratorio Nacional de Virología, Centro Nacional de Diagnóstico y Referencia, Ministry of

Health, Managua, Nicaragua

[e] Center for Immunization Research (CIR), Johns Hopkins Bloomberg School of Public Health,

Baltimore, MD 21205, USA

[f] Laboratory of Infectious Diseases, National Institute for Allergy and Infectious Diseases,

Bethesda, MD 20892, USA

[g] Department of Microbiology & Immunology, Stanford University School of Medicine, Stanford,

CA 94305, USA

[h] Veterans Affairs Palo Alto Health Care System, Palo Alto, CA 94304, USA

* These two authors contributed equally to the work

Correspondence: D. A. Relman, VA Palo Alto Health Care System 154T, 3801 Miranda Avenue,

Palo Alto, CA 94304; 650-736-6822 tel; 650-852-3291 fax; relman@stanford.edu

Current address: Center for Mental Health Research and Recovery, Montana State University,

Bozeman, MT 59717, USA

Abstract: 199 words

Text: 3450 words

Financial Support. This work was supported by the National Institute of Allergy and Infectious

Diseases Division of Intramural Research and Division of Microbiology and Infectious Diseases

(U19 AI109761, D.A.R.; U54 AI065359, A.B.), by the Thomas C. and Joan M. Merigan

Endowment at Stanford University (D.A.R.), and by a grant (VE-1) from the Pediatric Dengue

Vaccine Initiative of the Bill and Melinda Gates Foundation (E.H.).

Conflict of Interest. All authors: No reported conflicts of interest.

ABSTRACT

Background: Several promising live attenuated virus (LAV) dengue vaccines are in development, but information about innate immune responses and early correlates of protection are lacking.

Methods: We characterized human genome-wide transcripts in whole blood from 10 volunteers at 11 time-points after immunization with the dengue virus type 3 (DENV-3) component of the NIH dengue vaccine candidate TV003 and from 30 hospitalized children with acute primary DENV-3 infection. We compared day-specific gene expression patterns with subsequent neutralizing antibody (NAb) titers.

Results: The transcriptional response to vaccination was largely confined to days 5-20 and was dominated by an interferon-associated signature and a cell cycle signature that peaked on days 8 and 14, respectively. Changes in transcript abundance were much greater in magnitude and scope in symptomatic natural infection than following vaccination (maximum fold-change >200 versus 21 post-vaccination; 3,210 versus 286 transcripts with significant fold-change), but shared gene modules were induced in the same sequence. The abundance of 131 transcripts on days 8 and 9 post-vaccination was strongly correlated with NAb titers measured 6 weeks post-vaccination.

Conclusions: LAV dengue vaccination elicits early transcriptional responses that mirror those found in symptomatic natural infection and provide candidate early markers of protection against DENV infection.

Clinical Trial Registration Number: NCT00831012 (available at clinicaltrials.gov)

Keywords: dengue; vaccine; innate immune response; gene expression; microarray; correlates of protection; interferon; neutralizing antibody

BACKGROUND

Each year, the four dengue virus serotypes (DENV-1-4) infect an estimated 390 million individuals globally [1]. While most of these infections are asymptomatic, approximately 100 million individuals develop clinically apparent disease ranging from uncomplicated fever to life-threatening illness. Despite the high disease burden, there are no licensed therapeutics for DENV infection. Several promising candidate dengue vaccines are in Phase III clinical trials, and the live attenuated chimeric dengue vaccine Dengvaxia™ was recently licensed for use in children 9 years of age and older in DENV endemic areas. However, the efficacy and duration of protection were limited or uncertain, and DENV-naïve vaccine recipients were hospitalized for dengue and severe dengue at a higher rate than placebo recipients, possibly due to antibody-dependent enhancement (ADE) [2].

Studies of natural DENV infection and flavivirus LAVs have identified immune responses needed for protection against dengue disease. Pre-existing neutralizing antibody (NAb) titers correlated with a lack of symptomatic disease in subsequent infections [3–6] and are used as the primary measure of candidate vaccine immunogenicity. However, the risk of severe disease is elevated after a second infection with a heterotypic dengue virus [7]. The recognition of effective homotypic immunity after natural infection has led to a common vaccine development strategy of inducing homotypic NAbs to all four serotypes simultaneously.

Little is known about the role of early innate immune responses in enhancing NAb production and promoting protective immune memory against dengue. Studies of innate immunity have been hampered by the difficulty inherent in identifying individuals with early infection, when innate immune responses are most active, particularly those with mild or subclinical infections. Trials of LAVs provide a unique opportunity to examine early immune responses in a setting where the time, dose, and viral serotype are known. Genome-wide transcript responses to vaccines have provided important clues about early steps in the generation of humoral and cellular immunity [8–13]. Transcript profiling of peripheral blood also

incorporates information from cell populations that are difficult to examine in clinical settings,

and has led to signatures associated with dengue disease severity, identified links between

innate responses and humoral immunity in secondary DENV infection, and illustrated the

dynamic nature of these responses [14–20].

In this study, we characterized the transcript response to rDEN3Δ30/31, the DENV-3

component of TV003, a tetravalent live attenuated vaccine candidate developed by NIH. TV003

is a single-dose vaccine that has proven to be both safe and immunogenic and is being

evaluated in a Phase III efficacy trial [21,22]. We examined the temporal course of changes in

transcript abundance and identified early signatures correlated with NAb titers measured six

weeks post-vaccination. We also compared these results with the transcript patterns we

observed in patients with symptomatic wild-type primary DENV-3 infection. Despite the

anticipated differences in the magnitude of expression, we observed the induction of common

gene expression programs in the same temporal sequence, with a similar relationship to the

induction of NAb. These results reveal candidate biomarkers of early protective DENV immune

responses against dengue and suggest a path towards validation and deployment.

METHODS

Vaccine study population. Samples for this study were collected from a Phase I clinical trial of

the live attenuated dengue vaccine rDEN3Δ30/31-7164 (DENV-3), described previously [23].

Briefly, healthy, flavivirus-naïve adult volunteers were enrolled and randomized to receive a

single 0.5 ml subcutaneous dose of 1,000 PFU of DENV-3 vaccine or a placebo (0.5 ml of

vaccine diluent). Blood samples including whole blood for RNA profiling (PAXgene, Preanalytix)

were collected immediately prior to vaccination and on days 2, 5, 6, 8, 9, 12,14, 20, 29, 42 and

180, and stored at -80°C until used. Samples from each of these time-points were available

from nine of ten vaccinees and from all placebo recipients. Subject 9 had samples available for

all days except days 8 and 12; 166 samples in total were used for analysis. Serum virus titers

(viremia) were measured using a standard plaque assay as described previously [24]. Serum

NAb titer was determined by 60% plaque reduction ($PRNT_{60}$) [25]. Seroconversion was defined

by a $\geq$4-fold increase in $PRNT_{60}$ on study day 28 or 42 relative to day 0 and corresponds to a

post-vaccination titer >10 [23].

Dengue patient population. Patients presenting with fever and suspected dengue during the

2010 dengue season were enrolled at the Hospital Infantil Manuel de Jesús Rivera (HIMJR) in

Managua, Nicaragua. Inclusion criteria, recruitment, and laboratory testing have been described

previously [26]; a full description is available in the Supplementary Information. Blood samples

from healthy subjects were collected as part of a separate prospective cohort study in which

healthy children in the same general population were enrolled without regard to dengue status

[27].

Ethics statement. The trial of rDEN3Δ30/31 was approved by the Committee for Human

Research at the University of Vermont, and written informed consent was obtained from all

subjects following a review of risks and benefits and a comprehension assessment. The study in

Nicaragua was approved by the Institutional Review Boards of the University of California,

Berkeley, and the Nicaraguan Ministry of Health, and by the Stanford University Administrative

Panel on Human Subjects in Medical Research. All clinical research followed human

experimentation guidelines of the United States Department of Health and Human Services

and/or those of the authors' institutions. Parents or legal guardians of all subjects provided

written informed consent, and subjects 6 years of age and older provided assent.

RNA sample processing and transcriptome analysis. PAXgene RNA was amplified and

hybridized to Human Exonic Evidence Based Oligonucleotide (HEEBO) microarrays [14].

Microarray data were submitted to the Princeton University MicroArray (PUMA) database for

normalization and gene filtering and are deposited at Gene Expression Omnibus

(http://www.ncbi.nlm.nih.gov/geo/; accession numbers GSE96656 and GSE98053). Data

analysis was carried out using packages cited in the main text; a full description of both sample

processing and analysis steps is available in the Supplementary Information.

RESULTS

Temporal patterns of the transcriptional responses to live dengue vaccination

To identify the temporal pattern of the early human transcriptional response to dengue

vaccination, we examined changes in genome-wide transcript abundance in serial whole blood

samples from 10 volunteers infected with 1,000 plaque forming units (pfu) of rDEN3Δ30/31, the

dose included in TV003, and four volunteers inoculated with placebo (L-15 medium). Nine of ten

vaccinees seroconverted 28 days post-vaccination, defined as a 60% plaque reduction

neutralization titer (PRNT$_{60}$) >10 (Table 1). Four of the vaccinees had low-level viremia on one

or more days within the first 10 days post-vaccination, five developed a mild maculopapular

rash, and none were febrile. The four placebo recipients remained seronegative for DENV

serotypes.

We collected whole blood for isolation of RNA immediately before vaccination (day 0),

and on days 2, 5, 6, 8, 9, 12, 14, 20, 29, 42 and 180 post-vaccination from all volunteers and

measured genome-wide transcript abundance levels. Data were available for eight of the nine

participants who seroconverted. For each of these eight subjects, we compared transcript

abundances for each post-vaccination day with those for the matched pre-vaccination sample

(see Supplementary Information). Almost all significant changes in transcript abundance

occurred 5-20 days after vaccination, with a peak of 161 and 156 transcripts changing in

abundance (days 8 and 9, respectively), and 286 transcripts with a significant change in

abundance on at least one day (Figure 1). Fewer transcripts met criteria for significance when

comparing vaccinees to placebo recipients (n=131), but the direction of change for 271 of the

286 transcripts from vaccinees was the same whether the comparison was with day-matched

placebo recipients or each subject's baseline sample (Supplementary Figure 1).

To infer the functional implications of these changes in transcript abundance, we used

hierarchical clustering to organize the transcripts and compared gene membership in Gene

Ontology and the KEGG pathways using the DAVID bioinformatics resource [28]. Gene

transcripts were grouped in three clusters (Figure 2 and Supplementary Figure 2). Transcripts in

Cluster 1 were more abundant after vaccination (Figure 2C), peaked on days 8 and 9 post

vaccination, and included canonical interferon-stimulated gene (ISG) transcripts; IFI44, IFI44L,

IFI27, HERC5, IFIT1, USP18, and ISG15 transcripts all increased 10- to 22-fold compared to

baseline. Cluster 1 was strongly enriched for genes involved in the innate immune response to

viruses and highly enriched for genes we previously found to be expressed after treatment of

PBMCs with type I interferon (p<1E-36) [29].

Gene transcripts in Clusters 2 and 3 showed maximal changes on day 14, with Cluster 2

transcripts increasing and Cluster 3 transcripts decreasing in abundance from baseline (Figure

2A and 2B). Cluster 2 included TYMS, CEP55, CCNA2, and NEK2, whose genes products are

involved in DNA replication and cell division, and other genes associated with mitosis (p<2E-9,

Figure 2C). Genes in Cluster 3 were enriched in both reticulocytes (p=1E-20) and neutrophils

(p=2E-7) [30]. We did not measure reticulocyte counts, but we did measure neutrophils and the

relative neutrophil abundance in vaccinees did not change significantly with time (p=0.55, paired

t-test), suggesting that decreased expression of these genes was not due to decreased

neutrophil abundance.

Changes observed after vaccination are a subset of those observed in natural

symptomatic DENV-3 infection

To establish which features of the early response to vaccination are shared with the

response to natural symptomatic infection, we examined transcript responses in Nicaraguan

children hospitalized with acute dengue. We previously demonstrated that a history of previous

DENV exposure is the most prominent source of variation in gene expression in dengue

patients [14]. To ensure that DENV immune status, as well as serotype, did not confound our

analysis, we identified 30 children diagnosed with acute primary DENV-3 infection during a

single year (Supplementary Table 1), and compared transcript abundance in whole blood with

measurements from 9 healthy individuals. Principal components analysis confirmed previous

findings that there are significant day-to-day changes in the transcript response to natural

infection [14,31] (Supplementary Figure 4); thus, we subsequently performed analyses stratified

by day of fever.

Despite having fewer days available for comparison and lacking baseline samples for

each patient, we identified many more transcripts with significant changes in abundance post-

infection compared to those found in vaccinees: among the 20,623 transcripts measured in both

datasets, we identified 3,210 transcripts that differed significantly on at least one day of fever,

compared with 278 transcripts following vaccination (Figure 3A and Supplementary Figure 5A).

The magnitude of the maximum change in abundance post-infection was also nearly 10-fold

greater: there was a 200-fold difference post-infection compared to a maximum 21-fold

difference post-vaccination (Figure 3B). The transcripts with the greatest differences in relative

abundance during natural infection were MT2A (242-fold) and USP18 (183-fold), both of which

are interferon-induced; HESX1 (150-fold), which is expressed in activated dendritic cells; and

SPAT2SL (137-fold), which may be involved in activation and differentiation of multiple cell

types.

Despite differences in response magnitude and number, the response following natural

symptomatic infection included 90% (250/278) of transcripts that changed after vaccination, and

the direction of change was the same for 96% of these transcripts (240/250) (Figure 3C). The

transcripts that changed the most post-vaccination (IFI44, IFI44L, IFI27 and HERC5) were

among the 20 transcripts with the biggest differences in abundance following natural infection,

and relative increases in transcript abundance were strongly correlated across the two groups

(Spearman r^2 = 0.75).

Responses to dengue vaccination and symptomatic natural infection share a common

temporal sequence

We used gene set enrichment analysis and information from all measured transcripts to

identify 141 blood transcript gene modules that changed in abundance following either

immunization or infection [8] (FDR<1%). Many of these modules demonstrated similar changes

in both vaccinees and patients (Figure 4A). Modules enriched for ISG expression were elevated

on days 5-14 post-vaccination and were also persistently elevated after natural DENV infection.

Modules representing monocyte-associated transcripts were elevated on days 1-3 of natural

infection and on days 8-9 post-vaccination, while modules associated with the mitotic cell cycle

were elevated on later days in both groups, with the highest levels on day 5 of natural infection

and on day 14 post-vaccination. When we compared the overall profiles of the gene modules in

the two groups, we found that the responses to natural infection on fever days 1-3 were most

similar to responses to vaccination on days 8-9 (Pearson's r ≥0.60; peak on day 9), while fever

day 4 was most similar to vaccination day 12 (r>0.75, peak on day 12), and fever day 5 was

most similar to vaccination day 14 and subsequent time-points (r≥0.70, peak on 14) (Figure 4B,

Supplementary Dataset 1). Thus, the enrichment of common modules in the same sequence

indicates a similar progression in the early host response to vaccination and to natural infection.

We note there was also a cluster of 16 gene modules, six associated with platelet

activation and cytoskeletal remodeling, that were elevated in natural infection but not vaccinees

(Figure 4A and Supplementary Dataset 1). Previous studies have demonstrated that platelet

activation and TGFβ expression are elevated in DENV infection and higher in patients with more

severe disease [32]. TGFβ, which is expressed at high levels in platelets [33], was elevated on

fever days 1-2 in dengue patients but was never elevated post-vaccination (Supplementary

Figure 6).

Early transcriptional responses linked to neutralizing antibody production

DENV-specific NAbs are the primary endpoint for assessing vaccine responses in clinical trials

and are associated with protection from both symptomatic infection and severe disease [3–5].

To determine whether changes in host transcript patterns predicted differences in NAb titer we

calculated the correlation between the change in abundance of each transcript on each day and

the NAb titer on post-vaccination day 42, when NAbs are generally at peak titer (Table 1,

Supplementary Figure 7). During the first 6 days post-vaccination, we found no significant

correlations with NAb titer, but by day 8, expression of the ISGs in Cluster 1 positively correlated

with the day 42 NAb titer ($p<0.01$; Figure 5). This correlation was equally strong on day 9, and

131 transcripts were significantly correlated with day 42 NAb titer on both days. Among the

individual ISG transcripts most strongly correlated with day 42 NAb titer on both days 8 and 9

($r>0.8$) was IFI44, the transcript whose abundance changed the most post-vaccination. IFI44

was also elevated at one time-point in each of two placebo recipients, but the timing of elevated

expression was different and correlated with unrelated respiratory viral infections in each

instance (Supplementary Figure 8). Twelve of the 131 transcripts were also associated with

subsequent development of a rash, which was the only significant correlate with positive NAb

titer in a clinical trial of TV003 [21] (Supplementary Figure 9). Interestingly, the one vaccinee

who failed to develop neutralizing antibodies showed little evidence of increased abundance in

Cluster 1 genes (Supplementary Figure 3). The association of interferon-related transcript

abundance and later NAb titer diminished on days 12 and 14, but BUB1 ($r=0.9$) and other

transcripts associated with the mitotic cell cycle were correlated with subsequent NAb titers on

day 14 (Figure 5).

When we performed similar comparisons for naturally infected patients, we found no

transcript clusters significantly correlated with either convalescent or three month NAb titer

(Supplementary Figure 5B and 5C). However, the pattern of blood transcript module

enrichment indicated a similar relationship between day-specific gene expression and later

production of NAb; gene enrichment for both interferon-stimulated and cell cycle-associated

gene modules was associated with higher NAb titer in both vaccinees and patients (Figure 6),

albeit more weakly in patients, and cell cycle-associated modules were correlated with NAb titer

later in both groups.

There are at least three subpopulations of monocytes with distinct transcript profiles [34];

Kwissa et al. identified an increase in $CD14^+CD16^+$ intermediate-phenotype population after

secondary DENV infection, and showed that in vitro these cells stimulated formation of the

plasmablasts that secrete antibodies weeks after infection, mediated in part by secretion of the

ISG cytokine BAFF [19]. In our study, gene set enrichment analysis indicated enrichment of

transcripts for both intermediate and nonclassical monocytes at multiple time-points in both

vaccinees and patients, while BAFF transcripts were most abundant on fever days 1 and 2 in

the patients and days 8 and 9 in the vaccinees (Supplementary Figure 10).

DISCUSSION

In this study, we used intensive longitudinal sampling to characterize the transcriptional

response to dengue vaccination, compared results with those from natural infection with the

same DENV serotype, and identified early features that may predict a protective immune

response. We found that vaccination and natural infection induced common gene expression

programs, and the abundance of individual interferon-stimulated transcripts 8 days post-

vaccination was correlated with NAb titers measured five weeks later, representing the earliest

identified correlates of a protective adaptive immune response following dengue vaccination.

An interferon response signature has been observed in other studies profiling viral

vaccine transcriptional responses. Inactivated influenza and meningococcal vaccines both

induce a mild interferon response during the first week post-vaccination, but the response is

particularly strong after vaccination with live attenuated vaccines [9,12,35]. We reported that

ISG expression was much stronger in cynomolgous macaques infected with wild-type DENV

compared to live attenuated virus [35]. Here, we found that ISG expression was much stronger

in symptomatic dengue patients than vaccinees, presumably due to higher viral load after

infection with wild-type virus. Expression of ISGs was correlated with viral load in the patients,

as seen in other studies [19,36]. However, this association did not persist when patients were

stratified by day of fever, highlighting the importance of temporal variation in the innate immune

response and in viral load, and suggesting that factors in addition to viral replication influence

ISG expression. Several studies have found stable inter-individual differences in the response to

interferon, suggesting that genetic and environmental features may affect the relationship

between viral infection and the interferon response [37,38].

The links between type I interferon production and NAb production are likely to involve

multiple cell types. Plasmacytoid dendritic cells (pDCs) contribute to B cell differentiation and

antibody production after viral infection [39]. In this study, increases in monocyte-associated

gene expression coincided with ISG expression, and we found features related to multiple

monocyte phenotypes in both natural infection and vaccination (Supplementary Figure 10C).

Gene module analysis also suggested that T cells were responsible for the increase in cell

cycle-associated transcripts two weeks after vaccination that was linked to NAb titers. Future

targeted studies of pDCs, monocytes, and T cell populations during the first two weeks post-

vaccination will help clarify their role in establishing long-lasting antibody responses. In addition,

the link between an early interferon response and later NAb titer was only apparent in natural

infection when we used a module analysis approach. This may indicate a plateau, or saturation

effect, in the relationship between ISG expression and antibody titer. Alternatively, it may reflect

the variability in pathogen dose, prior health status and/or days of infection absent in clinical

trials but inherent in observational studies.

Comparison with LAV vaccination also provides a framework for identification of features

associated with pathogenic versus non-pathogenic infection. A recent study compared PBMC

gene expression in asymptomatic and clinically significant secondary DENV infection and

identified differences in antigen presentation and lymphocyte activation [36]. In this study

examining whole blood gene expression during primary infection, we found an increased

abundance of transcripts associated with platelet activation in natural (pathogenic) infection but

not vaccination (non-pathogenic infection), consistent with the hypothesis that platelet activation

contributes to dengue pathogenesis [40].

Neutralizing antibody titers were used as an endpoint for these vaccine studies because

many studies have shown that these antibodies play an important role in protective immunity.

However, recent work has demonstrated that NAbs measured in vitro are an imperfect correlate

of in vivo protection [37,38]. Immunity mediated by NAbs may be neither life-long nor sterilizing

[43,44] and will be affected by the quality as well as the quantity of NAbs [5,26,45]. Recent

studies also highlight a likely role for cytotoxic T cells in mediating protection against DENV

reinfection and severe disease [46–49]. The NIH tetravalent vaccine, of which rDEN3Δ30/31 is

a component, elicits CD4$^+$ T cell responses similar to those seen in natural infection [50]. It will

be important to establish whether the early transcript-based features we measured in this study

are associated with DENV-specific responses in memory T cell populations.

Our findings should be validated using a tetravalent dengue vaccine formulation. We

previously studied transcript-based responses to a different tetravalent dengue vaccine in

nonhuman primates and found that the interferon response was associated with antibody

formation [35]. We believe it is likely that the same relationship between early transcriptional

responses and neutralizing antibodies will exist in humans immunized with tetravalent LAV

dengue vaccines. The initiation of Phase 3 clinical trials of TetraVax-DV-TV003 provides the

opportunity to establish whether specific transcriptional profiles can be used as early surrogate

markers of both immunogenicity and protection.

Acknowledgements

The authors thank Cassandra Ventrone for assistance with sample collection and logistics,

Chunling Wang for providing viral load data, Ellen Sebastian for assistance with data

processing, and Elizabeth Costello for helpful suggestions and editing.

References

1. Bhatt S, Gething PW, Brady OJ, et al. The global distribution and burden of dengue.
Nature 2013; 496(7446):504–507.

2. Hadinegoro SR, Arredondo-García JL, Capeding MR, et al. Efficacy and Long-Term
Safety of a Dengue Vaccine in Regions of Endemic Disease. N Engl J Med 2015;
373(13):1195–1206.

3. Endy TP, Nisalak A, Chunsuttitwat S, et al. Relationship of Preexisting Dengue Virus
(DV) Neutralizing Antibody Levels to Viremia and Severity of Disease in a Prospective
Cohort Study of DV Infection in Thailand. J Infect Dis 2004; 189(6):990–1000.

4. Corbett KS, Katzelnick L, Tissera H, Amerasinghe A, Silva AD de, Silva AM de.
Preexisting Neutralizing Antibody Responses Distinguish Clinically Inapparent and
Apparent Dengue Virus Infections in a Sri Lankan Pediatric Cohort. J Infect Dis 2015;
211(4):590–599.

5. Katzelnick LC, Montoya M, Gresh L, Balmaseda A, Harris E. Neutralizing antibody titers
against dengue virus correlate with protection from symptomatic infection in a
longitudinal cohort. Proc Natl Acad Sci (USA) 2016; 113(3):728–733.

6. Sabin AB. Research on Dengue during World War II. Am J Trop Med Hyg 1952;
1(1):30–50.

7. Sangkawibha N, Rojanasuphot S, Ahandrik S, et al. Risk factors in dengue shock
syndrome: A prospective epidemiologic study in Rayong, Thailand. I. The 1980
outbreak. Am J Epidemiol 1984; 120(5):653–669.

8. Li S, Rouphael N, Duraisingham S, et al. Molecular signatures of antibody responses
derived from a systems biological study of 5 human vaccines. Nat Immunol 2014;
15(2):195–204.

9. Nakaya HI, Wrammert J, Lee EK, et al. Systems biology of vaccination for seasonal
influenza in humans. Nat Immunol 2011; 12(8):786–795.

10. Bucasas KL, Franco LM, Shaw CA, et al. Early patterns of gene expression correlate
with the humoral immune response to influenza vaccination in humans. J Infect Dis
2011; 203(7):921–929.

11. Obermoser G, Presnell S, Domico K, et al. Systems scale interactive exploration reveals
quantitative and qualitative differences in response to influenza and pneumococcal
vaccines. Immunity 2013; 38(4):831–844.

12. Querec TD, Akondy RS, Lee EK, et al. Systems biology approach predicts
immunogenicity of the yellow fever vaccine in humans. Nat Immunol 2009;
10(1):116–125.

13. Kazmin D, Nakaya HI, Lee EK, et al. Systems analysis of protective immune responses
to RTS,S malaria vaccination in humans. Proc Natl Acad Sci (USA) 2017; 114(9):2425–
2430.

14. Popper SJ, Gordon A, Liu M, Balmaseda A, Harris E, Relman DA. Temporal dynamics of
the transcriptional response to dengue virus infection in Nicaraguan children. PLoS
Negl Trop Dis 2012; 6(12):e1966.

15. Long HT, Hibberd ML, Hien TT, et al. Patterns of gene transcript abundance in the
blood of children with severe or uncomplicated dengue highlight differences in
disease evolution and host response to dengue virus infection. J Infect Dis 2009;
199(4):537–546.

16. Simmons CP, Popper S, Dolocek C, et al. Patterns of host genome-wide gene transcript
abundance in the peripheral blood of patients with acute dengue hemorrhagic fever. J
Infect Dis 2007; 195(8):1097–1107.

17. Nascimento EJM, Braga-Neto U, Calzavara-Silva CE, et al. Gene expression profiling
during early acute febrile stage of dengue infection can predict the disease outcome.
PloS One 2009; 4(11):e7892.

18. Loke P, Hammond SN, Leung JM, et al. Gene expression patterns of dengue virus-
infected children from nicaragua reveal a distinct signature of increased metabolism.
PLoS Negl Trop Dis 2010; 4(6):e710.

19. Kwissa M, Nakaya HI, Onlamoon N, et al. Dengue virus infection induces expansion of a
CD14(+)CD16(+) monocyte population that stimulates plasmablast differentiation.
Cell Host Microbe 2014; 16(1):115–127.

20. Weg CAM van de, Ham H-J van den, Bijl MA, et al. Time since onset of disease and
individual clinical markers associate with transcriptional changes in uncomplicated
dengue. PLoS Negl Trop Dis 2015; 9(3):e0003522.

21. Durbin AP, Kirkpatrick BD, Pierce KK, et al. A single dose of any of four different live
attenuated tetravalent dengue vaccines is safe and immunogenic in flavivirus-naive
adults: A randomized, double-blind clinical trial. J Infect Dis 2013; 207(6):957–965.

22. Kirkpatrick BD, Whitehead SS, Pierce KK, et al. The live attenuated dengue vaccine
TV003 elicits complete protection against dengue in a human challenge model. Sci
Transl Med 2016; 8(330):330ra36.

23. Lindow JC, Durbin AP, Whitehead SS, Pierce KK, Carmolli MP, Kirkpatrick BD.
Vaccination of volunteers with low-dose, live-attenuated, dengue viruses leads to

serotype-specific immunologic and virologic profiles. Vaccine 2013; 31(33):3347–
3352.

24.   Durbin AP, McArthur J, Marron JA, et al. The live attenuated dengue serotype 1 vaccine
rDEN1Delta30 is safe and highly immunogenic in healthy adult volunteers. Hum
Vaccin 2006; 2(4):167–173.

25.   Durbin AP, Karron RA, Sun W, et al. Attenuation and immunogenicity in humans of a
live dengue virus type-4 vaccine candidate with a 30 nucleotide deletion in its 3'-
untranslated region. Am J Trop Med Hyg 2001; 65(5):405–413.

26.   Puschnik A, Lau L, Cromwell EA, Balmaseda A, Zompi S, Harris E. Correlation between
dengue-specific neutralizing antibodies and serum avidity in primary and secondary
dengue virus 3 natural infections in humans. PLOS Negl Trop Dis 2013; 7(6):e2274.

27.   Balmaseda A, Standish K, Mercado JC, et al. Trends in patterns of dengue transmission
over 4 years in a pediatric cohort study in Nicaragua. J Infect Dis 2010; 201(1):5–14.

28.   Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. Nat Protoc 2009; 4(1):44–57.

29.   Waddell SJ, Popper SJ, Rubins KH, et al. Dissecting interferon-induced transcriptional
programs in human peripheral blood cells. PloS One 2010; 5(3):e9753.

30.   Kupershmidt I, Su QJ, Grewal A, et al. Ontology-based meta-analysis of global
collections of high-throughput public data. PloS One 2010; 5(9).

31.   Sun P, García J, Comach G, et al. Sequential waves of gene expression in patients with
clinically defined dengue illnesses reveal subtle disease phases and predict disease
severity. PLoS Negl Trop Dis 2013; 7(7):e2298.

32.   Pandey N, Jain A, Garg RK, Kumar R, Agrawal OP, Lakshmana Rao PV. Serum levels of
IL-8, IFNγ, IL-10, and TGF β and their gene expression levels in severe and non-severe
cases of dengue virus infection. Arch Virol 2015; 160(6):1463–1475.

33.   Assoian RK, Komoriya A, Meyers CA, Miller DM, Sporn MB. Transforming growth
factor-beta in human platelets. Identification of a major storage site, purification, and
characterization. J Biol Chem 1983; 258(11):7155–7160.

34.   Wong KL, Tai JJ-Y, Wong W-C, et al. Gene expression profiling reveals the defining
features of the classical, intermediate, and nonclassical human monocyte subsets.
Blood 2011; 118(5):e16-31.

35.   Strouts FR, Popper SJ, Partidos CD, Stinchcomb DT, Osorio JE, Relman DA. Early
transcriptional signatures of the immune response to a live attenuated tetravalent
dengue vaccine candidate in non-human primates. PLoS Negl Trop Dis 2016;
10(5):e0004731.

36. Simon-Lorière E, Duong V, Tawfik A, et al. Increased adaptive immune responses and
proper feedback regulation protect against clinical dengue. Sci Transl Med 2017;
9(405):eaal5088.

37. Whitney AR, Diehn M, Popper SJ, et al. Individuality and variation in gene expression
patterns in human blood. Proc Natl Acad Sci (USA) 2003; 100(4):1896–1901.

38. Rani MRS, Xu Y, Lee J, et al. Heterogeneous, longitudinally stable molecular signatures
in response to interferon-β. Ann N Y Acad Sci 2009; 1182(1):58–68.

39. Deal EM, Lahl K, Narváez CF, Butcher EC, Greenberg HB. Plasmacytoid dendritic cells
promote rotavirus-induced human and murine B cell responses. J Clin Invest 2013;
123(6):2464–2474.

40. Hottz ED, Medeiros-de-Moraes IM, Vieira-de-Abreu A, et al. Platelet activation and
apoptosis modulate monocyte inflammatory responses in dengue. J Immunol 2014;
193(4):1864–1872.

41. Sabchareon A, Wallace D, Sirivichayakul C, et al. Protective efficacy of the recombinant,
live-attenuated, CYD tetravalent dengue vaccine in Thai schoolchildren: a randomised,
controlled phase 2b trial. Lancet 2012; 380(9853):1559–1567.

42. Moodie Z, Juraska M, Huang Y, et al. Neutralizing antibody correlates analysis of
tetravalent dengue vaccine efficacy trials in Asia and Latin America. J Infect Dis 2018;
217(5):742–753.

43. Waggoner JJ, Balmaseda A, Gresh L, et al. Homotypic dengue virus reinfections in
Nicaraguan children. J Infect Dis 2016; .

44. Forshey BM, Reiner RC, Olkowski S, et al. Incomplete protection against dengue virus
Type 2 re-infection in Peru. PLOS Negl Trop Dis 2016; 10(2):e0004398.

45. Henein S, Swanstrom J, Byers AM, et al. Dissecting antibodies induced by a chimeric
Yellow Fever-Dengue, live-attenuated, tetravalent Dengue vaccine (CYD-TDV) in naive
and dengue-exposed individuals. J Infect Dis 2017; 215(3):351–358.

46. Weiskopf D, Bangs DJ, Sidney J, et al. Dengue virus infection elicits highly polarized
CX3CR1+ cytotoxic CD4+ T cells associated with protective immunity. Proc Natl Acad
Sci (USA) 2015; 112(31):E4256–E4263.

47. Weiskopf D, Angelo MA, Azeredo EL de, et al. Comprehensive analysis of dengue virus-
specific responses supports an HLA-linked protective role for CD8+ T cells. Proc Natl
Acad Sci (USA) 2013; 110(22):E2046–E2053.

48. Hatch S, Endy TP, Thomas S, et al. Intracellular cytokine production by dengue virus-
specific T cells correlates with subclinical secondary infection. J Infect Dis 2011;
203(9):1282–1291.

49. Duangchinda T, Dejnirattisai W, Vasanawathana S, et al. Immunodominant T-cell
responses to dengue virus NS3 are associated with DHF. Proc Natl Acad Sci (USA)
2010; 107(39):16922–16927.

50. Angelo MA, Grifoni A, O'Rourke PH, et al. Human CD4[+] T cell responses to an
attenuated tetravalent dengue vaccine parallel those induced by natural infection in
magnitude, HLA restriction, and antigen specificity. J Virol 2017; 91(5):e02147-16.

Figure Legends

Figure 1. Significant differences in transcript abundance post-vaccination (FDR<1%; minimum

2-fold change compared to pre-vaccination sample).

Figure 2. Changes in transcript abundances over time in vaccinees. A) Hierarchical clustering

of the 286 transcripts whose abundance was significantly different from baseline on more

than one day. Lines and numbers to the right of the heatmap mark sets of co-expressed

genes (average cluster r>0.5). B) Change over time in abundance for each transcript in

each gene cluster. Heavy line indicates median expression of all genes in each cluster. C)

Gene ontologies associated with gene clusters described in (A) and (B). There were no

significant gene ontologies for Cluster 3.

Figure 3. Comparison of post-vaccination and post-infection transcript abundance changes. A)

Transcripts with significant changes on days 2, 3, 4, or 5 of fever in patients with primary

DENV-3 infection (blue circle) and on any day post-vaccination (green circle). Numbers

indicate transcripts unique to vaccination, infection, or shared (overlap, n=246). B)

Maximum fold-change in transcript abundance following vaccination (red circles) or during

infection (blue diamonds). C) Maximum fold-change in abundance for transcripts with

significant changes post-vaccination or during infection. Dotted diagonal line at equal fold

change included for reference.

Figure 4. Gene modules affected by DENV vaccination and natural infection. A) Blood

transcript modules with transcripts that were significantly up- or down-regulated on at least

one day (FDR <1%) were hierarchically clustered. NES; normalized enrichment score.

Vertical lines on right denote module clusters described in the text. B) Hierarchical

clustering of each day post-vaccination or post-infection using the NES from (A). Days in

bold italics represent days of fever for infected patients; days preceded by "v" represent

days post-vaccination.

Figure 5. Correlation of transcript abundance and day 42 $PRNT_{60}$ among vaccine recipients. A)

Average fold change in abundance by day for all transcripts with significant differences

from baseline post-vaccination. Transcripts are ordered and clusters labeled as in Figure

2. Asterisk marks IFI44. B) Spearman correlation of each transcript and day 42 $PRNT_{60}$

using a moving average of window size 9. Solid lines indicate days post-vaccination on

which a significant correlation was identified ($p < 0.01$, indicated by vertical dotted grey line).

Figure 6. Gene modules correlated with subsequent neutralizing antibody response. A) Blood

Transcript Modules that were significantly enriched for transcripts positively correlated with

day 42 $PRNT_{60}$ (vaccinees) or convalescent NT_{50} (patients) on at least one day (FDR<1%)

were hierarchically clustered. NES; normalized enrichment score. Vertical lines delineate

module clusters described in the text. B) Significant modules (FDR<1%) are marked in red.

Modules and samples are organized as in (A). C) Hierarchical clustering of gene module

expression from each day post-vaccination or post-infection using the NES from (A). Day

labels in bold italics represent fever day for infected patients; day labels preceded by "v"

represent day post-vaccination.

Table 1. Characteristics of subjects in vaccine trial

Subject	Age	Sex	Viremia[a]	Rash[b]	Day 28 PRNT$_{60}$[c]	Day 42 PRNT$_{60}$[c]	Day 180 PRNT$_{60}$[c]
1 (Vaccine)	19	F	--	Days 12-20	54	70	30
2 (Vaccine)	26	F	--	--	22	15	<5
3 (Vaccine)	25	M	Days 8-9	--	52	106	22
4 (Vaccine)	20	M	Days 8-9	Days 12-16	26	32	<5
5 (Vaccine)	20	M	Day 6	Days 12-20	33	19	<5
6 (Vaccine)	22	M	--	--	<5	<5	<5
7 (Vaccine)	19	M	--	--	18	8	8
8 (Vaccine)	22	F	Days 5-8	Days 12-20	34	29	8
9 (Vaccine)	19	F	--	--	25	33	<5
10 (Vaccine)	46	F	--	Days 12-16	70	152	64
11 (Placebo)	18	F	--	--	<5	<5	<5
12 (Placebo)	19	M	--	--	<5	<5	<5
13 (Placebo)	45	M	--	--	<5	<5	<5
14 (Placebo)	21	F	--	--	<5	<5	<5

553   [a] Virus detected in serum from tissue culture plaque formation assay

554   [b] First and last day on which maculopapular rash observed

555   [c] Reciprocal serum dilution providing 60% reduction in plaque formation

A.

Day: 2 5 6 8 9 12 14 20 29 42 180

B.

C.

-log10(p-value)

18 16 14 12 10 8 6 4 2 0

response to virus
immune response
defense response
cytosolic DNA-sensing pathway
Cluster 1   RIG-I-like receptor signaling pathway
NAD-ADP-ribosyltransferase activity
JAK-STAT signaling cascade

chromosome segregation
mitotic cell cycle
condensed chromosome
Cluster 2   spindle
condensed nuclear chromosome
DNA replication

A.

B.

Infection Days 2-5	Vaccination Days 2-180

Cell cycle

Respiration
Protein folding

Viral sensing
IFN response

Monocytes
Ag presentation

Cytoskeletal remodeling
Platelet Activation

B cell
T cell activation/
differentiation

2.5

0   NES

-2.5

D3
D1,D2
vD9
vD8

D4
vD12
vD5

D5
vD14
vD42
vD29
vD20
vD6

vD180
vD2

A.

Day: 2 ⟶ 180

*

B. Spearman correlation coefficient

-1   -0.5   0.5   1

········ Day 2

········ Day 5

········ Day 6

—— Day 8

—— Day 9

—— Day 12

—— Day 14

A. Infection Days 3-5 | Vaccination Days 2-14

Monocyte

Anti-viral
Interferon

Cell cycle
Mitosis

T cell

NK cell

NES
2.5
0
-2.5

B.

C.

vD2
D4
D5
vD14
D3
vD8
vD9
vD6
vD5
vD12