1
2
3
4
5 **Real time portable genome sequencing for global food security**

15 Laura M. Boykin[1,*], Ammar Ghalab[1], Bruno Rossitto De Marchi[1,2], Anders Savill[1], James M. Wainaina[1],
16 Tonny Kinene[1], Stephen Lamb[1], Myriam Rodrigues[1], Monica A. Kehoe[3], Joseph Ndunguru[4], Fred
17 Tairo[4], Peter Sseruwagi[4], Charles Kayuki[4], Deogratius Mark[4], Joel Erasto[4], Hilda Bachwenkizi[4], Titus
18 Alicai[5], Geoffrey Okao-Okuja[5], Phillip Abidrabo[5], John Francis Osingada[5], Jimmy Akono[5], Elijah Ateka[6],
19 Brenda Muga[6], and Samuel Kiarie[6]
20
21
22

[1]School of Molecular Sciences and Australian Research Council Centre of Excellence in Plant Energy

Biology, University of Western Australia, Crawley, Perth, WA 6009 Australia.

[2]São Paulo State University (UNESP), School of Agriculture, Dept. of Plant Protection, CEP

18610-307, Botucatu (SP), Brazil

[3]Crop Protection Branch, Department of Agriculture and Food Western Australia, South Perth, WA

6151, Australia

[4]Mikocheni Agricultural Research Institute (MARI), P.O. Box 6226, Dar es Salaam, Tanzania

[5]National Crops Resources Research Institute (NaCRRI), P.O. Box 7084, Kampala, Uganda

[6]Jomo Kenyatta University of Agriculture and Technology (JKUAT), P.O. Box 62000 - 00200,

Nairobi, Kenya

*Author for correspondence: Laura Boykin, laura.boykin@uwa.edu.au

Abstract

The United Nations has listed Zero Hunger as one of the 17 global sustainable development goals to

end extreme poverty by 2030. Plant viruses are a major constraint to crop production globally causing

an estimated $30 billion in damage [1] leaving millions of people food insecure [2]. In Africa, agriculture

employs up to 50% of the workforce, yet only contributes 15% to the GDP on average [3], suggesting that

there is low productivity and limited value addition. This can be addressed through continued

innovation in the fields of science and technology as suggested in the Science Agenda for Agriculture in

Africa (S3A) [4]. Sustainable management of plant viruses and their associated vectors must include

efficient diagnostics for surveillance, detection and identification to inform disease management,

including the development and strategic deployment of virus resistant varieties. To date, researchers

have been utilizing conventional methods such as include; PCR, qPCR, high throughput sequencing

(RNA-Seq, DNA-Seq) and Sanger sequencing for pathogen identification. However, these methods are

both costly and time consuming, delaying timely control actions. The emergence of new tools for real-

time diagnostics, such as the Oxford Nanopore MinION, have recently proven useful for early detection

of Ebola [6] and Zika [7,8], even in low resourced laboratories. For the first time globally, the MinION

portable pocket DNA sequencer was used to sequence whole plant virus genomes. We used this

technology to identify the begomoviruses causing the devastating CMD which is ravaging smallholder

farmers' crops in sub-Saharan Africa. Cassava, a carbohydrate crop from which tapioca originates, is a

major source of calories for over eight hundred (800) million people worldwide. With this technology,

farmers struggling with diseased crops can take immediate, restorative action to improve their

livelihoods based on information about the health of their plants, generated using a portable, real-time

DNA sequencing device.

The portable DNA technology has great potential to reduce the risk of community crop failure and help improve livelihoods of millions of people, especially in low resourced communities. Plant diseases are a major cause of low crop productivity and viruses such as Tobacco mosaic virus, Tomato mosaic, Tomato spotted wilt, Potato leaf roll, Potato virus X and Y in Potato, Papaya mosaic, Citrus tristeza, Chilli leaf curl, and Banana bunchy top have been implicated. In particular, cassava viruses are among the world's greatest risk to food insecurity. Losses caused by cassava mosaic disease (CMD) and cassava brown streak disease are estimated at US $2-3 billion annually [5]. We therefore visited smallholder farmers in Tanzania, Uganda and Kenya (Table 1) who are suffering yield shortages due to cassava virus infections. We utilized the MinION to test infected material and farmers were informed within 48 hours of the specific strain of the virus that was infecting their cassava, and a resistant cassava variety was deployed. The advantages of adopting this technology far outweigh the challenges (see Table 2). Cassava mosaic begomoviruses were in high enough concentration that reads of whole genomes were obtained without an enrichment step (Table 1). As expected the viral reads increased with the severity of the symptoms observed (Table 1). We detected a dual infection for a leaf sample with the severity score of 5 in Uganda. In addition, one asymptomatic plant in Tanzania had one viral read detected. The shortest time to obtain a viral read was 15 seconds (severity score 5) and the longest was 4h11m15s (severity score 1).

Additionally, MinION sequencing is superior to traditional methods of PCR identification given its generation of whole genome sequences which enable the identification of the plant virus strain even if it becomes mutated or divergent, as it is not biased using primers that rely on known virus

sequences. With regards to cassava, there are three major advantages of this technology. Firstly,

improved diagnostics are required and real-time whole genome sequencing will help develop

diagnostic primers that are up-to-date. Secondly, this technology will assist with the development of

resistant cassava varieties and will allow breeders to immediately test the varieties they are developing

against different viral strains. Lastly, it ensures the delivery of the correct healthy uninfected planting

material to farmers. In addition, we could detect virus in a plant before it showed symptoms (Table 1).

Utilizing traditional PCR methods three samples collected from Asha's field in Tanzania tested positive

for EACMVs and none were positive for ACMV. The asymptomatic sample from MARI tested negative

for both ACMV and EACMVs. Eight fresh cassava leaf samples from Uganda were dually infected with

ACMV and EACMV-UG using conventional PCR primers for ACMV and EACMV-UG. The primers used in

this PCR yield products of 1000bp and 1500bp for ACMV and EACMV-UG respectively. Twelve Kenyan

samples were tested and all but two (barcode 2 and 3) were positive using conventional PCR (Table 1).

Further studies are needed to verify our results regarding the sensitivity of the protocol for early

detection of CMD in cassava, but these results are very promising for ensuring farmers receive clean

planting material with early detection of viral infection.

Nanopore sequencing technology has wide applications globally, but in East Africa these

include: (a) crop improvement by screening for virus resistant germplasm and genetic diversity during

breeding, (b) indexing of cassava planting materials for virus presence or absence to ensure that only

clean materials in multiplication fields are distributed to farmers, (c) detection and identification of

alternative plant species for cassava-infecting begomoviruses, so that farmers are advised to remove

and/or grow crops away from such plants as a management strategy, (d) virus and biodiversity studies.

Methods

Sample collection and DNA extraction: In Tanzania, three cassava mosaic disease (CMD) symptomatic

cassava leaf samples (Fig. 1, Table 1) were collected from the smallholder cassava farmer Asha

Muhammed's field in Bagamoyo. One more asymptomatic leaf sample was collected at Mikocheni

Agricultural Research Institute (MARI), Dar es Salaam. Seven CMD symptomatic plants were collected

from Naomi Kutesakwe's farm in Wakiso district in Uganda. Both Tanzanian and Ugandan samples

were collected in September 2017. Twelve samples from Kenya were collected in February 2018 from

various sources (Table 1). High quality DNA were isolated using CTAB method [9]. Each DNA sample

(Table 1) was quantified and purity checked using a NanoDrop 2000c UV–vis Spectrophotometer

(Thermo Scientific, Wilmington, DE, USA) was used to check the purity and quantity of DNA for each

sample and results were recorded in Table 1.

Nanopore library preparation and sequencing: In Tanzania and Uganda, the Rapid Barcoding kit SQK-

RBK001 and 9.4.1 flow cells were used to process genomic DNA extracted using a standard CTAB

method. We utilized the Rapid Barcoding kit SQK-RBK004 with 9.4.1 flow cells in Kenya. DNA was

diluted to 700 ng as specified in the library protocol. The SQK-RBK001 (Sept 2017) and/or the SQK-

RBK004 (Feb 2018) protocols were performed as described by the manufacturer. In Tanzania and

Uganda, the MinION was run for 24 hours instead of the recommended 48 hours, and in Kenya we had

a total run time of approximately 17 hours due to power interruptions.

Nanopore bioinformatics: In Tanzania and Uganda, Albacore 2.0.2 was used for base calling. In Kenya,

Albacore 2.1.10 was used and the scripts were modified to reflect the newest rapid barcoding kit

RBK004. Fastq files were imported into Geneious [10] and a local blast database of all known cassava

mosaic begomovirus whole genomes were downloaded from GenBank and a local blast was performed

on each of the reads generated using the nanopore device.

Verification of nanopore results: Traditional PCR was used to verify our nanopore results. In Tanzania

and Kenya, two primer pairs: EAB 555F/EAB 555F [11] and JSP001/JSP002 [12], which amplify 556bp and

774bp were used to detect East African cassava mosaic begomoviruses (EACMVs) and African cassava

mosaic begomoviruses (ACMVs), respectively. In Uganda, the presence of ACMV and EACMV in each

sample was detected using a pair of specific primers for ACMV, ACMV-AL1/F and ACMV-ARO/R and

specific for EACMV-UG2, UV-AL1/F and ACMC-CP/R3 [13].

Acknowledgements: Funding for the Kenyan trip was provided by the Crawford Fund. We also thank

the participants from the University of Eldoret who assisted in the preparation of libraries for the

Kenyan samples.

Author contributions: Designed the study: LMB, JN, TA, FT, PS, MK, AS, EA. Carried out experiments:

LMB, AG, BD, JW, MR, JN, PS, CK, DM, JE, HB, TA, GO, PA, JO, JA, EA, BM and SK. Analysed data: LMB,

AS, CK, DM, JE, HB, SL, JN, PS, TA, GO, PA, JO, JA, EA, BM. Contributed to the writing: All authors

contributed to the writing of the manuscript.

Competing interests: The authors have no competing interests.

References:

Subramanya Sastry, K. & Zitter, T. A. in *Plant Virus and Viroid Diseases in the Tropics Vol. 2 Epidemiology and Management* 149–480 (2014).

Legg, J. P. & Thresh, J. M. Cassava mosaic virus disease in East Africa: a dynamic disease in a changing environment. *Virus Res* **71**, 135-149 (2000).

AFDB. African Development Bank. Annual Report., (2016).

FARA. Science agenda for agriculture in Africa (S3A): "Connecting Science" to transform agriculture in Africa. . *Forum for Agricultural Research in Africa (FARA), Accra, Ghana* (2014).

Scholthof, K. B. *et al.* Top 10 plant viruses in molecular plant pathology. *Mol Plant Pathol* **12**, 938-954, doi:10.1111/j.1364-3703.2011.00752.x (2011).

Quick, J. *et al.* Real-time, portable genome sequencing for Ebola surveillance. *Nature* **530**, 228-232, doi:10.1038/nature16996 (2016).

Faria, N. R. *et al.* Mobile real-time surveillance of Zika virus in Brazil. *Genome Med* **8**, 97, doi:10.1186/s13073-016-0356-2 (2016).

Quick, J. *et al.* Multiplex PCR method for MinION and Illumina sequencing of Zika and other virus genomes directly from clinical samples. *Nat Protoc* **12**, 1261-1276, doi:10.1038/nprot.2017.066 (2017).

Lodhi, M. A., Guang-Ning, Y., Weeden, N. F. & Reisch, B. I. A simple and efficient method for DNA extraction from grapevine cultivars, Vitisspecies and Ampelopsis. *Plant Mol Biol Rep* **12**, 6-13 (1994).

Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647-1649, doi:10.1093/bioinformatics/bts199 (2012).

Fondong, V. N. *et al.* Evidence of synergism between African cassava mosaic virus and a new double-recombinant geminivirus infecting cassava in Cameroon. *J Gen Virol* **81**, 287-297, doi:10.1099/0022-1317-81-1-287 (2000).

Pita, J. S., Fondong, V. N., A, S., RNN, K. & Fauquet, C. Genomic and biological diversity of the African cassava geminiviruses. *Euphytica* **120**, 115-125 (2001).

Zhou, X. *et al.* Evidence that DNA-A of a geminivirus associated with severe cassava mosaic disease in Uganda has arisen by interspecific recombination. *J Gen Virol* **78 (Pt 8)**, 2101-2111, doi:10.1099/0022-1317-78-8-2101 (1997).

Table 1. Nanopore results of cassava mosaic begomovirus-infected cassava leaves from Tanzania, Uganda and Kenya.

Farmer	Disease Severity	Lab ID	Barcode	Location	Concentration ng/uL	Run Time	Time-to-result	# Nanopore reads	# Viral Reads	PCR Result	Length (nucleotide)	Closest Species Match
Tanzania												
Asha	3	3	1	Kiromo-Kitonga Bagamoyo	3,378.40	24 hrs	47s	63,653	950	+	969	EACMV-KE
Asha	2	5	2	Kiromo-Kitonga Bagamoyo	3,385.20	24 hrs	32m57s	8,844	43	+	971	EACMV-KE
Asha	4	6	3	Kiromo-Kitonga Bagamoyo	4,845.20	24 hrs	28s	32,540	338	+	775	EACMV-KE
Asha	Healthy	8	4	MARI	1,752.70	24 hrs	4h11m15s	20,005	1	-	2213	EACMV-KE
Uganda												
Naomi	4	1	1	Wakiso	603.2	24 hrs	15s	9,859	244	+	957	EACMV-UG
Naomi	3	2	2	Wakiso	711.3	24 hrs	43s	8,482	98	+	780	EACMV-UG
Naomi	5	3	3	Wakiso	603.6	24 hrs	17m14s	5,562	298	+	801	EACMV-UG AND ACMV-UG
Naomi	2	4	4	Wakiso	598.2	24 hrs	11s	17,146	371	+	789	ACMV-UG
Naomi	3	5	5	Wakiso	610	24 hrs	33s	7,999	250	+	2264	ACMV-UG
Naomi	1	6A	6	Wakiso	448	24 hrs	2h14m50s	9,972	18	+	920	ACMV-KE
Naomi	3	6B	7	Wakiso	490.9	24 hrs	2m52s	9,662	119	+	1418	ACMV-KE
Naomi	Healthy	7	8	NaCRRI	710.6	24 hrs	-	7,881	none	-	-	none
Kenya												
Rose	2	13	12	Eldoret	44.3	17hr 17min	3h4m9s	30	22	+	754	ACMV-KE
Lungo Mutunga	1	1	1	Ngoliba	2433.4	17hr 17min	20m17s	3,298	1,760	+	1145	ACMV-NG AND EACMV-UG
Lungo Mutunga	1	2	2	Ngoliba	2624.2	17hr 17min	37m23s	3,232	1,838	-	1778	ACMV-UG AND EACMV-UG
Joel Maina	1	3	3	Ngoliba	1812.5	17hr 17min	10m3s	3,787	2,108	-	2723	EACMV-UG
Rainbow Hotel	1	5	4	Ruiru	3066.3	17hr 17min	57m9s	2,536	1,423	+	905	EACMV-UG
JKUAT	3	NF19C2	5	Siaya	767.59	17hr 17min	50s	2,672	1,493	+	1985	ACMV-KE
JKUAT	4	CF13B2	6	Kilifi	1143.7	17hr 17min	28s	3,870	2,221	+	847	EACMV-KE AND ACMV-KE
JKUAT	3	EF23C2	7	Kitui	1213.8	17hr 17min	18m6s	2,964	1,629	+	766	EACMV-KE
JKUAT	3	EF20C1	8	Kitui	1023	17hr 17min	54m2s	5,120	2,906	+	1887	EACMV-UG
JKUAT	3	Wf2S1	9	Vihiga	1111.9	17hr 17min	23m7s	5,095	3,021	+	2233	ACMV-NG AND EACMV-UG
JKUAT	3	WF20C2	10	Busia	929.36	17hr 17min	36m1s	4,652	2,838	+	1966	ACMV-UG AND EACMV-UG
JKUAT	3	CF36C2	11	Taita taveta	1032.3	17hr 17min	55m	4,366	2,541	+	2089	EACMV-KE

Table 2. Advantages and challenges of using the MinION portable DNA sequencer in Tanzania, Uganda and Kenya.

Advantages	Challenges
1. Rapid – obtain results timely to support quick decision making on disease management	1. Power
2. Expenses on sending samples for sequencing abroad removed	2. Internet
3. High resolution of results and reliability	3. Cost
4. Detection of mixed infections	4. Computer access
5. Detection of virus in latently infected plants	5. Shipping
6. Discovery of unknown viruses	
7. Minimising sample degradation and loss during shipping	
8. Virus-indexing for safe movement of germplasm	
9. Better plant health regulation, inspection/phytosanitary – improved international trade	

Figure 1. Cassava plant infected with cassava mosaic begomoviruses. Photo credit: Dr. Ndunguru.