Genomic determinants of speciation and spread of the *Mycobacterium tuberculosis* complex

Álvaro Chiner-Oms[1,#], Leonor Sánchez-Busó[2,#], Jukka Corander[2,3,4], Sebastien Gagneux[5,6], Simon Harris[2], Douglas Young[7], Fernando González-Candelas[1,8] Iñaki Comas[8,9*]

[1]Unidad Mixta "Infección y Salud Pública" FISABIO-CSISP/Universidad de Valencia,

Instituto de Biología Integrativa de Sistemas-I2SysBio, Valencia, Spain.

[2]Pathogen Genomics, Wellcome Trust Sanger Institute, Cambridge CB10 1SA, UK.

[3]Department of Biostatistics, University of Oslo, 0317 Oslo, Norway.

[4]Helsinki Institute of Information Technology (HIIT), Department of Mathematics and

Statistics, University of Helsinki, 00014 Helsinki, Finland.

[5]Swiss Tropical and Public Health Institute, Basel, Switzerland.

[6]University of Basel, Basel, Switzerland.

[7]The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK.

[8]CIBER en Epidemiología y Salud Pública, Valencia, Spain

[9]Instituto de Biomedicina de Valencia, IBV-CSIC, Valencia, Spain.

[#] Contributed equally.
*Correspondence:icomas@ibv.csic.es

ABSTRACT

BACKGROUND

Models on how bacterial lineages differentiate increase our understanding on early bacterial speciation events and about the genetic loci involved. Here, we analyze the population genomics events leading to the emergence of the tuberculosis pathogen.

RESULTS

The emergence is characterized by a combination of recombination events involving core pathogenesis functions and purifying selection on early diverging loci. We identify the *phoR* gene, the sensor kinase of a two-component system involved in virulence, as a key functional player subject to pervasive positive selection after the divergence of the MTBC from its ancestor. Previous evidence showed that *phoR* mutations played a central role in the adaptation of the pathogen to different host species. Now we show that *phoR* have been under selection during the early spread of human tuberculosis, during later expansions and in on-going transmission events.

CONCLUSIONS

Our results show that linking pathogen evolution across evolutionary and epidemiological timescales point to past and present virulence determinants.

KEYWORDS

Bacterial evolution, pathogen genomics, recombination, transmission, two component system

BACKGROUND

The increasing availability of population genomics data has allowed an improved understanding of genotypic and ecological differentiation among closely related bacteria[1]. While a species concept *sensu stricto* cannot be applied to bacteria[2] models exists to understand how species can emerge in natural populations. Depending on the evolutionary forces involved, models range from differentiation driven by natural selection and adaptation to different ecological niches (Ecological Species Concept) to differentiation as a result of restricted gene flow that reinforces isolation (Biological Species Concept). In reality, most natural populations show a combination of both processes with certain overlap between habitats (Overlapping habitats model[3]). The study of natural populations and models shows that the emergence of new species is more common among bacterial groups sharing, partially or totally, their habitat, a process also known as sympatric speciation[4]. Processes of bacterial differentiation are often expected to leave measurable genetic signatures in extant genomes including "speciation islands" (regions of high divergence between the nascent species)[3,4]. These genetic signatures hold clues about the key genomic determinants responsible for ecological differentiation of a nascent species from a common genetic pool. However, how these models apply to professional pathogens, particularly those characterized by an obligate association with their host species, has been little explored.

Species of the *Mycobacterium tuberculosis* complex (MTBC) cause devastating morbidity and mortality in humans and animals, which also lead to important economic losses[5]. The MTBC comprises a group of bacteria with genome sequences having

an average nucleotide identity greater than 99% and sharing a single clonal ancestor.
This includes the predominantly human pathogens referred to as *Mycobacterium*
tuberculosis and *Mycobacterium africanum* as well as a series of pathogens isolated
from other mammalian species known as *M. bovis*, *M. pinnipedi*, *M. antelope*, *M.*
microti, etc. Human-adapted tuberculosis bacilli show a strong geographic association
with some lineages being globally distributed (e.g. lineage 4) and others being
geographically restricted (e.g., lineage 5, 6, 7)[6,7]. It is assumed that the causes of
this variable geographical distribution are both historical (e.g. trade, conquest,
globalization) and biological (e.g. interactions with different human genetic
backgrounds)[7]. There is limited transmissibility of animal-adapted strains in humans
and, conversely, human-adapted strains transmit poorly among animals[8]. Despite
the wide range of host species infected by the different members of the MTBC, there
is a maximum of ~2,500 single nucleotide polymorphisms (SNPs) separating any two
MTBC genomes[9]. The most closely-related bacteria that fall outside of the MTBC
include isolates known as *Mycobacterium canettii* (MCAN). MCAN strains differ from
MTBC isolates by tens of thousands of SNP, most of them contributed by
recombination between strains[10,11]. MCAN strains have been isolated from the
Horn of Africa, predominantly from children and often in association with
extrapulmonary tuberculosis[12]. Thus, it is assumed that MCAN represents an
opportunistic pathogen with an unidentified environmental reservoir[13] opposed to
the obligate MTBC pathogen. Genomic comparisons have identified gene-content
differences between MTBC, MCAN and other mycobacteria[10,14,15] as well as
genetic differences in virulence-related loci[16,17].
104
Two pieces of evidence suggest that MTBC and MCAN evolved from a common
genetic pool in Africa. Strains of the MTBC have an average nucleotide identity (ANI)
to MCAN strains of 98% (range 97.71%-99.30%)[10](our own data), suggesting
incomplete or recent speciation (an operational species concept identifies 95% as the
barrier to delineate species[18]). In addition, most reports suggest lack of on-going
recombination between MCAN and MTBC and within the MTBC[19,20] suggesting
complete separation (but see [21]). The second piece of evidence comes from
phylogeographic and genetic diversity analyses which identified the origin of the
tuberculosis bacilli in Africa[6,22], the likely place of origin of MCAN[10,23]. Taken
together, the data suggests that ancestral MTBC and MCAN strains at least shared
partially the same niche and genetic pool.
116
Our understanding about the population genomics events mediating the divergence of
the ancestor of the MTBC from a common ancestral pool with MCAN is far from
complete. The availability of genome sequences from thousands of MTBC clinical
strains as well as of close relatives like MCAN enables us to revisit previous evidence
about the evolution of MTBC and also to identify molecular signatures of MTBC
speciation events as well as new targets for biomedical research.
123
124

RESULTS

We first analyzed the differentiation between MTBC and MCAN by searching for any
hallmark of on-going recombination between and within these groups of strains.
Previous reports have suggested that there might be limited but significant
recombination among MTBC strains[21,24] while others failed to identify measurable
recombination events[25]. To revisit this question, we reasoned that including larger

available datasets may maximize the signal of recombination signal if exists. We
screened a dataset of complete genome sequences of strains from global
sources[26](n=1,591). These genomes are representative of the known geographic
and genetic diversity of the MTBC (Additional file 1: Figure S1). Among those
genomes, we identified all the variant positions and, more specifically, potential
homoplastic sites, i.e. polymorphic sites showing signs of convergent evolution. A total
of 96,143 variant positions were called in the 1,591 strains. Parsimony and likelihood
phylogenetic mapping of those positions revealed 3,439 positions as being
homoplastic (3,6%). Homoplasy can arise as a consequence of recombination but it
may be caused by other processes, such as positive selection, sequence gaps
contributing to homoplastic counts or mapping/calling errors. To increase the likelihood
for homoplastic positions to be due to recombination events, we filtered out non-
biallelic positions (1.8%) and potential mapping errors identified by generating
synthetic reads around each SNP position (0.2%). In addition, known drug-resistance
positions were filtered out due as these are well known instances of convergent
evolution[27–29](see Methods).

As a result, a total of 2,360 core homoplastic sites were identified across the 1,591
strains analyzed (2.5% of all variable sites). Homoplastic sites were not significantly
accumulating in any region of the genome, suggesting the absence of recombination
hotspots (Figure 1A). In addition, with these variant positions, we looked at
consecutive runs of homoplastic sites in the genomes. The 2,360 homoplastic
positions defined 97 homoplastic runs (see Methods for definition of a homoplastic
run, Additional file 1: Figure S2). If the accumulation of homoplasies was due to
recombination, we expect the variant positions involved in each consecutive run to
share the same phylogenetic history. From the 97 homoplastic runs, we detected only
2 cases in which two variant positions shared phylogenetic congruence. The two
regions accounted for 4 convergent variants (Additional file 2: Table S1) and affected
strains from different MTBC lineages. Variants in positions 2,195,896 and 2,195,899
fell in the primary regulatory region of *mazE5*[30]. Variants in positions 2,641,161 and
2,641,163 fell in the intergenic region of *glyS* and Rv2358. Although we cannot
completely discard the possibility that these represent recombination events, it is more
likely that the two regions have been under positive selection, a mechanism known to
lead to the accumulation of homoplastic accumulation in the MTBC[29]. In summary,
this large-scale variant-by-variant analysis was not able to identify significant on-going
recombination between any of the 1,591 MTBC strains analyzed.

Figure 1. **No ongoing recombination within the MTBC** A) Number of homoplasies (grey) as a function of the total number of variants detected (orange) in the MTBC dataset (n=1,591) B) Linkage disequilibrium as a function of genetic distance detected in a representative sample of *Mycobacterium tuberculosis* complex strains (n = 1,591). C) Site frequency spectrum of MTBC strains using the core variant positions.

167
Due to the low diversity within the MTBC, we also followed alternative approaches to
try to identify recombination events with a high statistical confidence. Using an
additional method, we evaluated linkage disequilibrium (LD) as a function of the
distance between the 94,780 core variant positions. R^2 has been used in a previous
publication to show on-going recombination at very short distances (less than 50
bp.)[21]. In our much larger dataset, R^2 values were also slightly higher at shorter
distances, which is compatible with recombination involving larger fragment sizes.

However, a close examination reveals that the peak at short distances is misleading,
as it is driven by only six points out of more than 11,000 comparisons (Figure 1B). In
addition, R^2 values are known to fail to reach the theoretical maximum of 1 when
variants compared have very different frequencies[31]. This is likely the case for
MTBC, in which there is a strong skew of the site frequency spectra towards low
frequency values (Figure 1C)[32]. Thus, as an alternative we calculated D'. In this
dataset, as expected for a mostly clonal organism, LD measured by D' remained at its
maximum value, even when focusing on distant variant positions more than 5 Kb apart,
suggesting very little or no ongoing recombination (Figure 1B).
184
To further validate these findings, we ran Gubbins with the same dataset and validated
them with RDP4 (see methods for details). Gubbins detects the accumulation of a
higher than expected number of variants in addition to homoplastic sites as a hallmark
of possible recombination. We partitioned the 1,591 strain dataset into the different
lineages and screened for possible tracks of recombination. Gubbins reported
potential recombining regions characterized by a higher than average number of SNPs
however the RDP4 methods did not confirmed any of them. Thus those events maybe
real but cannot be confirmed by alternative approaches.
193
Having established that recombination has little impact on the overall MTBC genetic
diversity, we compared a representative dataset of MTBC genomes[6] (n = 219) with
7 MCAN genomes to identify and quantify eventual ongoing recombination within
MCAN and between MCAN and the MTBC. Of the 93,922 polymorphic sites identified,
22,718 were biallelic homoplasies (24.2%). The genomic distribution of variant
positions and homoplasies in the MCAN group show a landscape different to the
MTBC group (Figure 2B). A total of 22,464 (98.9%) of those homoplasies were found
only among MCAN strains, representing almost half of the variability within this group
(22,464/52,392 biallelic sites, 42.9%) which points to recombination as a main source
of variability in MCAN. This is consistent with previous reports[10]. This profile is in
sharp contrast with the flat homoplastic profile for the MTBC described above (Figure
1A).
206
207 To test for ongoing recombination between MCAN and extant MTBC, we identified
208 homoplasies involving both groups. From the 93,922 total variants, 7,934 involved
209 MCAN and MTBC strains. We found 234 biallelic homoplasies involving extant MTBC

Figure 2. **Genome-wide variant profiles vary between *M. canettii*, *M. tuberculosis* and the MTBC ancestor** A) Schematic view of the phylogenetic relationships between the MCAN groups and the MTBC. In Additional file 1: Figure S3 a maximum-likelihood phylogeny of the MCAN group including the MTBC ancestor can be found. B) Number of homoplasies (grey) as a function of the total number of variants detected (orange) in the MCAN dataset and in the branch leading to the MTBC most recent common ancestor. Black dots indicate recombination events detected in the branch leading to the most recent common ancestor of the MTBC.

and MCAN strains, thus compatible with on-going recombination but also with
independent diversification. The vast majority of homoplasies detected (97%) mapped
to the branch leading to the MTBC clade (thus fixed within the MTBC but variable
within the MCAN group). These results indicate that measurable recombination events
were common between MCAN and the ancestral branch of the MTBC, but are unlikely
during subsequent diversification of the MTBC even when the sample size is greater
than previous reports.
217
Sympatric and stepwise emergence of the MTBC ancestor
Our results show that recombination with closely related mycobacteria occurred during
the emergence of the common ancestor of the MTBC. To gain a better insight on how
it occurred we reasoned that instead of comparing MCAN strains against extant MTBC
strain we should compare against a reconstructed most common ancestor of the
MTBC (derived in Comas et al., 2010[33]). This strategy allowed us to focus on those
changes specifically happening in the ancestral branch of the MTBC (see Figure 2A
and Additional file1: Figure S3, Figure S4). As described by others, the phylogeny
suggests a specific clone of the MCAN group diverged and resulted in the MTBC

[11,23]. To do so we extracted all the variant positions that were homoplastic between
the MTBC ancestor and any of the MCAN strains (7,700 positions). The SNPs
mapping to the branch leading to the MTBC ancestor genome showed a similar
homoplasy profile to that of the MCAN strains (Figure 2B), suggesting that there were
not hard barriers to gene flow between ancestral MTBC and MCAN ancestral strains,
thus supporting a model of sympatric speciation. Notably, both MCAN and the MTBC
ancestor shared a peak around the CRISPR region, highlighting the dynamic nature
of this region possibly as a result of common phage infections.
235
A Gubbins analysis including MCAN genomes and the most likely common ancestor
of the MTBC revealed a total of 65 recombination events mapping to the branch
leading to the MTBC (Additional file 2: Table S2). Mapping of variants into the
phylogeny revealed that those regions were coincident with a high number of
homoplastic variants between MCAN and the MTBC (Additional file 1: Figure S4). To
explore whether these fragments reflected real recombination, we constructed a
phylogeny for each of them. A comparison with the topology of the non-recombinant
alignment (whole genome alignment subtracting the recombinant regions) using those
recombinant regions with enough phylogenetic signal (Additional file 1: Figure S5)
revealed significant incongruence (SH test; p-value<0.05, Additional file 1: Figure S6,
Additional file 2: Table S3, Additional file 1: Supplementary text). Thus, both Gubbins
and phylogenetic approaches indicated that these 65 regions are likely recombinant
regions.
249
The potential recombinant regions were dated using BEAST and the results showed
that their inferred ages differ from the established time since the Most Recent Common
Ancestor (tMRCA) of the MTBC (see Additional file 1: Figure S7 and Additional file 1:
Supplementary text for details). Although the distribution of tMRCA for the fragments
represents a continuum, the analysis suggests a separation between "recent"
recombination events and "ancient" events closer to the time of divergence from the
MCAN group (Figure 3A). The large HPD intervals preclude any firm conclusion but
the results suggest that some regions in the MTBC ancestral genome were restricted
to gene flow before than others (Additional file 1: Figure S7).
259

Figure 3. **Past recombination between *M. canettii* strains and the MTBC ancestor** A)
Histogram distribution of the recombination fragments relative ages. The age was relativized
from 0 to 1, being 0 the age of the non-recombinant fraction of the genome and 1 the highest
value in the confidence intervals (Figure S5). B) Gene Ontology terms overrepresented in the
coding regions contained in the recombinant fragments.

If recombination played a major role in shaping the MTBC ancestral genome with
regards to pathogenesis, we would expect some functions related to the interaction
with the host to be affected. Indeed, we observed an enrichment in experimentally
confirmed essential genes in the regions involved in recombination events, suggesting
that recombination targeted important cell functions (Chi-square test; p-value < 0.01).
An enrichment analysis of Gene Ontology terms for the genes contained in these
regions identified functions related with growth, and most specifically with the category
"growth involved in symbiotic interactions inside a host cell" as significantly
overrepresented (Binomial test; adj. p-value < 0,05) (Figure 3B). This category can be
interpreted as genes involved in a strong association between the pathogen and the
host. Remarkably most of the genes involved have been implicated in virulence using
animal models of infection (see Discussion, Additional file 2: Table S4).
272
The recombination profile shown in Figure 2 suggests that the MTBC ancestor
recombined with MCAN ancestral strains and, thus, they shared a common niche. A
sympatric model of speciation predicts that some parts of the genome will be involved
in adaptation to a new niche[4]. The hallmark trace would be the accumulation of
variants differentiating the emerging species, at the genome-wide level or in a few loci,
as a consequence of reduced recombination between both groups. We identified all
the variants that mapped to the ancestral branch of the MTBC and that had a different
nucleotide in all the MCAN strains, the so called divergent variants (divSNPs). The
distribution of divSNPs per gene revealed that only few of them accumulated divSNPs
in the branch leading to the ancestor while most genes did not (Figure 4A). This pattern
is compatible with population differentiation models in which niche overlap between
emerging species is high[3]. The genome-wide landscape of divergent variants (n =
5,688, Figure 4B) revealed that a total of 120 genes harbored more divergent variants
than expected by chance (see Methods)(Figure 4B).
287

Figure 4. **Divergent positions between the MTBC ancestor and _M. canettii_ clade.** A) Average of divSNPs per 10 kb positions (green) as compared to the average of homoplastic variants (gray). Blue arrows above the distribution are genes that significantly accumulate more divSNPs. B) Accumulation of divSNPs per gene, corrected by gene length. A small number of genes accumulate a high amount of divSNPs while most of the genes have a low number of variants or even none. This pattern resembles those of high habitat overlap derived from Overlapping Habitat Models(3).

However, bacterial genomes are highly dynamic and different processes can contribute to the genetic make-up of extant species. Consequently, not all the detected regions necessarily result from pure divergence by accumulation of substitutions. A detailed comparative genomics and phylogenetic analysis using available genome databases of bacteria identified several genes in which divSNPs were introduced by horizontal gene transfer (n = 12) or by recombination to a MCAN not present in our dataset (n = 54) and one with uncertain phylogenetic origin (see Additional file 1: Supplementary Text for detailed results). Thus, a total of 53 genes in the MTBC ancestral genome were highly divergent with respect to MCAN due to substitution events (Additional file 2: Table S5). While the genome-wide analysis identified divSNPs that might result from genetic drift or hitch-hiking events associated with

selection on other loci, their accumulation in only 53 genes suggests that those regions
might have played an important role during the process of niche differentiation. In
agreement, those 53 genes are significantly more conserved than the rest of the
genome (dN/dS = 0.154 vs genome average dN/dS = 0.279, chi-squared p-value <=
0.001). This result suggests that, despite the increased divergence from the MCAN
strains, those 53 regions have been evolving under purifying selection. Alternatively,
the accumulation of divergent variants could also represent hotspot regions for
mutation. None of the genes showed a similar pattern of mutation accumulation in
other MCAN (no overlap between the divSNPs probabilities distributions for these 53
genes and the rest of the genomes, t-test p-value < 0.05).
318
Regions under positive selection after the transition to obligate pathogen
Having established that some divSNPs accumulate in genes under purifying selection,
we screened for positive selection patterns to identify additional genes relevant in the
transition from a newly emerged pathogen to a globally established pathogen. We first
revisited the evolution of antigenic proteins. Those regions are recognized by the
immune system and most of them are hyperconserved within the MTBC[33,34].
Interestingly, and in agreement with previous data from MCAN genomic analyses[10],
the dN/dS calculated in the branch leading to the ancestor showed a very similar
pattern, with essential genes being more conserved than non-essential ones and T-
cell epitopes being hyperconserved (Additional file 1: Figure S8). Only nine divSNPs
(5 synonymous and 4 non-synonymous) were found in T-cell epitope regions, which
is significantly less than expected by chance (Poisson distribution, p-value < 0.001).
331
Thus, antigenic regions do not show an altered pattern or intensity of selective
pressure. We then explored what other regions of the genome changed significantly
in selective pressure by comparing the MTBC ancestor dN/dS and the actual dN/dS
in extant populations using our global reference dataset of 4,598 MTBC strains. We
calculated a dN/dS for all the genes with at least one synonymous and one non-
synonymous mutation for each of the two sets (divSNPs versus within MTBC SNPs).
Due to the low number of divSNPs in individual genes, only 499 genes were evaluated.
Consequently, although additional genes to those shown in the ensuing analyses may
have changed the selection pattern or intensity, they cannot be evaluated properly
(Additional file 2: Table S6). We were particularly interested in those genes with a
drastic change from purifying (dN/dS < 1) to diversifying or positive selection (dN/dS
>1) or vice versa.
344
Most of the genes evaluated did not show any sign of changing selective pressure or
pattern. However, when looking at the dN/dS variation data, 14 genes appeared as
outliers (as defined by Tukey's method[35])(Figure 5A). Genes Rv1244 (*lpqZ)*,
Rv3910, Rv0166 (*fadD5)*, Rv0874c, Rv1152, Rv1678, Rv1951c, Rv2584c (*apt)*,
Rv3026c, Rv3276c (*purK)*, Rv3370, Rv3759c (*proX)* and Rv3900c were under a
stronger negative selective pressure following speciation. Many of them are
annotated[36] as hypothetical conserved proteins. On the other hand, only one gene

changed to evolve under positive selection after divergence from the MTBC ancestor
Rv0758, also known as *phoR.* Notably, PhoR forms part of the PhoP/PhoR virulence
regulation system[37]. In the branch leading to the MTBC ancestor, this gene was as
conserved at the amino acid level, as other essential genes (Chi-square test; p-value
0.4721), but when we looked within the extant MTBC diversity, the gene was
significantly less conserved at the amino acid level than essential genes (Chi-square
test; p-value < 0.001).
359

Figure 5. **Genes with differential selective pressures across the MTBC speciation stages.** A) Genes changing selective pressure in the branch of the MTBC ancestor as compared to extant MTBC strains. Red lines mark those genes being outliers of the dN/dS variation distribution. B) *phoR* and *phoP* show different selective pressure dynamycs. In both cases the accumulation of nonsynonmous (dN) or synonymous (dS) mutations through time is measure as the distance to the most common ancestor of the MTBC. The dN and dS values have been corrected by the number of branches in the phylogeny at each time point.

361 **Positive selection on *phoR* linked to on-going selective pressures**

Given the known central role of PhoPR in MTBC virulence, we focused our attention on the new mutations found in *phoR*. Previous work identified a basal phylogenetic mutation in *phoR* experimentally linked to animal host preferences. Here we report 193 nonsynonymous mutations and 31 synonymous mutations exclusive of human 351 isolates and mapping to very different phylogenetic depths (Figure 6A). The average dN/dS for this gene was well above 1 (dN/dS = 2.37), suggesting the action of positive selection. Furthermore, a plot of the dN and dS values over time reveals that the overall dN/dS remained high along the evolutionary history of the MTBC (Figure 5B) in comparison with the DNA binding response regulator (*phoP*), corroborating that *phoR* has been likely under pervasive positive selection. Codon-based maximum likelihood tests of positive selection are normally powerless in intraspecies comparisons however for the case of *phoR* the tests identified a higher dN/dS than expected by chance and at least two codons with strong evidence to be under positive selection (Additional file 2:Table S7). Additional evidence for the action of positive selection on this gene derives from nonsynonymous mutations, among which we found 34 homoplastic variants, which are strong predictors of positive selection in MTBC (Additional file 2: Table S8). Nonsynonymous mutations significantly accumulated in the sensor domain (Chi-square test, p-value < 0.01), further supporting the hypothesis that they could be involved in the fine-tuning of the PhoR sensitive function to the changing environment during infection (Figure 6C).

382
383 Figure 6. *phoR* **is under positive selection in human affecting strains.** A) Genome-based
384 phylogeny calculated from a total of 4,598 clinical samples obtained from different sources.
385 The synonymous and nonsynonymous variants found in *phoR* are mapped to the
386 corresponding branch. Variants in internal branches affect complete clades which are colored
387 in the phylogeny. Homoplasies are marked in the outer circle of the phylogeny. The star marks

All the new mutations identified in our analysis were found in human clinical isolates and mapped to relatively recent branches in the MTBC phylogeny (Figure 6A). Thus, we reasoned that most mutations were associated with recent selective pressures as opposed to the previously reported mutations found in *Mycobacterium africanum* lineages 5 and 6, and the animal clade[38] that map to deep branches in the phylogeny (Figure 6A). To prove it, we tested whether novel *phoR* mutations are also arising in clinical settings during infection and recent transmission events. We used a population-based dataset from Malawi[39] where more than 70% of the strains were collected during fifteen years and genome sequenced (n = 1,187). We found 14 mutations (13 nonsynonymous and 1 synonymous) in *phoR* exclusive of the Malawi dataset with *phoR* having a dN/dS of 3.93. Moreover, the mean relative age of the nonsynonymous *phoR* variants was significantly younger than that of other nonsynonymous variants in the dataset (Welch's t-test, p-value < 0.01) and the *phoR* variants from the Malawi dataset were more recent than those *phoR* mutations from the reference dataset (Welch's t-test, p-value = 0.01)(Figure 6B). From the 13 nonsynonymous mutations in the Malawi dataset, 8 were markers of recent transmission clusters. Moreover, *phoR* mutations in the Malawi dataset involved larger transmission clusters than other mutations (permutations test, p-value < 0.001). Taken together, these data indicate that novel *phoR* mutations arise during infection and propagate during on-going human-to-human transmissions in clinical settings.

DISCUSSION

We present evidence that the MTBC ancestor transitioned to an obligate pathogenic lifestyle in sympatry from a common genetic pool including the ancestors of extant MCAN strains. It was already shown before the high recombination rate in MCAN compared to MTBC [10,11]. However, our analyses is different as it specifically focus on the branch leading to the MTBC by comparing MCAN to a reconstructed MTBC common ancestor. Specifically, we found common patterns of genome-wide recombination in the branch leading to the MTBC ancestor and the extant MCAN strains. The high recombination rate between MCAN strains, including the MTBC ancestor, stands in sharp contrast to the strictly clonal population structure of extant MTBC strains. By analyzing events leading to the transition from a recombinogenic to a clonal organism, we have also been able to identify genomic regions under different selective pressures. The comparison between selective pressures before and after becoming an obligate pathogen also allow us to propose PhoR as an important player in the past evolutionary history of the MTBC as well as in current clinical settings.

Population genomics data has led to the development and testing of different models of how different genetic clusters of the same species can arise in sympatry[1,3,40]. In the case of *Vibrio cholerae*, an appropriate combination of certain virulence-associated variants, ecological opportunity and additional virulence factors mediated the successful transition of a particular clone from an environmental to a pathogenic lifestyle[41]. Other known cases such as pathogenic *Salmonella*[42] or *Yersinia* species[43] may have followed a similar trajectory. The MTBC represents an extreme case of clonal emergence associated to its obligate pathogenic lifestyle. Here, we have shown that, despite the high average nucleotide identity between MCAN and the MTBC, there is complete genomic isolation between these organisms. There is experimental evidence that genetic exchange among MCAN strains occurs easily but not between MCAN and the MTBC[17]. We have shown that there is no measurable on-going recombination among the MTBC strains based on our analysis of 1,591 genomes and which is in agreement with other recent reports[19,44]. It is important to note that due to the low divergence within the MTBC, most methods to detect recombination are limited. Hence, we cannot completely exclude the possibility that we might have missed some recombination events. It was previously suggested that recombination (or gene conversion) could be affecting PE/PPE genes disproportionally[45]. Unfortunately, short reads cannot be properly mapped to those regions so our approach does not allow to test this possibility. However, if recombination does occur in the MTBC, it seems to have a minor impact on the overall genetic diversity of the MTBC. Recombination in natural populations depends both on the capacity of chromosomal DNA exchange between the two groups involved and on the ecological opportunity. The mechanisms, if any, by which the MTBC bacilli lost their capacity to recombine while the ancestral genetic pool showed very similar recombination patterns to MCAN strains, remains to be elucidated. Ecological opportunity may also influence on the lack of opportunities of exchange between MTBC strains. Despite the occurrence of super-infections, the bacilli occupy mainly an intracellular lifestyle, thereby reducing the opportunities for genetic exchange.

We can only speculate on how the transition from a likely environmental or opportunistic pathogen to an obligate pathogen occurred, but our analysis has identified a series of non-random evolutionary events. Notably, those events involve core pathogenesis genes. We have identified highly divergent regions in the MTBC ancestor compared to MCAN. The pattern of SNP accumulation suggests that those regions were important in the transition to a closer association with the host. In addition, recombination events mapping to the branch leading to the MTBC ancestor affected essential genes as well as genic regions known to be involved in host-pathogen interaction. The *mymA* operon (Rv3083-Rv3089) is related to the production of mycolic acids and its disruption leads to an aberrant cell-wall structure. Importantly, knock-out studies[46] have shown that this operon is essential for growth in macrophages and the spleen of infected mice. Furthermore, the deletion of genes in this operon leads to a higher TNF-alpha production, highlighting their role on regulating

host-pathogen interactions[47]. The other major operon identified in our analysis is the
mce1 operon[48]. *mce1* knock-out mutants are hypervirulent in a mouse model of
infection and lose the capacity of a proper pro-inflammatory cytokine production that
is needed for the establishment of the infection[49] and granuloma[48]. How these
processes are mediated by *mce1* is still not clear pointing to *mce1* as a priority target
for biomedical research.
482
Our analysis identified one gene, *phoR*, which is under positive selection in extant
MTBC strains although it was under purifying selection in the MTBC ancestor. PhoR
is the sensor component of the PhoPR two-component system, which plays a major
role in MTBC pathogenesis[50]. Previous experimental data show that 1) PhoPR is a
major virulence determinant in MTBC[37]; 2) that deep phylogenetic branching
mutations in PhoPR were involved in the adaptation of the pathogen to different
mammalian hosts[38] and that there is at least one case in which natural
overexpression of PhoPR in a *Mycobacterium bovis* clinical isolate was linked to a
highly transmissible and virulent phenotype in humans[50]. In fact, mutations affecting
the whole animal clade in *phoR* have been proposed to be related with the fine-tune
MTBC virulence across different animal host species. We find some of these amino
acid changes experimentally tested by Gonzalo-Asensio et al. (2014) to have been
selected multiple times in unrelated human isolates. Based on these findings, we
speculate that recent *phoR* mutations help to fine-tune the immunogenicity of the
pathogen during infection, allowing it to manipulate the human host responses and
increase the chances of transmission. However, we still need to understand the stimuli
and the molecular pathways that are at the basis of the selective pressure driving the
evolution of *phoR*. Given that PhoPR is involved in membrane composition[51],
mutations in this regulator might also be involved in the susceptibility to some
antibiotics. However, antibiotic selection is an unlikely explanation for the oldest
mutations in PhoPR, as they likely predate antibiotic usage.
504
Based on our findings, a model can be proposed in which recombination, together with
the acquisition of new genetic material[14,52], generated a favorable genetic
background for the MTBC ancestor to occupy or increase its association with
mammalian hosts. We see this emergence only once in the MTBC, perhaps because
the right combination of multiple, fortuitous genetic events and the particular ecological
conditions has occurred only once. More provocative is the idea that MTBC might just
be part of a spectrum of association to the host occupied by the different MCAN-MTBC
groups. The fact that the so-called Clone A of MCAN strains are more common in the
clinic may suggest differences in ecological niches within the MCAN group[23]. In
agreement, previous publications[10,23] and our own analysis (Additional file 1: Figure
S4) have identified Clone A strains as the closest MCAN evolutionary group to MTBC.
516
517   **CONCLUSIONS**
518

In the MTBC, the strong and obligate association with new host(s) was accompanied by new selective pressures. In accordance, we identified genes in the MTBC genome highly diverging from MCAN and evolving under purifying selection, suggesting that they have become essential following MTBC's transition to an obligate pathogenic lifestyle. In the final stages of adaptation, positive selection on genes such as *phoR* and others[53–55] likely led to a narrowing of the host-range and later still to a further fine-tuning during the spread of the bacteria within the new host species.

MATERIALS AND METHODS
Datasets used

***Mycobacterium canettii* dataset.** The *M. canettii* dataset is composed by seven draft genomes downloaded from GenBank (CIPT 140010059, NC_015848.1; CIPT 140060008, NC_019950.1; CIPT 140070008, NC_019965.1; CIPT 140070002, NZ_CAOL00000000.1; CIPT 140070005, NZ_CAOM00000000.1; CIPT 140070013, NZ_CAON00000000.1 and CIPT 140070007, NZ_CAOO00000000.1).

MTBC global dataset. This dataset includes the global representatives published by Coll *et al.*[26] (n= 1,598), a transmission dataset from Malawi (see below), and the dataset from Walker *et al.*[56] with additional sequences from Europe. The total number of sequences in this dataset when putting together was 7,977 genomes. As the datasets from Walker *et al.* 2015 and Guerra-Assunção *et al.* 2015 are enriched in epidemiological clusters, we identified all clusters at a maximum distance of 15 SNPs (common threshold in MTB epidemiology), removed samples potentially coinfected with more than one strain and then kept just one representative from each cluster. Thus, the final number of genomes for this dataset was 4,598. Out of this total, we used different subsets of strains depending on the analysis to be performed. A 1,591 sequences subset (corresponding to Coll *et al.* 2014) was used for the recombination analyses within the MTBC. A specific subset of 219 sequences, corresponding also to global representatives, was used for Gubbins because it was not computationally feasible to run the program with more strains. The total 4,598 sequences dataset was used to calculate a dN/dS for *phoR* and other genes to increment the robustness of the measures and the number of variants per gene.

Transmission dataset. A dataset including samples taken over a 15-year period in the district of Malawi[39]. This dataset is enriched in transmission clusters and was used for the *phoR* transmission analysis.

MTBC most likely ancestral genome. The MTBC ancestor was derived in a previous publication by maximum parsimony and likelihood methods[33]. This ancestor is H37Rv-like in terms of genome structural variants, but H37Rv alleles were replaced by those present in the inferred common ancestor of all MTBC lineages.

FASTQ mapping and variant calling for the MTBC strains

Fastq files were trimmed to remove low-quality reads using prinseq[57] and aligned to
the MTBC most likely ancestral genome[33] using BWA-mem algorithm[58].
Alignments with less than 20x mean coverage per base were filtered out. The variant
calling was performed using samtools[59] and VarScan[60]. Due to the low variability
found in *M. tuberculosis*, to avoid mapping errors and false SNPs a variant was filtered
out if: i) it was supported by less than 20 reads; ii) it was found in a frequency of less
than 0.9; iii) it was found near indel areas (10 bp window); or iv) it was found in areas
of high accumulation of variants (more than 3 variants in a 10 bp defined window).
Variants were annotated using SnpEff[61]. Variants present in PE/PPE genes, phages
or repeated sequences were also filtered-out, as they tend to accumulate SNPs due
to mapping errors. High quality variant calls were combined in a non-redundant variant
list and used to retrieve the most likely allele at each strain to generate a variant
alignment.
577
578

Phylogenetic inference and parsimony mapping of SNPs
In the MTBC dataset we identified 140,239 variants following the steps defined above.
As we wanted to identify nucleotide variants due to recombination events, a stricter
filtering was applied to remove putative recombination signal due to polymorphisms
introduced by other causes. Variants related to antibiotic resistance were obtained
from PhyResSe[62] and were removed from the analysis. Also, non-biallelic variants
were removed from the analysis. To avoid false positives, we also removed positions
in which a variant was called in at least one strain but also with a gap in at least another
strain. To identify variants resulting from mapping errors we generated fragments of
50 bp downstream, upstream and midstream of the variant positions in the reference
genome. With these fragments, we performed a BLAST search over the reference
genome to check whether they mapped to other regions. Variants identified in reads
that mapped to more than one region of the reference genome (query coverage per
HSP over 98% and percentage of identical matches between the query and the
reference genome of 98%) were removed from the analysis.
594

The remaining variants (94,780) were used to infer a phylogenetic tree using
RAxML[63] with the GTRCATI (GTR + optimization of substitution rates + optimization
of site-specific evolutionary rates) model of evolution and represented with the iTOL
software[64]. Variants were mapped to the phylogeny using the Mesquite suite[65].
Homoplastic variants were identified based on parsimony criteria. Using these
homoplastic variant positions, we looked for homoplastic runs. A homoplastic run was
defined by two (or more) homoplastic variants found in the genome in correlative
positions or with at least one variant between them. Variants present in the same
homoplastic run were mapped on the phylogeny using Mesquite to look for coincident
phylogenetic patterns.
605
606

607 **Linkage-disequilibrium calculation**
608   Using the filtered variant positions (94,780), we used the PLINK software[66] to
609   calculate the linkage-disequilibrium statistics D' and R^2. To estimate these values, we
610   took into account variants with a minimum frequency of 0.01 and used a sliding window

of 10 Kb. The results obtained were processed with R[67]. To plot the D' and R² pattern by variant distance, we calculated average D' and R² values for 50 bp sliding windows.

Multiple alignment of *M. canettii* and MTBC
Seven *M. canettii* draft genomes were aligned to each other and to the ancestor of MTBC using progressiveMauve[68]. The segmented alignment obtained in XMFA format was converted to a plain Fasta format using the MTBC ancestor as reordering reference with a custom Perl script. Positions with gaps in the reference sequence were removed from the final alignment, so the resulting aligned genomes had the same size than the reconstructed MTBC ancestor (4,411,532 Mb). The MTBC pseudogenomes reconstructed from mapping to the MTBC ancestor from the different datasets described above were concatenated to the *M. canettii* alignment obtained in the previous step for further analyses.

From these alignments, homoplastic variants were identified using both, parsimony and maximum-likelihood approaches[69]. Both approaches agreed in identifying the same homoplastic variants.

Recombination analyses and phylogenetic evaluation
Besides SNPs, linkage-disequilibrium analysis and Gubbins, RDP4[70] was used to detect recombination signal in the MTBC dataset. To mark the regions reported by Gubbins as potentially recombinant we required at least three of the methods implemented in RDP4 to agree in showing a significant signal.

Recombination was evaluated in the alignment containing 219 strains from MTBC and 7 *M. canettii* and in the one containing the MTBC ancestor and 7 *M. canettii*. Firstly, repetitive regions (i.e. PPE/PGRS) were masked from both alignments and, secondly, recombination events were inferred using Gubbins[71], which identifies clusters of high SNP density as markers.

Gubbins identified 70 potential recombinant regions in the alignment containing the 7 *M. canettii* strains and the MTBC ancestor. Four of these regions were obviated because they fell in regions deleted in several *M. canettii* strains. One more region was removed from the analysis because it was extremely short (41 bp) and we did not obtain reliable results in the subsequent analyses.

For the remaining 65 fragments a phylogeny was calculated using RAxML[63] and applying the GTRCATI model. Also, a reference phylogeny was calculated with the same method using the complete genomes after subtracting these 65 regions. This reference phylogeny had the same topology as the one obtained from the complete genomes. To test for phylogenetic incongruence between the putative recombination fragments and the genome phylogeny, we applied the Shimodaira-Hasegawa and Expected Likelihood Weight tests implemented in TREE-PUZZLE[72].

Dating analyses

To infer the age of the 65 recombinant fragments we first reasoned that most of the
mutations found were contributed by recombination and not by mutation once the
fragment had been integrated in the genome. Thus, before dating the fragments we
first removed all the homoplastic variants with other MCAN strain found in the
fragments. The final alignments for the 65 fragments consisted of only those variants
accumulated after the recombination event. We then used the non-recombinant part
of the genome to infer a substitution rate assuming two different dating scenarios
published for the tMRCA[6,73]. We ran BEAST[74] for each fragment pre-specifying
monophyletic groups and substitution rate based on the non-recombinant genome
phylogenetic reconstruction. We used an uncorrelated log-normal distribution for the
substitution rate in all cases and a skyline model for population size changes. We ran
several chains of up to 10E6 generations sampling every 1E3 generations to ensure
independent convergence of the parameters. Convergence was assessed using
Tracer[75]. For both evolutionary scenarios, the results obtained were largely
congruent and proportional to the age limit imposed for the MTBC ancestor. As there
is controversy about the correct MTBC ancestor age, the results were transformed to
relative ages for plotting the final results.
676
677

Gene ontology enrichment analysis
Genes present in the recombinant regions between MCAN and the MTBC ancestor
were annotated using SnpEff[61]. A Gene Set Enrichment (GSE) analysis was
performed to look for enriched gene functions in these regions. The BiNGO tool[76]
was used to study the enrichment in certain functional categories comparing the most
abundant terms in the recombinant regions with those contained in the complete
annotation. The tool uses a hypergeometric test (sampling without replacement) and
the BH correction for multiple testing comparisons. The Cytoscape program[77] was
used to visualize the results.
687
688

divSNP analysis
From the *M. canettii* and MTBC ancestor alignment, we extracted those positions
having one variant in all the *M.canettii* strains and another variant in the MTB ancestor.
The divSNP frequency by nucleotide was calculated by dividing the total number of
divSNPs (5,688) by the total number of bases in the alignment. Next, the expected
abundance of divSNPs for each gene was calculated by multiplying the nucleotide
divSNP frequency by the number of nucleotides in each gene. From the expected and
observed divSNP abundances, we used a Poisson distribution to calculate the
probability of having the observed divSNPs by chance for each gene. We selected
genes having a pFDR <=0,01 using the q-value from Storey method[78].
699
Complete mycobacterial genomes for reference strains[79](Additional file 2: Table S9)
were downloaded from RefSeq and GenBank. The orthologous genes were obtained
from the amino acid sequences and using the Proteinortho tool[80]. A gene was
considered as orthologous based on reciprocal best hits in BLAST. BLAST analysis
required a minimum identity of 25%, a query coverage of 50% and a maximum e-value
of 1E-05. The orthologous genes were aligned using Clustal-omega[81] and the
phylogenies were constructed using RAxML and applying the PROTCATIAUTO

model. The reference phylogeny was constructed using only the core genome
(proteins having orthologous in all the mycobacterial genomes downloaded) with
RAxML using the same options as above. The reference and alternative phylogenies
calculated with the orthologous genes for the divSNPs enriched genes were manually
inspected to check for congruence.
712
713
714 **dN/dS analysis**
715 The dN/dS statistics were calculated using R[67]. The potential synonymous and non-
716 synonymous substitution sites for each region were calculated using the SNAP
717 tool[82,83]. The dN/dS ratio for each region was calculated as follows:
718

$$\frac{Non\ Synonymous\ variants \big/ Non\ Synonymous\ sites}{Synonymous\ variants \big/ Synonymous\ sites}$$

720
The dN/dS for the MTBC ancestor was calculated using the divSNPs while the dN/dS
for the MTBC were calculated using 208,238 variants detected in coding regions from
the 4,598 strains in the MTBC global dataset. To look for a robust comparison between
both ratios, only genes having at least 1 synonymous and 1 non-synonymous variants
were taken into account. To compare the dN/dS ratios, both were normalized by the
genomic dN/dS for each taxon(0,24 for the MTBC ancestor and 0,59 for the MTBC).
The difference between the dN/dS ratio was calculated by subtracting the MTBC
dN/dS to that of the MTBC ancestor. The genes that account for the largest differences
in the dN/dS were identified as outliers (Q2-1.5*IQR, Q3+1.5*IQR)[35] of the
differences distribution.
731
732
***phoR* positive selection analysis**
Positive selection on *phoR* was tested using FUBAR[84] and BUSTED[85]. FUBAR
was run with 5 MCMC chains of length 10,000,000. 1,000,000 states were used as
burn-in and a Dirichlet prior of 0.5. BUSTED was run with default parameters.
To study the potential effect of *phoR* mutations on transmission efficacy we used the
dataset from Guerra-Assunção *et al.*[39]. We identified SNPs in branches leading
either to leaves or to transmission clusters. Transmission clusters were categorized in
large, medium or small according to the number of isolates in the cluster (large = over
75[th] percentile, medium=between 25[th] and 75[th] percentile, small = under 25[th]
percentile). Each gene was scored to check for accumulation of mutations in branches
leading to large transmission clusters according to the expression:
744

$$Score = Large\ clusters * 3 + Medium\ clusters$$

746
747 Genes with high mutation rates have a higher number of polymorphisms that could
748 lead to a larger score by chance. To test the probability of obtaining the observed
749 score by chance, a permutation test was carried out 10,000 times. Each of the
750 identified SNP was randomly reassigned to the same branches and the score was
751 recalculated for each gene. The expected score distribution for each gene was

compared to the observed score to calculate the probability. This test was performed for transmission events defined at 10 SNPs.

The ages for the variant positions were calculated as node-to-tip distances. These distances were relativized to the maximum root-to-tip distance to obtain a relative age value in the [0 - 1] range. In order to have a common framework, a phylogeny was constructed including all the samples from the transmission and the reference datasets. The phylogeny was constructed using RAxML and applying the GTRCATI model. For each variant position we first identified the node in which the variant appeared. The node-to-tip distance was calculated afterwards for each node using the geiger package[86]. Distances were normalized to obtain a relative distance. Later, all the non-synonymous variants except the *phoR* polymorphisms were used as a reference set. The nonsynonymous *phoR* variants to be compared were categorized in two groups, those exclusive to the reference dataset[26] and those derived from the transmission dataset[39].

PhoR domains structure representation
The PhoR domains structure was inferred by using PFAM(Finn et al. 2016) and SMART(Letunic and Bork 2018).

DECLARATIONS

DATA AVAILABILITY
The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

ACKNOWLEDGEMENTS
We thank Alberto Marina for advice in the interpretation of the PhoR molecular structure. This work was funded by projects the European Research Council (ERC) (638553-TB-ACCELERATE) and Ministerio de Economía y Competitividad (Spanish Government) research grant SAF2016-77346-R (to IC). BFU2014-58656-R and BFU2017-89594-R from Ministerio de Economía y Competitividad (Spanish Government) and PROMETEO/2016/122 from Generalitat Valenciana (to FGC). ACO is recipient of a FPU fellowship from Ministerio de Educación y Ciencia FPU13/00913 (Spanish Government). Swiss National Science Foundation (grants 310030_166687, IZRJZ3_164171, IZLSZ3_170834 and CRSII5_177163), the European Research Council (309540-EVODRTB) and SystemsX.ch (to SG).

AUTHOR CONTRIBUTIONS
IC conceived this work. ACO, IC, LSB, SH, JC and FGC analyzed the data. ACO, IC, LSB, FGC wrote the first version of the draft. All authors critically reviewed and contributed to the final version of the manuscript.

COMPETING INTERESTS

799 The authors declare no competing interests
800

REFERENCES

1. Vos M. A species concept for bacteria based on adaptive divergence.
Trends Microbiol [Internet]. 2011 [cited 2018 Feb 5];19:1–7. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/21071229
2. Shapiro BJ. What microbial population genomics has taught us about
speciation. In: Polz M, Rajora OP, editors. Popul Genomics. Springer; 2018.
3. Marttinen P, Hanage WP. Speciation trajectories in recombining bacterial
species. Tanaka MM, editor. PLOS Comput Biol [Internet]. Public Library of
Science; 2017 [cited 2018 Feb 5];13:e1005640. Available from:
http://dx.plos.org/10.1371/journal.pcbi.1005640
4. Shapiro BJ, Polz MF. Microbial Speciation. Cold Spring Harb Perspect Biol
[Internet]. Cold Spring Harbor Laboratory Press; 2015 [cited 2018 Jun
26];7:a018143. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26354896
5. World Health Organization. WHO | Global tuberculosis report 2017. WHO
[Internet]. World Health Organization; 2017 [cited 2018 Jan 16]; Available
from: http://www.who.int/tb/publications/global_report/en/
6. Comas I, Coscolla M, Luo T, Borrell S, Holt KE, Kato-Maeda M, et al. Out-
of-Africa migration and Neolithic coexpansion of *Mycobacterium tuberculosis*
with modern humans. Nat Genet [Internet]. 2013;45:1176–82. Available from:
http://dx.doi.org/10.1038/ng.2744
7. Stucki D, Brites D, Jeljeli L, Coscolla M, Liu Q, Trauner A, et al.
Mycobacterium tuberculosis lineage 4 comprises globally distributed and
geographically restricted sublineages. Nat Genet [Internet]. Europe PMC
Funders; 2016 [cited 2018 Jan 16];48:1535–43. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/27798628
8. Davies PDO. Tuberculosis in humans and animals: are we a threat to each
other? J R Soc Med [Internet]. Royal Society of Medicine Press; 2006 [cited
2018 Feb 6];99:539–40. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/17021307
9. Coscolla M, Gagneux S. Consequences of genomic diversity in
Mycobacterium tuberculosis. Semin Immunol [Internet]. 2014;26:431–44.
Available from:
http://www.sciencedirect.com/science/article/pii/S1044532314000967
10. Supply P, Marceau M, Mangenot S, Roche D, Rouanet C, Khanna V, et al.
Genomic analysis of smooth tubercle bacilli provides insights into ancestry
and pathoadaptation of *Mycobacterium tuberculosis*. Nat Genet [Internet].
Europe PMC Funders; 2013 [cited 2018 Jan 17];45:172–9. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/23291586
11. Gutierrez MC, Brisse S, Brosch R, Fabre M, Omaïs B, Marmiesse M, et al.
Ancient origin and gene mosaicism of the progenitor of *Mycobacterium*
tuberculosis. PLoS Pathog [Internet]. Public Library of Science; 2005 [cited
2018 Nov 19];1:e5. Available from:
https://dx.plos.org/10.1371/journal.ppat.0010005
12. Aboubaker Osman D, Bouzid F, Canaan S, Drancourt M. Smooth tubercle

bacilli: neglected opportunistic tropical pathogens. Front public Heal [Internet]. Frontiers Media SA; 2015 [cited 2018 Jan 17];3:283. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26793699

13. Koeck J-L, Fabre M, Simon F, Daffé M, Garnotel É, Matan AB, et al. Clinical characteristics of the smooth tubercle bacilli *Mycobacterium canettii* infection suggest the existence of an environmental reservoir. Clin Microbiol Infect [Internet]. Elsevier; 2011 [cited 2018 Jan 16];17:1013–9. Available from: http://www.sciencedirect.com/science/article/pii/S1198743X1461380X?via%3 Dihub

14. Veyrier F, Pletzer D, Turenne C, Behr MA. Phylogenetic detection of horizontal gene transfer during the step-wise genesis of *Mycobacterium tuberculosis*. BMC Evol Biol [Internet]. BioMed Central; 2009 [cited 2018 Mar 2];9:196. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19664275

15. Levillain F, Poquet Y, Mallet L, Mazères S, Marceau M, Brosch R, et al. Horizontal acquisition of a hypoxia-responsive molybdenum cofactor biosynthesis pathway contributed to *Mycobacterium tuberculosis* pathoadaptation. Behr MA, editor. PLOS Pathog [Internet]. Public Library of Science; 2017 [cited 2018 Jun 27];13:e1006752. Available from: http://dx.plos.org/10.1371/journal.ppat.1006752

16. Brennan PJ. Bacterial evolution: Emergence of virulence in TB. Nat Microbiol [Internet]. Nature Publishing Group; 2016 [cited 2018 Mar 3];1:15031. Available from: http://www.nature.com/articles/nmicrobiol201531

17. Boritsch EC, Frigui W, Cascioferro A, Malaga W, Etienne G, Laval F, et al. *pks5*-recombination-mediated surface remodelling in *Mycobacterium tuberculosis* emergence. Nat Microbiol [Internet]. Nature Publishing Group; 2016 [cited 2018 Jan 17];1:15019. Available from: http://www.nature.com/articles/nmicrobiol201519

18. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. DNA–DNA hybridization values and their relationship to whole-genome sequence similarities. Int J Syst Evol Microbiol [Internet]. Microbiology Society; 2007 [cited 2018 Jan 17];57:81–91. Available from: http://www.microbiologyresearch.org/content/journal/ijsem/10.1099/ijs.0.64483 -0

19. Boritsch EC, Khanna V, Pawlik A, Honoré N, Navas VH, Ma L, et al. Key experimental evidence of chromosomal DNA transfer among selected tuberculosis-causing mycobacteria. Proc Natl Acad Sci [Internet]. 2016 [cited 2018 Feb 5];113:9876–81. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27528665

20. Gagneux S, Small PM. Global phylogeography of *Mycobacterium tuberculosis* and implications for tuberculosis product development. Lancet Infect Dis [Internet]. 2007 [cited 2018 Jun 22];7:328–37. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17448936

21. Namouchi A, Didelot X, Schöck U, Gicquel B, Rocha EPC. After the bottleneck: Genome-wide diversification of the *Mycobacterium tuberculosis* complex by mutation, recombination, and natural selection. Genome Res [Internet]. 2012 [cited 2017 Jun 19];22:721–34. Available from:

http://genome.cshlp.org/cgi/doi/10.1101/gr.129544.111

22. Comas I, Hailu E, Kiros T, Bekele S, Mekonnen W, Gumi B, et al.
Population genomics of *Mycobacterium tuberculosis* in Ethiopia contradicts
the virgin soil hypothesis for human tuberculosis in Sub-Saharan Africa. Curr
Biol [Internet]. Elsevier; 2015 [cited 2016 Nov 7];25:3260–6. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/26687624

23. Blouin Y, Cazajous G, Dehan C, Soler C, Vong R, Hassan MO, et al.
Progenitor *Mycobacterium canettii* clone responsible for lymph node
tuberculosis epidemic, Djibouti. Emerg Infect Dis [Internet]. 2014 [cited 2018
Feb 5];20:21–8. Available from: http://wwwnc.cdc.gov/eid/article/20/1/13-
0652_article.htm

24. Liu X, Gutacker MM, Musser JM, Fu Y-X. Evidence for recombination in
Mycobacterium tuberculosis. J Bacteriol [Internet]. American Society for
Microbiology (ASM); 2006 [cited 2017 Jun 19];188:8169–77. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/16997954

25. Pepperell CS, Casto AM, Kitchen A, Granka JM, Cornejo OE, Holmes EC,
et al. The role of selection in shaping diversity of natural *M. tuberculosis*
populations. PLoS Pathog [Internet]. Public Library of Science; 2013 [cited
2018 Jun 28];9:e1003543. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/23966858

26. Coll F, McNerney R, Guerra-Assunção JA, Glynn JR, Perdigão J, Viveiros
M, et al. A robust SNP barcode for typing *Mycobacterium tuberculosis*
complex strains. Nat Commun [Internet]. Nature Publishing Group; 2014 [cited
2018 Mar 28];5:4812. Available from:
http://www.nature.com/doifinder/10.1038/ncomms5812

27. Comas I, Borrell S, Roetzer A, Rose G, Malla B, Kato-Maeda M, et al.
Whole-genome sequencing of rifampicin-resistant *Mycobacterium tuberculosis*
strains identifies compensatory mutations in RNA polymerase genes. Nat
Genet [Internet]. 2012 [cited 2016 May 4];44:106–10. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3246538&tool=pmc
entrez&rendertype=abstract

28. Farhat MR, Shapiro BJ, Kieser KJ, Sultana R, Jacobson KR, Victor TC, et
al. Genomic analysis identifies targets of convergent positive selection in drug-
resistant *Mycobacterium tuberculosis*. Nat Genet [Internet]. Nature Publishing
Group; 2013 [cited 2018 Jun 28];45:1183–9. Available from:
http://www.nature.com/articles/ng.2747

29. Mortimer TD, Weber AM, Pepperell CS. Signatures of selection at drug
resistance loci in *Mycobacterium tuberculosis*. mSystems [Internet]. American
Society for Microbiology Journals; 2018 [cited 2018 Feb 5];3:e00108-17.
Available from: http://msystems.asm.org/content/3/1/e00108-17

30. Cortes T, Schubert OT, Rose G, Arnvig KB, Comas I, Aebersold R, et al.
Genome-wide mapping of transcriptional start sites defines an extensive
leaderless transcriptome in *Mycobacterium tuberculosis*. Cell Rep [Internet].
2013 [cited 2016 Mar 24];5:1121–31. Available from:
https://www.researchgate.net/publication/259171030_Genome-
wide_Mapping_of_Transcriptional_Start_Sites_Defines_an_Extensive_Leader

less_Transcriptome_in_Mycobacterium_tuberculosis

31. Hedrick P, Kumar S. Mutation and linkage disequilibrium in human mtDNA. Eur J Hum Genet [Internet]. Nature Publishing Group; 2001 [cited 2018 Feb 5];9:969–72. Available from: http://www.nature.com/articles/5200735

32. Hershberg R, Lipatov M, Small PM, Sheffer H, Niemann S, Homolka S, et al. High functional diversity in *Mycobacterium tuberculosis* driven by genetic drift and human demography. PLoS Biol [Internet]. 2008;6:e311---e311. Available from: http://dx.doi.org/10.1371%2Fjournal.pbio.0060311

33. Comas I, Chakravartti J, Small PM, Galagan J, Niemann S, Kremer K, et al. Human T cell epitopes of *Mycobacterium tuberculosis* are evolutionarily hyperconserved. Nat Genet [Internet]. NIH Public Access; 2010 [cited 2018 Jan 3];42:498–503. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20495566

34. Coscolla M, Copin R, Sutherland J, Gehre F, de Jong B, Owolabi O, et al. *M. tuberculosis* T cell epitope analysis reveals paucity of antigenic variation and identifies rare variable TB antigens. Cell Host Microbe [Internet]. NIH Public Access; 2015 [cited 2018 Feb 5];18:538–48. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26607161

35. Tukey JW (John W. Exploratory data analysis. Reading Mass.: Addison-Wesley Pub. Co; 1977.

36. Lew JM, Kapopoulou A, Jones LM, Cole ST. TubercuList - 10 years after. Tuberculosis [Internet]. 2011;91:1–7. Available from: http://www.sciencedirect.com/science/article/pii/S1472979210001113

37. Gonzalo-Asensio J, Mostowy S, Harders-Westerveen J, Huygen K, Hernández-Pando R, Thole J, et al. PhoP: a missing piece in the intricate puzzle of *Mycobacterium tuberculosis* virulence. PLoS One [Internet]. 2008;3:e3496–3496. Available from: http://dx.doi.org/10.1371%2Fjournal.pone.0003496

38. Gonzalo-Asensio J, Malaga W, Pawlik A, Astarie-Dequeker C, Passemar C, Moreau F, et al. Evolutionary history of tuberculosis shaped by conserved mutations in the PhoPR virulence regulator. Proc Natl Acad Sci [Internet]. 2014;111:11491–6. Available from: http://www.pnas.org/content/111/31/11491.abstract N2 - Although the bovine tuberculosis (TB) agent, Mycobacterium bovis, may infect humans and cause disease, long-term epidemiological data indicate that humans represent a spill-over host in which infectio

39. Guerra-Assunção JA, Crampin AC, Houben RMGJ, Mzembe T, Mallard K, Coll F, et al. Large-scale whole genome sequencing of *M. tuberculosis* provides insights into transmission in a high prevalence area. Elife [Internet]. eLife Sciences Publications, Ltd; 2015 [cited 2017 Jan 13];4. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25732036

40. Shapiro BJ, Friedman J, Cordero OX, Preheim SP, Timberlake SC, Szabó G, et al. Population genomics of early events in the ecological differentiation of bacteria. Science [Internet]. NIH Public Access; 2012 [cited 2018 Mar 27];336:48–51. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22491847

41. Shapiro BJ, Levade I, Kovacikova G, Taylor RK, Almagro-Moreno S. Origins of pandemic *Vibrio cholerae* from environmental gene pools. Nat Microbiol [Internet]. Nature Publishing Group; 2016 [cited 2018 Jan 18];2:16240. Available from: http://www.nature.com/articles/nmicrobiol2016240

42. Bäumler A, Fang FC. Host specificity of bacterial pathogens. Cold Spring Harb Perspect Med [Internet]. Cold Spring Harbor Laboratory Press; 2013 [cited 2018 Feb 5];3:a010041. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24296346

43. McNally A, Thomson NR, Reuter S, Wren BW. "Add, stir and reduce": Yersinia spp. as model bacteria for pathogen evolution. Nat Rev Microbiol [Internet]. 2016 [cited 2018 Feb 5];14:177–90. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26876035

44. Mortimer TD, Pepperell CS. Genomic signatures of distributive conjugal transfer among mycobacteria. Genome Biol Evol [Internet]. Oxford University Press; 2014 [cited 2018 Feb 5];6:2489–500. Available from: https://academic.oup.com/gbe/article-lookup/doi/10.1093/gbe/evu175

45. Phelan JE, Coll F, Bergval I, Anthony RM, Warren R, Sampson SL, et al. Recombination in pe/ppe genes contributes to genetic variation in *Mycobacterium tuberculosis* lineages. BMC Genomics [Internet]. 2016 [cited 2018 Jun 28];17:151. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26923687

46. Cheruvu M, Plikaytis BB, Shinnick TM. The acid-induced operon Rv3083–Rv3089 is required for growth of *Mycobacterium tuberculosis* in macrophages. Tuberculosis [Internet]. 2007 [cited 2018 Feb 5];87:12–20. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16893682

47. Olsen A, Chen Y, Ji Q, Zhu G, De Silva AD, Vilchèze C, et al. Targeting *Mycobacterium tuberculosis* tumor necrosis factor alpha-downregulating genes for the development of antituberculous vaccines. MBio [Internet]. American Society for Microbiology; 2016 [cited 2018 Feb 5];7:e01023-15. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27247233

48. Casali N, White AM, Riley LW. Regulation of the *Mycobacterium tuberculosis mce1* operon. J Bacteriol [Internet]. 2006 [cited 2018 Feb 5];188:441–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16385033

49. Shimono N, Morici L, Casali N, Cantrell S, Sidders B, Ehrt S, et al. Hypervirulent mutant of *Mycobacterium tuberculosis* resulting from disruption of the *mce1* operon. Proc Natl Acad Sci [Internet]. 2003 [cited 2018 Feb 5];100:15918–23. Available from: http://www.ncbi.nlm.nih.gov/pubmed/14663145

50. Broset E, Martín C, Gonzalo-Asensio J. Evolutionary landscape of the *Mycobacterium tuberculosis* complex from the viewpoint of PhoPR: implications for virulence regulation and application to vaccine development. MBio [Internet]. American Society for Microbiology; 2015 [cited 2018 Feb 5];6:e01289-15. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26489860

51. Walters SB, Dubnau E, Kolesnikova I, Laval F, Daffe M, Smith I. The

Mycobacterium tuberculosis PhoPR two-component system regulates genes essential for virulence and complex lipid biosynthesis. Mol Microbiol [Internet]. 2006 [cited 2018 Feb 5];60:312–30. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16573683

52. Orgeur M, Brosch R. Evolution of virulence in the *Mycobacterium tuberculosis* complex. Curr Opin Microbiol [Internet]. Elsevier Current Trends; 2018 [cited 2018 Apr 23];41:68–75. Available from: https://www.sciencedirect.com/science/article/pii/S1369527417300796?via%3Dihub

53. Ates LS, Dippenaar A, Ummels R, Piersma SR, van der Woude AD, van der Kuij K, et al. Mutations in *ppe38* block PE_PGRS secretion and increase virulence of Mycobacterium tuberculosis. Nat Microbiol [Internet]. Nature Publishing Group; 2018 [cited 2018 Mar 27];3:181–8. Available from: http://www.nature.com/articles/s41564-017-0090-6

54. Malone KM, Rue-Albrecht K, Magee DA, Conlon K, Schubert OT, Nalpas NC, et al. Comparative 'omics analyses differentiate *Mycobacterium tuberculosis* and *Mycobacterium bovis* and reveal distinct macrophage responses to infection with the human and bovine tubercle bacilli. Microb Genomics [Internet]. Microbiology Society; 2018 [cited 2018 Apr 4];4. Available from: http://www.microbiologyresearch.org/content/journal/mgen/10.1099/mgen.0.000163

55. Holt KE, McAdam P, Thai PVK, Thuong NTT, Ha DTM, Lan NN, et al. Frequent transmission of the *Mycobacterium tuberculosis* Beijing lineage and positive selection for the EsxW Beijing variant in Vietnam. Nat Genet [Internet]. Nature Publishing Group; 2018 [cited 2018 Jun 28];50:849–56. Available from: http://www.nature.com/articles/s41588-018-0117-9

56. Walker TM, Kohl TA, Omar S V, Hedge J, Del Ojo Elias C, Bradley P, et al. Whole-genome sequencing for prediction of *Mycobacterium tuberculosis* drug susceptibility and resistance: a retrospective cohort study. Lancet Infect Dis [Internet]. 2015;15:1193–202. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4579482/

57. Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets. Bioinformatics [Internet]. Oxford University Press; 2011 [cited 2018 Jan 3];27:863–4. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21278185

58. Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform. Bioinformatics [Internet]. Oxford University Press; 2010 [cited 2018 Jan 3];26:589–95. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20080505

59. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence alignment/map format and SAMtools. Bioinformatics [Internet]. Oxford University Press; 2009 [cited 2018 Jan 3];25:2078–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19505943

60. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: somatic mutation and copy number alteration discovery in cancer

by exome sequencing. Genome Res [Internet]. Cold Spring Harbor Laboratory Press; 2012 [cited 2018 Jan 3];22:568–76. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22300766

61. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. Fly (Austin) [Internet]. Taylor & Francis; 2012 [cited 2018 Jan 3];6:80–92. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22728672

62. Feuerriegel S, Schleusener V, Beckert P, Kohl TA, Miotto P, Cirillo DM, et al. PhyResSE: a web tool delineating *Mycobacterium tuberculosis* antibiotic resistance and lineage from whole-genome sequencing data. J Clin Microbiol [Internet]. American Society for Microbiology; 2015 [cited 2018 Jan 3];53:1908–14. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25854485

63. Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. Bioinformatics [Internet]. Oxford University Press; 2006 [cited 2018 Jan 3];22:2688–90. Available from: https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btl446

64. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. Nucleic Acids Res [Internet]. Oxford University Press; 2016 [cited 2018 Jan 3];44:W242-5. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27095192

65. Maddison WP, Maddison DR. Mesquite: a modular system for evolutionary analysis. [Internet]. 2017. Available from: http://mesquiteproject.org

66. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. Am J Hum Genet [Internet]. Elsevier; 2007 [cited 2018 Jan 3];81:559–75. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17701901

67. Team RC. R: a language and environment for statistical computing [Internet]. Vienna, Austria; 2015. Available from: http://www.r-project.org/

68. Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. Stajich JE, editor. PLoS One [Internet]. Public Library of Science; 2010 [cited 2018 Feb 6];5:e11147. Available from: http://dx.plos.org/10.1371/journal.pone.0011147

69. Ashkenazy H, Penn O, Doron-Faigenboim A, Cohen O, Cannarozzi G, Zomer O, et al. FastML: a web server for probabilistic reconstruction of ancestral sequences. Nucleic Acids Res [Internet]. Oxford University Press; 2012 [cited 2018 Jun 28];40:W580–4. Available from: https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gks498

70. Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. RDP4: Detection and analysis of recombination patterns in virus genomes. Virus Evol [Internet]. Oxford University Press; 2015 [cited 2018 Jun 28];1. Available from: https://academic.oup.com/ve/ve/article/2568683/RDP4:

71. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid phylogenetic analysis of large samples of recombinant bacterial

whole genome sequences using Gubbins. Nucleic Acids Res [Internet]. 2015 [cited 2016 May 6];43:e15. Available from: http://nar.oxfordjournals.org/content/43/3/e15

72. Schmidt HA, Strimmer K, Vingron M, von Haeseler A. TREE-PUZZLE: maximum likelihood phylogenetic analysis using quartets and parallel computing. Bioinformatics [Internet]. 2002 [cited 2018 Jan 3];18:502–4. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11934758

73. Bos KI, Harkins KM, Herbig A, Coscolla M, Weber N, Comas I, et al. Pre-Columbian mycobacterial genomes reveal seals as a source of New World human tuberculosis. Nature [Internet]. Nature Publishing Group; 2014 [cited 2018 Mar 5];514:494–7. Available from: http://www.nature.com/articles/nature13591

74. Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, et al. BEAST 2: a software platform for bayesian evolutionary analysis. Prlic A, editor. PLoS Comput Biol [Internet]. Public Library of Science; 2014 [cited 2018 Jan 3];10:e1003537. Available from: http://dx.plos.org/10.1371/journal.pcbi.1003537

75. Rambaut A, Suchard M, Xie D, Drummond A. Tracer v1.6. 2014.

76. Maere S, Heymans K, Kuiper M. BiNGO: a Cytoscape plugin to assess overrepresentation of Gene Ontology categories in biological networks. Bioinformatics [Internet]. 2005;21:3448–9. Available from: http://bioinformatics.oxfordjournals.org/content/21/16/3448.abstract N2 - Summary: The Biological Networks Gene Ontology tool (BiNGO) is an open-source Java tool to determine which Gene Ontology (GO) terms are significantly overrepresented in a set of gen

77. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res [Internet]. 2003;13:2498–504. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC403769/

78. Storey JD. The positive false discovery rate: a Bayesian interpretation and the q-value . 2003;2013–35. Available from: http://projecteuclid.org/euclid.aos/1074290335

79. Fedrizzi T, Meehan CJ, Grottola A, Giacobazzi E, Fregni Serpini G, Tagliazucchi S, et al. Genomic characterization of nontuberculous Mycobacteria. Sci Rep [Internet]. Nature Publishing Group; 2017 [cited 2018 Feb 5];7:45258. Available from: http://www.nature.com/articles/srep45258

80. Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ. Proteinortho: detection of (co-)orthologs in large-scale analysis. BMC Bioinformatics [Internet]. BioMed Central; 2011 [cited 2018 Jan 3];12:124. Available from: http://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-12-124

81. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol [Internet]. EMBO Press; 2011 [cited 2018 Jan 3];7:539. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21988835

82. Ota T, Nei M. Variance and covariances of the numbers of synonymous and nonsynonymous substitutions per site. Mol Biol Evol [Internet]. 1994 [cited 2018 Jan 3];11:613–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8078400

83. Korber-Irrgang B. HIV signature and sequence variation analysis. Computational analysis of HIV molecular sequences [Internet]. 2000 [cited 2018 Jun 28]. Available from: https://www.scienceopen.com/document?vid=f054582b-15a6-4ccc-be34-99d0e9d09004

84. Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, et al. FUBAR: a fast, unconstrained bayesian approximation for inferring selection. Mol Biol Evol [Internet]. 2013 [cited 2018 Feb 6];30:1196–205. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23420840

85. Murrell B, Weaver S, Smith MD, Wertheim JO, Murrell S, Aylward A, et al. Gene-wide identification of episodic selection. Mol Biol Evol [Internet]. Oxford University Press; 2015 [cited 2018 Mar 5];32:1365–71. Available from: https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msv035

86. Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, FitzJohn RG, et al. geiger v2.0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. Bioinformatics [Internet]. 2014 [cited 2018 Feb 6];30:2216–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24728855