1   **Refining sperm quality assessment by high-throughput nuclear morphometric analysis**

2
Abstract

Artificial insemination (AI) is commonplace in commercial pig breeding, and as such,

ensuring sperm sample quality is of utmost importance to avoid reduced farrowing rates

and litter sizes. Here, we have used high-throughput nuclear morphometric analysis to

compare pig sperm samples categorised as meeting the AI standard (AIS) or not meeting the

AI standard (N-AIS). We find that pig sperm nuclei are asymmetric, and that samples contain

phenotypic shape abnormalities that show continuous variation. Samples classed as N-AIS

have more abnormally shaped sperm than AIS samples, but conventional analysis misses

many. The specific phenotypes identified suggest aspects of spermiogenesis that may be

disrupted and indicate future avenues for improving pig sperm quality. This method reveals

a significant difference in sperm head morphology between AIS and N-AIS pig sperm

samples and has the potential to be further developed as a high-throughput tool for sperm

head morphology assessment both in the pig breeding industry and in other species.

Keywords: Artificial insemination, morphology, nucleus, pig, sperm

Background

Infertility, defined as the failure to achieve a pregnancy after one year of regular unprotected sexual intercourse, will affect around one in six people. Between 20 and 30% of these cases result solely from male factor infertility, with up to 50% of all instances being at least partly the result of male factor infertility [1]. The analysis of semen parameters is an important part of infertility investigations, with semen volume and pH being documented, and sperm parameters such as concentration, progressive motility and morphology all being routinely observed [2]. Many of these factors are determined during spermatogenesis, the complex process by which undifferentiated diploid spermatogonia become highly specialised haploid spermatozoa, with the process involving complicated cellular, proliferative, and developmental phases [3]. Which single sperm parameter is the most important for fertility is a source of conflict in the literature. Some studies, for example, suggest that sperm progressive motility is positively associated with pregnancy rates [4,5], whilst others do not [6]. Similarly, research has indicated that sperm morphology may be correlated with positive pregnancy outcomes [7–9], but others have suggested that this plays less of a role than previously thought [10].

In agriculturally important species such as pigs, cattle, and sheep, semen analysis followed by artificial insemination (AI) is commonplace in breeding programmes [11], with 90% of sows in the world's primary pork producing countries resulting from AI [12]. As such, an industry-led working code of practice, the AI standard for pigs has been developed to assure standards of AI are received by pork producers in the UK [13]. Generally, the goals of agricultural animal breeding are to maximise the production of meat at a low cost, and to

efficiently disseminate the genetics of particularly valuable animals. As such, the use of high-

quality semen doses is of paramount importance, for example in pigs the use of lower-

quality doses is associated with lower farrowing rates and smaller litter sizes [14,15]. The

fast identification of animals that consistently produce N-AIS samples is therefore important

to ensure their removal from breeding programmes; it has been shown for example that

boars with fertility problems in a nucleus herd have the potential to reduce litter sizes

throughout the entire breeding population [16], in part due to one boar ejaculate commonly

being used to serve between 15 and 25 sows [17] in routine AI. Semen parameters are

therefore of key importance both in human fertility assessment and treatment and in

agricultural practice.

As is the case for all mammals, an ejaculate from a boar, bull or ram does not contain a

homogenous population of sperm [18] making analysis and the subsequent identification of

subfertile individuals, challenging. In the general analysis of boar sperm morphology for

example, a sample is deemed to be "fertile" if the frequency of abnormal sperm heads does

not exceed 10%. Fertility can also be assumed if the frequency of abnormalities in

acrosomes, mid-pieces, tails, or proximal cytoplasmic droplets is less than either 5% each, or

15% when combined [19]. In humans, the percentage of normal forms (in both fertile and

subfertile men) is usually far less than 30% [20], and has been documented to be

consistently (and staggeringly) low, with only 3-5% normal forms being identified in assisted

reproductive technology (ART) cases [9,21,22].

There also remains much to learn about mammalian sperm biology. Only a limited number

of studies to date have focused on the analysis of sperm nuclear morphometry in

agricultural animals [23,23,24], and whilst it has previously been shown that sperm shape

differs between high and low fertility bulls [25], such observations are limited in pigs [26,27]

and differences in morphometric variables that may be related to pig fertility are yet to be

elucidated [28]. Whilst the goals of the livestock industry are in stark contrast to human

fertility treatment, furthering our understanding of the biology of sperm in general has

potential benefits for both fields. Here, we have used Nuclear Morphology Analysis (NMA)

[29–33], morphometrics software for microscopy images, to measure and compare the

shapes of sperm nuclei from 21,002 pig sperm cells. This is the largest reported number of

individual sperm nuclei that have been analysed at this level of detail in any agricultural

animal. Our objective is to understand how sperm shape relates to samples being classified

as AIS or N-AIS, and to determine if certain sperm features can be correlated with specific

biological processes. If these differences can subsequently be linked to fertility, such

analysis has the potential to provide an alternative to the livestock industry standard

practices and may provide transferrable insights into human sperm morphology.

Methods

Semen collection

Fresh ejaculated sperm samples from boars were collected using the gloved hand method

[34], by trained staff at JSR Genetics Ltd. Samples were stored in Duragen extender,

supplemented with no less than: 500 IU per ml streptomycin; 500 IU per ml penicillin; 150

mg per ml lincomycin; and 300 mg per ml spectinomycin, diluted to 2.3 billion sperm per

dose. Samples were stored at 17°C and were prepared within two days of collection. No

specific ethical approval was required for this study as all semen doses used were collected

as part of JSR Genetics Ltd.'s standard commercial procedures. Semen samples identified as

N-AIS by the industry were discarded, and those that were identified as AIS by the industry

were used commercially.

Sample preparation

Prior to preparation of samples for this study, semen samples were identified as either AIS

or N-AIS using computer assisted sperm analysis (CASA) followed by manual assessment at

JSR Genetics Ltd. Specifically, samples that had a morphology score of above 70% (obtained

from CASA) and a motility score of above 4 (1 to 5, 1 being dead and 5 being excellent

progressive forward movement) were graded as AIS and those falling below these criteria

were graded as N-AIS.

50 AIS and 44 N-AIS samples were used in this study. 2ml of each semen sample was

centrifuged at 300g for 5 minutes at 17°C. The supernatant was discarded, the pellet was

resuspended in 1.5ml of fixative solution (100% methanol and 100% acetic acid, added

dropwise at a 3:1 ratio) and centrifuged at 300g for 5 minutes at 17°C. The supernatant was

discarded, and the pellet was resuspended in 1.5ml of fixative solution. 10µl of each sample

was then dropped onto the centre of the surface of a labelled (sample ID, date), steam-

warmed slide, immediately followed by 10µl of fixative solution. Subsequently, slides were

air-dried for two minutes before one drop of fluorescent DAPI (4',6-diamidino-2-

phenylindole) was added to the centre of the slide. Prepared slides were air-dried in the

dark, for at least 20 minutes prior to microscopy.

Image acquisition

An Olympus IX83 inverted fluorescence microscope equipped with a XM10 monochrome

CCD camera and CellSens Dimension version 1.9 (expandable imaging software for Life

Science microscopy) was used for image capturing. A minimum of 200 nuclei per sample

were imaged at 1000x magnification.

Data analysis

Images were analysed using Nuclear Morphology Analysis (NMA) v2.2.1 [30,33], available

here). This software enables automated recognition of round or asymmetric nuclei within an

image of interest, and the subsequent morphological analysis of these nuclei. We adapted

the landmark recognition to detect and analyse pig sperm nuclei. The software generates

'profiles' describing the interior angles around the perimeter of the nucleus, and the

diameter and radii across the centre of mass (Figure 1). Pig sperm are (mostly) symmetrical

about the anterior-posterior axis, and the tail attachment region, characterised by a 'dimple'

in the nucleus, was chosen to anchor the profiles and orient the nuclei (Figure 1A, reference

point 1).

Figure 1 – Example morphometric profiles of pig spermatozoa. A) angle profiles measure the internal angle around the perimeter of the nucleus. Diameter profiles measure diameter across the centre of mass. The numbered points are indicated in the profiles, demonstrating key landmarks can be identified. B) Profiles show median and interquartile range of the angles and diameters of cells. C) schematic of a pig spermatozoon; the sperm nucleus (dark grey) fills most of the sperm head, with the remainder of the head comprising the acrosome (green) and the residual cytoplasm (lighter grey). The tail (purple) is not

shown to scale but is included to illustrate the attachment region.

For consistent alignment of the nuclei, the tail attachment region (as can be seen in Figure 1

A point 1 and 1C) was placed directly below the centre of mass of the nucleus.

Data was exported from NMA and further processed in R (4.4.0) [35] using a custom analysis

pipeline. Initially, k-means clustering on diameter profiles was used to identify distinct

groups of shapes, with dimensionality reduction using UMAP [36,37] employed to visualise

the shape profiles. Continuous phenotypes were then detected by selective focus on outlier

cells, detected using the angle, radius and diameter profiles (Figure 1B).

Cells were classed as normal or abnormal using the profile data. Each profile contains 100

indexes (Figure 1A and 1B); at each index, potentially abnormal cells were detected as those

cells with values outside 1.5x the interquartile range. Cells were then scored on how many

indexes met the criteria for abnormality. Consistent abnormal phenotypes have higher

values, while more normal phenotypes have lower values. For each profile, the cells with a

score higher than the mean number of abnormal indexes were determined to be

morphological outliers. This allowed us to identify continuous morphological variation

across cell populations. Cells were assigned a grouping of 'normal' (not identified as

outliers), 'intermediate abnormality' (the cell was an outlier across a single profile type) or

'extreme abnormality' (the was an outlier across multiple profile types).

Associations of breed, sample and date of collection with the severity of phenotype were

tested using a Scheirer-Ray-Hare test (a non-parametric equivalent to a two-factor ANOVA).

A significant association prompted the use of ordinal logistic regression to explore the

association further. The samples were filtered to only known breeds and collection dates for

which all samples had data (9321 cells). Ordinal logistic regression was implemented with

the *polr* function in MASS [38] using a negative log-log link to account for higher proportions

of normal phenotypes, modelling the phenotype cluster as the interaction of collection date

and sample. The proportional odds assumption of the model was tested with the Brant-

Wald test, and an intercept-only model was created to test the null hypothesis that the

predictor variables added no information.

The predictive ability of morphometric data to classify samples as either AIS or N-AIS was

tested using logistic regression via generalised linear models. Two models were created; the

first using standard industry measurements (width, perimeter, area, ellipticity, elongation,

roughness, regularity and height [19,39]), and the second using the morphometric profiles

(angle, radius and diameter), modelling a sperm class as dependent on the morphometric

parameters. The samples were split into training (70%) and test (30%) groups. The

probability of being in the AIS group was calculated for each sperm in a sample, and the

mean probability was taken as the overall sample value.

Figures were generated using ggplot2 [40],UMAPs were generated using the UMAP package

[36,37], Factoextra was used for heirarchicdal clustering and Caret was used for GLM

validation. Scripts used in this analysis are available here.

Results

Pig sperm samples contain continuous phenotypic abnormalities

Analysis of sperm nuclear morphology from 50 AIS ejaculates and 44 N-AIS ejaculates

yielded measures from 11,416 and 9,586 nuclei, respectively. Comparisons between these

groups of animals indicated that sperm heads differ, but that there is a large amount of

overlap and variation between individuals (Figures S1-4).

To identify different nuclear shape phenotypes in the samples, all samples were combined

into a single dataset to enable a full unbiased comparison of shapes. The first global

observation indicated that sperm fell clearly into two equal sizes groups with distinct shape

profiles (Figure 2A).

Figure 2. Cell shapes fall into two equally sized groups with mirroring around the vertical axis; A) diameter profiles for each cluster show the symmetrical clusters B) Consensus nucleus shapes show the main difference is at the tail attachment region with the more prominent side to the left or right respectively. C) the left cell in B mirrored across the vertical axis and overlaid on the right cell shows the two cell populations are mirrors of each other. D) Dimensionality reduction of diameter profiles by UMAP shows almost all

cells fall cleanly into these two groups.

These two groups of sperm cells correspond to the differences in the shape of the tail

attachment region, which is compressed on either the left or the right side (Figure 2B). The

profiles of the nuclei with compression on the right side were reversed to determine if the

shapes were mirrored and superimposable. The two groups are indeed mirror images of

each other around the tail attachment region (Figure 2C), and almost all cells in the dataset

fall neatly into these two major groups (Figure 2D). This, plus the general inflexibility of

sperm chromatin, suggests the differences are unlikely to be due to compression from the

angle of the tail at the time of fixation. Rather, the nucleus appears truly asymmetric, and

the difference we observe is caused by the sperm head falling on its dorsal or ventral

surface when the slide was prepared.

Since the shape around the tail attachment region was the primary factor that separated

nuclei of different shapes, and UMAP showed continuous variation, standard clustering *e.g.*

[41] to identify discrete shape groups was not feasible. Instead, we aggregated outlier nuclei

with similar shape profiles to identify continuous phenotypes, iteratively grouping nuclei

with the greatest differences to normal profiles. This amalgamation of outlier shapes

yielded a normal phenotype group (containing 85.2% of nuclei), an intermediate phenotype

group (12.7%) and an extreme phenotype group (2.1%). The progression from intermediate

to extreme involved condensation of the basal region and bulbing of the anterior region

(Figure 3) with two main pathways; one being the compression with failure to elongate

resulting a short pyriform phenotype (Figure 3A;7.75%) and the other being abaxial tail

attachment and basal compression resulting in an abaxial pyriform phenotype (Figure

3C;6.25%). The intermediate phenotypes are the hardest to categorise as the vary from with

minor tapering at the basal region that is difficult to detect to the relatively well-formed side

of the spectrum of pyriform cells. This spectrum makes up the largest portion of the

abnormality in pig sperm and is highly variable, containing cells that are either the

intermediates of a spectrum of abnormality or the end of more minor morphological

changes. There is also within the intermediate phenotypes minor tapering and formation of

spikes in the basal region (Figure 1B; 1%).

Figure 3. Example DAPI stained images of progressively abnormal sperm phenotypes. Three

main classes of abnormality can be seen: A) short (compression anterior-posterior, leading

to reduced bounding height) progressing to compression (tapering) around the base of the

nucleus and a pyriform phenotype; B) extrusion of nuclear DNA forming a spike near the

base of the nucleus with some tapering; C) abaxial compression (tapering) around the base

of the nucleus, leading to a pyriform phenotype.

AIS and N-AIS samples both contain abnormal sperm phenotypes

Our initial analysis indicated that *c.*15% of the sperm were abnormally shaped, but, as

samples were pooled, this did not indicate if these were enriched in the N-AIS samples or if

their abundance related to other characteristics of the samples. images showing the

progressive phenotypes from normal to extreme are shown in Figure 1. We therefore

looked at three characteristics: the sample's status as AIS or N-AIS, the breed of the boar

(where known, n= 11,448 sperm heads) and the date of sample collection (as a proxy for

environmental or technical factors). Intermediate and extreme sperm phenotypes were not

evenly distributed across all samples (Figure 4), and while the abnormal phenotypes were

more prevalent (1.96:1 N-AIS:AIS more extreme, 1.44:1 N-AIS:AIS more intermediate) in the

N-AIS samples, they were also found in AIS samples.

Figure 4 – Proportion of sperm in normal, intermediate or extreme groups for A) AIS samples (50 samples) and B) N-AIS samples (44 samples). Intermediate and extreme phenotypes are more frequent in N-AIS samples but are also common in a subset of the AIS samples.

Given that the pattern of data suggested there was variation between samples and between

sample collection dates (Figures S3 and S4), we tested whether the distribution of

phenotypes could be associated with the date of sample collection or individual samples

across the entire dataset. Significant associations were found for both variables and their

interaction (Scheirer-Ray-Hare test, $p<0.001$ for each). To explore the interactions in more

detail, we focussed on the samples for which breed was known. We then selected dates for

which all these breeds were present. Ordinal logistic regression did not reveal a global

association between the severity of the phenotype and the individual sample for the known

breeds (p>0.05) apart from White Duroc (p=0.01) in which severe phenotypes were more

common (Table S1). The odds that a sperm cell had an extreme phenotype versus

intermediate or normal was increased 2.1 times compared to other breeds.

The date of collection had some global associations with severity of phenotype; for

example, for samples collected on one date (6[th] July), the odds that a sperm cell had an

extreme phenotype versus intermediate or normal was increased 3.3 times compared to

other dates (p<0.001). Most other dates did not show global effects.

Interaction effects were detected between the date of collection and individual samples

(see Table S1 for a full breakdown), indicating that the environment has a larger impact on

the quality of the ejaculate than the breed of the boar. This was confirmed by testing for

differences in the shape profiles between breeds. Repeated Wilcoxon rank-sum tests,

multiple testing corrected with the Holm method showed no significant difference in the

angle, radius or diameter profiles between breeds suggested that there is subtle nuclear

morphological variation between breeds, with differences dominated by shared genetic or

environmental factors.

Classification of normal/abnormal cell types does not align with industry assigned AIS/N-AIS

classifications

We next considered if the extra morphometric information measured allowed for more

robust classification of samples as AIS/N-AIS than is currently possible. The samples were

originally classified based on motility and shape, with 70% morphologically normal forms

and a motility score of above 4 leading to a classification of AIS. Although motility

information was not available for these samples, the predictive power of conventional

morphometrics was compared. For this, a GLM was created to predict the probability of a

cell as originating from an AIS or N-AIS sample based on the width, perimeter, area,

ellipticity, elongation, roughness, regularity and height, as are commonly used in industry.

Each sample was scored as the mean of its individual cell predictions (Figure S2A). This score

was then compared to a GLM that used the shape profiles for prediction (Figure S2B). This

comparison indicated that greater discrimination between the AIS and N-AIS samples was

possible. The ability to classify samples directly by the proportion of normal, intermediate or

extreme phenotypes is in principle a useful improvement on the simpler morphometric

measures; AIS samples with a high rate of shape abnormalities in particular show that there

is morphological variation present that needs to be further explored.

Discussion

Continuous sperm shape phenotypes unify previous understanding of pig sperm variation

Studies of pig sperm shape, and indeed the sperm of other species, usually focus on simple

morphometric measurements such as area and circularity [19,28,42–44]. However, these

measures can miss subtle phenotypes, especially when large numbers of sperm are being

screened. Here, 21,002 sperm nuclei have been analysed at high resolution allowing in-

depth comparison of sperm nucleus shape. Previous pig sperm morphology studies report

the analysis of between 3,000 and 5,550 sperm heads [43,45]. To the best of our

knowledge, this is therefore the largest reported number of individual sperm nuclei that

have been analysed at this level of detail in any agricultural animal. The use of this large

sample size has allowed us to detect and characterise subtle phenotypes that have

previously been overlooked.

Previous studies measuring bright-field images have identified differences in sperm head

size between pig breeds with Saravia *et al.* concluding that Duroc sperm heads were 0.2μM

longer and 0.1μM wider than other tested breeds [19], and Kawarasaki, Enya and Otake

finding Duroc sperm heads to be 0.1μM longer and 0.03μM wider [45]. Given that such

differences were not observed here (Figure S1), the size difference seen in other studies

may be due to the structures outside of the nucleus, or due to sample variation. Of note

however, is that the Duroc pig samples analysed in this study were from crossbred white

Duroc animals. This crossbreeding may have resulted in the smaller sperm observed due to

a lack of genetic diversity or a potential founder effect (Figure S1).

As shown in Figure 2, asymmetry in pig sperm nuclei was identified, with one side of the tail

attachment being consistently larger than the other. Given that pig sperm nuclei are

flattened, we conject that this asymmetry reflects how the sperm cells fell on the slide

during preparation; either dorsally or ventrally. Evidence from scanning electron microscopy

also suggests that the asymmetry is present in the full sperm head, not just the nucleus [45].

The other shape phenotypes that were discovered in this study were continuous, with

progressive abnormalities involving compression around the base of the nucleus (Figure 3).

Some of these shape phenotypes are similar to those reported before, including the division

of intermediate and extreme phenotypes [43,45,46].

Basal nucleus compression causing a bulb-like shape has been observed in the sperm of

other species such as humans [46], and has been studied extensively in cattle [47–51].

Notably, *Kif3a* knockouts in mice have also been shown to generate similar sperm

phenotypes [52], although much of the characterisation of sperm phenotypes to date has

been either manual or using CASA systems measuring simple morphometric values, and

therefore lack the power required to detect subtle variations [23,24,28,43,45,53,54].

Some of the samples with the greatest proportion of abnormal sperm were collected on the

same day, suggesting that environmental factors, either pre- or post- sample collection, may

be as, if not more, important than genetic factors, as borne out by previous studies [55–57].

Purely technical factors in sample collection seem implausible to explain the range of

phenotypes we observed. Whilst breeding seasons are unlikely to be the cause (as pigs are

nonseasonal), it has been shown that stress and physical trauma can affect spermiogenesis,

with stress-induced infertility being previously been reported in mice, boars and humans

[56,58,59]. Although chromatin integrity is important for fertility [60], morphological

differences head shapes are not directly due to chromatin integrity in stud boar sperm [19]

or in humans [61] [62] except in the very round sperm examples seen in very few of in pig

samples. This suggests that chromatin integrity is not an important factor in the structure of

the main phenotypes, as there are very few other sperm that show known oxidative

damage-induced phenotypes. It is unlikely that there would be such a disparity in

phenotype representation if they were related.

Progressive pig sperm phenotypes match known defects in mechanisms of spermiogenesis

The sperm were organised into three groups that illustrate the progressive abnormalities

(Figure 3). These are (1) extrusion of a basal spike; (2) basal compression, developing

asymmetrically; and (3) basal compression with failure of the remainder of the nucleus to

elongate. Potential mechanisms that can generate these phenotypes have been described in

mice, and information from other mammals can be used as an analogue for boar

spermatogenesis [63,64].

The elongation of the nucleus is shaped by five core elements. First, the ectoplasmic

specialisation of the Sertoli cell, which provides compressive forces externally [65,66].

Second, the acroplaxome (a cytoskeletal plate composed of f-actin and keratin) upon which

the Sertoli cell pushes for nuclear elongation [65,67,68]. Third, the Sertoli cell-spermatid

linkers (afadin/nectin complexes) which transmit the forces from the Sertoli cell [69].

Fourth, the manchette (a network of microtubules around the base of the sperm head)

which directs the basal nuclear structure and tail attachment [52,70,71]. Fifth, the groove

belt (the connection between the machete and the nucleus where the acroplaxome ends;

[52,70,71]. These components are interconnected, and while not the only mediators of

sperm shape, they are important for understanding how perturbations to the process of

elongation in spermiogenesis can cause sperm defects. Many proteins are involved in

spermatid elongation; SUN3, SUN4 and SUN5 are of particular importance in the process of

manchette formation and the attachment of the sperm head to the tail. Both SUN3 and

SUN4 knockouts result in a phenotype that does not successfully elongate and have a round

nucleus with a mispositioned acrosome [70,72,73], with SUN3 knockout also causing

complete failure of manchette formation [74]. SUN5 anchors sperm heads to their tails [75]

and does not affect the acrosomal attachment [76]. SUN5 knockouts form acephalic

spermatozoa [75]; these are unlikely to be the cause of the abnormal sperm in our samples

as we did not see these phenotypes. Other proteins are more likely candidates to explain

these abnormalities. The dynein and kinesin motor proteins are involved in elongation and

the action of the Sertoli cell ectoplasmic specialisation. Dynein knockouts in mice result in

continuous deformation of sperm with malformation of the general structure with reduced

elongation and poor formation of the structures around the acrosome [77–80]. Kinesins are

also involved [65] and knock outs of the kinesin *KIF3A* give rise to bulb-like morphology with

over compression of the manchette [52].

Nectin-2/3 is part of the spermatid-Sertoli cell connections directing the majority of sperm

head elongation [65]. Nectin-2/3 disruption in spermatogenesis affects fertility [81], and is

linked to environmental changes [56]. Deficiency of Nectin-2 has also been shown to give a

diverse range of abnormal sperm in mice similar to the general structural abnormality

shown in the pig sperm [82].

Whilst Figure 3 showed the progression of phenotypes with increasing severity, we outline

the structural elements that influence sperm shape in Figure 5 and indicate the forces that

may be operating on the developing sperm to generate these phenotypes.

Figure 5 – Summary of the forces involved in shaping the sperm nucleus and the abnormal phenotypes we observe. (A) the normal forces of elongation of a pig sperm exerted on and generated within the cell by the Sertoli cell ectoplasmic specialisation. The Basal spike (B) section shows the forces happening to the cells with an anterior bulge around the tail attachment region that is possibly either from the force of a broken tail or malformation of the manchette/ ectoplasmic specialisation hoops slipping due to lack of acroplaxome attachment proteins below the manchette forcing out the bulge. The short with basal compression (C)phenotype has concentrated pressure around the base and a lack of intranuclear pressure which could come either from a lack of molecular motors or misalignment of ectoplasmic specialisation hoops that results in a lot of basal pressure causing constriction but not the elongation seen in other phenotypes. The elongation with increased tail asymmetry (D) is potentially due to manchette misattachment causing the asymmetry that is not severe enough to cause a full abaxial structure or abaxial misalignment of the hoops from the ectoplasmic specialisation without enough constricting force to cause full abaxial development. The elongation of this sperm phenotype is likely due to increased intranuclear pressure from the compression of the manchette and loops that are potentially due to increased molecular motor activity such as dyneins or kinesins or potentially moderated by the same acroplaxome attachment complex that allows for more force transfer due to more attachment. The Abaxial (E) phenotype with basal compression is likely due to a combination of manchette misalignment exacerbated by the basal compression due to the previously mentioned processes.

We see over compression of the manchette and a dysfunction of nuclear elongation;

potentially the absence or reduction in levels of one of these connector proteins or

molecular motors would result in over-compression of the manchette and improper

elongation [52,82]. One potential mechanism for this could involve the over-tightening of

the hoops of the Sertoli cell ectoplasmic specialisation via uneven pressure exerted on the

nucleus, either because the hoops lack constricting forces or because they lose their solid

connection to the acroplaxome and prevent its function as a modulator of elongation.

Although we cannot be certain that these are the mechanisms at play in these boar

samples, our analysis indicates that these structural changes are localised to specific regions

of the sperm. This suggests that further study of the proteins involved in these regions will

be important to understanding variation in boar sperm development. Particularly, if the

components are affected by environmental factors, there may be possible interventions to

improve sperm quality and ejaculate consistency.

Implications for the pig breeding industry

This work, and Barquero *et al.* [83] both show a consistent distribution of morphological

phenotypes. Barquero *et al.,* however, only used samples comparable to those in our study

categorised by breeders as AIS. Given the previously reported link between sperm head size

and motility with simple morphological parameters such as height and width [84], and an

association between the number of stillbirths in Piétrain boars and poor sperm motility [83],

we see clear support for the well-established idea that boar semen quality is significantly

affected by the environment [55,57,85]. Further investigation of this relationship is needed,

particularly focussing on candidate proteins that can explain the phenotypes; a deeper

understanding of sperm morphology may further explain the relationship between

morphology, motility and farrowing outcomes.

Evidence suggests that pig AI may be suboptimal [86–89], with this potentially due to the

methods used to classify sperm at AIS or N-AIS. This interpretation is supported in this

study, where a large overlap of morphology and the subtle intermediate phenotypes of

abnormal sperm between AIS and N-AIS samples is seen. It would therefore be an important

next step to test the fertility of samples classified as such using this new approach.

Conclusions

In this study we demonstrate that morphometric analysis of pig sperm using NMA software

effectively distinguishes morphologically distinct populations of nuclei. Our analysis reveals

differences in sperm nucleus morphology between animals classified as commercially

acceptable (AIS) and unacceptable (N-AIS) based on assessment to industry standards. Use

of NMA has offered additional insights into pig sperm. Importantly, the distribution of

sperm phenotypes between different collection dates and between breeds suggests that

environmental factors in sperm collection and transport influence can greatly influence

sperm morphology and fertility. Further study is needed to understand how these

phenotypes develop, and the functional consequences for boar fertility. These methods may

also be of use in the categorisation of the phenotypic distributions within human sperm and

have the potential to provide insights into the various complex and subtle morphological

differences that could be linked to environmental factors, fertility, and disease.

Acknowledgments

We would like to thank JSR Genetics Ltd. for supplying the samples that were used in this

study.

References

1. Agarwal A, Mulgund A, Hamada A, Chyatte MR. 2015 A unique view on male infertility

around the globe. *Reprod. Biol. Endocrinol. RBE* **13**, 37. (doi:10.1186/s12958-015-0032-1)

2. Organization WH, Geneva. 2021 WHO laboratory manual for the examination and

processing of human semen.

3. Sharma R, Agarwal A. 2011 Spermatogenesis: An Overview. In *Sperm Chromatin:*

Biological and Clinical Applications in Male Infertility and Assisted Reproduction (eds A

Zini, A Agarwal), pp. 19–44. New York, NY: Springer. (doi:10.1007/978-1-4419-6857-9_2)

4. Liu DY, Clarke GN, Baker HWG. 1991 Relationship Between Sperm Motility Assessed with

the Hamilton-Thorn Motility Analyzer and Fertilization Rates *In Vitro*. *J. Androl.* **12**, 231–

239. (doi:10.1002/j.1939-4640.1991.tb00258.x)

5. Donnelly ET, Lewis SEM, McNally JA, Thompson W. 1998 In vitro fertilization and

pregnancy rates: the influence of sperm motility and morphology on IVF outcome. *Fertil.*

Steril. **70**, 305–314. (doi:10.1016/S0015-0282(98)00146-0)

6. Moghadam KK, Nett R, Robins JC, Thomas MA, Awadalla SG, Scheiber MD, Williams DB.

2005 The Motility of Epididymal or Testicular Spermatozoa Does Not Directly Affect

IVF/ICSI Pregnancy Outcomes. *J. Androl.* **26**, 619–623. (doi:10.2164/jandrol.05018)

7. Kihaile P, Hirotsuru K, Kumasako Y, Misumi J, Utsunomiya T. 2003 Fertilization rates of

small-head sperm in conventional IVF and ICSI. *Arch Androl* **49**.

(doi:10.1080/01485010390219692))

8. Guzick DS *et al.* 2001 Sperm Morphology, Motility, and Concentration in Fertile and

Infertile Men. *N. Engl. J. Med.* **345**, 1388–1393. (doi:10.1056/NEJMoa003005)

9. Coetzee K, Kruge TF, Lombard CJ. 1998 Predictive value of normal sperm morphology: A

structured literature review. *Hum. Reprod. Update* **4**, 73–82.

10.   Danis RB, Samplaski MK. 2019 Sperm Morphology: History, Challenges, and Impact

on Natural and Assisted Fertility. *Curr. Urol. Rep.* **20**, 43. (doi:10.1007/s11934-019-0911-

7)

11.   Rodríguez AL, Rijsselaere T, Beek J, Vyt P, Soom A, Maes D. 2013 Boar seminal

plasma components and their relation with semen quality. *Syst Biol Reprod Med* **59**.

(doi:10.3109/19396368.2012.725120))

12.   Waberski D, Riesenbeck A, Schulze M, Weitze KF, Johnson L. 2019 Application of

preserved boar semen for artificial insemination: Past, present and future challenges.

Theriogenology **137**, 2–7. (doi:10.1016/j.theriogenology.2019.05.030)

13.   In press. Artificial Insemination (AI) Standard for pigs | AHDB. See

https://ahdb.org.uk/knowledge-library/artificial-insemination-ai-standard-for-pigs

(accessed on 3 March 2025).

14.   Roca J, Parrilla I, Rodriguez-Martinez H, Gil M, Cuello C, Vazquez J, Martinez E. 2011

Approaches Towards Efficient Use of Boar Semen in the Pig Industry. *Reprod. Domest.*

Anim. **46**, 79–83. (doi:10.1111/j.1439-0531.2011.01828.x)

15.   Tardif S, Laforest J-P, Cormier N, Bailey JL. 1999 The importance of porcine sperm

parameters on fertility in vivo. *Theriogenology* **52**, 447–459. (doi:10.1016/S0093-

691X(99)00142-9)

16.   O'Connor RE, Fonseka G, Frodsham R, Archibald AL, Lawrie M, Walling GA, Griffin DK.

2017 Isolation of subtelomeric sequences of porcine chromosomes for translocation

screening reveals errors in the pig genome assembly. *Anim. Genet.* **48**, 395–403.

(doi:10.1111/age.12548)

17.   Maes D, Lopez Rodriguez A, Rijsselaere T, Vyt P, Van Soom A. 2011 Artificial

Insemination in Pigs. In *Artificial Insemination in Farm Animals* (ed M Manafi), InTech.

(doi:10.5772/16592)

18.   Holt WV, Van Look KJW. 2004 Concepts in sperm heterogeneity, sperm selection and

sperm competition as biological foundations for laboratory tests of semen quality.

Reproduction **127**, 527–535. (doi:10.1530/rep.1.00134)

19.   Saravia F, Núñez-Martínez I, Morán JM, Soler C, Muriel A, Rodríguez-Martínez H,

Peña FJ. 2007 Differences in boar sperm head shape and dimensions recorded by

computer-assisted sperm morphometry are not related to chromatin integrity.

Theriogenology **68**, 196–203. (doi:10.1016/j.theriogenology.2007.04.052)

20.   Menkveld R, Wong WY, Lombard CJ, Wetzels AMM, Thomas CMG, Merkus HMWM,

Steegers-Theunissen RPM. 2001 Semen parameters, including WHO and strict criteria

morphology, in a fertile and subfertile population: An effort towards standardization of

in-vivo thresholds. *Hum. Reprod.* **16**. (doi:10.1093/humrep/16.6.1165))

21.   Van Der Merwe FH, Kruger TF, Oehninger SC, Lombard CJ. 2005 The Use of Semen

Parameters to Identify the Subfertile Male in the General Population. *Gynecol. Obstet.*

Invest. **59**, 86–91. (doi:10.1159/000082368)

22.   Van Waart J. 2001 Predictive value of normal sperm morphology in intrauterine

insemination (IUI): a structured literature review. *Hum. Reprod. Update* **7**, 495–500.

(doi:10.1093/humupd/7.5.495)

23.   Vicente-Fiel S, Palacín I, Santolaria P, Yániz JL. 2013 A comparative study of sperm

morphometric subpopulations in cattle, goat, sheep and pigs using a computer-assisted

fluorescence method (CASMA-F). *Anim. Reprod. Sci.* **139**, 182–189.

(doi:10.1016/j.anireprosci.2013.04.002)

24.   Yániz JL, Capistrós S, Vicente-Fiel S, Hidalgo CO, Santolaria P. 2016 A comparative

study of the morphometry of sperm head components in cattle, sheep, and pigs with a

computer-assisted fluorescence method. *Asian J. Androl.* **18**, 840–843.

(doi:10.4103/1008-682X.186877)

25.   Ostermeier GC, Sargeant GA, Yandell BS, Evenson DP, Parrish JJ. 2001 Relationship of

Bull Fertility to Sperm Nuclear Shape. *J. Androl.* **22**, 595–603. (doi:10.1002/j.1939-

4640.2001.tb02219.x)

26.   Genualdo V, Turri F, Pizzi F, Castiglioni B, Marletta D, Iannuzzi A. 2021 Sperm Nuclei

Analysis and Nuclear Organization of a Fertile Boar–Pig Hybrid by 2D FISH on Both Total

and Motile Sperm Fractions. *Animals* **11**, 738. (doi:10.3390/ani11030738)

27.   Hirai M, Boersma A, Hoeflich A, Wolf E, Föll J, Aumüller R, Braun J. 2001 Objectively

Measured Sperm Motility and Sperm Head Morphometry in Boars (Sus scrofa): Relation

to Fertility and Seminal Plasma Growth Factors. *J. Androl.* **22**, 104–110.

(doi:10.1002/j.1939-4640.2001.tb02159.x)

28.   Barquero V, Soler C, Sevilla F, Calderón-Calderón J, Valverde A. 2021 A Bayesian

analysis of boar spermatozoa kinematics and head morphometrics and their relationship

with litter size fertility variables. *Reprod. Domest. Anim.* **56**, 1024–1033.

(doi:10.1111/rda.13946)

29.   Rathje CC *et al.* 2019 Differential Sperm Motility Mediates the Sex Ratio Drive

Shaping Mouse Sex Chromosome Evolution. *Curr. Biol.* **29**, 3692-3698.e4.

(doi:10.1016/j.cub.2019.09.031)

30.   BM S. 2019 A high-throughput method for unbiased quantitation and categorization

of nuclear morphology. *Biol Reprod* **100**. (doi:10.1093/biolre/ioz013))

31.   Martinez G *et al.* 2022 Oligogenic heterozygous inheritance of sperm abnormalities

in mouse. *eLife* **11**, e75373. (doi:10.7554/eLife.75373)

32.   Stark-Dykema ER, Dulka EA, Gerlinger ER, Mueller JL. 2022 X-linked palindromic gene

families 4930567H17Rik and Mageb5 are dispensable for male mouse fertility. *Sci. Rep.*

12, 8554. (doi:10.1038/s41598-022-12433-9)

33.   Skinner BM. 2022 Nuclear Morphology Analysis 2.0.0: Improved image analysis

software for measuring nuclear shape. *J. Open Source Softw.* **7**, 4767.

(doi:10.21105/joss.04767)

34.   Hancock J, Hovell G. 1959 The collection of boar semen. *Vet. Rec.* **71**, 664–665.

35.   In press. R: The R Project for Statistical Computing. See https://www.r-project.org/

(accessed on 28 May 2024).

36.   McInnes L, Healy J, Melville J. 2020 UMAP: Uniform Manifold Approximation and

Projection for Dimension Reduction. (doi:10.48550/arXiv.1802.03426)

37.   Konopka T. 2023 umap: Uniform Manifold Approximation and Projection.

38.   Ripley B, Venables B, Bates DM, ca 1998) KH (partial port, ca 1998) AG (partial port,

polr) DF (support functions for. 2024 MASS: Support Functions and Datasets for Venables

and Ripley's MASS.

39.   Banaszewska D, Andraszek K. 2021 Assessment of the Morphometry of Heads of

Normal Sperm and Sperm with the Dag Defect in the Semen of Duroc Boars. *J. Vet. Res.*

65, 239–244. (doi:10.2478/jvetres-2021-0019)

40.   Wickham H. 2016 *ggplot2*. Cham: Springer International Publishing.

(doi:10.1007/978-3-319-24277-4)

41.   Rathje CC *et al.* 2019 A Conserved Requirement for Fbxo7 During Male Germ Cell

Cytoplasmic Remodeling. *Front. Physiol.* **10**, 1278. (doi:10.3389/fphys.2019.01278)

42.   Hancock JL. 1956 The Morphology of Boar Spermatozoa. *J. R. Microsc. Soc.* **76**, 84–

97. (doi:10.1111/j.1365-2818.1956.tb00443.x)

43.   Kondracki S, Bonaszewska D, Mielnicka C. 2005 The effect of age on the

morphometric sperm traits of domestic pigs (Sus scrofa domestica). *Cell. Mol. Biol. Lett.*

10, 3–13.

44.   Czubaszek M, Andraszek K, Banaszewska D, Walczak-Jędrzejowska R. 2019 The effect

of the staining technique on morphological and morphometric parameters of boar

sperm. *PloS One* **14**, e0214243. (doi:10.1371/journal.pone.0214243)

45.   Kawarasaki T, Enya S, Otake M. 2022 Sperm Characteristics in Microminipigs. *Vivo*

Athens Greece **36**, 731–737. (doi:10.21873/invivo.12759)

46.   Rousso D, Kourtis A, Mavromatidis G, Gkoutzioulis F, Makedos G, Panidis D. 2002

Pyriform Head: A Frequent but Little-Studied Morphological Abnormality of Sperm. *Arch.*

Androl. **48**, 267–272. (doi:10.1080/01485010290031574)

47.   Kawarsky SJ, Stubbings RB, Basrur PK, King WA. 1995 Pyriform bovine spermatozoa

and fertilization potential. *Can. Vet. J.* **36**, 434–436.

48.   Thundathil J, Palasz AT, Mapletoft RJ, Barth AD. 1999 An investigation of the

fertilizing characteristics of pyriform-shaped bovine spermatozoa. *Anim. Reprod. Sci.* **57**,

35–50. (doi:10.1016/S0378-4320(99)00058-5)

49.   Walters AH, Eyestone WE, Saacke RG, Pearson RE, Gwazdauskas FC. 2004 Sperm

Morphology and Preparation Method Affect Bovine Embryonic Development. *J. Androl.*

25, 554–563. (doi:10.1002/j.1939-4640.2004.tb02826.x)

50.   Shojaei Saadi HA. 2013 Proteins associated with critical sperm functions and sperm

head shape are differentially expressed in morphologically abnormal bovine sperm

induced by scrotal insulation'. *J. Proteomics* **82**, 64–80. (doi:10.1016/j.jprot.2013.02.027.)

51.   Boe-Hansen GB, Fortes MRS, Satake N. 2018 Morphological defects, sperm DNA

integrity, and protamination of bovine spermatozoa. *Andrology* **6**, 627–633.

(doi:10.1111/andr.12486)

52.   Lehti MS, Kotaja N, Sironen A. 2013 KIF3A is essential for sperm tail formation and

manchette function. *Mol. Cell. Endocrinol.* **377**, 44–55. (doi:10.1016/j.mce.2013.06.030)

53.   Kondracki S, Banaszewska D, Wysokińska A, Chomicz J. 2006 Sperm morphology of

cattle and domestic pigs. *Reprod. Biol.* **6 Suppl 2**, 99–104.

54.   García-Vázquez FA, HERNÁNDEZ-CARAVACA I, MATÁS C, SORIANO-ÚBEDA C, ABRIL-

SÁNCHEZ S, IZQUIERDO-RICO MJ. 2015 Morphological study of boar sperm during their

passage through the female genital tract. *J. Reprod. Dev.* **61**, 407–413.

(doi:10.1262/jrd.2014-170)

55.   Corcuera BD, Hernández-Gil R, De Alba Romero C, Martín Rillo S. 2002 Relationship

of environment temperature and boar facilities with seminal quality. *Livest. Prod. Sci.* **74**,

55–62. (doi:10.1016/S0301-6226(01)00286-X)

56.   LI T, YAO J, ZHANG Q, LI Q, LI J, WANG X, LI W, CHEN A, YAN J. 2020 Chronic Stress

Impairs Male Spermatogenesis Function and Nectin-3 Protein Expression in the Testis.

Physiol. Res. **69**, 297–306. (doi:10.33549/physiolres.934287)

57.   Kunavongkrit A, Suriyasomboon A, Lundeheim N, Heard TW, Einarsson S. 2005

Management and sperm production of boars under differing environmental conditions.

Theriogenology **63**, 657–667. (doi:10.1016/j.theriogenology.2004.09.039)

58.   Collodel G, Moretti E, Fontani V, Rinaldi S, Aravagli L, Saragò G, Capitani S, Anichini C.

2008 Effect of emotional stress on sperm quality. *Indian J. Med. Res.* **128**, 254–261.

59.   Lopez Rodriguez A, Van Soom A, Arsenakis I, Maes D. 2017 Boar management and

semen handling factors affect the quality of boar extended semen. *Porc. Health Manag.*

3, 15. (doi:10.1186/s40813-017-0062-5)

60.   Banaszewska D, Andraszek K, Biesiada-Drzazga B. 2015 Evaluation of sperm

chromatin structure in boar semen. *Bull. Vet. Inst. Pulawy* **59**, 271–277.

(doi:10.1515/bvip-2015-0040)

61.   Aziz N, Saleh RA, Sharma RK, Lewis-Jones I, Esfandiari N, Thomas AJ, Agarwal A. 2004

Novel association between sperm reactive oxygen species production, sperm

morphological defects, and the sperm deformity index. *Fertil. Steril.* **81**, 349–354.

(doi:10.1016/j.fertnstert.2003.06.026)

62.   Nguyen HTT, Dang HNT, Nguyen TTT, Nguyen TV, Dang TC, Nguyen QHV, Le MT.

2022 Correlations between abnormalities of morphological details and DNA

fragmentation in human sperm. *Clin. Exp. Reprod. Med.* **49**, 40–48.

(doi:10.5653/cerm.2021.04777)

63.   Frankenhuis MT, Kramer MF, De Rooij DG. 1982 Spermatogenesis in the boar. *Vet. Q.*

4, 57–61. (doi:10.1080/01652176.1982.9693840)

64.   França LR, Avelar GF, Almeida FFL. 2005 Spermatogenesis and sperm transit through

the epididymis in mammals with emphasis on pigs. *Theriogenology* **63**, 300–318.

(doi:10.1016/j.theriogenology.2004.09.014)

65.   Kierszenbaum AL, Tres LL. 2004 The acrosome-acroplaxome-manchette complex and

the shaping of the spermatid head. *Arch. Histol. Cytol.* **67**, 271–284.

(doi:10.1679/aohc.67.271)

66.   Yan HHN. 2007 Ectoplasmic specialization: a friend or a foe of spermatogenesis?'.

BioEssays News Rev. Mol. Cell. Dev. Biol. **29**, 36–48. (doi:10.1002/bies.20513.)

67.   Kierszenbaum AL, Tres LL, Rivkin E, Kang-Decker N, van Deursen JMA. 2004 The

acroplaxome is the docking site of Golgi-derived myosin Va/Rab27a/b- containing

proacrosomal vesicles in wild-type and Hrb mutant mouse spermatids. *Biol. Reprod.* **70**,

1400–1410. (doi:10.1095/biolreprod.103.025346)

68.   Zakrzewski P, Rędowicz MJ, Buss F, Lenartowska M. 2020 Loss of myosin VI

expression affects acrosome/acroplaxome complex morphology during mouse

spermiogenesis†. *Biol. Reprod.* **103**, 521–533. (doi:10.1093/biolre/ioaa071)

69.   Kierszenbaum A, Rivkin E, Tres L. 2007 Molecular biology of sperm head shaping.

Soc. Reprod. Fertil. Suppl. **65**, 33–43.

70.   Yang K. 2018 Ultra-structure of the sperm head-to-tail linkage complex in the

absence of the spermatid-specific LINC component SPAG4'. *Histochem. Cell Biol.* **150**, 49–

59. (doi:10.1007/s00418-018-1668-7.)

71.   Tapia Contreras C, Hoyer-Fender S. 2019 CCDC42 Localizes to Manchette, HTCA and

633   Tail and Interacts With ODF1 and ODF2 in the Formation of the Male Germ Cell

634   Cytoskeleton. *Front. Cell Dev. Biol.* **7**.

635   72.   Calvi A, Wong ASW, Wright G, Wong ESM, Loo TH, Stewart CL, Burke B. 2015 SUN4 is

636   essential for nuclear remodeling during mammalian spermiogenesis. *Dev. Biol.* **407**, 321–

637   330. (doi:10.1016/j.ydbio.2015.09.010)

638   73.   Pasch E, Link J, Beck C, Scheuerle S, Alsheimer M. 2015 The LINC complex component

639   Sun4 plays a crucial role in sperm head formation and fertility. *Biol. Open* **4**, 1792–1802.

640   (doi:10.1242/bio.015768)

641   74.   Gao Q. 2020 The testis-specific LINC component SUN3 is essential for sperm head

642   shaping during mouse spermiogenesis'. *J. Biol. Chem.* **295**, 6289–6298.

643   (doi:10.1074/jbc.RA119.012375.)

644   75.   Shang Y *et al.* 2017 Essential role for SUN5 in anchoring sperm head to the tail. *eLife*

645   **6**, e28199. (doi:10.7554/eLife.28199)

646   76.   Yassine S, Escoffier J, Nahed RA, Pierre V, Karaouzene T, Ray PF, Arnoult C. 2015

647   Dynamics of Sun5 Localization during Spermatogenesis in Wild Type and Dpy19l2 Knock-

648   Out Mice Indicates That Sun5 Is Not Involved in Acrosome Attachment to the Nuclear

649   Envelope. *PLOS ONE* **10**, e0118698. (doi:10.1371/journal.pone.0118698)

650   77.   Guttman JA, Kimel GH, Vogl AW. 2000 Dynein and plus-end microtubule-dependent

651   motors are associated with specialized Sertoli cell junction plaques (ectoplasmic

652   specializations). *J. Cell Sci.* **113**, 2167–2176. (doi:10.1242/jcs.113.12.2167)

78.   Wen Q. 2018 Dynein 1 supports spermatid transport and spermiation during

spermatogenesis in the rat testis'. *Am. J. Physiol.-Endocrinol. Metab.* **315**, 924–948.

(doi:10.1152/ajpendo.00114.2018.)

79.   Yang Z *et al.* 2019 Dynein axonemal intermediate chain 2 plays a role in

gametogenesis by activation of Stat3. *J. Cell. Mol. Med.* **23**, 417–425.

(doi:10.1111/jcmm.13945)

80.   Zhou S *et al.* 2023 DRC3 is an assembly adapter of the nexin-dynein regulatory

complex functional components during spermatogenesis in humans and mice. *Signal*

Transduct. Target. Ther. **8**, 1–4. (doi:10.1038/s41392-022-01293-4)

81.   Inagaki M, Irie K, Ishizaki H, Tanaka-Okamoto M, Miyoshi J, Takai Y. 2006 Role of cell

adhesion molecule nectin-3 in spermatid development. *Genes Cells* **11**, 1125–1132.

(doi:10.1111/j.1365-2443.2006.01006.x)

82.   Mueller S, Rosenquist TA, Takai Y, Bronson RA, Wimmer E. 2003 Loss of Nectin-2 at

Sertoli-Spermatid Junctions Leads to Male Infertility and Correlates with Severe

Spermatozoan Head and Midpiece Malformation, Impaired Binding to the Zona Pellucida,

and Oocyte Penetration1. *Biol. Reprod.* **69**, 1330–1340.

(doi:10.1095/biolreprod.102.014670)

83.   Barquero V, Roldan ERS, Soler C, Vargas-Leitón B, Sevilla F, Camacho M, Valverde A.

2021 Relationship between Fertility Traits and Kinematics in Clusters of Boar Ejaculates.

Biology **10**, 595. (doi:10.3390/biology10070595)

84.   Gu N-H, Zhao W-L, Wang G-S, Sun F. 2019 Comparative analysis of mammalian

sperm ultrastructure reveals relationships between sperm morphology, mitochondrial

functions and motility. *Reprod. Biol. Endocrinol.* **17**, 66. (doi:10.1186/s12958-019-0510-y)

85.   Colenbrander B, Kemp B. 1990 Factors influencingsemen quality in pigs. *Biosci. Proc.*

(doi:10.1530/biosciprocs.13.008)

86.   Dyck M, Foxcroft G, Novak S, Ruiz-Sanchez A, Patterson J, Dixon W. 2011 Biological

Markers of Boar Fertility. *Reprod. Domest. Anim.* **46**, 55–58. (doi:10.1111/j.1439-

0531.2011.01837.x)

87.   Sutovsky P. 2015 New Approaches to Boar Semen Evaluation, Processing and

Improvement. *Reprod. Domest. Anim.* **50**, 11–19. (doi:10.1111/rda.12554)

88.   Jung M, Rüdiger K, Schulze M. 2015 In Vitro Measures for Assessing Boar Semen

Fertility. *Reprod. Domest. Anim.* **50**, 20–24. (doi:10.1111/rda.12533)

89.   Zuidema D, Kerns K, Sutovsky P. 2021 An Exploration of Current and Perspective

Semen Analysis and Sperm Selection for Livestock Artificial Insemination. *Anim. Open*

Access J. MDPI **11**, 3563. (doi:10.3390/ani11123563)