Sperm morphology differences associated with pig fertility

Mandawala AA[1], Skinner BM[2], Walling GA[3], Harvey KE[1*^] and Harvey SC[1*]

[1] School of Psychology and Life Sciences, Canterbury Christ Church University, Canterbury, UK

[2] University of Essex, Colchester, UK

[3] Topigs Norsvin, The Netherlands

[*] Harvey KE and Harvey SC are joint last authors

^Corresponding author:

Dr Katie Harvey

School of Psychology and Life Sciences

Canterbury Christ Church University

Canterbury

UK.

E-mail: katie.harvey@canterbury.ac.uk

Abstract

Artificial insemination is routine in commercial pig breeding, and as such, the use of high-quality

semen samples is imperative. Here, we have developed a novel, semi-automated, software-based

approach to assess pig sperm nucleus morphology in greater detail than was previously possible.

This analysis identified subtle morphological differences between samples assessed by the industry

as normal and those assessed as abnormal. 50 normal and 50 abnormal samples that were initially

categorised using manual assessment to industry standards, were investigated using this new

method, with at least 200 fixed stained sperm heads analysed in each case. Differences in sperm

nuclear morphology were observed between normal and abnormal samples; specifically, normal

samples were associated with higher mean nuclear area, a consequence of a greater head width and

a lower variability between sperm heads. This novel, unbiased and fast analysis method

demonstrates a significant difference in sperm head morphology between normal and abnormal pig

sperm and has the potential to be further developed to be used as a tool for sperm morphology

assessment both in the pig breeding industry and potentially in human assisted reproductive

technologies.

Keywords: Artificial insemination, morphology, nucleus, pig, sperm
40

1. Introduction

Male fertility is a consequence of both the number and quality of sperm [1]. In humans, many issues with male infertility are addressed using one of a range of assisted reproductive technology procedures, such as *in vitro* fertilisation and intracytoplasmic sperm injection. In agriculturally significant species such as pigs, cattle, and sheep, where a key goal is to maximise the production of meat at a low cost, male fertility is also a challenge [2]. To this end, improving reproductive traits is paramount. In such species, the critical aim is often to identify sub-fertile animals quickly and cheaply so that they can be removed from breeding schemes; boars from a nucleus herd with fertility problems have the potential to reduce litter sizes throughout the breeding population [3].

Artificial insemination (AI) is the oldest [4] and most routinely used technique in commercial animal breeding, especially in livestock species [4–10]. Over the past three decades, the use of AI has greatly benefited the pig breeding industry, particularly in Europe where over 80% of sows are bred through AI [4]. In North America, the technique is also widely implemented, especially in large farming units [9]. The principal objective of AI in the pig breeding industry is to permit the dissemination of genetics from high genetic merit boars to as many sows as possible. Without AI, more boars would be needed and hence animals of lower genetic merit would be required in breeding programmes. Moreover, the technique enables the opportunity to introduce superior genetic traits into sows whilst reducing the incidence of disease transmission, an advantage that does not exist with natural mating [11]. AI is achieved by depositing spermatozoa into the female genital tract using artificial devices and processes. The standardised method of insemination is the intra-cervical insemination technique, with the semen dose deposited in the posterior region of cervical canal [4].

In humans, semen analysis is widely used to evaluate male fertility [12] and may also be used for the determination of reproductive toxicity in therapeutic and environmental agents [13,14]. Various

physical characteristics of semen are assessed and whilst parameters such as volume, appearance,

pH, and viscosity are considered important [15], several studies have shown that good sperm

morphology is critical when determining semen quality and hence quantifying male fertility [16–21].

Generally, the cut-off values of what is considered "normal" vary, and are dependent on the fertility

clinic. However, the following benchmarks were published in the World Health Organisation's 5th

edition of "normal semen analysis": morphology (≥4% normal forms), total motility (≥40%), vitality

(≥58% live), sperm concentration (≥15,000,000 per mL) and volume (≥1.5 mL) [12,22]. To date, a

number of studies have been performed to analyse semen composition [23,24] and to establish the

relationship between sperm quality and fertility in men [25–30]. One such study used several

comparative semen analyses of fertile and infertile men, to determine the most appropriate

measurements that could be used in the determination of fertility potential in men [31]. Here, it was

established that whilst threshold values for sperm motility, concentration and morphology could be

used in the classification of males into fertile, indeterminate fertility, or sub-fertile categories, these

measures cannot be used independently for the diagnosis of male infertility [31].

In livestock species, a key contribution to successful fertilisation following AI is also the use of high-

quality semen during insemination. As such, routine assessment of semen quality is a standard

process in the animal breeding industry [32]. An issue, however, is that the mammalian ejaculate

does not contain a homogenous population of spermatozoa [33]; in general analysis of boar semen,

semen quality is considered "normal" if the frequency of abnormal sperm heads does not exceed

10%. Normality can also be assumed if the frequency of abnormalities in acrosomes, mid-pieces,

tails, or proximal cytoplasmic droplets is less than either 5% each, or 15% when combined [34].

Some morphological abnormalities are regarded as significant due to their ability to hinder

fertilisation and cause infertility [34]; some of these include decapitated spermatozoa, acrosomal

plicae (knobbed defect), nuclear vacuoles (diadem defect), short tails (tail stump), coiled tails (dag

defect) and corkscrewed mid-pieces [34].

As well as studying the morphology of whole spermatozoa, a limited number of studies have focused

on analysis of sperm nuclear morphometry in agricultural animals [35,36]. A number of

measurements of sperm nuclei have been compared in cattle, sheep, goats, and pigs using

computer-assisted sperm morphometry analysis-F (CASMA-F) [35,36]. Whilst mean numerical values

for area, perimeter, length, and width were identified for each of the four species studied, the key

findings of this study were that drying and fixation only has a minimal effect on sperm nuclear

morphometry and that variations between morphometric parameters do exist between the sperm

nuclei of each species [35]. A second study [36] used a combination of the CASMA-F method and

multivariate cluster analysis to identify subpopulations of spermatozoa within the same four species.

Based on these nuclear morphometrics, three subpopulations, namely, large, small-elongated and

small-round were identified. Whilst it has previously been shown that sperm shape differed between

high and low fertility bulls [37], such observations are limited in pigs [38]. Here, we demonstrate the

use of a semi-automated software-based approach to assess sperm nucleus morphometrics in pig

semen samples deemed as normal and abnormal.

2. Material and Methods

2.1 Semen collection

Fresh ejaculated sperm samples from boars were collected using the 'gloved hand method' [47], by

trained staff at JSR Genetics Ltd. Samples were stored in Duragen extender, supplemented with no

less than: 500 IU per ml streptomycin; 500 IU per ml penicillin; 150 mg per ml lincomycin; and 300

mg per ml spectinomycin, diluted to 2.3 billion sperm per dose. Samples were stored at 17°C and

were prepared within two days following collection. No ethical approval was required for this study

as all semen doses used were collected as part of JSR Genetics Ltd.'s standard commercial

procedures. Semen samples identified as abnormal would have been discarded, those identified as

normal would have been used commercially.

2.2 Sample preparation

Prior to preparation of samples for this study, semen samples were identified as either normal or

abnormal using computer assisted sperm analysis (CASA) followed by manual assessment.

Specifically, samples that had a normal morphology score of above 70% (obtained from CASA) and a

motility score of above 4 (motility was graded from 1 to 5, 1 being dead and 5 being excellent) were

graded as normal and those falling below these criteria were graded as abnormal.

50 normal and 50 abnormal samples were used in this study. 2mL of each semen sample was

centrifuged at 300g for 5 minutes at 17°C. The supernatant was discarded, the pellet was re-

suspended in 1.5mL of fixative solution (100% methanol and 100% acetic acid, added dropwise at a

3:1 ratio) and centrifuged at 300g for 5 minutes at 17°C. The supernatant was discarded, and the

pellet was re-suspended in 1.5mL of fixative solution. 10μL of each sample was then dropped onto

the centre of the surface of a labelled (sample ID, date), steam-warmed slide, immediately followed

by 10μL of fixative solution. Subsequently, slides were air-dried for two minutes before one drop of

fluorescent DAPI (4',6-diamidino-2-phenylindole) was added to the centre of the slide. Prepared

slides were air-dried in the dark, for at least 20 minutes prior to microscopy.

2.3 Image acquisition

An Olympus IX83 inverted fluorescence microscope equipped with CellSens Dimension version 1.9

(expandable imaging software for Life Science microscopy) was used for image capturing. A

minimum of 200 nuclei were imaged (at 1000x magnification) per sample.

2.4 Data analysis

Images were analysed using the ImageJ plugin 'Nuclear Morphology Analysis' (NMA) (see [43]),

freely available under the GNU General Public License version 3 from

https://bitbucket.org/bmskinner/nuclear_morphology/wiki/Home). The analysis software enables

automated recognition of round or asymmetric nuclei within an image of interest, and subsequent

morphological analysis of these nuclei. Initially developed for the analysis of mouse sperm [43], we

adapted the feature recognition to analyse pig sperm. The software generates a range of measures;

in this study, sperm heads were measured for: Area; Perimeter; Bounding Height and Bounding

Width, the dimensions of the rectangle perfectly enclosing the nucleus when vertically oriented;

Circularity, a measure between 0 and 1 indicating how circular the heads are, with 1 indicating a

perfect circle; Minimum Width across the centre of mass; Aspect, the ratio produced from height

divided by width; and Variability, a per-nucleus measure calculated as the square root of the sum-of-

squares difference at each index between the nuclear profile and the dataset median profile,

normalised to the length of the median profile.

Pig sperm are (mostly) symmetrical about the anterior-posterior axis. Without a distinctive hook, as

in mouse sperm, to act as a reference point, the tail attachment point was chosen to anchor the

angle profiles generated and orient nuclei. The tail attachment point is characterised by a 'dimple' in

the nucleus (Figure 1A, point 1). For consistent alignment of the nuclei, we placed the tail

attachment region directly below the centre of mass of the nucleus. The software output prior to

statistical analyses is summarised in Figure 1. This indicates how we generate an angle profile for a

given sample and how a consensus sperm head shape for the population is produced. These

refolded sperm heads are then used both to trace any identified abnormalities on a particular

segment of the refolded nucleus back to the angle profile, and to calculate a range of sperm head

parameters for analysis. Analysis was carried out using software version 1.13.5, and initial

correlation analysis was used to identify redundant variables. Non-redundant variables were then

further analysed; for initial comparisons between normal and abnormal animals, means of each

variable were compared by ANOVA in R version 3.4.4 [48], with the normality of residual distribution

assessed using scatter and quantile-quantile plots.

For the cluster analysis, samples were randomly allocated into two groups; a training group of 30

normal and 30 abnormal animals and a test group of 20 normal and 20 abnormal animals. Sperm

heads from the training group were used to determine if there were sperm of detectably different

types. This was done by cluster analysis, with Ward linkage, squared Euclidean distance and

standardised variables in Mintab v17. The representation of sperm heads from normal and abnormal

animals within in each identified cluster was then compared to that expected by chance. The

training group was then used to investigate if identified clusters could be used to predict fertility

status. This was done by using the mean cluster value from the training set to define the starting

partitions for a K-means clustering.

A ROC analysis was then used to test the diagnostic capacity of these clusters in predicting whether

a given semen sample would have a normal or abnormal semen profile based on the proportion of

sperm that fall into the clusters 1 and 3. ROC curves were plotted in R version 3.6.0 via the pROC

package [49] using the predictions derived from the cluster output.

3. Results

Analysis of sperm nuclear morphology from 50 normal and 50 abnormal animals yielded measures

from 11,534 and 11,326 nuclei, respectively. Correlation analysis of measured sperm head

characters indicated that, as expected, many of the measures were highly correlated (Table S1), and

further analysis was undertaken using measures of Area, Circularity, Variability, Bounding Height and

Bounding Width. Given the orientation of the sperm heads, Bounding Height and Width are

subsequently referred to as Maximum Height and Width.

Comparisons between normal and abnormal animals indicated that sperm heads in these groups

differ, but that there is a large amount of variation between individuals (Figure 2, and Figures S1-4).

Analysis of mean trait values from each animal indicates that sperm heads from fertile animals have

a higher overall area (Area, $F_{1,98}$ = 34.55, $p < 0.001$), are wider (Width, $F_{1,98}$ = 11.58, $p = 0.001$), longer

(Height, $F_{1,98}$ = 21.68, $p < 0.001$) and are less variable in shape (Variability, $F_{1,98}$ = 24.98, $p < 0.001$)

than those from abnormal animals. At this level of analysis, no difference between the sperm heads

of normal and abnormal animals in Circularity was detected (Circularity, $F_{1,98}$ = 1.80, $p = 0.18$).

We hypothesised that individual sperm samples would contain different nucleus types, and that this

may allow the identification of abnormal animals. For example, a certain sperm head type might

only be found in abnormal animals or might be overrepresented within such animals. Such

differences might represent specific features within these abnormal animals, which could also be

used to investigate why their fertility was impaired, *i.e.,* this could be informative about the biology,

or the mechanism. Additionally, the presence of differences of this type might allow the fertility of

an animal to be predicted, thereby allowing the automated assessment of fertility. We therefore,

randomly allocated samples into two groups: a training group of 30 normal and 30 abnormal animals

in which we investigated the presence, or not, of different nuclei types; and a test group of 20

normal and 20 abnormal animals in which we investigated the ability of this type of analysis to

predict fertility status.

In the training group, cluster analysis (Ward linkage, squared Euclidean distance and standardised

variables) of 6,924 sperm heads from normal animals and 6,684 sperm heads from abnormal

animals supported the existence of three clusters (Figure 3). The identified clusters showed different

membership for the normal and abnormal sperm heads, with sperm heads from normal animals

overrepresented in the largest cluster and underrepresented in the smallest cluster.

Comparison of these clusters indicates that the cluster in which sperm heads from normal animals

are overrepresented is characterised by sperm heads with a low variability and a high area (Figure

4). In contrast, the cluster in which abnormal animals are overrepresented is characterised by sperm

heads with a lower area and a high variability (Figure 4).

The three clusters identified in the training group were also recovered in the test group (Figure S5),

with analysis of sperm head morphology showing the same differences between clusters (data not

shown). As in the training group, sperm heads from normal animals were overrepresented in cluster

1 and underrepresented in cluster 3 (Figure S5). This supports the idea that the frequency of certain

sperm head morphology types can be used to predict fertility status. Several approaches were

investigated to test this. Firstly, we used the mean trait values for the three clusters identified in the

training group to define the initial partitions for a K-means clustering of the sperm heads from the

test group of 20 normal and 20 abnormal animals. The proportion of nuclei from each animal that

was allocated to each cluster was then determined and compared to the proportions determined

from the training set (Figure 5). This indicates that this method does accurately predict fertility in

some animals, but that no scheme would correctly identify status for all animals. That is, if the aim

was to exclude any animal where abnormality was suggested, then 8 of the 20 normal animals and 6

of the 20 abnormal animals would be retained (Table S2). Conversely, when the aim was to retain

samples that were identified as normal through cluster membership, 13 of the 20 normal samples

and 6 of the 20 abnormal samples would be retained (Table S2). Similarly, attempts to predict

fertility using other approaches – for example defining thresholds based on rates of variability within

samples – also resulted in the inclusion of abnormal animals or the exclusion of normal ones. To

formally assess the diagnostic capacity of cluster membership, a receiver operating characteristic

(ROC) analysis was also performed using proportions in Table S1 as the range of cut-off values for

the predictor variables. This identified that using the proportion of sperm heads that do not fall into

cluster 3 results in the largest area under the curve (0.723, with 95% confidence interval of 0.537 –

0.868; Figure S6).

4. Discussion

Over the last decade, AI has become commonplace in the pig breeding industry, and as such it is an economical imperative to identify boars with prime fertility. To this end, various computer technologies have been developed, or adapted, to allow quantitative analysis of boar sperm characteristics. Examples of these are the various CASA systems that have been developed and improved over almost four decades that are now widely used in semen handling centres and spermatology laboratories. Currently, the Sperm Class Analyser (SCA) is considered to be the gold standard in automated sperm morphology analysis systems [39]; SCA has been shown to provide accurate measurements of head, mid-piece and tail morphometry in several mammalian species including humans [40], horses [41] and goats [42]. The absence of standardised morphometric parameters and corresponding values which could be used to identify both normal and abnormal sperm, however, is a limitation which is common to most species [39].

We previously developed Nuclear Morphology Analysis software for rapid and accurate assessment of nuclear morphology in mouse lines [43]. In contrast to other morphometric approaches, such as elliptic Fourier analysis [37], or Procrustes-based geometrics [44], our analysis can be run rapidly on many thousands of nuclei, using automatic detection of landmarks and semi-landmarks within the nucleus, and the results map cleanly back to the biological structure of the nucleus. Here, we have extended the capability of the software to recognise pig sperm for the first time, and have used this to compare sperm head morphometrics between semen samples previously identified as normal or abnormal by the industry. Results from these analyses support the hypothesis that subtle morphometric differences can be observed between sperm heads from normal and abnormal samples. The ability of the software to distinguish subtle morphological differences in pig sperm demonstrates the utility of this approach for other spatulate nuclei, as commonly found in other mammalian species, including humans [45].

Sperm head morphology in pigs is usually assessed as being either normal or abnormal based on

CASA data and manual assessment of morphology undertaken by the industry. These analyses have

identified differences between the sperm heads of these groups, with sperm heads from normal

samples having a higher overall area – a consequence of greater width and height – and being less

variable in shape than those from abnormal samples (Figures 2 and S1-4).

Our analysis also identifies three clusters of morphology types (Figure 3). These clusters group sperm

heads that have a low variability and a high area (cluster 1), those that are long and narrow (cluster

2), and those that have a low area and a high variability (cluster 3) (Figure 4). This mirrors the

clusters identified by previous work on sperm head morphology in pigs that used measures of area,

height (or length) and width [35,36,46]. Here however, our analysis also indicates that the variability

differs between these groups (Figure 4). We hypothesised that a certain type of sperm head might

only be present, or be overrepresented, in semen from abnormal samples. Whilst it is expected that

such differences in sperm head morphology would have an impact on the spermatozoa's fertilising

potential [46], there is limited research on which precise morphological parameters can in fact

impact fertility. Comparison of cluster membership indicated that sperm heads from normal samples

are overrepresented in cluster 1 and underrepresented in cluster 3, *i.e.*, abnormal samples have a

high incidence of sperm heads that have a low area and a high variability (Figure 4).

Given that automated approaches to identify abnormal semen samples would be of value in pig

production, we sought to determine if the clusters we identified could be used as a predictive tool

for semen analysis. Whilst grouping based on the previously identified clusters was able to

successfully distinguish some samples as normal or abnormal, it was found that cluster membership

alone could not accurately predict fertility in all 40 samples in the test group. To further test the

diagnostic capacity of these clusters, ROC analysis (Figure S6) was used to identify how suitable

cluster membership would be as a predictor variable in identifying whether a given semen sample

has a normal or abnormal profile. This analysis revealed that if the proportion of sperm heads that

do not fall into cluster 3 was used a predictor variable, there was a 72.3% chance of correctly

distinguishing a normal semen sample from an abnormal sample, with an AUC of between 0.7 and

0.8 generally being considered to be acceptable.

5. Conclusions

Here we show that morphometric analysis of pig sperm using NMA software distinguishes

morphologically distinct populations of nuclei. This analysis identifies differences in sperm nucleus

morphology between animals deemed to be commercially acceptable and unacceptable using

manual assessment to industry standards. These findings suggest that the use of NMA software

provides a high-throughput and more accurate method to identify pig sperm with a higher fertilising

potential. This approach may therefore have immediate utility for the pig breeding industry, and

longer-term implications for human assisted reproductive technologies.

References

[1]   Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HWG, Behre HM, et al. World Health
Organization reference values for human semen characteristics. Human Reproduction
Update 2009;16:231–45. https://doi.org/10.1093/humupd/dmp048.
[2]   Tardif S, Laforest JP, Cormier N, Bailey JL. The importance of porcine sperm parameters on
fertility in vivo. Theriogenology 1999;52:447–59. https://doi.org/10.1016/S0093-
691X(99)00142-9.
[3]   O'Connor RE, Fonseka G, Frodsham R, Archibald AL, Lawrie M, Walling GA, et al. Isolation of
subtelomeric sequences of porcine chromosomes for translocation screening reveals errors in
the pig genome assembly. Animal Genetics 2017;48:395–403.
https://doi.org/10.1111/age.12548.
[4]   Roca J, Vázquez JM, Gil MA, Cuello C, Parrilla I, Martínez EA. Challenges in pig artificial
insemination. Reproduction in Domestic Animals 2006;41 Suppl 2:43–53.
https://doi.org/10.1111/j.1439-0531.2006.00768.x.
[5]   Dziuk PJ, Henshaw G. Fertility of boar semen artificially inseminated following in vitro
Storage. Journal of Animal Science 1958;17:554–8. https://doi.org/10.2527/jas1958.173554x.
[6]   Polge C, Day B, Groves T. Synchronisation of ovulation and artificial insemination in pigs.
Veterinary Record 1968;83:136–42. https://doi.org/10.1136/vr.83.6.136.

[7]   Johnson LA, Aalbers JG, Willems CMT, Sybesma W. Use of boar spermatozoa for artificial
insemination. Fertilizing capacity of fresh and frozen spermatozoa in sows on 36 Farms.
Journal of Animal Science 1981;52:1130–6. https://doi.org/10.2527/jas1981.5251130x.

[8]   Singleton WL. State of the art in artificial insemination of pigs in the United States.
Theriogenology 2001;56:1305–10. https://doi.org/10.1016/S0093-691X(01)00631-8.

[9]   Gerrits RJ, Lunney JK, Johnson LA, Pursel VG, Kraeling RR, Rohrer GA, et al. Perspectives for
artificial insemination and genomics to improve global swine populations. Theriogenology
2005;63:283–99. https://doi.org/10.1016/j.theriogenology.2004.09.013.

[10]   Feitsma H. Artificial insemination in pigs , research and developments in The Netherlands, a
review. Acta Scientiae Veterinariae 2009;37:61–72.

[11]   Maes D, Nauwynck H, Rijsselaere T, Mateusen B, Vyt P, de Kruif A, et al. Diseases in swine
transmitted by artificial insemination: an overview. Theriogenology 2008;70:1337–45.
https://doi.org/10.1016/j.theriogenology.2008.06.018.

[12]   Rowe P, Comhaire F, Hargreave T, Mellows H. WHO manual for the standardized
investigation and diagnosis of the infertile couple. Cambridge, England: Cambridge University
Press 1993.

[13]   U.S. Environmental Protection Agency. Guidelines for Reproductive toxicity risk assessment.
Environmental Protection Agency 1996;61:56274–322.

[14]   Apostoli P, Kiss P, Porru S, Bonde JP, Vanhoorne M. Male reproductive toxicity of lead in
animals and humans. Occup Environ Med 1998;55:364–74.

[15]   Maree L, Du Plessis SS, Menkveld R, Van Der Horst G. Morphometric dimensions of the
human sperm head depend on the staining method used. Human Reproduction
2010;25:1369–82. https://doi.org/10.1093/humrep/deq075.

[16]   MacLeod J, Gold RZ. The male factor in fertility and infertility. Fertility and Sterility
1951;2:394–414. https://doi.org/http://dx.doi.org/10.1016/S0015-0282(16)30661-6.

[17]   Hartman CG, Schoenfeld C, Copeland E. Individualism in the semen picture of infertile men.
Fertility and Sterility 1964;15:231–53. https://doi.org/http://dx.doi.org/10.1016/S0015-
0282(16)35220-7.

[18]   Eliasson R. Standards for investigation of human semen. Andrologia 1971;3:49–64.
https://doi.org/10.1111/j.1439-0272.1971.tb01125.x.

[19]   Menkveld R, Kruger TF. Evaluation of sperm morphology by light microscopy. Human
Spermatozoa in Assisted Reproduction 1996:89–107.

[20]   Coetzee K, Kruge TF, Lombard CJ. Predictive value of normal sperm morphology: a structured
literature review. Human Reproduction Update 1998;4:73–82.
https://doi.org/10.1093/humupd/4.1.73.

[21]   Auger J, Jouannet P, Eustache F. Another look at human sperm morphology. Human
Reproduction 2016;31:10–23. https://doi.org/10.1093/humrep/dev251.

[22]   WHO. Examination and processing of human semen. World Health Organization 2010;5th
editio:286. https://doi.org/10.1038/aja.2008.57.

[23]   Huggins C, Scott WW, Heinen JH. Chemical composition of human semen and of the
secretions of the prostate and seminal vesicles. American Journal of Physiology -- Legacy
Content 1942;136:467 LP – 473.

[24]   Owen DH. A review of the physical and chemical properties of human semen and the
formulation of a semen simulant. Journal of Andrology 2005;26:459–69.
https://doi.org/10.2164/jandrol.04104.

[25]   Paz GF, Sofer A, Homonnai ZT, Kraicer PF. Human semen analysis: seminal plasma and
prostatic fluid compositions and their interrelations with sperm quality. International Journal
of Fertility 1977;22:140–7.

[26]   Overstreet JW, Katz DF. Semen analysis. The Urologic Clinics of North America 1987;14:441–
9.

[27] Martin RH, Rademaker A. The relationship between sperm chromosomal abnormalities and
sperm morphology in humans. Mutation Research Letters 1988;207:159–64.
https://doi.org/http://dx.doi.org/10.1016/0165-7992(88)90081-4.
[28] Perreault SD, Aitken RJ, Baker HWG, Evenson DP, Huszar G, Irvine DS, et al. Integrating new
tests of sperm genetic integrity into semen analysis: breakout group discussion. In: Robaire B,
Hales BF, editors. Advances in Male Mediated Developmental Toxicity, Boston, MA: Springer
US; 2003, p. 253–68. https://doi.org/10.1007/978-1-4419-9190-4_23.
[29] Jung A, Schuppe H-C. Influence of genital heat stress on semen quality in humans. Andrologia
2007;39:203–15. https://doi.org/10.1111/j.1439-0272.2007.00794.x.
[30] Agarwal A, Deepinder F, Sharma RK, Ranga G, Li J. Effect of cell phone usage on semen
analysis in men attending infertility clinic: an observational study. Fertility and Sterility
2008;89:124–8. https://doi.org/http://dx.doi.org/10.1016/j.fertnstert.2007.01.166.
[31] Guzick D, Overstreet J, Factor-Litvak P, Brazil C, Nakajima S, Coutifaris C, et al. Sperm
morphology, motility, and concentration in fertile and infertile men. The New England Journal
of Medicine 2001;345:1388–93.
[32] López Rodríguez A, Rijsselaere T, Beek J, Vyt P, Van Soom A, Maes D. Boar seminal plasma
components and their relation with semen quality. Systems Biology in Reproductive Medicine
2013;59:5–12. https://doi.org/10.3109/19396368.2012.725120.
[33] Holt W V., Van Look KJW. Concepts in sperm heterogeneity, sperm selection and sperm
competition as biological foundations for laboratory test of semen quality. Reproduction
2004;127:527–35. https://doi.org/10.1530/rep.1.00134.
[34] Saravia F, Nunez-Martınez I, Moran J, Soler C, Murielb A, Rodrıguez-Martıneza H, et al.
Differences in boar sperm head shape and dimensions recorded by computer-assisted sperm
morphometry are not related to chromatin integrity. Theriogenology 2007;68:196–203.
https://doi.org/10.1016/j.theriogenology.2007.04.052.
[35] Vicente-Fiel S, Palacín I, Santolaria P, Hidalgo CO, Silvestre MA, Arrebola F, et al. A
comparative study of the sperm nuclear morphometry in cattle, goat, sheep, and pigs using a
new computer-assisted method (CASMA-F). Theriogenology 2013;79:436–42.
https://doi.org/10.1016/j.theriogenology.2012.10.015.
[36] Vicente-Fiel S, Palacín I, Santolaria P, Yániz JL. A comparative study of sperm morphometric
subpopulations in cattle, goat, sheep, and pigs using a computer-assisted fluorescence
method (CASMA-F). Animal Reproduction Science 2013;139:182–9.
https://doi.org/10.1016/j.anireprosci.2013.04.002.
[37] Ostermeier GC, Sargeant GA, Yandell BS, Evenson DP, Parrish JJ. Relationship of bull fertility
to sperm nuclear shape. Journal of Andrology 2001;22:595–603.
https://doi.org/10.1002/j.1939-4640.2001.tb02219.x.
[38] Hirai M, Boersma A, Hoeflich A, Wolf E, Foll J, Aumuller R, et al. Objectively measured sperm
motility and sperm head morphometry in boars (*Sus scrofa*): Relation to fertility and seminal
plasma growth factors. Journal of Andrology 2001;22:104–10.
[39] Horst G Van Der. Automated sperm morphology analysis 2015;125:125-128
[40] Soler C, De Monserrat JJ, Gutiérrez R, Nuñez J, Nuñez M, Sancho M, et al. Use of the sperm-
class analyser® for objective assessment of human sperm morphology. International Journal
of Andrology 2003;26:262–70. https://doi.org/10.1046/j.1365-2605.2003.00422.x.
[41] Hidalgo M, Rodriguez I, Dorado J, Sanz J, Soler C. Effect of sample size and staining methods
on stallion sperm morphometry by the Sperm Class Analyzer. Veterinarni Medicina
2005;50:24–32.
[42] Hidalgo M, Rodríguez I, Dorado J. Influence of staining and sampling procedures on goat
sperm morphometry using the Sperm Class Analyzer. Theriogenology 2006;66:996–1003.
https://doi.org/10.1016/j.theriogenology.2006.02.039.

[43] Skinner BM, Rathje CC, Bacon J, Johnson EEP, Larson EL, Kopania EEK, et al. A high-
throughput method for unbiased quantitation and categorisation of nuclear morphology.
Biology of Reproduction 2019; 100:1250-1260. https://doi.org/10.1093/biolre/ioz013
[44] Varea Sánchez M, Bastir M, Roldan ERS. Geometric morphometrics of rodent sperm head
shape. PLOS ONE 2013;8:e80607.
[45] Skinner BM, Johnson EEP. Nuclear morphologies: their diversity and functional relevance.
Chromosoma 2017;126:195–212. https://doi.org/10.1007/s00412-016-0614-5.
[46] Curry M. Cryopreservation of semen from domestic livestock. Reviews of Reproduction
2000;5:46–52. https://doi.org/10.1007/978-1-4939-2193-5_10.
[47] King GJ, Macpherson JW. A comparison of two methods for boar semen collection. Journal of
Animal Science 1973;36:563–5. https://doi.org/10.2527/jas1973.363563x.
[48] R Core Team. R: A Language and Environment for Statistical Computing 2019.
[49] Robin, X. et al. (2011) 'pROC: an open-source package for R and S+ to analyze and compare
ROC curves', BMC Bioinformatics, 8, pp. 12–77. doi: 10.1007/s00134-009-1641-y.
447

Acknowledgments

We would like to thank JSR Genetics Ltd. for supplying the samples that were used in this study.

Authors' contributions

Conceptualisation, BMS and KH; Methodology, BMS, KH, SH; Software and Validation, BMS;

Investigation, AM; Data Curation and Formal Analysis, AM, SH; Visualisation, SH, BMS; Supervision

and Project Administration, KH, SH; Writing - Original Draft, AM, KH; Writing - Review and Editing,

BMS, KH, AM, SH; Resources, KH, GW. All authors gave final approval for publication.

Competing interests

None of the authors of this paper has a financial or personal relationship with other people or

organisations that could inappropriately influence or bias the content of the paper.

Funding

BMS was supported by the Leverhulme Trust (grant RPG337) and the Biotechnology and Biological

Sciences Research Council (BBSRC, grant BB/N000129/1).

Figure 1: A representative summary of analysis workflow. (A) DAPI stained nucleus from a normal

boar sample, captured using an Olympus IX83 fluorescence microscope with pre-installed CellSens

software at 1,000x magnification. The software measures the interior angles along the perimeter of

the nucleus as represented by point 1-6. (B) Schematic showing how these angles generate a profile.

The figure shows the median and interquartile rages for one normal sperm sample and has been

segmented at local maxima and minima. (C) The consensus normal pig sperm head shape, showing

positions of profile segments in the nucleus.

Figure 2: Comparison of sperm head area between 50 normal (blue) and 50 abnormal (red) samples.

(A) Area of 11,534 sperm heads from normal samples (blue) and 11,326 sperm heads from abnormal

samples (red).

Figure 3: Analysis of the training group identifies distinct morphological clusters. (A) Cluster analysis

of sperm from 30 normal and 30 abnormal individuals using measures of Area, Circularity,

Variability, Maximum Height and Maximum Width by Ward linkage using squared Euclidean distance

and standardised variables. This identifies three distinct clusters. Shown below each cluster is the

percentage of sperm heads within that cluster that are from fertile animals and the difference in

representation from that expected by chance (*i.e,.* a positive value indicates that fertile nuclei are

overrepresented in a cluster and *vice versa* for a negative value). (B) The consensus shapes of the

three clusters demonstrate the separation of nuclei on size and shape. Cluster 1 contains most of the

normal sperm; cluster 2 has equal representation of normal and abnormal sperm; cluster 3

predominantly contains abnormal sperm. The increasing compaction of the nuclei is readily

apparent overlaying consensus nuclei from clusters 1 and 3.

Figure 4: Sperm heads from the three clusters detected are morphologically distinct. Comparison of

(A) Area, with 1>2>3 ($p < 0.001$ by pairwise *post hoc* test), (B) Circularity, with 3>1>2, (C) Variability,

with 3>2>1, (D) Maximum Height, with 2>1>3, and (E) Maximum width, with 1>3>2, for the three

clusters.

Figure 5: Representation of how cluster membership can be used to predict the status of a semen

profile. The proportion of sperm heads from each sample in the test group that fall into (A) cluster 1

and (B) cluster 3 are illustrated. In both (A) and (B), the 95% confidence intervals from the training

set are represented by the dark shaded rectangles and the areas between these confidence intervals

and either 0 or 1 are represented by the light shaded rectangles. That is, blue points found in the

regions falling onto the red rectangles represent normal samples that have been incorrectly

classified as abnormal based on cluster membership and *vice versa* for red points falling into the

regions defined by blue rectangles.

Supplementary data

Supplementary tables

Table S1. Correlation matrix of measured parameters from the total dataset of 50 normal and 50

abnormal samples. Shown are the Pearson correlation values, all are significant at p < 0.001 except

for the correlation between Circularity and Variability where p = 0.10. Shaded boxes represent

Pearson correlation values > 0.8, and this was used to identify redundant variables.

	Area	Perimeter	Maximum feret	Circularity	Minimum diameter	Aspect	Variability	Maximum Height
Perimeter	0.93							
Maximum feret	0.78	0.93						
Circularity	-0.12	-0.47	-0.64					
Minimum diameter	0.64	0.35	0.09	0.58				
Aspect	0.16	0.47	0.70	-0.88	-0.62			
Variability	-0.29	-0.26	-0.16	-0.01	-0.05	-0.15		
Maximum Height	0.78	0.92	1.00	-0.64	0.07	0.71	-0.17	
Maximum Width	0.61	0.34	0.09	0.54	0.96	-0.64	0.08	0.07

Table S2: Cluster membership as a predictive tool. Shown are the proportions of sperm heads falling

into clusters 1 and 3 for each of the test individuals. Predictions are made based on comparison to the

relevant 95% confidence interval derived from the training set, e.g., to be considered normal based

on the proportion of sperm heads in cluster 1 an individual would need to have a proportion higher

than the lower confidence limit. Prediction (exclusive) indicates a scenario where the aim is to exclude

the maximum number of samples and hence a sample is excluded if either prediction indicates

'abnormal'. Prediction (inclusive) indicates a scenario where the aim is to include the maximum

number of samples and hence a sample is included if either prediction indicates 'normal'.

Sample	Type	Proportion in cluster 1	Prediction from cluster 1	Proportion in cluster 3	Prediction from cluster 3	Prediction (exclusive)	Prediction (inclusive)
N1	N	0.11	AB	0.27	AB		
N2	N	0.22	AB	0.44	AB		
N4	N	0.94	N	0.04	N	N	N
N5	N	0.60	N	0.32	AB		N
N7	N	0.48		0.38	AB		
N13	N	0.11	AB	0.77	AB		
N17	N	0.04	AB	0.02	N		N
N18	N	0.47		0.16			
N21	N	0.87	N	0.01	N	N	N
N22	N	0.88	N	0.02	N	N	N
N24	N	0.64	N	0.02	N	N	N
N29	N	0.32	AB	0.02	N		N
N34	N	0.04	AB	0.41	AB		
N36	N	0.68	N	0.10	N	N	N
N41	N	0.40		0.14	N	N	N
N42	N	0.78	N	0.09	N	N	N
N44	N	0.67	N	0.17		N	N
N46	N	0.35	AB	0.27	AB		
N48	N	0.23	AB	0.13	N		N
N49	N	0.39	AB	0.10	N		N
A4	AB	0.21	AB	0.48	AB		
A6	AB	0.15	AB	0.81	AB		
A8	AB	0.00	AB	0.38	AB		
A9	AB	0.08	AB	0.48	AB		
A14	AB	0.11	AB	0.77	AB		
A15	AB	0.61	N	0.22		N	N
A16	AB	0.06	AB	0.87	AB		
A19	AB	0.27	AB	0.58	AB		
A20	AB	0.13	AB	0.76	AB		
A22	AB	0.15	AB	0.58	AB		
A29	AB	0.78	N	0.04	N	N	N
A33	AB	0.04	AB	0.53	AB		
A34	AB	0.00	AB	0.57	AB		
A35	AB	0.06	AB	0.19			
A36	AB	0.70	N	0.03	N	N	N
A38	AB	0.47		0.04	N	N	N
A39	AB	0.47		0.05	N	N	N
A40	AB	0.48		0.07	N	N	N
A43	AB	0.31	AB	0.18			
A49	AB	0.11	AB	0.23			

526   *Supplementary figures*

528   **Figure S1**: Comparison of sperm head Circularity between 50 normal (blue) and 50 abnormal (red)

529   samples. Representation of the difference in Circularity between 11,534 normal nuclei (blue) and

530   11,326 abnormal nuclei (red). Means with the same letter are not significantly different at the 95%

531   confidence interval.

Figure S2: Comparison of sperm head Variability between 50 normal (blue) and 50 abnormal (red)

535 samples. Representation of the difference in Variability between 11,534 normal nuclei (blue) and

536 11,326 abnormal nuclei (red). Means with the same letter are not significantly different at the 95%

537 confidence interval.

Figure S3: Comparison of sperm head Maximum Height (µm) between 50 normal (blue) and 50

abnormal (red) samples. Representation of the difference in Maximum Height (µm) between 11,534

normal nuclei (blue) and 11,326 abnormal nuclei (red). Means with the same letter are not

significantly different at the 95% confidence interval.

Figure S4: Comparison of sperm head Maximum Width (μm) between 50 normal (blue) and 50

abnormal (red) samples. Representation of the difference in Maximum Width (μm) between 11,534

normal nuclei (blue) and 11,326 abnormal nuclei (red). Means with the same letter are not

significantly different at the 95% confidence interval.

Figure S5: Analysis of the test group recovers the same clusters seen in the training group. Cluster analysis of sperm from 4,610 sperm heads from 20 normal samples and 4,642 sperm heads from 20 abnormal samples in the test group using measures of Area, Height, Width, Circularity and Variability by Ward linkage, squared Euclidean distance and standardised variables identifies the same three clusters that were seen in the training group. Below each cluster is the percentage of nuclei within that cluster that are from normal samples and the difference in representation from that expected by chance of (here, positive values show that normal nuclei are overrepresented in a cluster and vice versa for negative values).

Figure S6: Empirical receiver operating curves based on the predictive functions of proportion in cluster 1 (blue), proportion not in cluster 3 (red) and the sum of proportion in cluster 1 and proportion not in cluster 3 (green). The figure also shows the respective values for the area under the curve (AUC) for each of the three predictor variables.