Prevalence of subclinical ketosis and production diseases in dairy cows in

Central and South America, Africa, Asia, Australia and New Zealand, and

Eastern Europe

Nikolaus Brunner[1], Stephan Groeger[1], Jose Raposo[1], Rupert M. Bruckmaier[2], Josef J. Gross[2]*

[1] Bayer Animal Health GmbH, Leverkusen, Germany

[2] Veterinary Physiology, Vetsuisse Faculty, University of Bern, Bern, Switzerland

* Corresponding author

Email: josef.gross@vetsuisse.unibe.ch (JJG)

Abstract

Subclinical ketosis (SCK) and periparturient diseases considerably account for economic and welfare losses in dairy cows. The majority of scientific reports investigating the prevalence of SCK and production diseases are based on empirical studies conducted in Western Europe and North America. The present study surveyed the prevalence of SCK and production-related clinical diseases in early lactating cows in various countries across the world other than those in North America and Western Europe. Twelve countries of South and Central America (Argentina, Brazil, Chile, Colombia, Mexico), Africa (South Africa), Asia (Thailand, China), Eastern Europe (Russia, Ukraine), Australia and New Zealand were assessed, and data from a total of 8,902 cows kept at 541 commercial dairy farms were obtained. A minimum of 5 cows per farm were blood sampled and examined once after parturition up to day 21 of lactation. Blood concentration of β-hydroxybutyrate (BHBA) was measured (threshold for SCK: 1.2 mmol/l) and the presence of production-related diseases such as milk fever, retained placenta, mastitis, metritis, displaced abomasum, claw disease and clinical ketosis was recorded. More than 95% of all cows were examined in their second week of lactation. Across all investigated countries, the SCK prevalence was 24.1%, ranging from 8.3% up to 40.1%. The prevalence of production-related diseases detected during the first 21 days of lactation was relatively low (< 5%). Calculated odds ratios did not indicate an elevated risk for production diseases in cows with SCK. Despite differences in production systems across countries and variation between individual farms within a region, the present study data on SCK prevalence align with observations in Western European and North American dairy herds. At the very early stage of sampling and clinical examination for detection of SCK, it cannot be excluded that certain production diseases such as DA, lameness and mastitis have developed later.

Introduction

At the onset of lactation, dairy cows experience a marked metabolic load due to the prevailing negative energy balance, which makes them susceptible towards infectious and metabolic diseases [1, 2]. Increased concentrations of circulating ketone bodies, predominantly β-hydroxybutyrate (BHBA), without the presence of clinical signs of ketosis are considered as subclinical ketosis (SCK; [3]). The blood BHBA thresholds for SCK diagnosis in literature range between 1.2 and 1.4 mmol/l [4-8]. Although symptoms of clinical ketosis such as reduced milk production, lethargy and loss of appetite are commonly observed at higher concentrations of BHBA (>3.0 mmol/l), some cows, however, may be exposed to high BHBA concentrations without showing any clinical signs, whereas others develop ketosis characteristics already at lower BHBA levels [2, 9, 10].

SCK affects performance and is obviously related to an increased risk of production-related diseases such as clinical ketosis, displaced abomasum (DA), retained placenta, and metritis [11, 12]. Concomitantly, production efficiency decreases (e.g. lower milk production, poor fertility, and increased culling rates) which results in economic losses [11-15]. Serum BHBA concentrations ≥1.2 mmol/l in the first week after calving were associated with a several fold increased risk of subsequently developing DA and metritis, respectively [11]. Ospina et al. [16] found increased risk ratios of 4.9 for clinical ketosis, 2.3 for metritis and 6.9 for DA when plasma BHBA concentrations exceeded a threshold of 1.0 mmol/l during days 3 to 14 post partum. A negative impact of SCK on milk yield of early lactating cows was reported by several authors [10-12, 17]. Furthermore, cows with BHBA concentrations above 1.0 mmol/l in the first week after calving were significantly less likely to be diagnosed pregnant after first insemination [13]. The economic impact of SCK is therefore indisputable, although exact figures quantifying the actual costs of SCK are difficult to collect. Based on an earlier investigation on Canadian dairy farms including more than 2,600 cows, McLaren et al.

[18] estimated that a reduction of 1% in SCK incidence would amount to a saving of 584

USD per year. Recently, McArt et al. [14] calculated the average total costs of

hyperketonemia (blood BHBA concentrations ≥ 1.2 mmol/l) to be 289 USD per case

diagnosed.

Prevalence of SCK in the existing literature ranges between 10 and 40% in early

lactation with highest values occurring within the first three weeks of lactation [11, 17, 19].

Most of these studies, however, included only a small number of dairy herds and cows

(mainly research stations), and were performed in North America and Western Europe [5-8,

12].

The objective of the present study was to survey the prevalence of SCK and the

concomitant occurrence of health disorders in early lactating dairy cows in various countries

across the world other than those in North America and Western Europe. Dairy farms in 12

countries of South and Central America (Argentina, Brazil, Chile, Colombia and Mexico),

Africa (South Africa), Asia (Thailand, China), Eastern Europe (Russia, Ukraine), Australia

and New Zealand were investigated. To our knowledge, this is the first large-scale approach

to investigate the prevalence of SCK beyond Western Europe and North America.

83 Materials and methods

84 Ethics statement

85   All procedures in terms of blood sampling and veterinary examinations followed the

86   animal care and welfare legislation criteria of the involved countries. The approval was

87   provided by the local responsible institutions of the following countries: Argentina, Australia,

88   Brazil, Chile, China, Colombia, Mexico, New Zealand, Russia, South Africa, Thailand and

89   Ukraine.

Animals, blood sampling and analysis

The study was conducted from June 2011 to September 2013 at different commercial dairy farms in 12 different countries worldwide (Argentina, Australia, Brazil, Chile, China, Colombia, Mexico, New Zealand, Russia, South Africa, Thailand and Ukraine). Breeds included were dairy breeds (e.g. predominantly Holstein Friesian, Jersey), local crossbred lines (e.g. Girolondo in Brazil, Simmental x Red Steppe in Russia) and others (e.g. Simmental). Management systems varied from grazing systems to indoor housing of animals throughout the year. In total, 8,902 early lactating dairy cows housed on 541 farms were evaluated. At least five randomly selected cows of a herd in early lactation (day 2 to 21 post partum (p.p.)) inconspicuous of obvious health disorders and without any previous treatments were clinically examined and blood was sampled for diagnostic purposes and health monitoring during regular herd health management visits by veterinarians in the field. Each cow was only tested once and no further restrictions in selection criteria such as parity number, breed, or performance were implied. In all cows, blood samples were taken from the coccygeal vein, and BHBA concentrations were directly determined on site using a handheld meter (Precision Xceed, Abbott Diabetes Care Inc., Alameda, CA, USA), which was previously validated for the use in cows [20, 21]. Concomitantly, each sampled cow was clinically examined for the presence of metabolic and infectious health disorders (milk fever, retained placenta, displaced abomasum (DA), claw disease, mastitis, metritis and clinical ketosis (CK)) following standardized diagnostic definitions (Table 1).

Table 1. Definitions used for disease diagnosis during veterinary examinations of dairy cows in early lactation.

Postpartum disease	Definition	Reference
Milk fever	Cow requires Ca injections due to hypocalcaemia, based on clinical	[29-31]

	signs such as muscular weakness or lying cow unable to rise.	
Retained placenta	Failure to pass placenta within 24 h after calving	[29, 30]
Mastitis	Visually abnormal milk appearance and/or changes in the appearance of the udder (inflammatory signs, swollen or hard quarters)	[29, 30, 32]
Displaced abomasum	Presence of gas-filled abomasum on left or upper right flank, based on a characteristic "ping" sound at auscultation or percussion, confirmed by surgery	[29, 30]
Claw disease	Abnormal findings during claw examinations (such as interdigital and digital dermatitis) and/or lameness with a locomotion score[a] of ≥3 (scale of 1-5).	[29, 30, 33]
Metritis	Enlarged uterus and/or purulent, smelly uterine discharge associated with systemic signs (fever[b], inappetence/anorexia, inactivity, decreased milk yield)	[29, 30, 34]
Clinical ketosis	Decreased milk production, reduced feed intake or inappetence, reduced activity, positive blood, milk or urine ketone test, absence of DA or other primary disease, ketone odor in breath or milk	[29, 30]

[a]scale 1 to 5.

[b]body temperature >39.5°C.

Statistical analysis

Cows sampled between day 2 and 21 p.p. were categorized "SCK negative" if blood BHBA concentrations were <1.2 mmol/l, or "SCK positive" when BHBA concentrations were ≥1.2 mmol/l). The prevalence of SCK was calculated as proportion of animals with BHBA concentrations ≥1.2 mmol/l relative to all investigated cows. Prevalence data were estimated at country and average global level. In addition, overall and country-specific prevalence of postpartum diseases was calculated. The occurrence of postpartum diseases in relation to SCK was evaluated by odds ratio (OR) analyses using the statistical package TESTIMATE (version 6.5, idv-Data Analysis & Study Planning, Krailling, Germany) according to following formula: $OR = (p_1/(1-p_1)) / (p_2/(1-p_2))$, where p_1 indicates the

probability of postpartum diseases under SCK conditions and p_2 the probability of postpartum diseases under non-SCK conditions.

Results

Data analyses were based on a total of 8,902 dairy cows from 541 different commercial dairy farms. The number of farms investigated per country and their herd size ranged from two farms with 83 cows in Thailand to 102 farms housing 2,989 cows in New Zealand. On average, cows of the present dataset were in their 3rd lactation (range of parity number: 1 - 14, 26.9% primiparous and 73.1% multiparous dairy cows; Table 2). The majority (95.1%) of the cows was examined and sampled between days 7 and 15 p.p..

Table 2. Information on animals and farms of the different participating countries in the present study.

Country	Farms (n)	Cows (n)	Percentage of cows		Cattle breed	Housing	Parity number[a]	Milk yield (kg/d per cow)
			primiparous	multiparous				
Argentina	27	720	26.0	74.0	Holstein-Friesian	Pasture and indoors	2.8 ± 1.8	28.5
Australia	22	208	15.0	85.0	Holstein-Friesian	Pasture	3.4 ± 2.0	18.4
Brazil	26	159	35.5	64.5	50% Holstein-Friesian, 50% others (Girolondo, Gyr, Brown Swiss)	Pasture and indoors	2.5 ± 1.7	13 - 30
Chile	17	183	24.6	75.4	Holstein-Friesian	Pasture and indoors	2.6 ± 1.3	23.0
China	41	404	63.1	36.9	Holstein-Friesian	Indoors	1.3 ± 1.5	25 - 35
Colombia	79	791	27.1	72.9	Holstein-Friesian, Brahman, Brahman crossbred	Pasture	2.9 ± 1.8	17 - 21
Mexico	63	2,060	37.6	62.4	Holstein-Friesian or	Pasture and	2.3 ± 1.7	18 - 22

					HF crossbred	indoors		
New Zealand	102	2,989	13.7	86.3	Holstein-Friesian, Jersey, HF/Jersey crossbred	Pasture	4.0 ± 2.4	13 - 19
Russia	77	764	39.3	60.7	Holstein-Friesian, Simmental, Red Steppe	Indoors	2.2 ± 1.2	13 - 23
South Africa	54	377	18.1	81.9	Holstein-Friesian	Pasture	3.3 ± 1.9	20 - 25
Thailand	2	83	32.5	67.5	88% Holstein-Friesian	Indoors	3.3 ± 2.3	12 - 20
Ukraine	31	164	21.3	78.7	50% Holstein-Friesian, 50% others	Indoors	2.6 ± 1.3	13 - 25
Overall	541	8,902	26.9	73.1			3.0 ± 2.1	

[a]Data are expressed as mean values ± SD.

Concentration of blood BHBA and subclinical ketosis

Blood BHBA concentrations observed between day 2 and day 21 p.p. were 1.0 ± 0.8 mmol/l, ranging from an average value of 0.7 mmol/l (Australia, Brazil, Colombia and Russia) up to 1.5 mmol/l (Ukraine). In two thirds of the investigated countries, mean BHBA concentrations were below 1.0 mmol/l (0.7 - 0.9 mmol/l) while in one third of the countries (namely China, New Zealand, Thailand and Ukraine) mean BHBA concentrations of 1.0 mmol/l and above were observed (Table 3). Blood BHBA concentrations of 3.0 mmol/l and beyond during the first 21 days of lactation were predominately observed in China and Ukraine (Table 3). Overall, SCK (BHBA concentrations ≥1.2 mmol/l) was diagnosed in 24.1% of all cows examined, ranging from 8.3% (Colombia) up to 40.1% (New Zealand; Fig 1). In only two (Australia and Colombia) out of the 12 investigated countries, SCK prevalence on the participating farms was less than 10%. In four countries (Brazil, Chile, Mexico, Russia) SCK prevalence varied between 10.7% and 14.8%, whereas in the remaining countries (Argentina, China, New Zealand, South Africa, Thailand, Ukraine) SCK prevalence

was above 15% (between 17.0% and 40.1%). More than 80% of all cows diagnosed SCK were multiparous. In contrast, we observed SCK occurring more frequently in primiparous cows in three countries (Chile, South Africa, Ukraine; Fig 2).

Table 3. Blood BHBA concentrations in early lactating dairy cows (2- 21 days postpartum) in different countries and proportion of cows with highly elevated BHBA concentrations.

Country	BHBA concentration (mmol/l)			Cows with BHBA ≥3.0 mmol/l	
	n	mean ± SD	range	n	%
Argentina	720	0.9 ± 0.9	0.1 - 6.2	36	5.0
Australia	208	0.7 ± 0.7	0.1 - 7.0	4	1.9
Brazil	159	0.7 ± 0.6	0.1 - 4.8	3	1.9
Chile	183	0.8 ± 0.7	0.0 - 4.6	3	1.6
China	404	1.2 ± 1.0	0.0 - 5.8	33	8.2
Colombia	791	0.7 ± 0.4	0.1 - 4.2	4	0.5
Mexico	2,060	0.8 ± 0.7	0.0 - 6.0	53	2.6
New Zealand	2,990	1.2 ± 0.7	0.0 - 8.0	101	3.4
Russia	764	0.7 ± 0.7	0.1 - 6.4	18	2.4
South Africa	376	0.9 ± 1.0	0.1 - 7.2	20	5.3
Thailand	83	1.0 ± 0.8	0.1 - 5.8	2	2.4
Ukraine	164	1.5 ± 1.5	0.0 - 6.7	22	13.4
Overall	8,902	1.0 ± 0.8	0.0 - 8.0	299	3.4

Fig 1. Prevalence of subclinical ketosis worldwide. Prevalence of subclinical ketosis (SCK, blood BHBA concentrations ≥1.2 mmol/l) in early lactating dairy cows studied in 12 countries worldwide.

Fig 2. Subclinical ketosis in primi- and multiparous dairy cows. Occurrence of subclinical ketosis (SCK, blood BHBA concentrations ≥1.2 mmol/l) in primiparous and multiparous cows during early lactation in different countries worldwide.

Prevalence of concomitant production diseases and their association with SCK

The overall prevalence of production diseases was 4.3% for milk fever, 4.0% for

retained placenta, 3.4% for mastitis, 1.7% for claw disease and 5.3% for metritis (Table 4).

Displaced abomasum (DA) and clinical ketosis showed a very low prevalence (0.3% and

0.7%, respectively). During the first 21 days p.p., DA was not seen in 50% (6/12) of the

investigated countries and CK was diagnosed in only 7 out of 12 countries. We further

investigated the relationship of SCK with the concomitant presence of health disorders in

early lactation by performing an odds ratio analysis. However, due to the generally low

prevalence of diseases, odds ratios (OR) were <1, and an increased risk of disease in

association with SCK could not be identified in the present study (data not shown). The

highest odds ratio was calculated for clinical ketosis as cows diagnosed SCK had a 1.062

higher probability of having subsequently clinical ketosis.

Table 4. Prevalence of production diseases between 2 and 21 days postpartum.

Country	Milk fever %	Milk fever n	Retained placenta %	Retained placenta n	Mastitis %	Mastitis n	DA[a] %	DA[a] n	Claw disease %	Claw disease n	Metritis %	Metritis n	CK[b] %	CK[b] n
Argentina	0.7	(5/720)	2.1	(15/720)	5.1	(37/720)	1.0	(7/720)	2.2	(16/720)	15.7	(113/720)	4.0	(29/720)
Australia	10.1	(21/208)	7.2	(15/208)	3.8	(8/208)	0	(0/208)	0.5	(1/207)	6.3	(13/208)	1.9	(4/208)
Brazil	0	(0/159)	8.8	(14/159)	3.8	(6/159)	0.6	(1/159)	1.3	(2/159)	0	(0/159)	0	(0/159)
Chile	7.1	(13/182)	5.5	(10/183)	9.3	(17/183)	1.1	(2/183)	5.5	(10/183)	9.8	(18/183)	2.2	(4/183)
China	1.7	(7/403)	4.0	(16/404)	2.0	(8/404)	2.7	(14/404)	1.0	(4/404)	3.7	(15/404)	1.2	(5/404)
Colombia	1.5	(12/791)	4.3	(34/791)	4.6	(36/791)	0.1	(1/791)	1.0	(8/791)	0.9	(7/791)	0	(0/791)
Mexico	0.6	(13/2060)	4.7	(97/2060)	1.4	(29/2060)	0.2	(4/2060)	1.3	(27/2060)	5.7	(118/2059)	0.6	(12/2060)
New Zealand	0.7	(20/2990)	0.2	(7/2990)	1.2	(36/2990)	0	(0/2990)	0.1	(3/2990)	0.3	(10/2990)	0.1	(2/2989)
Russia	37.4	(286/764)	17.0	(130/764)	14.9	(114/764)	0	(0/764)	10.5	(80/764)	13.5	(103/764)	0.9	(7/764)
South Africa	0.8	(3/377)	1.3	(5/377)	0.8	(3/377)	0	(0/377)	0	(0/377)	6.6	(25/377)	0	(0/377)
Thailand	1.2	(1/83)	0	(0/83)	4.8	(4/83)	0	(0/83)	1.2	(1/83)	19.3	(16/83)	0	(0/83)
Ukraine	0.6	(1/164)	7.3	(12/164)	3.7	(6/164)	0	(0/164)	0.6	(1/164)	19.5	(32/164)	0	(0/164)
Overall	4.3	(382/8901)	4.0	(355/8903)	3.4	(304/8903)	0.3	(26/8903)	1.7	(153/8902)	5.3	(470/8902)	0.7	(63/8902)
Range	0 - 37.4 %		0 - 17.0 %		0.8 - 14.9 %		0 - 2.7 %		0 - 10.5 %		0 - 19.5 %		0 - 4.0 %	

[a]DA = displaced abomasum.

[b]CK = clinical ketosis.

Discussion

In the last three decades, research results addressing SCK and periparturient diseases

represented primarily published data from the United States, Canada and Western Europe [6,

12, 22]. The majority of reports usually based on investigations conducted in regional and

small-sized dairy farms of individual countries [8, 16, 18, 19]. Studies involving a higher

number of animals and covering wider geographic areas were presented earlier by Chapinal et

al. [23] and Suthar et al. [6], but still considered only SCK occurrence within Europe and the

United States, respectively. Different study designs and scopes (such as pre- or postpartum

reclassification of animals, herd or cow level investigations, sampling of blood, milk or urine,

and defining SCK at various BHBA thresholds) do not always allow a straight forward

comparison of literature data across farms, regions and countries, as recently described in a

meta-analysis evaluating 23 SCK publications by Raboisson et al. [22]. Furthermore, data on

global SCK prevalence obtained by a standardized protocol for sampling and defining SCK

are scarce. Therefore, we have investigated the prevalence of SCK and typical production-

related disorders in early lactating dairy cows in various countries across the world other than

those in North America and Western Europe. By using the same validated tool for on-site

BHBA testing in all cows, diagnosing SCK at the generally accepted blood BHBA cut-off

concentration of 1.2 mmol/l and sampling cows in the very early lactation (days 2 to 21 p.p.)

make the results presented here comparable to studies performed in other countries.

The overall SCK prevalence of the present survey was in a comparable range to

observations from a European evaluation of Suthar et al. [6]. In addition, the variation in the

global SCK rates presented here highly scatters (approximately 8 - 40%) and is similar to

findings reported for US and Europe [6, 8, 12]. The majority of our investigated cows with

SCK were multiparous confirming the assumptions that the risk of developing SCK increases

with parity number as milk production rises concomitantly [24]. Based on an analysis of a

variety of publications, Oetzel [19] considered a prevalence rate of 15% representative for

SCK in dairy herds, but emphasized to consider a SCK prevalence of 10% already as an alert

level. In the present study only 2 out of 12 countries (Australia and Colombia) had an average

SCK prevalence of less than 10%, whereas the remaining countries had higher rates. Four

countries (China, New Zealand, Ukraine, and Thailand) revealed even SCK rates up to 40%.

However, this does not exclude that SCK may also occur at higher rates in the countries with

low SCK prevalence in the present study which might be attributed to the random selection of

farms and animals. In general SCK in early lactating dairy cows is globally present. The

highest average SCK prevalence of all countries was observed in New Zealand. One

explanation might be that dairy cows in New Zealand are mostly kept under extensive pasture

based production conditions with low or even zero concentrate supplementation [7]. Along

with the genetic progress in dairy cow breeding, milk production in export-oriented countries

such as New Zealand likely increased and aggravated the energy deficiency in early lactating

cows [2, 25]. In another study, Compton et al. [7] observed a lower SCK prevalence of

approximately 17% in New Zealand dairy cows, which would be still in the upper range when

compared to the present evaluations. The difference just implies the possible variation due to

the selection of farms and animals.

Interestingly, the occurrence of postpartum diseases of the present study was much

lower compared to literature reports. Our observations on average prevalence rates of retained

placenta, mastitis, claw diseases and metritis were approximately half of the values reported

in a European study by Suthar et al. [6]. At first sight, these results were surprising

considering the fact that SCK prevalence rates of this investigation were in agreement with

other reports published. The cows diagnosed SCK in the present study did obviously not

develop health disorders at this early sampling stage. However, results of the concomitant

presence of SCK and further production diseases are inconsistent. While we observed the

highest SCK prevalence on farms in New Zealand, postpartum diseases occurred only at less

than 1% in cows diagnosed SCK. On the other hand, the SCK prevalence data in Russia were

only moderately high (14.1%), but many health disorders were diagnosed at the same time

(milk fever 37.4%, retained placenta 17.0%, mastitis 14.9%, claw disease 10.5%). In a recent

study by Zbinden et al. [10], high-yielding cows fed only herbage showed blood BHBA

concentrations approaching that representative for clinical ketosis, however, not necessarily

followed by the occurrence of diseases. Irrespective of the concomitant presence of

production diseases in cows with SCK, elevated BHBA concentrations ≥1.2 mmol/l

undoubtedly increase the risk of subsequently developing further health disorders [11, 16] at

simultaneously reduced performance in terms of milk yield and reproduction [11-13, 17].

Management and environmental conditions are crucial factors that affect the ability of cows to

cope successfully with the imposed metabolic load. These circumstances cannot be effectively

recorded in a field study like the present one. Furthermore, one might speculate on regional

differences in terms of breed and feeding system affecting the development of health

disorders. The occurrence of DA was reported to be relatively rare in dairy cattle in New

Zealand and Australia [7, 26], which is assumed to be associated with the predominantly

extensive farming on pasture where important risk factors for DA such as low locomotion

activity are not important. However, prevalence of DA concomitant with SCK in the present

study was in particular low in countries with a more intensive milk production in confined

housing systems (e.g. China, Russia, Thailand, and Ukraine). Similar observations were made

for mastitis and the occurrence of lameness. Although lameness may also occur in cattle on

pasture, it appears to be less frequent and to be usually based on a different aetiology [27, 28].

In contrast, we observed that mastitis prevalence in cows with SCK can be elevated

independent if cows were kept on pasture or indoors.

Conclusions

In conclusion, findings of the present study showed that SCK in dairy cows is a global issue and can be observed at various rates in many countries over the world. Possibly due to the diagnosis of SCK and clinical examination at a very early stage of lactation, concomitant postpartum diseases in cows diagnosed for SCK have not yet developed at this stage. An appropriate management is crucial for avoiding a deterioration of animals' metabolic status with the associated negative impacts on health and performance. The results of the present study support the hypothesis that the association between SCK and postpartum diseases is complex and that based on an occasional single determination of blood BHBA concentrations alone, a reliable and robust prediction of further production diseases is not possible.

Acknowledgements

Authors are grateful to all involved veterinarians, farmers and staff on the different farms and countries. Furthermore we would like to thank Mrs. Marion Ocak for her contribution to the statistical analysis and Mrs. Tanja Knoppe for data processing and literature research. This study was supported by Bayer Animal Health GmbH, 51368 Leverkusen, Germany.

References

1. Gross J, van Dorland HA, Bruckmaier RM, Schwarz FJ. Performance and metabolic profile of dairy cows during a lactational and deliberately induced negative energy balance by feed restriction with subsequent realimentation. J Dairy Sci. 2011; 94: 1820–30. https://doi.org/10.3168/jds.2010-3707 PMID: 21426971

2. Bruckmaier RM, Gross JJ. Lactational challenges in transition dairy cows. Anim Prod Sci. 2017; 57: 1471-81. https://doi.org/10.1071/AN16657

3. Duffield T. Subclinical ketosis in lactating dairy cattle. Vet Clin North Am Food Anim

Pract. 2000; 16: 231-53. PMID: 11022338

4. LeBlanc SJ. Monitoring metabolic health of dairy cattle in the transition period. J Reprod

Dev. 2010; 56(suppl): S29-35. PMID: 20629214

5. Rollin E, Berghaus RD, Rapnicki P, Godden SM, Overton MW. The effect of injectable

butaphosphan and cyanocobalamin on postpartum serum beta-hydroxybutyrate, calcium,

and phosphorus concentrations in dairy cattle. J Dairy Sci. 2010; 93: 978-87.

https://doi.org/10.3168/jds.2009-2508 PMID: 20172218

6. Suthar VS, Canelas-Raposo J, Deniz A, Heuwieser W. Prevalence of subclinical ketosis

and relationships with postpartum diseases in European dairy cows. J Dairy Sci. 2013; 96:

2925-38. https://doi.org/10.3168/jds.2012-6035 PMID: 23497997

7. Compton C, McDougall S, Young L, Bryan M. Prevalence of subclinical ketosis in mainly

pasture-grazed dairy cows in New Zealand in early lactation. N Z Vet J. 2014; 62: 30-7.

https://doi.org/10.1080/00480169.2013.823829 PMID: 23981014

8. Garro CJ, Mian L, Cobos Roldán M. (2014). Subclinical ketosis in dairy cows: prevalence

and risk factors in grazing production system. J Anim Physiol Anim Nutr (Berl). 2014; 98:

838-44. https://doi.org/10.1111/jpn.12141 PMID: 24236545

9. Andersson L. Subclinical ketosis in dairy cows. Vet Clin North Am Food Anim Pract.

1988; 4: 233-51. PMID: 3061609

10. Zbinden RS, Falk M, Münger A, Dohme-Meier F, van Dorland HA, Bruckmaier RM,

Gross JJ. Metabolic load in dairy cows kept in herbage-based feeding systems and

suitability of potential markers for compromised well-being. J Anim Physiol Anim Nutr

(Berl). 2017; 101: 767-78. https://doi.org/10.1111/jpn.12498 PMID: 26959798

11. Duffield TF, Lissemore KD, McBride BW, Leslie KE. Impact of hyperketonemia in early

lactation dairy cows on health and production. J Dairy Sci. 2009; 92: 571-80.

https://doi.org/10.3168/jds.2008-1507 PMID: 19164667

12. McArt JA, Nydam DV, Oetzel GR. Epidemiology of subclinical ketosis in early lactation

dairy cattle. J Dairy Sci. 2012; 95: 5056-66. https://doi.org/10.3168/jds.2012-5443 PMID:

22916909

13. Walsh RB, Walton JS, Kelton DF, LeBlanc SJ, Leslie KE, Duffield TF. The effect of

subclinical ketosis in early lactation on reproductive performance of postpartum dairy

cows. J Dairy Sci. 2007; 90: 2788-96. https://doi.org/10.3168/jds.2006-560 PMID:

17517719

14. McArt JA, Nydam DV, Overton MW. Hyperketonemia in early lactation dairy cattle: A

deterministic estimate of component and total cost per case. J Dairy Sci. 2015; 98: 2043-

54. https://doi.org/10.3168/jds.2014-8740 PMID: 25622874

15. Rutherford AJ, Oikonomou G, Smith RF. The effect of subclinical ketosis on activity at

estrus and reproductive performance in dairy cattle. J Dairy Sci. 2016; 99: 4808-15.

https://doi.org/10.3168/jds.2015-10154 PMID: 26995121

16. Ospina PA, Nydam DV, Stokol T, Overton TR. Evaluation of nonesterified fatty acids and

beta-hydroxybutyrate in transition dairy cattle in the northeastern United States: Critical

thresholds for prediction of clinical diseases. J Dairy Sci. 2010; 93: 546-54.

https://doi.org/10.3168/jds.2009-2277 PMID: 20105526

17. Dohoo IR, Martin SW. Subclinical ketosis: prevalence and associations with production

and disease. Can J Comp Med. 1984; 48: 1-5. PMID: 6713247

18. McLaren CJ, Lissemore KD, Duffield TF, Leslie KE, Kelton DF, Grexton B. The

relationship between herd level disease incidence and a return over feed index in Ontario

dairy herds. Can Vet J. 2006; 47: 767-73. PMID: 16933554

19. Oetzel GR. Monitoring and testing dairy herds for metabolic disease. Vet Clin North Am

Food Anim Pract. 2004; 20: 651-674. https://doi.org/10.1016/j.cvfa.2004.06.006 PMID:

15471629

20. Iwersen M, Falkenberg U, Voigtsberger R, Forderung D, Heuwieser W. Evaluation of an

electronic cowside test to detect subclinical ketosis in dairy cows. J Dairy Sci. 2009; 92:

2618-24. https://doi.org/10.3168/jds.2008-1795 PMID: 19447994

21. Pineda A, Cardoso FC. Technical note: Validation of a handheld meter for measuring β-

hydroxybutyrate concentrations in plasma and serum from dairy cows. J Dairy Sci. 2015;

98: 8818-24. https://doi.org/10.3168/jds.2015-9667 PMID: 26506547

22. Raboisson D, Mounié M, Maigné E. Diseases, reproductive performance, and changes in

milk production associated with subclinical ketosis in dairy cows: a meta-analysis and

review. J Dairy Sci. 2014; 97: 7547-63. https://doi.org/10.3168/jds.2014-8237 PMID:

25306269

23. Chapinal N, Carson M, Duffield TF, Capel M, Godden S, Overton M, Santos JE, LeBlanc

SJ. The association of serum metabolites with clinical disease during the transition period.

J Dairy Sci. 2011; 94: 4897-903. https://doi.org/10.3168/jds.2010-4075 PMID: 21943741

24. Duffield TF, Kelton DF, Leslie KE, Lissemore KD, Lumsden JH. Use of test day milk fat

and milk protein to detect subclinical ketosis in dairy cattle in Ontario. Can Vet J. 1997;

38: 713-18. PMID: 9360791

25. Capper JL, Cady RA, Bauman DE. The environmental impact of dairy production: 1944

compared with 2007. J Anim Sci. 2009; 87: 2160-7. https://doi.org/10.2527/jas.2009-1781

PMID: 19286817

26. Jubb TF, Malmo J, Davis GM, Vawser AS. Left-side displacement of the abomasum in

dairy cows at pasture. Aust Vet J. 1991; 68: 140-2. PMID: 2069542

27. Chesterton RN, Pfeiffer DU, Morris RS, Tanner CM. Environmental and behavioural

factors affecting the prevalence of foot lameness in New Zealand dairy herds - a case-

control study. N Z Vet J. 1989; 37: 135-42. PMID: 16031547

28. Laven RA, Holmes CW. A review of the potential impact of increased use of housing on

the health and welfare of dairy cattle in New Zealand. N Z Vet J. 2008; 56: 151-7.

https://doi.org/10.1080/00480169.2008.36827 PMID: 18690250

29. Duffield TF, Leslie KE, Sandals D, Lissemore K, McBride BW, Lumsden JH, Dick P,

Bagg R. Effect of a monensin-controlled release capsule on cow health and reproductive

performance. J Dairy Sci. 1999; 82: 2377-84. PMID: 10575604

30. LeBlanc SJ, Duffield TF, Leslie KE, Bateman KG, TenHag J, Walton JS, Johnson WH.

The effect of prepartum injection of vitamin E on health in transition dairy cows. J Dairy

Sci. 2002; 85: 1416-26. https://doi.org/10.3168/jds.S0022-0302(02)74209-4 PMID:

12146472

31. DeGaris PJ, Lean IJ. Milk fever in dairy cows: a review of pathophysiology and control

principles. Vet J. 2008; 176: 58-69. https://doi.org/10.1016/j.tvjl.2007.12.029 PMID:

18329301

32. Pérez-Cabal MA, de los Campos G, Vazquez AI, Gianola D, Rosa GJ, Weigel KA,

Alenda R. Genetic evaluation of susceptibility to clinical mastitis in Spanish Holstein

cows. J Dairy Sci. 2009; 92: 3472-80. https://doi.org/10.3168/jds.2008-1978 PMID:

19528625

33. Thomsen PT, Munksgaard L, Tøgersen FA. Evaluation of a lameness scoring system for

dairy cows. J Dairy Sci. 2008; 91: 119-26. https://doi.org/10.3168/jds.2007-0496 PMID:

18096932

34. Sheldon IM, Lewis GS, LeBlanc S, Gilbert RO. Defining postpartum uterine disease in

cattle. Theriogenology. 2006; 65: 1516-30.

https://doi.org/10.1016/j.theriogenology.2005.08.021 PMID: 16226305