Title: Multisite phosphorylation regulates phenotypic variability in antibiotic tolerance

Authors: Elizabeth A. Libby [1,2,3], Shlomi Reuveni [2,4,5,6], and Jonathan Dworkin [*,1]

Affiliations:

[1]Department of Microbiology & Immunology, College of Physicians and Surgeons, Columbia

University, New York, NY 10032 USA

[2]Department of Systems Biology, Harvard Medical School, Boston, MA 02115

[3]Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, MA 02115

[4]School of Chemistry, Tel-Aviv University, Tel-Aviv 6997801, Israel

[5]Center for the Physics and Chemistry of Living Systems. Tel Aviv University, 6997801, Tel

Aviv, Israel.

[6]The Sackler Center for Computational Molecular and Materials Science, Tel Aviv University,

6997801, Tel Aviv, Israel.

*Corresponding author, jonathan.dworkin@columbia.edu, 212.342.3731 (tel)

Keywords: antibiotic tolerance, noise, (p)ppGpp, phenotypic variability, Ser/Thr kinase, kinase

inhibitors, Gram-positive bacteria, eSTK, eSTP, two-component systems.

Abstract

Isogenic populations of cells exhibit phenotypic variability that has specific physiological consequences. For example, individual bacteria within a population can differ in their sensitivity to an antibiotic, but whether this variability can be regulated or is generally an unavoidable consequence of stochastic fluctuations is unclear. We observed that a bacterial stress response gene, the (p)ppGpp synthetase *sasA*, exhibits high levels of extrinsic noise in expression, suggestive of a regulatory process. We traced this variability to the convergence of two signaling systems that together control the multisite phosphorylation of a transcription factor, an event largely unexplored in bacteria, This regulatory intersection between a Ser/Thr kinase and a prototypical two component system is crucial for controlling the appearance of outliers, rare cells with unusually high levels of *sasA* expression. Additionally, by examining the full distributions of gene expression we calculated the contribution of the additional Ser/Thr kinase-dependent phosphorylation in setting the relative abundance of cells with a given a level of SasA. We then created a predictive model for the probability of a given cell surviving antibiotic treatment as a function of *sasA* expression. Therefore, our data show that multisite phosphorylation can be used to strongly regulate variations in phenotypes across a bacterial population.

Introduction

Many bacterial phenotypes, including antibiotic tolerance and virulence, often reflect the phenotype of a subset of the population rather than the average behavior [1,2]. Subpopulations of bacteria can arise through purely stochastic processes as well as by regulatory and signaling pathways [3]. Theoretically, one way to create phenotypic diversity via a signaling pathway is multisite phosphorylation, in which each successive phosphorylation changes the activity of a protein [4,5]. However, it has not been experimentally shown in bacterial populations that multisite phosphorylation regulates variation in gene expression between cells, and subsequently, the emergence of phenotypic diversity. Recently, multisite phosphorylation of transcription factors have been observed in pathways involved in antibiotic tolerance and virulence [6], suggesting that dynamics of multisite phosphorylation could have particular physiological relevance.

Bacterial signaling is often characterized in the context of two-component signal transduction systems (TCS) that generally consist of a histidine kinase that phosphorylates a response regulator on a single residue, which then acts as a transcription factor [7]. The stimulus-dependent response of this type of signaling system architecture has been analyzed theoretically [8,9] and experimentally [10,11], with little cell-to-cell variability observed (as quantified by CV), regardless of inducer level. This suggests that extensive cell-to-cell variability is not a general feature of bacterial TCS. However, some notable exceptions have been found for two-component systems with more complex architectures, such as the broad distribution of gene expression in the *E. coli* TorS/TorR regulon [12] which has recently been shown to be an important factor for cell survival during oxygen depletion [13]. The network architecture of bacterial signal transduction systems may therefore play an underappreciated role in the dynamics and survival of bacterial populations.

In addition to TCS, bacteria also have eukaryotic-like (also called Hank's type) Ser/Thr kinase – phosphatase pairs with homology to eukaryotic kinase systems that perform reversible

phosphorylation on Ser and Thr residues [14]. One particular subfamily of these systems appears

to be universally conserved across Gram-positive bacteria and plays key roles in growth and

virulence for many clinically important pathogens including the streptococci, *S. aureus*, *M.*

tuberculosis, *E. faecalis*, and others [6,15]. Genetic and proteomic studies indicate that these

Ser/Thr kinases can perform transcriptional regulation of key cellular processes involved in

antibiotic tolerance and persistence through multisite phosphorylation of transcription factors.

However, to date, the consequences of multisite phosphorylation for gene regulation at the

single-cell-level has not been quantified. In this context the model gram-positive bacterium *B.*

subtilis presents a comparatively straightforward system to quantify the contribution of the

additional Ser/Thr phosphorylation *in vivo*: the homologous kinase-phosphatase pair is

PrkC/PrpC, and it has been verified to regulate gene expression through additional

phosphorylation of a response regulator.

It has been apparent for over 60 years that bacterial populations contain rare cells that

display increased phenotypic resistance to antibiotics [16]. These cells, presumed to be quiescent,

have been implicated in antibiotic treatment failure in genetically susceptible bacterial infections

[17]. To date, it remains unclear to what extent the appearance of these rare cells, is subject to

regulation. Emerging evidence strongly implicates elevated levels of the nucleotide second

messenger (p)ppGpp as a causative agent of quiescence in many bacterial species [18-21].

(p)ppGpp downregulates essential cellular processes such as transcription, translation, and

DNA replication [22]. Although the precise mechanism of (p)ppGpp synthesis and its direct cellular

targets vary between bacterial species, highly elevated levels of (p)ppGpp confer a quiescent

state to the bacterial cell. As many antibiotics target active cellular processes, the resulting cells

exhibit increased antibiotic tolerance, suggesting that cell-to-cell variability in (p)ppGpp may be

involved in phenotypic resistance to antibiotics [21,23-25].

The mechanistic origin of cell-to-cell variability in (p)ppGpp levels across bacterial

populations remains a major open question. To date, this has been best studied in *E. coli*, which

has the RelA (p)ppGpp synthetase and the SpoT hydrolase [26]. In contrast, other bacterial

species often possess dedicated (p)ppGpp synthetases, termed small alarmone synthetases

(SAS), in addition to bi-functional synthetase-hydrolases [27]. These SAS proteins can be

activated transcriptionally [22], suggesting that cell-to-cell variability in (p)ppGpp levels could

originate in the transcriptional regulation of the synthetases themselves. In the Gram-positive

bacterium *B. subtilis*, (p)ppGpp synthesis is regulated by three distinct proteins: RelA, SasA,

and SasB [28]. *B. subtilis* RelA is a bi-functional (p)ppGpp synthetase-hydrolase, and both SasA

and SasB are dedicated synthetases. Although *relA* and *sasB* transcripts are both readily

detectable during log phase growth, *sasA* (formerly *ywaC*) transcripts are found at considerably

lower levels. However, *sasA* is inducible by certain classes of cell-wall-active antibiotics [29,30],

and its induction by alkaline shock increases the cellular levels of ppGpp [28]. Since *sasA*

expression stops growth [31], SasA-mediated (p)ppGpp synthesis provides a mechanism to

induce cellular quiescence in response to environmental stresses. To date, SasA is only known

to be regulated transcriptionally, so significant cell-to-cell variability in *sasA* expression could

produce physiologically relevant cell-to-cell variability in (p)ppGpp levels. The pre-existing

distribution of *sasA* expression may therefore be critical in predicting the relative survival of cells

under conditions that do not specifically induce *sasA*.

In this work, we demonstrate that *sasA* expression displays physiologically relevant amounts

of extrinsic noise, although the average level of *sasA* expression is very low during growth

under non-inducing conditions. Furthermore, we find that both the distribution of *sasA*

expression and the frequency of outliers are strongly regulated by the activity of a highly

conserved eukaryotic-like Ser/Thr kinase system and its subsequent multisite phosphorylation

of a transcription factor. Using quantitative analysis of the full distributions of *sasA* expression,

we find that multisite phosphorylation is responsible for exponentially regulating the abundance

of cells with a given level of SasA and generate a predictive model for *sasA*-expression-

dependent antibiotic tolerance.

Results

The (p)ppGpp synthetase *sasA* exhibits high levels of extrinsic noise in expression

While the population average level of *sasA* expression during growth is extremely low [29], the

average behavior may mask important phenotypic variation between cells. We therefore

generated a transcriptional reporter for *sasA* (P$_{sasA}$-*yfp*) to study the population at the single-cell

level. Surprisingly, there was considerable cell-to-cell variability in P$_{sasA}$-*yfp* (coefficient of

variation, CV~4.95 ± 0.42, mean ± SEM), with most cells having very low expression, and rare

cells showing significantly higher levels of expression (Fig. 1A). Quantification of YFP

fluorescence revealed that a small fraction of the population had much higher (>~10x) levels of

fluorescence than the mean, and rare cells had ~100x. Note that a typical bacterial gene has a

CV in the range 0.1-1 [32-34].

The high levels of cell-to-cell variability in *sasA* expression could be caused by intrinsic

noise from the promoter itself, or by extrinsic noise originating in an upstream process [35]. To

differentiate between these mechanisms, we used a strain with dual fluorescent reporters, P$_{sasA}$-

yfp and P$_{sasA}$-*mcherry* (Fig. 1B). Expression of the dual reporters in individual cells was highly

correlated (Pearson's correlation coefficient, r~0.90±0.08, mean ± SEM), demonstrating that the

noise was largely extrinsic to the promoter (Fig. 1C, D).

To determine if variation in YFP levels were simply caused by global changes in the

population that result in the accumulation of large amounts of fluorescent protein [36], P$_{sasA}$-*yfp*

was compared to a presumably unrelated promoter known to be constitutively active during log

phase growth, P$_{veg}$-*mcherry* [37] (Fig. S1). We found that YFP and mCherry levels were not highly

correlated, suggesting that the high levels of variability in *sasA* expression are largely caused by

a *sasA*-specific pathway. We then tested a previously characterized *sasA* regulator, the sigma

factor σ^M (SigM) [38], that is required for *sasA* expression [30] (Fig. S2A, B). However, *sasA*

expression also did not correlate strongly with *sigM* expression (Fig. S2C, D) demonstrating that

sigM levels alone do not predict variability in *sasA* expression.

sasA is repressed by the Ser/Thr kinase PrkC through the response regulator WalR

Another potential regulator of *sasA* is the WalR transcription factor observed to bind the

sasA promoter in a genome-wide screen [39]. WalR is the response regulator of the essential

WalRK two-component system and is activated by phosphorylation of Asp-53 by WalK [40]. Once

phosphorylated, WalR can either activate or repress genes in its regulon. A reversible second

phosphorylation on WalR Thr-101 by the eukaryotic-like Ser/Thr kinase-phosphatase pair

PrkC/PrpC [41] further increases WalR activity at both activating and repressing sites [42]. In rich

media (LB), the multisite phosphorylation of WalR affects gene expression (*e.g.*, enhanced

activation of *yocH*) specifically in post-log phase [42]. However, in commonly used defined

minimal media (S7-glucose), there is a consistent PrkC-dependent effect on the population

average level of *yocH* expression throughout log phase (Fig. S3). *sasA* is known to be activated

by antibiotics such as bacitracin [29] through σ^M activation. We first tested whether the PrkC/PrpC

– WalR system regulates *sasA* at the population level to determine if WalR activates or

represses *sasA*. We found that PrkC activity represses *sasA* expression through WalR

Thr101~P during bacitracin treatment (Fig. S4). Based on these bulk measurements, we

developed a model for *sasA* regulation (Fig. 2A) in which PrkC activity further potentiates WalR-

repressing activity at *sasA* through a second phosphorylation of WalR at Thr-101. However, it

remained unclear to what extent multisite phosphorylation of WalR affects pre-existing cell-to-

cell variability in *sasA* expression under non-inducing conditions.

PrkC regulates noise in sasA expression through WalR Thr-101 phosphorylation

Cell-to-cell variability in gene expression can be tuned by changing repressor-binding

affinities [43,44], suggesting that multisite phosphorylation of WalR may play a critical role in setting

the observed distribution of *sasA* expression across the population. To test this, we measured

the distribution of *sasA* expression in wild type (WT) cells and compared it to genetic

backgrounds that alter the phosphorylation state of WalR: Δ*prpC* (no phosphatase, high levels

of T101~P), and Δ*prkC* (no kinase, no detectable T101~P) (Fig. 2B). Qualitatively, in the Δ*prpC*

background, the frequency of cells with high levels of *sasA* expression was strongly reduced,

whereas it was strongly increased in the Δ*prkC* background. The PrpC-dependent effect on

sasA expression requires PrkC, since the distribution of *sasA* expression in a strain lacking both

the kinase and phosphatase (Δ(*prpC-prkC*)) is very similar to a strain lacking just the kinase.

We first sought to quantify the effect of WalR multisite phosphorylation on the frequency of

"outliers": cells with a level of *sasA* expression above a fixed threshold in each population. We

therefore compared independent measurements of the distribution of *sasA* expression in WT,

Δ*prpC*, and Δ(*prpC-prkC*) backgrounds (Fig. 2C, left) and found that PrkC significantly affects

the mean frequency of outliers >8 fold by this measure (*walR* ^{WT}, Δ*prpC* vs Δ(*prpC-prkC*): **p-

value~0.004, Kolmogorov-Smirnov test). We repeated the measurements in a *walR* ^{T101A}

background (Fig. 2C, right) and found that PrkC no longer has a significant effect on the mean

frequency of outliers in the phosphosite mutant background (*walR* ^{T101A}, Δ*prpC* vs Δ(*prpC-prkC*):

p-value~0.56, ns, Kolmogorov-Smirnov test). These results are consistent with increased WalR

activity by Thr-101 phosphorylation causing increased repression of *sasA*, and thereby

regulating the frequency of *sasA* outliers (Fig. 2D). Furthermore, heterologous expression of

PrkC was sufficient to reduce the frequency of outliers observed in the Δ*prkC* background (Fig.

S5A). Heterologous expression of PrkC was also able to further reduce the variability to below

that observed in the Δ*prpC* background, approaching the level of cellular autofluorescence (Fig.

S5B). This suggests that at least some of the remaining variability in the Δ*prpC* background

arises due to incomplete saturation of WalR T101~P.

This definition of outliers, however, relies on the definition of a cutoff threshold and therefore

does not fully address how multisite phosphorylation affects the entire distribution of *sasA*

expression across the population. To quantify the effect of PrkC on the distribution of cell-to-cell

variability in *sasA* expression, we deconvolved the measured data from the cellular

autofluorescence (SI). This resulted in autofluorescence-free distributions of *sasA* expression,

allowing better quantitative comparison of expression between genetic backgrounds (Fig. 3A).

This deconvolution method uses only the first two moments (*i.e.*, the mean and variance) of the

observed distributions of fluorescence. As such, the auto-fluorescence free distributions are

relatively insensitive to the observed "outliers" in each distribution, but makes a statistical

prediction for those frequencies. To verify the accuracy of the predictions, these calculated

distributions were re-convolved with the cellular autofluorescence and the reconstructed data

set compared to the original data (Fig. S6). Calculation of the relative enrichment of cells with a

given level of *sasA* in each genetic background revealed that maximal T101~P (Δ*prpC*)

compared to the absence of T101~P (Δ*prkC*) results in exponential changes in the relative

abundance of cells at a given level of *sasA* (Fig. 3B).

Together, the model and the outlier analysis in Fig. 2 suggest that PrkC-dependent

regulation of the distribution of *sasA* expression requires the second WalR phosphosite at Thr-

101. To test this, we repeated the deconvolution procedure for a strain expressing a WalR

mutant that lacks the Thr-101 phosphosite, *walR T101A*, and found that the PrkC-dependent

effect on *sasA* expression is indeed WalR Thr-101-dependent (Fig. 3C, Fig. S7A). Thus,

multisite phosphorylation is responsible for the exponential depletion of cells with medium to

high levels of *sasA* expression in the Δ*prpC* background (Fig. 3B).

We then measured how intermediate levels of multisite phosphorylation regulate the

distribution of *sasA* expression using the kinase inhibitor staurosporine to progressively inhibit

PrkC activity [45] (Figs. S7B; S8). The distributions of *sasA* expression were again deconvolved

(Fig. S7C), and we calculated the relative enrichment of cells with a given level of *sasA*

fluorescence at increasing concentrations of staurosporine (Fig. 3D). Titration of PrkC activity

resulted in exponential enrichment of cells with a given level of *sasA*. Therefore, even small

changes in PrkC activity result in large changes in the abundance of "outliers", cells with

unusually high levels of *sasA*.

sasA expression level continuously predicts the probability of surviving antibiotic

treatment

Cell-to-cell variability in (p)ppGpp production has been proposed to result in cell-to-cell

variability in antibiotic survival [4748,49]. However, a direct and quantitative relationship between the

expression of a transcriptionally regulated (p)ppGpp synthetase and the probability of survival

for an individual cell has not been demonstrated. We therefore sought to determine if cells with

pre-existing high levels of *sasA* preferentially survive antibiotic exposure, and if so, provide a

model for how the level of *sasA* expression influences the probability of survival for a given cell.

We used ciprofloxacin, a DNA gyrase inhibitor that does not significantly increase the

population average level of *sasA* expression [29]. We measured (Fig. S9) and deconvolved (Fig.

4A,B) the distributions of *sasA* expression (P*sasA*-*yfp*) both pre-and post-ciprofloxacin treatment

that results in ~99% killing in both WT and Δ*sasA* backgrounds. We note that, importantly, the

starting distributions of P*sasA*-*yfp* are very similar in both genetic backgrounds, allowing a direct

comparison. Using these distributions, we calculated the relative enrichment of cells with a

given level of *sasA* following antibiotic treatment (Fig. 4C), yielding a simple model for the effect

of antibiotic treatment on the distribution of *sasA* expression (SI). Because survival after

antibiotic treatments can be affected by many processes, we separated out the size of the *sasA*-

dependent effect by using a Δ*sasA* mutant as a control. WT populations exhibited a significant

increase in the fraction of cells with elevated levels of *sasA* (Fig. 4C). This effect is strongly

reduced in the Δ*sasA* background, demonstrating that *sasA* has a significant contribution to

survival after ciprofloxacin treatment.

We then sought to determine whether our enrichment model reflects the probability of

survival for cells as a function of *sasA* expression. Since SasA is a (p)ppGpp synthetase, the

sasA-dependent enrichment we observe post-ciprofloxacin treatment could be due to an

expression- dependent probability of surviving antibiotic treatment. Alternately, since the

measured increase in mean fluorescence was relatively small (~2 fold), it is also possible that

ciprofloxacin acts in a complex, expression-dependent, manner to generate the observed post-

treatment distribution of *sasA* expression but does not affect survival. To differentiate between

these hypotheses, we used FACS to sort bacteria prior to ciprofloxacin treatment from both WT

and Δ*sasA* populations into "high" (upper ~1%) and "low" (~average) P$_{sasA}$-*yfp* expression

groups (Fig. 4D). Following ciprofloxacin treatment (as in Figs. 4A,B, S9), the relative survival of

"high" and "low" expression cells, or the survival ratio, was assayed by plating for CFUs (Fig.

4E). The average fluorescence cutoff values used in the FACS experiments, low: 4.1±1.8 and

high: 779.5±134.9 (mean ± SEM, 3 experiments), were then used as inputs for a model where

the enrichment of cells with increased levels of *sasA* (Fig. 4C) is caused by increased survival

(SI). The model yielded good agreement with the results of the FACS experiments: it predicted

relative survival ratios of ~9 for wild-type, and ~2 for Δ*sasA*, respectively, compared to the

measured values of ~9.5 ± 0.6 (WT) and 1.8 ± 0.7 (Δ*sasA*) (Fig. 4E). Therefore, the enrichment

of cells with elevated levels of *sasA* post-ciprofloxacin treatment can be largely attributed to the

increased survival probability of pre-existing cells in the population with elevated *sasA*

expression.

Taken together, our results demonstrate that an important consequence of PrkC-dependent

multisite phosphorylation of WalR is the regulation of cell-to-cell variability, or noise, in the WalR

regulon gene *sasA*. By comparing the full distributions of gene expression, we demonstrate that

this effect is not just confined to the regulation of outliers in gene expression above an arbitrary

threshold within the population, but has an exponential effect on the relative abundance of cells

with a given level of expression. By analyzing the full distributions of expression, we are also

able to demonstrate that *sasA* expression also continuously affects the antibiotic tolerance of

individual cells: specifically, the survival probability during a fixed course antibiotic treatment.

This model (SI) is consistent with cell sorting experiments that explicitly demonstrate that the

observed distributions are a consequence of survival probability.

Discussion:

Antibiotic tolerance is believed to be an important factor in the failure of antibiotic treatments

and a key step toward the development of antibiotic resistance [50]. We therefore sought to trace

the origin of the cell-to-cell variability in expression of *sasA* and determine if it can be regulated

by genetic or chemical means. Noise in gene expression can be conceptually separated into

intrinsic and extrinsic noise. Although it is difficult to design strategies to specifically target

events generated by intrinsic noise, extrinsic noise may have upstream regulatory pathways that

can be modulated. Therefore, it is significant that the cell-to-cell variability in *sasA* was

dominated by extrinsic noise at high levels of expression (Fig. 1) that have the strongest effect

on antibiotic tolerance (Fig. 4). Furthermore, since multisite phosphorylation is responsible for

setting the observed distribution of cell-to-cell variability (Figs. 2, 3), this regulatory pathway

could be a novel antibiotic target.

Multisite phosphorylation can expand the range of a protein's function, generating both

switch-like [51,52], and graded [53,54] changes in average activity. In contrast, here we observed only

minimal changes in the average levels of *sasA* expression as a function of PrkC activity, but

measured up to a ~100-fold effect on the frequency of "outliers", cells with particularly high

levels of expression (Fig. 3B). This response was shown to be graded, rather than switch-like,

likely arising as a consequence of the integration of signals from two distinct signaling systems.

A single phosphorylation at WalR Asp-53 strongly, but imperfectly, represses the *sasA*

promoter. The addition of the second phosphorylation at Thr-101 by a distinct signaling system

then acts as a second input to further regulate WalR. Interestingly, even small changes in

activity of the second system result in marked changes in the frequency of outliers.

Heterologous expression of PrkC is capable of reducing the variability observed to close to

cellular autofluorescence, but does not eliminate it completely. This remaining variability in *sasA*

may be due to PrkC overexpression still being unable to completely saturate WalR

phosphorylation, intrinsic noise at the promoter, or as yet unidentified sources.

Transcriptional regulation of outliers in eukaryotes has been shown to be predictive of which

cancer cells survive drug treatment [55]. Here we found that transcriptional regulation by multisite

phosphorylation is also critical for setting the pre-existing distribution of survival probabilities for

cells within a bacterial population. Distinct from bacterial persistence, which is characterized by

bi-phasic killing, these survival probabilities reflect antibiotic tolerance or the killing kinetics

during a relatively short, fixed time-course, antibiotic treatment [56]. In the Δ*sasA* background, we

observed a much weaker dependence of antibiotic tolerance on *sasA* expression. This residual

dependence is consistent with previous results that have implicated many global processes in

antibiotic tolerance including heterogeneity in growth state [24,57,58] and enhanced expression of

drug efflux pumps [59,60]. This is also consistent with the relatively weak correlation in expression

between *sasA* and the constitutive promoter *veg* (Fig. S2). Indeed, it remains to be seen

precisely how cellular physiology changes in a *sasA*-expression dependent manner. SasA has

been shown to be important for ribosome assembly in *B. subtilis* [31] and for survival during

envelope stress in *S. aureus* [61]. More generally, various cellular processes are known to be

directly and indirectly affected by rising (p)ppGpp levels including inhibition of DNA primase

activity [62], and reduction in intracellular GTP pools [63] thereby downregulating rRNA transcription

[64]. As our results show that antibiotic survival increases continuously with *sasA* expression, it

suggests that SasA exerts a continuous effect proportional to its level on physiological

processes that mediate ciprofloxacin killing. Therefore, multisite phosphorylation may provide a

"bet-hedging" strategy to regulate the phenotypic diversity of a bacterial population, serving as a

broadly useful mechanism to tune the frequency of rare phenotypes that facilitate survival under

adverse conditions.

Acknowledgements:

We thank Eric Brown for the strain EB1385, Isaac Plant (Silver lab) for pIP384 and IP563, and Amir Figueroa at the Microbiology & Immunology Core Facility at Columbia University for assistance with flow cytometry and cell sorting. This work was supported by NIH GM114213 and a BWF Investigators in the Pathogenesis of Infectious Disease award to JD, a grant from the Department of Systems Biology at Harvard Medical School to EAL, and SR was supported by NIH GM095784 and gratefully acknowledges support from the Azrieli Foundation. Author Contributions: Conceived and designed the experiments EAL and JD. EAL performed the experiments. EAL and SR analyzed data. Contributed reagents/materials/analysis tools: EAL and SR. Wrote the paper: EAL, SR and JD.

References:

Arnoldini, M. *et al.* Bistable expression of virulence genes in salmonella leads to the formation of an antibiotic-tolerant subpopulation. *PLoS Biol* **12**, e1001928, doi:10.1371/journal.pbio.1001928 (2014).

Norman, T. M., Lord, N. D., Paulsson, J. & Losick, R. Stochastic Switching of Cell Fate in Microbes. *Annual review of microbiology* **69**, 381-403, doi:10.1146/annurev-micro-091213-112852 (2015).

Maamar, H., Raj, A. & Dubnau, D. Noise in gene expression determines cell fate in Bacillus subtilis. *Science* **317**, 526-529, doi:10.1126/science.1140818 (2007).

Gunawardena, J. Multisite protein phosphorylation makes a good threshold but can be a poor switch. *Proc Natl Acad Sci U S A* **102**, 14617-14622, doi:10.1073/pnas.0507322102 (2005).

Ortega, F., Garces, J. L., Mas, F., Kholodenko, B. N. & Cascante, M. Bistability from double phosphorylation in signal transduction. Kinetic and structural requirements. *FEBS J* **273**, 3915-3926, doi:10.1111/j.1742-4658.2006.05394.x (2006).

Wright, D. P. & Ulijasz, A. T. Regulation of transcription by eukaryotic-like serine-threonine kinases and phosphatases in Gram-positive bacterial pathogens. *Virulence* **5**, 863-885, doi:10.4161/21505594.2014.983404 (2014).

Laub, M. T. & Goulian, M. Specificity in two-component signal transduction pathways. *Annu Rev Genet* **41**, 121-145, doi:10.1146/annurev.genet.41.042007.170548 (2007).

Batchelor, E. & Goulian, M. Robustness and the cycle of phosphorylation and dephosphorylation in a two-component regulatory system. *Proc Natl Acad Sci U S A* **100**, 691-696, doi:10.1073/pnas.0234782100 (2003).

Shinar, G., Milo, R., Martinez, M. R. & Alon, U. Input output robustness in simple bacterial signaling systems. *Proc Natl Acad Sci U S A* **104**, 19931-19935, doi:10.1073/pnas.0706792104 (2007).

Batchelor, E., Silhavy, T. J. & Goulian, M. Continuous control in bacterial regulatory circuits. *J Bacteriol* **186**, 7618-7625, doi:10.1128/JB.186.22.7618-7625.2004 (2004).

Miyashiro, T. & Goulian, M. High stimulus unmasks positive feedback in an autoregulated bacterial signaling circuit. *Proc Natl Acad Sci U S A* **105**, 17457-17462, doi:10.1073/pnas.0807278105 (2008).

Roggiani, M. & Goulian, M. Oxygen-Dependent Cell-to-Cell Variability in the Output of the Escherichia coli Tor Phosphorelay. *J Bacteriol* **197**, 1976-1987, doi:10.1128/JB.00074-15 (2015).

Carey, J. N. *et al.* Regulated Stochasticity in a Bacterial Signaling Network Permits Tolerance to a Rapid Environmental Change. *Cell*, doi:10.1016/j.cell.2018.02.005 (2018).

Pereira, S. F., Goss, L. & Dworkin, J. Eukaryote-like serine/threonine kinases and phosphatases in bacteria. *Microbiol Mol Biol Rev* **75**, 192-212, doi:10.1128/MMBR.00042-10 (2011).

15   Manuse, S., Fleurie, A., Zucchini, L., Lesterlin, C. & Grangeasse, C. Role of eukaryotic-
like serine/threonine kinases in bacterial cell division and morphogenesis. *FEMS*
Microbiol Rev **40**, 41-56, doi:10.1093/femsre/fuv041 (2016).

16   Bigger, J. W. Treatment of staphylococcal infections with penicillin by intermittent
sterilisation. *Lancet* **244**, 497–500 (1944).

17   Fauvart, M., De Groote, V. N. & Michiels, J. Role of persister cells in chronic infections:
clinical relevance and perspectives on anti-persister therapies. *J Med Microbiol* **60**, 699-
709, doi:10.1099/jmm.0.030932-0 (2011).

18   Abranches, J. *et al.* The molecular alarmone (p)ppGpp mediates stress responses,
vancomycin tolerance, and virulence in Enterococcus faecalis. *J Bacteriol* **191**, 2248-
2256, doi:10.1128/JB.01726-08 (2009).

19   Gaca, A. O. *et al.* Basal levels of (p)ppGpp in Enterococcus faecalis: the magic beyond
the stringent response. *MBio* **4**, e00646-00613, doi:10.1128/mBio.00646-13 (2013).

20   Dordel, J. *et al.* Novel determinants of antibiotic resistance: identification of mutated loci
in highly methicillin-resistant subpopulations of methicillin-resistant Staphylococcus
aureus. *MBio* **5**, e01000, doi:10.1128/mBio.01000-13 (2014).

21   Maisonneuve, E. & Gerdes, K. Molecular mechanisms underlying bacterial persisters.
Cell **157**, 539-548, doi:10.1016/j.cell.2014.02.050 (2014).

22   Liu, K., Bittner, A. N. & Wang, J. D. Diversity in (p)ppGpp metabolism and effectors. *Curr*
Opin Microbiol **24**, 72-79, doi:10.1016/j.mib.2015.01.012 (2015).

23   Nguyen, D. *et al.* Active starvation responses mediate antibiotic tolerance in biofilms and
nutrient-limited bacteria. *Science* **334**, 982-986, doi:10.1126/science.1211037 (2011).

24   Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. Bacterial persistence as a
phenotypic switch. *Science* **305**, 1622-1625, doi:10.1126/science.1099390 (2004).

25   Maisonneuve, E., Castro-Camargo, M. & Gerdes, K. (p)ppGpp Controls Bacterial
Persistence by Stochastic Induction of Toxin-Antitoxin Activity. *Cell* **154**, 1140-1150,
doi:10.1016/j.cell.2013.07.048 (2013).

26   Cashel, M., Gentry, D., Hernandez, V. & Vinella, D. in *Esherichia coli and Salmonella;*
Cellular and Molecular Biology Vol. 2   1458-1496 (ASM Press, Washington DC, 1996).

27   Atkinson, G. C., Tenson, T. & Hauryliuk, V. The RelA/SpoT homolog (RSH) superfamily:
distribution and functional evolution of ppGpp synthetases and hydrolases across the
tree of life. *PLoS One* **6**, e23479, doi:10.1371/journal.pone.0023479 (2011).

28   Nanamiya, H. *et al.* Identification and functional analysis of novel (p)ppGpp synthetase
genes in Bacillus subtilis. *Mol Microbiol* **67**, 291-304, doi:10.1111/j.1365-
2958.2007.06018.x (2008).

29   D'Elia, M. A. *et al.* Probing teichoic acid genetics with bioactive molecules reveals new
interactions among diverse processes in bacterial cell wall biogenesis. *Chemistry &*
biology **16**, 548-556, doi:10.1016/j.chembiol.2009.04.009 (2009).

30   Eiamphungporn, W. & Helmann, J. D. The Bacillus subtilis sigma(M) regulon and its
contribution to cell envelope stress responses. *Mol Microbiol* **67**, 830-848,
doi:10.1111/j.1365-2958.2007.06090.x (2008).

31　Tagami, K. *et al.* Expression of a small (p)ppGpp synthetase, YwaC, in the (p)ppGpp(0)
mutant of Bacillus subtilis triggers YvyD-dependent dimerization of ribosome.
MicrobiologyOpen **1**, 115-134, doi:10.1002/mbo3.16 (2012).

32　Pedraza, J. M. & van Oudenaarden, A. Noise propagation in gene networks. *Science*
307, 1965-1969, doi:10.1126/science.1109090 (2005).

33　Milo, R., Jorgensen, P., Moran, U., Weber, G. & Springer, M. BioNumbers--the database
of key numbers in molecular and cell biology. *Nucleic Acids Res* **38**, D750-753,
doi:10.1093/nar/gkp889 (2010).

34　Taniguchi, Y. *et al.* Quantifying E. coli proteome and transcriptome with single-molecule
sensitivity in single cells. *Science* **329**, 533-538, doi:10.1126/science.1188308 (2010).

35　Elowitz, M. B., Levine, A. J., Siggia, E. D. & Swain, P. S. Stochastic gene expression in
a single cell. *Science* **297**, 1183-1186, doi:10.1126/science.1070919 (2002).

36　Leveau, J. H. & Lindow, S. E. Predictive and interpretive simulation of green fluorescent
protein expression in reporter bacteria. *J Bacteriol* **183**, 6752-6762,
doi:10.1128/JB.183.23.6752-6762.2001 (2001).

37　Radeck, J. *et al.* The Bacillus BioBrick Box: generation and evaluation of essential
genetic building blocks for standardized work with Bacillus subtilis. *J Biol Eng* **7**, 29,
doi:10.1186/1754-1611-7-29 (2013).

38　Thackray, P. D. & Moir, A. SigM, an extracytoplasmic function sigma factor of Bacillus
subtilis, is activated in response to cell wall antibiotics, ethanol, heat, acid, and
superoxide stress. *J Bacteriol* **185**, 3491-3498 (2003).

39　Salzberg, L. I. *et al.* The WalRK (YycFG) and sigma(I) RsgI regulators cooperate to
control CwlO and LytE expression in exponentially growing and stressed Bacillus subtilis
cells. *Mol Microbiol* **87**, 180-195, doi:10.1111/mmi.12092 (2013).

40　Dubrac, S., Bisicchia, P., Devine, K. M. & Msadek, T. A matter of life and death: cell wall
homeostasis and the WalKR (YycGF) essential signal transduction pathway. *Mol
Microbiol* **70**, 1307-1322 (2008).

41　Gaidenko, T. A., Kim, T. J. & Price, C. W. The PrpC serine-threonine phosphatase and
PrkC kinase have opposing physiological roles in stationary-phase Bacillus subtilis cells.
J Bacteriol **184**, 6109-6114 (2002).

42　Libby, E. A., Goss, L. A. & Dworkin, J. The Eukaryotic-Like Ser/Thr Kinase PrkC
Regulates the Essential WalRK Two-Component System in Bacillus subtilis. *PLoS
Genet* **11**, e1005275, doi:10.1371/journal.pgen.1005275 (2015).

43　Jones, D. L., Brewster, R. C. & Phillips, R. Promoter architecture dictates cell-to-cell
variability in gene expression. *Science* **346**, 1533-1536, doi:10.1126/science.1255301
(2014).

44　Sanchez, A., Garcia, H. G., Jones, D., Phillips, R. & Kondev, J. Effect of promoter
architecture on the cell-to-cell variability in gene expression. *PLoS Comput Biol* **7**,
e1001100, doi:10.1371/journal.pcbi.1001100 (2011).

45　Shah, I. M., Laaberki, M. H., Popham, D. L. & Dworkin, J. A eukaryotic-like Ser/Thr
kinase signals bacteria to exit dormancy in response to peptidoglycan fragments. *Cell*
135, 486-496 (2008).

46   Kriel, A. *et al.* GTP dysregulation in Bacillus subtilis cells lacking (p)ppGpp results in
phenotypic amino acid auxotrophy and failure to adapt to nutrient downshift and regulate
biosynthesis genes. *J Bacteriol* **196**, 189-201, doi:10.1128/JB.00918-13 (2014).

47   Fisher, R. A., Gollan, B. & Helaine, S. Persistent bacterial infections and persister cells.
Nat Rev Microbiol **15**, 453-464, doi:10.1038/nrmicro.2017.42 (2017).

48   Hahn, J., Tanner, A. W., Carabetta, V. J., Cristea, I. M. & Dubnau, D. ComGA-RelA
interaction and persistence in the Bacillus subtilis K-state. *Mol Microbiol* **97**, 454-471,
doi:10.1111/mmi.13040 (2015).

49   Germain, E., Castro-Roa, D., Zenkin, N. & Gerdes, K. Molecular mechanism of bacterial
persistence by HipA. *Mol Cell* **52**, 248-254, doi:10.1016/j.molcel.2013.08.045 (2013).

50   Levin-Reisman, I. *et al.* Antibiotic tolerance facilitates the evolution of resistance.
Science **355**, 826-830, doi:10.1126/science.aaj2191 (2017).

51   Ferrell, J. E., Jr. & Ha, S. H. Ultrasensitivity part II: multisite phosphorylation,
stoichiometric inhibitors, and positive feedback. *Trends Biochem Sci* **39**, 556-569,
doi:10.1016/j.tibs.2014.09.003 (2014).

52   Nash, P. *et al.* Multisite phosphorylation of a CDK inhibitor sets a threshold for the onset
of DNA replication. *Nature* **414**, 514-521, doi:10.1038/35107009 (2001).

53   Zaytsev, A. V. *et al.* Multisite phosphorylation of the NDC80 complex gradually tunes its
microtubule-binding affinity. *Mol Biol Cell* **26**, 1829-1844, doi:10.1091/mbc.E14-11-1539
(2015).

54   Pufall, M. A. *et al.* Variable control of Ets-1 DNA binding by multiple phosphates in an
unstructured region. *Science* **309**, 142-145, doi:10.1126/science.1111915 (2005).

55   Shaffer, S. M. *et al.* Rare cell variability and drug-induced reprogramming as a mode of
cancer drug resistance. *Nature* **546**, 431-435, doi:10.1038/nature22794 (2017).

56   Kester, J. C. & Fortune, S. M. Persisters and beyond: mechanisms of phenotypic drug
resistance and drug tolerance in bacteria. *Crit Rev Biochem Mol Biol* **49**, 91-101,
doi:10.3109/10409238.2013.869543 (2014).

57   Conlon, B. P. *et al.* Persister formation in Staphylococcus aureus is associated with ATP
depletion. *Nature microbiology* **1**, 16051, doi:10.1038/nmicrobiol.2016.51 (2016).

58   Gutierrez, A. *et al.* Understanding and Sensitizing Density-Dependent Persistence to
Quinolone Antibiotics. *Mol Cell* **68**, 1147-1154 e1143, doi:10.1016/j.molcel.2017.11.012
(2017).

59   Adams, K. N. *et al.* Drug tolerance in replicating mycobacteria mediated by a
macrophage-induced efflux mechanism. *Cell* **145**, 39-53, doi:10.1016/j.cell.2011.02.022
(2011).

60   Pu, Y. *et al.* Enhanced Efflux Activity Facilitates Drug Tolerance in Dormant Bacterial
Cells. *Mol Cell* **62**, 284-294, doi:10.1016/j.molcel.2016.03.035 (2016).

61   Geiger, T., Kastle, B., Gratani, F. L., Goerke, C. & Wolz, C. Two small (p)ppGpp
synthases in Staphylococcus aureus mediate tolerance against cell envelope stress
conditions. *J Bacteriol* **196**, 894-902, doi:10.1128/JB.01201-13 (2014).

62   Wang, J. D., Sanders, G. M. & Grossman, A. D. Nutritional control of elongation of DNA
replication by (p)ppGpp. *Cell* **128**, 865-875, doi:10.1016/j.cell.2006.12.043 (2007).

63 Kriel, A. *et al.* Direct regulation of GTP homeostasis by (p)ppGpp: a critical component of
viability and stress resistance. *Mol Cell* **48**, 231-241, doi:10.1016/j.molcel.2012.08.009
(2012).

64 Krasny, L. & Gourse, R. L. An alternative strategy for bacterial ribosome synthesis:
Bacillus subtilis rRNA transcription regulation. *EMBO J* **23**, 4473-4483,
doi:10.1038/sj.emboj.7600423 (2004).

Figure Legends:

Figure 1: *sasA* exhibits cell-to-cell variability in expression.

A) P$_{sasA}$-*yfp* expression in log phase culture. **Left:** YFP image. Color scale indicates the fold change relative to the population average. **Right:** Phase contrast. Scale bar indicates 10 μm.

B) Schematic of dual-color transcriptional reporter strain to characterize the noise in *sasA* expression. Two copies of the *sasA* promoter driving different reporters (P$_{sasA}$-*yfp*, P$_{sasA}$-*mcherry*) are inserted at separate ectopic loci.

C) Evidence for extrinsic noise in *sasA* expression**.** Images of single cells from the dual reporter strain with significant fluorescence shown in mCherry (**Left**), YFP (**Center**), and phase contrast (**Right**) channels. Scale bar indicates 5 μm.

D) Quantification of cellular fluorescence intensities for single cells in the experiment described in **B,C**. Values were normalized relative to the population mean value for each channel. Data shown was measured on ~ 580 cells with detectable fluorescence in both channels. The fluorescence of the two reporters have a Pearson's correlation coefficient of r ~ 0.90±0.08 (mean ± SEM, 4 experiments). For each reporter, the CV ~ 4.95±0.42 (mean ± SEM, 4 experiments).

Figure 2: The Ser/Thr kinase PrkC and phosphatase PrpC regulate cell-to-cell variability in *sasA*

A) Model for PrkC-dependent regulation of *sasA*. WalR binding to the *sasA* promoter represses *sasA*. WalR activity is primarily controlled by phosphorylation on Asp-53 by its cognate histidine kinase WalK, and secondarily by phosphorylation on Thr-101 by the Ser/Thr kinase PrkC. WalR Thr-101~P can be dephosphorylated by the phosphatase PrpC. Thr-101~P further enhances the repressing activity of WalR Asp-53~P, resulting in lower expression of *sasA* under conditions with high Thr-101~P (e.g., Δ*prpC*). Under conditions lacking Thr-101~P (*e.g.,* Δ*prkC*), the increased repression of *sasA* is relieved.

B) Histograms demonstrating PrkC-dependent cell-to-cell variability in *sasA*. **Top:** P*sasA*-*yfp* reporter activity was quantified by flow cytometry in wild type (WT, gray), Δ*prpC* (orange), Δ*prkC* (green), and Δ(*prpC-prkC*) (blue) backgrounds. Shaded region of each plot indicates the range of cellular autofluorescence observed. Each histogram was computed from data on ~3.0×10^4 events. **Bottom:** Smoothed, overlaid histograms of the data shown above for comparison.

C) Percentage of outliers in each genetic background. At least 4 independent experiments, similar to and including the representative one shown in **B**, were performed in *walR* ^{WT} (**Left**) and *walR* ^{T101A} (**Right**) backgrounds. Each experiment was normalized to a control and outliers were defined as cells above a fixed threshold level of normalized fluorescence (~1250 A.U.). Dots represent the percentage of each population that is above the threshold; bars and lines represent the mean and SEM, respectively.

D) Summary of each genotype, its effect on the maximal occupancy of the two known WalR phosphosites, and the expected effect on WalR activity relative to WT. For each

541   genotype the mean percentage of outliers in *sasA* expression (shown in **C**) are also

542   summarized. Note that in the Δ*prpC* background, the Thr-101 phosphorylation is

543   stabilized (denoted as **T101***).

Figure 3: Multisite phosphorylation causes exponential changes in abundance of cells with a given level of *sasA* expression

A) Functional fits of autofluorescence-free distributions of *sasA* expression. A deconvolution algorithm was used to remove the contribution of autofluorescence from the measured distributions of *sasA* expression shown in Figure 2B. Validation of the fits are shown in Fig. S6.

B) Effect of PrkC on the distribution of *sasA* expression. For each genotype in **A** the frequency of cells with a given level of *sasA* expression was normalized against the equivalent frequency in the WT population. On this semi-logarithmic plot, straight lines indicate that the enrichment factor changes ~exponentially with fluorescence.

C) PrkC-dependent regulation of cell-to-cell variability in *sasA* is WalR Thr-101-dependent. P$_{sasA}$-*yfp* reporter activity was quantified by flow cytometry in a Δ*prpC walR T101A* (yellow, solid) background and compared to Δ*prpC* (orange), Δ(*prpC-prkC*) (blue), and Δ(*prpC-prkC*) *walR T101A* (yellow, dashed) backgrounds in the same experiment. Plots were generated by applying the deconvolution method described in **A** to the measured data in Fig. S7A.

D) Gradual inhibition of PrkC results in progressive enrichment of cells with elevated levels of *sasA*. P$_{sasA}$-*yfp* reporter activity was quantified by flow cytometry during treatment with increasing concentrations of staurosporine: 0 (solvent only; black), 50 nM, 100 nM, and 1 µM (shades of magenta), in otherwise WT populations. For reference, a Δ(*prpC-prkC*) (blue) population treated with solvent only is shown; note that the effect of staurosporine saturates at this level. Plots were generated by applying the deconvolution method described in **A** to the measured data in Fig. S7B,C. For each condition the frequency of cells with a given level of *sasA* expression was normalized against the equivalent frequency in the WT population. As in **B**,

569   straight lines indicate that enrichment increases ~exponentially with fluorescence.

570   Enrichment becomes more pronounced as the concentration of staurosporine

571   increases and saturates at the level of complete inhibition ~Δ(*prpC-prkC*).

Figure 4: Cells with elevated *sasA* expression preferentially survive antibiotic treatment

A) Autofluorescence-free distributions of *sasA* expression before (dark gray) and after (light gray) ciprofloxacin treatment in an otherwise WT background based on the data shown in Fig. S9A.

B) Autofluorescence-free distributions of *sasA* expression before (dark red) and after (light red) ciprofloxacin treatment in a Δ*sasA* background based on the data shown in Fig. S9B.

C) Enrichment plot following antibiotic treatment for WT and Δ*sasA* populations. For each genotype in **A** and **B** the frequency of cells with a given level of P$_{sasA}$-*yfp* after antibiotic treatment was normalized against the corresponding frequency prior to antibiotic treatment. Dashed lines and dots indicate the average values used in **D** and **E**.

D) Histograms of P$_{sasA}$-*yfp* fluorescence in WT (black) and Δ*sasA* (red) backgrounds measured by flow cytometry prior to sorting into "low" (4.1 ± 1.8) or "high" (779.5 ± 134.9) expression groups (relative mean ± SEM, 3 experiments). Each histogram is comprised of data obtained from ~$3.0*10^4$ events and is from a representative experiment used in **E**.

E) Relative survival of the "high" and "low" expression populations in **D** after treatment with ciprofloxacin for ~3.5 h as in **A,B**. Bars indicate the mean survival ratio of the "high" to "low" expressing populations. The data shown represents 3 independent experiments (paired t-test, p-value ~ 0.009, **). The red dashed line represents a survival ratio of 1, indicating no advantage.

a P_*sasA*-YFP   Phase

b P_*sasA* *mcherry*   P_*sasA* *yfp*
Chromosome

c P_*sasA*-mCherry   P_*sasA*-YFP   Phase

d P_*sasA*-*mcherry* Relative mCherry Fluorescence
Relative YFP Fluorescence
P_*sasA*-*yfp*

a WT
Frequency

— Post-treatment
— Pre-treatment

Fluorescence (A.U.)

b Δ*sasA*
Frequency

— Post-treatment
— Pre-treatment

Fluorescence (A.U.)

c
Enrichment Factor

— WT post/pre-treatment
— Δ*sasA* post/pre-treatment

"High"
"Low"

Fluorescence (A.U.)

d
Frequency

— WT
— Δ*sasA*

"Low"
"High"

Fluorescence (A.U.)
P$_{sasA}$-*yfp*

e
Survival Ratio (High/Low Expression)

**

WT   Δ*sasA*