1

Drug Target Identification and Virtual Screening in Pursuit of Phytochemical Intervention
of *Mycobacterium chelonae*

Zarrin Basharat[a, b], Shumaila Zaib[c], Azra Yasmin[c*], Yigang Tong[d]

[a] Jamil-ur-Rahman Center for Genome Research, Dr. Panjwani Center for Molecular Medicine and Drug Research, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan.

[b] Laboratoire Génomique, Bioinformatique et Applications, Conservatoire National des Arts et Métiers, 75003 Paris, France.

[c] Microbiology & Biotechnology Research Lab, Department of Environmental Sciences, Fatima Jinnah Women University, Rawalpindi 46000, Pakistan.

[d] State Key Laboratory of Pathogen and Biosecurity, Beijing Institute of Microbiology and Epidemiology, Beijing 100071, China.

*Corresponding author

Email: azrayasmin@fjwu.edu.pk

ABSTRACT

Mycobacterium chelonae is a rapidly growing mycobacterium present in the environment. It is associated with skin and soft tissue infections including abscess, cellulitis and osteomyelitis. Other infections by this bacterium are post-operative/transplant-associated, catheter, prostheses and even concomitant to haemodialytic procedures. In this study, we employ a subtractive genomics approach to predict the potential therapeutic candidates, intended for experimental research against this bacterium. A computational workflow was devised and executed to procure core proteome targets essential to the pathogen but with no similarity to the human host. Initially, essential *Mycobacterium chelonae* proteins were predicted through homology searching of core proteome content from 19 different bacteria. Druggable proteins were then identified and N-acetylglucosamine-1-phosphate uridyltransferase (GlmU) was chosen as a case study from identified therapeutic targets, based on its important bifunctional role. Structure modeling was followed by virtual screening of phytochemicals (N > 10,000) against it. 4,4'-[(1E)-5-hydroxy-4-(methoxymethyl)-1-pentene-1,5-diyl]diphenol, apigenin-7-O-beta-gluconopyranoside and methyl rosmarinate were screened as compounds having best potential for binding GlmU. Phytotherapy helps curb the menace of antibiotic resistance so treatment of *Mycobacterium chelonae* infection through this method is recommended.

Key words:

Mycobacterium chelonae, drug design, phytotherapy, GlmU, virtual screening.

INTRODUCTION

Mycobacteria are categorized into two major groups, tubercular and non-tubercular mycobacteria. Nontuberculous mycobacteria are further divided into two groups, rapidly growing and slow growing mycobacteria, depending upon duration of their reproduction in suitable medium (Tortoli, 2014). *Mycobacterium chelonae* is a member of rapidly growing group, which takes less than a week for its reproduction in/on medium. It is the most commonly isolated organism among all rapidly growing mycobacteria. It is mostly found in water sources and on medical instruments such as bronchoscopes (Gonzalez-Santiago and Drage, 2015), and has been isolated from environmental, animal and human sources. *Mycobacterium chelonae* infections in human hosts have increased over time, with reports of both haematogenous and localized occurrences in the recent past (Hay, 2009).

Outbreaks due to *Mycobacterium chelonae* contaminated water and compromised injections are a rising problem. Infections have been linked to cosmetic and surgical procedures, such as trauma, surgery, injection (botulinum toxin, biologics, dermal fillers), liposuction, breast augmentation, under skin flaps, laser resurfacing, skin biopsy, tattoos, acupuncture, body piercing, pedicures, mesotherapy and contaminated foot bath (Gonzalez-Santiago and Drage, 2015). *Mycobacterium chelonae* may also colonize skin wounds, as result of which patients form abscess, skin nodules and sinus tracts (Patnaik *et al.,* 2013).

Recent era has observed a trend for search of drug targets in pathogens using computational methods, with a focus on genomic and proteomic data (Shanmugham and Pan, 2013). Comparative/differential and subtractive genomics along with proteomics has been used by many researchers for the identification of drug targets in various pathogenic bacteria like *Pseudomonas aeruginosa, Helicobacter pylori*, *Brugia malayi*, *Leptospira interrogans*, *Listeria monocytogenes*, *Mycobacterium leprae* (Amineni *et al.,* 2010; Shanmugam and Natarajan, 2010; Sarangi *et al.,* 2015) etc. Determination of potential drug targets has been made possible by the availability of whole genome and their inferred protein complement sequences in public domain databases (Sarangi *et al*., 2015). Numerous anti-tubercular drug targets, even though mostly for *Mycobacterium tuberculosis* have been reported previously. Some of these include proteins essential for survival or the ones that could be targeted by anti-microbial peptides. Common ones include sulphur metabolic pathway proteins (Reviewed by Bhave *et al.* 2007) and folate pathway

proteins (Rengarajan *et al.,* 2004) but due to mutations, folate pathway proteins are prone to drug

resistance as well.

Parenteral antibiotics against *Mycobacterium chelonae* include tobramycin, amikacin, imipenem,

and tigecycline, but it has demonstrated resistance to antibiotics and disinfectants. This property

assists *Mycobacterium chelonae* in colonizing water systems and allows its access to humans

(Jaén-Luchoro *et al.,* 2016). Successful resolution of *Mycobacterium chelonae* infection with

antibiotics has been reported in different scenarios (Brown-Elliott *et al.,* 2001; Choi *et al*., 2018),

but resistance to beta-lactams (Jarlier *et al.,* 1991) and multiple drugs (Churgin *et al*., 2018) has

also been observed. Till now, no specific guidelines for the treatment of *Mycobacterium*

chelonae have been defined in the literature (Gonzalez-Santiago and Drage, 2015). Antibiotic

resistance also illustrates the need to search out new drug targets for design of better therapies

against infection by this bacterium, especially using naturally existing metabolites from microbes

and plants. In the current study, subtractive proteomics was applied to identify essential

druggable proteins in *Mycobacterium chelonae*. Docking of the selected protein GlmU with

phytochemicals was carried out for identification of a candidate which might bind it and stop its

normal cellular function, leading to bacterial lysis/death.

MATERIAL AND METHODS

Prediction of *Mycobacterium chelonae* essential proteome

Complete proteome sequence of *Mycobacterium chelonae* CCUG 47445 with accession no:

NZ_CP007220 was downloaded from the NCBI database. For the prediction of essential

proteins, Geptop (Wei *et al.,* 2013) was installed on computer and a search was carried out to

align the *Mycobacterium chelonae* protein sequences against the essential or core protein

sequences from defined set of 19 bacteria with an essentiality score cut-off value range of 0.15

(Wei *et al.,* 2013). Results were saved and analyzed further according to the workflow (Fig. 1).

Prediction of non-homologous host proteins

In order to find the bacterial proteins which do not have similarity with human host, the set of

essential protein sequences of *Mycobacterium chelonae* was subjected to BLASTP against the

human proteome database (Uniprot release 2014). The standalone BLAST software was used for

this purpose. For identification of non-homologous proteins, an expectation (E-value) cut-off of

10^{-2}, gap penalty of 11 and gap extension penalty of 1 was set as the standard. E-value cut-off

(10^{-2}), based on reported research protocols (Sarangi *et al.,* 2015) was considered.

Identification of putative drug targets

There are several molecular and structural properties which have been explored by researchers for selecting suitable therapeutic targets in pathogenic microorganisms. These properties include determination of molecular weight, sub-cellular localization, 3D structure and druggability analysis. These properties were evaluated for selection of the therapeutic targets in *Mycobacterium chelonae*. Molecular weight was calculated by using computational tools and drug target-associated literature available in the Swiss-Prot database. Subcellular localization of therapeutic targets was predicted using PSORTb (Nancy *et al*., 2010). It uses feature support vector machine-based method and suffix tree algorithm for downstream analysis. Predictions are grouped through a Bayesian scheme into one final (consensus) result. Druggable targets were identified with BLAST hits through unified protocol from the DrugBank. Parameters were: gap cost: -1 in case of extension or opening, mismatch penalty: -3, E-value: $1*10^{-5}$, match: 1, filter algorithm: DUST and SEG (Azam and Shamim, 2014). The targets were subjected to KEGG blast for identification of associated pathways.

Virtual screening of ligand against selected target

Keeping in view the results of sub-cellular localization, molecular weight determination and druggability analysis, an essential protein (GlmU) was chosen for further downstream processing. Swiss Model was used for the prediction of 3D structure of the selected target protein (Biasini et al., 2014). This tool constructs structure model by recognizing structural templates from the PDB using multiple threading alignment approaches (Wang *et al*., 2016). The top structure used for structure prediction was that of GlmU from *Mycobacterium tuberculosis* (PDB ID: 3D8V). The structure was validated and analyzed for quality using SAVES (https://services.mbi.ucla.edu/SAVES/), consisting of ERRAT, VERIFY3D and Ramachandran plot analysis. TM-score was calculated using TM-align (Zhang and Skolnick, 2005).

Phytochemical libraries consisting of three groups were downloaded and docked with GlmU. group I consisted of ~2000 ayurvedic pharmacopia compounds (downloaded from http://ayurveda.pharmaexpert.ru/) (Lagunin et al., 2015), group II had ~6000 North African flora compounds (Ntie-Kang et al., 2017) (downloaded from http://african-compounds.org/nanpdb/) and group III had 2266 phytochemical compounds from various other medicinal plants (available from http://bioinform.info/ in .sdf format) (Mumtaz et al., 2016). Blind docking was carried out using Molecular Operating Environment (MOE) with the parameters: placement:

triangle matcher, rescoring 1: London dG, refinement: forcefield, rescoring 2: affinity dG. MOE

provides fast and accurate docking results based on dedicated algorithms and accurate scoring

functions. Structural preparation program embedded in MOE added the missing hydrogen atoms,

corrected the charges and assigned near precise hybridization state to each residue (Basharat *et*

al., 2017). Compounds with least energy and fulfilling the criteria of Lipinski's drug like test

were considered for further analysis. ADME/Tox analysis was also performed for the top

compounds by inputting the structure at the webserver https://preadmet.bmdrc.kr/.

RESULTS AND DISCUSSION

Essential proteome prediction

Initially, total proteome of *Mycobacterium chelonae* was subjected to core or essential proteins

prediction. Geptop identified these proteins by screening against 19 bacteria (Supplementary

Table 1), based upon orthology and phylogeny features. In the process of essential proteome

mining through homology-based methods, a query protein is considered as essential if it is also

present in another bacterium and experimentally identified as essential for survival. There are

various methods for the prediction of essential proteins, for example single-gene knockout,

transposon mutagenesis and RNA interference but all these methods are time consuming and

laborious. A good alternative is high-efficiency computational methods designed specifically for

this type of work (Wei *et al.,* 2013). Predicted essential proteins were 305 in number

(Supplementary Table 2), linked with significant metabolic pathways in the pathogen life cycle

and necessary for its survival. In order to disrupt the function and existence of pathogen it is

most important to attack those bacterial proteins which regulate important functions (e.g. nutrient

uptake) in the host environment (Butt *et al*., 2012). Latest antimicrobial drugs are designed on

the principle of the inhibition of the pathogen's metabolic pathway. Therefore, such protein

sequences may be considered as possible therapeutic targets.

Identification of non-host proteins

Non-host proteins refer to those bacterial proteins which do not have homology with human

proteins. If the homologous proteins are targeted, they could badly affect the metabolism of host

due to similarity with host proteins. Therefore, non-host proteins could be preferred better drug

targets, as side effects and cross-reactivity caused by the use of antibiotics could be evaded for

harming host (Sarangi *et al*., 2015). Among the core proteome of *Mycobacterium chelonae*, 117

proteins (Supplementary Table 3) indicated 'no hit' against the human proteome according to the

set criteria. These proteins were then used for subsequent analysis.

Drug Target mining and analysis

BLASTp was performed to identify significant drug target from newly selected essential

proteins. Only 17 proteins had significant hits against druggable proteins present in the

DrugBank (Table 1).

Molecular weight determination and druggability analysis could improve the screening process

for therapeutic targets, as observed previously for numerous pathogenic bacteria and fungi

(Abadio *et al.,* 2011). The molecular weight for each potential drug target was calculated (Table

1) and based upon previous studies, it is suggested that smaller proteins are readily soluble and

easier to purify (Duffield *et al.,* 2010).

Sub-cellular localization is a critical factor as it helps in accessing the target gene. Cellular

functions are compartment specific, so if the location of unknown protein is predicted then its

function could also be known which help in selection of proteins for further study. Membrane

proteins are reported as more useful target and more than 60% of the currently known drug

targets are membrane proteins (Tsirigos *et al.,* 2015). Cytoplasm is the site of proteins synthesis

and most of these proteins remain there to carry out their specific functions after synthesis.

However, some proteins need to be transported to different cellular compartments for their

specific function (Strzyz, 2016). The subcellular localization of non-host proteins of

Mycobacterium chelonae was predicted and majority were demarcated as cytoplasmic (Table 1).

GlmU analysis and phytochemical screening

After the characterization of all druggable proteins of *Mycobacterium chelonae,* GlmU was

selected for further analysis (out of 305 essential and 117 non-homologous proteins). It is a

bifunctional enzyme, exhibiting both acetyltransferase and uridyltransferase activities (Sharma *et*

al., 2016) and involved in biosynthesis of peptidoglycan. Peptidoglycan in complex with

mycolyl-arabinogalactan is important for the viability of Mycobacteria as it upkeeps the basal

structure necessary for upper "myco-membrane" support. GlmU has been analyzed for various

bacterial species, including *Escherichia coli*, *Streptococcus pneumonia*, *Haemophilus influenzae*

and *Mycobacterium tuberculosis* (Patin *et al*., 2015). Predicted structure of GlmU had an

estimated RMSD of 0.5 Å from the reference structure. A TM-score of 0.97 was obtained and

since a score of 0.5-1 means same fold, with greater value meaning more similarity. 0.97 meant a

very similar fold structure. Structure was found to be a homotrimer. Ramachandran plot showed

97.2% residues in favored regions (90.4% in core, 9.4% in allowed, 0.3% in generously allowed)

and no residue in disallowed region (Fig. 2). According to VERIFY3D program, at least 80% of

the amino acids should have value >= 0.2 in the 3D/1D profile and 96.55% of predicted GlmU

residues had an averaged 3D-1D score >= 0.2, thus passing the quality check. An ERRAT score

of 90.66 was obtained.

Each monomer of GlmU consists of two domains: N and C-terminal domains. N-terminal

domain has α/β like fold, similar to dinucleotide-binding Rossmann fold topology. C-terminal

domain exhibits a regular left-handed β-helix conformation and a long α-helical arm connecting

both domains (Sharma *et al*., 2016). N-terminal domain is essential for uridyltransferase activity

as it catalyses the transfer of uridine monophosphate from uridine-triphosphate to N-

acetylglucosamine-1-phosphate (GlcNAc1P). C-terminal domain has acetyltransferase activity as

it catalyzes the transfer of an acetyl group from acetyl-CoA coenzyme to GlcN1P, in order to

produce GlcNAc1P (Moraes *et al.,* 2015).

GlmU plays fundamental role in the formation of bacterial cell wall by carrying out catalysis of

uridine-diphospho-N-acetylglucosamine, an important precursor in bacterial peptidoglycan cell

wall (Sharma *et al*., 2016). Activity of acetyltransferase of eukaryotic cells differ from the

activity of GlmU in the way that, eukaryotic acetyltransferase occurs on GlcN6P and not on

GlcN1P. These properties make GlmU suitable as the drug target (Moraes *et al.,* 2015) and it has

been used for drug targeting in bacteria such as *Haemophilus influenza* (Mochalkin *et al.,* 2007)

apart from designing inhibitors against it for Mycobacterial species (Mehra *et al.,* 2016). It is

also known that proteins that are involved in more than one pathway of pathogen, in addition to

that they are non-host proteins, could be more effective drug targets (Sarangi *et al*., 2015).

Inactivation of bifunctional GlmU enzyme leads to loss of mycobacterial viability, therefore

GlmU was used for docking against phytochemicals.

The possible interaction between protein and the ligand is understood computationally through

molecular docking therefore receptor centric docking approach was employed for screening of

prospective phytochemical library of compounds from more than 10,000 plant derived

compounds against GlmU of *Mycobacterium chelonae.* Whole surface of the protein was

scanned for best docking pose. A comparative analysis of structural shape and chemical

complementarity to GlmU was ranked, based on S value and Lipinski's drug likeliness in MOE.

Ones with least S-value and Lipinski's drug likeliness=1 were considered as best inhibitors.
Results of GlmU binding with compounds from group I, II and II showed several interactions
(Fig. 3, Table 2). 4,4'-[(1E)-5-Hydroxy-4-(methoxymethyl)-1-pentene-1,5-diyl]diphenol from
group I, apigenin-7-O-beta-guconopyranoside (also commonly known as apigetrin) from group
II and methyl rosmarinate was obtained as the best anti-microbial against *Mycobacterium*
chelonae from group III compounds. Common source of 4,4'-[(1E)-5-Hydroxy-4-
(methoxymethyl)-1-pentene-1,5-diyl]diphenol is *Alpinia galanga* while apigetrin is sourced from
dandelion coffee and *Teucrium gnaphalodes.* Methyl rosmarinate is a constituent of *Rabdosia*
serra, Ehretia thyrsiflora, Dracocephalum heterophyllum, Perilla frutescens, Nepeta deflersiana,
Cordia sinensis and *Hyptis atrorubens* Poit. ADME and toxicity parameters were also calculates
and it was found that apigenin-7-O-beta-gluconopyranoside had lowest blood brain penetration
but predicted as a mutagen while methyl rosmarinate and 4,4'-[(1E)-5-Hydroxy-4-
(methoxymethyl)-1-pentene-1,5-diyl]diphenol were found to be non-mutagenic and had slightly
higher blood brain barrier penetration (Table 3). Analogs of apigenin have shown activity against
Mycobacterial species (Tsouh Fokou *et al*., 2015; Evina *et al*., 2017) while other compounds
have not been reported in literature as anti-mycobacterials yet.
We focused on phytochemical screening against *Mycobacterium chelonae* GlmU because plant
derived/natural compounds could be used as antibacterial therapeutics for treatment of bacterial
infections. Further Lab testing is proposed to know about minimum inhibitory concentration
value and other parameters for these compounds against *Mycobacterium chelonae.*

CONCLUSION
In this study, we proposed GlmU as one of the important therapeutic drug targets for
Mycobacterium chelonae as it is bifunctional, essential protein for pathogen and has no
homology with human proteome. We have provided putative model for phytotherapy against
Mycobacterium chelonae through virtual screening-based identification of potent metabolites
from three data sets, comprising of more than 10,000 compounds. This study could be taken as
an initiative for screening and quick designing of phytotherapy against microbes using a
computational *modus operandi*. It is expected that our study will also facilitate selection and
screening of other *Mycobacterium chelonae* therapeutic target proteins against phytochemical
and other relevant compounds for Lab testing and successful entry into drug design pipeline.

ACKNOWLEDGEMENTS

The authors are thankful to Dr. Waqasuddin Khan (Jamil-ur-Rahman Center for Genome
Research, PCMD, ICCBS, University of Karachi) for useful comments and insights that
facilitated the analysis. The authors are also thankful to Dr. Alexandra Elbakyan for help with
literature.

DECLARATION OF INTEREST

The authors declare that they have no conflict of interest.

FUNDING

This research did not receive any specific grant from funding agencies in the public, commercial,
or not-for-profit sectors.

FIGURE/TABLE LEGENDS

Fig. 1. Workflow describing subtractive proteomics with respective number of obtained

sequences. *In silico* parameter analysis included hydrophobicity, sub-cellular location prediction,

molecular weight etc.

Fig. 2. Ramachandran plot of the predicted GlmU structure.

Fig. 3. (A) Predicted 3D structure of GlmU **(B)** Docked GlmU with 4,4'-[(1E)-5-Hydroxy-4-

(methoxymethyl)-1-pentene-1,5-diyl]diphenol showing residue interactions (2D conformation)

(C) apigenin-7-O-beta-guconopyranoside and GlmU docked complex(2D conformation) **(D)**

Methyl rosmarinate and GlmU docked complex(2D conformation). Blue arrowhead represents

backbone donor/acceptor while green arrowhead depicts sidechain donor/acceptor, dotted line

represents proximity contour.

Table 1. Predicted druggable proteome of *Mycobacterium chelonae*. GRAVY is average

hydrophobicity of the protein measured by Kyte-Doolittle algorithm. Hydrophobicity value

below 0 are indicates globularity (means protein is hydrophilic), while score value of above 0

indicates protein to be membranous (hydrophobic).

Table 2. Top 5 compounds against GlmU of *Mycobacterium chelonae* (from each docked group
obtained with best inhibition score).

Table 3. ADME and toxicity profile of top drug-like molecules from studied phytochemical

group compounds.

Supplementary Table 1. List of bacterial species used for essential proteome prediction of

Mycobacterium chelonae.

Supplementary Table 2. Essential proteome of *Mycobacterium chelonae.*

Supplementary Table 3. Non-homologous core proteome of *Mycobacterium chelonae.*

Fig. 1.

324   Fig. 2.

326   **Fig. 3.**

343 **Table 1. Predicted druggable proteome of *Mycobacterium chelonae*. GRAVY is average**

344 **hydrophobicity of the protein measured by Kyte-Doolittle algorithm. Hydrophobicity value**

345 **below 0 are indicates globularity (means protein is hydrophilic), while score value of above**

346 **0 indicates protein to be membranous (hydrophobic).**

S. No.	Accession #	Protein name	No. of Matches	Mol. Wt.	Length (amino acids)	GRAVY	Sub cellular localization	KEGG Pathway	Experimental evidence for druggability in bacteria
	WP_0300 93515	6,7-dimethyl-8-ribityllumazine synthase/riboflavin synthase	1	16775.9	162	0.173	Unknown	Riboflavin metabolism, biosynthesis of secondary metabolites	*Zhao et al.*, 2009
	WP_0300 95371	acyl carrier protein	1	10805.0	99	-0.207	Cytoplasmic	Fatty acid biosynthesis	*Luckner et al.*, 2010
	WP_0300 97637	replicative DNA helicase	1	50372.3	458	-0.323	Cytoplasmic membrane	DNA replication	*Aiello et al.*, 2009
	WP_0462 51970	DNA topoisomerase IV subunit B	3	74147.8	675	-0.371	Cytoplasmic	DNA repair and recombination	*Li et al.*, 2016
	WP_0462 52393	dihydropteroate synthase	2	37365.3	350	-0.165	Cytoplasmic	Folate biosynthesis	*Zhao et al.*, 2016
	WP_0462 52395	2-amino-4-hydroxy-6-hydroxymethyldihydropteridine diphosphokinase	1	19332.9	178	-0.054	Cytoplasmic	Tetrahydrofolate biosynthesis	*Shi et al.*, 2001
	WP_0462 52399	pantoate - beta-alanine ligase/pantothenate synthetase	1	33527.3	314	-0.057	Cytoplasmic	Pantothenate biosynthesis	*Yang et al.*, 2011
	WP_0462 52764	3-oxoacyl-ACP synthase/beta-ketoacyl acyl carrier protein synthase	2	35485.1	340	0.070	Cytoplasmic	Lipid synthesis	*Lee et al.*, 2012
	WP_0462 52769	bifunctional N-acetylglucosamine-1-phosphate uridyltransferase (GlmU)/glucosami	1	50208.3	482	-0.026	Cytoplasmic	Amino sugar and nucleotide sugar metabolism, Biosynthesis	*Mehra et al.*, 2015

	ne-1-phosphate acetyltransferase						of antibiotics	
WP_0462 53413	cell division protein FtsI	1	68842.9	642	-0.300	Cytoplas mic membrane	Peptidoglycan biosynthesis, beta-Lactam resistance	Slayden and Belisle, 2009
WP_0462 53421	cell division protein FtsZ	1	38795.6	383	0.076	Cytoplas mic	Chromosome partition, cytoskeleton, toxin-anti-toxin	Park *et al.*, 2014
WP_0462 53835	enoyl-[acyl-carrier-protein] reductase (InhA)	3	28724.0	269	0.152	Cytoplas mic membrane	Lipid biosynthesis	Luckner *et al.*, 2010
WP_0462 54273	DNA ligase (NAD(+)) LigA	1	74430.9	684	-0.226	Cytoplas mic	DNA replication, Base excision repair, nucleotide excision repair, mismatch repair	Mills *et al.*, 2011
WP_0462 54500	thymidylate kinase	1	23403.1	212	-0.394	Cytoplas mic	Pyrimidine metabolism	Kosinska *et al.*, 2005
WP_0462 54518	biotin--[acetyl-CoA-carboxylase] ligase	1	27934.5	264	-0.084	Cytoplas mic	Biotin metabolism	Soares da Costa *et al.*, 2012
WP_0462 54614	alanine racemase	1	41861.3	397	-0.008	Cytoplas mic	D-Alanine metabolism, Vancomycin resistance	Anthony *et al.*, 2011
WP_0462 55923	ribonucleotide-diphosphate reductase subunit beta/ ribonucleoside diphosphate reductase-β	2	36899.2	320	-0.437	Unknown	Purine and pyrimidine metabolism	Tholander and Sjöberg, 2012

347
348
349

Table 2. Top 5 compounds against GlmU of *Mycobacterium chelonae* (from each docked group obtained with best inhibition score).

Group I		Group II		Group III	
Compound	S-value	Compound	S-value	Compound	S-value
4,4'-[(1E)-5-Hydroxy-4-(methoxymethyl)-1-pentene-1,5-diyl]diphenol	-6.63	Apigenin-7-O-beta-gluconopyranoside	-6.35	Methyl rosmarinate	-12.33
1-(3,4-dihydroxy-5-prenylphenyl)-2-(3,5-dihydroxyphenyl)ethane	-6.49	Cosmetin	-5.88	5,7,3',4',5'-Pentahydroxyflavanone	-12.23
4,4',7,9-tetrahydroxy-8->9'-neolign-7'-ene	-6.35	Anastatin B	-5.77	Peonidin	-11.95
1,7-bis(3,4-dihydroxyphenyl)-1,4,6-heptatrien-3-one	-5.79	Ixerisoside D	-5.66	Flowerone	-11.92
Nimocinolide	-5.77	Quercetin	-5.59	Hovenitin III	-11.82

367 **Table 3.** ADME and toxicity profile of top drug-like molecules from studied phytochemical
368 group compounds.

Parameter	4,4'-[(1E)-5-Hydroxy-4-(methoxymethyl)-1-pentene-1,5-diyl]diphenol	Apigenin-7-O-beta-gluconopyranoside	Methyl rosmarinate
In vivo blood-brain barrier penetration	1.90	0.03	0.17
CYP450 2D6 inhibition	None	None	None
Human intestinal absorption (%)	90.47	47.10	76.88
Plasma protein binding (%)	89.08	73.43	86.57
Ames test	Non-mutagen	mutagen	Non-mutagen
Skin permeability (logKp, cm/hour)	-2.79	-4.52762	-3.30
hERG inhibition	Medium risk	High risk	Medium risk
Carcinogenicity in mouse	Negative	Positive	Negative
Carcinogenicity in rat	Negative	Negative	positive

REFERENCES

Abadio, A. K. R., Kioshima, E. S., Teixeira, M. M., Martins, N. F., Maigret, B. & Felipe, M.

S. (2011). Comparative genomics allowed the identification of drug targets against human

fungal pathogens. *BMC Genomics,* 12(1),1.

Aiello, D., Barnes, M. H., Biswas, E. E., Biswas, S. B., Gu, S., Williams, J. D., Bowlin, T. L.

& Moir, D. T. (2009). Discovery, characterization and comparison of inhibitors of *Bacillus*

anthracis and *Staphylococcus aureus* replicative DNA helicases. *Bioorganic and Medical*

Chemistry, 17(13), 4466-76.

Amineni, U., Pradhan, D. & Marisetty, H. (2010) *In silico* identification of common putative

drug targets in *Leptospira interrogans. Journal of Chemical Biology,* 3, 165–73.

Anthony, K. G., Strych, U., Yeung, K. R., Shoen, C. S., Perez, O., Krause, K. L., Cynamon,

M. H., Aristoff, P. A. & Koski, R. A. (2011). New classes of alanine racemase inhibitors

identified by high-throughput screening show antimicrobial activity against *Mycobacterium*

tuberculosis. PLoS One, 6(5), e20374.

Azam, S.S. & Shamim, A. (2014). An insight into the exploration of druggable genome of

Streptococcus gordonii for the identification of novel therapeutic candidates. *Genomics,*

104(3), 203-14.

Basharat, Z., Zaib, S. & Yasmin A. (2017). Computational study of some amoebicidal

phytochemicals against heat shock protein of *Naegleria fowleri. Gene Reports,* 1(6), 158-62.

Biasini, M., Bienert, S., Waterhouse, A., Arnold, K., Studer, G., Schmidt, T., Kiefer, F.,

Cassarino, T.G., Bertoni, M., Bordoli, L. & Schwede, T. (2014). SWISS-MODEL: modelling

protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids*

Research, 42(W1), W252-W258.

Bhave, D. P., Wilson III, B., & Carroll, K. S. (2007). Drug targets in mycobacterial sulfur

metabolism. *Infectious Disorders-Drug Targets*, 7(2), 140-158.

Brown-Elliott, B.A., Wallace Jr, R.J., Blinkhorn, R., Crist, C.J. & Mann, L.B. (2001).

Successful treatment of disseminated *Mycobacterium chelonae* infection with linezolid.

Clinical Infectious Diseases, 33(8),1433-4.

Butt, A. M., Tahir, S., Nasrullah, I., Idrees, M., Lu, J. & Tong, Y. (2012). *Mycoplasma*

genitalium: A comparative genomics study of metabolic pathways for the identification of

drug and vaccine targets. *Infection Genetics and Evolution,* 12, 53–62.

Choi, Y.J., Oh, E.K., Lee, M.J., Kim, N., Choung, H.K., Khwarg, S.I. (2018). Atypical

mycobacterial infection in anophthalmic sockets with porous orbital implant exposure.

American Journal of Ophthalmology, In press.

Churgin, D.S., Tran, K.D., Gregori, N.Z., Young, R.C., Alabiad, C. & Flynn Jr H.W. (2018).

Multi-drug resistant *Mycobacterium chelonae* scleral buckle infection. American Journal of

Ophthalmology Case Reports, 10:276-8.

Duffield, M., Cooper, I., McAlister, E., Bayliss, M., Ford, D. & Oyston, P. (2010). Predicting

conserved essential genes in bacteria: *in silico* identification of putative drug targets.

Molecular Biosystems, 6, 2482–89.

Evina, J.N., Bikobo, D.S., Zintchem, A.A., Nyemeck II, Mbabi, N., Ndedi, E.D., Diboué,

P.H., Nyegue, M.A., Atchadé, A.D., Pegnyemb, D.E. & Koert, U. (2017). *In vitro*

antitubercular activity of extract and constituents from the stem bark of *Disthemonanthus*

benthamianus. *Revista Brasileira de Farmacognosia*, 27(6), 739-43.

Gonzalez-Santiago, T. M. & Drage, L. A. (2015). Nontuberculous *Mycobacteria:* Skin and

soft tissue infections. *Dermatologic Clinics,* 33(3), 563-577.

Hay, R. J. (2009). *Mycobacterium chelonae* -a growing problem in soft tissue infection.

Current Opinion in Infectious Diseases, 22, 99–101.

Jaén-Luchoro, D., Salvà-Serra, F., Aliaga-Lozano, F., Seguí, C., Busquets, A., Ramírez, A.,

Ruíz, M., Gomila, M., Lalucat, J. & Bennasar-Figueras, A. (2016). Complete genome

sequence of *Mycobacterium chelonae* type strain CCUG 47445, a rapidly growing species of

nontuberculous mycobacteria. *Genome Announcements,* 4(3), e00550-16.

Jarlier VI, Gutmann, L. & Nikaido, H. (1991). Interplay of cell wall barrier and beta-

lactamase activity determines high resistance to beta-lactam antibiotics in *Mycobacterium*

chelonae. *Antimicrobial Agents and Chemotherapy.* 35(9), 1937-9.

Kosinska, U., Carnrot, C., Eriksson, S., Wang, L. & Eklund, H. (2005). Structure of the

substrate complex of thymidine kinase from *Ureaplasma urealyticum* and investigations of

possible drug targets for the enzyme. *FEBS Journal,* 272(24), 6365-6372.

Lagunin, A.A., Druzhilovsky, D.S., Rudik, A.V., Filimonov, D.A., Gawande, D., Suresh, K.,

Goel, R., Poroikov, V.V. (2015). Computer evaluation of hidden potential of phytochemicals

of medicinal plants of the traditional Indian ayurvedic medicine. *Biomeditsinskaia Khimiia*,

61(2), 286-97.

Lee, J. Y., Jeong, K. W., Shin, S., Lee, J. U. & Kim, Y. (2012). Discovery of novel selective

inhibitors of *Staphylococcus aureus* β-ketoacyl acyl carrier protein synthase III. *European*

Journal of Medical Chemistry, 47, 261-269.

Li, Y., Wong, Y. L., Ng, F. M., Liu, B., Wong, Y. X., Poh, Z. Y., Liu, S., Then, S. W., Lee,

M. Y., Ng, H. Q. & Huang, Q. (2016). *Escherichia coli* topoisomerase IV E subunit and an

inhibitor binding mode revealed by NMR spectroscopy. *Journal of Biological Chemistry,*

291(34), 17743-17753.

Luckner, S. R., Liu, N., Am, Ende, C.W., Tonge, P.J. & Kisker, C. (2010). A slow, tight

binding inhibitor of InhA, the enoyl-acyl carrier protein reductase from *Mycobacterium*

tuberculosis. Journal of Biological Chemistry, 285(19), 14330-14337.

Mehra, R., Rani, C., Mahajan, P., Vishwakarma, R. A., Khan, I. A & Nargotra, A. (2016).

Computationally guided identification of novel *Mycobacterium tuberculosis* GlmU inhibitory

leads, their optimization and *in vitro* validation. *ACS Combinatorial Science*, 18(2), 100-116.

Mehra, R., Sharma, R., Khan, I. A. & Nargotra, A. (2015). Identification and optimization of

Escherichia coli GlmU inhibitors: an *in silico* approach with validation thereof. *European*

Journal of Medical Chemistry, 92, 78-90.

Mills, S. D., Eakin, A. E., Buurman, E. T., Newman, J. V., Gao, N., Huynh, H., Johnson,

K.D., Lahiri, S., Shapiro, A. B., Walkup, G. K. & Yang, W. (2011). Novel bacterial NAD+-

dependent DNA ligase inhibitors with broad-spectrum activity and antibacterial efficacy *in*

vivo. *Antimicrobial Agents Chemotherapy*, 55(3),1088-1096.

Mochalkin, I., Lightle, S., Narasimhan, L., Bornemeier, D., Melnick, M., VanderRoest, S.,

McDowell L (2008). Structure of a small-molecule inhibitor complexed with GlmU from

Haemophilus influenzae reveals an allosteric binding site. *Protein Science*, 17, 577-82.

Moraes, G. L., Gomes, G. C., deSousa, P. R., Alves, C. N., Govender, T., Kruger, H. G.,

Maguire, G. E., Lamichhane, G. & Lameira, J. (2015). Structural and functional features of

enzymes of *Mycobacterium tuberculosis* peptidoglycan biosynthesis as targets for drug

development. *Tuberculosis*, 95, 95-111.

Mumtaz, A., Ashfaq, U. A., ulQamar, M. T., Anwar, F., Gulzar, F., Ali, M. A., Saari, N.&

Pervez, M. T. (2016). MPD3: a useful medicinal plants database for drug designing. *Natural*

Product Research, 1-9.

Nancy, Y. Y., Wagner, J. R., Laird, M. R., Melli, G., Rey, S., Lo, R., Dao, P., Sahinalp, S.

C., Ester, M., Foster, L. J. & Brinkman, F. S. (2010). PSORTb 3.0: improved protein

subcellular localization prediction with refined localization subcategories and predictive

capabilities for all prokaryotes. *Bioinformatics*, 26, 1608–15.

Ntie-Kang, F., Telukunta, K.K., Do☐ring, K., Simoben, C.V.A., Moumbock, A.F., Malange,

Y.I., Njume, L.E., Yong, J.N., Sippl, W. & Gu☐nther, S. (2017). NANPDB: A resource for

natural products from Northern African sources. *Journal of Natural Products.* 80(7), 2067-

76.

Park, B., Awasthi, D., Chowdhury, S. R., Melief, E. H., Kumar, K., Knudson, S. E., Slayden,

R. A. & Ojima, I. (2014). Design, synthesis and evaluation of novel 2, 5, 6-trisubstituted

benzimidazoles targeting FtsZ as antitubercular agents. *Bioorganic and Medicinal Chemistry*,

22(9), 2602-2612.

Patin, D., Bayliss, M., Lecreulx, D. M., Oyston, P. & Blanot, D. (2015). Purification and

biochemical characterisation of GlmU from *Yersinia pestis.* *Archives Microbiology*,

197(3),371-78.

Patnaik, S., Mohanty, I., Panda, P., Sahu, S. & Dash, M. (2013). Disseminated

Mycobacterium chelonae infection: Complicating a case of hidradenitis suppurativa. *Indian*

Dermatology Online Journal, 4(4), 336-396.

Rengarajan, J., Sassetti, C. M., Naroditskaya, V., Sloutsky, A., Bloom, B. R., & Rubin, E. J.

(2004). The folate pathway is a target for resistance to the drug para-aminosalicylic acid

(PAS) in mycobacteria. *Molecular Microbiology*, 53(1), 275-282.

Sarangi, A. N., Lohani, M. & Aggarwal, R. (2015). Proteome mining for drug target

identification in *Listeria monocytogenes* EGD-e and structure based virtual screening of a

candidate drug target penicillin binding protein. *Journal of Microbiological Methods*, 111,9-

18.

Shanmugam, A. & Natarajan, J. (2010). Computational genome analyses of metabolic

enzymes in *Mycobacterium leprae* for drug target identification. *Bioinformation*, 4,392–95.

Shanmugham, B. & Pan, A. (2013). Identification and characterization of potential

therapeutic candidates in emerging human pathogen *Mycobacterium abscessus:* A novel

hierarchical *in silico* approach. *PLOS One,* 8(3), 59126.

Sharma, R., Lambu, M. R., Jamwa, U., Rani, C., Chib, R. & Wazir, P. (2016). *Escherichia*

coli N-Acetylglucosamine-1- Phosphate-Uridyltransferase/ Glucosamine-1- Phosphate-

Acetyltransferase (GlmU) inhibitory activity of terreic acid isolated from *Aspergillus terreus.*

Journal of Biomolecular Screening, 1-12.

Shi, G., Blaszczyk, J., Ji, X. & Yan, H. (2001). Bisubstrate analogue inhibitors of 6-

hydroxymethyl-7, 8-dihydropterin pyrophosphokinase: synthesis and biochemical and

crystallographic studies. *Journal of Medicinal Chemistry*, 44(9), 1364-1371.

Slayden, R. A. & Belisle, J. T. (2009). Morphological features and signature gene response

elicited by inactivation of FtsI in Mycobacterium tuberculosis. *Journal of Antimicrobial*

Chemotherapy, 63(3), 451-457.

Soares da Costa, T. P., Tieu, W., Yap, M. Y., Zvarec, O., Bell, J. M., Turnidge, J.D., Wallace, J. C., Booker, G. W, Wilce, M. C., Abell, A. D. & Polyak, S.W. (2012). Biotin analogues with antibacterial activity are potent inhibitors of biotin protein ligase. *ACS Medicinal Chemistry Letters*, 3(6), 509-514.

Strzyz, P. (2016). Protein translocation channelling the route for ER misfolded proteins. *Nature Reviews Molecular Cell Biology*, 17, 462–463.

Tholander, F.& Sjöberg, B. M. (2012). Discovery of antimicrobial ribonucleotide reductase inhibitors by screening in microwell format. *Proceedings of the National Academy of Sciences*, 109(25), 9798-9803.

Tortoli, E. (2014). Microbiological features and clinical relevance of new species of the genus *Mycobacterium. Clinical Microbiology Reviews*, 27(4),727-752.

Tsirigos, K. D., Peters, C., Shu, N., Käll, L. & Elofsson, A. (2015). The TOPCONS web server for consensus prediction of membrane protein topology and signal peptides. *Nucleic Acids Research*, 485

Tsouh Fokou, P. V., Nyarko, A. K., Appiah-Opong, R., Yamthe, T., Rachel, L., Ofosuhene, M., & Boyom, F. F. (2015). Update on medicinal plants with potency on *Mycobacterium ulcerans. BioMed Research International,* 917086, 1-16.

Wang, Z., Sun, H., Yao, X., Li, D., Xu, L., Li, Y., Tian, S. & Hou, T. (2016). Comprehensive evaluation of ten docking programs on a diverse set of protein–ligand complexes: the prediction accuracy of sampling power and scoring power. *Journal of Physical Chemistry*, 18, 12964-75.

Wei, W., Ning, L. W., Ye, Y. N. & Guo, F. B. (2013). Geptop: A Gene Essentiality

Prediction Tool for Sequenced Bacterial Genomes Based on Orthology and Phylogeny.

PLOS One, 8(8), 72343.

Zhang, Y. & Skolnick, J. (2005). TM-align: a protein structure alignment algorithm based on

the TM-score. *Nucleic Acids Research*, *33*(7), 2302-2309.

Zhao, Y., Bacher, A., Illarionov, B., Fischer, M., Georg, G., Ye, Q. Z., Fanwick , P. E.,

Franzblau, S. G., Wan, B. & Cushman, M. (2009). Discovery and development of the

covalent hydrates of trifluoromethylated pyrazoles as riboflavin synthase inhibitors with

antibiotic activity against *Mycobacterium tuberculosis. Journal of Organic Chemistry*,

74(15),5297-5303.

Zhao, Y., Shadrick, W. R., Wallace, M. J., Wu, Y., Griffith, E.C., Qi, J., Yun, M. K., White,

S. W. & Lee, R. E. (2016). Pterin–sulfa conjugates as dihydropteroate synthase inhibitors and

antibacterial agents. *Bioorganic and Medicinal Chemistry Letters*, 26(16), 3950-3954.

Zhuo, H. S., Ni, H. H., Yun, M. Q., He, M. M., Liang, Z. M., Dai Hao Fu, J. Y., Hai, W. Q.

& Xing, Z. Y. (2015). The phytochemicals with antagonistic activities toward pathogens of a

disease complex caused by *Meloidogyne incognita* and *Ralstonia solanacearum. Journal of*

Pure and Applied Microbiology, 9, 209-13.