Title: Spatial variation in exploited metapopulations obscures risk of collapse

Authors: Daniel K Okamoto[a,b,c]*, Margot Hessing-Lewis[b], Jameal F Samhouri[d], Andrew O Shelton[d], Adrian Stier[e], Philip S Levin[f,g] and Anne K Salomon[b,c]

a: Department of Biological Science, Florida State University

b: Hakai Institute

c: School of Resource and Environmental Management, Simon Fraser University

d: Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration

e: Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara

f: The Nature Conservancy

g: School of Environment and Forestry Sciences, University of Washington

*corresponding author: dokamoto@bio.fsu.edu

Running Head: Cryptic collapses in metapopulations

Author contributions:

DKO designed, built and conducted analyses. AS conducted the home-range literature review. DKO and AKS wrote the initial manuscript draft. DKO, MHL, JFS, AOS, AS, PSL, and AKS initiated the research focus, refined analyses and contributed to revisions.

Abstract

Unanticipated declines among exploited species have commonly occurred despite harvests that

appeared sustainable prior to collapse. This is particularly true in the oceans where spatial scales

of management are often mismatched with spatially complex metapopulations. We explore

causes, consequences and potential solutions for spatial mismatches in harvested

metapopulations in three ways. First, we generate novel theory illustrating when and how

harvesting metapopulations increases spatial variability and in turn masks local scale volatility.

Second, we illustrate why spatial variability in harvested metapopulations leads to negative

consequences using an empirical example of a Pacific herring metapopulation. Finally, we

construct a numerical management strategy evaluation model to identify and highlight potential

solutions for mismatches in spatial scale and spatial variability. Our results highlight that spatial

complexity can promote stability at large scales, however ignoring spatial complexity produces

cryptic and negative consequences for people and animals that interact with resources at small

scales. Harvesting metapopulations magnifies spatial variability, which creates discrepancies

between regional and local trends while increasing risk of local population collapses. Such

effects asymmetrically impact locally constrained fishers and predators, which are more exposed

to risks of localized collapses. Importantly, we show that dynamically optimizing harvest can

minimize local risk without sacrificing yield. Thus, multiple nested scales of management may

be necessary to avoid cryptic collapses in metapopulations and the ensuing ecological, social and

economic consequences.

Keywords: Sustainable fisheries, Metapopulation, Population Dynamics, Conservation, Natural

Resource Management

Introduction

Mismatches in spatial scale create pervasive problems in ecology and natural resource management (Cumming et al. 2006, Cope and Punt 2011). This problem occurs in part because the spatial extent of management or conservation units is often defined by history, jurisdictional, or institutional criteria rather than the scale of social and ecological processes at play (Levin 1992, Chesson 1998, Cumming et al. 2006). Such choices concerning the scale of management can result in spatial mismatches, where feedbacks controlling interactions among groups occur at different scales. In managed ecosystems like forestry and fisheries, mismatches may occur when harvest recommendations are based on trends in large-scale abundance without accounting for localized collapses (Johnson et al. 2012) or spatial variation in population structure and harvest rates (Cope and Punt 2011). Yet these spatially isolated collapses can have far-reaching consequences when the species play an indispensable role in local social-ecological systems, including human communities with limited capacity to forage over wide geographic scales. Empirical identification of appropriate spatial scales of management remains difficult for spatially structured populations, but can be a pre-requisite for diagnosing and reconciling challenges that spatial mismatches impose on the sustainable and equitable use of natural resources.

In metapopulations – plant or animal populations connected through dispersal – it is well established that the dynamics of individual populations can differ substantially from the aggregate metapopulation (Chesson 1998, Mangel and Levin 2005, Melbourne and Chesson 2006). A combination of movement, shared climate drivers, and compensatory processes determine whether dynamics of individual populations reflect the dynamics of the aggregate

metapopulation (Kendall et al. 2000). Consideration of metapopulation structure has improved

the management of the spotted owl (*Strix occidentalis*) (Lande 1988), salmon (*Onchorhyncus*

spp.) (Stephenson 1999, Rieman and Dunham 2000, Schtickzelle and Quinn 2007, Peterson et al.

2014), amphibians (Marsh and Trenham 2001), and mosquitoes (Adams and Kapan 2009). To

date, efforts to integrate metapopulation dynamics into natural resource management have

largely focused on either minimizing the risk of localized extinction (Chadès et al. 2011),

characterizing productivity of the metapopulation as a whole (sensu Takashina and Mougi 2015),

or valuing benefits of portfolio effects (i.e. stabilizing effects of spatial asynchrony sensu

Schindler et al. 2010). Less understood, both in theory and in practice, is if and when harvesting

metapopulations can increase spatial variability that yield mismatches in spatial scales of

management and population dynamics.

In this study, we assess how harvest dynamics interact with animal movement and recruitment to

shape spatial population variability and risk of collapse at different spatial scales. Our results

illustrate the challenge in managing spatially complex populations using three complementary

approaches. First, we develop new theory using a stochastic analytical model to examine when

and how harvesting in a metapopulation amplifies spatial variability that can create mismatches

in spatial scale. Second, we contextualize the problem of spatial variability and mismatches in

spatial scale by presenting historical analyses from spatial Pacific herring (*Clupea pallasii*)

fisheries in British Columbia, Canada and home-ranges of associated predators and fishers that

may be impacted by localized collapses. Finally, we evaluate how and when different harvest

management approaches can ameliorate such mismatches using a numerical risk analysis in

Pacific herring fisheries.

Pacific herring case study in British Columbia's Central Coast

Pacific herring exemplify the challenges inherent to managing metapopulations that exhibit

spatial variation in population trends. In British Columbia (BC), Canada, herring return annually

to nearshore coastlines in the late winter/early spring to reproduce. During this annual migration,

they are harvested and preyed upon by a range of consumers. Mobile commercial fishing fleets

harvest adult herring, largely for their roe, in the days prior to spawning. In contrast, Indigenous

fishers are constrained to a local area and largely harvest eggs after spawning events (though

some adults as well) as an important food, trade, and cultural resource (Lepofsky and Caldwell

2013, McKechnie et al. 2014, Department of Fisheries and Oceans 2015, von der Porten et al.

2016, Okamoto et al. 2019). These activities create trade-offs among commercial roe fisheries

that remove spawning adults, which truncates adult age structure and reduces abundance, versus

those that remove only eggs from shorelines (Shelton et al. 2014). Unfortunately, a core

uncertainty for herring management, as for many species, is the extent of movement between

areas (Flostrand et al. 2009, Benson et al. 2015, Jones et al. 2016, Levin et al. 2016)**.** Similar

uncertainty surrounds spatial variation in spawning biomass (Siple and Francis 2016), which

may result in part from the degree of demographic synchrony between areas (e.g. synchrony in

recruitment). Pacific herring in British Columbia are currently managed as stocks at regional

scales (100s of kilometers) by Canada's federal fisheries agency. Within these stocks, multiple

spawning aggregations (substocks) seasonally occupy individual stretches of coastline, many of

which are of important traditional and cultural value to Indigenous groups. Thus, Pacific herring

fisheries present a valuable system in which to explore how spatial population dynamics, scales

of management, and spatial constraints of fishers and predators interact to influence differential

risk exposure to population collapses.

Methods

Methodological Overview

We used three distinct modeling approaches in this study to explore how harvest can affect

spatial dynamics in metapopulations. In **Model 1,** we developed a novel analytical approach to

modeling stochastic age-structured metapopulations to illustrate how and when harvest can

interact with migration and both the magnitude and spatial synchrony of environmental

stochasticity. We illustrated these important interactions using this model because of its

simplicity and interpretability relative to more complicated nonlinear numerical models. In

Model 2, we analyzed Pacific herring data from British Columbia using a Bayesian hierarchical

model to estimate spatiotemporal variation in spawning biomass and harvest rates. In **Model 3,**

to evaluate potential solutions for spatial mismatches and conditions for their success, we

developed and applied a spatially explicit stochastic numerical model of metapopulations and

their interaction with the fishery. Together, these approaches (Models 1-3) generate theoretical,

empirical, and numerical results for understanding the complex, spatially-structured interactions

among populations, harvest, and environmental variability. The collective results are then

applied to evaluate the consequences for the availability of important natural resources for both

human and non-human user groups.

Theoretical effects of harvest on spatial variability in metapopulations (Model 1)

We used a simple metapopulation model to evaluate how harvest alters spatial variability,

conditional on the underlying properties of the metapopulation. For our purposes, spatial

variability is the degree to which volatility of the populations are masked by the observed

volatility of the metapopulation.

We considered a simple metapopulation consisting of two populations, linked through the annual

fraction migrating between populations (δ). We assumed both populations have the same

dynamics (identical density independent adult total mortality rate (Z), maturity at age 2,

Gompertz stock-recruit relationship, and symmetric adult annual migration) and trends in

recruitment variability are controlled by a spatially correlated lognormal environmental

stochasticity.

Let represent the abundance of age class a at location i at time t as $Y_{a,i,t}$. Adult dynamics are

shaped by mortality (Z - the sum of mortality from natural causes and harvest) and migration

($0 \leq \delta \leq 0.5$):

Eq. 1: $Y_{a,i,t+1} = \begin{cases} e^{-Z}\left[(1-\delta)Y_{a-1,i,t} + \delta Y_{a-1,j,t}\right] & 2 \leq a \leq n-1 \\ e^{-Z}\left[(1-\delta)(Y_{a-1,i,t} + Y_{a,i,t}) + \delta(Y_{a-1,j,t} + Y_{a,j,t})\right] & a = n \text{ (plus group)} \end{cases}$

Total reproduction is the sum of adults across all adult age classes multiplied by 0.5 to account

for an equal sex ratio:

Eq. 2: $Y_{a=0,i,t+1} = 0.5e^{-z}\sum_{a=1}^{n}\left((1-\delta)Y_{a,i,t} + \delta Y_{a,j,t}\right)$

Both Eq. 1 and 2 are identical for both subpopulations (i.e. the system is symmetrical). We used

a stochastic Gompertz model as the compensatory function that determines how zygotes translate

into one-year-olds:

Eq. 3: $Y_{a=1,i,t+1} = \alpha(Y_{a=0,i,t})^{1-\beta}e^{\zeta_{i,t}}$

α, β and $\zeta_{i,t}$ represent, respectively, the density independent productivity parameter, the within-

location compensatory parameter, and environmental stochasticity that operates on post-

dispersing larvae. The vector $\boldsymbol{\xi}_t = [\zeta_{1,t}, \zeta_{2,t}]'$ follows a multivariate normal distribution with

mean zero and covariance controlled by the common within-site variance (σ_R^2) and spatial

correlation (ρ_R) yielding $\text{cov}[\zeta_{1,t}, \zeta_{2,t}] = \begin{bmatrix} \sigma_R^2 & \sigma_R^2\rho_R \\ \sigma_R^2\rho_R & \sigma_R^2 \end{bmatrix}$.

To analyze the stochastic metapopulation model, we converted the model to a first order vector

autoregressive model where statistical properties of stochastic forcing in multivariate systems are

well described (Lütkepohl 2005). We first vectorized the model by aligning variables from both

populations in a single vector:

$\mathbf{Y}_t = [Y_{a=0,L=1,t} \quad \cdots \quad Y_{n,L=1,t} \quad Y_{a=0,L=2,t} \quad \cdots \quad Y_{n,L=2,t}]'$

We then cast the model in terms of log-scale deviations from the equilibrium (i.e. $x_{a,i,t} =$

$\ln Y_{a,i,t} - \ln Y_{a,i}^*$ where $Y_{a,i}^*$ is the equilibrium) and linearized about the equilibrium (Nisbet and

Gurney 1982, Bjørnstad et al. 2004) to create a first order vector autoregressive model. This

approach approximates the deterministic dynamics of the nonlinear model with a Jacobian matrix

(**J**) of first order dependencies and an environmental covariance matrix.

The matrix of Jacobian coefficients (with plus groups summed at age n) are partial derivatives

for each age class within each subpopulation with respect to each other age class in each

subpopulation. $\mathbf{J}$ can be represented by the block matrix comprised of matrices describing

among and within population age transitions:

Eq. 4: $\mathbf{J} = \begin{bmatrix} J_1 & J_2 \\ J_2 & J_1 \end{bmatrix}$

Where J_1 is within population dynamics and J_2 is among population dynamics defined by:

Eq. 5: $J_i =$

162
$$
\begin{bmatrix}
0 & \frac{D}{1+\sum_{a=1}^{n-1}(e^{-Z})^a} & \frac{D(e^{-Z})}{1+\sum_{a=1}^{n-1}(e^{-Z})^a} & \frac{D(e^{-Z})^2}{1+\sum_{a=1}^{n-1}(e^{-Z})^a} & \cdots & \frac{D(e^{-Z})^{n-2}}{1+\sum_{a=1}^{n-1}(e^{-Z})^a} & \frac{D(e^{-Z})^{n-1}}{1+\sum_{a=1}^{n-1}(e^{-Z})^a} \\
1-\beta & 0 & 0 & 0 & \cdots & 0 & 0 \\
0 & D & 0 & 0 & \cdots & 0 & 0 \\
0 & 0 & D & 0 & \cdots & 0 & 0 \\
\vdots & \vdots & \vdots & \vdots & \ddots & 0 & 0 \\
0 & 0 & 0 & 0 & \cdots & \frac{D}{(e^{-Z})+1} & \frac{D(e^{-Z})}{(e^{-Z})+1}
\end{bmatrix}
$$

If $J_i = J_1,\ D = (1-\delta)$ and if $J_i = J_2,\ D = \delta$. Z is the annual total mortality rate and n is the

number of age classes. See Appendix S1 for full derivation of the Jacobian and VAR(1)

properties. The resulting VAR(1) model is:

Eq. 6: $\mathbf{x}_t = \mathbf{J}\mathbf{x}_{t-1} + \mathbf{Z}\boldsymbol{\xi}_t$

Eq. 7: $\mathbf{x}_t = [x_{a=0,L=1,t} \quad \cdots \quad x_{n,1,t} \quad x_{a=0,L=2,t} \quad \cdots \quad x_{n,2,t}]'$

$\mathbf{Z}$ is a 2n x 2 binary matrix (2 age classes for each location, two locations) that controls which of

the entries in $\mathbf{x}_t$ are subject to stochasticity from $\boldsymbol{\xi}_t$ (i.e. translating the 2x1 vector of location

specific stochasticity to a 2n x 1 sparse vector). In this case, only age class 1 (corresponding to

the second column of J_1) for each subpopulation is subject to stochasticity.

We used the known statistical properties of a VAR(1) model (Lütkepohl 2005) in tandem with

the moments of a multivariate lognormal to derive the coefficient of variation for the

subpopulations and metapopulation that yields the spatial variation in the metapopulation.

Finally, to evaluate how local environmental sensitivity of population growth changes with

mortality rate, we used a first-order impulse response analysis (Lütkepohl 2005) to estimate the

annual intrinsic growth rate response to a recruitment pulse. The impulse response in our case

describes how a recruitment perturbation affects the intrinsic growth rate of each subpopulation.

Specifically, it is given by the entry of the 2^{nd} column of the 1^{st} row of the Jacobian matrix in Eq.

5. See Appendix S1 for full derivations and details.

Pacific herring case-study: spatial variability in biomass and catch (Model 2)

To estimate patterns of spatial and temporal variability in herring biomass and harvest rates, we

used spawn deposition and harvest data for six major Pacific Herring spawning units that

comprise the Central Coast stock in British Columbia, Canada. Briefly, SCUBA surveys are

used to estimate herring egg abundance which is converted to total spawning biomass with the

assumed conversion of ~100 eggs/g and an equal sex ratio (DFO 2015). For consistency we use

the same index as in the DFO stock assessment but in the spatially disaggregated form. Catch

(which occurs in the days prior to spawning) is reported by geographic section that are delineated

and aggregated geographically. For full description of the time series see DFO (2015) and

references therein.

To estimate spatial biomass trends from the survey and harvest time series observations, we used

a nonlinear model in a Bayesian hierarchical state-space framework. We assumed survival,

reproduction, and competition were location specific. We modeled expected change in biomass

in year t+1 using a combined growth and survival model (individual growth and survival from

year t) and Gompertz recruitment model (from year t-1, because fish mature at or after age 2 -

Table 1, Eq. 8).

We estimated both process error and observation error in the model.   We estimated spatially

correlated process error that represents deviations from the expected log-scale pre-harvest

biomass within each location. Process error may arise from a diverse combination of factors

including immigration or emigration and temporal variation in growth, recruitment, or mortality

(Table 1, Eq. 9). We also estimated location-specific observation error (Objective function -

Table 1, Eq. 10), and a common survey bias (a "catchability" coefficient quantifying the mean

proportion of eggs that are observed in fishery-independent surveys - Table 1, Eq. 10). Thus, we

estimated trends conditional on estimated survival, biomass growth, process error covariance,

observation error, and survey bias.

Because we estimated log-scale pre-harvest biomass but observations are of post-harvest

biomass and harvested biomass, the model requires a change of variables and thus the posterior

requires a Jacobian adjustment of the inverse transform (Gelman et al. 2013, Carpenter et al.

2016, Table 1, Eqs 10, 11). See Table 1 for all model equations, parameter definitions, and prior

distributions. For details of estimation, posteriors, and model validation, see Appendix S2. We

estimated the models using Stan (Stan Development Team 2016b, a) with 3 independent Markov

Chains with 1000 iteration chains after 1000 iteration burn-in. We confirmed chain mixing and

convergence using Gelman-Rubin statistic ($R<1.01$, Gelman and Rubin 1992) and for mdel

adequacy and model fit using posterior predictive checks (see below). Posteriors compared with

priors for core parameters are shown in Appendix S2: Fig. S1.

We used the full model posterior of post-harvest biomass to calculate the following metrics: 1)

temporal and spatial variation in biomass (see Appendix S2: Fig. S1), and 2) the exploitation

rate. We compared the estimated local harvest rates from the model posterior to the theoretical

proportional allocation where fishing mortality is constant in space and an optimized allocation

of harvest given the quota (see Appendix S2: Fig. S3 for results, Appendix S3 for methods) and

the posterior mean. Here, we defined "optimized allocation" of catch as one that distributes

catches in space according to the ideal free distribution (see Appendix S3 for methods). This

distribution removes proportionally more biomass from subpopulations with higher biomass and

is the spatial allocation of catch that minimizes effort to achieve the overall quota (assuming

catch per unit effort is linearly related to biomass).

Solutions for spatial scale mismatches in fished herring metapopulations (Model 3)

We used a stochastic model to simulate how spatial metapopulation dynamics and alternative

management scenarios interact to influence risk of collapse at the subpopulation and

metapopulation scales under a diverse suite of scenarios. The scenarios we examined include a

factorial gradient of 1) annual movement (see Annual adult migration among spawning areas

below), 2) environmental recruitment synchrony (the degree of correlation in recruitment in

space – see Spatiotemporal Recruitment Dynamics below), 3) a range of harvest rates (see Stock

Harvest Quota below), and 4) allocation of harvest in space (see Spatial Harvest Prosecution

below). For this analysis, we defined a "collapse" as years with spawning biomass below 20%

of unfished biomass. We used this definition for herring because, 1) it is generally seen as a

conservative estimate of biomass below which traditional Indigenous harvest becomes difficult

and, 2) it lies below the current closure threshold of 25% unfished biomass. Alternative

thresholds defining collapse were also assessed but yielded qualitatively similar results (data not

shown).

For all analyses, we simulated 10 populations placed around a hypothetical circular island of

arbitrary size, where distances among adjacent populations were equal. Fish spawning at any

site in a given year were able to move to any other site in the next year. The probability of

migration from one site to another declines as alongshore distance between the locations

increases and is controlled by a periodic kernel. Likewise, synchrony in stochastic recruitment

among sites decays with distance between locations controlled by a periodic covariance kernel.

Stochasticity in dynamics was included in recruitment (spatial and temporal variation), survival

(temporal variation only) and movement probabilities (spatial and temporal variance). The order

of operations mathematically was 1) recruitment, 2) survival, 3) movement, 4) roe fishery

harvest, and 5) spawning. Details of the simulation are outlined below with core equations

listed in Table 2 and definitions in Table 3.

Spatiotemporal Recruitment Dynamics

Following the assumptions of the current British Columbia herring assessments (DFO 2015),

recruitment of age 2 individuals at location i is a function of eggs produced 2-years prior and

local density dependence via a Beverton-Holt model. We allowed recruitment to exhibit both

spatial and temporal variability and autocorrelation with a first-order vector autoregressive

model. Spatial correlations in recruitment followed a Gaussian spatial decay with distance (d),

and location specific recruitment variability was tuned such that net recruitment variability was

approximately constant across scenarios (CV of metapopulation recruitment = 0.8; DFO 2015).

Annual Adult Survival and Migration Among Spawning Areas

All adult survival occurred prior to movement into spawning locations, was constant in space

and was randomly drawn from a beta-binomial with mean $\lambda = 0.6$ with coefficient of variation of

0.2 (DFO 2015). Survivors migrated to new spawning locations with a probability that decays

with distance from the previous site, controlled by a periodic kernel, tuned to achieve the desired

retention rate.

Stock Harvest Quota

We followed the existing harvest control rules of Pacific herring in British Columbia (DFO

2015). The annual biomass harvest quota for the stock ($\hat{Q}_{t+1}$) was designed to achieve a target

harvest rate (H_{Target}) and a minimum spawning escapement (25% of B_0 – the average steady

state biomass without fishing). For simplicity and to avoid evaluating stock assessment model

performance (which is out of the scope of this study) we assume forecasts are unbiased with no

observation noise.

Spatial Harvest Prosecution

We used and compared two spatial fleet allocation scenarios to generate the distribution of

spatial commercial roe harvest given the stock quota: 1) the fleet prosecuted the fishery equally

in space whereby the absolute harvest was directly proportional to spawning biomass within a

given year (***proportional allocation***) or 2) the fleet was allocated according to the ideal free

distribution (IFD) that in theory would optimize catch efficiency if fleets are not spatially

constrained (***optimized allocation -*** See Appendix S3 for details in generating the IFD). Here,

realized catch was nonlinearly related to spawning biomass, harvesting more from areas with

higher biomass and leaving alone areas with lower biomass. See Appendix S3 for methods and

results from a third allocation strategy, a random spatial allocation that is more similar to the

fleet allocation in the empirical case study. In all cases, fishery selectivity was identical to

maturity-at-age reflecting that harvest occurs on mature fish at the spawning grounds.

Simulation details

For each simulation, we 1) evaluated deterministic equilibria without fishing, 2) initiated

stochastic forcing of recruitment from the equilibrium for 12 years (allowing the full suite of age

classes to be influenced by environmental stochasticity), 3) initiated the fishery in year 13, and 4)

recorded performance metrics for years 23-52. We ran 100 replicate simulations for each

combination of migration probability and recruitment synchrony. Primary performance metrics

summarized for each simulation included a) mean number of years below 20% B_0 at both stock

and substock scales to assess risk of collapse, b) mean stock and substock level catch, c) mean

stock and substock biomass, d) stock and substock temporal variability (coefficient of variation)

in biomass, and e) mean spatial variation in biomass (difference between squared coefficient of

variation at the substock versus stock scale).

Results

Theoretical effects of harvest on spatial variability in metapopulations (Model 1)

Our model illustrates that spatial variation among subpopulations increases with higher harvest

rates (Fig. 1a) and decreases with synchronizing forces of migration and environmental

correlation (difference in surfaces in Fig. 1b). As a result, the metapopulation trend and

coefficient of variation are less reflective of trends and variation in its subpopulations as harvest

increases (Fig. 1b). Biologically, this higher mortality rate decreases longevity and increases

local-scale sensitivity to spatially explicit recruitment pulses. The amplification of fluctuations

occurs at a higher rate at local scales than on aggregate. This response decreases spatial coupling

and predictability. Reduced longevity of adults (via higher mortality) reduces the abundance of

adults in each subpopulation that buffers against local stochasticity through survival and

migration.

Mathematically, this result emerges for several related reasons. Increases in mortality (constant

across space in this case) reduce local subpopulation inertia (predictability, measured as 1[st] order

temporal autocorrelation, thick black line in Fig. 1c). This reduction in inertia increases

subpopulation sensitivity to temporal variation in local subpopulation recruitment (red line in

Fig. 1c). This sensitivity is illustrated by the primary impulse response at a single lag which in

this case always increases with total mortality (Z) derived from Eq. 5:

Eq. 22: $(1 - \delta)/(1 + \sum_{a=1}^{n-1}(e^{-Z})^a)$

Such increases in local environmental sensitivity reduce spatial coupling (measured as the spatial

autocorrelation, dotted black line in Fig. 1c) and thereby increase spatial variability in

abundance.

Importantly these patterns of spatial variability can persist even in the presence of modest

migration rates (Fig. 1b). While spatial variability decreases with synchronizing forces of

migration (δ) and environmental correlations (ρ_R) (Fig. 1b, see Appendix S1 for solutions),

spatial variability only diminishes as migration and environmental correlations are substantially

high (i.e. as migration probabilities approach 0.5 to produce 100% mixing).   This phenomenon

is illustrated by the discrepancies between temporal variability of the metapopulation and

component subpopulations that produce spatial variation (Fig. 1b).

Next we illustrate challenges imposed by spatial variation in exploited metapopulations using an

empirical case study of Pacific herring and present solutions using a numerical management

strategy evaluation.

Pacific herring case-study: spatial variability in biomass and catch (Model 2)

Pacific herring subpopulations on the Central Coast of BC exhibit substantial spatial variability

in subpopulation trends (Fig. 2a, b, c). The estimated biomass of individual ("local")

subpopulations has varied by more than an order of magnitude over the past three decades, and

similar differences in biomass are evident among subpopulations in the same year. As a

consequence, aggregate ("regional") stock biomass at any one point in time is bolstered by a few

subpopulations, while others linger at low levels (Fig. 2b, Appendix S2: Fig. S2). Synchrony in

biomass among subpopulations is low (0.29), with much higher variability at the subpopulations

scale (average CV = 0.88) than in aggregate (CV = 0.54). This discrepancy results from high

spatial variability in subpopulations trends (Fig. 1c - also known as β variability; (Wang and

Loreau 2014). In fact as much as an estimated 91% of an individual subpopulation's biomass is

harvested annually, though the aggregate exploitation rate fluctuates around the target of 20%

(Fig. 2d, e, Appendix S2: Fig. S3). Counterproductively, this can result in occurrences where

subpopulations experiencing periods of lower-than-average biomass are heavily exploited

preceding collapse (e.g. Fig. 2d, in 2006). Harvest rates generally differ in magnitude among

subpopulations in any given year (Appendix S2: Fig. S3) and higher harvest rarely focus on the

subpopulations with highest biomass (Appendix S2: Fig. S3). This is illustrated directly by

spatial harvest distributions that deviate substantially from baselines of spatial harvest evenness

used here and in the simulation model (proportional vs optimized allocation - Appendix S2: Fig.

S3). Next we explore the potential consequences of different spatial harvest distributions via a

closed-loop simulation model.

Solutions for spatial scale mismatches in fished herring metapopulations (Model 3)

We used numerical simulation of a metapopulation to determine whether and under what

conditions exploitation rates that appear sustainable in aggregate can risk collapse of local

subpopulations, and by extension, adversely affect predators and the fishers who target them at

this scale. The divergence in risk among scales, an effect of the mismatch in spatial scale, is

controlled by the magnitude of harvest rates, allocation of harvest in space, annual migration,

and spatial recruitment synchrony. Our simulations show that risk of collapse can be 10 times

greater at local subpopulation scales than at aggregate metapopulation scales (Fig. 3) for the 20%

harvest rule. While it may seem intuitive that relatively modest adult migration would minimize

differences in risk to subpopulations and the metapopulation, our results do not support this

supposition. Even relatively high migration rates can impose substantial discrepancies in the risk

of collapse between subpopulation and metapopulation scales (Fig. 3). This principle holds so

long as spatial synchrony in recruitment is not exceedingly high (Fig. 3 upper portions of

heatmaps). For most scenarios, high spatial variability (i.e. from low annual migration among

subpopulations or low spatial recruitment synchrony) leads to low apparent risk at the aggregate

metapopulation scale, despite high risk of collapse for local subpopulations (Fig. 3; risk increases

towards the lower left quadrants).

We find that risk to local subpopulations greatly exceeds risk to the aggregate metapopulation

with both a simulated 10% and 20% target harvest rate. Under the 20% target harvest, risk of

collapse at the local scale exceeds 10%, even with annual migration among subpopulations

approaching 50% (Fig. 3 h). When harvest is optimized in space or target harvest rates are

reduced to 10%, risks of collapse at local scales are substantially reduced (Fig. 3 d,f,j versus Fig.

3 h) and limited to scenarios where migration is ~10-15% and spatial synchrony in recruitment is

low. Local scale risks of collapse are even worse under a random (or opportunistic) spatial

allocation, but also ameliorated by reductions in harvest (Appendix S3: Fig. S2). Importantly, a

10% target harvest with optimized spatial allocation never exceeded 10% risk of collapse in our

simulations.

The discrepancy in risk of collapse at local subpopulation versus aggregate metapopulation

scales is shaped by both overall harvest rates and spatial variance in subpopulation trend (Fig.

S5). Risk at the subpopulation scale matches that of metapopulation scales when subpopulations

exhibit little spatial variance (i.e. when there is high recruitment synchrony or high migration,

top vs bottom panels in Fig. 3 c-j, SI Fig. S6). In contrast, risk diverges with increases in spatial

variance (Fig. S5).   Spatial variance is shaped not only by spatial recruitment synchrony and

migration, but also harvest rates and allocation in space. Fig. 4b-c illustrates how different

harvest allocations impact spatial variation in trends (Fig. 4 d, e). Higher harvest rates increase

risk at subpopulation scales in part because of higher depletion at the metapopulation scale (Fig.

3 d,f,h,i) and also because harvest can increase spatial variance (Fig. 4d, also explored more

generally in Fig. 1) when efforts are allocated proportionally. This result demonstrates how low

demographic synchrony creates a portfolio effect at the metapopulation scale but simultaneously

masks risks of collapse at the subpopulation scale which can be exacerbated with higher harvest

rates.

However, dynamically optimizing harvest rates in space to match local scale variability lowers

the risk of subpopulation and aggregate metapopulation collapse (Fig. 3 d vs f and Fig. 3 h vs i)

with no cost to aggregate catch (Appendix S3: Fig. S3). These results emerge from simulating a

form of spatially optimized fishing effort that leaves underperforming locations unexploited and

heavily targets overperforming local subpopulations, in accordance with predictions of ideal free

distribution theory. Optimal spatial allocation of harvest rates reduces the effects of harvest

intensity on spatial variability (Fig. 4d) which is strongly related to the bias in estimating risk

among scales (Appendix S3: Fig. S1) and also reduces length of individual collapses (Appendix

S3: Fig. S3). Thus, approaches that optimize spatial harvest to account for subpopulation

dynamics can minimize the spatially destabilizing effect of harvest as well as disparity in

exposure to risk of collapse among spatial scales. The benefits of spatially optimized harvest in

terms of risk to subpopulations diminishes as migration and spatial synchrony in recruitment

productivity decline to low levels (i.e. approach the lower left quadrant of panels in Fig. 3 d,f,h,j,

where subpopulations are nearly autonomous with low connectivity).

Discussion

Harvest strategies that appear appropriately prescribed at large spatial scales can, at local scales,

lead to declines or even effective extirpation of local subpopulations. We call these small-scale

declines "cryptic collapses". Specifically, regional harvest strategies can create a "gilded trap"

(Steneck et al. 2011) where, in this case, management focuses on the aspects of metapopulations

that can benefit conservation and economics at the aggregate scale, but neglect social-ecological

inequity in the exposure to risk at local scales. Our results show that spatial mismatch among

scales is not merely an esoteric concern. Indeed, they occur in current management situations,

and are supported by ecological theory we develop here. Our multiscale risk analysis highlights

the impacts of scale mismatch on consumers and the potential value of optimized spatial

management for sustainability and equity. Previous studies have also investigated spatial

mismatches in fisheries contexts to understand consequences stemming from spatial mismatches

among biological processes and available data, as well as the spatial implementation of fisheries

for yield and measure of population status assessments (Cope and Punt 2011, McGilliard et al.

2011, Spies et al. 2015, McGilliard et al. 2017). Our work expands on previous investigations by

considering how spatial dynamics of fish and fisheries affect resource sustainability at multiple

spatial scales that are relevant to different species and fishing communities. Our models address

this gap by building on existing research with increased biological realism to allow for 1) adult

migration rates among subpopulations rather than only dispersal associated with recruitment (e.g.

Cope and Punt 2011, McGilliard et al. 2011, Spies et al. 2015, McGilliard et al. 2017) ; 2) a

range of complex spatio-temporal correlations in the stochastic populations dynamics (but see

McGilliard et al. 2011 for an implementation of spatio-temporal variation in adult mortality); and

3) a novel suite of fisheries spatial harvest strategies.

Insights into the benefits of population portfolios to manage risk for different groups

Our work adds critical resolution and understanding to the literature on portfolio effects that is

focused on the benefits of spatial variability. Recent work viewing multiple populations as a

"portfolio" of assets has shown benefits of maintaining a diversity of subpopulations with high

asynchrony. These benefits include reducing local extinctions through so called 'rescue effects'

(Hill et al. 2002, Secor et al. 2009, Fox et al. 2017), providing increased stability in the form of

food security for people or animals (Nesbitt and Moore 2016) and minimizing economic risks

over large scales by minimizing variance in harvestable abundance (Schindler et al. 2010). Yet

these portfolio analyses typically focus on the attributes of the aggregate metapopulation,

whereas the risk of localized subpopulation collapse (e.g. depletion below a threshold of

ecological functionality or socioeconomic value) affects the interests of locally constrained

fishers and spatially constrained organisms with small home ranges. Using theory and data we

show that the same spatial variation that leads to resilience at the metapopulation scale, when left

unaccounted for in management strategies, can also produce unforeseen negative consequences

in the form of magnified spatial variation and local risk of collapse (see also Spies et al. 2015).

In the case of Pacific herring in the Central Coast of British Columbia, local reductions in some

subpopulations occurred well before the entire stock showed substantial declines in the mid-

2000s. These collapses had greater impact on spatially constrained groups; namely Indigenous

communities for whom herring is a source of cultural and economic vitality (Brown and Brown

2009, Gavreau et al. 2017). In contrast, mobile fishing fleets and transient predators should be

less vulnerable to local depletion events in the short term. Such context dependent effects of

ignoring spatial variation are exemplified through considering the dramatic differences in the

spatial scale at which predators and fishers operate. Indigenous fishers are spatially constrained

by boat size, fuel costs, and political/cultural boundaries (Fig 2a, Table 4, Harris 2000, von der

Porten et al. 2016). In contrast, the commercial fleet of seine and gillnet fishers are highly mobile

and can pursue fish throughout the region. Similarly, non-human predators of herring and

herring roe have a diversity of home-ranges and therefore interact with herring at multiple scales

(Table 4). Many predators rely on herring when they move inshore around spawning season.

Many crustaceans (Hines 1982, Stone and O'Clair 2001), echinoderms (Mattison et al. 1976,

Cieciel et al. 2009), rockfishes and lingcod (Jorgensen et al. 2006, Mitamura et al. 2009,

Tolimieri et al. 2009, Beaudreau and Essington 2011, Green and Starr 2011, Freiwald 2012),

harbor seals (Peterson et al. 2012, Ward et al. 2012), some seabirds (Peery et al. 2009, Barbaree

et al. 2015, Lorenz et al. 2017), and some flatfishes (Moser et al. 2013), exhibit restricted

patterns of movement and are likely to exploit one to several major subpopulations, but generally

not the entire spatial distribution of the metapopulation (stock). In contrast, humpback whales

(Dalla Rosa et al. 2008, Kennedy et al. 2014), orcas (Hauser et al. 2007, Fearnbach et al. 2014),

some seabirds (Pearce et al. 2005), sea lions (Merrick and Loughlin 1997, Fearnbach et al. 2014,

Kuhn and Costa 2014), fur seals (Kuhn et al. 2014), halibut (Loher 2008, Seitz et al. 2011,

Nielsen et al. 2014), and gadiforms (Wespestad et al. 1983, Hanselman et al. 2014, Rand et al.

2014) can, given ranges reported, access the geographic area covered by the stock (Table 4, see

DataS1:Appendix S4). Thus, herring provide food resources to groups with varying movement

constraints. As a result, herring collapses that range from small-scale subpopulations to

metapopulation-wide phenomena may have differential impacts on the diverse predators that

depend on this resource. These spatial scale dependencies affect which fishing communities or

species bear the brunt of management risks and who reaps the benefits from the portfolio payoff

of regional metapopulation stability. These outcomes highlight that ignoring space can exclude

critical social, ecological, and economic responses central to the triple bottom line (Elkington

1994, Okamoto et al. 2019).

Pacific herring fisheries on Canada's west coast provide an empirical case where the scale of

regional stock assessments masked episodic local overexploitation and subpopulation collapses.

Here, high local harvest rates appear to be commonplace even when local subpopulations are

depleted presumably because of at least two key factors. First, schooling fish are easy to catch

even at low abundances (Mackinson et al. 1997) and thus the quota is likely to be achieved even

if the abundance of spawning fish in a particular location is small. Second, managers are

challenged with fulfilling quotas with imperfect spatial information about spawning abundances.

While high local harvest rates may have contributed to subsequent local collapses observed in

this case study, quantifying other confounding demographics (i.e. spatial variation in stochastic

adult mortality) are necessary to explicitly test this hypothesis. While recent local collapses may

have occurred in the absence of fishing, fishing is very likely imposing higher subpopulation

sensitivity to any environmental or biotic influence on recruitment or survival by reducing adult

longevity thereby eroding an important buffer against recruitment volatility (Essington et al.

2015) that we show can contribute to spatial variation. Importantly, spatial variation may not

only affect the localized groups. In the long term, assumptions of spatial homogeneity can

generate biased estimates of total productivity (Takashina and Mougi 2015) that may lead to

overly optimistic harvest strategies. Thus, high local harvest rates may produce sequential

depletion over time that eventually erodes the principal of the stock with potential to generate

collapse of the portfolio as a whole (Spies et al. 2015).

Linking harvest dynamics to spatial variation in population dynamics

Importantly, spatial variation in population dynamics is not independent of harvest dynamics.

Rather, we demonstrate numerically and analytically that spatial variation is likely to increase

with harvest. This can occur in part because harvesting reduces spatial coupling. Higher

mortality is known to reduce the abundance of older age classes (Barnett et al. 2017) and

increase sensitivity to fluctuations in recruitment (Beddington and May 1977, Bjørnstad et al.

2004, Hsieh et al. 2006, Shelton and Mangel 2011, Okamoto et al. 2016). In spatially structured

systems, such reduced longevity diminishes the synchronizing effect of migration and elevates

sensitivity of subpopulations to localized environmental effects. However, if the distribution of

harvest in space can be optimized to more adequately accommodate the spatial distribution of

fish, we show how the overall portfolio can benefit at multiple scales. Here, demographic

asynchrony (in this case asynchrony in recruitment productivity) can be maintained while

minimizing subpopulation risk. Translated into practice, our simulation and analytical results

highlight two non-mutually exclusive solutions that provide more equitable spreading of risk

among scales. First, reductions in overall harvest rates can ameliorate biases in risk among

scales. This occurs by reducing baseline levels of risk and reducing effects of harvest on spatial

variability. Second, spatially optimizeing harvest allocations can minimize spatial variance and

ease pressure on at-risk subpopulations, thereby reducing risk of local scale depletion without

sacrificing catch. The first solution creates a trade-off between commercial yield and local risk;

the second solution between the costs of management and fleet transportation, and local risk of

subpopulation collapse. Specifically, achieving something similar to the latter (second solution)

in a realistic setting is likely to require either some combination of greater investment in spatial

monitoring, spatial stock assessments (Punt et al. 2018), and in season-management. Thus,

moving in the direction of spatial optimization is likely to require substantial investment in costs

and personnel for research, stock assessment, and management.

Addressing spatial inequity in risk exposure requires confronting these economic and logistical

trade-offs. For species such as Pacific herring that have volatile spatiotemporal dynamics and

complex migratory phenology (Benson et al. 2015), polycentric governance structures where

governing authorities are nested at different spatial scales may help balance these trade-offs by

addressing the problems of fit between ecosystems, social systems, and management agencies

(Young 2002, Berkes 2006, Borgström et al. 2006, Folke et al. 2007, Biggs et al. 2012, von der

Porten et al. 2016). Such systems can capitalize on scale-specific ecological knowledge

(including local, traditional, and scientific knowledge), scientific capacity and socioeconomic

experience to 1) guide decision analyses, 2) co-coordinate data collection and harvest allocation

in space, and 3) test policies (e.g. in the Maine Lobster fishery (Acheson 2003)). Polycentric

management schemes, however, are not a silver bullet. For systems like Pacific herring where

placed-based Indigenous fishing communities often object to purely centralized scales of

management for social, ecological, and economic reasons (Brown and Brown 2009, Thornton

and Kitka 2015, von der Porten et al. 2016, Gavreau et al. 2017, von der Porten et al. 2019),

opportunities to integrate knowledge and objectives into management strategies (and their

evaluation) at smaller spatial scales is well placed in managing these fisheries, and may be

critical to their perpetuity (Okamoto et al. 2019, Salomon et al. 2019).

The models used to generate inference in this study are simple in comparison to the nature of

complex stochastic systems in space. The analytical model (Model 1) makes numerous

simplifications (linearization, biological simplicity) in order to generate analytical and

generalizeable solutions but may ignore more nuanced nonlinearities. The numerical model

(Model 3) on the other hand is more detailed but outcomes are context dependent. Moreover,

both models ignore many biological and management scenarios that are likely to further

complicate spatial patterns (e.g. behaviorally or geographically complex migration (MacCall et

al. 2018, Rogers et al. 2018), spatial and age specific mortality (McGilliard et al. 2011), spatially

complex density dependence (McGilliard et al. 2017), Allee effects, cost, and data limitations for

spatial management). We also ignore many alternative spatial allocation strategies that may be

explicitly designed to maximize long-term yields or minimize spatiotemporal variability in part

because these approaches would require a spatial assessment model, which is out of the scope of

this study. However, the principles from our simulation and analytical models should generalize

regardless of the degree of complexity in the system: a precautionary approach cognizant of

resource users across multiple spatial scales may necessitate the incorporation of some degree of

locally based management to minimize spatial discrepancies in risk exposure. Our results

highlight the need to consider diverse scenarios and incorporate fundamental biological attributes

that may impact resource dynamics and people at different scales. Overall, our models suggest a

mixture of management scales may be key to selecting and coordinating harvest levels in space

to navigate towards sustainable and equitable outcomes (Cope and Punt 2011, Biggs et al. 2012).

Conclusion

Our analyses illustrate the importance of considering spatial dynamics for determining how to

most effectively balance equity in management and conservation strategies aimed at achieving

social, economic, and ecological outcomes (Halpern et al. 2013, Law et al. 2017). These issues

are often ignored by centralized management and conservation initiatives focused on larger

spatial scales. For over half a century, fisheries scientists have debated how best to exploit and

conserve "mixed stocks" that have separate dynamics but are inseparable in space, with the aim

of balancing conservation of weak stocks and maximizing total yields (Ricker 1958). Our work

inverts this focus to metapopulations where subpopulations have inseparable dynamics but are

separated in space. We show how, in these settings, exploitation can magnify local scale

fluctuations and spatial variability. As a result, aggregate metrics poorly represent local scale

dynamics and the risk of collapse at the local scale increases at a greater rate than the aggregate

large scale. This phenomenon creates cryptic collapses with ensuing discrepancies in risk

exposure. Fortunately, the magnitude of these discrepancies can potentially be controlled by

lowering harvest rates or seeking harvest dynamics that are spatially optimized. Overall, these

conclusions are relevant not only to Pacific herring fisheries, but also to the great number of

exploited natural resources that exhibit spatial structure and are valuable to species and people

that operate on multiple scales.

Acknowledgements:

DKO was supported by a Strategic Partnership Grant from the National Sciences and

Engineering Research Council (NSERC) of Canada to AKS and MHL and a first-year assistant

professor (FYAP) award to DKO from the FSU Council on Research and Creativity. We thank

the Gordon and Betty Moore Foundation for their support of AS, JSF, PSL, the Ocean Tipping

Points project and the David and Lucille Packard Foundation for their support of PSL, the Tula

Foundation for their support of MHL, and the Ocean Modeling Forum. We thank J. Cleary, S.

Harper, M. Reid, K. Gladstone, B. Gladstone, K. Brown, and D. Neasloss for discussions that

initiated and refined the research, A. Frid and A. Rassweiler for input on the manuscript, and S.

Cox, L. Dee, L. Hauser, B. Hunt, S. Miller, I. McKechnie, S. Pau, T. Pitcher, E. Petrou, T.

Francis, A. Punt and W. Smith for discussions that improved the research. We also thank the

Heiltsuk and Haida First Nations for their partnership in the NSERC Strategic Partnership Grant

and the Department of Fisheries and Oceans for providing herring spatial data.

References

References

Acheson, J. M. 2003. Capturing the commons: Devising institutions to manage the maine lobtser

industry. University Press New England, Lebanon, NH.

Adams, B., and D. D. Kapan. 2009. Man bites mosquito: understanding the contribution of

human movement to vector-borne disease dynamics. PloS one 4:e6763.

Barbaree, B. A., S. Nelson, and B. D. Dugger. 2015. Marine space use by marbled murrelets

Brachyramphus marmoratus at a mainland fjord system in southeast Alaska. Marine

Ornithology 116:173-184.

Barnett, L. A., T. A. Branch, R. A. Ranasinghe, and T. E. Essington. 2017. Old-Growth Fishes

Become Scarce under Fishing. Current Biology 27:2843-2848. e2842.

Beaudreau, A., and T. Essington. 2011. Use of pelagic prey subsidies by demersal predators in

rocky reefs: insight from movement patterns of lingcod. Marine biology 158:471-483.

Beddington, J. R., and R. M. May. 1977. Harvesting natural populations in a randomly

fluctuating environment. Science 197:463-465.

Benson, A. J., S. P. Cox, and J. S. Cleary. 2015. Evaluating the conservation risks of aggregate

harvest management in a spatially-structured herring fishery. Fisheries Research 167:101-

113.

Berkes, F. 2006. From community-based resource management to complex systems: The scale

issue and marine commons. Ecology and Society 11.

Biggs, R., M. Schluter, D. Biggs, E. L. Bohensky, S. BurnSilver, G. Cundill, V. Dakos, T. M.

Daw, L. S. Evans, K. Kotschy, A. M. Leitch, C. Meek, A. Quinlan, C. Raudsepp-Hearne,

M. D. Robards, M. L. Schoon, L. Schultz, and P. C. West. 2012. Toward Principles for

Enhancing the Resilience of Ecosystem Services. Annual Review of Environment and

Resources 37:421-448.

Bjørnstad, O. N., R. M. Nisbet, and J. Fromentin. 2004. Trends and cohort resonant effects in

age-structured populations. Journal of animal ecology 73:1157-1167.

Borgström, S., T. Elmqvist, P. Angelstam, and C. Alfsen-Norodom. 2006. Scale mismatches in

management of urban landscapes. Ecology and Society 11.

Brown, F., and Y. K. Brown. 2009. Staying the course, staying alive - coastal First Nations

fundamental truths: biodiversity, stewardship and sustainability. Biodiversity BC.,

Victora, BC.

Carpenter, B., A. Gelman, M. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. A. Brubaker, J.

Guo, P. Li, and A. Riddell. 2016. Stan: A probabilistic programming language. Journal of

Statistical Software 20.

Chadès, I., T. Martin, S. Nicol, M. Burgman, H. Possingham, and Y. Buckley. 2011. General

rules for managing and surveying networks of pests, diseases, and endangered species.

Proceedings of the National Academy of Sciences 108:8323-8328.

Chesson, P. 1998. Spatial scales in the study of reef fishes: a theoretical perspective. Australian

Journal of Ecology 23:209-215.

Cieciel, K., B. J. Pyper, and G. L. Eckert. 2009. Tag retention and effects of tagging on

movement of the giant red sea cucumber Parastichopus californicus. North American

Journal of Fisheries Management 29:288-294.

Cope, J. M., and A. E. Punt. 2011. Reconciling stock assessment and management scales under

conditions of spatially varying catch histories. Fisheries Research 107:22-38.

Cumming, G., D. H. Cumming, and C. Redman. 2006. Scale mismatches in social-ecological

672   systems: causes, consequences, and solutions. Ecology and Society 11.

673   Dalla Rosa, L., E. Secchi, Y. Maia, A. Zerbini, and M. Heide-Jørgensen. 2008. Movements of

674   satellite-monitored humpback whales on their feeding ground along the Antarctic

675   Peninsula. Polar Biology 31:771-781.

676   Department of Fisheries and Oceans. 2015. Stock Assessment and Management Advice for BC

677   Pacific Herring: 2015 Status and 2016 Forecast. . Can. Sci. Adv. Sec. Res. Doc 2015/038.

678   Elkington, J. 1994. Towards the sustainable corporation: Win-win-win business strategies for

679   sustainable development. California management review 36:90-100.

680   Essington, T. E., P. E. Moriarty, H. E. Froehlich, E. E. Hodgson, L. E. Koehn, K. L. Oken, M. C.

681   Siple, and C. C. Stawitz. 2015. Fishing amplifies forage fish population collapses.

682   Proceedings of the National Academy of Sciences 112:6648-6652.

683   Fearnbach, H., J. W. Durban, D. K. Ellifrit, J. M. Waite, C. O. Matkin, C. R. Lunsford, M. J.

684   Peterson, J. Barlow, and P. R. Wade. 2014. Spatial and social connectivity of fish-eating

685   "Resident" killer whales (*Orcinus orca*) in the northern North Pacific. Marine biology

686   161:459-472.

687   Flostrand, L. A., J. F. Schweigert, K. S. Daniel, and J. S. Cleary. 2009. Measuring and modelling

688   Pacific herring spawning-site fidelity and dispersal using tag-recovery dispersal curves.

689   ICES Journal of Marine Science 66:1754-1761.

690   Folke, C., L. Pritchard, F. Berkes, J. Colding, and U. Svedin. 2007. The problem of fit between

691   ecosystems and institutions: Ten years later. Ecology and Society 12.

692   Fox, J. W., D. Vasseur, M. Cotroneo, L. Guan, and F. Simon. 2017. Population extinctions can

693   increase metapopulation persistence. Nature ecology & evolution 1:1271.

694 Freiwald, J. 2012. Movement of adult temperate reef fishes off the west coast of North America.

695   canadian Journal of Fisheries and aquatic Sciences 69:1362-1374.

696 Gavreau, A., D. Lepofsky, M. Rutherford, and M. Reid. 2017. "Everything revolves around the

697   herring": The Heiltsuk-herring relationship through time. Ecology and Society 22.

698 Gelman, A., and D. B. Rubin. 1992. Inference from iterative simulation using multiple

699   sequences. Statistical science 7:457-472.

700 Gelman, A., H. S. Stern, J. B. Carlin, D. B. Dunson, A. Vehtari, and D. B. Rubin. 2013. Bayesian

701   data analysis. Chapman and Hall/CRC.

702 Green, K. M., and R. M. Starr. 2011. Movements of small adult black rockfish: implications for

703   the design of MPAs. Marine Ecology Progress Series 436:219-230.

704 Halpern, B. S., C. J. Klein, C. J. Brown, M. Beger, H. S. Grantham, S. Mangubhai, M.

705   Ruckelshaus, V. J. Tulloch, M. Watts, C. White, and H. P. Possingham. 2013. Achieving

706   the triple bottom line in the face of inherent trade-offs among social equity, economic

707   return, and conservation. Proceedings of the National Academy of Sciences 110:6229-

708   6234.

709 Hanselman, D. H., J. Heifetz, K. B. Echave, and S. C. Dressel. 2014. Move it or lose it:

710   movement and mortality of sablefish tagged in Alaska. Canadian Journal of Fisheries and

711   Aquatic Sciences 72:238-251.

712 Harris, D. C. 2000. Territoriality, aboriginal rights, and the Heiltsuk spawn-on-kelp fishery. U.

713   Brit. Colum. L. Rev. 34:195.

714 Hauser, D. D., M. G. Logsdon, E. E. Holmes, G. R. VanBlaricom, and R. W. Osborne. 2007.

715   Summer distribution patterns of southern resident killer whales *Orcinus orca*: core areas

716   and spatial segregation of social groups. Marine Ecology Progress Series 351:301-310.

Hill, M. F., A. Hastings, and L. W. Botsford. 2002. The effects of small dispersal rates on

extinction times in structured metapopulation models. The American Naturalist 160:389-

402.

Hines, A. H. 1982. Coexistence in a kelp forest: size, population dynamics, and resource

partitioning in a guild of spider crabs (Brachyura, Majidae). Ecological Monographs

52:179-198.

Hsieh, C., C. S. Reiss, J. R. Hunter, J. R. Beddington, R. M. May, and G. Sugihara. 2006.

Fishing elevates variability in the abundance of exploited species. Nature 443:859-862.

Johnson, T., J. Wilson, C. Cleaver, and R. Vadas. 2012. Social-ecological scale mismatches and

the collapse of the sea urchin fishery in Maine, USA. Ecology and Society 17.

Jones, R., C. Rigg, and E. Pinkerton. 2016. Strategies for assertion of conservation and local

management rights: A Haida Gwaii herring story. Marine Policy.

Jorgensen, S. J., D. M. Kaplan, A. P. Klimley, S. G. Morgan, M. R. O'Farrell, and L. W.

Botsford. 2006. Limited movement in blue rockfish *Sebastes mystinus*: internal structure

of home range. Marine Ecology Progress Series 327:157-170.

Kendall, B. E., O. N. Bjørnstad, J. Bascompte, T. H. Keitt, and W. F. Fagan. 2000. Dispersal,

environmental correlation, and spatial synchrony in population dynamics. The American

Naturalist 155:628-636.

Kennedy, A. S., A. N. Zerbini, B. K. Rone, and P. J. Clapham. 2014. Individual variation in

movements of satellite-tracked humpback whales *Megaptera novaeangliae* in the eastern

Aleutian Islands and Bering Sea. Endangered Species Research 23:187-195.

Kuhn, C., R. Ream, J. Sterling, J. Thomason, and R. Towell. 2014. Spatial segregation and the

influence of habitat on the foraging behavior of northern fur seals (Callorhinus ursinus).

Canadian journal of zoology 92:861-873.

Kuhn, C. E., and D. P. Costa. 2014. Interannual variation in the at-sea behavior of California sea

lions (*Zalophus californianus*). Marine Mammal Science 30:1297-1319.

Lande, R. 1988. Demographic models of the northern spotted owl (Strix occidentalis caurina).

Oecologia 75:601-607.

Law, E. A., N. J. Bennett, C. D. Ives, R. Friedman, K. J. Davis, C. Archibald, and K. A. Wilson.

2017. Equity trade-offs in conservation decision making. Conservation Biology.

Lepofsky, D., and M. Caldwell. 2013. Indigenous marine resource management on the

Northwest Coast of North America. Ecological Processes 2:1-12.

Levin, P. S., T. B. Francis, and N. G. Taylor. 2016. Thirty-two essential questions for

understanding the social–ecological system of forage fish: the case of Pacific Herring.

Ecosystem Health and Sustainability 2.

Levin, S. A. 1992. The problem of pattern and scale in ecology: the Robert H. MacArthur award

lecture. Ecology 73:1943-1967.

Lewandowski, D., D. Kurowicka, and H. Joe. 2009. Generating random correlation matrices

based on vines and extended onion method. Journal of multivariate analysis 100:1989-

2001.

Loher, T. 2008. Homing and summer feeding site fidelity of Pacific halibut (*Hippoglossus*

stenolepis) in the Gulf of Alaska, established using satellite-transmitting archival tags.

Fisheries Research 92:63-69.

Lorenz, T. J., M. G. Raphael, T. D. Bloxton, and P. G. Cunningham. 2017. Low breeding

propensity and wide-ranging movements by marbled murrelets in Washington. Journal of

Wildlife Management 81:306-321.

Lütkepohl, H. 2005. New introduction to multiple time series analysis. Springer, Berlin.

MacCall, A., T. Francis, D. Armitage, J. Cleary, S. Dressel, R. Jones, H. Kitka, L. Lee, J.

McIsaac, P. Levin, D. Okamoto, M. Poe, A. Punt, S. Reifenstuhl, A. Shelton, M. Miple,

J. Silver, J. Schmidt, T. Thornton, R. Voss, and J. Woodruff. 2018. A heuristic model of

learned migration behavior exhibits distinctive spatial and reproductive dynamics. ICES

Journal of Marine Science. 76:598-608.

Mackinson, S., U. R. Sumaila, and T. J. Pitcher. 1997. Bioeconomics and catchability: fish and

fishers behaviour during stock collapse. Fisheries Research 31:11-17.

Mangel, M., and P. S. Levin. 2005. Regime, phase and paradigm shifts: making community

ecology the basic science for fisheries. Philosophical Transactions of the Royal Society

B: Biological Sciences 360:95-105.

Marsh, D. M., and P. C. Trenham. 2001. Metapopulation dynamics and amphibian conservation.

Conservation biology 15:40-49.

Martell, S., J. Schweigert, V. Haist, and J. Cleary. 2012. Moving towards the sustainable

fisheries framework for Pacific herring: data, models, and alternative assumptions; Stock

Assessment and Management Advice for the British Columbia Pacific Herring Stocks:

2011 Assessment and 2012 Forecasts. Can. Sci. Adv. Sec. Res. Doc 136:163.

Mattison, J., J. Trent, A. Shanks, T. Akin, and J. Pearse. 1976. Movement and feeding activity of

red sea urchins (*Strongylocentrotus franciscanus*) adjacent to a kelp forest. Marine

Biology 39:25-30.

McGilliard, C. R., A. E. Punt, and R. Hilborn. 2011. Spatial structure induced by marine reserves

shapes population responses to catastrophes in mathematical models. Ecological

applications 21:1399-1409.

McGilliard, C. R., A. E. Punt, R. Hilborn, and T. Essington. 2017. Modeling the impacts of two

age-related portfolio effects on recruitment variability with and without a marine reserve.

Ecological applications 27:1985-2000.

McKechnie, I., D. Lepofsky, M. L. Moss, V. L. Butler, T. J. Orchard, G. Coupland, F. Foster, M.

Caldwell, and K. Lertzman. 2014. Archaeological data provide alternative hypotheses on

Pacific herring (*Clupea pallasii*) distribution, abundance, and variability. Proceedings of

the National Academy of Sciences 111:E807-E816.

Melbourne, B. A., and P. Chesson. 2006. The scale transition: scaling up population dynamics

with field data. Ecology 87:1478-1488.

Merrick, R. L., and T. R. Loughlin. 1997. Foraging behavior of adult female and young-of-the-

year Steller sea lions in Alaskan waters. Canadian Journal of Zoology 75:776-786.

Mitamura, H., K. Uchida, Y. Miyamoto, N. Arai, T. Kakihara, T. Yokota, J. Okuyama, Y.

Kawabata, and T. Yasuda. 2009. Preliminary study on homing, site fidelity, and diel

movement of black rockfish *Sebastes inermis* measured by acoustic telemetry. Fisheries

Science 75:1133-1140.

Moser, M. L., M. S. Myers, J. E. West, S. M. O'Neill, and B. J. Burke. 2013. English sole

spawning migration and evidence for feeding site fidelity in Puget Sound, USA, with

implications for contaminant exposure. Northwest Science 87:317-325.

Nesbitt, H. K., and J. W. Moore. 2016. Species and population diversity in Pacific salmon

fisheries underpin indigenous food security. Journal of Applied Ecology 53:1489-1499.

Nielsen, J. K., P. N. Hooge, S. J. Taggart, and A. C. Seitz. 2014. Characterizing Pacific halibut

movement and habitat in a Marine Protected Area using net squared displacement

analysis methods. Marine Ecology Progress Series 517:229-250.

Nisbet, R. M., and W. Gurney. 1982. Modelling fluctuating populations. Wiley, New York.

Okamoto, D. K., M. R. Poe, T. B. Francis, A. E. Punt, P. S. Levin, A. O. Shelton, D. R.

Armitage, J. S. Cleary, S. C. Dressell, R. Jones, H. Kitka, L. C. Lee, A. D. MacCall, J. A.

McIsaac, S. Reifenstuhl, J. J. Silver, J. O. Schmidt, T. F. Thornton, R. Voss, and J.

Woodruff. 2019. Attending to spatial social–ecological sensitivities to improve trade-off

analysis in natural resource management. Fish and Fisheries.

Okamoto, D. K., R. J. Schmitt, and S. J. Holbrook. 2016. Stochastic density effects on adult fish

survival and implications for population fluctuations. Ecology letters 19:153-162.

Pearce, J. M., J. A. Reed, and P. L. Flint. 2005. Geographic variation in survival and migratory

tendency among North American Common Mergansers. Journal of Field Ornithology

76:109-118.

Peery, M. Z., S. H. Newman, C. D. Storlazzi, and S. R. Beissinger. 2009. Meeting reproductive

demands in a dynamic upwelling system: foraging strategies of a pursuit-diving seabird,

the marbled murrelet. The Condor 111:120-134.

Peterson, D. A., R. Hilborn, and L. Hauser. 2014. Local adaptation limits lifetime reproductive

success of dispersers in a wild salmon metapopulation. Nature communications 5.

Peterson, S. H., M. M. Lance, S. J. Jeffries, and A. Acevedo-Gutiérrez. 2012. Long distance

movements and disjunct spatial use of harbor seals (*Phoca vitulina*) in the inland waters

of the Pacific Northwest. PloS one 7:e39046.

Punt, A. E., D. K. Okamoto, A. D. MacCall, A. O. Shelton, D. R. Armitage, J. S. Cleary, I. P.

Davies, S. C. Dressel, T. B. Francis, and P. S. Levin. 2018. When are estimates of

spawning stock biomass for small pelagic fishes improved by taking spatial structure into

account? Fisheries research 206:65-78.

Rand, K. M., P. Munro, S. K. Neidetcher, and D. G. Nichol. 2014. Observations of seasonal

movement from a single tag release group of Pacific cod in the eastern Bering Sea.

Marine and Coastal Fisheries 6:287-296.

Ricker, W. 1958. Maximum sustained yields from fluctuating environments and mixed stocks.

Journal of the Fisheries Board of Canada 15:991-1006.

Rieman, B., and J. Dunham. 2000. Metapopulations and salmonids: a synthesis of life history

patterns and empirical observations. Ecology of Freshwater Fish 9:51-64.

Rogers, L. A., A. K. Salomon, B. Connors, and M. Krkošek. 2018. Collapse, tipping points, and

spatial demographic structure arising from the adopted migrant life history. The

American Naturalist 192:49-61.

Salomon, A., A. Quinlan, G. Pang, D. Okamoto, and L. Vazquez-Vera. 2019. Measuring social-

ecological resilience reveals opportunities for transforming environmental governance.

Ecology and Society 24.

Schindler, D. E., R. Hilborn, B. Chasco, C. P. Boatright, T. P. Quinn, L. A. Rogers, and M. S.

Webster. 2010. Population diversity and the portfolio effect in an exploited species.

Nature 465:609-612.

Schtickzelle, N., and T. P. Quinn. 2007. A metapopulation perspective for salmon and other

anadromous fish. Fish and Fisheries 8:297-314.

Secor, D. H., L. A. Kerr, and S. X. Cadrin. 2009. Connectivity effects on productivity, stability,

and persistence in a herring metapopulation model. ICES Journal of Marine Science

66:1726-1732.

Seitz, A. C., T. Loher, B. L. Norcross, and J. L. Nielsen. 2011. Dispersal and behavior of Pacific

halibut *Hippoglossus stenolepis* in the Bering Sea and Aleutian Islands region. Aquatic

Biology 12:225-239.

Shelton, A., J. Samhouri, A. Stier, and P. Levin. 2014. Assessing trade-offs to inform ecosystem-

based fisheries management of forage fish. Scientific Reports 4.

Shelton, A. O., and M. Mangel. 2011. Fluctuations of fish populations and the magnifying

effects of fishing. Proceedings of the National Academy of Sciences 108:7075-7080.

Siple, M. C., and T. B. Francis. 2016. Population diversity in Pacific herring of the Puget Sound,

USA. Oecologia 180:111-125.

Spies, I., P. D. Spencer, and A. E. Punt. 2015. Where do we draw the line? A simulation

approach for evaluating management of marine fish stocks with isolation-by-distance

stock structure. Canadian journal of fisheries and aquatic sciences 72:968-982.

Stan Development Team. 2016a. RStan: the R interface to Stan. Version 2.10.1. <http://mc-

stan.org/rstan.html>

Stan Development Team. 2016b. Stan: A C++ library for probability and sampling. Version

2.12.0. <http://mc-stan.org>

Steneck, R., T. Hughes, J. Cinner, W. Adger, S. Arnold, F. Berkes, S. Boudreau, K. Brown, C.

Folke, and L. Gunderson. 2011. Creation of a gilded trap by the high economic value of

the Maine lobster fishery. Conservation biology 25:904-912.

872 Stephenson, R. L. 1999. Stock complexity in fisheries management: a perspective of emerging

873   issues related to population sub-units. Fisheries Research 43:247-249.

874 Stone, R. P., and C. E. O'Clair. 2001. Seasonal movements and distribution of Dungeness crabs

875   Cancer magister in a glacial southeastern Alaska estuary. Marine Ecology Progress Series

876   214:167-176.

877 Takashina, N., and A. Mougi. 2015. Maximum sustainable yields from a spatially-explicit

878   harvest model. Journal of Theoretical Biology 383:87-92.

879 Tanasichuk, R., A. Kristofferson, and D. Gillman. 1993. Comparison of some life history

880   characteristics of Pacific herring (*Clupea pallasi*) from the Canadian Pacific Ocean and

881   Beaufort Sea. Canadian Journal of Fisheries and Aquatic Sciences 50:964-971.

882 Thornton, T. F., and H. Kitka. 2015. An indigenous model of a contested Pacific Herring fishery

883   in Sitka, Alaska. International Journal of Applied Geospatial Research 6:94-117.

884 Tolimieri, N., K. Andrews, G. Williams, S. Katz, and P. Levin. 2009. Home range size and

885   patterns of space use by lingcod, copper rockfish and quillback rockfish in relation to diel

886   and tidal cycles. Marine Ecology Progress Series 380:229-243.

887 von der Porten, S., J. Corntassel, and D. Mucina. 2019. Indigenous nationhood and herring

888   governance: strategies for the reassertion of Indigenous authority and inter-Indigenous

889   solidarity regarding marine resources. AlterNative: An International Journal of

890   Indigenous Peoples:1177180118823560.

891 von der Porten, S., D. Lepofsky, D. McGregor, and J. Silver. 2016. Recommendations for marine

892   herring policy change in Canada: Aligning with Indigenous legal and inherent rights.

893   Marine Policy 74:68-76.

Wang, S., and M. Loreau. 2014. Ecosystem stability in space: α, β and γ variability. Ecology

letters 17:891-901.

Ward, E. J., P. S. Levin, M. M. Lance, S. J. Jeffries, and A. Acevedo-Gutiérrez. 2012. Integrating

diet and movement data to identify hot spots of predation risk and areas of conservation

concern for endangered species. Conservation Letters 5:37-47.

Wespestad, V. G., K. Thorsen, and S. Mizroch. 1983. Movement of sablefish, *Anoplopoma*

fimbria, in the northeastern Pacific Ocean as determined by tagging experiments (1971-

80). Fishery Bulletin 81:415-420.

Young, O. R. 2002. The institutional dimensions of environmental change: fit, interplay, and

scale. MIT press.

Table 1: Equations used in Model 2 (Pacific herring case-study: spatial variability in biomass and

catch)

Eq	Description	Equation
Eq. 8	Expected pre-spawn biomass	$\hat{b}_{t+1,l}^+ = \exp\left(\alpha_l + (1 - \beta_l)\ln(b_{t-1,l})\right) + \lambda_l b_{t,l}$
Eq. 9	Estimated log-biomass	$\ln \boldsymbol{b}_t^+ = \ln(e^{\ln \boldsymbol{b}_t} + \boldsymbol{h}_t) \sim \text{MVN}(\ln \hat{\boldsymbol{b}}_t^+, \text{diag}(\boldsymbol{\sigma})\Omega)$
Eq. 10	Objective function	$\ell = \sum_{t=1}^{T} \sum_{l=1}^{L} \frac{\left(\ln(\text{obs}[b_{t,l}]) - \left(\ln b_{t,l} + \ln q\right)\right)^2}{(2\sigma_{obs}^2)} - \ln(|d|)$
Eq. 11	Jacobian adjustment for change of variables in Eq. 10	$d = \frac{\partial}{\partial \ln b_{t,l}} \ln(e^{\ln b_{t,l}} + h_{t,l}) = \frac{b_{t,l}}{b_{t,l} + h_{t,l}}$
Symbol	**Description**	**Prior or Data Source**
α_l	Site-specific Gompertz productivity	
β_l	Site-specific Gompertz compensation	$[\alpha_l \quad \text{logit}(\beta_l) \quad \text{logit}(\lambda_l)]' \sim \text{MVN}(\boldsymbol{\Theta}, \boldsymbol{\sigma}_\Theta \Omega_\Theta);$
λ_l	Site-specific aggregate mortality & somatic growth	
Θ	Mean vector for $\alpha, \text{logit}(\beta)$, and $\text{logit}(\lambda)$	$\boldsymbol{\Theta} \sim \text{MVN}(0, \text{diag}(1.5)\Omega_{\bar{\Theta}});$
$\sigma_{\Theta l}$	Among site variances in $\alpha, \text{logit}(\beta)$, and $\text{logit}(\lambda)$	$\sigma_{\Theta l} \sim \text{half-cauchy}(0, 2.5)$
Ω	Process error spatial corr. matrix	
Ω_Θ	parameter corr. matrix	LKJ prior (Lewandowski et al. 2009): scale = 2
$\Omega_{\bar{\Theta}}$	mean parameter corr. matrix	
$\boldsymbol{\sigma}$	Vector of site-specific lognormal process variances	$\sigma_l \sim \text{normal}\left(\sigma_\mu, 0.15\right); \ \sigma_\mu \sim \text{normal}(0, 0.15)$
σ_{obs}	Lognormal obs. error	$\sigma_{obs} \sim \text{half-cauchy}(0, 2.5)$
$\ln q$	Survey bias (egg "catchability")	$\ln q \sim \text{normal}(0, 0.05)$ (Martell et al. 2012)
$\text{obs}[b_{t,l}]$	Input Data	observed spawning biomass at time t at location l
$h_{t,l}$	Input Data	harvest at time t at location l

907   Table 2: Equations used in Model 3 (Solutions for spatial scale mismatches in fished herring

908   metapopulations). See Table 3 for parameter values and definitions.

Eq	Description	Equation
Eq. 12	Recruits (age 2) prior to movement	$n^{++}_{a=2,l,t+2} = \text{eggs}_{l,t} \exp\left[\epsilon_{l,t} - \dfrac{\sigma_r{}^2}{2}\right] / (\alpha + \beta \text{eggs}_{l,t})$
Eq. 13	Total eggs lain	$\text{eggs}_{l,t} = \sum_{a=2}^{10} \text{m}_a \text{f}_a [0.5 n_{a,l,t}]$
Eq. 14	Environmental recruitment stochasticity	$\boldsymbol{\epsilon_t} = \phi \boldsymbol{\epsilon_t} + \text{MVN}(0, \sigma_r{}^2[1 - \phi^2]\boldsymbol{\Omega_r})$
Eq. 15	Spatial correlation in recruitment stochasticity	$\rho_{i,j} = \exp\left(-\dfrac{2\sin^2\pi\left\lvert dist_{i\to j}\right\rvert}{\eta^2}\right)$
Eq. 16	Pre-harvest abundance	$n^{++}_{a,l,t+1} = \begin{cases} \lambda_t\, n_{a-1,l,t} & 2 < a < 10 \\ \lambda_t\, n_{a-1,l,t} + \lambda_t\, n_{a,l,t} & a = 10 \text{ (plus group)} \end{cases}$
Eq. 17	Post-migration abundance	$\boldsymbol{N^+_t} = \boldsymbol{N^{++}_t S};\ \ S_{i,j} = \text{P}(i \to j)$
Eq. 18	Recruit or Adult migration by distance function	$\text{P}(i \to j) = \exp\left(-\dfrac{2\sin^2\pi\left\lvert dist_{i\to j}\right\rvert}{\gamma^2}\right) \sum_{j=1}^{N} \exp\left(-\dfrac{2\sin^2\pi\left\lvert dist_{i\to j}\right\rvert}{\gamma^2}\right)^{-1}$
Eq. 19	Quota	$\widehat{Q}_{t+1} = \begin{cases} min\left(H_{Target} * \widehat{B}_{forecast,t},\ \widehat{B}_{forecast,t} - L_{crit} * \widehat{B}_{0,t}\right) & \text{if } \widehat{B}_{forecast,t} \geq L_{crit} * \widehat{B}_{0,t} \\ 0 & \text{otherwise} \end{cases}$
Eq. 20	Post-harvest abundance at age in year t	$n_{a,l,t} = (1 - \text{m}_a)n^+_{a,l,t} - \text{m}_a n^+_{a,l,t} e^{-F_{t,l}};\ \ F_{t,l} = \ln h_{t,l} / \sum_{a=2}^{10} \text{w}_a \text{m}_a\, n^+_{a,l,t}$
Eq. 21	Biomass at location l in year t	$B_{l,t} = \sum_{a=2}^{10} \text{w}_a \text{m}_a\, n^+_{a,l,t}$

911   Table 3: List of key parameters and state-variables in Model 3 with definitions, values and

912   citations. Note that equilibrium considerations are generated by simulating with no harvest and

913   no stochasticity ($\sigma_r = 0$, cv in $\lambda_t = 0$, and harvest = 0). Selectivity-at-age in all cases is equal to

914   maturity-at-age.

Parameter	Description	Value(s)	Source
α, β	Beverton-Holt stock-recruit parameters	583.43, 2.089 x 10^{-8}	1
ϕ	1st order partial autocorrelation	0.6	Chosen
m_a	Maturity-at-age (=Selectivity-at-age)	Age 1 = 0, Age 2= 0.2, Age 3=0.9, Age 4 + = 1	2
w_a	Weight-at-age	4.5x10^{-6} [27(1-Exp(-0.48*age))]$^{3.127}$	3
f_a	Fecundity-at-age	$\exp(4.69)(1000w_a)^{1.13}$	3
λ_t	Annual pre-harvest survival rate	Beta-binomial: mean = 0.6, cv= 0.2	Derived from 1
σ_r	Stock-scale standard deviation of rec-devs	0.8	1
η	Scale parameter regulating distance decay in rec-dev spatial covariance	Numerically tuned to achieve desired spatial correlations	Chosen
L_{crit}	Lower biomass threshold for fishing closure relative to $\hat{B}_0$	0.25	1
$\overline{P}(i \rightarrow j)$	Mean migration probability	0.01 to 0.8	Chosen
$\bar{\rho}_{i,j}$	Spatial synchrony in recruitment	0.15 to 0.85	Chosen
State Variable	**Description**		
$n^{++}_{a,l,t}$	Abundance at age a in time t at location l prior to movement and harvest (Eq. 16)		
$n^{+}_{a,l,t}$	Abundance after movement and prior to harvest (Eq. 17)		
$F_{t,l}$	Fishing mortality of susceptible fish at time t at location l after movement and natural mortality (Eq. 20)		

915   1: (Department of Fisheries and Oceans 2015)

916   2: (Martell et al. 2011)

917   3: (Tanasichuk et al. 1993)

918   Table 4: Home-range categories of different Pacific herring predator groups. Details of sources

919   and home-ranges are provided in the DataS1:Appendix S4.

Home-Range Category	Home-Range Description	Herring Predator Groups
Localized	single subpopulation *(< 10 km radius)*	Indigenous fishers**, Seabirds*, Flatfishes*, Crustaceans*, Urchins, Cucumbers, Reef Fishes, Rockfishes
Centralized	multiple subpopulations *(~10-40 km-radius)*	Indigenous fishers**, Seabirds*, Flatfishes*, Crustaceans*, Gadiforms*, Salmonids**, Halibut**, Harbor Seals
Transient	full stock or greater	Seine & Gillnet Fishers, Cetaceans, Sea Lions, Seabirds*, Gadiforms*, Salmonids**, Halibut**

*Varies by species, ** Varies by migration phase, location or individual

List of Figures:

Figure 1: Effect of elevating harvest mortality rate on asynchrony related spatial variability in the

metapopulation from Model 1. **(a)** Spatial variation increases as a function of total mortality for

different levels of spatial synchrony in recruitment productivity. Here migration is held at 0.25

and log-scale recruitment variability (σ_R) is 0.7 (i.e. CV = 0.8). Note spatial variation is a result

of the discrepancy in population versus metapopulation temporal variation in panel

(b); **(b)** Temporal variability (squared coefficient of variation) of the population and

metapopulation. The arrows illustrate spatial variability as the difference between

surfaces. **(c)** Measures of local population predictability (measured by first order within

population autocorrelation, thick black line), among population coupling (measured by among

population correlations, thin black line), and local environmental sensitivity (measured by the

response at the population level to a unit environmental impulse affecting recruitment

productivity, red line).

Figure 2: **(a)** Map of major Pacific Herring spawning substocks ("Sections" - denoted by

different symbols) as defined by Canada's Federal Fisheries Agency in the Central Coast of

British Columbia. Indigenous communities noted with *. **(b)** Trends for the six main

supopulations (substocks - thin colored lines & points) and the metapopulation (stock) mean

(thick blue line). Trends were estimated using multivariate hierarchical Bayesian state-space

model integrating spawn surveys and catch information (model 2). Individual plots for each

substock are shown, along with catch, spawn observations, and 95% posterior credible intervals

in Appendix S2. **(c)** Temporal coefficients of variation estimated from the Bayesian state-space

mode over 25 years for each substock, the mean of each substock CV (large black symbol), the

regional stock coefficient of variation (blue point), and the spatial variability (black point with

error bar) which is the standardized variance of the system after accounting for the variance of

the stock and the variance of the substock means. Error bars are 95% credibility intervals. (d)

Trends in annual harvest rate for 2 of the 6 local substocks (Higgins and Lower Spiller) with

points, 95% credibility intervals and the mean for the aggregate stock (thick blue line). For full

results see Appendix S2. The dotted line represents the target harvest rate of 20%.

Figure 3: Mean risk of collapse (%) at metapopulation and subpopulation scales for different

harvest rates and spatial harvest allocation strategies under varying spatial scenarios. Results are

derived from simulation-based risk analysis of hypothetical Pacific herring fisheries. Axes

present simulations with different levels of annual migration and spatial synchrony in

recruitment productivity, columns represent different harvest strategies and rows represent the

spatial scales of inference. Risk of collapse is the probability of falling below 20% of unfished

equilibrium (i.e. the dotted lines in Fig. 4 a-c); at the subpopulation scale, risk is measured as the

mean probability of collapse for each population. *Optimized allocation* harvests proportionally

more from subpopulations with higher biomass; *Proportional allocation* removes the same

proportion of biomass from each subpopulation. Annual migration is the mean percent of each

subpopulation that emigrates each year and spatial synchrony is the mean pairwise correlation in

recruitment productivity. For full results see Appendix S3: Fig. S2. Note that the allocation

strategies produce equivalent average yields (Figure 3c vs. 3e and 3g vs. 3i; see Appendix S3:

Fig. S3 for yield comparisons). For an illustration of how alternative strategies affect the mean

duration of collapses, see Appendix S3: Fig. S4.

Figure 4: **(a-c)** Example simulations (a single simulation run) showing spatial variation in

biomass measured at the subpopulation level (colored lines) or metapopulation mean (thick blue)

for **(a)** an unexploited metapopulation, **(b)** a 20% target harvest allocated proportional to

spawning biomass in space, or **(c)** a 20% target harvest allocated optimally in space according to

the ideal free distribution (see Appendix S3: Fig. S4 for summaries across runs). B_0 =

deterministic biomass in the absence of fishing, dotted line = lower conservation threshold of

20% of B_0. **(d)** change in spatial variation for different harvest rates and spatial harvest allocation

across different spatial parameter scenarios for migration rates and spatial synchrony. Each point

represents the spatial variance from 100 runs from a single 52-year simulation using a unique

migration-synchrony parameter combination.

Figure 1

Figure 2

Figure 3

Figure 4