Spatial variation in exploited metapopulations obscures risk of collapse

Daniel K Okamoto[a,b,c], Margot Hessing-Lewis[b], Jameal F Samhouri[d],
Andrew O Shelton[d], Adrian Stier[e], Philip S Levin[f,g], and Anne K Salomon[b,c]

[a]Department of Biological Science, Florida State University, Tallahassee, Florida 32303, USA
[b]Hakai Institute, PO Box 309, Heriot Bay, BC V0P 1H0, Canada
[c]School of Resource and Environmental Management, Simon Fraser University, 8888 University Drive, Burnaby, B.C. V5A 1S6, Canada
[d]Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Blvd. East,Seattle, WA 98112, USA
[e]Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, CA 93106, USA
[f]The Nature Conservancy, 74 Wall St, Seattle, WA98121, USA
[g]School of Environment and Forestry Sciences, University of Washington, Seattle, Washington, USA

Corresponding Author
Daniel K Okamoto
Department of Biological Science
The Florida State University
Tallahassee, Florida 32303
Phone: (850) 645-9532
emaildokamoto@bio.fsu.edu

Author contributions
DKO designed, built and conducted analyses. AS conducted the home-range literature review. DKO and AKS wrote the initial manuscript draft. DKO, MHL, JFS, AOS, AS, PSL, and AKS initiated the research focus, refined analyses and contributed to revisions

1 **Abstract**

Unanticipated declines among exploited species frequently occur despite harvests that appear
sustainable. This is common in the oceans, where spatial scales of management can be
mismatched with spatially dynamic metapopulations. We explore causes, consequences and
potential solutions for spatial mismatches in metapopulations of an important forage fish by
combining empirical analysis, numerical modeling and novel theory. We show that spatial
complexity can promote stability at large scales, however ignoring complexity produces cryptic
and negative consequences at small scales. Harvesting metapopulations magnifies spatial
variability, which creates discrepancies between regional and local trends while increasing risk
of local population collapses. Such effects asymmetrically impact locally constrained fishers and
predators, which are more exposed to risks of cryptic collapses. Importantly, we show that
dynamically optimizing harvest can minimize local risk without sacrificing yield. Thus, multiple
nested scales of management can help avoid cryptic collapses and the ensuing ecological, social
and economic consequences.

Keywords: Metapopulation, Fisheries, Population Dynamics, Conservation, Natural Resource
Management

Introduction

Mismatches in spatial scale create pervasive problems in ecology and natural resource
management (1). This occurs in part because the spatial extent of management or conservation
units are often defined by history, jurisdictional, or institutional criteria rather than the scale of
social and ecological processes at play (1-6). Such choices about the scale of management can
result in mismatches, where feedbacks controlling interactions among groups occur at different
spatial scales. In managed ecosystems like forestry and fisheries, mismatches may occur when
harvest recommendations are based on trends in large-scale abundance without accounting for
localized collapses (7). Yet these spatially isolated collapses can have far-reaching consequences
when that species plays an indispensable role in local social-ecological systems, including
human communities with limited capacity to forage over wide geographic scales. Empirical
identification of appropriate spatial scales of management remains difficult for spatially
structured populations, but can be a pre-requisite for diagnosing and reconciling challenges that
spatial mismatches impose on the sustainable and equitable use of natural resources.

In metapopulations - plant or animal populations connected through dispersal - the dynamics
of individual populations can differ substantially from the aggregate metapopulation (3, 4, 8-11).
A combination of movement, shared climate drivers, and compensatory processes determine
whether dynamics of individual populations reflect the dynamics of the aggregate
metapopulation (12). Consideration of metapopulation structure has improved the management
of the spotted owl (*Strix occidentalis*) (13), salmon (*Onchorhyncus* spp.) (14-16), amphibians
(17), and mosquitoes (18). To date, efforts to integrate metapopulation dynamics into natural
resource management have largely focused on either minimizing extirpations (19), characterizing
productivity of the metapopulation as a whole (sensu 20), or valuing benefits of portfolio effects
(i.e. stabilizing effects of spatial asynchrony sensu 21). Less understood, both in theory and in
practice, is how mismatches in spatial scale, governed by the interaction between resource
exploitation and metapopulation properties, affect spatial variability, frequency of localized
collapses, and their ecological and social consequences across spatial scales.

In this study, we assess how spatial differences in harvest rates interact with animal
movement and recruitment to shape patterns of population variability and risk of collapse at
different spatial scales. Our results illustrate the challenge in managing spatially complex
populations using three complementary approaches. Our empirical case study of a managed
stock of Pacific herring (*Clupea pallasii*) in Canada's Pacific coast highlights the discrepancies
among scales in both population dynamics and harvest rates. Further, for this system, we use
multiscale risk analysis using closed-loop simulation modeling to highlight the impacts of scale
mismatch. In doing so we demonstrate the potential value of optimized spatial management in
reduction of risk and inequity. We contextualize such spatial risks by conducting a review of
herring predator home-range size. Finally, we generalize these results by developing new theory

to evaluate how exploitation of metapopulations can alter spatial and temporal variability at

different spatial scales and thus affect spatial inequities in risk exposure.

Empirical Pacific herring case study in British Columbia's Central Coast

Pacific herring exemplify the challenges inherent to managing metapopulations. In British

Columbia (BC), Canada (Fig 1a), herring return annually to nearshore coastlines in the late

winter/early spring to reproduce. During this annual migration, they are harvested and preyed

upon by a range of consumers. Mobile commercial fishing fleets harvest adult herring for their

roe in the days prior to spawning; in contrast, indigenous fishers, constrained to a local area,

harvest both adult herring and their eggs after spawning events as an important food, trade and

cultural resource (22-26). Unfortunately, a core uncertainty for herring management, as for many

species, is the extent of movement between areas (27-32)**.** Similar uncertainty surrounds spatial

variation in spawning biomass (i.e. 33), which may result in part from the degree of demographic

synchrony between areas (e.g.. synchrony in recruitment). Pacific herring in British Columbia

are currently managed as stocks at regional (100s of kilometers) scales by Canada's federal

fisheries agency. Within these stocks, multiple spawning aggregations (substocks) seasonally

occupy individual stretches of coastline, many of which are of important traditional and cultural

value to indigenous groups. Thus, Pacific herring fisheries present a valuable system in which to

explore how spatial population dynamics, scales of management, and spatial constraints of

fishers and predators interact to influence differential risk exposure to population collapses.

Results

Empirical Pacific Herring Case Study: British Columbia's Central Coast

Spatiotemporal variability in Pacific herring biomass and catch

Pacific herring subpopulations on the Central Coast of BC exhibit substantial temporal and

spatial variability (Fig 1a, b, c). The estimated biomass of individual ("local") subpopulations

has varied by more than an order of magnitude over the past three decades, and similar

differences in population size are evident among subpopulations in the same year. As a

consequence, aggregate ("regional") stock biomass at any one point in time is bolstered by a few

subpopulations, while others linger at low levels (Fig1b, SI Fig S.1). Synchrony in biomass

among subpopulations is low (0.29), with much higher variability at the subpopulations scale

(avg. CV = 0.88) than in aggregate (CV = 0.54). This discrepancy results from high spatial

variability in subpopulations trends [Fig 1c - also known as β variability (34)].

This discrepancy between variability at the metapopulation (stock) and local subpopulation

scales challenges the management objective of maintaining a target harvest rate of 20%. In fact

as much as an estimated 91% of an individual subpopulation's biomass is harvested annually,

though the aggregate exploitation rate fluctuates around the target of 20% (Fig 1d, e, SI Fig S.2)**.**

Counterproductively, this can result in occurrences where subpopulations experiencing periods

of lower-than-average biomass are heavily exploited preceding collapse (e.g. Fig 1d, in 2006).
Harvest rates generally differ in magnitude among subpopulations in any given year (SI Fig S.2)
and higher harvest rarely focus on the subpopulations with highest biomass (SI Fig S.2). This is
illustrated directly by uneven spatial harvest distributions (SI Fig S.2), which we measured as the
deviations from two baselines of spatial harvest evenness used here and in the simulation model.
These are: proportional allocation[*] or an optimized allocation[†]. We explore the potential
consequences of different spatial harvest distributions via a closed-loop simulation model (see
Spatial risk analysis section below).

Foraging-ranges of select Pacific herring predators

Consumers that depend on Pacific herring for food vary dramatically in the spatial scales at
which they operate. Indigenous fishers are spatially constrained by boat size, fuel costs and
political/cultural boundaries (Fig 1a, Table 1, 26, 35). In contrast, the commercial fleet of seine
and gillnet fishers are highly mobile and can pursue fish throughout the region. Similarly, non-
human predators of herring and herring roe have a diversity of home-ranges and therefore
interact with herring at multiple scales (Table 1). Many predators rely on herring when they
move inshore around spawning season. Many crustaceans (36, 37), echinoderms (38, 39),
rockfishes and lingcod (40-45), harbor seals (46, 47) and some seabirds (48-50) and flatfishes
(51), exhibit restricted patterns of movement and are likely to exploit one to several major
subpopulations, but generally not the entire spatial distribution of the metapopulation (stock). In
contrast, humpback whales (52, 53), orcas (54, 55), some seabirds (56), sea lions (54, 57, 58), fur
seals (59), halibut (60-62) and gadiforms (63-65) can, given ranges reported, access the
geographic area covered by the stock (Table 1, see SI Appendix D). Thus, herring provide food
resources to groups with varying movement constraints. As a result, herring population collapses
that range from small-scale subpopulations to metapopulation-wide phenomena may have
differential impacts on the diverse predators that depend on this resource.

Spatial risk analysis with simulation models of Pacific herring fisheries

116   We used numerical simulation of a metapopulation to determine whether and under what
117   conditions exploitation rates that appear sustainable in aggregate can risk collapse of local
118   subpopulations, and by extension, adversely affect predators and the fishers who target them at
119   this scale. The divergence in risk among scales, an effect of the mismatch in spatial scale, is

---

[*]**proportional allocation**: proportional harvest rates are equal among substocks in a given year.
[†] **optimized allocation**: proportional harvest rates are higher on substocks with more biomass via the ideal free distribution.

controlled by the magnitude of harvest rates, allocation of harvest in space, annual migration,
and spatial recruitment synchrony. Our simulations show that risk of collapse[‡] can be 10 times
greater at local subpopulation scales than at aggregate metapopulation scales (Fig 2) for the 20%
harvest rule. While it may seem intuitive that relatively modest adult migration would minimize
differences in risk to subpopulations and the metapopulation, our results do not support this
supposition. Even relatively high migration rates can impose substantial discrepancies in the risk
of collapse between subpopulation and metapopulation scales (Fig 2). This principle holds so
long as spatial synchrony in recruitment is not exceedingly high (Fig 2 upper portions of
heatmaps). For most scenarios, high spatial variability (i.e. from low annual migration among
subpopulations or low spatial recruitment synchrony) leads to low apparent risk at the aggregate
metapopulation scale, despite high risk of collapse for local subpopulations (Fig 2; risk increases
towards the lower left quadrants).

We find that risk to local subpopulations greatly exceeds risk to the aggregate
metapopulation with both a simulated 10% and 20% target harvest rate. Under the 20% target
harvest risk of collapse at the local scale exceeds 10% even with annual migration among
subpopulations approaching 50% (Fig 2 h). When harvest is optimized in space or target harvest
rates are reduced to 10%, risks of collapse at local scales are dramatically reduced (Fig 2 d,f,i
versus Fig 2 h) and limited to scenarios where migration is ~10-15% and spatial synchrony in
recruitment is low. Local scale risks of collapse are even worse under a random (or
opportunistic) spatial allocation, but also ameliorated by reductions in harvest (SI Fig S.6).
Importantly, a 10% target harvest with optimized spatial allocation never exceeded 10% risk of
collapse in our simulations.

The discrepancy in risk of collapse at local subpopulation versus aggregate metapopulation
scales is shaped by both overall harvest rates and spatial variance in subpopulation trends. Risk
at the subpopulation scale matches that of metapopulation scales when subpopulations exhibit
little spatial variance (i.e. when there is high recruitment synchrony or high migration, top vs
bottom panels in Fig 2 c-j, SI Fig S.4). In contrast, risk diverges with increases in spatial
variance. Spatial variance is shaped not only by spatial recruitment synchrony and migration, but
also harvest rates and allocation in space. Fig 3b-c illustrates how different harvest allocations
impact spatial variation in trends (Fig 3 d, e). Higher harvest rates increase risk at subpopulation
scales in part because of higher depletion at the metapopulation scale (Fig 2 c,e,g,i) and also
because harvest can increase spatial variance (Fig 3d, explored more generally below with
analytical model) when efforts are allocated proportionally. This result demonstrates how low
demographic synchrony creates a portfolio effect at the metapopulation scale but simultaneously

[‡] **risk of collapse**: frequency of years with spawning biomass below 20% of unfished biomass

masks risks of collapse at the subpopulation scale which can be exacerbated with higher harvest
rates.

However, spatially optimizing harvest rates in space to match local scale variability lowers
the risk of subpopulation and aggregate metapopulation collapse (Fig 2 d vs f and Fig 2 h vs i)
with no cost to aggregate catch (SI Fig S.5). These results emerge from simulating spatially
optimized fishing effort that leaves underperforming locations unexploited and heavily targets
overperforming local subpopulations, in accordance with predictions of ideal free distribution
theory. Optimal spatial allocation of harvest rates reduces the effects of harvest intensity on
spatial variability (Fig 3d) which is strongly related to local risk of collapse (SI Appendix Fig
S.4). Thus, approaches that optimize spatial harvest to account for subpopulation dynamics can
minimize the spatially destabilizing effect of harvest as well as disparity in exposure to risk of
collapse among spatial scales. The benefits of spatially optimized harvest in terms of risk to
subpopulations diminishes as migration and spatial synchrony in recruitment productivity decline
to low levels (i.e. approach the lower left quadrant of panels in Fig 2c-h, where populations are
nearly autonomous with low connectivity).

Theoretical expectations of adult spatial & temporal variability in a metapopulation

To generalize findings from the Pacific herring example, our analytical model explores
metapopulation spatial variation mechanistically, starting with changing adult mortality. We find
that spatial variation among subpopulations increases with higher total mortality rates (via
harvest or otherwise) (Fig 4a) and decreases with migration and synchrony (SI Appendix C), in
line with the numerical simulation model results (Fig 3d). As a result, the metapopulation mean
and coefficient of variation are less reflective of its subpopulations (Fig 4b). In biological terms,
higher mortality decreases mean adult longevity (66) and thereby elevates sensitivity in
population fluctuations to recruitment pulses (67-70). The amplification of fluctuations occurs at
a higher rate at local scales than on aggregate. This response decreases spatial coupling and
predictability. Reduced longevity of adults (via higher mortality) reduces the abundance of adults
in each subpopulation that buffers against local stochasticity through survival and migration.

Mathematically, this result emerges for several related reasons. Increases in mortality
(constant across space in this case) reduce local population inertia (predictability, measured as 1^{st}
order temporal autocorrelation, thick black line in Fig 4c). This reduction in inertia increases
population sensitivity to temporal variation in local subpopulation recruitment (red line in Fig
4c). This sensitivity is illustrated by the primary impulse response (Equation 1), which describes
how a recruitment perturbation affects the intrinsic growth rate of each subpopulation (red line in
Fig 4c). In this case, the response always increases with total mortality (z).

Eq. 1: (Environmental impulse response): $(1 - \delta)/(1 + e^{-z} + e^{-2z})$

Such increases in local environmental sensitivity reduce spatial coupling (measured as the spatial
correlation, dotted black line in Fig 4c) and thereby increase spatial variability in abundance.

These patterns of asynchrony can persist even in the presence of modest migration rates (Fig
4b), another key mechanism explored in our analytical model. In this simple model, temporal
variability of local populations is controlled by the combination of migration (δ), mortality (z)
and recruitment synchrony in recruitment productivity (synchrony = $(\rho_R +1)/2$) (Fig 4b, see SI
Appendix for solutions). Spatial variation increases directly with decreases in recruitment
correlations (ρ_R) and with increases in recruitment variability. We illustrate this graphically for
modest migration rates ($\delta = 0.25$) in Fig 4a. Temporal variability at the local subpopulation scale
is highly sensitive to movement— spatial variance of adults approaches zero when the
population converges to 100% mixing (i.e. as migration approaches 0.5 for two subpopulations)
–while aggregate temporal variability of the metapopulation is not.

As a consequence of these combined properties, the risk of collapse at the metapopulation
level can obscure or underestimate local scale conservation risks. As with the empirical data and
numerical simulation model, aggregate trends can poorly reflect local trends and can appear
more predictable than local scale populations. Both analytical and empirical results demonstrate
that differences between individual subpopulations and the metapopulation are exacerbated with
increases in mortality (due to harvest or otherwise), which increase spatial variation across the
metapopulation.

Discussion

Harvest strategies that appear appropriately prescribed at large spatial scales can, at local
scales, lead to declines or even effective extirpation of local populations. We call these small-
scale declines "cryptic collapses". Specifically, regional harvest strategies can create a "gilded
trap" (71) where, in this case, management focuses on the aspects of metapopulations that can
benefit conservation and economics at the aggregate scale, but neglect social-ecological inequity
in the exposure to risk at local scales. Our results show that spatial mismatch among scales is not
merely an esoteric concern. Indeed, they occur in current management situations, and are
supported by ecological theory we develop here. Our multiscale risk analysis highlights the
impacts of scale mismatch on consumers and the potential value of optimized spatial
management for sustainability and equity.

Our work adds critical resolution and understanding to the literature on portfolio effects that
is focused on the benefits of spatial variability. Recent work viewing multiple populations as a
"portfolio" of assets has shown benefits of maintaining a diversity of populations with high
asynchrony. These benefits include reducing local extinctions through so called 'rescue effects'
(72-74), providing increased stability in the form of food security for people or animals (75) and
minimizing economic risks over large scales by minimizing variance in harvestable abundance

(21). Yet these portfolio analyses typically focus on the attributes of the aggregate
metapopulation, whereas the risk of localized population collapse (e.g. depletion below threshold
of ecological functionality or socioeconomic value) affects the interests of locally constrained
fishers and spatially constrained organisms with small home ranges. Using theory and data we
show that the same spatial variation that leads to resilience at the metapopulation scale, when left
unaccounted for in management strategies, can also produce unforeseen negative consequences
in the form of magnified local risk of collapse.

The consequences of ignoring spatial variation are most severe for spatially constrained
communities and predators. In the case of Pacific herring in the Central Coast of British
Columbia, local reductions in some subpopulations occurred well before the entire stock showed
substantial declines in the mid-2000s. This had greater impact on spatially constrained groups;
namely indigenous communities for whom herring is a source of cultural and economic vitality
(27, 76, 77), as well as animals with small home ranges that rely on herring. In contrast, mobile
fishing fleets and transient predators should be less vulnerable to local depletion events in the
short term. Yet in the long term, assumptions of spatial homogeneity can also lead to biased
estimates of total productivity (20) that may lead to overly optimistic harvest strategies.
Moreover, high local harvest rates may produce sequential depletion over time that eventually
erodes the principal of the stock leading to a collapse of the portfolio as a whole.

Pacific herring fisheries on Canada's west coast provide an empirical case where the scale of
regional stock assessments masked episodic local overexploitation and subpopulation collapses.
Our work exposes inequity among scales in risk exposure, with differentiation among predator
groups depending on their home-ranges (i.e. mobile versus non-mobile fishing fleets). Here,
high local harvest rates appear to be commonplace even when local subpopulations are depleted
(Fig 1) presumably because schooling fish are easy to catch even at low abundances (78). These
spatial, scale-dependencies affect which groups or species bear the brunt of management risks,
and who reaps the benefits from the portfolio payoff of regional metapopulation stability. Such
inequities highlight that ignoring space also excludes critical social, ecological and economic
responses central to the triple bottom line (79). While high local harvest rates may have
contributed to subsequent local collapses observed in this case study, quantifying other
confounding demographics (i.e. spatial variation in stochastic adult mortality) are necessary to
explicitly test this hypothesis. While recent local collapses may have occurred in the absence of
fishing, fishing is very likely imposing higher population sensitivity to any environmental or
biotic influence on recruitment or survival by reducing adult longevity thereby eroding an
important buffer against recruitment volatility (80) that we show can contribute to spatial
variation.

Importantly, spatial variation in population dynamics is not independent of harvest dynamics.
Rather, we demonstrate numerically and analytically that spatial variation is likely to increase
with harvest. However, if the distribution of harvest in space can be optimized, we show how the

overall portfolio benefits at multiple scales. Here, demographic asynchrony (in this case
asynchrony in recruitment productivity), can be maintained while minimizing subpopulation risk.
Translated into practice, our simulation and analytical results highlight two non-mutually
exclusive potential solutions for equitably spreading risk among scales: 1) reductions in overall
harvest rates can ameliorate biases in risk among scales by reducing baseline levels of risk and
reducing effects of harvest on spatial variability. 2) spatially optimizing harvest allocation can
minimize spatial variance and ease pressure on at-risk subpopulations, thereby reducing risk of
local scale depletion without sacrificing catch. The former creates a trade-off between
commercial yield and local risk; the latter between the cost of management and local risk.

Addressing spatial inequity in risk exposure requires confronting these economic and
logistical trade-offs. For species such as Pacific herring that have volatile spatiotemporal
dynamics and complex migratory phenology (32), polycentric governance structures where
governing authorities are nested at different spatial scales may help balance these trade-offs by
addressing the problems of fit between ecosystems, social systems and management agencies
(26, 81-85). Such systems can capitalize on scale-specific ecological knowledge (including local,
traditional and scientific knowledge), scientific capacity and socioeconomic experience to 1)
guide decision analyses, 2) co-coordinate data collection and harvest allocation in space, and 3)
test policies [e.g. in the Maine Lobster fishery (86)]. Polycentric management schemes are not a
silver bullet. However, for systems like Pacific herring where placed-based indigenous fishing
communities often object to purely centralized scales of management for social, ecological and
economic reasons (26, 27, 76, 77, 87), opportunities to integrate knowledge and objectives into
management strategies (and their evaluation) at smaller spatial scales is well placed in managing
these fisheries, and may be critical to their perpetuity.

The models used to generate inference in this study are simple in comparison to the nature of
complex stochastic systems in space. We ignore many scenarios that are likely to further
complicate spatial patterns (i.e. behaviorally or geographically complex migration, spatial and
age specific mortality, spatially complex density dependence, Allee effects, cost and data
limitations for spatial management, etc.). However, the principles from our simulation and
analytical models should generalize regardless of the degree of complexity in the system: a
precautionary approach cognizant of resource users across multiple spatial scales may necessitate
the incorporation of some degree of locally based management to minimize spatial discrepancies
in risk exposure. Overall, our models suggest a mixture of management scales may be key to
selecting and implementing harvest levels, and informing how regional harvest is coordinated in
space to navigate towards both sustainable and equitable outcomes (83).

Our analyses illustrate the importance of considering spatial dynamics for determining how
to most effectively balance equity in management and conservation strategies aimed at achieving
social, economic and ecological outcomes (88, 89). These issues are often ignored by centralized
management and conservation initiatives focused on larger spatial scales. For over half a

century, fisheries scientists have debated how best to exploit and conserve "mixed stocks" that
have separate dynamics but are inseparable in space with the aim of balancing conservation of
weak stocks and maximizing total yields (90). Our work inverts this focus to metapopulations,
populations with inseparable dynamics that are separated in space. We show how, in these
settings, exploitation can magnify local scale fluctuations and spatial variability. As a result
aggregate metrics poorly represent local scale dynamics (Fig 3d,e, Fig 4) and the risk of collapse
at the local scale increases at a greater rate than the aggregate large scale (Fig 2). This
phenomenon creates cryptic collapses with ensuing discrepancies in risk exposure. Fortunately,
the magnitude of these discrepancies can potentially be controlled by lowering harvest rates or
seeking harvest dynamics that are spatially optimized (Fig 3). Overall, these conclusions are
relevant not only to Pacific herring fisheries, but also to the great number of exploited resources
that exhibit spatial structure and are valuable to species and people that operate on multiple
scales.

Methods

Empirical Pacific Herring Case Study in the British Columbia's Central Coast

Foraging-ranges of select Pacific herring predators

References were searched for using a combination of Web of Science and Google Scholar.
Searches were conducted for papers published before April 2017 using various combinations of
the following terms: The scientific and common name of target species, range, foraging range,
home range, migration, telemetry, tracking, and movement. Range sizes were determined based
on results directly reported in the text of each study, or derived from data included in
supplementary materials.

Spatiotemporal variability in Pacific herring biomass and catch

To estimate patterns of spatial and temporal variability in herring biomass and harvest rates,
we used spawn deposition and harvest data for six major Pacific Herring spawning units
(groupings of spawn sections) that comprise the Central Coast stock in British Columbia,
Canada. In the Central Coast, sections are delineated by spatially isolated spawning
aggregations. The remaining sections comprised less than 1% of the biomass across the time
series duration. Briefly, SCUBA surveys are used to estimate herring egg abundance which is
converted to total spawning biomass with the assumed conversion of ~100 eggs/g and an equal
sex ratio (25, 91). Catch (which occurs in the days prior to spawning) is reported by geographic
section. For full description of the time series see (25) and references therein.

To reconstruct spatial biomass from the biomass and harvest time series observations, we use
a nonlinear model in a Bayesian hierarchical state-space framework. We modeled biomass as:

Eq. 2: $\hat{b}^{+}_{t+1,l} = \exp\left[\alpha_l + (1 - \beta_l)\ln\left[b_{t-1,l}\right] + \lambda_l b_{t,l}\right]$

where, $\hat{b}^{+}_{t+1,l}$ is the expected pre-spawn biomass at site l in year t+1 given the estimate of

spawning biomass after harvest from previous years ($b_{t,l}$), growth parameters of a Gompertz

model (α_l & β_l) and a survivorship term (λ_l). We estimate latent spawning biomass following

harvest ($b_{t,l}$) and convert to pre-harvest biomass by adding known local harvest ($h_{t,l}$).

The latent biomass vector $\boldsymbol{b}_t = [b_{1,t} \; b_{2,t} \ldots b_{L,t}]'$ has a multivariate normal conditional

probability:

Eq. 3: $\ln[\boldsymbol{b}_t + \boldsymbol{h}_t] \sim \mathrm{MVN}(\ln \hat{\boldsymbol{b}}^{+}_t , \boldsymbol{\sigma}\Omega)$

We adjust the log-probability for the change of variables by the log-Jacobian of the

transformed left-hand side of the equation (92). Ω is the process error correlation matrix and $\boldsymbol{\sigma}$ is

the vector of within-site process standard deviations.

We assume the observed spawning biomass ($\mathrm{obs}[b_{t,l}]$) is lognormally distributed:

Eq. 4: $\mathrm{obs}[b_{t,l}] \sim \mathrm{lognormal}(\ln b_{t,l} + \ln q , \sigma_{obs})$

where ln q is the log-proportion of spawning biomass that is observable, with a normal prior

borrowed from the herring assessment (25). The lognormal observation error (σ_{obs}) has a half-

Cauchy prior with scale parameter 2.5.

We modeled the location specific parameter vectors in a multivariate, hierarchical framework

as:

Eq. 5: $\Theta_l \sim \mathrm{MVN}(\Theta_\mu , \boldsymbol{\sigma}_\Theta \Omega_\Theta)$

Where $\Theta_l = [\alpha_l \quad \mathrm{logit}(\beta_l) \quad \mathrm{logit}(\lambda_l)]'$. Elements of the mean parameter vector Θ_μ were

given a normal prior with mean 0 and sd = 1.5 (which is approximately uniform on the

untransformed scale). Ω_Θ is the process error correlation matrix and $\boldsymbol{\sigma}_\Theta$ is the vector of within-

site process standard deviations. We used a Lewandowski-Dorota-Joe LKJ prior (93) for the

Cholesky decomposition of the correlation matrices (Ω and Ω_Θ) and the vectors of process

standard deviations ($\boldsymbol{\sigma}$ and $\boldsymbol{\sigma}_\Theta$). We estimated the models using Stan (92, 94-96) via R (97) with

3, 1000 iteration chains after burn-in.

We used the model posterior to calculate 1) total biomass at each site at each year, 2) metrics

of biomass temporal and spatial variation (see SI Appendix A), and 3) the exploitation rate ($h_{t,l}$/

$b_{t,l}$). We compared the observed local harvest rates to the theoretical proportional allocation

[sum($h_{t,l}$)/sum($b_{t,l}$)] and optimized allocation of harvest given the quota. The optimized

allocation of catch distributes catches in space according to the ideal free distribution. This

distribution removes proportionally more biomass from subpopulations with higher biomass and

is the spatial allocation of catch that minimizes effort to achieve the overall quota assuming catch

per unit effort is linearly related to biomass (see SI Appendix A for mathematical details).

Risk analysis with spatial closed-loop simulation of Pacific herring fisheries

We use a spatial, stochastic model to simulate how spatial metapopulation dynamics and
fishing behavior interact to influence risk of overfishing (frequency of years with spawning
biomass below 20% of unfished biomass) at the subpopulation and metapopulation scales under
a diverse suite of scenarios that include a factorial gradient of 1) annual movement (see Annual
adult migration among spawning areas below), 2) environmental recruitment synchrony (the
degree of correlation in recruitment in space – see Spatiotemporal Recruitment Dynamics
below), 3) a range of harvest rates (see Stock Harvest Quota below), and 4) allocation of harvest
in space (see Spatial Harvest Prosecution below).

For all analyses, we simulate 10 subpopulations placed equidistant around a hypothetical
island of arbitrary size such that oceanographic distance from subpopulation 1 to 2 is the same as
1 to 10. Fish spawning at any site in a given year can move to any other site in the next year,
with a probability that declines with distance. Likewise, synchrony in stochastic recruitment
among sites decays with distance. Stochasticity in dynamics is included in recruitment (spatial
and temporal variation), survival (temporal variation only) and movement probabilities (spatial
and temporal variance).

Abundance of fish in location l at age a in year t is represented by $n_{a,l,t}$. The order of
operations mathematically is 1) recruitment, 2) survival, 3) movement, 4) roe fishery harvest,
and 5) spawning. The abundance is respectively denoted $n_{a,l,t}^{++}$, (after recruitment and survival
[for age 2] or survival [for age 2+]), $n_{a,l,t}^{+}$ (after movement) and $n_{a,l,t}$ (after harvest).

Spatiotemporal Recruitment Dynamics

Following the assumptions of the current British Columbia herring assessments (23, 25),
recruitment of age 2 individuals at location l is a function of eggs produced 2-years prior at
location l (eggs$_{l,t}$) and local density dependence. Recruitment of age 2 individuals is
characterized by the Beverton-Holt model:

Eq. 6: $n_{a=2,l,t+2}^{++} = \text{eggs}_{l,t} \exp\left[\epsilon_{l,t} - \frac{\sigma_r^2}{2}\right]/(\alpha + \beta \text{eggs}_{l,t})$

Where α and β are stock-recruit parameters [595.25 & 1.79×10^{-8} derived from (23)], $\epsilon_{l,t}$ is
the error at time t at location l with standard devation σ_ϵ , eggs$_{l,t}$ is the product of abundance at
age ($n_{a,l,t}$), maturity [m_a = 0.25 (age 2), 0.9 (3), 1 (4+) from (25)] and female fecundity $f_a =$
$\exp[4.69](w_a)^{1.13}$ where w_a (grams at age) = $4.5 \times 10^{-3}(27(1 - \exp[-0.48 * \text{age}])^{3.127}$ from
(25, 98):

Eq. 7: eggs$_{l,t} = \sum_{a=2}^{10} m_a f_a [0.5 n_{a,l,t}]$

We allow recruitment to exhibit both spatial and temporal variability and autocorrelation
with a first-order vector autoregressive model [VAR(1)]:

Eq. 8: $\epsilon_t = \phi\epsilon_t + \text{MVN}(0, \sigma_r{}^2[1-\phi^2]\Omega_r)$

ϕ is a 1st order autoregressive term, and $\mathbf{e}_t$ is a multivariate normal error vector with spatial

correlation matrix Ω_r. Spatial correlations follow a Gaussian spatial decay with distance (d),

controlled by the standard deviation σ_Ω. For each scenario, σ_r is numerically tuned such that net

recruitment variability is approximately constant across scenarios [$\sigma_{r.mu}$ = metapopulation level

coefficient of variation = 0.8 (23)].

Adult Survival

All adult survival occurs prior to movement into spawning locations, yielding pre-movement,

pre-harvest abundance:

Eq. 9: $n_{a,l,t+1}^{++} = \begin{cases} \lambda_t\, n_{a-1,l,t} & 2 < a < 10 \\ \lambda_t\, n_{a-1,l,t} + \lambda_t\, n_{a,l,t} & a = 10 \text{ (plus group)} \end{cases}$

Survival in any given year is constant in space and is randomly drawn from a beta-binomial

with mean $\lambda_t = 0.6$ with coefficient of variation of 0.2, derived from (25).

Annual Adult Migration Among Spawning Areas

Survivors migrate to new spawning areas to yield a matrix of pre-spawn, pre-harvest matrix

of abundance ($\mathbf{N}_t{}^+$) describing abundance at each age given by the matrix product:

Eq. 10: $\mathbf{N}^+{}_t = \mathbf{N}^{++}{}_t\, \mathbf{S}_t$

$\mathbf{S}_t$ is a square matrix of linear migration rates. Row elements of $\mathbf{S}_t$ sum to one and are

stochastic Dirichlet realizations with shared scale parameter (scale =10) and a mean migration

probability controlled by a scaled Cauchy decay given a distance from site l to site i ($dist_{l \to i}$)

and a migration scaling parameter (γ):

Eq. 11: $\text{P}(migration_{l \to i} \mid dist_{l \to i}, \gamma) = \left[\gamma\left(1 + \frac{dist_{l \to i}}{\gamma}\right)^2\right]^{-1} / \sum_{j=1}^{L}\left[\gamma\left(1 + \frac{dist_{l \to i}}{\gamma}\right)^2\right]^{-1}$

γ is numerically tuned to achieve a desired retention rate for each scenario. For reference

regarding stochasticity, if the mean migration rate is 0.2, the coefficient of variation in migration

among years with scale =10 is 0.1.

Stock Harvest Quota

We follow the existing harvest control rules of Pacific herring in British Columbia (23, 25).

The annual biomass harvest quota for the stock (Q_{t+1}) is designed to achieve a target harvest rate

(H_{Target}) and a minimum spawning escapement (25% of B_0 - the average steady state biomass

without fishing).

Eq. 12: $Q_{t+1} = \begin{cases} \min(\ H_{Target}\ B_{Forecast,t}, B_{Forecast,t} - 0.25B_0\ ) & \text{if } B_{Forecast,t} > 0.25B_0 \\ 0 & \text{otherwise} \end{cases}$

For simplicity, forecast biomass made in year t ($B_{Forecast,t}$) for year ($t+1$) is the expected

biomass assuming deterministic recruitment (i.e. without stochastic variability). In order to avoid

evaluating stock assessment model performance (which is out of the scope of this study) we

avoid assessments explicitly within the closed loop. Rather, we assume forecasts are unbiased

with no observation noise.

Spatial Harvest Prosecution

We use two spatial fleet allocation scenarios to generate the distribution of spatial harvest

given the stock quota (Q_t): 1) the fleet prosecutes the fishery in direct proportion to spawning

biomass (proportional allocation) or 2) the fleet is allocated according to the ideal free

distribution (IFD) that in theory maximizes catch efficiency. Here, realized catch is nonlinearly

related to spawning biomass, harvesting more from areas with higher abundance and leaving

alone areas with lower abundance. See SI Appendix B for details in generating the IFD and an

additional case with a random spatial allocation more similar to the empirical case study. In all

cases, fishery selectivity is identical to maturity-at-age.

Given the local harvest ($h_{l,t}$) produced by these scenarios the post-harvest abundance is:

Eq. 13: $n_{a,l,t} = n_{a,l,t}^+ - m_a n_{a,l,t}^+ h_{l,t} / \sum_{a=2}^{10} w_a m_a n_{a,l,t}^+$

Simulation details

For each simulation, we ran the following simulation algorithm, which: 1) evaluated

deterministic equilibria without fishing, 2) initiated stochastic forcing of recruitment from the

equilibrium for 12 years, 3) initiated the fishery, and 4) recorded statistics for years 23-52. We

ran 100 replicate simulations for each parameter combination, recording a) mean number of

years below 20% B_0 at both stock and substock scales (risk of collapse), b) stock and substock

level catch, c) stock and substock biomass, d) stock and substock temporal variability

(coefficient of variation) in biomass, and e) spatial variation in biomass (difference between

squared coefficient of variation at the substock versus stock scale).

Theoretical impact of mortality, migration and synchrony on spatial and temporal variability

in stochastic metapopulations

To evaluate how mortality, migration and demographic synchrony interact to impact

temporal variability, spatial variability and sensitivity to a single perturbation for local

populations and the metapopulation, we use a stochastic metapopulation model with two

populations in discrete time. We convert the model to a first order Vector Autoregressive model

(VAR(1)) where statistical properties of stochastic forcing in multivariate systems are well

described (99) . We explore this simple model for generality across the diversity of species and

management applications discussed above. Using this model we calculate 1) temporal variability
(temporal coefficient of variation) of the populations and metapopulation, 2) spatial variability
(the differences between the squared temporal coefficient of variation at the populations and
metapopulation), 3) the sensitivity to recruitment via an impulse response analysis that measures
how a recruitment pulse propagates through the population, 4) the predictability of the
populations as measured by the first-order autocorrelation for the populations, and 5) the spatial
coupling measured as the spatial autocorrelation in the metapopulation.

We assume that individuals in the population are mature at age 2, adults move among
populations by linear diffusion, all adults are identical, and environmental stochasticity only
affects survival of recruits (survival to pre-movement age 2 fish).

Mathematically, the simple model is represented as follows. Let $Y_{a,i,t}$ represent the adults of
age a at location i in year t and δ is the per capita annual probability of moving among sites and
e^{-M} is the survival rate. Then the equations for dynamics in site i are given as follows.

Total reproduction is the sum of adults across all adult age classes.

Eq. 14: $Y_{0,i,t+1} = 0.5e^{-M} \sum_{a=1}^{n}(1-\delta)Y_{a,i,t} + \delta Y_{a,j,t}\,,\; 0 \le \delta \le 0.5$

The stock recruit function that determines how zygotes translate into one-year-olds is a
Gompertz function:

Eq. 15: $Y_{1,i,t+1} = \alpha(Y_{0,i,t})^{1-\beta}\exp\left[\zeta_{i,t}\right]$

Survival and migration are given by:

Eq. 16: $Y_{a,i,t+1} = \begin{cases} e^{-M}\left[(1-\delta)Y_{a-1,i,t} + \delta Y_{a-1,j,t}\right] & 2 \le a \le n-1 \\ e^{-M}\left[(1-\delta)(Y_{a-1,i,t} + Y_{a,i,t}) + \delta(Y_{a-1,j,t} + Y_{a,j,t})\right] & a = n \text{ (plus group)} \end{cases}$

The vector $\boldsymbol{\xi}_t = [\zeta_{1,t},\zeta_{2,t}]'$ follows a multivariate normal distribution [i.e. a multivariate
extension of the model of Bjornstad et al. (70)] with mean zero and variance-covariance matrix
$\boldsymbol{\Sigma}_{\boldsymbol{\xi}}$, whose entries are controlled by variance given by σ_R^2 and covariance controlled by the
spatial correlation ρ_R. Thus, synchrony among the two populations is regulated by the spatial
correlation in environmental noise and the degree of migration between populations (with a
maximum migration probability of 0.5 providing a single, fully mixed population).

To evaluate how the nature of variability is shaped by correlation in recruitment stochasticity
(controlled by covariance in $\boldsymbol{\xi}_t$) and how migration probability (δ) influences variability at the
local and metapopulation scales, we can cast the model in terms of log-scale deviations from the
equilibrium. To do so, we vectorize the variables and convert the log-scale model (i.e $x_{a,i,t} =$
$\ln Y_{a,i,t} - \ln Y_{a,i}^*$ where $Y_{a,i}^*$ is the equilibrium) into a VAR(1) model by linearizing about the
equilibrium (70, 100). This approach approximates the deterministic dynamics of the nonlinear

model with a Jacobian matrix (**J**) of first order dependencies and an environmental covariance

matrix. As a result, the matrix of coefficients (with plus group summed at age *n*) is given by the

block matrix comprised of among and within population age transitions:

Eq. 17:
$$\mathbf{J} = \begin{bmatrix} J_1 & J_2 \\ J_2 & J_1 \end{bmatrix}$$

502   defined by within and population dynamics given by:

503   **Eq. 18:**
$$J_i = \begin{bmatrix} 0 & \frac{D}{1+\sum_{n=1}^{n-1}\lambda^n} & \frac{(D)\lambda}{1+\sum_{n=1}^{n-1}\lambda^n} & \frac{(D)\lambda^2}{1+\sum_{n=1}^{n-1}\lambda^n} & \cdots & \frac{(D)\lambda^{n-2}}{1+\sum_{n=1}^{n-1}\lambda^n} & \frac{(D)\lambda^{n-1}}{1+\sum_{n=1}^{n-1}\lambda^n} \\ 1-\beta & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & D & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & D & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & \frac{D}{\lambda+1} & \frac{(D)\lambda}{\lambda+1} \end{bmatrix}$$

Where $D = (1-\delta)$ if $J_i = J_2$ and δ if $J_i = J_2$.

The resulting VAR(1) model is:

Eq. 19: $\mathsf{x}_t = \mathsf{J}\,\mathsf{x}_{t-1} + \mathsf{Z}\boldsymbol{\xi}_t$

Eq. 20: $\mathbf{x}_t = [x_{L=1,a=1,t} \quad \cdots \quad x_{1,n,t} \quad x_{2,1,t} \quad \cdots \quad x_{2,n,t}]'$

Z is a 2n x L binary matrix that controls which of the entries in $\mathbf{x}_t$ are subject to stochasticity

terms.

We use the known statistical properties of a VAR(1) model (99) in tandem with the moments

of a multivariate lognormal to derive the coefficient of variation for the population and

metapopulation as well as spatial variation in the metapopulation. See SI Appendix C for

derivation. Finally, to evaluate how local environmental sensitivity changes with mortality rate

we analyze the model using impulse response analysis for a VAR(1) (99). The impulse response

function in this case gives the log-scale adult response to a stochastic environmental shock in $\zeta_{1,t}$

(log-scale unit deviation in recruitment) and is interpreted as the annual intrinsic growth rate

response to the shock. See SI Appendix C for details.

Data along with R, Stan and Mathematica code used for this study are available for download

from GitHub.

Acknowledgements:

DKO was supported by a Strategic Grant from the National Sciences and Engineering

Research Council of Canada to Simon Fraser University. PSL thanks the Gordon and Betty

Moore Foundation for their support of the Ocean Tipping Points project and the David and

Lucille Packard Foundation for their support of the Ocean Modeling Forum. We thank J. Cleary,

A. Frid and lab group of A. Rassweiler for input on the manuscript, and S. Cox, L. Dee, L.

Hauser, B. Hunt, S. Miller, I. McKechnie, S. Pau, T. Pitcher, E. Petrou and W. Smith for
discussions that improved the research. We also thank the Heiltsuk and Haida First Nations for
their partnership in the NSERC Strategic Grant and the Department of Fisheries and Oceans for
providing herring spatial data.

References:

1. Cumming G, Cumming DH, Redman C (2006) Scale mismatches in social-ecological
systems: causes, consequences, and solutions. *Ecol Soc* 11(1).

2. Levin SA (1992) The problem of pattern and scale in ecology: the Robert H. MacArthur
award lecture. *Ecology* 73(6):1943-1967.

3. Chesson P (1996) Matters of scale in the dynamics of populations and communities.
Frontiers of population ecology, eds Floyd R, Sheppard A, de Barro P (CSIRO, Victoria,
Australia), pp 353-368.

4. Chesson P (1998) Spatial scales in the study of reef fishes: a theoretical perspective. *Aust
J Ecol* 23(3):209-215.

5. Englund G, Cooper SD (2003) Scale effects and extrapolation in ecological experiments.
Adv Ecol Res 33:161-213.

6. Wiens JA, Chr N, Van Horne B, Ims RA (1993) Ecological mechanisms and landscape
ecology. *Oikos*:369-380.

7. Johnson T, Wilson J, Cleaver C, Vadas R (2012) Social-ecological scale mismatches and
the collapse of the sea urchin fishery in Maine, USA. *Ecol Soc* 17(2).

8. Melbourne BA, Chesson P (2006) The scale transition: scaling up population dynamics
with field data. *Ecology* 87(6):1478-1488.

9. Melbourne BA, Chesson P (2005) Scaling up population dynamics: integrating theory
and data. *Oecologia* 145(2):178-186.

10. Chesson P, Donahue MJ, Melbourne BA, Sears AL (2005) Scale transition theory for
understanding mechanisms in metacommunities. *Metacommunities: spatial dynamics and
ecological communities*, eds Holyoak M, Leibold M, Holt R (University of Chicago
Press), pp 279-306.

11. Mangel M, Levin PS (2005) Regime, phase and paradigm shifts: making community
ecology the basic science for fisheries. *Phil Trans Roy Soc B* 360(1453):95-105.

12. Kendall BE, Bjørnstad ON, Bascompte J, Keitt TH, Fagan WF (2000) Dispersal,
environmental correlation, and spatial synchrony in population dynamics. *Am Nat*
155(5):628-636.

13. Lande R (1988) Demographic models of the northern spotted owl (Strix occidentalis
caurina). *Oecologia* 75(4):601-607.

14. Peterson DA, Hilborn R, Hauser L (2014) Local adaptation limits lifetime reproductive
success of dispersers in a wild salmon metapopulation. *Nat Commun* 5.

15.   Rieman B, Dunham J (2000) Metapopulations and salmonids: a synthesis of life history
564   patterns and empirical observations. *Ecol Freshwat Fish* 9(1-2):51-64.

565   16.   Schtickzelle N, Quinn TP (2007) A metapopulation perspective for salmon and other
566   anadromous fish. *Fish Fish* 8(4):297-314.

567   17.   Marsh DM, Trenham PC (2001) Metapopulation dynamics and amphibian conservation.
568   *Conserv Biol* 15(1):40-49.

569   18.   Stephenson RL (1999) Stock complexity in fisheries management: a perspective of
570   emerging issues related to population sub-units. *Fish Res* 43(1):247-249.

571   19.   Chadès I, *et al.* (2011) General rules for managing and surveying networks of pests,
572   diseases, and endangered species. *Proc Natl Acad Sci* 108(20):8323-8328.

573   20.   Takashina N, Mougi A (2015) Maximum sustainable yields from a spatially-explicit
574   harvest model. *J Theor Biol* 383:87-92.

575   21.   Schindler DE, *et al.* (2010) Population diversity and the portfolio effect in an exploited
576   species. *Nature* 465(7298):609-612.

577   22.   McKechnie I, *et al.* (2014) Archaeological data provide alternative hypotheses on Pacific
578   herring (*Clupea pallasii*) distribution, abundance, and variability. *Proc Natl Acad Sci*
579   111(9):E807-E816.

580   23.   Department of Fisheries and Oceans (2015) Stock Assessment and Management Advice
581   for BC Pacific Herring: 2015 Status and 2016 Forecast. . *Can. Sci. Adv. Sec. Res. Doc*
582   2015/038.

583   24.   Lepofsky D, Caldwell M (2013) Indigenous marine resource management on the
584   Northwest Coast of North America. *Ecol Process* 2(1):1-12.

585   25.   Martell S, Schweigert J, Haist V, Cleary J (2011) Moving towards the sustainable
586   fisheries framework for Pacific herring: data, models, and alternative assumptions; Stock
587   Assessment and Management Advice for the British Columbia Pacific Herring Stocks:
588   2011 Assessment and 2012 Forecasts. *Can. Sci. Adv. Sec. Res. Doc* 136:163.

589   26.   von der Porten S, Lepofsky D, McGregor D, Silver J (2016) Recommendations for
590   marine herring policy change in Canada: Aligning with Indigenous legal and inherent
591   rights. *Mar Policy* 74:68-76.

592   27.   Powell M (2012) Divided waters: Heiltsuk spatial management of herring fisheries and
593   the politics of native sovereignty. *West Hist Q* 43(4):463-484.

594   28.   Flostrand LA, Schweigert JF, Daniel KS, Cleary JS (2009) Measuring and modelling
595   Pacific herring spawning-site fidelity and dispersal using tag-recovery dispersal curves.
596   *ICES J Mar Sci* 66(8):1754-1761.

597   29.   Hay D, McCarter P, Daniel K (2001) Tagging of Pacific herring *Clupea pallasi* from
598   1936 1992: a review with comments on homing, geographic fidelity, and straying. *Can J*
599   *Fish Aquat Sci* 58(7):1356-1370.

600   30.   Levin PS, Francis TB, Taylor NG (2016) Thirty-two essential questions for
601   understanding the social–ecological system of forage fish: the case of Pacific Herring.
602   *Ecosyst Health Sustain* 2(4).

31. Jones R, Rigg C, Pinkerton E (2016) Strategies for assertion of conservation and local
management rights: A Haida Gwaii herring story. *Mar Policy*.

32. Benson AJ, Cox SP, Cleary JS (2015) Evaluating the conservation risks of aggregate
harvest management in a spatially-structured herring fishery. *Fish Res* 167:101-113.

33. Siple MC, Francis TB (2016) Population diversity in Pacific herring of the Puget Sound,
USA. *Oecologia* 180(1):111-125.

34. Wang S, Loreau M (2014) Ecosystem stability in space: α, β and γ variability. *Ecol Lett*
17(8):891-901.

35. Harris DC (2000) Territoriality, aboriginal rights, and the Heiltsuk spawn-on-kelp
fishery. *U. Brit. Colum. L. Rev.* 34:195.

36. Hines AH (1982) Coexistence in a kelp forest: size, population dynamics, and resource
partitioning in a guild of spider crabs (Brachyura, Majidae). *Ecol Monogr* 52(2):179-198.

37. Stone RP, O'Clair CE (2001) Seasonal movements and distribution of Dungeness crabs
Cancer magister in a glacial southeastern Alaska estuary. *Mar Ecol Prog Ser* 214:167-
176.

38. Cieciel K, Pyper BJ, Eckert GL (2009) Tag retention and effects of tagging on movement
of the giant red sea cucumber Parastichopus californicus. *N Am J Fish Manage*
29(2):288-294.

39. Mattison J, Trent J, Shanks A, Akin T, Pearse J (1976) Movement and feeding activity of
red sea urchins (*Strongylocentrotus franciscanus*) adjacent to a kelp forest. *Mar Biol*
39(1):25-30.

40. Beaudreau A, Essington T (2011) Use of pelagic prey subsidies by demersal predators in
rocky reefs: insight from movement patterns of lingcod. *Mar Biol* 158(2):471-483.

41. Green KM, Starr RM (2011) Movements of small adult black rockfish: implications for
the design of MPAs. *Mar Ecol Prog Ser* 436:219-230.

42. Jorgensen SJ*, et al.* (2006) Limited movement in blue rockfish *Sebastes mystinus*:
internal structure of home range. *Mar Ecol Prog Ser* 327:157-170.

43. Mitamura H*, et al.* (2009) Preliminary study on homing, site fidelity, and diel movement
of black rockfish *Sebastes inermis* measured by acoustic telemetry. *Fish Sci* 75(5):1133-
1140.

44. Tolimieri N, Andrews K, Williams G, Katz S, Levin P (2009) Home range size and
patterns of space use by lingcod, copper rockfish and quillback rockfish in relation to diel
and tidal cycles. *Mar Ecol Prog Ser* 380:229-243.

45. Freiwald J (2012) Movement of adult temperate reef fishes off the west coast of North
America. *Can J Fish Aquat Sci* 69(8):1362-1374.

46. Peterson SH, Lance MM, Jeffries SJ, Acevedo-Gutiérrez A (2012) Long distance
movements and disjunct spatial use of harbor seals (*Phoca vitulina*) in the inland waters
of the Pacific Northwest. *PloS one* 7(6):e39046.

47.   Ward EJ, Levin PS, Lance MM, Jeffries SJ, Acevedo-Gutiérrez A (2012) Integrating diet
and movement data to identify hot spots of predation risk and areas of conservation
concern for endangered species. *Cons Lett* 5(1):37-47.

48.   Lorenz TJ, Raphael MG, Bloxton TD, Cunningham PG (2017) Low breeding propensity
and wide-ranging movements by marbled murrelets in Washington. *J Wildl Manage*
81(2):306-321.

49.   Peery MZ, Newman SH, Storlazzi CD, Beissinger SR (2009) Meeting reproductive
demands in a dynamic upwelling system: foraging strategies of a pursuit-diving seabird,
the marbled murrelet. *The Condor* 111(1):120-134.

50.   Barbaree BA, Nelson S, Dugger BD (2015) Marine space use by marbled murrelets
Brachyramphus marmoratus at a mainland fjord system in southeast Alaska. *Mar*
Ornithol 116:173-184.

51.   Moser ML, Myers MS, West JE, O'Neill SM, Burke BJ (2013) English sole spawning
migration and evidence for feeding site fidelity in Puget Sound, USA, with implications
for contaminant exposure. *Northwest Sci* 87(4):317-325.

52.   Dalla Rosa L, Secchi E, Maia Y, Zerbini A, Heide-Jørgensen M (2008) Movements of
satellite-monitored humpback whales on their feeding ground along the Antarctic
Peninsula. *Polar Biol* 31(7):771-781.

53.   Kennedy AS, Zerbini AN, Rone BK, Clapham PJ (2014) Individual variation in
movements of satellite-tracked humpback whales *Megaptera novaeangliae* in the eastern
Aleutian Islands and Bering Sea. *Endanger Species Res* 23(2):187-195.

54.   Fearnbach H*, et al.* (2014) Spatial and social connectivity of fish-eating "Resident" killer
whales (*Orcinus orca*) in the northern North Pacific. *Mar Biol* 161(2):459-472.

55.   Hauser DD, Logsdon MG, Holmes EE, VanBlaricom GR, Osborne RW (2007) Summer
distribution patterns of southern resident killer whales *Orcinus orca*: core areas and
spatial segregation of social groups. *Mar Ecol Prog Ser* 351:301-310.

56.   Pearce JM, Reed JA, Flint PL (2005) Geographic variation in survival and migratory
tendency among North American Common Mergansers. *J Field Ornithol* 76(2):109-118.

57.   Kuhn CE, Costa DP (2014) Interannual variation in the at-sea behavior of California sea
lions (*Zalophus californianus*). *Mar Mamm Sci* 30(4):1297-1319.

58.   Merrick RL, Loughlin TR (1997) Foraging behavior of adult female and young-of-the-
year Steller sea lions in Alaskan waters. *Can J Zool* 75(5):776-786.

59.   Kuhn C, Ream R, Sterling J, Thomason J, Towell R (2014) Spatial segregation and the
influence of habitat on the foraging behavior of northern fur seals (Callorhinus ursinus).
Can J Zool 92(10):861-873.

60.   Loher T (2008) Homing and summer feeding site fidelity of Pacific halibut
(*Hippoglossus stenolepis*) in the Gulf of Alaska, established using satellite-transmitting
archival tags. *Fish Res* 92(1):63-69.

61.   Nielsen JK, Hooge PN, Taggart SJ, Seitz AC (2014) Characterizing Pacific halibut
680   movement and habitat in a Marine Protected Area using net squared displacement
681   analysis methods. *Mar Ecol Prog Ser* 517:229-250.

682   62.   Seitz AC, Loher T, Norcross BL, Nielsen JL (2011) Dispersal and behavior of Pacific
683   halibut *Hippoglossus stenolepis* in the Bering Sea and Aleutian Islands region. *Aquat Biol*
684   12(3):225-239.

685   63.   Hanselman DH, Heifetz J, Echave KB, Dressel SC (2014) Move it or lose it: movement
686   and mortality of sablefish tagged in Alaska. *Can J Fish Aquat Sci* 72(2):238-251.

687   64.   Rand KM, Munro P, Neidetcher SK, Nichol DG (2014) Observations of seasonal
688   movement from a single tag release group of Pacific cod in the eastern Bering Sea. *Mar*
689   *Coast Fish* 6(1):287-296.

690   65.   Wespestad VG, Thorsen K, Mizroch S (1983) Movement of sablefish, *Anoplopoma*
691   *fimbria*, in the northeastern Pacific Ocean as determined by tagging experiments (1971-
692   80). *Fish Bull* 81(2):415-420.

693   66.   Barnett LA, Branch TA, Ranasinghe RA, Essington TE (2017) Old-Growth Fishes
694   Become Scarce under Fishing. *Curr Biol* 27(18):2843-2848. e2842.

695   67.   Beddington JR, May RM (1977) Harvesting natural populations in a randomly fluctuating
696   environment. *Science* 197(4302):463-465.

697   68.   Hsieh C*, et al.* (2006) Fishing elevates variability in the abundance of exploited species.
698   *Nature* 443(7113):859-862.

699   69.   Shelton AO, Mangel M (2011) Fluctuations of fish populations and the magnifying
700   effects of fishing. *Proc Natl Acad Sci* 108(17):7075-7080.

701   70.   Bjørnstad ON, Nisbet RM, Fromentin J (2004) Trends and cohort resonant effects in age-
702   structured populations. *J Anim Ecol* 73(6):1157-1167.

703   71.   Steneck R*, et al.* (2011) Creation of a gilded trap by the high economic value of the
704   Maine lobster fishery. *Conserv Biol* 25(5):904-912.

705   72.   Hill MF, Hastings A, Botsford LW (2002) The effects of small dispersal rates on
706   extinction times in structured metapopulation models. *Am Nat* 160(3):389-402.

707   73.   Secor DH, Kerr LA, Cadrin SX (2009) Connectivity effects on productivity, stability, and
708   persistence in a herring metapopulation model. *ICES J Mar Sci* 66(8):1726-1732.

709   74.   Fox JW, Vasseur D, Cotroneo M, Guan L, Simon F (2017) Population extinctions can
710   increase metapopulation persistence. *Nature Ecol Evol* 1(9):1271.

711   75.   Nesbitt HK, Moore JW (2016) Species and population diversity in Pacific salmon
712   fisheries underpin indigenous food security. *J Appl Ecol* 53(5):1489-1499.

713   76.   Brown F, Brown YK (2009) *Staying the course, staying alive: coastal First Nations*
714   *fundamental truths: biodiversity, stewardship and sustainability* (Biodiversity Bc
715   Victoria).

716   77.   Gavreau A, Lepofsky D, Rutherford M, Reid M (2017) "Everything revolves around the
717   herring": The Heiltsuk-herring relationship through time. *Ecol Soc* 22(2).

78. Mackinson S, Sumaila UR, Pitcher TJ (1997) Bioeconomics and catchability: fish and fishers behaviour during stock collapse. *Fish Res* 31(1):11-17.

79. Elkington J (1994) Towards the sustainable corporation: Win-win-win business strategies for sustainable development. *Calif Manag Rev* 36(2):90-100.

80. Essington TE*, et al.* (2015) Fishing amplifies forage fish population collapses. *Proc Natl Acad Sci* 112(21):6648-6652.

81. Borgström S, Elmqvist T, Angelstam P, Alfsen-Norodom C (2006) Scale mismatches in management of urban landscapes. *Ecol Soc* 11(2).

82. Berkes F (2006) From community-based resource management to complex systems: The scale issue and marine commons. *Ecol Soc* 11(1).

83. Biggs R*, et al.* (2012) Toward Principles for Enhancing the Resilience of Ecosystem Services. *Annu Rev Environ Resour* 37:421-448.

84. Folke C, Pritchard L, Berkes F, Colding J, Svedin U (2007) The problem of fit between ecosystems and institutions: Ten years later. *Ecol Soc* 12(1).

85. Young OR (2002) *The institutional dimensions of environmental change: fit, interplay, and scale* (MIT press).

86. Acheson JM (2003) *Capturing the commons: Devising institutions to manage the maine lobtser industry* (University Press New England, Lebanon, NH).

87. Thornton TF, Kitka H (2015) An indigenous model of a contested Pacific Herring fishery in Sitka, Alaska. *Int J Appl Geospatial Res* 6(1):94-117.

88. Halpern BS*, et al.* (2013) Achieving the triple bottom line in the face of inherent trade-offs among social equity, economic return, and conservation. *Proc Natl Acad Sci* 110(15):6229-6234.

89. Law EA*, et al.* (2017) Equity trade-offs in conservation decision making. *Conserv Biol.*

90. Ricker W (1958) Maximum sustained yields from fluctuating environments and mixed stocks. *Journal of the Fisheries Board of Canada* 15(5):991-1006.

91. Hay D (1985) Reproductive biology of Pacific herring (*Clupea harengus pallasi*). *Can J Fish Aquat Sci* 42(S1):s111-s126.

92. Carpenter B*, et al.* (2016) Stan: A probabilistic programming language. *J Stat Softw* 20.

93. Lewandowski D, Kurowicka D, Joe H (2009) Generating random correlation matrices based on vines and extended onion method. *J Multivar Anal* 100(9):1989-2001.

94. Hoffman MD, Gelman A (2014) The No-U-Turn Sampler: Adaptively Setting Path Lengths in Hamiltonian Monte Carlo. *J Mach Learn Res* 15:1593-1623.

95. Stan Development Team (2016) Stan: A C++ library for probability and sampling. Version 2.12.0.).

96. Stan Development Team (2016) RStan: the R interface to Stan. Version 2.10.1.).

97. R Core Team (2017) R: A language and environment for statistical computing (R Foundation for Statistical Computing, Vienna, Austria).

756 98. Tanasichuk R, Kristofferson A, Gillman D (1993) Comparison of some life history
757   characteristics of Pacific herring (*Clupea pallasi*) from the Canadian Pacific Ocean and
758   Beaufort Sea. *Can J Fish Aquat Sci* 50(5):964-971.

759 99. Lütkepohl H (2005) *New introduction to multiple time series analysis* (Springer, Berlin).

760 100. Nisbet RM, Gurney W (1982) *Modelling fluctuating populations* (Wiley, New York).

Table 1: Home-range categories of different Pacific herring predator groups. Details of sources and home-ranges are provided in the SI Appendix.

Home-Range Category	Home-Range Description	Herring Predator Groups
Localized	single local substock *(< 10 km radius)*	Indigenous fishers**, Seabirds*, Flatfishes*, Crustaceans*, Urchins, Cucumbers, Reef Fishes, Rockfishes
Centralized	multiple substocks *(~10-40 km-radius)*	Indigenous fishers**, Seabirds*, Flatfishes*, Crustaceans*, Gadiforms*, Salmonids**, Halibut**, Harbor Seals
Transient	full stock or greater	Seine & Gillnet Fishers, Cetaceans, Sea Lions, Seabirds*, Gadiforms*, Salmonids**, Halibut**

*Varies by species, ** Varies by migration phase, location or individual

Figure 1: **(a)** Map of major Pacific Herring spawning substocks ("Sections" - denoted by different symbols) as defined by Canada's Federal Fisheries Agency in the Central Coast of British Columbia. Indigenous communities noted with *. **(b)** Estimated trends for the six main substocks (thin colored lines & points) and the metapopulation (stock) mean (thick blue line). Trends were estimated using multivariate hierarchical Bayesian state-space models integrating spawn surveys and catch information. **(c)** Temporal coefficients of variation over 25 years for each substock (points and errorbars), the mean of each substock CV (large black symbol), the regional stock mean (coefficient of variation, blue line), and the spatial variability (black point with error bar) which is the standardized variance of the system after accounting for the variance of the stock and the variance of the substock means. Individual plots for each substock area are shown, along with catch, spawn observations, and 95% posterior credible intervals in SI Appendix A. **(d)** Trends in annual harvest rate for 2 of the 6 local substocks (Higgins and Lower Spiller) with points, 95% credibility intervals and the mean for the aggregate stock (thick blue line). The dotted line represents the target harvest rate of 20%.

Figure 2: Results of simulation-based risk analysis of hypothetical Pacific herring fisheries under different management and spatial scenarios. Heatmaps show risk of collapse at metapopulation and population scales against annual migration and spatial synchrony in recruitment productivity for different harvest rates and spatial harvest allocation strategies. Risk of collapse is the probability of falling below 20% of unfished equilibrium (i.e. the dotted lines in Fig 3 a-c); at the subpopulation scale, risk is measured as the mean probability of collapse for each population. Optimized spatial allocation removes proportionally less biomass from subpopulations with lower biomass; proportional allocation removes the same proportion of biomass from each subpopulation. Note that differences in mean annual stock level catch among optimized versus proportional allocation are minimal (2c vs. 2e and 2g vs. 2i; SI Appendix B). Annual migration is the mean percent of each subpopulation that emigrates each year. Spatial synchrony in recruitment productivity is $1 + (n - 1)\bar{r}$ where $\bar{r}$ is the mean of all unique pairwise correlations in subpopulation recruitment deviations and n is the number of populations. Heatmaps for 30% harvest rules are not shown here as they produce uniformly high risk at both scales. For full results see SI Appendix B.

Figure 3: (a-c) Example simulations showing spatial variation in biomass measured at the subpopulation level (colored lines) or metapopulation mean (thick blue) for **(a)** an unexploited metapopulation, **(b)** a 20% target harvest allocated proportional to spawning biomass in space, or **(c)** a 20% target harvest allocated optimally in space according to the ideal free distribution. B0 = deterministic biomass in the absence of fishing, dotted line = lower conservation threshold of 20% of B0. **(d)** change in spatial variation for different harvest rates and spatial harvest allocation across different spatial parameter scenarios for migration rates and spatial synchrony. Each point represents the spatial variance from 100 runs from a single 50-year simulation using a unique migration-synchrony parameter combination.

Figure 4: Effect of mortality rate on asynchrony related spatial variability in the metapopulation from the simple two subpopulation model (used for analytical tractability). **(a)** Solutions for spatial variation among populations as a function of total mortality for different levels of spatial synchrony in recruitment productivity when annual migration is 0.25 and log-scale recruitment standard deviation (σ_R) is 0.7 (i.e. CV = 0.8); **(b)** Temporal variability (squared coefficient of variation) of the population and metapopulation. The arrows illustrate spatial variability as the difference between surfaces. **(c)** Measures of local population predictability (measured by first order within population autocorrelation, thick black line), among population coupling (measured by among population correlations, thin black line), and local environmental sensitivity (measured by the response at the population level to a unit environmental impulse affecting recruitment productivity, red line).

Supplementary Information - Appendices for "Spatial variation in exploited metapopulations obscures risk of collapse"

Daniel K Okamoto[a,b,c], Margot Hessing-Lewis[b], Jameal F Samhouri[d], Andrew O Shelton[d], Adrian Stier[e], Philip S Levin[f,g], and Anne K Salomon[b,c]

[a]Department of Biological Science, Florida State University, Tallahassee, Florida 32303, USA
[b]Hakai Institute, PO Box 309, Heriot Bay, BC V0P 1H0, Canada
[c]School of Resource and Environmental Management, Simon Fraser University, 8888 University Drive, Burnaby, B.C. V5A 1S6, Canada
[d]Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Blvd. East,Seattle, WA 98112, USA
[e]Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, CA 93106, USA
[f]The Nature Conservancy, 74 Wall St, Seattle, WA98121, USA
[g]School of Environment and Forestry Sciences, University of Washington, Seattle, Washington, USA

1 Substock trends in Central Coast herring

2 *Residual spatial variability*

3   The residual spatial variation (RSV, also called β variation or the spatial variation due
4 to asynchrony) is the difference between the variance of the mean metapopulation trend and
5 the expected metapopulation mean variance if populations are perfectly correlated. In other
6 words, residual spatial variability is the additional temporal variability at the population
7 level left unexplained by the variability of the metapopulation mean.

    While the actual variability of a metapopulation mean given individual variability and
covariance of each of N individual population time series (X_i):

$$\text{Var}\left(\frac{1}{N}\sum_{i=1}^{N}X_i\right) = \frac{1}{N^2}\left[\sum_{i=1}^{N}\text{Var}\left(X_i\right) + \sum_{i=1}^{N}\sum_{j\neq 1}^{N-1}\rho_{i,j}\sqrt{\text{Var}\left(X_i\right)\text{Var}\left(X_j\right)}\right] \tag{S.1}$$

$$= \frac{1}{N^2}\left[\sum_{i=1}^{N}\text{Var}\left(X_i\right) + \sum_{i=1}^{N}\sum_{j\neq 1}^{N-1}\text{Cov}\left(X_i, X_j\right)\right] \tag{S.2}$$

$$\tag{S.3}$$

8   The maximum potential variance of a metapopulation mean given individual variance of
9 each of N populations is:

$$\text{Var}_{\text{max}}\left(\bar{X}\right) = \frac{1}{N^2}\left[\sum_{i=1}^{N}\text{Var}\left(X_i\right) + \sum_{i=1}^{N}\sum_{j\neq 1}^{N-1}\sqrt{\text{Var}\left(X_i\right)\text{Var}\left(X_j\right)}\right] = \left(\sum_{i=1}^{N}\frac{\sqrt{\text{Var}\left(X_i\right)}}{N}\right)^2 \tag{S.4}$$

10   Hence, the residual spatial variance (RSV) is:

$$\text{RSV}\left(\bar{X}\right) = \text{Var}_{\text{max}}\left(\bar{X}\right) - \text{Var}\left(\bar{X}\right) \tag{S.5}$$

$$= \frac{1}{N^2}\left[\sum_{i=1}^{N}\sum_{j\neq 1}^{N-1}\sqrt{\text{Var}\left(X_i\right)\text{Var}\left(X_j\right)} - \sum_{i=1}^{N}\sum_{j\neq 1}^{N-1}\text{Cov}\left(X_i, X_j\right)\right] \tag{S.6}$$

$$= \frac{1}{N^2}\left[\sum_{i=1}^{N}\sum_{j\neq 1}^{N-1}\left(1 - \rho_{i,j}\right)\sqrt{\text{Var}\left(X_i\right)\text{Var}\left(X_j\right)}\right] \tag{S.7}$$

$$= \left(\sum_{i=1}^{N}\frac{\sqrt{\text{Var}\left(X_i\right)}}{N}\right)^2 - \text{Var}\left(\bar{X}\right) \tag{S.8}$$

11   Similarly, population synchrony as defined by X is given by:

$$Synchrony = \frac{\text{Var}\left(\bar{X}\right)}{\text{Var}\left(\bar{X}\right) + \text{RSV}\left(\bar{X}\right)} \tag{S.9}$$

12 *Full time series results*

Figure S.1: Trends in pre-spawning biomass (spawn plus catch) and associated catch for the six main spawning locations of Pacific herring the Central Coast of British Columbia. Open points represent the observed pre-spawn biomass, black lines represent posterior estimate of biomass from the model, blue lines represent the overall stock mean, and grey bands are the 95% credible interval of estimated biomass.

Figure S.2: (a-f) Trends in annual harvest rate for 2 of the 6 local substocks (points and 95% credibility intervals) and the mean for the aggregate stock (thick blue line). The dotted line represents the target harvest rate of 20%. (g-j) Harvest rate and 95% credible intervals for each section in four select years compared to the optimized strategy via the ideal free distribution of harvest (IFD, grey band) or stock allocation with uniform proportional harvest (solid blue line). Points above both the IFD and the equal stock allocation may represent a suboptimal harvest strategy. In contrast, points along the IFD line, such as that in 1999, represent a strategy may reduce local risks. (k) Spatial harvest inequity (measured as the absolute mean relative error from perfect equity). Note that in 1999 equity differs depending on which benchmark is used (IFD or uniform proportional harvest).

13 Spatial allocation of harvest

14   The fishing fleet either distributes itself in space or a manager forces spatial allocation. We
15 use three scenarios for spatial allocation: 1) diffuse effort in space (i.e harvest is proportional
16 to abundance), 2) ideal distribution of effort in space, and 3) random spatial allocation.

17 Proportional allocation of harvest

For a proportional allocation of harvest, the calculation of biomass caught in each location
($H_{l,t}$) is simply:

$$H_{\text{adults}_{l,t}} = \frac{\sum_{a=2}^{10} w_a m_a n_{a,l,t}}{\sum_{l=1}^{L} \sum_{a=2}^{10} w_a m_a n_{a,l,t}} \times \text{Quota}_t \tag{S.10}$$

18 Ideal free distribution

For the Ideal Free Distribution of Effort in Space, the fishing fleet would ideally allocate
themselves to optimize efficiency of catch (i.e. minimal effort for the return). We can
numerically solve for the spatial harvest strategy that optimizes efficiency using a system
of ordinary differential equations (ODEs). First, we 1) simplify the fishery to a very short
time period, 2) assume vessels can move instantaneously among locations with no cost, 3)
assume vessels have perfect information and 4) assume fishing effort (i.e. sets) can be broken
into infinitesimally small units. Consider $F_{l,\tau}$ the rate of fishing effort (i.e. number of sets
per unit time) allocated at time τ in location l, q is the catchability of fish (i.e. ratio of
proportional capture rate to effort), B_τ is the biomass available for capture at time τ D_e
is the rate at which vessels respond to spatial disparities in efficiency, where efficiency =
effort x catchability x biomass. To maximize efficiency, vessels sense disparities in efficiency
and move instantaneously to neighboring locations with higher efficiency (i.e. catch per unit
time per unit effort).

$$\frac{dB_l}{d\tau} = -F_l(\tau)qB_l(\tau) \tag{S.11a}$$

$$\frac{dF_l}{d\tau} = \sum_{j=1}^{L} [F_l(\tau)qB_l(\tau) - F_j(\tau)qB_j(\tau)] \tag{S.11b}$$

$$\frac{dH_l}{d\tau} = F_l(\tau)qB_l(\tau) \tag{S.11c}$$

$$H(\tau_Q, l) = \int_0^{\tau_Q} F_l(\tau)qB_l(\tau)d\tau \tag{S.12a}$$

$$\text{StockH}(\tau_Q) = \sum_{l=1}^{L} H(\tau_Q, l) \tag{S.12b}$$

32 We numerically evaluate this system of ODEs until the quota is achieved to produce the
33 final, optimal distribution of catch in space.

34 Random spatial allocation distribution

For random distribution of the harvest, the proportion of the quota removed from each
location within each year is a random draw from a Dirichlet distribution, with mean equal
to the actual proportional biomass distribution.

random proportional harvest rates are assigned to each site, with a maximum proportion
of biomass removed at any given spawning site.the fleet randomly chooses a site, prosecutes a
fishery, and moves on until the quota is achieved. Functionally, the fleet effort is distributed
randomly in space, with mean allocation probabilities given by the distribution of biomass
in space. Potential, proportional harvest rates are randomly assigned to each spawning area
(though cannot exceed the maximum) and prosecuted in order from highest to lowest until
the quota is achieved. The initial random proportional harvest rates are set by a Dirichlet
distribution where means are proportional to the distribution of stock biomass in space and a
count parameter (K) that sets the degree of randomness (set as $K = 5$) to achieve a spatial
harvest inequity of approximately 0.20. Lower count parameters defy the proportionality
more (i.e. harvests are unlikley to reflect biomass distributions in space) while very high
count parameters approximate harvests that are porportional to biomass distributions in
space. An example of random harvest allocation is given by Figure S.3.

Figure S.3: Examples of implementation of random harvest allocation for individual years
in the simulation model for a target harvest of 20%. Each panel represents one year with
10 substocks. Vertical bars with points represent substock harvest rates compared to the
relative pre-spawn (and pre-harvest) biomass in that substock.

Figure S.4: Discrepancy in risk among spatial scales (mean risk at subpopulation scale minus mean risk at metapopulation scale) versus spatial a variance by harvest rate and spatial harvest allocation strategy

.

Figure S.5: Relative frequency of the % difference in catch between proportional allocation and ideal free distribution of harvest, calculated for each individual simulation scenario as $100 * \left(\overline{catch}_{PA} - \overline{catch}_{IDF} \right) / \overline{catch}_{PA}$
.

Figure S.6: Simulation results including target harvest rates ranging from 0 to 20% for each spatial allocation scenario and measured at the metapopulation (stock) and population (substock) scales. 30% harvest rate not shown due to all risk panels above 10%

51 **Statistical properties of the VAR(1) model**

52   The long-term temporal variability of the individual populations and the metapopulation
53 is determined by the long-run covariance matrix:

$$\text{vec}(\boldsymbol{\Sigma}_x) = [\boldsymbol{I}_{p^2} - \boldsymbol{J} \otimes \boldsymbol{J}]^{-1} \text{vec}(\boldsymbol{\Sigma}_\zeta) \tag{S.13}$$

Where p is the total number of variables (i.e. reproducers, recruits, and adult stages in both
populations), $\boldsymbol{I}_{p^2}$ is a $p^2 \times p^2$ identity matrix, $\otimes$ is the Kronecker product, and vec is the
vector operator (i.e. turning a matrix into a vector for operational sake). The population
and metapopulation coefficient of variation (on the natural scale), as well as the among
population spatial variation and spatial correlations the can be derived from $\boldsymbol{\Sigma}_x$ using the
definition of the mean, variance and covariance of a multivariate lognormal given the mean
and covariance matrix of the log-scale system (see appendix for derivations).

If we assume the system conforms to a multivariate lognormal, the temporal variance of
spawners at site i is:

$$\text{Var}(e^{[x_{i,a_0}]}) = e^{[2\bar{x}_{i,a_0}+\sigma_{i,i}^2]}e^{[\sigma_{i,i}^2-1]} = e^{[\sigma_{i,i}^2]}e^{[\sigma_{i,i}^2-1]} \tag{S.14}$$

63 where $\sigma_{i,i}^2$ is the diagonal element of the log-scale variance covariance matrix $\boldsymbol{\Sigma}_x$ associated
64 with spawners at site i (x_{i,a_0}), and $\bar{x}_{i,a_0}$ is the mean of x_{i,a_0} which is set to zero. From the
65 lognormal, the mean of $e^{[x_{i,a_0}]}$ is $e^{[0.5\sigma_{i,i}^2]}$.

66   The resulting coefficient of variation of each spawning population is:

$$\text{CV}(e^{[x_{i,a_0}]}) = \frac{\sqrt{e^{[\sigma_{i,i}^2]}e^{[\sigma_{i,i}^2-1]}}}{e^{[0.5\sigma_{i,i}^2]}} = \sqrt{e^{[\sigma_{i,i}^2-1]}} \tag{S.15}$$

67   The temporal covariance of scaled spawners among sites is:

$$\text{Cov}(e^{[x_{i,a_0}]}, e^{[x_{j,a_0}]}) = e^{[0.5\sigma_{i,i}^2+0.5\sigma_{j,j}^2]}e^{[\sigma_{i,j}^2-1]} \tag{S.16}$$

68 where $\sigma_{i,j}^2$ is the off-diagonal (covariance) element of $\boldsymbol{\Sigma}_x$.

69   Considering a metapopulation with two component populations, we set the log-
70 population variances equal (i.e. $\sigma_{i,i} = \sigma_{j,j} = \sigma$), the resulting variance of the spatial mean
71 on the natural scale is:

$$\text{Var}\left[\frac{e^{[x_{i,a_0}]}+e^{[x_{j,a_0}]}}{2}\right] = \text{Var}\left(\frac{e^{[x_{i,a_0}]}}{2}\right) + \text{Var}\left(\frac{e^{[x_{j,a_0}]}}{2}\right) + 2\text{Cov}\left(\frac{e^{[x_{i,a_0}]}}{2}, \frac{e^{[x_{j,a_0}]}}{2}\right) \tag{S.17}$$

$$= 0.5e^{[\sigma^2]}e^{[\sigma^2-1]} + 0.5e^{[\sigma^2]}e^{[\sigma_{i,j}^2-1]} \tag{S.18}$$

72   Thus, the temporal coefficient of variation of the spatial mean is:

$$\text{CV}\left[\frac{e^{[x_{i,a_0}]}+e^{[x_{j,a_0}]}}{2}\right] = \frac{\sqrt{0.5e^{[\sigma^2]}e^{[\sigma^2-1]} + 0.5e^{[\sigma^2]}e^{[\sigma_{i,j}^2-1]}}}{e^{[0.5\sigma^2]}} = \sqrt{0.5e^{[\sigma^2-1]} + 0.5e^{[\sigma_{i,j}^2-1]}} \tag{S.19}$$

73   In this case, the residual spatial variance (taken from S.7) is:

$$\text{residual spatial variance} = 0.5e^{[\sigma^2]}e^{[\sigma^2-1]} - 0.5e^{[\sigma^2]}e^{[\sigma_{i,j}^2-1]} \tag{S.20}$$

74
75 Which when standardized by the square of the population mean ($e^{[0.5\sigma^2]}$) yields:

$$\text{standardized residual spatial variance} = 0.5e^{[\sigma^2-1]} - 0.5e^{[\sigma_{i,j}^2-1]} \tag{S.21}$$

76 In this case this is the same as the variance of the difference between either population and
77 the population mean trend (because both populations are identical).

$$\text{Var}\left[e^{[x_{i,a_0}]} - \left(\frac{e^{[x_{i,a_0}]} + e^{[x_{j,a_0}]}}{2}\right)\right] = \text{Var}\left[\frac{e^{[x_{i,a_0}]}}{2} - \frac{e^{[x_{j,a_0}]}}{2}\right] \tag{S.22}$$

$$= \text{Var}\left[\frac{e^{[x_{i,a_0}]}}{2}\right] + \text{Var}\left[\frac{e^{[x_{j,a_0}]}}{2}\right] - 2\text{Cov}\left(\frac{e^{[x_{i,a_0}]}}{2}, \frac{e^{[x_{j,a_0}]}}{2}\right) \tag{S.23}$$

$$= 0.5e^{[\sigma^2]}e^{[\sigma^2-1]} - 0.5e^{[\sigma^2]}e^{[\sigma_{i,j}^2-1]} \tag{S.24}$$

As a result, we can combine the solutions from C.13 and C.21 to generate the residual
spatial variation for the metapopulation and how it is influenced by mortality and environ-
mental correlations.

Autocorrelations

The matrix of first-order autocorrelations that yields within population correlation and
spatial coupling is given by:

$$\mathbf{R} = (\text{diag}(\mathbf{\Sigma}_x)^{\circ\frac{1}{2}})^{-1}\mathbf{\Sigma}_x(\text{diag}(\mathbf{\Sigma}_x)^{\circ\frac{1}{2}})^{-1} \tag{S.25}$$

Impulse response of the VAR(1) model

The impulse response (IR) of adults at location i after a lag t to a single pulse log-scale
recruitment deviation is the first coefficient of the nth column of the moving average (MA)
representation of the VAR model. Specifically, the matrix of MA coefficients for the VAR(1)
at lag t is produced by $\mathbf{J}^t$. In this case, the IR for adults is the first entry of the corresponding
MA vector. The greatest IR of adults is at lag 1 (when a recruitment pulse first enters into
the adult stage) and is thus conveniently represented by the first entry of the jacobian matrix
$\mathbf{J}$. Because this represents the log-scale additive response, the IR on the natural scale is a
multiplicative effect.