Title: Observed Antibody Space: a resource for data mining next generation sequencing antibody

repertoires.

Aleksandr Kovaltsuk[1], Jinwoo Leem[1], Sebastian Kelm[2], James Snowden[2], Charlotte M. Deane[1,*], Konrad

Krawczyk[1,*]

1.   University of Oxford, Department of Statistics, Oxford, UK

2.   UCB Pharma, Slough, UK

* to whom the correspondence should be addressed,

deane@stats.ox.ac.uk

konrad@proteincontact.org

Abstract. Antibodies are immune system proteins that recognize noxious molecules for elimination.

Their sequence diversity and binding versatility have made antibodies the primary class of

biopharmaceuticals. Recently it has become possible to query their immense natural diversity using

next-generation sequencing of immunoglobulin gene repertoires (Ig-seq). However, Ig-seq outputs are

currently fragmented across repositories and tend to be presented as raw nucleotide reads, which

means nontrivial effort is required to reuse the data for analysis. To address this issue, we have

collected Ig-seq outputs from 53 studies, covering more than half a billion antibody sequences across

diverse immune states, organisms and individuals. We have sorted, cleaned, annotated, translated and

numbered these sequences and make the data available via our Observed Antibody Space (OAS)

resource at antibodymap.org. The data within OAS will be regularly updated with newly released Ig-seq

datasets. We believe OAS will facilitate data mining of immune repertoires for improved understanding

of the immune system and development of better biotherapeutics.

1. Introduction

Antibodies (or B-cell receptors) are protein products of B-cells and primary actors of adaptive immunity

in jawed vertebrates [1]. They are highly malleable molecules that can bind to virtually any antigen. An

organism holds a great variety of these molecules increasing the probability that an arbitrary antigen

can be recognized by an antibody, initiating an immune response [2]. Owing to their binding malleability

they are the most prominent class of reagents and biotherapeutics [3,4]. Continued successful exploitation

of these molecules relies on our ability to discern the functional diversity of antibody repertoires [5–7].

Next-generation sequencing of immunoglobulin gene repertoires (Ig-seq) has enabled researchers to

take snapshots of millions of sequences at a time across individuals, diverse organisms and different

immune states [8,9]. The ability to analyze millions of antibody sequences has the potential to uncover the

mechanics of the immune response to any antigen [10,11] and dysfunctions of the immune system itself [12].

Many previous studies have addressed the issue of antibody diversity, contributing invaluable evidence

to the dynamics of human immune systems [13]. Numerous analyses have focused on the frequencies of

V(D)J gene usages, which can offer insights into creating biased antibody libraries for therapeutic

applications [14–16]. Another therapeutic application of antibody repertoire analysis is advancing vaccine

design by comparative longitudinal studies of pre- and post-antigen challenge experiments [10,11,17–22].

Such comparative studies have shown that different individuals can converge on the same antibody

sequence against a given vaccine [11,19]. Due to sequencing limitations, these analyses have focused on

heavy or light chains separately, whereas one ought to study the paired repertoire to obtain a deeper

insight of the antibody diversity [23].

Technical advances in sequencing technology have outpaced storage and analysis pipelines [24,25]. This has

meant that the outputs of Ig-seq studies are fragmented across repositories making it difficult to

perform large-scale data mining of antibody repertoires [25]. Metadata such as isotype, age or subject

identifiers are not typically standardized, therefore extraction of specific subsets of antibody repertoires

for comparative analyses is challenging. Furthermore, the data are typically deposited as raw nucleotide

reads. It requires non-trivial *ad hoc* effort to convert such raw reads to amino acid sequences that

ultimately dictate the molecular structure and antigen-recognition. Some of these issues are addressed

by services that provide Ig-seq-specific data deposition and analysis pipelines such as Immport

(http://immport.org) [26,27], or ImmunoseqAnalyzer (http://clients.adaptivebiotech.com/), IReceptor

(http://ireceptor.irmacs.sfu.ca/) or VDJServer (http://vdjserver.org) [28]. The IReceptor and the VDJServer

are the main resources that fall under the umbrella of the organized effort by the Adaptive Immune

Receptor Repertoire (AIRR) Community to provide standardized deposition and analysis pipelines for the

Ig-seq outputs [24]. These services chiefly focus on facilitating bulk deposition of raw data to perform

standardized sequencing analyses. Ultimately, because immunoinformatics is not the chief focus of such

services, bulk data download from such websites is limited and converting the raw nucleotide data

obtained into format suitable for analysis still requires non-trivial effort.

To address these issues, we have created the Observed Antibody Space (OAS) resource that allows large-

scale data mining of antibody repertoires. We have collected the raw outputs of 53 Ig-seq experiments

covering over half a billion sequences. We have organized the sequences by metadata such as organism,

isotype, B-cell type and source, and the immune status of B-cell donors to facilitate bulk retrieval of

specific subsets for comparative analyses. We have converted all of the Ig-seq sequences to amino acids

and numbered them using the IMGT scheme. The data is available for querying or bulk download at

http://antibodymap.org. We believe that OAS will facilitate data mining antibody repertoires for

improved understanding of the dynamics of the immune system and thus better engineering of

biotherapeutics.

2. Materials and Methods

A list of study accession codes of publically available Ig-seq datasets were obtained via a literature

review. The majority of raw reads were downloaded from the European Nucleotide Archive (ENA) [29] and

the National Center for Biotechnology Information (NCBI) websites [30]. In a small number of cases,

another public Ig-seq repository was specified e.g. [14,31–33]. Metadata were manually extracted from the

deposited datasets and arranged in a reproducible format.

The downloaded FASTQ files were processed depending on the sequencing platform. Paired raw

Illumina reads were assembled with FLASH [34]. The assembled antibody sequences were converted to the

FASTA format using FASTX-toolkit [35]. As raw reads from Roche 454 are not paired, these FASTQ files

were directly converted to the FASTA format with the FASTX-toolkit.

The heavy chain sequences were automatically annotated with isotype information unless such data was

given in the corresponding publication. Automatic isotype annotation was performed by aligning the

constant heavy domain 1 (CH1) of any given antibody sequence against the IMGT isotype reference [36] of

the respective species using the Smith-Waterman algorithm [37]. We assigned score of two for a

nucleotide match, and a score of minus one for a nucleotide mismatch or a gap in our Smith-Waterman

scoring function. The IMGT isotype references comprised 21 nucleotide-long fragments of the CH1

domain of the antibody isotypes. To ensure a high confidence of correct isotype identification, we

employed a conservative threshold of 30 in the Smith-Waterman algorithm scoring function. Sequences

whose Smith-Waterman algorithm score was below the threshold for all isotypes were assigned as

'bulk'. The robustness of this protocol was confirmed on the author-annotated Ig-seq datasets [38–40]

where it resulted in 99% accurate annotations. Around 1% of the Ig-seq data had a very short (or

missing) sequence of CH1 domain. Such sequences were also assigned as 'bulk'.

IgBlastn [41] was used to convert the FASTA files of antibody nucleotide sequences to amino acids. The

amino acid sequences were then numbered with ANARCI [42] using the IMGT scheme [43]. ANARCI does not

number a sequence if it does not align to a suitable Hidden Markov Model [44]. ANARCI therefore ensures

that antibody sequences do not harbor unusual indels or stop codons in the antibody regions, that V and

J genes align to respective species amino acid IMGT germlines [36], and that the length of CDR-H3 is not

greater than 37 residues in human, mouse, rat, rabbit, alpaca and rhesus antibodies. Due to technical

limitations of sequencing platforms, certain reads were missing significant portions of the variable

region (e.g. portions of CDR1), sequences that did not have all three CDRs were discarded as incomplete.

The V and J genes are identified during the ANARCI numbering step.

Using the protocol above we annotated Ig-seq results of 53 independent studies. In order to streamline

updating OAS with new data, we have generated a procedure to automatically identify Ig-seq datasets

from raw sequence read archives. We apply our antibody annotation protocol to each raw nucleotide

dataset deposited in the NCBI/ENA repositories, if we find more than 10,000 antibody sequences in any

given dataset, it is set aside for manual inspection. It is still necessary to manually inspect raw datasets

to efficiently assign metadata as these are currently deposited in a non-standardized manner. This

procedure allows for automatic identification of new Ig-seq datasets, which will enable us to semi

automatically update OAS.

3. Results

We collected raw sequencing outputs from 53 Ig-seq studies. All raw nucleotide reads were converted

into amino acids using IgBlastn [41]. The full amino acid sequences were then IMGT numbered using

ANARCI [42]. As well as providing IMGT and gene annotations, ANARCI acts as a broad-brush filter of

antibody sequences that are likely to be erroneous (see Materials and Methods). Applying the same

retrieval, amino acid conversion, gene annotation and numbering protocol to all sequences assures the

same point of reference across the 53 heterogeneous Ig-seq datasets [45]. This protocol produces the full

IMGT-numbered sequences together with gene annotations for each of the 53 datasets.

Figure 1. T**he Observed Antibody Space database**. The data from 53 studies is sorted into Data

Units. Each Data Unit is a set of antibody sequences that share the same set of meta-data. Each

sequence in a Data Unit is further associated with sequence-specific annotations.

The numbered amino acid sequences in each dataset are sorted by metadata e.g. individuals, age,

vaccination regime, B-cell type and source *etc*. (Figure 1). Deposition of such metadata is currently not

standardized and requires *ad hoc* manual curation for each dataset. In an effort to organize the antibody

sequences using such metadata, we have grouped the sequences within each dataset into Data Units.

Each Data Unit represents a group of sequences within a given dataset with a unique combination of

metadata values. The metadata values are summarized in Table 1.

Metadata name	Metadata description
Chain	Heavy/light chain annotation.
Isotype	Identified or deposited isotype information.
Age	Information on the age of the human B-cell donors.
Disease	Indication of whether the donor was sick at the time of B-cell extraction.
Vaccine	Indication if the B-cell donor was purposely immunized prior to B-cell extraction.
B-cell subset	Indication if a particular B-cell subset was sorted for Ig-seq
Species	Organism of the B-cell donor
B-cell source	Which organ/tissue the B-cells were extracted from.
Subject	Indication of a particular B-cell donor the B-cells were sourced from.
Longitudinal	If the study was longitudinal, an indicator of the time point.
Size	Number of non-redundant sequences in the dataset.
Link	Link to the source publication.

145   <u>Table 1.</u> Metadata descriptors of each Data Unit in OAS. Each Data Unit is uniquely identified by the

146   study and a collection of the metadata values.

As of April 29th 2018, 53 Ig-seq studies are included in OAS totaling 608,651,423 sequences (552,824,460

VH and 55,826,963 VL sequences). The majority of these sequences are murine (~50.4%) and human

(~47.4%). Twenty-two of the Ig-seq studies interrogate the immune system of diseased individuals, the

most common ailment being HIV (13 studies). The database also contains 22 Ig-seq studies of the naive

antibody gene repertoires (the collection of B-cells from donors who are healthy and not purposefully

vaccinated). The main source of B-cells in the OAS database is peripheral blood (~231m of sequences)

followed by spleen/splenocytes (~198m) and bone marrow (~124m). The database holds isotype

information for each individual heavy sequence and the two most common isotypes are IgM (~312m)

and IgG (~139m). For ~65m sequences we were not able to assign isotypes with high confidence. The

median total redundant size of the Ig-seq studies in the OSA database is 2,006,196 sequences, while the

largest Ig-seq study was that by Greiff et al., (246,449,189 redundant sequences) [14]. Detailed statistics on

each dataset are given in Table 2. All the data may be bulk downloaded or individual Data Units queried,

at http://antibodymap.org.

Study	Species	Disease	Vaccine	B-cell source	B-cell subset	Total ANARCI parsed sequences
Banerjee et al., [46]	Rabbit	None	HIV	PBMC	Unsorted	4,334,088 (2,926,727)
Bashford-Rogers et al., [47]	Human	CLL/None	None	PBMC	Unsorted	129,013 (86,166)
Bhiman et al., [48]	Human	HIV	None	PBMC	Unsorted	785,751 (187,067)
Bonsignori et al., [49]	Human	HIV/None	None	PBMC	Memory/ Unsorted	210,377 (57,374)
Collins et al., [50]	Mouse	None	None	Splenocytes	Unsorted	812,439 (194,752)
Corcoran et al., [51]	Human/ Mouse/ Rhesus	None	None	PBMC	Unsorted	5,307,880 (2,840,877)
Cui et al., [52]	Mouse	None	NP-CGG/None	Splenocytes	Memory	5,513,816 (935,646)
Doria-Rose et al., [13]	Human	HIV	None	PBMC	Unsorted	2,164,901 (549,544)
Fisher et al., [53]	Mouse	None	Plasmodium	Spleen	Unsorted	175,015 (113,594)
Galson et al., [38]	Human	None	Hepatitis-B	PBMC	Unsorted/ Plasma cells/ HepB-specific	21,755,739 (10,442,291)
Galson et al., [39]	Human	None	Hepatitis-B	PBMC	Unsorted/ Plasma cells/ HepB-specific	26,687,394 (14,343,236)
Galson et al., [21]	Human	None	Meningitis	PBMC	Naïve/ Plasma cells/ Memory/ Marginal zone	7,918,197 (3,282,907)
Galson et al., [17]	Human	None	Flu	PBMC	Plasma cells	13,685,210 (5,065,786)
Greiff et al., [40]	Mouse	None	NP-CGG	Bone marrow/ Spleen	Plasma cells/ Plasmablasts	7,955,739 (2,891,649)
Greiff et al., [33]	Mouse	None	NP-CGG	Spleen	ASCs/Plasma cells/ Naive	788,787 (523,716)
Greiff et al., [14]	Mouse	None	OVA/Hepatitis-B/ NP-HEL/None	Spleen/ Bone marrow	Plasma cells/ Pre-B-cells/ Naive	246,449,189 (129,417,638)
Gupta et al., [54]	Human	None	Flu/Hepatitis-A/ Hepatitis-B	PBMC	Unsorted	25,134,322 (9,966,175)
Halliley et al., [55]	Human	None	Flu/Tetanus	Bone marrow	Plasma cells	2,348,164 (1,208,616)
Huang et al., [56]	Human	HIV	None	PBMC	Memory	11,693,783 (5,701,433)
Jiang et al., [57]	Human	None	Flu	PBMC	Naïve/ Plasmablasts	3,199,271 (1,809,306)
Joyce et al., [58]	Human	None	None	PBMC	Unsorted	2,747,688 (1,463,421)
Khan et al., [59]	Mouse	None	OVA	Spleen	Unsorted	24,175,033 (7,113,411)
Levin et al., [60]	Human	Allergy	None	PBMC/ Nasal biopsy	Unsorted	528,173 (370,465)
Levin et al., [61]	Human	Allergy	None	PBMC/	Unsorted	29,643,305

				Bone marrow		(9,557,586)
Li et al., [62]	Camel	None	None	PBMC	Unsorted	1,152,359 (1,127,651)
Liao et al., [63]	Human	HIV	None	PBMC	Unsorted	1,420,314 (619,492)
Lindner et al., [64]	Mouse	None	E.Coli/ Clostridia/ Lactobacillus	Biopsy of small intestine	Unsorted	1,686,350 (544,061)
Meng et al., [65]	Human	CMV/ EBV/None	None	PBMC/Lung/ Spleen/Bone marrow/Colon/ Jejunum/Lymph node/Ileum	Unsorted	45,576,606 (21,738,501)
Menzel et al., [66]	Mouse	None	NP-CGG	Spleen/Bone marrow	ASCs	14,355,151 (6,058,480)
Mroczek et al., [67]	Human	None	None	PBMC	Immature/ Transitional/ Mature/ Plasmacytes/ Memory	104,154 (85,525)
Ota et al., [68]	Mouse	None	None	Spleen/Lymph	Unsorted	21,505 (9,619)
Palanichamy et al., [69]	Human	MS	None	CSF/PBMC	Unsorted	776,895 (292,801)
Parameswaran et al., [11]	Human	Dengue/None / Non-dengue febrile illness/	None	PBMC	Unsorted	26,584 (23,606)
Prohaska et al., [70]	Mouse	None	None	Spleen/ Peritoneum	B-1/ B-2/ Marginal Zone/ Follicular	336,723 (198,983)
Rettig et al., [32]	Mouse	None	None	Spleen/ Splenocytes	Unsorted	41,908 (24,908)
Rubelt et al., [71]	Human	None	None	PBMC	Naïve/Memory	2,320,947 (1,719,507)
Schanz et al., [31]	Human	HIV/None	None	PBMC	Unsorted	12,734,958 (5,412,549)
Zhu et al., [72]	Human	HIV	None	PBMC	Unsorted	1,962,643 (532,350)
Stern et al., [73]	Human	MS	None	Cervical lymph node/ White matter lesion/ Pia mater/ Choroid plexus/ Cortex/ Spleen	Unsorted	8,550,247 (3,321,530)
Sundling et al., [74]	Rhesus	None	HIV	PBMC	Unsorted	40,960 (26,298)
Tipton et al., [75]	Human	SLE/None	Flu/Tetanus	PBMC	Unsorted	28,204,742 (13,301,396)
Turchaninova et al., [76]	Human	None	None	PBMC	Memory/Plasma cells/Naive	183,949 (176,441)
Vander Heiden et al., [77]	Human	MG/None	None	PBMC	Memory/Naïve/ Unsorted	13,939,166 (5,170,299)
VanDuijn et al., [78]	Rat	None	DNP/HuD	Splenocytes	Unsorted	6,359,396 (4,234,597)
Vergani et al., [79]	Human	None	None	PBMC	Unsorted	14,161,949 (5,987,086)

Wasemann et al., [80]	Mouse	None	NP-CGG	Lamina propria/ Bone marrow/ Spleen	Unsorted	146,370 (40,132)
Wu et al., [81]	Human	HIV	None	PBMC	Unsorted	394,144 (198,468)
Wu et al., [82]	Human	HIV	None	PBMC	Unsorted	5,545,910 (1,370,109)
Wu et al., [83]	Human	Allergy/ None	None	PBMC/ Nasal biopsy	Unsorted	35,034 (23,923)
Zhou et al., [22]	Human	HIV	None	PBMC	Unsorted	1,541,645 (458,227)
Zhou et al., [84]	Human	HIV	None	PBMC	Unsorted	722,112 (291,670)
Zhu et al., [85]	Human	HIV	None	PBMC	Unsorted	874,930 (174,435)
Zhu et al., [86]	Human	HIV	None	PBMC	Unsorted	1,290,499 (699,828)

Table 2. **Summary of Ig-seq studies that are incorporated into the Observed Antibody Space database.** We organized the datasets among studies related to a given Ig-seq experiment. Each study in the OAS database is subdivided into Data Units. Each Data Unit is a collection of IMGT-numbered amino acid sequences uniquely identified by the metadata descriptors given in Table 1, five of which (species, disease, vaccine, B-cell source and B-cell type) are given in this Table. The 'total ANARCI parsed sequences' field indicates the total number of redundant sequences in our database, with the non-redundant numbers in parentheses. Abbreviations: PBMC, peripheral blood mononuclear cell; CLL, chronic lymphocytic leukemia; NP-CGG, chicken gamma globulin; ASC, antibody secreting cell; OVA, ovalbumin; NP-HEL, hen egg lysozyme; CSF, cerebrospinal fluid; MS, multiple sclerosis; SLE, systemic lupus erythematosus; MG, myasthenia gravis; DNP, dinitrophenyl; HuD, paraneoplastic encephalomyelitis antigen.

209 **4. Discussion**
210

Here, we describe the Observed Antibody Space (OAS) database, a unified repository to facilitate large-

scale data mining of antibody repertoires in both their amino acid and nucleotide forms. Absence of well

established repositories in Ig-seq deposition space required us to perform a combination of literature

search and manual curation of the datasets in order to organize the data into OAS. The current lack of

widely-adopted deposition standards hampers automatic updating of OAS, as datasets where we find

large number of antibodies still require manual curation to perform metadata annotation correctly.

Hopefully, efforts such as that by the AIRR community will result in standardization of Ig-seq outputs

and will further streamline deposition procedures facilitating large-scale data mining of antibody

repertoires [24]. Devising unified antibody repertoire repositories is challenging due to both the size of the

datasets as well as the diverse data descriptors and analytical pipelines desired by bioinformaticians,

wetlab scientists and clinicians [33].

OAS is the first organized collection of a large body of Ig-seq outputs. In order to allow comparative

bioinformatics analyses across different subsets of the antibody repertoires, we have annotated the

datasets by commonly used metadata descriptions such as organism, isotype, B-cell type and source,

and the immune state of B-cell donors. To facilitate research about particular antibody sequences or

regions, we make full IMGT-numbered high-quality amino acid sequences available together with gene

annotations, as well as raw nucleotide data.

This data should aid in-depth comparative analyses across different studies to discern the commonalities

observed between independent samples as well as directing Ig-seq experiments on not yet interrogated

antibody repertoires. Revealing shared preferences can be invaluable in identifying the portions of the

theoretically allowed antibody space that are strategically used to start an immune response [6].

Furthermore, such comparative studies can offer a way of deconvoluting the various degrees of freedom

of immune repertoires such as differences between diversity of isotypes [67] or organisms [87]. Charting the

differences between repertoires of human/mouse is of particular interest for engineering better

humanized biotherapeutics [88].

Beyond identifying broad commonalities across repertoires, data mining Ig-seq outputs provides novel

avenues for designing better antibody-based therapeutics. The plethora of currently available Ig-seq

data offers a glimpse at a set of sequences that should be able to fold and function in an organism.

Aligning therapeutic candidates to sequences in Ig-seq repertoires can reveal mutational choices that

might be naturally acceptable hence providing shortcuts for antibody engineering such as humanization

[89]. Furthermore, contrasting the naturally observed antibodies with therapeutic ones can offer insight as

to the naturally favored biophysical properties of these molecules [4,90,91]. All such future applications rely

on the availability of well-structured datasets that can offer a unified point of reference for

bioinformatics analyses. We hope that OAS will aid data mining antibody repertoires, help identify

strategic preferences of our immune systems and will ultimately improve how we engineer antibodies

into better therapeutics.

Acknowledgment:
We would like to thank all members of Oxford Protein Informatics Group for testing our OAS resource.

In particular, we are grateful to Garret M. Morris and Matthew Raybould for their comments, which

significantly improved the quality of our work.

Funding:
This work was supported by funding from Biotechnology and Biological Sciences Research Council

(BBSRC) [BB/M011224/1] and UCB Pharma Ltd awarded to AK.

References:
260
1.   Kindt, T. J., Goldsby, R. A., Osborne, B. A. & Kuby, J. *Kuby immunology*. (WH Freeman & Company,
2007).
2.   Glanville, J. *et al.* Precise determination of the diversity of a combinatorial antibody library gives
insight into the human immunoglobulin repertoire. *Proc. Natl. Acad. Sci. U. S. A.* **106,** 20216–
20221 (2009).
3.   Kaplon, H. & Reichert, J. M. Antibodies to watch in 2018. *mAbs* 1–21 (2018).
doi:10.1080/19420862.2018.1415671
4.   Jain, T. *et al.* Biophysical properties of the clinical-stage antibody landscape. *Proc. Natl. Acad. Sci.*
114, (2017).
5.   Miho, E. *et al.* Computational Strategies for Dissecting the High-Dimensional Complexity of
Adaptive Immune Repertoires. *Front. Immunol.* **9,** (2018).

6.   Greiff, V. *et al.* Learning the High-Dimensional Immunogenomic Features That Predict Public and

273   Private Antibody Repertoires. *J. Immunol.* ji1700594 (2017). doi:10.4049/jimmunol.1700594

274   7.   Kovaltsuk, A. *et al.* How B-Cell Receptor Repertoire Sequencing Can Be Enriched with Structural

275   Antibody Data. *Front. Immunol.* **8,** 1753 (2017).

276   8.   Georgiou, G. *et al.* The promise and challenge of high-throughput sequencing of the antibody

277   repertoire. *Nat. Biotechnol.* **32,** 158–68 (2014).

278   9.   Friedensohn, S., Khan, T. A. & Reddy, S. T. Advanced Methodologies in High-Throughput

279   Sequencing of Immune Repertoires. *Trends in Biotechnology* **35,** 203–214 (2017).

280   10.   Galson, J., Pollard, A. J., Trück, J. & Kelly, D. F. Studying the antibody repertoire after vaccination:

281   Practical applications. *Trends in Immunology* **35,** 319–331 (2014).

282   11.   Parameswaran, P. *et al.* Convergent antibody signatures in human dengue. *Cell Host Microbe* **13,**

283   691–700 (2013).

284   12.   Ghraichy, M., Galson, J. D., Kelly, D. F. & Trück, J. B-cell receptor repertoire sequencing in patients

285   with primary immunodeficiency: A review. *Immunology* 145–160 (2017). doi:10.1111/imm.12865

286   13.   Doria-Rose, N. A. *et al.* Developmental pathway for potent V1V2-directed HIV-neutralizing

287   antibodies. *Nature* **508,** 55–62 (2014).

288   14.   Greiff, V. *et al.* Systems Analysis Reveals High Genetic and Antigen-Driven Predetermination of

289   Antibody Repertoires throughout B Cell Development. *Cell Rep.* **19,** 1467–1478 (2017).

290   15.   Hoi, K. H. & Ippolito, G. C. Intrinsic bias and public rearrangements in the human immunoglobulin

291   Vλ light chain repertoire. *Genes Immun.* 1–6 (2013). doi:10.1038/gene.2013.10

292   16.   Dekosky, B. J. *et al.* In-depth determination and analysis of the human paired heavy- and light-

293   chain antibody repertoire. *Nat. Med.* **21,** 1–8 (2014).

294   17.   Galson, J., Trück, J., Kelly, D. F. & van der Most, R. Investigating the effect of AS03 adjuvant on

295   the plasma cell repertoire following pH1N1 influenza vaccination. *Sci. Rep.* **6,** 37229 (2016).

296   18.   Galson, J. *et al.* B cell repertoire dynamics after sequential Hepatitis B vaccination, and evidence

297   for cross-reactive B cell activation. *Submitt. Manuscr.* 1–13 (2016). doi:10.1186/s13073-016-

298   0322-z

299   19.   Jackson, K. J. L. *et al.* Human responses to influenza vaccination show seroconversion signatures

300   and convergent antibody rearrangements. *Cell Host Microbe* **16,** 105–114 (2014).

301   20.   Lee, J. *et al.* Molecular-level analysis of the serum antibody repertoire in young adults before and

302   after seasonal influenza vaccination. *Nat Med* **22,** 1456–1464 (2016).

21.   Galson, J. *et al.* BCR repertoire sequencing: Different patterns of B-cell activation after two

Meningococcal vaccines. *Immunol. Cell Biol.* **93,** 885–895 (2015).

22.   Zhou, T. *et al.* Multidonor analysis reveals structural elements, genetic determinants, and

maturation pathway for HIV-1 neutralization by VRC01-class antibodies. *Immunity* **39,** 245–258

(2013).

23.   DeKosky, B. J. *et al.* High-throughput sequencing of the paired human immunoglobulin heavy and

light chain repertoire. *Nat. Biotechnol.* **31,** 166–9 (2013).

24.   Rubelt, F. *et al.* Adaptive Immune Receptor Repertoire Community recommendations for sharing

immune-repertoire sequencing data. *Nature Immunology* **18,** 1274–1278 (2017).

25.   Breden, F. *et al.* Reproducibility and Reuse of Adaptive Immune Receptor Repertoire Data. *Front.*

Immunol. **8,** 1418 (2017).

26.   Bhattacharya, S. *et al.* ImmPort: Disseminating data to the public for the future of immunology.

Immunol. Res. **58,** 234–239 (2014).

27.   Bhattacharya, S. *et al.* ImmPort, toward repurposing of open access immunological assay data for

translational and clinical research. *Sci. Data* **5,** (2018).

28.   Cowell, L. G. VDJServer: A Cloud-Based Analysis Portal and Data Commons for Immune

Repertoire Sequences and Rearrangements. *Front. Immunol.* **9,** 976 (2018).

29.   Leinonen, R. *et al.* The European nucleotide archive. *Nucleic Acids Res.* **39,** (2011).

30.   NCBI Resource Coordinators. Database Resources of the National Center for Biotechnology

Information. *Nucleic Acids Res.* **45,** D12–D17 (2017).

31.   Schanz, M. *et al.* High-throughput sequencing of human immunoglobulin variable regions with

subtype identification. *PLoS One* **9,** (2014).

32.   Rettig, T. A., Ward, C., Bye, B. A., Pecaut, M. J. & Chapes, S. K. Characterization of the naive

murine antibody repertoire using unamplified high-throughput sequencing. *PLoS One* **13,** (2018).

33.   Greiff, V. *et al.* A bioinformatic framework for immune repertoire diversity profiling enables

detection of immunological status. *Genome Med.* **7,** 49 (2015).

34.   Magoč, T. & Salzberg, S. L. FLASH: Fast length adjustment of short reads to improve genome

assemblies. *Bioinformatics* **27,** 2957–2963 (2011).

35.   HannonLab. FASTX toolkit. *Cold Spring Harbor Laboratory, Cold Spring Harbor, NY* (2014).

36.   Giudicelli, V., Chaume, D. & Lefranc, M.-P. IMGT/GENE-DB: a comprehensive database for human

and mouse immunoglobulin and T cell receptor genes. *Nucleic Acids Res.* **33,** D256-61 (2005).

37.   Smith, T. F. & Waterman, M. S. Identification of common molecular subsequences. *J. Mol. Biol.*

147, 195–197 (1981).

38.   Galson, J. D. *et al.* B-cell repertoire dynamics after sequential hepatitis B vaccination and

evidence for cross-reactive B-cell activation. *Genome Med.* **8,** 68 (2016).

39.   Galson, J. *et al.* Analysis of B Cell Repertoire Dynamics Following Hepatitis B Vaccination in

Humans, and Enrichment of Vaccine-specific Antibody Sequences. *EBioMedicine* **2,** 2070–2079

(2015).

40.   Greiff, V. *et al.* Quantitative assessment of the robustness of next-generation sequencing of

antibody variable gene repertoires from immunized mice. *BMC Immunol.* **15,** 40 (2014).

41.   Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable domain

sequence analysis tool. *Nucleic Acids Res.* **41,** (2013).

42.   Dunbar, J. & Deane, C. M. ANARCI: Antigen receptor numbering and receptor classification.

Bioinformatics btv552 (2015). doi:10.1093/bioinformatics/btv552

43.   Lefranc, M.-P. *et al.* IMGT unique numbering fro immunoglobulin and T cell receptor variable

domains and Ig superfamily V-like domains. *Dev. Comp. Immunol.* **27,** 55–77 (2003).

44.   Eddy, S. R. Multiple alignment using hidden Markov models. *Proc. Int. Conf. Intell. Syst. Mol. Biol.*

3, 114–120 (1995).

45.   Shugay, M. *et al.* Towards error-free profiling of immune repertoires. *Nat. Methods* **11,** 653–5

(2014).

46.   Banerjee, S. *et al.* Evaluation of a novel multi-immunogen vaccine strategy for targeting

4E10/10E8 neutralizing epitopes on HIV-1 gp41 membrane proximal external region. *Virology*

505, 113–126 (2017).

47.   Bashford-Rogers, R. J. M. *et al.* Network properties derived from deep sequencing of human B-

cell receptor repertoires delineate B-cell populations. *Genome Res.* **23,** 1874–1884 (2013).

48.   Bhiman, J. N. *et al.* Viral variants that initiate and drive maturation of V1V2-directed HIV-1

broadly neutralizing antibodies. *Nat. Med.* **21,** 1332–1336 (2015).

49.   Bonsignori, M. *et al.* Maturation Pathway from Germline to Broad HIV-1 Neutralizer of a CD4-

Mimic Antibody. *Cell* **165,** 449–463 (2016).

50.   Collins, A. M. *et al.* The mouse antibody heavy chain repertoire is germline-focused and highly

variable between inbred strains. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **370,** S41–S52 (2015).

51.   Corcoran, M. M. *et al.* Production of individualized v gene databases reveals high levels of

immunoglobulin genetic diversity. *Nat. Commun.* **7,** (2016).

52.   Cui, A. *et al.* A Model of Somatic Hypermutation Targeting in Mice Based on High-Throughput Ig

Sequencing Data. *J. Immunol.* **197,** 3566–3574 (2016).

53.   Fisher, C. R. *et al.* T-dependent B cell responses to Plasmodium induce antibodies that form a
high-avidity multivalent complex with the circumsporozoite protein. *PLoS Pathog.* **13,** (2017).

54.   Gupta, N. T. *et al.* Hierarchical Clustering Can Identify B Cell Clones with High Confidence in Ig
Repertoire Sequencing Data. *J. Immunol.* **198,** 2489–2499 (2017).

55.   Halliley, J. L. *et al.* Long-Lived Plasma Cells Are Contained within the CD19(-)CD38(hi)CD138(+)
Subset in Human Bone Marrow. *Immunity* **43,** 132–45 (2015).

56.   Huang, J. *et al.* Identification of a CD4-Binding-Site Antibody to HIV that Evolved Near-Pan
Neutralization Breadth. *Immunity* **45,** 1108–1121 (2016).

57.   Jiang, N. *et al.* Lineage structure of the human antibody repertoire in response to influenza
vaccination. *Sci. Transl. Med.* **5,** 171ra19 (2013).

58.   Joyce, M. G. *et al.* Vaccine-Induced Antibodies that Neutralize Group 1 and Group 2 Influenza A
Viruses. *Cell* **166,** 609–623 (2016).

59.   Khan, T. A. *et al.* Accurate and predictive antibody repertoire profiling by molecular amplification
fingerprinting. *Sci. Adv.* **2,** e1501371–e1501371 (2016).

60.   Levin, M. *et al.* Persistence and evolution of allergen-specific IgE repertoires during subcutaneous
specific immunotherapy. *J. Allergy Clin. Immunol.* **137,** 1535–1544 (2016).

61.   Levin, M., Levander, F., Palmason, R., Greiff, L. & Ohlin, M. Antibody-encoding repertoires of
bone marrow and peripheral blood—a focus on IgE. *J. Allergy Clin. Immunol.* **139,** 1026–1030
(2017).

62.   Li, X. *et al.* Comparative analysis of immune repertoires between bactrian Camel's conventional
and heavy-chain antibodies. *PLoS One* **11,** (2016).

63.   Liao, H. X. *et al.* Co-evolution of a broadly neutralizing HIV-1 antibody and founder virus. *Nature*
496, 469–476 (2013).

64.   Lindner, C. *et al.* Diversification of memory B cells drives the continuous adaptation of secretory
antibodies to gut microbiota. *Nat. Immunol.* **16,** 880–888 (2015).

65.   Meng, W. *et al.* An atlas of B-cell clonal distribution in the human body. *Nat. Biotechnol.* **35,** 879–
886 (2017).

66.   Menzel, U. *et al.* Comprehensive evaluation and optimization of amplicon library preparation
methods for high-throughput antibody sequencing. *PLoS One* **9,** (2014).

67.   Mroczek, E. S. *et al.* Differences in the composition of the human antibody repertoire by B cell
subsets in the blood. *Front Immunol* **5,** 96 (2014).

68.   Ota, M. *et al.* Regulation of the B cell receptor repertoire and self-reactivity by BAFF. *J. Immunol.*

185, 4128–36 (2010).

69. Palanichamy, A. *et al.* Immunoglobulin class-switched B cells form an active immune axis between CNS and periphery in multiple sclerosis. *Sci. Transl. Med.* **6,** (2014).

70. Prohaska, T. A. *et al.* Massively Parallel Sequencing of Peritoneal and Splenic B Cell Repertoires Highlights Unique Properties of B-1 Cell Antibodies. *J. Immunol.* **200,** 1702–1717 (2018).

71. Rubelt, F. *et al.* Individual heritable differences result in unique cell lymphocyte receptor repertoires of naïve and antigen-experienced cells. *Nat. Commun.* **7,** (2016).

72. Zhu, J. *et al.* Mining the antibodyome for HIV-1-neutralizing antibodies with next-generation sequencing and phylogenetic pairing of heavy/light chains. *Proc. Natl. Acad. Sci. U. S. A.* **110,** 6470–5 (2013).

73. Stern, J. N. H. *et al.* B cells populating the multiple sclerosis brain mature in the draining cervical lymph nodes. *Sci. Transl. Med.* **6,** (2014).

74. Sundling, C. *et al.* Single-Cell and Deep Sequencing of IgG-Switched Macaque B Cells Reveal a Diverse Ig Repertoire following Immunization. *J. Immunol.* **192,** 3637–3644 (2014).

75. Tipton, C. M. *et al.* Diversity, cellular origin and autoreactivity of antibody-secreting cell population expansions in acute systemic lupus erythematosus. *Nat. Immunol.* **16,** 755–765 (2015).

76. Turchaninova, M. A. *et al.* High-quality full-length immunoglobulin profiling with unique molecular barcoding. *Nat. Protoc.* **11,** 1599–1616 (2016).

77. Vander Heiden, J. A. *et al.* Dysregulation of B Cell Repertoire Formation in Myasthenia Gravis Patients Revealed through Deep Sequencing. *J. Immunol.* **198,** 1460–1473 (2017).

78. VanDuijn, M. M., Dekker, L. J., van IJcken, W. F. J., Sillevis Smitt, P. A. E. & Luider, T. M. Immune repertoire after immunization as seen by next-generation sequencing and proteomics. *Front. Immunol.* **8,** (2017).

79. Vergani, S. *et al.* Novel method for high-throughput full-length IGHV-D-J sequencing of the immune repertoire from bulk B-cells with single-cell resolution. *Front. Immunol.* **8,** (2017).

80. Wesemann, D. R. *et al.* Microbial colonization influences early B-lineage development in the gut lamina propria. *Nature* **501,** 112–115 (2013).

81. Wu, X. *et al.* Focused evolution of HIV-1 neutralizing antibodies revealed by structures and deep sequencing. *Science* **333,** 1593–602 (2011).

82. Wu, X. *et al.* Maturation and diversity of the VRC01-antibody lineage over 15 years of chronic HIV-1 infection. *Cell* **161,** 480–485 (2015).

83. Wu, Y. C. B. *et al.* Influence of seasonal exposure to grass pollen on local and peripheral blood IgE

repertoires in patients with allergic rhinitis. *J. Allergy Clin. Immunol.* **134,** 604–612 (2014).

84. Zhou, T. *et al.* Structural repertoire of HIV-1-neutralizing antibodies targeting the CD4 supersite in

14 donors. *Cell* **161,** 1280–1292 (2015).

85. Zhu, J. *et al.* Somatic populations of PGT135-137 HIV-1-neutralizing antibodies identified by 454

pyrosequencing and bioinformatics. *Front. Microbiol.* **3,** (2012).

86. Zhu, J. *et al.* De novo identification of VRC01 class HIV-1-neutralizing antibodies by next-

generation sequencing of B-cell transcripts. *Proc. Natl. Acad. Sci.* **110,** E4088–E4097 (2013).

87. Schroeder Jr, H. W. Similarity and divergence in the development and expression of the mouse

and human antibody repertoires. *Dev. Comp. Immunol.* **30,** 119–135 (2006).

88. Zemlin, M. *et al.* Expressed murine and human CDR-H3 intervals of equal length exhibit distinct

repertoires that differ in their amino acid composition and predicted range of structures. *J. Mol.*

Biol. **334,** 733–749 (2003).

89. Olimpieri, P. P., Marcatili, P. & Tramontano, A. Tabhu: Tools for antibody humanization.

Bioinformatics **31,** 434–435 (2014).

90. Lowe, D. *et al.* Aggregation, stability, and formulation of human antibody therapeutics. in 41–61

(2011). doi:10.1016/B978-0-12-386483-3.00004-5

91. Lauer, T. M. *et al.* Developability index: A rapid in silico tool for the screening of antibody

aggregation propensity. *J. Pharm. Sci.* **101,** 102–115 (2012).

451