Title: Observed Antibody Space: a resource for data mining next generation sequencing of antibody

repertoires.

Aleksandr Kovaltsuk[1], Jinwoo Leem[1], Sebastian Kelm[2], James Snowden[2], Charlotte M. Deane[1],*, Konrad

Krawczyk[1],*

1.   University of Oxford, Department of Statistics, Oxford, UK

2.   UCB Pharma, Slough, UK

* to whom the correspondence should be addressed,

deane@stats.ox.ac.uk

konrad@proteincontact.org

Abstract. Antibodies are immune system proteins that recognize noxious molecules for elimination.

Their sequence diversity and binding versatility have made antibodies the primary class of

biopharmaceuticals. Recently it has become possible to query their immense natural diversity using

next-generation sequencing of immunoglobulin gene repertoires (Ig-seq). However, Ig-seq outputs are

currently fragmented across repositories and tend to be presented as raw nucleotide reads, which

means nontrivial effort is required to reuse the data for analysis. To address this issue, we have

collected Ig-seq outputs from 53 studies, covering more than half a billion antibody sequences across

diverse immune states, organisms and individuals. We have sorted, cleaned, annotated, translated and

numbered these sequences and make the data available via our Observed Antibody Space (OAS)

resource at antibodymap.org. The data within OAS will be regularly updated with newly released Ig-seq

datasets. We believe OAS will facilitate data mining of immune repertoires for improved understanding

of the immune system and development of better biotherapeutics.

1. Introduction

Antibodies (or B-cell receptors) are protein products of B-cells and primary actors of adaptive immunity

in jawed vertebrates (1). They are highly malleable molecules that can bind to virtually any antigen. An

organism holds a great variety of these molecules increasing the probability that an arbitrary antigen

can be recognized by an antibody, initiating an immune response (2). Owing to their binding malleability

they are the most prominent class of reagents and biotherapeutics (3, 4). Continued successful

exploitation of these molecules relies on our ability to discern the functional diversity of antibody

repertoires (5–7).

Next-generation sequencing of immunoglobulin gene repertoires (Ig-seq) has enabled researchers to

take snapshots of millions of sequences at a time across individuals, diverse organisms and different

immune states (8, 9). The ability to sequence and analyze millions of antibody sequences has the

potential to uncover the mechanics of the immune response to any antigen (10, 11) and dysfunctions of

the immune system itself (12).

Many previous studies have addressed the issue of antibody diversity, contributing invaluable evidence

to understanding the dynamics of human immune systems (13). Numerous analyses have focused on

the frequencies of V(D)J gene usages, which can offer insights into creating biased therapeutic antibody

libraries (14–16). Another therapeutic application of antibody repertoire analysis is advancing vaccine

design by comparative longitudinal studies of pre- and post-antigen challenge experiments (10, 11, 17–

22). Such comparative studies have shown that different individuals can converge on the same antibody

sequence against a given vaccine (11, 19). Due to sequencing limitations, these analyses have focused

on heavy or light chains separately, whereas one ought to study the paired repertoire to obtain deeper

insights of antibody diversity (23).

Technical advances in sequencing technology have outpaced storage and analysis pipelines (24, 25). This

has meant that the outputs of Ig-seq studies are fragmented across repositories making it difficult to

perform large-scale data mining of antibody repertoires (25). Metadata such as isotype, age or subject

identifiers are not typically standardized, therefore extraction of specific subsets of antibody repertoires

for comparative analyses is challenging. Furthermore, the data are typically deposited as raw nucleotide

reads. It requires non-trivial *ad hoc* effort to convert such raw reads to amino acid sequences that

ultimately dictate the molecular structure and antigen-recognition. Some of these issues are addressed

by services that provide Ig-seq-specific data deposition and analysis pipelines such as Immport

(http://immport.org) (26, 27), ImmunoseqAnalyzer (http://clients.adaptivebiotech.com/), IReceptor

(http://ireceptor.irmacs.sfu.ca/) or VDJServer (http://vdjserver.org) (28). The IReceptor and the

VDJServer are the main resources that fall under the umbrella of the organized effort of the Adaptive

Immune Receptor Repertoire (AIRR) Community to provide standardized deposition and analysis

pipelines for the Ig-seq outputs (24). These services chiefly focus on facilitating bulk deposition of raw

data to perform standardized sequencing analyses. Ultimately, because immunoinformatics is not the

chief focus of such services, bulk data download from such websites is limited and converting the raw

nucleotide data obtained into a format suitable for analysis still requires non-trivial effort.

To address these issues, we have created the Observed Antibody Space (OAS) resource that allows large-

scale data mining of antibody repertoires. We have so far collected the raw outputs of 53 Ig-seq

experiments covering over half a billion sequences. We have organized the sequences by metadata such

as organism, isotype, B-cell type and source, and the immune status of B-cell donors to facilitate bulk

retrieval of specific subsets for comparative analyses. We have converted all of the Ig-seq sequences to

amino acids and numbered them using the IMGT scheme. The data is available for querying or bulk

download at http://antibodymap.org. We believe that OAS will facilitate data mining antibody

repertoires for improved understanding of the dynamics of the immune system and thus better

engineering of biotherapeutics.

2. Materials and Methods
80

A list of study accession codes of publically available Ig-seq datasets were obtained via a literature

review. The majority of raw reads were downloaded from the European Nucleotide Archive (ENA) (29)

and the National Center for Biotechnology Information (NCBI) websites (30). In a small number of cases,

another public Ig-seq repository was specified e.g. (14, 31–33). Metadata were manually extracted from

the deposited datasets and arranged in a reproducible format.

The downloaded FASTQ files were processed depending on the sequencing platform. Paired raw

Illumina reads were assembled with FLASH (34). The assembled antibody sequences were converted to

the FASTA format using FASTX-toolkit (35). As raw reads from Roche 454 are not paired, these FASTQ

files were directly converted to the FASTA format with the FASTX-toolkit.

The heavy chain sequences were automatically annotated with isotype information unless such data was

given in the corresponding publication. Automatic isotype annotation was performed by aligning the

constant heavy domain 1 (CH1) of any given antibody sequence against the IMGT isotype reference (36)

of the respective species using the Smith-Waterman algorithm (37). We assigned a score of two for a

nucleotide match, and a score of minus one for a nucleotide mismatch or a gap. The IMGT isotype

references comprised 21 nucleotide-long fragments of the CH1 domain of the antibody isotypes. To

ensure a high confidence of correct isotype identification, we employed a conservative threshold of 30

in the Smith-Waterman algorithm scoring function. Sequences whose Smith-Waterman algorithm score

was below the threshold for all isotypes were assigned as 'bulk'. The robustness of this protocol was

confirmed on the author-annotated Ig-seq datasets (38–40) where it resulted in 99% accurate

annotations. Around 1% of the Ig-seq data had a very short (or missing) of CH1 domain sequence. Such

sequences were also assigned as 'bulk'.

IgBlastn (41) was used to convert the FASTA files of antibody nucleotide sequences to amino acids. The

amino acid sequences were then numbered with ANARCI (42) using the IMGT scheme (43). ANARCI does

not number a sequence if it does not align to a suitable Hidden Markov Model (44). ANARCI therefore

ensures that the antibody sequences do not harbor unusual indels or stop codons in the antibody

regions, that the V and J genes align to the respective species amino acid IMGT germlines (36), and that

the length of CDR-H3 is not greater than 37 residues in human, mouse, rat, rabbit, alpaca and rhesus

antibodies. Due to technical limitations of sequencing platforms, certain reads were missing significant

portions of the variable region (e.g. portions of CDR1), sequences that did not have all three CDRs were

discarded as incomplete. The V and J genes are identified during the ANARCI numbering step.

Using the protocol above we annotated Ig-seq results of 53 independent studies. In order to streamline

updating OAS with new data, we have generated a procedure to automatically identify Ig-seq datasets

from raw sequence read archives. We apply our antibody annotation protocol to each raw nucleotide

dataset deposited in the NCBI/ENA repositories, if we find more than 10,000 antibody sequences in any

given dataset, it is set aside for manual inspection. Manual inspection is still necessary to efficiently

assign metadata as these are currently deposited in a non-standardized manner. This procedure allows

for automatic identification of new Ig-seq datasets and semi automatically updating of OAS.

3. Results

We have so far collected raw sequencing outputs from 53 Ig-seq studies. All raw nucleotide reads were

converted into amino acids using IgBlastn (41). The full amino acid sequences were then IMGT

numbered using ANARCI (42). As well as providing IMGT and gene annotations, ANARCI acts as a broad-

brush filter for antibody sequences that are likely to be erroneous (see Materials and Methods).

Applying the same retrieval, amino acid conversion, gene annotation and numbering protocol to all

sequences assures the same point of reference across the 53 heterogeneous Ig-seq datasets (45). This

protocol produces the full IMGT-numbered sequences together with gene annotations for each of the

datasets.

Figure 1. T**he Observed Antibody Space database**. The data from 53 studies is sorted into Data

Units. Each Data Unit is a set of antibody sequences that share the same set of meta-data. Each

sequence in a Data Unit is further associated with sequence-specific annotations.

The numbered amino acid sequences in each dataset are sorted by metadata e.g. individuals, age,

vaccination regime, B-cell type and source *etc*. (Figure 1). Deposition of such metadata is currently not

standardized and requires *ad hoc* manual curation for each dataset. In an effort to organize the antibody

sequences using such metadata, we have grouped the sequences within each dataset into Data Units.

Each Data Unit represents a group of sequences within a given dataset with a unique combination of

metadata values. The metadata values are summarized in Table 1.

Metadata name	Metadata description
Chain	Heavy/light chain annotation.
Isotype	Identified or deposited isotype information.
Age	Information on the age of the human B-cell donors.
Disease	Indication of whether the donor was sick at the time of B-cell extraction.
Vaccine	Indication if the B-cell donor was purposely immunized prior to B-cell extraction.
B-cell subset	Indication if a particular B-cell subset was sorted for Ig-seq
Species	Organism of the B-cell donor
B-cell source	Which organ/tissue the B-cells were extracted from.
Subject	Indication of a particular B-cell donor the B-cells were sourced from.
Longitudinal	If the study was longitudinal, an indicator of the time point.
Size	Number of non-redundant sequences in the dataset.
Link	Link to the source publication.

147   <u>Table 1.</u> Metadata descriptors of each Data Unit in OAS. Each Data Unit is uniquely identified by the

148   study and a collection of the metadata values.

As of April 29th 2018, 53 Ig-seq studies are included in OAS totaling 608,651,423 sequences

(552,824,460 VH and 55,826,963 VL sequences). The majority of these sequences are murine

(~50.4%) and human (~47.4%). Twenty-two of the Ig-seq studies interrogate the immune system of

diseased individuals, the most common ailment being HIV (13 studies). The database also contains 22 Ig-

seq studies of naive antibody gene repertoires (the collection of B-cells from donors who are healthy

and not purposefully vaccinated). The main source of B-cells in the OAS database is peripheral blood

(~231m of sequences) followed by spleen/splenocytes (~198m) and bone marrow (~124m). The

database holds isotype information for each individual heavy sequence and the two most common

isotypes are IgM (~312m) and IgG (~139m). For ~65m sequences we were not able to assign isotypes

with high confidence. The median total redundant size of the Ig-seq studies in the OSA database is

2,006,196 sequences, while the largest Ig-seq study was that by Greiff et al., (246,449,189 redundant

sequences) (14). Detailed statistics on each dataset are given in Table 2. All the data may be bulk

downloaded or individual Data Units queried, at http://antibodymap.org.

Study	Species	Disease	Vaccine	B-cell source	B-cell subset	Total ANARCI parsed sequences
Banerjee et al., (46)	Rabbit	None	HIV	PBMC	Unsorted	4,334,088 (2,926,727)
Bashford-Rogers et al., (47)	Human	CLL/None	None	PBMC	Unsorted	129,013 (86,166)
Bhiman et al., (48)	Human	HIV	None	PBMC	Unsorted	785,751 (187,067)
Bonsignori et al., (49)	Human	HIV/None	None	PBMC	Memory/ Unsorted	210,377 (57,374)
Collins et al., (50)	Mouse	None	None	Splenocytes	Unsorted	812,439 (194,752)
Corcoran et al., (51)	Human/ Mouse/ Rhesus	None	None	PBMC	Unsorted	5,307,880 (2,840,877)
Cui et al., (52)	Mouse	None	NP-CGG/None	Splenocytes	Memory	5,513,816 (935,646)
Doria-Rose et al., (13)	Human	HIV	None	PBMC	Unsorted	2,164,901 (549,544)
Fisher et al.,(53)	Mouse	None	Plasmodium	Spleen	Unsorted	175,015 (113,594)
Galson et al., (38)	Human	None	Hepatitis-B	PBMC	Unsorted/ Plasma cells/ HepB-specific	21,755,739 (10,442,291)
Galson et al., (39)	Human	None	Hepatitis-B	PBMC	Unsorted/ Plasma cells/ HepB-specific	26,687,394 (14,343,236)
Galson et al., (21)	Human	None	Meningitis	PBMC	Naïve/ Plasma cells/ Memory/ Marginal zone	7,918,197 (3,282,907)
Galson et al., (17)	Human	None	Flu	PBMC	Plasma cells	13,685,210 (5,065,786)
Greiff et al., (40)	Mouse	None	NP-CGG	Bone marrow/ Spleen	Plasma cells/ Plasmablasts	7,955,739 (2,891,649)
Greiff et al., (33)	Mouse	None	NP-CGG	Spleen	ASCs/Plasma cells/ Naive	788,787 (523,716)
Greiff et al., (14)	Mouse	None	OVA/Hepatitis-B/ NP-HEL/None	Spleen/ Bone marrow	Plasma cells/ Pre-B-cells/ Naive	246,449,189 (129,417,638)
Gupta et al., (54)	Human	None	Flu/Hepatitis-A/ Hepatitis-B	PBMC	Unsorted	25,134,322 (9,966,175)
Halliley et al., (55)	Human	None	Flu/Tetanus	Bone marrow	Plasma cells	2,348,164 (1,208,616)
Huang et al., (56)	Human	HIV	None	PBMC	Memory	11,693,783 (5,701,433)
Jiang et al., (57)	Human	None	Flu	PBMC	Naïve/ Plasmablasts	3,199,271 (1,809,306)
Joyce et al., (58)	Human	None	None	PBMC	Unsorted	2,747,688 (1,463,421)
Khan et al., (59)	Mouse	None	OVA	Spleen	Unsorted	24,175,033 (7,113,411)
Levin et al., (60)	Human	Allergy	None	PBMC/ Nasal biopsy	Unsorted	528,173 (370,465)

Levin et al., (61)	Human	Allergy	None	PBMC/ Bone marrow	Unsorted	29,643,305 (9,557,586)
Li et al., (62)	Camel	None	None	PBMC	Unsorted	1,152,359 (1,127,651)
Liao et al., (63)	Human	HIV	None	PBMC	Unsorted	1,420,314 (619,492)
Lindner et al., (64)	Mouse	None	E.Coli/ Clostridia/ Lactobacillus	Biopsy of small intestine	Unsorted	1,686,350 (544,061)
Meng et al., (65)	Human	CMV/ EBV/None	None	PBMC/Lung/ Spleen/Bone marrow/Colon/ Jejunum/Lymph node/Ileum	Unsorted	45,576,606 (21,738,501)
Menzel et al., (66)	Mouse	None	NP-CGG	Spleen/Bone marrow	ASCs	14,355,151 (6,058,480)
Mroczek et al., (67)	Human	None	None	PBMC	Immature/ Transitional/ Mature/ Plasmacytes/ Memory	104,154 (85,525)
Ota et al., (68)	Mouse	None	None	Spleen/Lymph	Unsorted	21,505 (9,619)
Palanichamy et al., (69)	Human	MS	None	CSF/PBMC	Unsorted	776,895 (292,801)
Parameswaran et al., (11)	Human	Dengue/None / Non-dengue febrile illness/	None	PBMC	Unsorted	26,584 (23,606)
Prohaska et al., (70)	Mouse	None	None	Spleen/ Peritoneum	B-1/ B-2/ Marginal Zone/ Follicular	336,723 (198,983)
Rettig et al., (32)	Mouse	None	None	Spleen/ Splenocytes	Unsorted	41,908 (24,908)
Rubelt et al., (71)	Human	None	None	PBMC	Naïve/Memory	2,320,947 (1,719,507)
Schanz et al., (31)	Human	HIV/None	None	PBMC	Unsorted	12,734,958 (5,412,549)
Zhu et al., (72)	Human	HIV	None	PBMC	Unsorted	1,962,643 (532,350)
Stern et al., (73)	Human	MS	None	Cervical lymph node/ White matter lesion/ Pia mater/ Choroid plexus/ Cortex/ Spleen	Unsorted	8,550,247 (3,321,530)
Sundling et al., (74)	Rhesus	None	HIV	PBMC	Unsorted	40,960 (26,298)
Tipton et al., (75)	Human	SLE/None	Flu/Tetanus	PBMC	Unsorted	28,204,742 (13,301,396)
Turchaninova et al., (76)	Human	None	None	PBMC	Memory/Plasma cells/Naive	183,949 (176,441)
Vander Heiden et al., (77)	Human	MG/None	None	PBMC	Memory/Naïve/ Unsorted	13,939,166 (5,170,299)
VanDuijn et al., (78)	Rat	None	DNP/HuD	Splenocytes	Unsorted	6,359,396 (4,234,597)

Vergani et al., (79)	Human	None	None	PBMC	Unsorted	14,161,949 (5,987,086)
Wasemann et al., (80)	Mouse	None	NP-CGG	Lamina propria/ Bone marrow/ Spleen	Unsorted	146,370 (40,132)
Wu et al., (81)	Human	HIV	None	PBMC	Unsorted	394,144 (198,468)
Wu et al., (82)	Human	HIV	None	PBMC	Unsorted	5,545,910 (1,370,109)
Wu et al., (83)	Human	Allergy/ None	None	PBMC/ Nasal biopsy	Unsorted	35,034 (23,923)
Zhou et al., (22)	Human	HIV	None	PBMC	Unsorted	1,541,645 (458,227)
Zhou et al., (84)	Human	HIV	None	PBMC	Unsorted	722,112 (291,670)
Zhu et al., (85)	Human	HIV	None	PBMC	Unsorted	874,930 (174,435)
Zhu et al., (86)	Human	HIV	None	PBMC	Unsorted	1,290,499 (699,828)

Table 2. **Summary of Ig-seq studies that are incorporated into the Observed Antibody Space database.**

The datasets are organized into studies related to a given Ig-seq experiment. Each study in the OAS

database is subdivided into Data Units. Each Data Unit is a collection of IMGT-numbered amino acid

sequences uniquely identified by the metadata descriptors given in Table 1, five of which (species,

disease, vaccine, B-cell source and B-cell type) are given in this Table. The 'total ANARCI parsed

sequences' field indicates the total number of redundant sequences in our database, with the non-

redundant numbers in parentheses. Abbreviations: PBMC, peripheral blood mononuclear cell; CLL,

chronic lymphocytic leukemia; NP-CGG, chicken gamma globulin; ASC, antibody secreting cell; OVA,

ovalbumin; NP-HEL, hen egg lysozyme; CSF, cerebrospinal fluid; MS, multiple sclerosis; SLE, systemic

lupus erythematosus; MG, myasthenia gravis; DNP, dinitrophenyl; HuD, paraneoplastic

encephalomyelitis antigen.

4. Discussion

Here, we describe the Observed Antibody Space (OAS) database, a unified repository to facilitate large-scale data mining of antibody repertoires in both their amino acid and nucleotide forms. Absence of well established repositories in Ig-seq deposition space required us to perform a combination of literature search and manual curation of the datasets in order to organize the data into OAS. The current lack of widely-adopted deposition standards hampers automatic updating of OAS, as datasets where we find large number of antibodies still require manual curation to perform metadata annotation correctly. Hopefully, efforts such as that by the AIRR community will result in standardization of Ig-seq outputs and will further streamline deposition procedures facilitating large-scale data mining of antibody repertoires (24). Devising unified antibody repertoire repositories is challenging due to both the size of the datasets as well as the diverse data descriptors and analytical pipelines desired by bioinformaticians, wetlab scientists and clinicians (33).

OAS is the first organized collection of a large body of Ig-seq outputs. In order to allow comparative bioinformatics analyses across different subsets of antibody repertoires, we have annotated the datasets by commonly used metadata descriptions such as organism, isotype, B-cell type and source, and the immune state of B-cell donors. To facilitate research about particular antibody sequences or regions, we make full IMGT-numbered high-quality amino acid sequences available together with gene annotations, as well as raw nucleotide data.

This data should aid in-depth comparative analyses across different studies to discern the commonalities observed between independent samples as well as directing Ig-seq experiments on not yet interrogated antibody repertoires. Revealing shared preferences can be invaluable in identifying the portions of the theoretically allowed antibody space that are strategically used to start immune responses (6). Furthermore, such comparative studies can offer a way of deconvoluting the various degrees of freedom of immune repertoires such as differences between diversity of isotypes (67) or organisms (87). Charting the differences between repertoires of human/mouse is of particular interest for engineering better humanized biotherapeutics (88).

Beyond identifying broad commonalities across repertoires, data mining Ig-seq outputs provides novel avenues for designing better antibody-based therapeutics. The plethora of currently available Ig-seq data offers a glimpse at a set of sequences that should be able to fold and function in an organism. Aligning therapeutic candidates to sequences in Ig-seq repertoires can reveal mutational choices that might be naturally acceptable hence providing shortcuts for antibody engineering such as humanization (89). Furthermore, contrasting the naturally observed antibodies with therapeutic ones can offer insight as to the naturally favored biophysical properties of these molecules (4). All such future applications rely on the availability of well-structured datasets that can offer a unified point of reference for bioinformatics analyses. We hope that OAS will aid data mining antibody repertoires, help identify strategic preferences of our immune systems and will ultimately improve how we engineer antibodies into better therapeutics.

Acknowledgment:
We would like to thank all members of Oxford Protein Informatics Group for testing our OAS resource. In particular, we are grateful to Garret M. Morris and Matthew Raybould for their comments, which significantly improved the quality of our work.

Funding:
This work was supported by funding from Biotechnology and Biological Sciences Research Council (BBSRC) [BB/M011224/1] and UCB Pharma Ltd awarded to AK.

References:

1. Kindt, T. J., R. A. Goldsby, B. A. Osborne, and J. Kuby. 2007. *Kuby immunology*,. WH Freeman & Company.

2. Glanville, J., W. Zhai, J. Berka, D. Telman, G. Huerta, G. R. Mehta, I. Ni, L. Mei, P. D. Sundar, G. M. R. Day, D. Cox, A. Rajpal, and J. Pons. 2009. Precise determination of the diversity of a combinatorial antibody library gives insight into the human immunoglobulin repertoire. *Proc. Natl. Acad. Sci. U. S. A.* 106: 20216–20221.

3. Kaplon, H., and J. M. Reichert. 2018. Antibodies to watch in 2018. *MAbs* 1–21.

4. Jain, T., T. Sun, S. Durand, A. Hall, N. R. Houston, J. H. Nett, B. Sharkey, B. Bobrowicz, I. Caffry, Y. Yu, Y. Cao, H. Lynaugh, M. Brown, H. Baruah, L. T. Gray, E. M. Krauland, Y. Xu, M. Vásquez, and K. D. Wittrup.

2017. Biophysical properties of the clinical-stage antibody landscape. *Proc. Natl. Acad. Sci.* 114.

5. Miho, E., A. Yermanos, C. R. Weber, C. T. Berger, S. T. Reddy, and V. Greiff. 2018. Computational

Strategies for Dissecting the High-Dimensional Complexity of Adaptive Immune Repertoires. *Front.*

Immunol. 9.

6. Greiff, V., C. R. Weber, J. Palme, U. Bodenhofer, E. Miho, U. Menzel, and S. T. Reddy. 2017. Learning

the High-Dimensional Immunogenomic Features That Predict Public and Private Antibody Repertoires. *J.*

Immunol. ji1700594.

7. Kovaltsuk, A., K. Krawczyk, J. D. Galson, D. F. Kelly, C. M. Deane, and J. Trück. 2017. How B-Cell

Receptor Repertoire Sequencing Can Be Enriched with Structural Antibody Data. *Front. Immunol.* 8:

1753.

8. Georgiou, G., G. C. Ippolito, J. Beausang, C. E. Busse, H. Wardemann, and S. R. Quake. 2014. The

promise and challenge of high-throughput sequencing of the antibody repertoire. *Nat. Biotechnol.* 32:

158–68.

9. Friedensohn, S., T. A. Khan, and S. T. Reddy. 2017. Advanced Methodologies in High-Throughput

Sequencing of Immune Repertoires. *Trends Biotechnol.* 35: 203–214.

10. Galson, J., A. J. Pollard, J. Trück, and D. F. Kelly. 2014. Studying the antibody repertoire after

vaccination: Practical applications. *Trends Immunol.* 35: 319–331.

11. Parameswaran, P., Y. Liu, K. M. Roskin, K. K. L. Jackson, V. P. Dixit, J. Y. Lee, K. L. Artiles, S. Zompi, M.

J. Vargas, B. B. Simen, B. Hanczaruk, K. R. McGowan, M. A. Tariq, N. Pourmand, D. Koller, A. Balmaseda,

S. D. Boyd, E. Harris, and A. Z. Fire. 2013. Convergent antibody signatures in human dengue. *Cell Host*

Microbe 13: 691–700.

12. Ghraichy, M., J. D. Galson, D. F. Kelly, and J. Trück. 2017. B-cell receptor repertoire sequencing in

patients with primary immunodeficiency: A review. *Immunology* 145–160.

13. Doria-Rose, N. A., C. A. Schramm, J. Gorman, P. L. Moore, J. N. Bhiman, B. J. DeKosky, M. J. Ernandes,

I. S. Georgiev, H. J. Kim, M. Pancera, R. P. Staupe, H. R. Altae-Tran, R. T. Bailer, E. T. Crooks, A. Cupo, A.

Druz, N. J. Garrett, K. H. Hoi, R. Kong, M. K. Louder, N. S. Longo, K. McKee, M. Nonyane, S. O'Dell, R. S.

Roark, R. S. Rudicell, S. D. Schmidt, D. J. Sheward, C. Soto, C. K. Wibmer, Y. Yang, Z. Zhang, J. C. Mullikin,

J. M. Binley, R. W. Sanders, I. A. Wilson, J. P. Moore, A. B. Ward, G. Georgiou, C. Williamson, S. S. A.

Karim, L. Morris, P. D. Kwong, L. Shapiro, and J. R. Mascola. 2014. Developmental pathway for potent

V1V2-directed HIV-neutralizing antibodies. *Nature* 508: 55–62.

14. Greiff, V., U. Menzel, E. Miho, C. Weber, R. Riedel, S. Cook, A. Valai, T. Lopes, A. Radbruch, T. H.

Winkler, and S. T. Reddy. 2017. Systems Analysis Reveals High Genetic and Antigen-Driven

Predetermination of Antibody Repertoires throughout B Cell Development. *Cell Rep.* 19: 1467–1478.

15. Hoi, K. H., and G. C. Ippolito. 2013. Intrinsic bias and public rearrangements in the human

immunoglobulin Vλ light chain repertoire. *Genes Immun.* 1–6.

16. Dekosky, B. J., T. Kojima, A. Rodin, W. Charab, G. C. Ippolito, A. D. Ellington, and G. Georgiou. 2014.

In-depth determination and analysis of the human paired heavy- and light-chain antibody repertoire.

Nat. Med. 21: 1–8.

17. Galson, J., J. Trück, D. F. Kelly, and R. van der Most. 2016. Investigating the effect of AS03 adjuvant

on the plasma cell repertoire following pH1N1 influenza vaccination. *Sci. Rep.* 6: 37229.

18. Galson, J., J. Trück, E. A. Clutterbuck, A. Fowler, V. Cerundolo, A. J. Pollard, G. Lunter, and D. F. Kelly.

2016. B cell repertoire dynamics after sequential Hepatitis B vaccination, and evidence for cross-reactive

B cell activation. *Submitt. Manuscr.* 1–13.

19. Jackson, K. J. L., Y. Liu, K. M. Roskin, J. Glanville, R. A. Hoh, K. Seo, E. L. Marshall, T. C. Gurley, M. A.

Moody, B. F. Haynes, E. B. Walter, H. X. Liao, R. A. Albrecht, A. García-Sastre, J. Chaparro-Riggers, A.

Rajpal, J. Pons, B. B. Simen, B. Hanczaruk, C. L. Dekker, J. Laserson, D. Koller, M. M. Davis, A. Z. Fire, and

S. D. Boyd. 2014. Human responses to influenza vaccination show seroconversion signatures and

convergent antibody rearrangements. *Cell Host Microbe* 16: 105–114.

20. Lee, J., D. R. Boutz, V. Chromikova, M. G. Joyce, C. Vollmers, K. Leung, A. P. Horton, B. J. DeKosky, C.-

H. Lee, J. J. Lavinder, E. M. Murrin, C. Chrysostomou, K. H. Hoi, Y. Tsybovsky, P. V Thomas, A. Druz, B.

Zhang, Y. Zhang, L. Wang, W.-P. Kong, D. Park, L. I. Popova, C. L. Dekker, M. M. Davis, C. E. Carter, T. M.

Ross, A. D. Ellington, P. C. Wilson, E. M. Marcotte, J. R. Mascola, G. C. Ippolito, F. Krammer, S. R. Quake,

P. D. Kwong, and G. Georgiou. 2016. Molecular-level analysis of the serum antibody repertoire in young

adults before and after seasonal influenza vaccination. *Nat Med* 22: 1456–1464.

21. Galson, J., E. A. Clutterbuck, J. Trück, M. N. Ramasamy, M. Münz, A. Fowler, V. Cerundolo, A. J.

Pollard, G. Lunter, and D. F. Kelly. 2015. BCR repertoire sequencing: Different patterns of B-cell

activation after two Meningococcal vaccines. *Immunol. Cell Biol.* 93: 885–895.

22. Zhou, T., J. Zhu, X. Wu, S. Moquin, B. Zhang, P. Acharya, I. S. Georgiev, H. R. Altae-Tran, G. Y. Chuang,

M. G. Joyce, Y. DoKwon, N. S. Longo, M. K. Louder, T. Luongo, K. McKee, C. A. Schramm, J. Skinner, Y.

Yang, Z. Yang, Z. Zhang, A. Zheng, M. Bonsignori, B. F. Haynes, J. F. Scheid, M. C. Nussenzweig, M. Simek,

D. R. Burton, W. C. Koff, J. C. Mullikin, M. Connors, L. Shapiro, G. J. Nabel, J. R. Mascola, and P. D. Kwong.

2013. Multidonor analysis reveals structural elements, genetic determinants, and maturation pathway

for HIV-1 neutralization by VRC01-class antibodies. *Immunity* 39: 245–258.

23. DeKosky, B. J., G. C. Ippolito, R. P. Deschner, J. J. Lavinder, Y. Wine, B. M. Rawlings, N. Varadarajan, C.

Giesecke, T. Dörner, S. F. Andrews, P. C. Wilson, S. P. Hunicke-Smith, C. G. Willson, A. D. Ellington, and G.

Georgiou. 2013. High-throughput sequencing of the paired human immunoglobulin heavy and light

chain repertoire. *Nat. Biotechnol.* 31: 166–9.

24. Rubelt, F., C. E. Busse, S. A. C. Bukhari, J. P. Bürckert, E. Mariotti-Ferrandiz, L. G. Cowell, C. T. Watson,

N. Marthandan, W. J. Faison, U. Hershberg, U. Laserson, B. D. Corrie, M. M. Davis, B. Peters, M. P.

Lefranc, J. K. Scott, F. Breden, E. T. Luning Prak, and S. H. Kleinstein. 2017. Adaptive Immune Receptor

Repertoire Community recommendations for sharing immune-repertoire sequencing data. *Nat.*

Immunol. 18: 1274–1278.

25. Breden, F., E. T. Luning Prak, B. Peters, F. Rubelt, C. A. Schramm, C. E. Busse, J. A. Vander Heiden, S.

Christley, S. A. C. Bukhari, A. Thorogood, F. A. Matsen IV, Y. Wine, U. Laserson, D. Klatzmann, D. C.

Douek, M.-P. Lefranc, A. M. Collins, T. Bubela, S. H. Kleinstein, C. T. Watson, L. G. Cowell, J. K. Scott, and

T. B. Kepler. 2017. Reproducibility and Reuse of Adaptive Immune Receptor Repertoire Data. *Front.*

Immunol. 8: 1418.

26. Bhattacharya, S., S. Andorf, L. Gomes, P. Dunn, H. Schaefer, J. Pontius, P. Berger, V. Desborough, T.

Smith, J. Campbell, E. Thomson, R. Monteiro, P. Guimaraes, B. Walters, J. Wiser, and A. J. Butte. 2014.

ImmPort: Disseminating data to the public for the future of immunology. *Immunol. Res.* 58: 234–239.

27. Bhattacharya, S., P. Dunn, C. G. Thomas, B. Smith, H. Schaefer, J. Chen, Z. Hu, K. A. Zalocusky, R. D.

Shankar, S. S. Shen-Orr, E. Thomson, J. Wiser, and A. J. Butte. 2018. ImmPort, toward repurposing of

open access immunological assay data for translational and clinical research. *Sci. Data* 5.

28. Cowell, L. G. 2018. VDJServer: A Cloud-Based Analysis Portal and Data Commons for Immune

Repertoire Sequences and Rearrangements. *Front. Immunol.* 9: 976.

29. Leinonen, R., R. Akhtar, E. Birney, L. Bower, A. Cerdeno-Tárraga, Y. Cheng, I. Cleland, N. Faruque, N.

Goodgame, R. Gibson, G. Hoad, M. Jang, N. Pakseresht, S. Plaister, R. Radhakrishnan, K. Reddy, S.

Sobhany, P. Ten Hoopen, R. Vaughan, V. Zalunin, and G. Cochrane. 2011. The European nucleotide

archive. *Nucleic Acids Res.* 39.

30. NCBI Resource Coordinators. 2017. Database Resources of the National Center for Biotechnology

Information. *Nucleic Acids Res.* 45: D12–D17.

31. Schanz, M., T. Liechti, O. Zagordi, E. Miho, S. T. Reddy, H. F. Günthard, A. Trkola, and M. Huber. 2014.

High-throughput sequencing of human immunoglobulin variable regions with subtype identification.

PLoS One 9.

32. Rettig, T. A., C. Ward, B. A. Bye, M. J. Pecaut, and S. K. Chapes. 2018. Characterization of the naive

murine antibody repertoire using unamplified high-throughput sequencing. *PLoS One* 13.

33. Greiff, V., P. Bhat, S. C. Cook, U. Menzel, W. Kang, and S. T. Reddy. 2015. A bioinformatic framework

for immune repertoire diversity profiling enables detection of immunological status. *Genome Med.* 7:

49.

34. Magoč, T., and S. L. Salzberg. 2011. FLASH: Fast length adjustment of short reads to improve genome

assemblies. *Bioinformatics* 27: 2957–2963.

35. HannonLab. 2014. FASTX toolkit. *Cold Spring Harb. Lab. Cold Spring Harb. NY* .

36. Giudicelli, V., D. Chaume, and M.-P. Lefranc. 2005. IMGT/GENE-DB: a comprehensive database for

human and mouse immunoglobulin and T cell receptor genes. *Nucleic Acids Res.* 33: D256-61.

37. Smith, T. F., and M. S. Waterman. 1981. Identification of common molecular subsequences. *J. Mol.*

Biol. 147: 195–197.

38. Galson, J. D., J. Trück, E. A. Clutterbuck, A. Fowler, V. Cerundolo, A. J. Pollard, G. Lunter, and D. F.

Kelly. 2016. B-cell repertoire dynamics after sequential hepatitis B vaccination and evidence for cross-

reactive B-cell activation. *Genome Med.* 8: 68.

39. Galson, J., J. Trück, A. Fowler, E. A. Clutterbuck, M. Münz, V. Cerundolo, C. Reinhard, R. van der Most,

A. J. Pollard, G. Lunter, and D. F. Kelly. 2015. Analysis of B Cell Repertoire Dynamics Following Hepatitis B

Vaccination in Humans, and Enrichment of Vaccine-specific Antibody Sequences. *EBioMedicine* 2: 2070–

2079.

40. Greiff, V., U. Menzel, U. Haessler, S. C. Cook, S. Friedensohn, T. A. Khan, M. Pogson, I. Hellmann, and

S. T. Reddy. 2014. Quantitative assessment of the robustness of next-generation sequencing of antibody

variable gene repertoires from immunized mice. *BMC Immunol.* 15: 40.

41. Ye, J., N. Ma, T. L. Madden, and J. M. Ostell. 2013. IgBLAST: an immunoglobulin variable domain

sequence analysis tool. *Nucleic Acids Res.* 41.

42. Dunbar, J., and C. M. Deane. 2015. ANARCI: Antigen receptor numbering and receptor classification.

Bioinformatics btv552.

43. Lefranc, M.-P., C. Pommié, M. Ruiz, V. Giuducelli, E. Foulquier, L. Truong, V. Thouvenin-Contet, and

G. Lefranc. 2003. IMGT unique numbering fro immunoglobulin and T cell receptor variable domains and

Ig superfamily V-like domains. *Dev. Comp. Immunol.* 27: 55–77.

44. Eddy, S. R. 1995. Multiple alignment using hidden Markov models. *Proc. Int. Conf. Intell. Syst. Mol.*

Biol. 3: 114–120.

45. Shugay, M., O. V Britanova, E. M. Merzlyak, M. a Turchaninova, I. Z. Mamedov, T. R. Tuganbaev, D. a

Bolotin, D. B. Staroverov, E. V Putintseva, K. Plevova, C. Linnemann, D. Shagin, S. Pospisilova, S.

Lukyanov, T. N. Schumacher, and D. M. Chudakov. 2014. Towards error-free profiling of immune

repertoires. *Nat. Methods* 11: 653–5.

46. Banerjee, S., H. Shi, M. Banasik, H. Moon, W. Lees, Y. Qin, A. Harley, A. Shepherd, and M. W. Cho.

2017. Evaluation of a novel multi-immunogen vaccine strategy for targeting 4E10/10E8 neutralizing

epitopes on HIV-1 gp41 membrane proximal external region. *Virology* 505: 113–126.

47. Bashford-Rogers, R. J. M., A. L. Palser, B. J. Huntly, R. Rance, G. S. Vassiliou, G. A. Follows, and P.

Kellam. 2013. Network properties derived from deep sequencing of human B-cell receptor repertoires

delineate B-cell populations. *Genome Res.* 23: 1874–1884.

48. Bhiman, J. N., C. Anthony, N. A. Doria-Rose, O. Karimanzira, C. A. Schramm, T. Khoza, D. Kitchin, G.

Botha, J. Gorman, N. J. Garrett, S. S. A. Karim, L. Shapiro, C. Williamson, P. D. Kwong, J. R. Mascola, L.

Morris, and P. L. Moore. 2015. Viral variants that initiate and drive maturation of V1V2-directed HIV-1

broadly neutralizing antibodies. *Nat. Med.* 21: 1332–1336.

49. Bonsignori, M., T. Zhou, Z. Sheng, L. Chen, F. Gao, M. G. Joyce, G. Ozorowski, G. Y. Chuang, C. A.

Schramm, K. Wiehe, S. M. Alam, T. Bradley, M. A. Gladden, K. K. Hwang, S. Iyengar, A. Kumar, X. Lu, K.

Luo, M. C. Mangiapani, R. J. Parks, H. Song, P. Acharya, R. T. Bailer, A. Cao, A. Druz, I. S. Georgiev, Y. D.

Kwon, M. K. Louder, B. Zhang, A. Zheng, B. J. Hill, R. Kong, C. Soto, J. C. Mullikin, D. C. Douek, D. C.

Montefiori, M. A. Moody, G. M. Shaw, B. H. Hahn, G. Kelsoe, P. T. Hraber, B. T. Korber, S. D. Boyd, A. Z.

Fire, T. B. Kepler, L. Shapiro, A. B. Ward, J. R. Mascola, H. X. Liao, P. D. Kwong, and B. F. Haynes. 2016.

Maturation Pathway from Germline to Broad HIV-1 Neutralizer of a CD4-Mimic Antibody. *Cell* 165: 449–

463.

50. Collins, A. M., Y. Wang, K. M. Roskin, C. P. Marquis, K. J. L. Jackson, H. Schroeder, L. Cavacini, S.

Tonegawa, F. Alt, D. Baltimore, K. Jackson, M. Kidd, Y. Wang, A. Collins, M. Atkinson, M. Cowan, A.

Feeney, D. Granoff, P. Shackelford, S. Holmes, A. Lucas, B. Nadel, L. Liu, A. Lucas, L. Pappas, Y. Wang, K.

Jackson, B. Gaeta, W. Pomat, P. Siba, W. Sewell, A. Collins, C. Watson, C. Scheepers, M. Kidd, E. Houpt, K.

Breitbach, P. Wongprompitak, I. Steinmetz, G. Peltz, E. Marino, S. Grey, M. Potter, R. Riblet, C. Johnston,

M. Lefranc, I. Retter, K. Jackson, Y. Wang, A. Collins, J. Ye, N. Ma, T. Madden, J. Ostell, R. Edgar, V. Greiff,

U. Menzel, U. Haessler, S. Cook, S. Friedensohn, T. Khan, M. Pogson, I. Hellmann, S. Reddy, S. Boyd, I.

Retter, J. Meier, S. Lewis, D. Gadala-Maria, G. Yaari, M. Uduman, S. Kleinstein, B. Haines, C. Angeles, A.

Parmelee, P. McLean, P. Brodeur, J. Ye, A. Feeney, R. Riblet, K. Jackson, B. Gaeta, A. Collins, B. Blomberg,

W. Geckeler, M. Weigert, E. Jouvin-Marche, M. Morgado, P. Brodeur, R. Riblet, A. Collins, M. Ikutani, D.

Puiu, G. Buck, A. Nadkarni, B. Gaeta, K. Larimore, M. McCormick, H. Robins, P. Greenberg, K. Jackson, B.

Gaeta, W. Sewell, A. Collins, J. Glanville, H. Yang, C. Scott, M. Potter, L. Reidl, C. Kinoshita, and L. Steiner.

2015. The mouse antibody heavy chain repertoire is germline-focused and highly variable between

inbred strains. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 370: S41–S52.

51. Corcoran, M. M., G. E. Phad, N. V. Bernat, C. Stahl-Hennig, N. Sumida, M. A. A. Persson, M. Martin,

and G. B. K. Hedestam. 2016. Production of individualized v gene databases reveals high levels of

immunoglobulin genetic diversity. *Nat. Commun.* 7.

52. Cui, A., R. Di Niro, J. A. Vander Heiden, A. W. Briggs, K. Adams, T. Gilbert, K. C. O'Connor, F.

Vigneault, M. J. Shlomchik, and S. H. Kleinstein. 2016. A Model of Somatic Hypermutation Targeting in

Mice Based on High-Throughput Ig Sequencing Data. *J. Immunol.* 197: 3566–3574.

53. Fisher, C. R., H. J. Sutton, J. A. Kaczmarski, H. A. McNamara, B. Clifton, J. Mitchell, Y. Cai, J. N. Dups,

N. J. D'Arcy, M. Singh, A. Chuah, T. S. Peat, C. J. Jackson, and I. A. Cockburn. 2017. T-dependent B cell

responses to Plasmodium induce antibodies that form a high-avidity multivalent complex with the

circumsporozoite protein. *PLoS Pathog.* 13.

54. Gupta, N. T., K. D. Adams, A. W. Briggs, S. C. Timberlake, F. Vigneault, and S. H. Kleinstein. 2017.

Hierarchical Clustering Can Identify B Cell Clones with High Confidence in Ig Repertoire Sequencing Data.

J. Immunol. 198: 2489–2499.

55. Halliley, J. L., C. M. Tipton, J. Liesveld, A. F. Rosenberg, J. Darce, I. V Gregoretti, L. Popova, D.

Kaminiski, C. F. Fucile, I. Albizua, S. Kyu, K.-Y. Chiang, K. T. Bradley, R. Burack, M. Slifka, E. Hammarlund,

H. Wu, L. Zhao, E. E. Walsh, A. R. Falsey, T. D. Randall, W. C. Cheung, I. Sanz, and F. E.-H. Lee. 2015. Long-

Lived Plasma Cells Are Contained within the CD19(-)CD38(hi)CD138(+) Subset in Human Bone Marrow.

Immunity 43: 132–45.

56. Huang, J., B. H. Kang, E. Ishida, T. Zhou, T. Griesman, Z. Sheng, F. Wu, N. A. Doria-Rose, B. Zhang, K.

McKee, S. O'Dell, G. Y. Chuang, A. Druz, I. S. Georgiev, C. A. Schramm, A. Zheng, M. G. Joyce, M. Asokan,

A. Ransier, S. Darko, S. A. Migueles, R. T. Bailer, M. K. Louder, S. M. Alam, R. Parks, G. Kelsoe, T. Von

Holle, B. F. Haynes, D. C. Douek, V. Hirsch, M. S. Seaman, L. Shapiro, J. R. Mascola, P. D. Kwong, and M.

Connors. 2016. Identification of a CD4-Binding-Site Antibody to HIV that Evolved Near-Pan

Neutralization Breadth. *Immunity* 45: 1108–1121.

57. Jiang, N., J. He, J. a Weinstein, L. Penland, S. Sasaki, X.-S. He, C. L. Dekker, N.-Y. Zheng, M. Huang, M.

Sullivan, P. C. Wilson, H. B. Greenberg, M. M. Davis, D. S. Fisher, and S. R. Quake. 2013. Lineage structure

of the human antibody repertoire in response to influenza vaccination. *Sci. Transl. Med.* 5: 171ra19.

58. Joyce, M. G., A. K. Wheatley, P. V. Thomas, G. Y. Chuang, C. Soto, R. T. Bailer, A. Druz, I. S. Georgiev,

R. A. Gillespie, M. Kanekiyo, W. P. Kong, K. Leung, S. N. Narpala, M. S. Prabhakaran, E. S. Yang, B. Zhang,

Y. Zhang, M. Asokan, J. C. Boyington, T. Bylund, S. Darko, C. R. Lees, A. Ransier, C. H. Shen, L. Wang, J. R.

Whittle, X. Wu, H. M. Yassine, C. Santos, Y. Matsuoka, Y. Tsybovsky, U. Baxa, J. C. Mullikin, K. Subbarao,

D. C. Douek, B. S. Graham, R. A. Koup, J. E. Ledgerwood, M. Roederer, L. Shapiro, P. D. Kwong, J. R.

Mascola, and A. B. McDermott. 2016. Vaccine-Induced Antibodies that Neutralize Group 1 and Group 2

Influenza A Viruses. *Cell* 166: 609–623.

59. Khan, T. A., S. Friedensohn, A. R. G. de Vries, J. Straszewski, H.-J. Ruscheweyh, and S. T. Reddy. 2016.

Accurate and predictive antibody repertoire profiling by molecular amplification fingerprinting. *Sci. Adv.*

2: e1501371–e1501371.

60. Levin, M., J. J. King, J. Glanville, K. J. L. Jackson, T. J. Looney, R. A. Hoh, A. Mari, M. Andersson, L.

Greiff, A. Z. Fire, S. D. Boyd, and M. Ohlin. 2016. Persistence and evolution of allergen-specific IgE

repertoires during subcutaneous specific immunotherapy. *J. Allergy Clin. Immunol.* 137: 1535–1544.

61. Levin, M., F. Levander, R. Palmason, L. Greiff, and M. Ohlin. 2017. Antibody-encoding repertoires of

bone marrow and peripheral blood—a focus on IgE. *J. Allergy Clin. Immunol.* 139: 1026–1030.

62. Li, X., X. Duan, K. Yang, W. Zhang, C. Zhang, L. Fu, Z. Ren, C. Wang, J. Wu, R. Lu, Y. Ye, M. He, C. Nie,

N. Yang, J. Wang, H. Yang, X. Liu, and W. Tan. 2016. Comparative analysis of immune repertoires

between bactrian Camel's conventional and heavy-chain antibodies. *PLoS One* 11.

63. Liao, H. X., R. Lynch, T. Zhou, F. Gao, S. Munir Alam, S. D. Boyd, A. Z. Fire, K. M. Roskin, C. A.

Schramm, Z. Zhang, J. Zhu, L. Shapiro, J. C. Mullikin, S. Gnanakaran, P. Hraber, K. Wiehe, G. Kelsoe, G.

Yang, S. M. Xia, D. C. Montefiori, R. Parks, K. E. Lloyd, R. M. Scearce, K. A. Soderberg, M. Cohen, G.

Kamanga, M. K. Louder, L. M. Tran, Y. Chen, F. Cai, S. Chen, S. Moquin, X. Du, M. Gordon Joyce, S.

Srivatsan, B. Zhang, A. Zheng, G. M. Shaw, B. H. Hahn, T. B. Kepler, B. T. M. Korber, P. D. Kwong, J. R.

Mascola, B. F. Haynes, J. Becker, B. Benjamin, R. Blakesley, G. Bouffard, S. Brooks, H. Coleman, M.

Dekhtyar, M. Gregory, X. Guan, J. Gupta, J. Han, A. Hargrove, S. L. Ho, T. Johnson, R. Legaspi, S. Lovett, Q.

Maduro, C. Masiello, B. Maskeri, J. McDowell, C. Montemayor, J. Mulliki, M. Park, N. Riebow, K.

Schandler, B. Schmidt, C. Sison, M. Stantripop, J. Thomas, P. Thomas, M. Vemulapalli, and A. Young.

2013. Co-evolution of a broadly neutralizing HIV-1 antibody and founder virus. *Nature* 496: 469–476.

64. Lindner, C., I. Thomsen, B. Wahl, M. Ugur, M. K. Sethi, M. Friedrichsen, A. Smoczek, S. Ott, U.

Baumann, S. Suerbaum, S. Schreiber, A. Bleich, V. Gaboriau-Routhiau, N. Cerf-Bensussan, H. Hazanov, R.

Mehr, P. Boysen, P. Rosenstiel, and O. Pabst. 2015. Diversification of memory B cells drives the

continuous adaptation of secretory antibodies to gut microbiota. *Nat. Immunol.* 16: 880–888.

65. Meng, W., B. Zhang, G. W. Schwartz, A. M. Rosenfeld, D. Ren, J. J. C. Thome, D. J. Carpenter, N.

Matsuoka, H. Lerner, A. L. Friedman, T. Granot, D. L. Farber, M. J. Shlomchik, U. Hershberg, and E. T.

Luning Prak. 2017. An atlas of B-cell clonal distribution in the human body. *Nat. Biotechnol.* 35: 879–886.

66. Menzel, U., V. Greiff, T. A. Khan, U. Haessler, I. Hellmann, S. Friedensohn, S. C. Cook, M. Pogson, and

S. T. Reddy. 2014. Comprehensive evaluation and optimization of amplicon library preparation methods

for high-throughput antibody sequencing. *PLoS One* 9.

67. Mroczek, E. S., G. C. Ippolito, T. Rogosch, K. H. Hoi, T. A. Hwangpo, M. G. Brand, Y. Zhuang, C. R. Liu,

D. A. Schneider, M. Zemlin, E. E. Brown, G. Georgiou, and H. W. Schroeder. 2014. Differences in the

composition of the human antibody repertoire by B cell subsets in the blood. *Front Immunol* 5: 96.

68. Ota, M., B. H. Duong, A. Torkamani, C. M. Doyle, A. L. Gavin, T. Ota, and D. Nemazee. 2010.

Regulation of the B cell receptor repertoire and self-reactivity by BAFF. *J. Immunol.* 185: 4128–36.

69. Palanichamy, A., L. Apeltsin, T. C. Kuo, M. Sirota, S. Wang, S. J. Pitts, P. D. Sundar, D. Telman, L. Z.

Zhao, M. Derstine, A. Abounasr, S. L. Hauser, and H. C. Von B??dingen. 2014. Immunoglobulin class-

switched B cells form an active immune axis between CNS and periphery in multiple sclerosis. *Sci. Transl.*

Med. 6.

70. Prohaska, T. A., X. Que, C. J. Diehl, S. Hendrikx, M. W. Chang, K. Jepsen, C. K. Glass, C. Benner, and J.

L. Witztum. 2018. Massively Parallel Sequencing of Peritoneal and Splenic B Cell Repertoires Highlights

Unique Properties of B-1 Cell Antibodies. *J. Immunol.* 200: 1702–1717.

71. Rubelt, F., C. R. Bolen, H. M. McGuire, J. A. V. Heiden, D. Gadala-Maria, M. Levin, G. M. Euskirchen,

M. R. Mamedov, G. E. Swan, C. L. Dekker, L. G. Cowell, S. H. Kleinstein, and M. M. Davis. 2016. Individual

heritable differences result in unique cell lymphocyte receptor repertoires of naïve and antigen-

experienced cells. *Nat. Commun.* 7.

72. Zhu, J., G. Ofek, Y. Yang, B. Zhang, M. K. Louder, G. Lu, K. McKee, M. Pancera, J. Skinner, Z. Zhang, R.

Parks, J. Eudailey, K. E. Lloyd, J. Blinn, S. M. Alam, B. F. Haynes, M. Simek, D. R. Burton, W. C. Koff, N. C. S.

NISC Comparative Sequencing Program, J. C. Mullikin, J. R. Mascola, L. Shapiro, and P. D. Kwong. 2013.

Mining the antibodyome for HIV-1-neutralizing antibodies with next-generation sequencing and

phylogenetic pairing of heavy/light chains. *Proc. Natl. Acad. Sci. U. S. A.* 110: 6470–5.

73. Stern, J. N. H., G. Yaari, J. A. Vander Heiden, G. Church, W. F. Donahue, R. Q. Hintzen, A. J. Huttner, J.

D. Laman, R. M. Nagra, A. Nylander, D. Pitt, S. Ramanan, B. A. Siddiqui, F. Vigneault, S. H. Kleinstein, D. A.

Hafler, and K. C. O'Connor. 2014. B cells populating the multiple sclerosis brain mature in the draining

cervical lymph nodes. *Sci. Transl. Med.* 6.

74. Sundling, C., Z. Zhang, G. E. Phad, Z. Sheng, Y. Wang, J. R. Mascola, Y. Li, R. T. Wyatt, L. Shapiro, and

G. B. Karlsson Hedestam. 2014. Single-Cell and Deep Sequencing of IgG-Switched Macaque B Cells

Reveal a Diverse Ig Repertoire following Immunization. *J. Immunol.* 192: 3637–3644.

75. Tipton, C. M., C. F. Fucile, J. Darce, A. Chida, T. Ichikawa, I. Gregoretti, S. Schieferl, J. Hom, S. Jenks, R.

J. Feldman, R. Mehr, C. Wei, F. E. H. Lee, W. C. Cheung, A. F. Rosenberg, and I. Sanz. 2015. Diversity,

cellular origin and autoreactivity of antibody-secreting cell population expansions in acute systemic

lupus erythematosus. *Nat. Immunol.* 16: 755–765.

76. Turchaninova, M. A., A. Davydov, O. V Britanova, M. Shugay, V. Bikos, E. S. Egorov, V. I. Kirgizova, E.

M. Merzlyak, D. B. Staroverov, D. A. Bolotin, I. Z. Mamedov, M. Izraelson, M. D. Logacheva, O. Kladova,

K. Plevova, S. Pospisilova, and D. M. Chudakov. 2016. High-quality full-length immunoglobulin profiling

with unique molecular barcoding. *Nat. Protoc.* 11: 1599–1616.

77. Vander Heiden, J. A., P. Stathopoulos, J. Q. Zhou, L. Chen, T. J. Gilbert, C. R. Bolen, R. J. Barohn, M. M.

Dimachkie, E. Ciafaloni, T. J. Broering, F. Vigneault, R. J. Nowak, S. H. Kleinstein, and K. C. O'Connor.

2017. Dysregulation of B Cell Repertoire Formation in Myasthenia Gravis Patients Revealed through

Deep Sequencing. *J. Immunol.* 198: 1460–1473.

78. VanDuijn, M. M., L. J. Dekker, W. F. J. van IJcken, P. A. E. Sillevis Smitt, and T. M. Luider. 2017.

Immune repertoire after immunization as seen by next-generation sequencing and proteomics. *Front.*

Immunol. 8.

79. Vergani, S., I. Korsunsky, A. N. Mazzarello, G. Ferrer, N. Chiorazzi, and D. Bagnara. 2017. Novel

method for high-throughput full-length IGHV-D-J sequencing of the immune repertoire from bulk B-cells

with single-cell resolution. *Front. Immunol.* 8.

80. Wesemann, D. R., A. J. Portuguese, R. M. Meyers, M. P. Gallagher, K. Cluff-Jones, J. M. Magee, R. A.

Panchakshari, S. J. Rodig, T. B. Kepler, and F. W. Alt. 2013. Microbial colonization influences early B-

lineage development in the gut lamina propria. *Nature* 501: 112–115.

81. Wu, X., T. Zhou, J. Zhu, B. Zhang, I. Georgiev, C. Wang, X. Chen, N. S. Longo, M. Louder, K. McKee, S.

O'Dell, S. Perfetto, S. D. Schmidt, W. Shi, L. Wu, Y. Yang, Z.-Y. Yang, Z. Yang, Z. Zhang, M. Bonsignori, J. A.

Crump, S. H. Kapiga, N. E. Sam, B. F. Haynes, M. Simek, D. R. Burton, W. C. Koff, N. A. Doria-Rose, M.

Connors, N. C. S. NISC Comparative Sequencing Program, J. C. Mullikin, G. J. Nabel, M. Roederer, L.

Shapiro, P. D. Kwong, and J. R. Mascola. 2011. Focused evolution of HIV-1 neutralizing antibodies

revealed by structures and deep sequencing. *Science* 333: 1593–602.

82. Wu, X., Z. Zhang, C. A. Schramm, M. G. Joyce, Y. Do Kwon, T. Zhou, Z. Sheng, B. Zhang, S. O'Dell, K.

McKee, I. S. Georgiev, G. Y. Chuang, N. S. Longo, R. M. Lynch, K. O. Saunders, C. Soto, S. Srivatsan, Y.

Yang, R. T. Bailer, M. K. Louder, J. C. Mullikin, M. Connors, P. D. Kwong, J. R. Mascola, and L. Shapiro.

2015. Maturation and diversity of the VRC01-antibody lineage over 15 years of chronic HIV-1 infection.

Cell 161: 480–485.

83. Wu, Y. C. B., L. K. James, J. A. Vander Heiden, M. Uduman, S. R. Durham, S. H. Kleinstein, D. Kipling,

and H. J. Gould. 2014. Influence of seasonal exposure to grass pollen on local and peripheral blood IgE

repertoires in patients with allergic rhinitis. *J. Allergy Clin. Immunol.* 134: 604–612.

84. Zhou, T., R. M. Lynch, L. Chen, P. Acharya, X. Wu, N. A. Doria-Rose, M. G. Joyce, D. Lingwood, C. Soto,

R. T. Bailer, M. J. Ernandes, R. Kong, N. S. Longo, M. K. Louder, K. McKee, S. O'Dell, S. D. Schmidt, L. Tran,

Z. Yang, A. Druz, T. S. Luongo, S. Moquin, S. Srivatsan, Y. Yang, B. Zhang, A. Zheng, M. Pancera, T. Kirys, I.

S. Georgiev, T. Gindin, H. P. Peng, A. S. Yang, J. C. Mullikin, M. D. Gray, L. Stamatatos, D. R. Burton, W. C.

Koff, M. S. Cohen, B. F. Haynes, J. P. Casazza, M. Connors, D. Corti, A. Lanzavecchia, Q. J. Sattentau, R. A.

Weiss, A. P. West, P. J. Bjorkman, J. F. Scheid, M. C. Nussenzweig, L. Shapiro, J. R. Mascola, and P. D.

Kwong. 2015. Structural repertoire of HIV-1-neutralizing antibodies targeting the CD4 supersite in 14

donors. *Cell* 161: 1280–1292.

85. Zhu, J., S. O'Dell, G. Ofek, M. Pancera, X. Wu, B. Zhang, Z. Zhang, J. C. Mullikin, M. Simek, D. R.

Burton, W. C. Koff, L. Shapiro, J. R. Mascola, and P. D. Kwong. 2012. Somatic populations of PGT135-137

HIV-1-neutralizing antibodies identified by 454 pyrosequencing and bioinformatics. *Front. Microbiol.* 3.

86. Zhu, J., X. Wu, B. Zhang, K. McKee, S. O'Dell, C. Soto, T. Zhou, J. P. Casazza, J. C. Mullikin, P. D. Kwong,

J. R. Mascola, L. Shapiro, J. Becker, B. Benjamin, R. Blakesley, G. Bouffard, S. Brooks, H. Coleman, M.

Dekhtyar, M. Gregory, X. Guan, J. Gupta, J. Han, A. Hargrove, S. -l. Ho, T. Johnson, R. Legaspi, S. Lovett,

Q. Maduro, C. Masiello, B. Maskeri, J. McDowell, C. Montemayor, J. Mullikin, M. Park, N. Riebow, K.

Schandler, B. Schmidt, C. Sison, M. Stantripop, J. Thomas, P. Thomas, M. Vemulapalli, and A. Young.

2013. De novo identification of VRC01 class HIV-1-neutralizing antibodies by next-generation sequencing

of B-cell transcripts. *Proc. Natl. Acad. Sci.* 110: E4088–E4097.

87. Schroeder Jr, H. W. 2006. Similarity and divergence in the development and expression of the mouse

and human antibody repertoires. *Dev. Comp. Immunol.* 30: 119–135.

88. Zemlin, M., M. Klinger, J. Link, C. Zemlin, K. Bauer, J. A. Engler, H. W. Schroeder, and P. M. Kirkham.

2003. Expressed murine and human CDR-H3 intervals of equal length exhibit distinct repertoires that

differ in their amino acid composition and predicted range of structures. *J. Mol. Biol.* 334: 733–749.

89. Olimpieri, P. P., P. Marcatili, and A. Tramontano. 2014. Tabhu: Tools for antibody humanization.

Bioinformatics 31: 434–435.

586