The development of individual differences in cooperative behaviour: maternal

glucocorticoid hormones alter helping behaviour of offspring in wild meerkats

Ben Dantzer[1,2,3]*, Constance Dubuc[1,2], Ines Braga Goncalves[2,4], Dominic L. Cram[1,2],

Nigel C. Bennett[5], Andre Ganswindt[5,6], Michael Heistermann[7], Chris Duncan[2], David

Gaynor[2,5], Tim H. Clutton-Brock[1,2,5]

[1]Department of Zoology, University of Cambridge, Cambridge, United Kingdom

[2]Kalahari Meerkat Project, Kuruman River Reserve, Northern Cape, South Africa

[3]Department of Psychology, Department of Ecology and Evolutionary Biology,

University of Michigan, Ann Arbor, MI 48109, USA

[4]Department of Evolutionary Biology and Environmental Studies, Animal Behaviour,

University of Zurich, Switzerland

[5]Mammal Research Institute, Department of Zoology and Entomology, University of

Pretoria, Pretoria, South Africa

[6]Endocrine Research Laboratory, Department of Anatomy and Physiology, University of

Pretoria, 0110, Onderstepoort, South Africa

[7]Endocrinology Laboratory, German Primate Center, Leibniz Institute for Primate

Research, Kellnerweg 4, D-37077 Göttingen, Germany

*Corresponding Author: dantzer@umich.edu

Abstract

The phenotype of parents can have long-lasting effects on the development of offspring as well as on their behaviour, physiology, and morphology as adults. In some cases, these changes may increase offspring fitness but, in others, they can elevate parental fitness at a cost to the fitness of their offspring. We show that in Kalahari meerkats (*Suricata suricatta*), the circulating glucocorticoid (GC) hormones of pregnant females affect the growth and cooperative behaviour of their offspring. We performed a 3-year experiment in wild meerkats to test the hypothesis that GC-mediated maternal effects reduce the potential for offspring to reproduce directly and therefore cause them to exhibit more cooperative behaviour. Daughters (but not sons) born to mothers treated with cortisol during pregnancy grew more slowly early in life and exhibited significantly more of two types of cooperative behaviour (pup rearing and feeding) once they were adults compared to offspring from control mothers. They also had lower measures of GCs as they aged, which could explain the observed increases in cooperative behaviour. Because early life growth is a crucial determinant of fitness in female meerkats, our results indicate that GC-mediated maternal effects may reduce the fitness of offspring, but may elevate parental fitness as a consequence of increasing the cooperative behaviour of their daughters.

Keywords: Cooperation, Early life adversity, Glucocorticoids, Growth, Maternal stress

Introduction

Parental effects are a mechanism of trans-generational phenotypic plasticity that occurs when the parental phenotype or parental environment modifies offspring characteristics (1). Parental effects can increase the survival or reproduction of offspring, thereby elevating the direct fitness of both offspring and parents (2-6). Alternatively, parental effects can increase parental fitness, but at some cost to the fitness of their offspring (7-8) – a process regarded as a type of parental manipulation (9-12) or 'selfish parental effect' (13). For example, in mammals, the optimal birth weight or litter size often differs between mothers and offspring (14) and pregnant females experiencing stressful environments may reallocate resources away from offspring and towards themselves, so that their offspring are smaller or grow more slowly before weaning (15). Despite these observations, it has been suggested that selfish parental effects may be rare and unstable because selection would be expected to favour the evolution of resistance mechanisms in offspring (7, 11, 13, 16, 17).

Selfish parental effects may in fact be more likely in cooperatively breeding species where philopatric offspring (subordinates) help to rear the subsequent offspring of their parents or other close relatives. This could be especially likely under low food or high stress conditions as parents may gain substantial direct fitness benefits from delaying the development of their offspring if this causes them to invest in alloparental care directed at the parent's subsequent offspring (9-10). In addition, the costs of selfish parental effects to offspring could be reduced in these circumstances, as offspring will gain indirect fitness benefits by contributing to raising the subsequent offspring of their parents (18). For example, laboratory studies of eusocial insects suggest the possibility

that selection will favour the evolution of alleles that enable mothers to increase the

helping behaviour of their offspring while simultaneously reducing their capabilities of

reproducing on their own (19-20; but see 21).

To date, empirical field tests of how parental effects shape the helping behaviour

of offspring are rare (23) and studies of selfish parental effects have mostly focused on

non-social species (13, 24). Here, we report the results of experiments designed to test

the hypothesis that elevated maternal glucocorticoid levels (GCs) reduce the potential

for offspring to have direct reproductive opportunities and causes them to exhibit more

cooperative behaviour. In a 3-year field study, we experimentally elevated maternal

GCs by treating pregnant dominant female meerkats with cortisol and tracking the

growth, stress physiology, and cooperative behaviour of their offspring from birth until

~18 months of age, compared to those from control litters. We manipulated maternal

GCs because they are known to cause mothers to reallocate energy away from

offspring and towards themselves (15), indicating that they may function as a mediator

of selfish maternal effects. Changes in maternal GCs have also previously been shown

to delay the dispersal of offspring as well as influence the parental care behaviour of

offspring (25-26), both traits that are important in cooperative breeders where philopatric

offspring exhibit alloparental care behaviour towards juveniles.

To identify if the exposure of mothers to heightened GCs reduced reproductive

success of their offspring, we examined if offspring from mothers treated with cortisol

during pregnancy grew more slowly early in life. In meerkats, the rate of early life growth

and body mass is closely linked to future direct fitness through its effects on survival,

foraging success, adult body mass (27-29), as well as the probability of acquiring

dominance (30-31) and other direct reproductive opportunities (32). As elevated

exposure to maternal GCs in some mammals may reduce offspring size and growth

early in life (15), we predicted that offspring from mothers treated with cortisol during

pregnancy would be smaller or grow more slowly early in life. Because the rate of early

life growth is predictive of future direct fitness in meerkats (27-32), we predicted that if

offspring from mothers treated with cortisol did grow more slowly, they would

consequently invest more in indirect fitness opportunities by contributing more to

cooperative activities than controls.

Secondly, we determined if offspring from mothers treated with cortisol during

pregnancy subsequently increased their contributions to two types of cooperative

behaviours: pup rearing ("babysitting": 33) and food provisioning during the period when

the pups are foraging with their natal group, but are not yet nutritionally independent

("pup feeding": 34). We chose these two behaviours as they appear to be most costly

from an energetic perspective (35) and are most closely tied to the probability of parents

successfully rearing offspring. If offspring from mothers treated with cortisol during

pregnancy exhibit more of either of these two types of alloparental care, this should

increase both parental direct fitness (the number of offspring that they subsequently

produce) and the indirect fitness of offspring, because subsequent offspring that receive

more alloparental care should grow faster or have higher early life survival (27, 32, 36,

37). Previous studies in meerkats show that offspring with more helpers or those that

receive more alloparental care grow faster or have early life survival (27, 32, 36, 37).

To assess the mechanism by which elevated exposure to maternal stress may

affect the alloparental care behaviour of offspring, we repeatedly measured plasma

cortisol and faecal glucocorticoid metabolite (fGCM) concentrations of offspring from

when they were approximately 1 to 18 months of age to identify how our manipulations

affected their neuroendocrine stress axes (GC output). Elevated maternal GCs can

cause long-term changes in the neuroendocrine stress axis of offspring (38) and

elevated activity of the neuroendocrine stress axis in meerkats can reduce their

contributions to alloparental care (39). We therefore predicted that if offspring born to

mothers treated with cortisol during pregnancy exhibited more alloparental care

behaviour compared to controls, they would also have reduced plasma cortisol and

fGCM concentrations.

Methods

122   *Study site & basic data collection*

We studied free-living meerkats at the Kuruman River Reserve (26° 58' S, 21°

49' E) in the Northern Cape, South Africa from 2014-2017. Individuals were marked

uniquely with PIT tags (Identipet®, Johannesburg, South Africa) as well as dye marks so

that they could be identified. Groups were visited for ~4-8 hours per day ~4-6 times per

week throughout each year of study and sometimes more frequently such as when

there were pups being babysat. Groups were visited at sunrise before meerkats

emerged from their sleeping burrow. After all the meerkats had emerged, but prior to

when they started going foraging, we counted the total number of meerkats in the group

(to get estimates of group size) and recorded which individuals were present (using their

unique combinations of dye marks). We recorded their body mass on a portable

balance each morning before foraging, 2-4 hrs after foraging was initiated, and

immediately prior to when foraging ended (40). These measures of body mass provided

our estimates of growth, body mass, and foraging success that are used in our analyses

described below.

137   *Experimental manipulations of dominant females*

138   Dominant females in each group were identified via behavioural observations

(41). The pregnancy status of dominant females was determined visually (distended

abdomen) as well as noting a constant increase in their body mass. Dominant females

were treated with either a cortisol solution or a control oil vehicle when they were

pregnant by feeding them food containing one of these two treatments. We initially

offered experimental animals hard boiled eggs with added cortisol but found that they

rejected all foods that contained added cortisol with the exception of scorpions. We

consequently fed experimental females with cortisol (10 mg/kg of hydrocortisone, Sigma

H4126), that were dissolved in 100 μl of 100% coconut oil and injected into a dead

scorpion (*Opistophthalmus* spp.). Control females were fed a dead scorpion that was

injected with 100 μl of 100% coconut oil. A previous study using the same protocol

showed that meerkats that were fed cortisol had significantly higher plasma cortisol and

fGCM concentrations than control animals and these increases were within a

biologically relevant range (39). This indicated that our treatment causes the exogenous

glucocorticoids that we feed the meerkats to enter their bloodstream and leads to

sustained increases in their circulating glucocorticoid concentrations.

154   Females were randomly allocated to the treatments. Across the three years of

this study, we produced a total of 13 cortisol-treated litters from 10 females and 7

control litters produced by 6 females (Table 1). Three of the females experienced both

the control and cortisol treatments at different time points of the experiment, whereas

one female experienced the control treatment once and the cortisol treatments twice.

For these latter females treated twice, the order of treatments was randomly selected.

We conducted these experiments over the course of three years: 13 litters from 10

females in 2014 (April-December 2014, 3 litters aborted), 5 litters from 5 females in

2015 (February-July 2015, 1 litter aborted), and 3 litters from 3 females in 2016 (July

2016). Three cortisol-treated mothers and one control mother aborted their offspring

prior to birth and were excluded from any analyses except for assessing differences in

the frequency of abortion between control and cortisol-treated mothers (Table 2). This

provided final sample sizes of 31 pups from 10 litters from 9 cortisol-treated females

and 25 pups from 6 litters from 6 control females (Table 1).

We aimed to experimentally increase the glucocorticoid concentrations of

pregnant dominant females from when they were first confirmed to be pregnant (second

half of gestation) until parturition. Gestation in meerkats is ~70 days so we aimed to

treat them with glucocorticoids from approximately 35-70 d during gestation. In reality,

females that successfully produced a litter where pups emerged from the natal burrow

were treated with cortisol for 12-36 days prior to birth (n=10 litters from 9 females, mean

= 23.7 d, median = 23.5 d), whereas controls were fed for 12-58 d prior to birth (n=6

litters from 6 females, mean = 30 d, median = 20.5 d). Although controls were treated

for slightly longer, there was no significant difference in treatment duration between

control and cortisol-treated females (general linear model, t = 1.05, P = 0.31).

To provide an additional comparison group to investigate how our treatments (fed

during pregnancy or fed cortisol during pregnancy) affected offspring survival, growth,

and cooperative behaviour, we also monitored these traits in offspring produced by

dominant females that were untreated during pregnancy (n = 52 litters from 21 dominant

females, Table 1). These females were not fed or treated with cortisol (hereafter,

"untreated mothers"). For our analyses of how the treatments affected offspring survival

and growth, the untreated offspring were those from litters produced by dominant

females in other meerkat groups in our same study area and were born during our

study. We assessed the contributions of offspring from mothers treated with cortisol

during pregnancy to two cooperative behaviours (babysitting and pup feeding)

compared to those from control mothers, but also to other group members from

untreated mothers. We did not have data from offspring from untreated mothers when

we assessed how our treatments affected their plasma cortisol or fGCM concentrations.

191   *Quantifying early life growth of offspring*

Meerkat pups typically first emerge from their natal burrow approximately 21-30 d

after birth. Meerkat groups and dominant females were monitored daily around the

estimated date of parturition and birth dates were estimated according to the change in

the physical appearance of the dominant female, a large drop in body mass overnight,

and group members exhibiting babysitting behaviour at the sleeping burrow. Burrows

containing pups were monitored each day and, when pups emerged, they were uniquely

marked by trimming small sections of hair before permanent PIT tags could be applied.

Pups were weighed each time we visited the groups on a portable balance in the

morning after group members emerged from their sleeping burrow (as above).

201   *Quantifying cooperative behaviour of offspring*

We measured the babysitting (controls: 195-655 d; cortisol: 184-655 d; untreated:

155-655 d) and pup feeding (controls: 220-635 d; cortisol: 184-655 d; untreated: 155-

626 d) contributions of offspring from cortisol-treated and control mothers when they

were >6 months of age until death or disappearance. We visited sleeping burrows

containing pups every day in the morning and recorded the identity of the attending

babysitters. As we have done previously (33, 36, 39, 42), we calculated relative

babysitting contributions of each individual meerkat for each litter by dividing the total

number of days an individual babysat a litter over the total number of days that this

specific litter had a babysitter. Pup feeding behaviour for each pup produced by the

dominant females in the different treatment groups was estimated using *ad libitum*

sampling (34, 39). When the social group contained pups (up to 90 d of age), we

recorded all pup-feeding events from all individuals, which are visually and acoustically

conspicuous to observers (43). We then used these data to estimate the proportion of

pup-feeding events exhibited by an individual relative to all others in the group (i.e.,

relative pup feeding). Because the total amount of time devoted to the *ad libitum*

recording sessions varied, we corrected for variation in observation time (see below).

218   *Quantifying plasma cortisol concentrations from offspring*

We obtained plasma samples from offspring from cortisol-treated and control

mothers approximately every 3 months from first emergence from the burrow (~1

month) until ~18 months of age (controls: 20-548 d; cortisol-treated: 25-559 d). Capture

and blood processing procedures are described elsewhere (44-45). The amount of time

it took to obtain the blood samples varied (median = 10.6 min, SD = 7.2 min), but we

included co-variates for sampling time and sampling time^2 to control for effects of

sampling time (described in 45). We measured total plasma cortisol concentrations

using a previously validated assay (Coat-a-Count, Siemens Diagnostic Products

Corporation, Los Angeles, USA: validation described in 44). The sensitivity of the assay

was 1.9 ng/ml and cross-reactivity to other hormones was 76% with prednisolone,

11.4% with 11-deoxycortisol, 2.3% with prednisone and <1% with aldosterone,

corticosterone, cortisone, oestriol, estrone and pregnenolone. Intra-assay coefficient of

variation (CV) was 7% (n = 20 samples). Inter-assay CV for a low control (78.5 ± 6.3

ng/ml n = 5 assays) was 8% and 2.8% for a high control (187 ± 5.3 ng/ml, n = 5

assays).

234   *Quantifying fGCM concentrations from offspring*

We collected faecal samples from offspring of cortisol-treated and control

mothers opportunistically during behavioural observations over the course of the study

(controls: 25-356 d; cortisol-treated: 32-326 d). Faecal samples were processed as

described previously using a methanol solution to extract fGCMs for analysis (46-47).

Immunoreactive fGCM concentrations were determined using a group-specific enzyme

immunoassay measuring cortisol metabolites with a 5β-3α,11β-diol-structure (11β-

hydroxyetiocholanolone), already validated and established for monitoring fGCM

alterations in meerkats (47). Faecal GCMs measured reflect average adrenal cortisol

production over the previous ~24 to 48 hr period (47). Detailed assay characteristics,

including full descriptions of the assay components and cross-reactivities, are found

elsewhere (48). The sensitivity of the assay was 1.2 ng/g dry weight and intra-assay CV

determined by repeated measurements of high and low value quality controls were

6.9% and 7.4% and inter-assay CV values were 11.5% and 15.9% (n = 29 assays),

respectively.

249   *Statistical analyses*

We used generalized (binomial errors) or linear mixed-effects models (LMMs) to

examine how our treatments affected the probability that the litter was aborted, litter size

and sex ratio at emergence from the burrow, and the proportion of the litter that survived

to emergence from the burrow, independence (~90 d of age: 29), and 6 or 12 months of

age. We focused on addressing whether the offspring from cortisol-treated mothers

differed from control or untreated mothers. These models included a fixed effect for date

of birth of the litter and random intercept terms for dominant female identity and year (as

the experiments were conducted over 3 years). None of the GLMMs were

overdispersed (Table 2).

We used a LMM to investigate how the maternal treatments affected offspring

growth from first emergence from their natal burrow (~1 month) to 3 months of age

when the pups are typically foraging independently (29, 34). Morning body mass (in

grams) was the response variable with the fixed effects of maternal treatment (cortisol-

treated, control, or untreated), pup sex, pup age, litter size at burrow emergence, first

measure of body mass when the pups first emerged from the burrow (to control for

possible differences in age or development when they entered our study population),

group size, group size^2, total rainfall in the previous 60 days, two measures of

seasonality (sine and co-sine functions of day of weight measure: see 40), and two

three-way interactions between sex, treatment, age or age^2. Group size was defined as

the average number of subordinate meerkats >6 months of age in the group during the

entire period of offspring growth. Random intercept terms for year and the identity of the

individual nested in litter, nested in dominant female identity, nested in group were also

included in this model. Fixed and random effects included in these models were based

upon previous studies investigating meerkat body mass and/or growth from 1-3 months

(28, 31, 40). To prevent any issues associated with selective disappearance of specific

individuals, only individuals that survived to 90 d were included in these analyses.

We assessed how the treatments affected the relative babysitting and pup

feeding contributions of subordinates when they were >6 months (as they rarely do

alloparental care behaviour when <6 months: 36) from cortisol-treated, control, and

untreated mothers. Relative babysitting and pup feeding contributions are defined as

the proportion of babysitting or pup feeding contributions exhibited by a specific

individual compared to the total number of babysitting or pup feeding contributions for

that litter exhibited by all individuals in the group that were >6 months of age at the time

of the birth of the litter (36, 39, 42). In these generalized linear mixed-effects models

(GLMM, binomial errors), we included a three-way interaction between treatment, sex,

and age of the subordinate to assess if the effects of the treatments on babysitting or

pup feeding varied according to the sex or age of the subordinate, as contributions to

cooperative behaviour in meerkats are known to vary according to subordinate sex and

age (36). To account for differences in observation time, we included a co-variate for the

number of days the litter was babysat (babysitting length) and the number of days the

subordinate was observed in the group during babysitting as well as the total time spent

observing the group during pup feeding (observation time). We included a range of co-

variates (see Tables 4-6) that have been previously documented to affect relative

contributions to babysitting and pup feeding, including age, foraging success, body

mass, and group size (34-36, 42, 49; 50). Group size was defined as the average

number of subordinate meerkats >6 months of age in the group while the litter was

being babysat or pup fed. Foraging success was defined as the average weight gained

per hour estimated as the change in body mass from morning weight to evening weight

over the total number of hours that had elapsed since those two weights (45).

Relatedness between the subordinate and the litter being babysat was not included as it

has not been shown to impact babysitting or pup feeding contributions (27, 42) and

nearly all of the litters in our dataset were produced by the mother or full sibling of the

subordinate. Random intercept terms for year and the identity of the individual, and litter

being babysat or pup fed were nested within the group where the litter was being

babysat or pup fed. Overdispersion was not an issue for our GLMM for babysitting as

indicated by the goodness of fit test (Pearson χ^2 = 147.1, df = 154, P = 0.64, using

package aods3: 51) but our GLMM for pup feeding was initially overdispersed (Pearson

χ^2 = 310, df=165, P < 0.0001) so we included an observation level random intercept

term.

We used two separate LMMs to assess how our manipulations affected plasma

cortisol and fGCMs in offspring from cortisol-treated and control mothers (we did not

have these data from offspring from untreated mothers). Each model included fixed

effects for maternal treatment, pup sex and age, time of day and year that the sample

was acquired (2014 or 2015), and random intercept terms for identity of individual

nested in their birth litter and group. In the model for plasma cortisol concentrations, we

also included a linear and second order fixed effect for the time it took to acquire the

blood sample to control for any variation in plasma cortisol concentrations due to

restraint stress (45). Year was included as a fixed effect because we only had samples

from two separate years. We included covariates associated with the individual meerkat

and weather or social group characteristics that are known to affect plasma cortisol (45)

or fGCM (47) concentrations (see Tables 6-7).

We used R (version 3.4.3: 52) for all of our statistical analyses. R package lme4

(version 1.1-14: 53) was used for LMMs and P values were estimated using lmerTest

(version 2.0-33: 54). A graphical approach was used to confirm normality and

homoscedasticity of residuals and to confirm there were no observations with high

leverage (55). Collinearity among predictor variables included in our models was

assessed by calculating variance inflation factors (55) or generalized variance inflation

factors (for variables that had a second order term or those included in an interaction:

56). Collinearity was not a problem as indicated by our variance inflation factors (VIFs)

as VIFs or generalized VIFs were less than ~4 for all variables. In our model for how our

treatments affected offspring growth (Table 3), the generalized VIF for the two

measures of seasonality (sine and co-sine functions of day of weight measure) were <6

but these two variables were included a priori given their previously documented effects

on body mass and growth in meerkats (40). All continuous variables were standardized

to a mean of 0 and SD of 1.

Results

Effects of treatments on litter characteristics and offspring survival

There was no evidence that the treatment of pregnant females with cortisol

affected their ability to maintain litters to term or the survival of their pups prior to

emergence from the natal burrow (Tables 1-2). The number of pups surviving to

emergence from the natal burrow or 3, 6, or 12 months of age and the litter sex ratio

were not different among litters from cortisol-treated, control, or untreated females

(Tables 1-2).

344   *Effects of treatments on offspring early life growth*

The effects of the treatments on offspring growth from initial emergence to

nutritional independence (1-3 months) differed between daughters and sons, as

reflected in the significant three-way interaction between treatment, sex, and age (Table

3). Daughters (but not sons) from cortisol treated mothers grew more slowly from 1-3

months compared to those from control (fed) mothers (daughters: age x treatment, t = -

4.17, P < 0.0001; sons: t = -1.48, P = 0.14), but exhibited similar growth compared to

those from untreated (unfed) mothers (daughters: age x treatment, t = 0.65, P = 0.51;

sons: t = -0.52, P = 0.6, Table 3, Fig. 1). Daughters, but not sons, from control (fed)

mothers grew faster than those from untreated mothers (daughters: age x treatment, t =

-4.24, P < 0.0001; sons: t = 1.35, P = 0.18).

355   *Effects of treatments on offspring cooperative behaviour*

The effects of the maternal treatments on babysitting behaviour of offspring

depended upon the age and sex of the offspring (Table 4). Babysitting contributions in

daughters from mothers treated with cortisol during pregnancy were slightly, but

significantly higher with increasing age of the babysitter compared to those from control

mothers (age x treatment, z = -2.89, P = 0.0039) but not untreated mothers (age x

treatment, z = 1.88, P = 0.06; Table 4, Fig. 2). Babysitting contributions in sons from

mothers treated with cortisol during pregnancy showed a similar tendency to slightly

increase with age compared to those from control mothers, but this difference was not

significant (age x treatment, z = -1.92, P = 0.055). Further, age-related increases in

babysitting contributions between males from mothers treated with cortisol during

pregnancy and untreated mothers did not differ (age x treatment, z = -0.03, P = 0.97;

Table 4, Fig. 2). Comparisons of the magnitude of effect sizes showed that the

interaction between age and maternal treatment had a larger effect on babysitting

contributions in daughters but not sons than other variables known to impact babysitting

contributions, such as foraging success, age-related body mass, or group size (Table

4).

The effects of the maternal treatments on pup feeding depended upon the sex of

the offspring, but not their age (Table 5). Daughters, but not sons from mothers treated

with cortisol during pregnancy exhibited significantly more pup feeding contributions

than those from control mothers (females: z = -3.12, P = 0.00018; males: z = -1.14, P =

0.25) or untreated mothers (females: z = -3.49, P = 0.0005, sons: z = -1.03, P = 0.3,

Table 5, Fig. 3). Notably, the magnitude of effect size of maternal treatment for

daughters was much larger than other variables known to impact babysitting

contributions such as foraging success, age-related body mass, and group size (Table

5).

381   *Effects of treatments on offspring stress physiology*

Daughters from mothers treated with cortisol during pregnancy had lower

plasma cortisol concentrations (age x treatment, t = -1.76, P = 0.08, Table 6, Fig. 4A)

and lower fGCM concentrations (age x treatment, t = -2.9, P = 0 .004, Table 7, Fig. 5A)

as they became older compared to those from control mothers but these differences

were only significant for fGCM concentrations. Sons from mothers treated with cortisol

during pregnancy had significantly lower plasma cortisol concentrations as they became

older compared to those from control mothers (age x treatment t = -2.68, P = 0.008,

Table 6, Fig. 4B) but similar fGCM concentrations compared to those from control

mothers as they became older (age x treatment, t = -0.1, P = 0.49, Table 7, Fig. 5B).

Discussion

We found some support for our hypothesis that elevated maternal GCs would

reduce the potential for offspring to have direct reproductive opportunities and would

therefore shift them towards exhibiting more cooperative behaviour that could increase

their indirect fitness. Daughters, but not sons, from mothers treated with cortisol during

pregnancy grew more slowly early in life and exhibited more babysitting and pup

feeding behaviour as they became older compared to controls. Other than offspring

survival (Table 2), we were unable to quantify the direct and indirect fitness of offspring

from control or cortisol-treated mothers, but early life growth or body mass (which we

measured here) is closely linked to direct fitness opportunities in daughters (27-32).

Previous studies in meerkats show that female, but not male, offspring that grow faster

from 1-3 months are more likely to acquire the dominant breeding position (31), perhaps

because offspring that grow faster in their first 3 months of life are heavier later in life

(32, 57, 58), and heavier females are more likely to acquire a vacant dominant breeding

position (30, 32). As such, daughters, but not sons, from mothers treated with cortisol

levels during pregnancy should have reduced future direct fitness opportunities and

therefore increase their investment in behaviours that elevate their indirect fitness. Our

results are consistent with studies in other taxa that suggest that individuals adjust their

contributions to cooperative behaviour according to their future reproductive potential.

For example, in cooperatively breeding birds, when the chances of direct reproduction

are elevated, subordinates often stop helping at the nest (59). Studies of social wasps

show that individuals whose probability of acquiring the dominant breeding position was

experimentally increased exhibited significantly less helping behaviour (60, 61). Finally,

in cooperatively breeding fish, subordinates will reduce their helping investment

immediately prior to dispersal from their natal group where they attempt to reproduce on

their own rather than stay in their natal group and queue for dominance (62).

Our results show that increases in maternal GCs can increase the cooperative

behaviour of daughters, which should lead to substantial direct fitness benefits to

mothers. Daughters from mothers treated with cortisol during pregnancy exhibited more

alloparental care compared to controls, such that subsequent offspring produced in

groups with offspring from cortisol-treated mothers should have received more

alloparental care. Because offspring that receive more alloparental care grow faster

early in life or are larger later in life (32, 57), the presence of offspring from cortisol-

treated mothers should increase the direct fitness of dominant breeders and the indirect

fitness of the offspring from cortisol-treated mothers. Taken together, our results

suggest that this GC-mediated maternal effect reduced the direct fitness opportunities of

daughters by reducing their early life growth, but they compensated by increasing their

investment in indirect fitness opportunities (helping to rear non-descendent offspring).

This is in line with theoretical predictions that parental manipulation of the cooperative

behaviour of offspring can evolve if the costs of resisting the parental effect are high and

inclusive fitness benefits of helping rear subsequent offspring are increased (18), as is

the case in cooperative breeders.

Control females that were fed during pregnancy produced daughters that grew

faster during early development (1-3 months) compared to daughters from cortisol-

treated or untreated mothers. Although mothers that were treated with cortisol during

pregnancy received the same amount of supplemental food as controls, daughters and

sons from mothers fed cortisol during pregnancy did not differ in early life growth

compared to those from untreated mothers. This indicates that the additional food

provided to dominant females during pregnancy had the potential to increase growth,

but the added cortisol prevented those gains in body mass. This has implications for

understanding the fitness consequences of maternal stress on offspring growth

trajectories (15, 63) because our results show that elevated circulating GC levels in

pregnant females in the absence of energetic constraints induced reductions in the early

life growth of offspring. This supports the hypothesis that maternal GC levels during

offspring development act as a cue that induces plasticity in offspring growth rather than

simply mediating the effects of energetic constraints. Alternatively, elevated maternal

GCs could alter patterns of maternal investment in offspring. Identifying whether

offspring or mothers are driving these effects is a major challenge in studies of maternal

stress effects in wild animals.

The reductions in the activity of the neuroendocrine stress axis of daughters may

have potentiated the increased alloparental care behaviour that we observed.

Compared to daughters from control mothers, daughters from mothers treated with

cortisol during pregnancy exhibited more babysitting as they became older, more overall

pup feeding, and they also had lower plasma cortisol and fGCM concentrations. Males

from mothers treated with cortisol during pregnancy had significantly lower plasma

cortisol concentrations, but not fGCM concentrations as they got older and also tended

to exhibit more babysitting as they aged. The activity of the neuroendocrine stress axis

is closely linked to an array of social behaviours (64) and our recent work shows that

elevated activity of the neuroendocrine stress axis reduces babysitting in both females

and males and decreases pup feeding in females (39). Together, this supports the

hypothesis that the mechanism by which early life stress increases the cooperative

behaviour of daughters is by dampening the activity of their neuroendocrine stress axis.

Our results show that the effects of maternal GCs on offspring growth,

physiology, and behaviour were greater in daughters than in sons, which adds to

biomedical (65-66) and ecological (67-69) studies that highlight how early life conditions

or maternal GC levels can have sex-specific consequences for offspring. In meerkats,

there may be added benefits for the dominant female for altering the cooperative

behaviour of daughters compared to sons; daughters exhibit more cooperative

behaviour than sons (36) and are more responsive to the begging calls of subsequent

offspring that they provision with food (70). More broadly, sex-differences in natal

dispersal may cause these differential responses to parental effects. In meerkats,

subordinate males voluntarily disperse from their natal group to look for receptive

females but can return to their natal group whereas subordinate females rarely

voluntarily disperse from their natal group (71). In our case and in others (63), the more

philopatric sex (females) is more sensitive to early life conditions, which may be due to

differential costs of parental modification between the philopatric and dispersing sex. If

parental effects have long-term consequences on offspring characteristics, as we show

here, there may be an increased degree of mismatch between the phenotype of the

dispersing sex and the postnatal environment where individuals eventually settle. If this

mismatch has fitness costs, this should select for individuals from the dispersing sex to

be less responsive to cues from the parental phenotype or environment.

Our results provide some support for the hypothesis that parents may alter the

cooperative tendencies of their offspring by manipulating the characteristics of their

offspring (9-10), though we note that it is uncertain if the transfer of maternal GCs to

offspring was passive or active. Explanations regarding the evolutionary origins of

cooperative behaviour involve nepotism or kin selection (72), mutualisms or reciprocity

(73), but few studies have tested the "parental manipulation" hypothesis proposed by

Alexander (9). Some studies show that alleles that increase maternal fitness at the

expense of the direct fitness of offspring can evolve (19) and that cooperative breeders

may bias investment towards offspring that exhibit more cooperative behaviour (74).

Our study supports the hypothesis that environmental stressors may induce a parental

effect that can modify the cooperative tendencies of their offspring.

Finally, our results have two implications for theoretical models examining the

evolution of parental effects. First, given the sex-specificity of parental effects, our

results challenge the conclusions of models examining the evolution of parental effects

that assume that all offspring are equally sensitive to the parental effect (16), or those

that assume that the benefits of exhibiting the phenotype resulting from the parental

effect are equal for all offspring (18). Second, selfish parental effects are thought to be

relatively rare (8, 13) and theory (16-17) and empirical studies showing sex-specific

responses to early life stress (65-66) indicate that offspring can become resistant to

such selfish parental effects. However, some models indicate that the evolution of

selfish parental effects may be dependent upon the social environment (24), especially

if the selfish parental effect influences the expression of alloparental care behaviour of

offspring and therefore increases the indirect fitness of offspring. Our results provide an

example whereby a GC-mediated maternal effect should decrease the direct fitness of

daughters (by reducing their early life growth), but should increase the direct fitness of

mothers and indirect fitness of daughters by elevating their cooperative behaviour.

Ethics: All protocols used in our experiments were approved by the Animal Ethics

Committee at the University of Pretoria (Pretoria, South Africa: #EC031-13, #EC047-16)

and the Northern Cape Department of Environment and Nature Conservation Research

(South Africa: FAUNA 050/2013, FAUNA 192/2014, FAUNA 1020/2016).

Acknowledgements: We thank M. Manser for her contributions to maintain the long-

term Kalahari Meerkat Project (KMP) and her contributions to this specific project.

Thanks to P. Roth, S. Bischoff-Mattson, L. Howell, J. Samson, N. Thavarajah, and

numerous KMP volunteers who contributed to data collection. Thanks to M. Manser and

S. Guindre-Parker for helpful comments. Thanks to the Northern Cape Conservation

Authority for permission to conduct research and the Kotze family for allowing us to

study meerkats on their property.

Authors' Contributions: B.D. and T.H.C-B planned the study, B.D. designed the

experiments, B.D., C. Dubuc, D.L.C, T.H.C-B, D.G., I.BG., N.B., M.H., A.G., and C.

Duncan coordinated and collected data, B.D. conducted analyses and produced figures,

B.D. and T.H.C-B authored manuscript with contributions from all authors.

Date, code and materials: All data will be archived on Data Dryad. All code for

statistical analyses is available from first author.

Competing interests: We have no competing interests.

Funding: Research was supported by grants from National Environment Research

Council (RG53472) and the European Research Council (294494) to T.H.C-B. I.B.G.

was funded by the Swiss National Science Foundation (grant no. 31003A_13676 to M.

Manser). NCB was supported by a DST-NRF SARCHI chair of Mammal Behavioural

Ecology and Physiology. The KMP has also been financed by the University of

Cambridge (T.H.C.-B) and University of Zurich (M. Manser).

References

1) Wolf JB, Wade MJ (2009) What are maternal effects (and what are they not)?

Philosophical Transactions of the Royal Society B: Biological

Sciences, *364*(1520), 1107-1115.

2) Kirkpatrick M, Lande R (1989) The evolution of maternal characters. *Evolution*, *43*(3),

485-503.

3) Mousseau TA, Fox CW (Eds.) (1998) *Maternal Effects as Adaptations*. Oxford

University Press.

4) Agrawal AA, Laforsch C, Tollrian R (1999) Transgenerational induction of defences in

animals and plants. *Nature* 401, 60-63

5) Dantzer B, Newman AE, Boonstra R, Palme R, Boutin S, Humphries MM, McAdam

AG (2013) Density triggers maternal hormones that increase adaptive offspring

growth in a wild mammal. *Science*, 1235765.

6) Duckworth RA, Belloni V, Anderson SR (2015) Cycles of species replacement

emerge from locally induced maternal effects on offspring behavior in a

passerine bird. *Science*, *347*(6224), 875-877.

7) Uller T (2008) Developmental plasticity and the evolution of parental effects. *Trends*

in Ecology & Evolution, *23*(8), 432-438.

8) Uller T, Nakagawa S, English S (2013) Weak evidence for anticipatory parental

effects in plants and animals. *Journal of Evolutionary Biology*, *26*(10), 2161-2170.

9) Alexander, RD, (1974) The evolution of social behavior. *Annual Review of Ecology*

and Systematics, *5*(1), 325-383.

10) Michener CD, Brothers DJ 1974 Were workers of eusocial Hymenoptera initially

altruistic or oppressed? *Proceedings of the National Academy of Science* USA

71, 671–674

11) Müller W, Lessells CKM, Korsten P, Engelhardt NV (2007) Manipulative signals in

family conflict? On the function of maternal yolk hormones in birds. *The American*

Naturalist, *169*(4), E84-E96.

12) Schwarzkopf L, Andrews RM (2012) Are moms manipulative or just selfish?

Evaluating the "maternal manipulation hypothesis" and implications for life-history

studies of reptiles. *Herpetologica*, *68*(2), 147-159.

13) Marshall DJ, Uller T (2007) When is a maternal effect adaptive?. *Oikos*, *116*(12),

1957-1963.

14) Wilson AJ, Pilkington JG, Pemberton JM, Coltman DW, Overall AD, Byrne KA,

Kruuk, LEB (2005) Selection on mothers and offspring: whose phenotype is it

and does it matter? *Evolution*, *59*(2), 451-463.

15) Berghänel A, Heistermann M, Schülke O, Ostner J (2017) Prenatal stress

accelerates offspring growth to compensate for reduced maternal investment

across mammals. *Proceedings of the National Academy of Sciences*, *114*(50),

E10658-E10666.

16) Uller T, Pen I (2011) A theoretical model of the evolution of maternal effects under

parent–offspring conflict. *Evolution*, *65*(7), 2075-2084.

17) Kuijper B, Johnstone RA (2018) Maternal effects and parent–offspring

conflict. *Evolution*, *72*(2), 220-233.

18) González-Forero M, Gavrilets S (2013) Evolution of manipulated behavior. *The*

American Naturalist, *182*(4), 439-451.

19) Wade MJ (2001) Maternal effect genes and the evolution of sociality in haplo-diploid

organisms. *Evolution*, *55*(3), 453-458.

20) Linksvayer TA, Wade MJ (2005) The evolutionary origin and elaboration of sociality

in the aculeate Hymenoptera: maternal effects, sib-social effects, and

heterochrony. *The Quarterly Review of Biology*, *80*, 317-336.

21) Bourke AF, Franks NR (1995). *Social Evolution in Ants*. Princeton University Press.

22) Clutton-Brock TH (2016) *Mammal Societies*. Wiley-Blackwell

23) Russell A F, Lummaa V (2009) Maternal effects in cooperative breeders: from

hymenopterans to humans. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *364*(1520), 1143-1167.

24) Taborsky B (2017) Developmental plasticity: preparing for life in a complex

world. *Advances in the Study of Behavior*, *49*, 49-99.

25) Carter CS, Roberts RL. (1997) The psychobiological basis of cooperative breeding

in rodents. In *Cooperative Breeding in Mammals* (Eds.: NG Solomon, JA

French), Cambridge University Press, Pp. 231-266.

26) De Fraipont M, Clobert J, Alder HJ, Meylan S (2000) Increased pre-natal maternal

corticosterone promotes philopatry of offspring in common lizards *Lacerta*

vivipara. *Journal of Animal Ecology*, *69*(3), 404-413.

27) Clutton-Brock TH, Brotherton PNM, O'riain MJ, Griffin AS, Gaynor D, Kansky R,

Sharpe L, McIlrath GM (2001) Contributions to cooperative rearing in

meerkats. *Animal Behaviour*, *61*(4), 705-710.

28) Russell AF, Clutton-Brock TH, Brotherton PN, Sharpe LL, Mcilrath G, Dalerum FD,

Cameron EZ, Barnard JA (2002). Factors affecting pup growth and survival in co-

operatively breeding meerkats *Suricata suricatta*. *Journal of Animal*

Ecology, *71*(4), 700-709.

29) Thornton A (2008) Social learning about novel foods in young meerkats. *Animal*

Behaviour, *76*(4), 1411-1421.

30) Hodge SJ, Manica A, Flower TP, Clutton-Brock TH (2008) Determinants of

reproductive success in dominant female meerkats. *Journal of Animal*

Ecology, *77*(1), 92-102.

31) English S, Huchard E, Nielsen JF, Clutton-Brock TH (2013) Early growth,

dominance acquisition and lifetime reproductive success in male and female

cooperative meerkats. *Ecology and Evolution*, *3*(13), 4401-4407.

32) Russell AF, Young AJ, Spong G, Jordan NR, Clutton-Brock TH (2007) Helpers

increase the reproductive potential of offspring in cooperative

meerkats. *Proceedings of the Royal Society of London B: Biological*

Sciences, *274*(1609), 513-520.

33) Clutton-Brock TH, Gaynor D, Kansky R, MacColl ADC, McIlrath G, Chadwick P,

Brotherton PNM, O'rian JM, Manser M, Skinner JD (1998) Costs of cooperative

behaviour in suricates (*Suricata suricatta*). *Proceedings of the Royal Society of*

London B: Biological Sciences, *265*(1392), 185-190.

34) Brotherton PNM, Clutton-Brock TH, O'riain MJ, Gaynor D, Sharpe L, Kansky R,

McIlrath GM (2001) Offspring food allocation by parents and helpers in a

cooperative mammal. *Behavioral Ecology*, *12*(5), 590-599.

35) Russell AF, Sharpe LL, Brotherton PNM, Clutton-Brock TH (2003) Cost

631   minimization by helpers in cooperative vertebrates. Proceedings of the National

632   Academy of Sciences USA 100(6), 3333-3338.

633 36) Clutton-Brock TH, Russell AF, Sharpe LL, Young AJ, Balmforth Z, McIlrath GM

634   (2002). Evolution and development of sex differences in cooperative behavior in

635   meerkats. *Science*, *297*(5579), 253-256.

636 37) Bell MBV, Cant MA, Borgeaud C, Thavarajah N, Samson J, Clutton-Brock TH

637   (2014) Suppressing subordinate reproduction provides benefits to dominants in

638   cooperative societies of meerkats. *Nature Communications* 5, 4499

639 38) Harris A, Seckl J (2011) Glucocorticoids, prenatal stress and the programming of

640   disease. *Hormones and Behavior*, *59*(3), 279-289.

641 39) Dantzer B, Goncalves IB, Spence-Jones HC, Bennett NC, Heistermann M,

642   Ganswindt A, Dubuc C, Gaynor D, Manser MB, Clutton-Brock TH 2017. The

643   influence of stress hormones and aggression on cooperative behaviour in

644   subordinate meerkats. *Proc. R. Soc. B*, *284*(1863), 20171248.

645 40) English S, Bateman AW, Mares R, Ozgul A, Clutton-Brock TH (2014) Maternal,

646   social and abiotic environmental effects on growth vary across life stages in a

647   cooperative mammal. *Journal of Animal Ecology*, *83*(2), 332-342.

648 41) Kutsukake N, Clutton-Brock TH (2006) Aggression and submission reflect

649   reproductive conflict between females in cooperatively breeding meerkats

650   *Suricata suricatta*. *Behav. Ecol. Sociobiol.* **59**, 541-548. (doi: 10.1007/s00265-

651   005-0079-7)

652 42) Clutton-Brock TH, Brotherton PNM, O'Riain MJ, Griffin AS, Gaynor D, Sharpe L,

Kansky R, Manser MB, McIlrath GM (2000). Individual contributions to

babysitting in a cooperative mongoose, *Suricata suricatta. Proceedings of the*

Royal Society of London B: Biological Sciences, *267*(1440), 301-305.

43) Kunc HP, Madden JR, Manser MB (2007) Begging signals in a mobile feeding

system: The evolution of different call types. *The American Naturalist* 170, 617-

624.

44) Carlson AA, Young AJ, Russell AF, Bennett NC, McNeilly AS, Clutton-Brock T

(2004). Hormonal correlates of dominance in meerkats (*Suricata*

suricatta). *Hormones and Behavior*, *46*(2), 141-150.

45) Dantzer B, Bennett NC, Clutton-Brock TH (2017) Social conflict and costs of

cooperation in meerkats are reflected in measures of stress

hormones. *Behavioral Ecology*, *28*(4), 1131-1141.

46) Heistermann M, Finke M, Hodges JK (1995) Assessment of female reproductive

status in captive-housed Hanuman langurs (*Presbytis entellus*) by measurement

of urinary and fecal steroid excretion patterns. *Am. J. Primatol.* 37, 275-284. (doi:

10.1002/ajp.1350370402)

47) Braga Goncalves I, Heistermann M, Santema P, Dantzer B, Mausbach J,

Ganswindt A, Manser, MB (2016) Validation of a fecal glucocorticoid assay to

assess adrenocortical activity in meerkats using physiological and biological

stimuli. *PloS One*, *11*(4), e0153161.

48) Heistermann M, Palme R, Ganswindt A (2006) Comparison of different

enzyme immunoassays for assessment of adrenocortical activity in primates

based on fecal analysis. *American Journal of Primatology* 68, 257-273.

49) Clutton-Brock TH, Russell AF, Sharpe LL (2003) Meerkat helpers do not specialize

in particular activities. *Animal Behaviour* 66(3), 531-540.

50) Carlson AA, Manser MB, Young AJ, Russell AF, Jordan NR, McNeilly AS, Clutton-

Brock T (2006) Cortisol levels are positively associated with pup-feeding rates in

male meerkats Proceedings of the Royal Society B 273(1586), 571-577.

51) Lesnoff M, Lancelot R (2013) aods3: analysis of overdispersed data using S3

methods. aods3 package version 0.4-1.

52) R Development Core Team. 2013. R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria; available at

www.r-project.org/.

53) Bates D, Maechler M, Bolker B, Walker S (2014) lme4: Linear mixed-effects models

using Eigen and S4. *R Package Version 1*(7), 1-23.

54) Kuznetsova A, Brockhoff PB, Christensen RHB (2016) lmerTest: Tests in Linear

Mixed Effects Models. R package version 2.0-30; available at http://CRAN.R-

project.org/package=lmerTest

55) Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid

common statistical problems. *Meth. Ecol. Evol.* **1**, 3-14. (doi: 10.1111/j.2041-

210X.2009.00001.x)

56) Fox J, Monette G (1992) Generalized collinearity diagnostics. *Journal of the*

American Statistical Association 87(417), 178-183.

57) Clutton-Brock TH, Russell AF, Sharpe LL, Brotherton PNM, McIlrath GM, White S,

Cameron EZ (2001) Effects of helpers on juvenile development and survival in

meerkats. *Science, 293*(5539), 2446-2449.

58) Huchard E, English S, Bell MB, Thavarajah N, Clutton-Brock T (2016) Competitive

growth in a cooperative mammal. *Nature*, *533*(7604), 532.

59) Hatchwell BJ, Komdeur J (2000) Ecological constraints, life history traits and the

evolution of cooperative breeding. *Animal Behaviour* 59(6), 1079-1086.

60) Cant MA, Field J (2001) Helping effort and future fitness in cooperative animal

societies. Proceedings of the Royal Society B 268, 1959-1964.

61) Field J, Cronin A, Bridge C (2006) Future fitness and helping in social

queues. *Nature*, *441*(7090), 214.

62) Zöttl M, Chapuis L, Freiburghaus M, Taborsky M (2013). Strategic reduction of help

before dispersal in a cooperative breeder. *Biology Letters*, *9*(1), 20120878.

63) Sheriff MJ, Bell A, Boonstra R, Dantzer B, Lavergne SG, McGhee KE, MacLeod KJ,

Winandy L, Zimmer C, Love OP (2017) Integrating ecological and evolutionary

context in the study of maternal stress. *Integrative and Comparative*

Biology, *57*(3), 437-449.

64) Raulo A, Dantzer B. (2018) Associations between glucocorticoids and sociality

across a continuum of vertebrate social behaviour. *Ecology and Evolution* in

press (doi: 10.1002/ece3.4059).

65) Weinstock M (2007) Gender differences in the effects of prenatal stress on brain

development and behaviour. *Neurochemical Research*, *32*(10), 1730-1740.

66) Mueller, BR, Bale, TL (2008) Sex-specific programming of offspring emotionality

after stress early in pregnancy. *Journal of Neuroscience*, *28*(36), 9055-9065.

67) Wilkin TA, Sheldon BC (2009) Sex differences in the persistence of natal

environmental effects on life histories. *Current Biology*, *19*(23), 1998-2002.

68) Metzger DC, Schulte PM (2016) Maternal stress has divergent effects on gene

expression patterns in the brains of male and female threespine

stickleback. *Proc. R. Soc. B*, *283*(1839), 20161734.

69) Marshall HH, Vitikainen EI, Mwanguhya F, Businge R, Kyabulima S, Hares MC,

Inzani E, Kalema-Zikusoka G, Mwesige K, Nichols HJ, Sanderson JL, Thompson

FJ, Cant MA (2017) Lifetime fitness consequences of early-life ecological

hardship in a wild mammal population. *Ecology and Evolution*, *7*(6), 1712-1724.

70) English S, Kunc HP, Madden JR, Clutton-Brock TH (2008) Sex differences in

responsiveness to begging in a cooperative mammal. *Biology Letters*, *4*(4), 334-

337.

71) Clutton-Brock TH, Manser M (2016) Meerkats: cooperative breeding in the Kalahari.

Cooperative Breeding in Vertebrates (Eds.: WD Koenig, JL Dickinson).

Cambridge University Press, Pp. 294-317

72) Hamilton WD (1964) The genetical evolution of social behaviour. II. *Journal of*

Theoretical Biology, *7*(1), 17-52.

73) Trivers RL (1971) The evolution of reciprocal altruism. *The Quarterly Review of*

Biology, *46*(1), 35-57.

74) Griffin AS, Sheldon BC, West SA (2005) Cooperative breeders adjust offspring sex

ratios to produce helpful helpers. *The American Naturalist*, *166*(5), 628-632.

741
742

Figure 1. (A) Daughters but not (B) sons from mothers treated with cortisol during

pregnancy were significantly smaller from initial emergence from their natal burrow to

nutritional independence (~1-3 months) compared to those from control mothers

(daughters: age x treatment, t = -4.17, P < 0.0001; sons: t = -1.48, P = 0.14), but not

untreated mothers (daughters: age x treatment, t = 0.65, P = 0.51; sons: t = -0.52, P =

0.6, Table 3). Data are body mass measures from offspring from cortisol-treated

(females: n = 373 estimates; males: n = 488), control (females: n = 215; males: n =

241), and untreated mothers (females: n = 1121; males: n = 2238). Raw data and

regression lines are shown (full results in Table 3).

Figure 2. Babysitting contributions of (A) daughters and (B) sons from mothers treated

with cortisol during pregnancy increased with age at a faster rate than those from

control (females: z = -2.89, P = 0.0039; males: z = -1.92, P = 0.055), but not untreated

("None") mothers (females: z = 1.88, P = 0.06; males: z = -0.03, P = 0.97, Table 4).

Data are relative babysitting contributions from offspring from cortisol-treated (females:

n = 15 estimates; males: n = 24), control (females: n = 15; males: n = 10), and untreated

mothers (females: n = 49; males: n = 69). Raw data and regression lines are shown (full

results in Table 4).

Figure 3. Pup feeding contributions of (A) daughters, but not (B) sons, from mothers

treated with cortisol during pregnancy were significantly higher compared to from control

(females: z = -3.09, P = 0.0004; males: z = -1.15, P = 0.25) or untreated ("None")

mothers (females: z = -3.47, P = 0.0005; males: z = -0.89, P = 0.37, Table 5). Data are

relative pup feeding contributions from offspring from cortisol-treated (females: n = 16

estimates; males: n = 26), control (females: n = 16; males: n = 10), and untreated

mothers (females: n = 64; males: n = 71). Raw data are shown (full results in Table 5).

Boxplots show median (solid horizontal line), mean (grey diamonds), and first (25%)

and third (75%) quartiles.

Figure 4. (A) Daughters and (B) sons from mothers treated with cortisol during

pregnancy had lower plasma cortisol concentrations as they became older compared to

those from control mothers, though the difference was only significant in males

(daughters: age x treatment, t = -1.76, P = 0.08; sons: age x treatment t = -2.68, P =

0.008, Table 6). Data are residual plasma cortisol concentrations from offspring from

cortisol-treated (females: n = 64 samples; males: n = 92) and control (females: n = 89;

males: n = 104) mothers. Residuals from a linear mixed-effects model (Table 6) are

shown on y-axis.

Figure 5. Faecal glucocorticoid metabolite (fGCM) concentrations in (A) daughters but

not (B) sons from mothers treated with cortisol during pregnancy were significantly

lower than those from control mothers as they became older (daughters: age x

treatment, t = -2.9, P = 0.004; sons: age x treatment, t = -0.1, P = 0.49, Table 7. Data

are residual fGCM concentrations from offspring from cortisol-treated (females: n = 79

samples; males: n = 118) and control (females: n = 154; males: n = 201) mothers.

Residuals from a linear mixed-effects model (Table 7) are shown on y-axis.

796 **Table 1. Summary of effects of dominant female treatments on litter characteristics and**
797 **offspring survival.** Number of pups emerged correspond to those that emerged from the natal
798 burrow and in some cases these pups died before their sex could be determined (shown as
799 "Unk"). Three of the 13 litters treated with cortisol and one of the 7 control (fed) litters were
800 aborted prior to birth.

Treatment	Total # litters & females treated	Total # pups emerged (F, M, Unk)	Avg. Pups emerged	Avg. Pups Surviving to 3 months	Avg. Pups Surviving to 6 months	Avg. Pups Surviving to 12 months
Untreated	52 (21 females)	49 F, 84 M, 52 Unk	3.78 ± 1.23	2.68 ± 1.58	2.1 ± 1.63	1.62 ± 1.54
Control	7 (6 females)	12 F, 13 M	4.17 ± 0.98	3.83 ± 1.17	3.83 ± 1.17	2.83 ± 1.94
Cortisol	13 (10 females)	13 F, 18 M	3.87 ± 0.83	3.25 ± 1.03	2.75 ± 1.28	1.75 ± 1.75

801
802

Table 2. Effects of dominant female treatments (cortisol or control) on litter characteristics
and pup survival. Results are from a linear mixed-effects model (# pups emerged) or
generalized linear mixed-effects models (GLMMs, all other response variables) that each
contained random intercept terms for dominant female identity and year. No GLMM was
overdispersed as indicated by goodness of fit tests (R package aods3, P-values from Pearson χ^2
tests ranged from 0.13 to 1). The number of litters aborted by untreated females was not known
so we only assessed the effects of cortisol vs. control treatments on the number of litters aborted.
Litter sex ratio is the proportion of males in the litter.

Response variable	Fixed effect	b	SE	t or z	P-value
# Litters aborted					
	Intercept	-1.21	0.66	-1.83	0.07
	Birthdate	-0.17	0.59	-0.29	0.77
	Treatment				
	Control	-0.59	1.27	-0.46	0.64
# Pups emerged					
	Intercept	**3.9**	**0.42**	**9.36**	**<0.0001**
	Birthdate	0.02	0.16	0.11	0.91
	Treatment				
	Control	0.3	0.62	0.49	0.63
	Untreated	-0.11	0.45	-0.25	0.8
Litter sex ratio					
	Intercept	-0.54	0.3	-1.84	0.066
	Birthdate	0.02	0.12	0.15	0.88
	Treatment				
	Control	-0.11	0.45	-0.23	0.81
	Untreated	0.09	0.33	0.27	0.78
Prop. litter surviving to 3 months					
	Intercept	-0.16	0.27	-0.62	0.53
	Birthdate	-0.12	0.11	-1.09	0.27
	Treatment				
	Control	0.06	0.39	0.16	0.87
	Untreated	-0.19	0.3	-0.64	0.52
Prop. litter surviving to 6 months					
	Intercept	-0.32	0.28	-1.16	0.24
	Birthdate	-0.19	0.11	-1.65	0.099
	Treatment				
	Control	0.21	0.4	0.52	0.6
	Untreated	-0.28	0.31	-0.91	0.36
Prop. litter surviving to 12 months					
	Intercept	**-0.91**	**0.46**	**-1.999**	**0.046**
	Birthdate	-0.22	0.14	-1.58	0.11
	Treatment				
	Control	0.36	0.46	0.79	0.43
	Untreated	0.096	0.38	0.25	0.8

811 Reference for Treatment was cortisol-treated mothers. Data other than # litters aborted are based upon an
812 initial sample size of offspring from untreated (195 pups from 52 litters produced by 21 dominant
813 females), control (25 pups from 6 litters produced by 6 females), or cortisol-treated (31 pups from 10
814 litters produced by 9 females) litters that produced pups that emerged from the burrow.
815
816
817

Table 3. Effect of dominant female treatments on offspring growth from emergence to nutritional independence (1-3 months of age). Data are from a linear mixed-effects model where the response variable was morning body mass that contained random intercept terms for individual identity nested in birth litter nested in mother nested in natal group ($\sigma^2 = 116.7$) and year ($\sigma^2 = 0$). If fixed effects by themselves were involved in significant higher order interactions with other variables, only parameter estimates are shown.

Fixed Effect	b	SE	t	df	P-value
Intercept					
Females	**184.6**	**9.75**	**18.93**	**53**	**<0.0001**
Males	**189.1**	**9.5**	**19.8**	**49**	**<0.0001**
Litter size	-5.31	3.9	-1.37	48	0.18
First weight	**15.96**	**2.52**	**6.344**	**166**	**<0.0001**
Sex (M)	4.54	4.32			
Age	53.3	1.06			
Age^2	-4.76	1.15			
Rainfall	**2.81**	**0.61**	**4.57**	**4550**	**<0.0001**
Season (Sine)	**-67.2**	**3.3**	**-20.6**	**4238**	**<0.0001**
Season (Co-sine)	**66.22**	**3.4**	**19.2**	**4191**	**<0.0001**
Group size	2.3	2.66	0.86	109	0.39
Group size^2	-0.09	2.21	-0.04	178	0.97
Treatment (Control)					
Females	22.6	15.6			
Males	12.9	15.5			
Treatment (None)					
Females	23.6	10.6			
Males	22.9	10.4			
Age x Sex (M)	1.64	1.37	1.19	4567	0.23
Age^2 x Sex (M)	1.02	1.53	0.67	4543	0.5
Sex (M) x Treatment (Control)	-9.73	7.7	-1.27	164	0.21
Sex (M) x Treatment (None)	-0.63	5.1	-0.12	172	0.9
Age x Treatment (Control)					
Females	**7.6**	**1.8**	**4.17**	**4595**	**<0.0001**
Males	-2.3	1.5	-1.48	4548	0.14
Age x Treatment (None)					
Females	0.79	1.2	0.65	4574	0.51
Males	-0.53	1.01	-0.52	4580	0.6
Age^2 x Treatment (Control)					
Females	-0.13	1.93	-0.07	4521	0.94
Males	-1.05	1.6	-0.65	4547	0.52
Age^2 x Treatment (None)					
Females	-2.6	1.3	-1.96	4534	0.051
Males	**-6.9**	**1.1**	**-6.23**	**4556**	**<0.0001**
Age x Treatment (Control) x Sex (M)	**-9.9**	**2.3**	**-4.23**	**4574**	**<0.0001**
Age x Treatment (None) x Sex (M)	-1.3	1.6	-0.84	4581	0.4
Age^2 x Treatment (Control) x Sex (M)	-0.93	2.5	-0.37	4552	0.71
Age^2 x Treatment (None) x Sex (M)	**-4.3**	**1.7**	**-2.5**	**4552**	**0.01**

Data used in these analyses were 4676 measures of body mass from 195 meerkats produced by 21 dominant females across 53 different litters in 16 different social groups in three different years. Only offspring that survived to 90 days of age were included in these analyses.

Table 4. Effect of dominant female treatments on relative babysitting contributions. Data
are from a generalized linear mixed-effects model where the response variable is the proportion
of babysitting exhibited by the subordinate meerkat relative to the total babysitting contributions
the litter received. The model contained random intercept terms for individual ($\sigma^2 = 0.12$), litter
nested within group ($\sigma^2 = < 0.0001$), and year ($\sigma^2 = 0.000$). If fixed effects by themselves were
involved in significant higher order interactions with other variables, only parameter estimates
are shown.

Fixed Effect	b	SE	z	P-value
Intercept				
Females	**-2.1**	**0.22**	**-9.37**	**< 0.0001**
Males	**-2.14**	**0.19**	**-11.01**	**< 0.0001**
Babysitting length	**0.24**	**0.08**	**2.77**	**0.0056**
Observation time	**-0.28**	**0.08**	**-3.4**	**0.0007**
Litter size	0.015	0.05	0.32	0.75
Mixed Litter?	-0.04	0.12	-0.31	0.76
Sex (M)	-0.03	0.27	-0.13	0.9
Age				
Females	0.13	0.17		
Males	0.4	0.19		
Foraging success	-0.05	0.05	-0.99	0.32
Mass				
Females	-0.32	0.12		
Males	-0.098	0.14		
Group size				
Females	**-0.35**	**0.09**	**-3.97**	**< 0.0001**
Males	**-0.27**	**0.08**	**-3.28**	**0.001**
Treatment (Control)				
Females	-0.67	0.31		
Males	-0.12	0.28		
Treatment (None)				
Females	-0.06	0.23		
Males	-0.16	0.18		
Foraging success x Mass	0.036	0.06	0.57	0.57
Age x Mass				
Females	**-0.47**	**0.09**	**-5.08**	**< 0.0001**
Males	**-0.29**	**0.08**	**-3.79**	**0.0001**
Age x Sex (M)	0.28	0.23	1.18	0.24
Mass x Sex (M)	0.22	0.17	1.26	0.21
Group size x Sex (M)	0.07	0.1	0.77	0.44
Treatment (Control) x Sex (M)	0.55	0.42	1.31	0.19
Treatment (None) x Sex (M)	-0.1	0.28	-0.36	0.72
Age x Treatment (Control)				
Females	**-0.62**	**0.21**	**-2.89**	**0.0039**
Males	-0.52	0.27	-1.92	0.055
Age x Treatment (None)				
Females	0.34	0.18	1.88	0.059
Males	-0.006	0.16	-0.03	0.97
Age x Mass x Sex	0.18	0.12	1.5	0.13
Age x Treatment (Control) x Sex (M)	0.09	0.34	0.28	0.78
Age x Treatment (None) x Sex (M)	-0.34	0.24	-1.45	0.14

835   Data used in these analyses were 182 observations of relative babysitting contributions to 28
836 litters produced in 9 groups across 3 years recorded from 105 subordinate meerkats.
837
838

840 **Table 5. Effect of dominant female treatments on relative pup feeding contributions.** Data
are from a generalized linear mixed-effects model where the response variable is the proportion
of pup feeds exhibited by the subordinate meerkat relative to the total pup feeds the litter
received. The model contained random intercept terms for individual ($\sigma^2 = 0.000$), litter nested
within group ($\sigma^2 = 0.2$), year ($\sigma^2 = 0.08$), and an observational level random intercept term to
control for overdispersion ($\sigma^2 = 0.19$). If fixed effects by themselves were involved in significant
higher order interactions with other variables, only parameter estimates are shown.

Fixed Effect	b	SE	z	P-value
Intercept				
Females	**-2.27**	**0.28**	**-8.2**	**<0.0001**
Males	**-2.66**	**0.25**	**-10.45**	**<0.0001**
Observation time	**0.59**	**0.06**	**9.41**	**<0.0001**
Litter size	-0.055	0.1	-0.55	0.58
Mixed litter (Y)	-0.03	0.29	-0.1	0.92
Sex (M)	-0.39	0.2		
Age				
Females	-0.37	0.16		
Males	0.29	0.14		
Foraging success	**0.14**	**0.06**	**2.12**	**0.033**
Mass				
Females	0.1	0.11		
Males	-0.37	0.11		
Group size				
Females	**-0.48**	**0.13**	**-3.57**	**0.0003**
Males	**-0.32**	**0.13**	**-2.46**	**0.014**
Treatment (Control)				
Females	**-0.73**	**0.23**	**-3.12**	**0.0018**
Males	-0.24	0.21	-1.14	0.25
Treatment (None)				
Females	**-0.64**	**0.18**	**-3.49**	**0.0005**
Males	-0.14	0.13	-1.03	0.3
Foraging success x Mass	0.06	0.06	0.97	0.33
Age x Mass				
Females	-0.04	0.07	-0.62	0.54
Males	**-0.19**	**0.07**	**-2.68**	**0.007**
Age x Sex (M)	**0.66**	**0.19**	**3.49**	**0.00048**
Mass x Sex (M)	**-0.47**	**0.13**	**-3.63**	**0.0003**
Group size x Sex (M)	0.16	0.08	1.89	0.059
Treatment (Control) x Sex (M)	0.49	0.31	1.58	0.11
Treatment (None) x Sex (M)	**0.5**	**0.22**	**2.23**	**0.023**
Age x Treatment (Control)				
Females	-0.11	0.22	-0.5	0.61
Males	0.03	0.29	0.11	0.91
Age x Treatment (None)				
Females	0.17	0.16	1.03	0.3
Males	-0.02	0.14	-0.12	0.9
Age x Mass x Sex	-0.14	0.09	-1.51	0.13
Age x Treatment (Control) x Sex (M)	0.14	0.34	0.42	0.67
Age x Treatment (None) x Sex (M)	-0.19	0.21	-0.91	0.36

846   Data used in these analyses were 192 observations of relative pup feeding contributions to 26
847 litters produced in 7 groups across 3 years recorded from 101 subordinate meerkats.
848

Table 6. Effect of dominant female treatments on plasma cortisol concentrations. Data are
from a linear mixed-effects model where the response variable is plasma cortisol concentrations
(ln transformed) of the subordinate meerkat. The model contained random intercept terms for
individual nested within their birth litter ($\sigma^2 = 0.034$), and capture group ($\sigma^2 = 0.000$). If fixed
effects by themselves were involved in significant higher order interactions with other variables,
only parameter estimates are shown.

Fixed Effect	b	SE	t	df	P-value
Intercept					
Females	**3.18**	**0.29**	**11.31**	**128**	**<0.0001**
Males	**3.37**	**0.25**	**13.37**	**113**	**<0.0001**
Sampling time	**1.16**	**0.10**	**11.95**	**259**	**<0.0001**
Sampling time^2	**-0.24**	**0.03**	**-7.24**	**269**	**<0.0001**
Time of day	-0.2	0.14	-1.38	276	0.17
Sample year (2015)	0.41	0.24	1.72	228	0.09
Sex (M)	0.2	0.2	0.97	39	0.34
Age					
Females	-0.24	0.18			
Males	-0.01	0.14			
Foraging success					
Females	0.13	0.11	-1.22	231	0.22
Males	-0.15	0.1	-1.42	269	0.16
Group size	0.07	0.11	0.62	242	0.54
Group size^2	0.05	0.08	0.63	211	0.53
Pups in group	-0.31	0.18	-1.79	276	0.075
Group sex ratio	**0.2**	**0.08**	**2.53**	**180**	**0.01**
Relatedness	-0.09	0.18	-0.5	40	0.62
Weather (PC1)	-0.08	0.1	-0.75	276	0.45
Treatment (Cortisol)					
Females	-0.02	0.24			
Males	-0.32	0.19			
Sex (M) x Age	0.22	0.22	1.04	272	0.3
Sex (M) x Foraging success	-0.013	0.14	-0.09	257	0.93
Sex (M) x Treatment (Cortisol)	-0.3	0.3	-1.01	35	0.32
Age x Treatment (Cortisol)					
Females	-0.42	0.24	-1.76	276	0.08
Males	**-0.5**	**0.19**	**-2.68**	**274**	**0.008**
Group size x Pups Present (Yes)	-0.05	0.15	-0.36	216	0.72
Group size^2 x Pups Present (Yes)	0.07	0.12	0.6	263	0.55
Group size x Weather (PC1)	-0.02	0.08	-0.30	273	0.76
Age x Sex (M) x Treatment (Cortisol)	-0.08	0.31	-0.26	275	0.8

855   Data used in these analyses were 299 measures of plasma cortisol concentrations from 49
856 subordinate meerkats produced in 14 litters from 10 different groups. Reference levels (intercept) for
857 "Sex" was female, "Pups in group" was Yes, and for Relatedness was "No parent was dominant".
858

Table 7. Effect of dominant female treatments on faecal glucocorticoid metabolite (fGCM) concentrations. Data are from a linear mixed-effects model where the response variable fGCM concentrations (ln+1 transformed) of the subordinate meerkat. The model contained random intercept terms for individual nested within their birth litter ($\sigma^2 = 0.094$) and collection group ($\sigma^2 = 0.000$). If fixed effects by themselves were involved in significant higher order interactions with other variables, only parameter estimates are shown.

Fixed Effect	b	SE	t	df	P-value
Intercept					
Females	**5.4**	**0.25**	**21.36**	**50**	**<0.0001**
Males	**5.33**	**0.23**	**23.4**	**23**	**<0.0001**
Time of day	**-0.14**	**0.04**	**-3.29**	**516**	**0.001**
Sample year (2015)	0.07	0.17	0.43	120	0.67
Sex (M)	-0.06	0.2	-0.31	22	0.76
Age					
Females	0.48	0.16			
Males	0.06	0.1			
Foraging success					
Females	0.16	0.13	1.29	355	0.2
Males	0.04	0.06	0.81	520	0.42
Group size	**0.41**	**0.12**	**3.52**	**96**	**0.0007**
Group size^2	-0.05	0.11	-0.78	267	0.43
Pups in group	-0.13	0.16	-0.78	267	0.43
Group sex ratio	**0.16**	**0.07**	**2.27**	**158**	**0.024**
Relatedness	0.28	0.25	1.1	14	0.29
Weather (PC1)	0.09	0.07	1.4	203	0.16
Treatment (Cortisol)					
Females	-0.12	0.29			
Males	-0.03	0.24			
Sex (M) x Age	**-0.42**	**0.17**	**-2.47**	**221**	**0.014**
Sex (M) x Foraging success	-0.12	0.13	-0.87	375	0.38
Sex (M) x Treatment (Cortisol)	0.1	0.33	0.29	18	0.77
Age x Treatment (Cortisol)					
Females	**-0.63**	**0.22**	**-2.9**	**447**	**0.004**
Males	-0.1	0.14	-0.69	491	0.49
Group size x Pups Present (Yes)	-0.17	0.12	-1.46	216	0.14
Group size^2 x Pups Present (Yes)	0.11	0.12	456	0.97	0.33
Group size x Weather (PC1)	0.04	0.05	0.69	129	0.49
Age x Sex (M) x Treatment (Cortisol)	**0.53**	**0.25**	**2.15**	**413**	**0.032**

Data used in these analyses were 542 faecal samples (n= 355 from controls, n = 187 from cortisol treated litters) from 34 subordinate meerkats (control: n = 12 females, n = 11 males; cortisol-treated: n = 5 females, n = 6 males) produced in 10 litters from 7 different groups. Reference levels (intercept) for "Sex" was female, "Pups in group" was Yes, and for Relatedness was "No parent was dominant".