1 **Maternal stress increases cooperative behaviour of offspring in wild meerkats**

3   Ben Dantzer[1,2,3*], Constance Dubuc[1,2], Ines Braga Goncalves[2,4], Dominic L. Cram[1,2],

4   Nigel C. Bennett[5], Andre Ganswindt[5,6], Chris Duncan[2], David Gaynor[2,5], Tim H. Clutton-

5   Brock[1,2,5]

[1]Department of Zoology, University of Cambridge, Cambridge, United Kingdom

[2]Kalahari Meerkat Project, Kuruman River Reserve, Northern Cape, South Africa

[3]Department of Psychology, Department of Ecology and Evolutionary Biology,

University of Michigan, Ann Arbor, MI 48109, USA

[4]Department of Evolutionary Biology and Environmental Studies, Animal Behaviour,

University of Zurich, Switzerland

[5]Mammal Research Institute, Department of Zoology and Entomology, University of

Pretoria, Pretoria, South Africa

[6]Endocrine Research Laboratory, Department of Anatomy and Physiology, University of

Pretoria, 0110, Onderstepoort, South Africa

*Corresponding Author: dantzer@umich.edu

Abstract

The phenotype of parents can have long-lasting effects on the development of offspring as well as on their behaviour, physiology, and morphology as adults. In some cases, these changes may increase offspring fitness but, in others, they can elevate parental fitness at a cost to the fitness of their offspring. Here, we show that in Kalahari meerkats (*Suricata suricatta*) the circulating glucocorticoid hormones of pregnant females affect the growth and cooperative behaviour of their offspring. We performed a 3-year experiment in wild meerkats to test the hypothesis that stress-mediated maternal effects reduce the potential for offspring to reproduce directly and therefore cause them to exhibit more cooperative behaviour. Daughters (but not sons) born to mothers treated with cortisol during pregnancy were smaller early in life and exhibited significantly more of two types of cooperative behaviour (pup rearing and feeding) once they were adults compared to offspring from control mothers. They also had lower measures of stress hormones as they aged, which could explain the observed increases in cooperative behaviour. Because early growth is a crucial determinant of fitness in female meerkats, our results indicate that stress-mediated maternal effects may reduce the fitness of offspring but may elevate parental fitness as a consequence of increasing the cooperative behaviour of their daughters.

Significance Statement

Maternal characteristics can have a profound influence on offspring, which are referred to as maternal effects. It is widely assumed that selfish maternal effects, where mothers modify offspring to benefit their own fitness and decrease the fitness of offspring, are rare but this assumption is rarely tested. We show that a stress-mediated maternal effect in meerkats altered the growth, behaviour, and stress physiology of daughters in such a way that benefitted mothers at the expense of their daughters. Daughters from stressed mothers grew more slowly early in life, which should reduce their future reproductive opportunities, but exhibited more cooperative behaviour that should directly benefit their mother. This highlights the important role of sociality in the evolution of maternal effects.

Introduction

Parental effects are a mechanism of trans-generational phenotypic plasticity that

occurs when the parental phenotype or parental environment modifies offspring

characteristics (1). Parental effects can increase the survival or reproduction of

offspring, thereby elevating the direct fitness of both offspring and parents (2-6).

Alternatively, parental effects can increase parental fitness, but at some cost to the

fitness of their offspring (7-8) – a process regarded as a type of parental manipulation

(9-12) or 'selfish parental effect' (13). For example, in mammals, the optimal birth weight

or litter size often differs between mothers and offspring (14) and pregnant females

experiencing stressful environments may reallocate resources away from offspring and

towards themselves, so that their offspring are smaller or grow more slowly before

weaning (15). Despite these observations, it has also been suggested that selfish

parental effects may be rare and unstable because selection would be expected to

favour the evolution of resistance mechanisms in offspring (7, 11, 13, 16, 17).

Selfish parental effects may in fact be more likely in cooperatively breeding

species where philopatric offspring (subordinates) help to rear the subsequent offspring

of their parents or other close relatives. This is because parents may gain substantial

direct fitness benefits from delaying the development of their offspring if this causes

them to invest in alloparental care directed at the parent's subsequent offspring (9-10).

In addition, the costs of selfish parental effects to offspring could be reduced in these

circumstances, as offspring will gain indirect fitness benefits by contributing to raising

the subsequent offspring of their parents (18). For example, laboratory studies of

eusocial insects suggest the possibility that selection will favour the evolution of alleles

that enable mothers to increase the helping behaviour of their offspring while

simultaneously reducing their capabilities of reproducing on their own (19-20; but see

21). In addition, it is often the case that the probability that parents will rear young

successfully is substantially higher than the probability that inexperienced and

subordinate females will do so, increasing the inclusive fitness benefits of helping

versus breeding to offspring (22).

To date, empirical field tests of how parental effects shape the helping behaviour

of offspring are rare (23) and studies of selfish parental effects have mostly focused on

non-social species (13, 24). Here, we report the results of experiments designed to test

the hypothesis that elevated maternal glucocorticoid (stress hormone) levels reduce the

potential for offspring to have direct reproductive opportunities and causes them to

exhibit more cooperative behaviour. In a 3-year field study, we experimentally elevated

maternal stress by treating pregnant dominant female meerkats with glucocorticoids

(cortisol) and tracking the growth, stress physiology, and cooperative behaviour of their

offspring from birth until ~18 months of age, compared to those from control litters. We

manipulated maternal glucocorticoids because they are known to cause mothers to

reallocate energy away from offspring and towards themselves (15), indicating that they

may function as a mediator of selfish maternal effects. Changes in maternal

glucocorticoids have also previously been shown to delay the dispersal of offspring as

well as influence the parental care behaviour of offspring (25-26), both traits that are

important in cooperative breeders where philopatric offspring exhibit alloparental care

behaviour towards juveniles.

To identify if the exposure of mothers to heightened glucocorticoids reduced

reproductive success of their offspring, we examined if offspring from mothers treated

with cortisol during pregnancy grew slower or were smaller early in life. In meerkats, the

rate of early life growth and body mass is closely linked to future direct fitness through

its effects on survival, foraging success, adult body mass (27-29), as well as the

probability of acquiring dominance (30-31) and other direct reproductive opportunities

(32). As elevated exposure to maternal glucocorticoids in some mammals may reduce

offspring size and growth early in life (15), we predicted that offspring from mothers

treated with cortisol during pregnancy would be smaller early in life and would

consequently contribute more to cooperative activities than controls.

Secondly, we determined if offspring from mothers treated with cortisol during

pregnancy subsequently increased their contributions to two types of cooperative

behaviours: pup rearing ("babysitting": 33) and food provisioning during the period when

the pups are foraging with their natal group but are not yet nutritionally independent

("pup feeding": 34). If offspring from mothers treated with cortisol during pregnancy

exhibit more of either of these two types of alloparental care, this should increase both

parental fitness and the indirect fitness of offspring because subsequent offspring that

receive more alloparental care should grow faster or have higher early life survival (27,

32, 35, 36).

To assess the mechanism by which elevated exposure to maternal stress may

affect the alloparental care behaviour of offspring, we repeatedly measured plasma

cortisol and faecal glucocorticoid metabolite (fGCM) concentrations of offspring from

when they were approximately 1 to 18 months of age to identify how our manipulations

affected their neuroendocrine stress axes (glucocorticoid output). Elevated maternal

glucocorticoids can cause long-term changes in the neuroendocrine stress axis of

offspring (37) and elevated activity of the neuroendocrine stress axis in meerkats can

reduce their contributions to alloparental care (38). We therefore predicted that if

offspring born to mothers treated with cortisol during pregnancy exhibited more

alloparental care behaviour compared to controls, they would also have reduced plasma

cortisol and fGCM concentrations.

Results

127   *Effects of treatments on litter characteristics and offspring survival*

128   There was no evidence that the treatment of pregnant females with cortisol

affected their ability to maintain litters to term or the survival of their pups prior to

emergence from the natal burrow (Tables S1-S2). The number of pups surviving to

emergence from the natal burrow or 3, 6, or 12 months of age and the litter sex ratio

were not different among litters from cortisol-treated (31 pups from 10 litters produced

by 9 dominant females), control (25 pups from 6 litters produced by 6 females), or

untreated (195 pups from 52 litters produced by 21 dominant females) females (Table

S1-S2).

136   *Effects of treatments on offspring early life body mass*

137   The effects of the treatments on body mass from initial emergence to nutritional

independence (1-3 months) differed between daughters and sons and depended upon

the age of offspring (Table S3). Daughters (but not sons) from cortisol treated litters

were significantly smaller from 1-3 months than those from control litters especially as

they approached nutritional independence (daughters: age × treatment, t = -4.17, P <

0.0001; sons: t = -1.48, P = 0.14), but were of similar body mass compared to those

from untreated litters (daughters: age × treatment, t = 0.65, P = 0.51; sons: t = -0.52, P

= 0.6, Table S3, Fig. 1). This difference in body mass between daughters from mothers

treated with cortisol during pregnancy and those from control litters increased from one

to three months, as indicated by the significant interaction between age and control

treatment (Table S3). Daughters, but not sons, from control litters were larger than

those from untreated litters, especially as they got older (daughters: age × treatment, t =

-4.24, P < 0.0001; sons: t = 1.35, P = 0.18).

150   *Effects of treatments on offspring cooperative behaviour*

151   The effects of the maternal treatments on babysitting behaviour of offspring

depended upon the age and sex of the offspring (Table S4). Babysitting contributions in

daughters from mothers treated with cortisol during pregnancy slightly, but significantly

increased and rose more rapidly with increasing age of the babysitter than those from

control litters (age × treatment, z = -2.89, P = 0.0039) but not untreated litters (age ×

treatment, z = 1.88, P = 0.06; Table S4, Fig. 2). Babysitting contributions in sons from

mothers treated with cortisol during pregnancy showed a similar tendency to slightly

increase more rapidly with age compared to those from control litters, but this difference

was not significant (age × treatment, z = -1.92, P = 0.055). Further, age-related

increases in babysitting contributions between males from mothers treated with cortisol

during pregnancy and untreated litters did not differ (age × treatment, z = -0.03, P =

0.97; Table S4, Fig. 2). Comparisons of the magnitude of effect sizes showed that the

interaction between age and litter treatment had a larger effect on babysitting

contributions in daughters but not sons than other variables known to impact babysitting

contributions, such as foraging success, age-related body mass, or group size (Table

S4).

The effects of the maternal treatments on pup feeding depended upon the sex of

the offspring, but not their age (Table S5). Daughters but not sons from mothers treated

with cortisol during pregnancy exhibited significantly more pup feeding contributions

than those from control litters (females: z = -3.12, P = 0.00018; males: z = -1.14, P =

0.25) or untreated litters (females: z = -3.49, P = 0.0005, sons: z = -1.03, P = 0.3, Table

S5, Fig. 3). Notably, the magnitude of effect size of litter treatment for daughters was

much larger than other variables known to impact babysitting contributions such as

foraging success, age-related body mass, and group size (Table S5).

175   *Effects of treatments on offspring stress physiology*

Daughters and sons from mothers treated with cortisol during pregnancy had

lower plasma cortisol and fGCM concentrations as they became older compared to

those from control litters. Plasma cortisol concentrations in daughters from mothers

treated with cortisol during pregnancy had lower plasma cortisol concentrations as they

became older compared to those from control litters but these differences were not

significant (age $\times$ treatment, t = -1.61, P = 0.11, Table S6, Fig. 3A). fGCM

concentrations in daughters were significantly lower than those from control litters as

they became older (age $\times$ treatment, t = -2.95, P = 0 .0033, Table S7, Fig. 4A). Sons

from mothers treated with cortisol during pregnancy had significantly lower plasma

cortisol concentrations as they became older compared to those from control litters (age

$\times$ treatment t = -2.59, P = 0.01, Table S6, Fig. 3B) but similar fGCM concentrations

compared to those from control litters as they became older (age × treatment, t = -0.49,

P = 0.62, Table S7, Fig. 4B).

Discussion

We found some support for our hypothesis that elevated maternal stress would

reduce the potential for offspring to have direct reproductive opportunities and would

therefore shift them towards exhibiting more cooperative behaviour that should increase

their indirect fitness. Daughters, but not sons, from mothers treated with cortisol during

pregnancy grew more slowly early in life and exhibited more babysitting and pup

feeding behaviour as they became older compared to controls. Other than offspring

survival (Table S2), we were unable to quantify the direct and indirect fitness of

offspring from control or cortisol-treated litters, but early life growth or body mass (which

we measured here) is closely linked to direct fitness opportunities in daughters.

Previous studies in meerkats show that female, but not male, offspring that grow faster

from 1-3 months are more likely to acquire the dominant breeding position (31), perhaps

because offspring that grow faster from 1-3 months are heavier later in life (32, 39, 40),

and heavier females are more likely to acquire a vacant dominant breeding position (30,

32). As such, daughters, but not sons, from mothers with elevated cortisol levels during

pregnancy should have reduced future direct fitness opportunities and therefore

increase their investment in behaviours that elevate their indirect fitness. Our results are

consistent with studies in other taxa that suggest that individuals adjust their

contributions to cooperative behaviour according to their future reproductive potential.

For example, wasps whose probability of acquiring the dominant breeding position was

experimentally increased exhibited significantly less helping behaviour (41). Similarly, in

cooperatively breeding fish, subordinates will reduce their helping investment

immediately prior to dispersal from their natal group where they attempt to reproduce on

their own rather than stay in their natal group and queue for dominance (42).

The effects of maternal stress on offspring should lead to substantial fitness

benefits to mothers. Daughters from mothers treated with cortisol during pregnancy

exhibited more alloparental care compared to controls, such that subsequent offspring

produced in groups with offspring from cortisol-treated litters should have received more

alloparental care behaviours. Because offspring that receive more alloparental care

grow faster early in life or are larger later in life (32, 39), the presence of offspring from

cortisol-treated litters should increase the direct fitness of dominant breeders and the

indirect fitness of the cortisol-treated offspring. Taken together, our results suggest that

this stress-mediated maternal effect reduced the direct fitness opportunities of

daughters by reducing their early life growth but they compensated by increasing their

investment in indirect fitness opportunities (helping to rear non-descendent offspring).

This is in line with theoretical predictions that parental manipulation of the cooperative

behaviour of offspring can evolve if the costs of resisting the parental effect are high and

inclusive fitness benefits of helping rear subsequent offspring are high (18).

Control dominant females that were fed during pregnancy produced daughters

that were larger during early development (1-3 months) compared to daughters from

cortisol-treated or untreated litters. Although mothers that were treated with cortisol

during pregnancy received the same amount of food as controls, daughters and sons

from mothers fed cortisol during pregnancy did not differ in body mass compared to

those from untreated litters. This indicates that the additional food provided to dominant

females during pregnancy had the potential to increase growth, but the added cortisol

prevented those gains in body mass. This has implications for understanding the fitness

consequences of maternal stress on offspring growth trajectories (15, 43) because our

results show that elevated circulating glucocorticoid levels in pregnant females in the

absence of energetic constraints induced reductions in the early life growth of offspring.

This supports the hypothesis that maternal stress levels during offspring development

act as a cue that induces plasticity in offspring growth rather than simply mediating the

effects of energetic constraints.

The reductions in the activity of the neuroendocrine stress axis of daughters may

have potentiated the increased alloparental care behaviour that we observed.

Compared to daughters from control litters, daughters from mothers treated with cortisol

during pregnancy exhibited more babysitting as they became older, more overall pup

feeding, and they also had lower plasma cortisol and fGCM concentrations. Males from

mothers treated with cortisol during pregnancy had significantly lower plasma cortisol

concentrations, but not fGCM concentrations as they got older and also tended to

exhibit more babysitting as they aged. The activity of the neuroendocrine stress axis is

closely linked to an array of social behaviours (44) and our recent work shows that

elevated activity of the neuroendocrine stress axis reduces babysitting in both females

and males and decreases pupfeeding in females (38). Together, this supports the

hypothesis that the mechanism by which early life stress increases the cooperative

behaviour of daughters is by dampening the activity of their neuroendocrine stress axis.

Our results show that the effects of maternal stress on offspring growth,

physiology, and behaviour were greater in daughters than in sons, which adds to

biomedical (45-46) and ecological (47-49) studies that highlight how early life conditions

or maternal stress levels can have sex-specific consequences for offspring. In

meerkats, there may be added benefits for the dominant female for altering the

cooperative behaviour of daughters compared to sons; daughters exhibit more

cooperative behaviour than sons (35) and are more responsive to the begging calls of

subsequent offspring that they provision with food (50). More broadly, sex-differences in

natal dispersal may cause these differential responses to parental effects. In meerkats,

subordinate males voluntarily disperse from their natal group to look for receptive

females but can return to their natal group whereas subordinate females rarely

voluntarily disperse from their natal group (51). In our case and in others (43), the more

philopatric sex (females) is more sensitive to early life conditions, which may be due to

differential costs of parental modification between the philopatric and dispersing sex. If

parental effects have long-term consequences on offspring characteristics, as we show

here, there may be an increased degree of mismatch between the phenotype of the

dispersing sex and the postnatal environment where individuals eventually settle. If this

mismatch has fitness costs, this should select for individuals from the dispersing sex to

be less responsive to cues from the parental phenotype or environment.

Our results provide some support for the hypothesis that parents may alter the

cooperative tendencies of their offspring by manipulating the characteristics of their

offspring (9-10). Explanations regarding the evolutionary origins of cooperative

behaviour involve nepotism or kin selection (52), mutualisms or reciprocity (53), but few

studies have tested the "parental manipulation" hypothesis proposed by Alexander (9).

Some studies show that alleles that increase maternal fitness at the expense of the

direct fitness of offspring can evolve (19) and that cooperative breeders may bias

investment towards offspring that exhibit more cooperative behaviour (54). Our study

provides evidence that environmental stressors and parental effects can modify the

cooperative tendencies of their offspring.

Finally, our results have two implications for theoretical models examining the

evolution of parental effects. First, given the sex-specificity of parental effects, our

results challenge the conclusions of models examining the evolution of parental effects

that assume that all offspring are equally sensitive to the parental effect (16), or those

that assume that the benefits of exhibiting the phenotype resulting from the parental

effect are equal for all offspring (18). Second, selfish parental effects are thought to be

relatively rare (8, 13) and theory (16-17) and empirical studies showing sex-specific

responses to early life stress (45-46) indicate that offspring can become resistant to

such selfish parental effects. However, the evolution of selfish parental effects may be

dependent upon the social environment (24) especially if the selfish parental effect

influences the expression of alloparental care behaviour of offspring and therefore

increases the indirect fitness of offspring. Our results provide an example whereby a

stress-mediated maternal effect reduces the direct fitness of daughters (by reducing

their early life growth), but presumably increases the direct fitness of mothers and

indirect fitness of daughters by elevating their cooperative behaviour.

Methods

Study site

We studied free-living meerkats at the Kuruman River Reserve (26° 58' S, 21°

49' E) in the Northern Cape, South Africa from 2014-2017. Individuals were marked

uniquely (with PIT tags: Identipet®, Johannesburg, South Africa) as well as dye marks

so that they could be identified. Groups were visited for ~4-8 hours per day ~4-6 times

per week throughout each year of study and sometimes more frequently such as when

there were pups being babysat.

Experimental manipulations of dominant females

Dominant females in each group were identified via behavioural observations

(55). During each visit, we recorded which individuals were present as well as their body

mass. Meerkats were weighed on a portable balance each morning before foraging, 2-4

hrs after foraging was initiated, and immediately prior to when foraging ended (56). The

pregnancy status of dominant females was determined visually (distended abdomen) as

well as noting a constant increase in their body mass.

Dominant females were treated with either a cortisol solution or a control oil

vehicle when they were pregnant by feeding them food containing one of these two

treatments. We initially offered experimental animals hard boiled egg with added cortisol

but found that they rejected all foods that contained added cortisol with the exception of

scorpions. We consequently fed experimental females with cortisol (10 mg/kg of

hydrocortisone, Sigma H4126), that were dissolved in 100 μl of 100% coconut oil and

injected into a dead scorpion (*Opistophthalmus* spp.). Control females were fed a dead

scorpion that was injected with 100 μl of 100% coconut oil. A previous study using the

same protocol showed that meerkats that were fed cortisol had significantly higher

plasma cortisol and fGCM concentrations than control animals and these increases

were within a biologically-relevant range (38). This indicated that our treatment causes

the exogenous glucocorticoids that we feed the meerkats to enter their bloodstream and

leads to sustained increases in their circulating glucocorticoid concentrations.

We aimed to experimentally increase the glucocorticoid concentrations of

pregnant dominant females from when they were first confirmed to be pregnant (second

half of gestation) until parturition. Meerkats have a ~70 d gestation period so we aimed

to treat them with glucocorticoids from approximately 35-70 d during gestation. In

reality, females that successfully produced a litter where pups emerged from the natal

burrow were treated with cortisol for 12-36 days prior to birth (n=10 litters from 9

females, mean = 23.7 d, median = 23.5 d) whereas controls were fed for 12-58 d prior

to birth (n=6 litters from 6 females, mean = 30 d, median = 20.5 d).

Females were randomly allocated to the treatments. Across the three years of

this study, we produced a total of 16 cortisol-treated or control litters produced by 12

dominant females in 12 different social groups. Three of the females experienced both

the control and cortisol treatments at different time points of the experiment, whereas

one female experienced the control treatment once and the cortisol treatments twice.

For these latter females treated twice, the order of treatments was randomly selected.

We conducted these experiments over the course of three years: 13 litters from 10

females in 2014 (April-December 2014, 3 litters aborted), 5 litters from 5 females in

2015 (February-July 2015, 1 litter aborted), and 3 litters from 3 females in 2016 (July

2016).

To provide an additional comparison group to investigate how our treatments (fed

during pregnancy or fed cortisol during pregnancy) affected offspring survival, body

mass, and cooperative behaviour, we also monitored these traits in offspring produced

by dominant females that were untreated during pregnancy (n = 52 litters from 21

dominant females). For our analyses of how the treatments affected offspring survival

and body mass, the untreated offspring were those from litters produced by dominant

females in other meerkat groups in our same study area and were born during our study

(hereafter called 'untreated' litters). We assessed the contributions of offspring from

mothers treated with cortisol during pregnancy to two cooperative behaviours

(babysitting and pup feeding) compared to those from control litters but also to other

group members from untreated litters. We did not have data from offspring from

untreated litters when we assessed how our treatments affected their plasma cortisol or

fGCM concentrations.

Quantifying early life growth of offspring

Meerkat pups typically first emerge from their natal burrow approximately 21-30 d

after birth. Meerkat groups and dominant females were monitored daily around the

estimated date of parturition and birth dates were estimated according to the change in

the physical appearance of the dominant female, a large drop in body mass overnight,

and group members exhibiting babysitting behaviour at the sleeping burrow. Burrows

containing pups were monitored each day and, when pups emerged, they were uniquely

marked by trimming small sections of hair before permanent PIT tags could be applied.

Pups were weighed each time we visited the groups on a portable balance in the

morning after group members emerged from their sleeping burrow (as above).

Quantifying cooperative behaviour of offspring

We measured the babysitting (controls: 195-655 d; cortisol: 184-655 d; untreated:

155-655 d) and pup feeding (controls: 220-635 d; cortisol: 184-655 d; untreated: 155-

626 d) contributions of offspring from cortisol-treated and control litters when they were

>6 months of age until death or disappearance. We visited sleeping burrows containing

pups every day in the morning and recorded the identity of the attending babysitters. As

we have done previously (33, 57), we calculated relative babysitting contributions of

each individual meerkat for each litter by dividing the total number of days an individual

babysat a litter over the total number of days that this specific litter had a babysitter.

Pup feeding behaviour for each pup produced by the dominant females in the different

treatment groups was estimated using *ad libitum* sampling (34). When the social group

contained pups (up to 90 d of age), we recorded all pup-feeding events, which are

visually and acoustically conspicuous to observers (58). We used the probability of pup

feeding occurring and corrected for variation in observation time (see below).

Quantifying plasma cortisol concentrations from offspring

We obtained plasma samples from offspring from cortisol-treated and control

litters approximately every 3 months from first emergence from the burrow (~1 month)

until ~18 months of age (controls: 20-548 d; cortisol-treated: 25-559 d). Capture and

blood processing procedures are described elsewhere (59-60). We measured total

plasma cortisol concentrations using a previously validated assay (Coat-a-Count,

Siemens Diagnostic Products Corporation, Los Angeles, USA: validation described in

59). The sensitivity of the assay was 1.9 ng/ml and cross reactivity to other hormones

was 76% with prednisolone, 11.4% with 11-deoxycortisol, 2.3% with prednisone and

<1% with aldosterone, corticosterone, cortisone, estriol, estrone and pregnenolone.

Intra-assay coefficient of was 7% (n = 20 samples). Inter-assay coefficient of variation

for a low control was 8% (78.5 ± 6.3 nmols/l, n = 5 assays) and 2.8% for a high control

(187 ± 5.3 ng/ml, n = 5 assays).

Quantifying fGCM concentrations from offspring

We collected faecal samples from offspring from cortisol-treated and control

litters opportunistically during behavioural observations over the course of the study

(controls: 25-356 d; cortisol-treated: 32-326 d). Faecal samples were processed as

described previously using a methanol extraction to remove fGCMs (61-62).

Immunoreactive fGCM concentrations were determined using an enzyme immunoassay

measuring cortisol metabolites with a 5β-3α,11β-diol-structure (11β-

hydroxyetiocholanolone) already established for monitoring fGCM alterations in

meerkats (62). fGCMs measured reflect average adrenal cortisol production over the

previous ~24 to 48 hr period (62). Detailed assay characteristics, including full

descriptions of the assay components and cross-reactivities, are found elsewhere (63)

The sensitivity of the assay was 1.2 ng/g dry weight and intra-assay coefficients of

variation (CV) determined by repeated measurements of high and low value quality

controls were 6.9% and 7.4%, respectively, and inter-assay CV values were 11.5% and

15.9% (n = 29 assays). fGCM concentrations below are expressed as ln ng/g dry faecal

matter.

Statistical analyses

We used generalized (binomial errors) or linear mixed-effects models (LMMs) to

examine how our treatments affected the probability that the litter was aborted, litter size

and sex ratio at emergence from the burrow, and the proportion of the litter that survived

to emergence from the burrow, independence (90 d of age: 29), and 6 or 12 months of

age. We focused on addressing whether the cortisol-treated litters differed from our

control or untreated litters. These models included a fixed effect for date of birth of the

litter and random intercept terms for dominant female identity and year (as the

experiments were conducted over 3 years). None of the GLMMs were overdispersed

(Table S2).

We used a LMM to investigate how the maternal treatments affected offspring

body mass from first emergence from their natal burrow (~1 month) to 3 months of age

when the pups are typically foraging independently (29, 34). Morning body mass (in

grams) was the response variable with the fixed effects of maternal treatment (cortisol-

treated, control, or untreated), pup sex, pup age, litter size at burrow emergence, group

size, group size^2, total rainfall in the previous 60 days, two measures of seasonality

(sine and co-sine functions of day of weight measure: see 56), and two three-way

interactions between sex, treatment, age or age^2. Random intercept terms for year and

the identity of the individual nested in litter, nested in dominant female identity, nested in

group were also included in this model. Fixed and random effects included in these

models were based upon previous studies investigating meerkat body mass and growth

from 1-3 months (28, 31, 56). Only individuals that survived to 90 d were included in

these analyses.

We assessed how the treatments affected the relative babysitting and pup

feeding contributions of subordinates when they were >6 months (as they rarely do

alloparental care behaviour <6 months: 35) from cortisol-treated, control, and untreated

litters. In these generalized linear mixed-effects models (GLMM, binomial errors), we

included a three-way interaction between treatment, sex, and age of the subordinate to

assess if the effects of the treatments on babysitting or pup feeding varied according to

the sex or age of the subordinate, as contributions to cooperative behaviour in meerkats

are known to vary according to subordinate sex and age (35). To account for

differences in observation time, we included a co-variate for the number of days the

litter was babysat (babysitting length) and the number of days the subordinate was

observed in the group during babysitting as well as the total time spent observing the

group during pup feeding (observation time). We included a range of co-variates (see

Tables S4-S6) that have been previously documented to affect contributions to

babysitting and pup feeding, including age, foraging success, body mass, and group

size (Clutton-Brock et al., 2000, 2002, 2003; Carlson et al., 2006*a*; Dantzer et al.,

2017*a*). Relatedness between the subordinate and the litter being babysat was not

included as it has not been shown to impact babysitting or pup feeding contributions

(27, 57) and most of the litters in our entire dataset were produced by the mother or full

sibling of the subordinate. Random intercept terms for year and the identity of the

individual, and litter being babysat or pup fed nested within the group where the litter

was being babysat or pup fed. Overdispersion was not an issue for our GLMM for

babysitting as indicated by the goodness of fit test (Pearson χ^2 = 147.1, df = 154, P =

0.64, using package aods3: 64) but our GLMM for pup feeding was initially

overdispersed (Pearson χ^2 = 310, df=165, P < 0.0001) so we included an observation

level random intercept term.

We used two separate LMMs to assess how our manipulations affected plasma

cortisol and fGCMs in offspring from cortisol-treated and control litters. Each model

included fixed effects for maternal treatment, pup sex and age, time of day and year that

the sample was acquired (2014 or 2015), and random intercept terms for identity of

individual nested in their birth litter and group. In the model for plasma cortisol

concentrations, we also included a linear and second order fixed effect for the time it

took to acquire the blood sample to control for any variation in plasma cortisol

concentrations due to restraint stress (60). Year was included as a fixed effect here

because we only had samples from two separate years. We included covariates

associated with the individual meerkat and weather or social group characteristics that

are known to affect plasma cortisol (60) or fGCM (62) concentrations (see Tables S6-

S7).

We used R (version 3.4.3: 65) for all of our statistical analyses. R package lme4

(version 1.1-14: 66) was used for LMMs and P values were estimated using lmerTest

(version 2.0-33: 67). A graphical approach was used to confirm normality and

homoscedasticity of residuals and to confirm there were no observations with high

leverage (68). All continuous variables were standardized to a mean of 0 and SD of 1.

Ethics: All protocols used in our experiments were approved by the Animal Ethics

Committee at the University of Pretoria (Pretoria, South Africa: #EC031-13, #EC047-16)

and the Northern Cape Department of Environment and Nature Conservation Research

(South Africa: FAUNA 050/2013, FAUNA 192/2014, FAUNA 1020/2016).

Acknowledgements: We thank M. Manser for her contributions to maintain the long-

term Kalahari Meerkat Project and her contributions to this specific project. Thanks to P.

Roth, S. Bischoff-Mattson, L. Howell, J. Samson, D. Cram, and N. Thavarajah and staff

at the Kalahari Meerkat Project for helping to organize the project, and numerous

volunteers who contributed to data collection. Thanks to M. Manser and S. Guindre-

Parker for helpful comments. Thanks to the Northern Cape Conservation Authority for

permission to conduct research and the Kotze family for allowing us to study meerkats

on their property.

Authors' Contributions: B.D. and T.H.C-B planned the study, B.D. designed the

experiments, B.D., C. Dubuc, D.L.C, T.H.C-B, D.G., I.BG., N.B., M.H., A.G., and C.

Duncan coordinated and collected data, B.D. conducted analyses and produced figures,

B.D. and T.H.C-B authored manuscript with contributions from all authors.

Date, code and materials: All data will be archived on Data Dryad. All code for

statistical analyses is available from first author.

Competing interests: We have no competing interests.

Funding: Research was supported by grants from National Environment Research

Council (RG53472) and the European Research Council (294494) to T.H.C-B. I.B.G.

was funded by the Swiss National Science Foundation (grant no. 31003A_13676 to M.

Manser). NCB was supported by a DST-NRF SARCHI chair of Mammal Behavioural

References

1) Wolf JB, Wade MJ (2009) What are maternal effects (and what are they not)?

Philosophical Transactions of the Royal Society B: Biological

Sciences, *364*(1520), 1107-1115.

2) Kirkpatrick M, Lande R (1989) The evolution of maternal characters. *Evolution*, *43*(3),

485-503.

3) Mousseau TA, Fox CW (Eds.) (1998) *Maternal Effects as Adaptations*. Oxford

University Press.

4) Agrawal AA, Laforsch C, Tollrian R (1999) Transgenerational induction of defences in

animals and plants. *Nature* 401, 60-63

5) Dantzer B, Newman AE, Boonstra R, Palme R, Boutin S, Humphries MM, McAdam

AG (2013) Density triggers maternal hormones that increase adaptive offspring

growth in a wild mammal. *Science*, 1235765.

6) Duckworth RA, Belloni V, Anderson SR (2015) Cycles of species replacement

emerge from locally induced maternal effects on offspring behavior in a

passerine bird. *Science*, *347*(6224), 875-877.

7) Uller T (2008) Developmental plasticity and the evolution of parental effects. *Trends*

in Ecology & Evolution, *23*(8), 432-438.

8) Uller T, Nakagawa S, English S (2013) Weak evidence for anticipatory parental

effects in plants and animals. *Journal of Evolutionary Biology*, *26*(10), 2161-2170.

9) Alexander, RD, (1974) The evolution of social behavior. *Annual Review of Ecology*

and Systematics, *5*(1), 325-383.

10) Michener CD, Brothers DJ 1974 Were workers of eusocial Hymenoptera initially

altruistic or oppressed? *Proceedings of the National Academy of Science* USA

71, 671–674

11) Müller W, Lessells CKM, Korsten P, Engelhardt NV (2007) Manipulative signals in

family conflict? On the function of maternal yolk hormones in birds. *The American

Naturalist*, *169*(4), E84-E96.

12) Schwarzkopf L, Andrews RM (2012) Are moms manipulative or just selfish?

Evaluating the "maternal manipulation hypothesis" and implications for life-history

studies of reptiles. *Herpetologica*, *68*(2), 147-159.

13) Marshall DJ, Uller T (2007) When is a maternal effect adaptive?. *Oikos*, *116*(12),

1957-1963.

14) Wilson AJ, Pilkington JG, Pemberton JM, Coltman DW, Overall AD, Byrne KA,

Kruuk, LEB (2005) Selection on mothers and offspring: whose phenotype is it

and does it matter? *Evolution*, *59*(2), 451-463.

15) Berghänel A, Heistermann M, Schülke O, Ostner J (2017) Prenatal stress

accelerates offspring growth to compensate for reduced maternal investment

across mammals. *Proceedings of the National Academy of Sciences*, *114*(50),

E10658-E10666.

16) Uller T, Pen I (2011) A theoretical model of the evolution of maternal effects under

parent–offspring conflict. *Evolution*, *65*(7), 2075-2084.

17) Kuijper B, Johnstone RA (2018) Maternal effects and parent–offspring

conflict. *Evolution*, *72*(2), 220-233.

18) González-Forero M, Gavrilets S (2013) Evolution of manipulated behavior. *The

American Naturalist*, *182*(4), 439-451.

19) Wade MJ (2001) Maternal effect genes and the evolution of sociality in haplo-diploid

organisms. *Evolution*, *55*(3), 453-458.

20) Linksvayer TA, Wade MJ (2005) The evolutionary origin and elaboration of sociality

in the aculeate Hymenoptera: maternal effects, sib-social effects, and

heterochrony. *The Quarterly Review of Biology*, *80*, 317-336.

21) Bourke AF, Franks NR (1995). *Social Evolution in Ants*. Princeton University Press.

22) Clutton-Brock TH (2016) *Mammal Societies*. Wiley-Blackwell

23) Russell A F, Lummaa V (2009) Maternal effects in cooperative breeders: from

hymenopterans to humans. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *364*(1520), 1143-1167.

24) Taborsky B (2017) Developmental plasticity: preparing for life in a complex

world. *Advances in the Study of Behavior*, *49*, 49-99.

25) Carter CS, Roberts RL. (1997) The psychobiological basis of cooperative breeding

in rodents. In *Cooperative Breeding in Mammals* (Eds.: NG Solomon, JA

French), Cambridge University Press, Pp. 231-266.

26) De Fraipont M, Clobert J, Alder HJ, Meylan S (2000) Increased pre-natal maternal

corticosterone promotes philopatry of offspring in common lizards *Lacerta*

vivipara. *Journal of Animal Ecology*, *69*(3), 404-413.

27) Clutton-Brock TH, Brotherton PNM, O'riain MJ, Griffin AS, Gaynor D, Kansky R,

Sharpe L, McIlrath GM (2001) Contributions to cooperative rearing in

meerkats. *Animal Behaviour*, *61*(4), 705-710.

28) Russell AF, Clutton-Brock TH, Brotherton PN, Sharpe LL, Mcilrath G, Dalerum FD,

Cameron EZ, Barnard JA (2002). Factors affecting pup growth and survival in co-

operatively breeding meerkats *Suricata suricatta*. *Journal of Animal*

Ecology, *71*(4), 700-709.

29) Thornton A (2008) Social learning about novel foods in young meerkats. *Animal*

Behaviour, *76*(4), 1411-1421.

30) Hodge SJ, Manica A, Flower TP, Clutton-Brock TH (2008) Determinants of

reproductive success in dominant female meerkats. *Journal of Animal*

Ecology, *77*(1), 92-102.

31) English S, Huchard E, Nielsen JF, Clutton-Brock TH (2013) Early growth,

dominance acquisition and lifetime reproductive success in male and female

cooperative meerkats. *Ecology and Evolution*, *3*(13), 4401-4407.

32) Russell AF, Young AJ, Spong G, Jordan NR, Clutton-Brock TH (2007) Helpers

increase the reproductive potential of offspring in cooperative

meerkats. *Proceedings of the Royal Society of London B: Biological*

Sciences, *274*(1609), 513-520.

33) Clutton-Brock TH, Gaynor D, Kansky R, MacColl ADC, McIlrath G, Chadwick P,

Brotherton PNM, O'rian JM, Manser M, Skinner JD (1998) Costs of cooperative

behaviour in suricates (*Suricata suricatta*). *Proceedings of the Royal Society of*

London B: Biological Sciences, *265*(1392), 185-190.

34) Brotherton PNM, Clutton-Brock TH, O'riain MJ, Gaynor D, Sharpe L, Kansky R,

McIlrath GM (2001) Offspring food allocation by parents and helpers in a

cooperative mammal. *Behavioral Ecology*, *12*(5), 590-599.

35) Clutton-Brock TH, Russell AF, Sharpe LL, Young AJ, Balmforth Z, McIlrath GM

(2002). Evolution and development of sex differences in cooperative behavior in

meerkats. *Science*, *297*(5579), 253-256.

36) Bell MBV, Cant MA, Borgeaud C, Thavarajah N, Samson J, Clutton-Brock TH

(2014) Suppressing subordinate reproduction provides benefits to dominants in

cooperative societies of meerkats. *Nature Communications* 5, 4499

37) Harris A, Seckl J (2011) Glucocorticoids, prenatal stress and the programming of

disease. *Hormones and Behavior*, *59*(3), 279-289.

38) Dantzer B, Goncalves IB, Spence-Jones HC, Bennett NC, Heistermann M,

Ganswindt A, Dubuc C, Gaynor D, Manser MB, Clutton-Brock TH 2017. The

influence of stress hormones and aggression on cooperative behaviour in

subordinate meerkats. *Proc. R. Soc. B*, *284*(1863), 20171248.

39) Clutton-Brock TH, Russell AF, Sharpe LL, Brotherton PNM, McIlrath GM, White S,

Cameron EZ (2001) Effects of helpers on juvenile development and survival in

meerkats. *Science*, *293*(5539), 2446-2449.

40) Huchard E, English S, Bell MB, Thavarajah N, Clutton-Brock T (2016) Competitive

growth in a cooperative mammal. *Nature*, *533*(7604), 532.

41) Field J, Cronin A, Bridge C (2006) Future fitness and helping in social

queues. *Nature*, *441*(7090), 214.

42) Zöttl M, Chapuis L, Freiburghaus M, Taborsky M (2013). Strategic reduction of help

before dispersal in a cooperative breeder. *Biology Letters*, *9*(1), 20120878.

43) Sheriff MJ, Bell A, Boonstra R, Dantzer B, Lavergne SG, McGhee KE, MacLeod KJ,

Winandy L, Zimmer C, Love OP (2017) Integrating ecological and evolutionary

context in the study of maternal stress. *Integrative and Comparative*

Biology, *57*(3), 437-449.

44) Raulo A, Dantzer B. (2018) Associations between glucocorticoids and sociality

across a continuum of vertebrate social behaviour. *Ecology and Evolution* in

press (doi: 10.1002/ece3.4059).

45) Weinstock M (2007) Gender differences in the effects of prenatal stress on brain

development and behaviour. *Neurochemical Research*, *32*(10), 1730-1740.

46) Mueller, BR, Bale, TL (2008) Sex-specific programming of offspring emotionality

after stress early in pregnancy. *Journal of Neuroscience*, *28*(36), 9055-9065.

47) Wilkin TA, Sheldon BC (2009) Sex differences in the persistence of natal

environmental effects on life histories. *Current Biology*, *19*(23), 1998-2002.

48) Metzger DC, Schulte PM (2016) Maternal stress has divergent effects on gene

expression patterns in the brains of male and female threespine

stickleback. *Proc. R. Soc. B*, *283*(1839), 20161734.

49) Marshall HH, Vitikainen EI, Mwanguhya F, Businge R, Kyabulima S, Hares MC,

Inzani E, Kalema-Zikusoka G, Mwesige K, Nichols HJ, Sanderson JL, Thompson

FJ, Cant MA (2017) Lifetime fitness consequences of early-life ecological

hardship in a wild mammal population. *Ecology and Evolution*, *7*(6), 1712-1724.

50) English S, Kunc HP, Madden JR, Clutton-Brock TH (2008) Sex differences in

responsiveness to begging in a cooperative mammal. *Biology Letters*, *4*(4), 334-

337.

51) Clutton-Brock TH, Manser M (2016) Meerkats: cooperative breeding in the Kalahari.

Cooperative Breeding in Vertebrates (Eds.: WD Koenig, JL Dickinson).

Cambridge University Press, Pp. 294-317

52) Hamilton WD (1964) The genetical evolution of social behaviour. II. *Journal of*

Theoretical Biology, *7*(1), 17-52.

53) Trivers RL (1971) The evolution of reciprocal altruism. *The Quarterly Review of*

Biology, *46*(1), 35-57.

54) Griffin AS, Sheldon BC, West SA (2005) Cooperative breeders adjust offspring sex

ratios to produce helpful helpers. *The American Naturalist*, *166*(5), 628-632.

55) Kutsukake N, Clutton-Brock TH (2006) Aggression and submission reflect

reproductive conflict between females in cooperatively breeding meerkats

Suricata suricatta. *Behav. Ecol. Sociobiol.* **59**, 541-548. (doi: 10.1007/s00265-

005-0079-7)

56) English S, Bateman AW, Mares R, Ozgul A, Clutton-Brock TH (2014) Maternal,

social and abiotic environmental effects on growth vary across life stages in a

cooperative mammal. *Journal of Animal Ecology*, *83*(2), 332-342.

57) Clutton-Brock TH, Brotherton PNM, O'Riain MJ, Griffin AS, Gaynor D, Sharpe L,

Kansky R, Manser MB, McIlrath GM (2000). Individual contributions to

babysitting in a cooperative mongoose, *Suricata suricatta*. *Proceedings of the*

Royal Society of London B: Biological Sciences, *267*(1440), 301-305.

58) Kunc HP, Madden JR, Manser MB (2007) Begging signals in a mobile feeding

system: The evolution of different call types. *The American Naturalist* 170, 617-

624.

59) Carlson AA, Young AJ, Russell AF, Bennett NC, McNeilly AS, Clutton-Brock T

(2004). Hormonal correlates of dominance in meerkats (*Suricata*

suricatta). *Hormones and Behavior, 46*(2), 141-150.

60) Dantzer B, Bennett NC, Clutton-Brock TH (2017) Social conflict and costs of

cooperation in meerkats are reflected in measures of stress

hormones. *Behavioral Ecology, 28*(4), 1131-1141.

61) Heistermann M, Finke M, Hodges JK (1995) Assessment of female reproductive

status in captive-housed Hanuman langurs (*Presbytis entellus*) by measurement

of urinary and fecal steroid excretion patterns. *Am. J. Primatol.* 37, 275-284. (doi:

10.1002/ajp.1350370402)

62) Goncalves IB, Heistermann M, Santema P, Dantzer B, Mausbach J, Ganswindt A,

Manser, MB (2016) Validation of a fecal glucocorticoid assay to assess

adrenocortical activity in meerkats using physiological and biological stimuli. *PloS*

One, 11(4), e0153161.

63) Heistermann M, Palme R, Ganswindt A (2006) Comparison of different

enzymeimmunoassays for assessment of adrenocortical activity in primates

based on fecal analysis. *American Journal of Primatology* 68, 257-273.

64) Lesnoff M, Lancelot R (2013) aods3: analysis of overdispersed data using S3

methods. aods3 package version 0.4-1.

65) R Development Core Team. 2013. R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria; available at

www.r-project.org/.

66) Bates D, Maechler M, Bolker B, Walker S (2014) lme4: Linear mixed-effects models

using Eigen and S4. *R Package Version 1*(7), 1-23.

67) Kuznetsova A, Brockhoff PB, Christensen RHB (2016) lmerTest: Tests in Linear

Mixed Effects Models. R package version 2.0-30; available at http://CRAN.R-

project.org/package=lmerTest

68) Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid

common statistical problems. *Meth. Ecol. Evol.* **1**, 3-14. (doi: 10.1111/j.2041-

210X.2009.00001.x)

 Figure Legends

Figure 1. (A) Daughters but not (B) sons from mothers treated with cortisol during

pregnancy were significantly smaller from initial emergence from their natal burrow to

nutritional independence (~1-3 months) compared to those from control litters

(daughters: age $\times$ treatment, t = -4.17, P < 0.0001; sons: t = -1.48, P = 0.14), but not

untreated litters (daughters: age $\times$ treatment, t = 0.65, P = 0.51; sons: t = -0.52, P = 0.6,

Table S3). Data are body mass measures from offspring from cortisol-treated (females:

n = 373 estimates; males: n = 488), control (females: n = 215; males: n = 241), and

untreated litters (females: n = 1121; males: n = 2238). Raw data and regression lines

are shown (full results in Table S3).

Figure 2. Babysitting contributions of (A) daughters and (B) sons from mothers treated

with cortisol during pregnancy increased with age at a faster rate than those from

control (females: z = -2.89, P = 0.0039; males: z = -1.92, P = 0.055), but not untreated

("None") litters (females: z = 1.88, P = 0.06; males: z = -0.03, P = 0.97, Table S4). Data

are relative babysitting contributions from offspring from cortisol-treated (females: n = 15

estimates; males: n = 24), control (females: n = 15; males: n = 10), and untreated litters

(females: n = 49; males: n = 69). Raw data and regression lines are shown (full results

in Table S4).

Figure 3. Pup feeding contributions of (A) daughters, but not (B) sons, from mothers

treated with cortisol during pregnancy were significantly higher compared to from control

(females: z = -3.09, P = 0.0004; males: z = -1.15, P = 0.25) or untreated ("None") litters

(females: z = -3.47, P = 0.0005; males: z = -0.89, P = 0.37, Table S5). Data are relative

pup feeding contributions from offspring from cortisol-treated (females: n = 16

estimates; males: n = 26), control (females: n = 16; males: n = 10), and untreated litters

(females: n = 64; males: n = 71). Raw data are shown (full results in Table S5). Boxplots

show median (solid horizontal line), mean (grey diamonds), and first (25%) and third

(75%) quartiles.

Figure 4. (A) Daughters and (B) sons from mothers treated with cortisol during

pregnancy had lower plasma cortisol concentrations as they became older compared to

those from control litters, though the difference was only significant in males (daughters:

age x treatment, t = -1.61, P = 0.11; sons: age x treatment t = -2.59, P = 0.01, Table

S6). Data are residual plasma cortisol concentrations from offspring from cortisol-

treated (females: n = 64 samples; males: n = 92) and control (females: n = 89; males: n

= 104) litters. Residuals from a linear mixed-effects model (Table S6) are shown on y-

axis.

Figure 5. Faecal glucocorticoid metabolite (fGCM) concentrations in (A) daughters but

not (B) sons from mothers treated with cortisol during pregnancy were significantly

lower than those from control litters as they became older (daughters: age x treatment, t

= -2.95, P = 0 .0033; sons: age x treatment, t = -0.49, P = 0.62, Table S7. Data are

residual fGCM concentrations from offspring from cortisol-treated (females: n = 79

samples; males: n = 118) and control (females: n = 154; males: n = 201) litters.

Residuals from a linear mixed-effects model (Table S7) are shown on y-axis.

Figure 1. (A) Daughters but not (B) sons from mothers treated with cortisol during

pregnancy were significantly smaller from initial emergence from their natal burrow to

nutritional independence (~1-3 months) compared to those from control litters

(daughters: age × treatment, t = -4.17, P < 0.0001; sons: t = -1.48, P = 0.14), but not

untreated litters (daughters: age × treatment, t = 0.65, P = 0.51; sons: t = -0.52, P = 0.6,

Table S3). Data are body mass measures from offspring from cortisol-treated (females:

n = 373 estimates; males: n = 488), control (females: n = 215; males: n = 241), and

untreated litters (females: n = 1121; males: n = 2238). Raw data and regression lines

are shown (full results in Table S3).

Figure 2. Babysitting contributions of (A) daughters and (B) sons from mothers treated

with cortisol during pregnancy increased with age at a faster rate than those from

control (females: z = -2.89, P = 0.0039; males: z = -1.92, P = 0.055), but not untreated

("None") litters (females: z = 1.88, P = 0.06; males: z = -0.03, P = 0.97, Table S4). Data

are relative babysitting contributions from offspring from cortisol-treated (females: n = 15

estimates; males: n = 24), control (females: n = 15; males: n = 10), and untreated litters

(females: n = 49; males: n = 69). Raw data and regression lines are shown (full results

in Table S4).

Figure 3. Pup feeding contributions of (A) daughters, but not (B) sons, from mothers

treated with cortisol during pregnancy were significantly higher compared to from control

(females: z = -3.09, P = 0.0004; males: z = -1.15, P = 0.25) or untreated ("None") litters

(females: z = -3.47, P = 0.0005; males: z = -0.89, P = 0.37, Table S5). Data are relative

pup feeding contributions from offspring from cortisol-treated (females: n = 16

estimates; males: n = 26), control (females: n = 16; males: n = 10), and untreated litters

(females: n = 64; males: n = 71). Raw data are shown (full results in Table S5). Boxplots

show median (solid horizontal line), mean (grey diamonds), and first (25%) and third

(75%) quartiles.

Figure 4. (A) Daughters and (B) sons from mothers treated with cortisol during

pregnancy had lower plasma cortisol concentrations as they became older compared to

those from control litters, though the difference was only significant in males (daughters:

age x treatment, t = -1.61, P = 0.11; sons: age x treatment t = -2.59, P = 0.01, Table

S6). Data are residual plasma cortisol concentrations from offspring from cortisol-

treated (females: n = 64 samples; males: n = 92) and control (females: n = 89; males: n

= 104) litters. Residuals from a linear mixed-effects model (Table S6) are shown on y-

axis.

Figure 5. Faecal glucocorticoid metabolite (fGCM) concentrations in (A) daughters but

not (B) sons from mothers treated with cortisol during pregnancy were significantly

lower than those from control litters as they became older (daughters: age x treatment, t

= -2.95, P = 0 .0033; sons: age x treatment, t = -0.49, P = 0.62, Table S7. Data are

residual fGCM concentrations from offspring from cortisol-treated (females: n = 79

samples; males: n = 118) and control (females: n = 154; males: n = 201) litters.

Residuals from a linear mixed-effects model (Table S7) are shown on y-axis.

Electronic Supplementary Material

Maternal stress increases cooperative behaviour of offspring in wild meerkats

Ben Dantzer, Constance Dubuc, Ines Braga Goncalves, Dominic L. Cram, Nigel C. Bennett, Andre Ganswindt, Chris Duncan, David Gaynor, Marta B. Manser, Tim H. Clutton-Brock

Contents:
   Tables S1-S7

Table S1. Sample sizes and effects of dominant female treatments on litter characteristics and offspring survival. Means and standard deviations are shown. The number of litters aborted for untreated females was not known. Number of pups emerged correspond to those that emerged from the natal burrow. Three cortisol-treated litters and 1 control litter were aborted prior to birth and are not shown below. Litter sex ratio is the proportion of males in the litter. In some of the untreated litters, sex of the pups was not known (they died before sexing) and so are listed as unknown ("Unk").

Treatment	Total # litters & females treated	Total # pups emerged (F, M, Unk)	Avg. Pups emerged	Avg. Pups Surviving to 3 months	Avg. Pups Surviving to 6 months	Avg. Pups Surviving to 12 months
Untreated	52 (21 females)	49 F, 84 M, 52 Unk	3.78 ± 1.23	2.68 ± 1.58	2.1 ± 1.63	1.62 ± 1.54
Control	7 (6 females)	12 F, 13 M	4.17 ± 0.98	3.83 ± 1.17	3.83 ± 1.17	2.83 ± 1.94
Cortisol	13 (10 females)	13 F, 18 M	3.87 ± 0.83	3.25 ± 1.03	2.75 ± 1.28	1.75 ± 1.75

Table S2. Effects of dominant female treatments on litter characteristics and pup survival. Results are from a linear mixed-effects model (# pups emerged) or generalized linear mixed-effects models (GLMMs, all other response variables) that each contained random intercept terms for dominant female identity and year. No GLMM was overdispersed as indicated by goodness of fit tests (R package aods3, P-values from Pearson χ^2 tests ranged from 0.13 to 1).

Response variable	Fixed effect	b	SE	t or z	P-value
# Litters aborted					
	Intercept	-1.21	0.66	-1.83	0.07
	Birthdate	-0.17	0.59	-0.29	0.77
	Treatment				
	Control	-0.59	1.27	-0.46	0.64
# Pups emerged					
	Intercept	**3.9**	**0.42**	**9.36**	**<0.0001**
	Birthdate	0.02	0.16	0.11	0.91
	Treatment				
	Control	0.3	0.62	0.49	0.63
	Untreated	-0.11	0.45	-0.25	0.8
Litter sex ratio					
	Intercept	-0.54	0.3	-1.84	0.066
	Birthdate	0.02	0.12	0.15	0.88
	Treatment				
	Control	-0.11	0.45	-0.23	0.81
	Untreated	0.09	0.33	0.27	0.78
Prop. litter surviving to 3 months					
	Intercept	-0.16	0.27	-0.62	0.53
	Birthdate	-0.12	0.11	-1.09	0.27
	Treatment				
	Control	0.06	0.39	0.16	0.87
	Untreated	-0.19	0.3	-0.64	0.52
Prop. litter surviving to 6 months					
	Intercept	-0.32	0.28	-1.16	0.24
	Birthdate	-0.19	0.11	-1.65	0.099
	Treatment				
	Control	0.21	0.4	0.52	0.6
	Untreated	-0.28	0.31	-0.91	0.36
Prop. litter surviving to 12 months					
	Intercept	**-0.91**	**0.46**	**-1.999**	**0.046**
	Birthdate	-0.22	0.14	-1.58	0.11
	Treatment				
	Control	0.36	0.46	0.79	0.43
	Untreated	0.096	0.38	0.25	0.8

Reference for Treatment was cortisol-treated litters. Data other than # litters aborted are based upon an initial sample size of offspring from untreated (195 pups from 52 litters produced by 21 dominant females), control (25 pups from 6 litters produced by 6 females), or cortisol-treated (31 pups from 10 litters produced by 9 females) litters that produced pups that emerged from the burrow.

Table S3. Effect of dominant female treatments on offspring body mass from emergence to nutritional independence (1-3 months of age). Data are from a linear mixed-effects model where the response variable was morning body mass that contained random intercept terms for individual identity nested in birth litter nested in mother nested in natal group ($\sigma^2 = 116.7$) and year ($\sigma^2 = 0$).

Fixed Effect	b	SE	t	df	P-value
Intercept					
Females	**184.6**	**9.75**	**18.93**	**53**	**<0.0001**
Males	**189.1**	**9.5**	**19.8**	**49**	**<0.0001**
Litter size	-5.31	3.9	-1.37	48	0.18
First weight	**15.96**	**2.52**	**6.344**	**166**	**<0.0001**
Sex (M)	4.54	4.32	1.05	171	0.29
Age	**53.3**	**1.06**	**50.4**	**4569**	**<0.0001**
Age^2	**-4.76**	**1.15**	**-4.12**	**4516**	**<0.0001**
Rainfall	**2.81**	**0.61**	**4.57**	**4550**	**<0.0001**
Season (Sine)	**-67.2**	**3.3**	**-20.6**	**4238**	**<0.0001**
Season (Co-sine)	**66.22**	**3.4**	**19.2**	**4191**	**<0.0001**
Group size	2.3	2.66	0.86	109	0.39
Group size^2	-0.09	2.21	-0.04	178	0.97
Treatment (Control)					
Females	22.6	15.6	1.45	52	0.15
Males	12.9	15.5	0.83	50	0.41
Treatment (None)					
Females	**23.6**	**10.6**	**2.23**	**54**	**0.029**
Males	**22.9**	**10.4**	**2.22**	**50**	**0.031**
Age x Sex (M)	1.64	1.37	1.19	4567	0.23
Age^2 x Sex (M)	1.02	1.53	0.67	4543	0.5
Sex (M) x Treatment (Control)	-9.73	7.7	-1.27	164	0.21
Sex (M) x Treatment (None)	-0.63	5.1	-0.12	172	0.9
Age x Treatment (Control)					
Females	**7.6**	**1.8**	**4.17**	**4595**	**<0.0001**
Males	-2.3	1.5	-1.48	4548	0.14
Age x Treatment (None)					
Females	0.79	1.2	0.65	4574	0.51
Males	-0.53	1.01	-0.52	4580	0.6
Age^2 x Treatment (Control)					
Females	-0.13	1.93	-0.07	4521	0.94
Males	-1.05	1.6	-0.65	4547	0.52
Age^2 x Treatment (None)					
Females	-2.6	1.3	-1.96	4534	0.051
Males	**-6.9**	**1.1**	**-6.23**	**4556**	**<0.0001**
Age x Treatment (Control) x Sex (M)	**-9.9**	**2.3**	**-4.23**	**4574**	**<0.0001**
Age x Treatment (None) x Sex (M)	-1.3	1.6	-0.84	4581	0.4
Age^2 x Treatment (Control) x Sex (M)	-0.93	2.5	-0.37	4552	0.71
Age^2 x Treatment (None) x Sex (M)	**-4.3**	**1.7**	**-2.5**	**4552**	**0.01**

Data used in these analyses were 4676 measures of body mass from 195 meerkats produced by 21 dominant females across 53 different litters in 16 different social groups in three different years. Only offspring that survived to 90 days of age were included in these analyses.

Table S4. Effect of dominant female treatments on relative babysitting contributions. Data are from a generalized linear mixed-effects model where the response variable is the proportion of babysitting exhibited by the subordinate meerkat relative to the total babysitting contributions the litter received. The model contained random intercept terms for individual ($\sigma^2 = 0.12$), litter nested within group ($\sigma^2 = <0.0001$), and year ($\sigma^2 = 0.000$).

Fixed Effect	b	SE	z	P-value
Intercept				
Females	**-2.1**	**0.22**	**-9.37**	**< 0.0001**
Males	**-2.14**	**0.19**	**-11.01**	**< 0.0001**
Babysitting length	**0.24**	**0.08**	**2.77**	**0.0056**
Observation time	**-0.28**	**0.08**	**-3.4**	**0.0007**
Litter size	0.015	0.05	0.32	0.75
Mixed Litter?	-0.04	0.12	-0.31	0.76
Sex (M)	-0.03	0.27	-0.13	0.9
Age				
Females	0.13	0.17	0.75	0.45
Males	**0.4**	**0.19**	**2.77**	**0.0056**
Foraging success	-0.05	0.05	-0.99	0.32
Mass				
Females	**-0.32**	**0.12**	**-2.61**	**0.009**
Males	-0.098	0.14	-0.7	0.48
Group size				
Females	**-0.35**	**0.09**	**-3.97**	**< 0.0001**
Males	**-0.27**	**0.08**	**-3.28**	**0.001**
Treatment (Control)				
Females	**-0.67**	**0.31**	**-2.14**	**0.032**
Males	-0.12	0.28	-0.43	0.67
Treatment (None)				
Females	-0.06	0.23	-0.25	0.8
Males	-0.16	0.18	-0.85	0.39
Foraging success x Mass	0.036	0.06	0.57	0.57
Age x Mass				
Females	**-0.47**	**0.09**	**-5.08**	**< 0.0001**
Males	**-0.29**	**0.08**	**-3.79**	**0.0001**
Age x Sex (M)	0.28	0.23	1.18	0.24
Mass x Sex (M)	0.22	0.17	1.26	0.21
Group size x Sex (M)	0.07	0.1	0.77	0.44
Treatment (Control) x Sex (M)	0.55	0.42	1.31	0.19
Treatment (None) x Sex (M)	-0.1	0.28	-0.36	0.72
Age x Treatment (Control)				
Females	**-0.62**	**0.21**	**-2.89**	**0.0039**
Males	-0.52	0.27	-1.92	0.055
Age x Treatment (None)				
Females	0.34	0.18	1.88	0.059
Males	-0.006	0.16	-0.03	0.97
Age x Mass x Sex	0.18	0.12	1.5	0.13
Age x Treatment (Control) x Sex (M)	0.09	0.34	0.28	0.78
Age x Treatment (None) x Sex (M)	-0.34	0.24	-1.45	0.14

Data used in these analyses were 182 observations of relative babysitting contributions to 28 litters produced in 9 groups across 3 years recorded from 105 subordinate meerkats.

Table S5. Effect of dominant female treatments on relative pup feeding contributions. Data are from a generalized linear mixed-effects model where the response variable is the proportion of pup feeds exhibited by the subordinate meerkat relative to the total pup feeds the litter received. The model contained random intercept terms for individual ($\sigma^2 = 0.000$), litter nested within group ($\sigma^2 = 0.2$), year ($\sigma^2 = 0.08$), and an observational level random intercept term to control for overdispersion ($\sigma^2 = 0.19$).

Fixed Effect	b	SE	z	P-value
Intercept				
Females	**-2.27**	**0.28**	**-8.2**	**<0.0001**
Males	**-2.66**	**0.25**	**-10.45**	**<0.0001**
Observation time	**0.59**	**0.06**	**9.41**	**<0.0001**
Litter size	-0.055	0.1	-0.55	0.58
Mixed litter (Y)	-0.03	0.29	-0.1	0.92
Sex (M)	-0.39	0.2	-1.93	0.053
Age				
Females	**-0.37**	**0.16**	**-2.33**	**0.019**
Males	**0.29**	**0.14**	**2.07**	**0.039**
Foraging success	**0.14**	**0.06**	**2.12**	**0.033**
Mass				
Females	0.1	0.11	0.91	0.36
Males	**-0.37**	**0.11**	**-3.4**	**0.0007**
Group size				
Females	**-0.48**	**0.13**	**-3.57**	**0.0003**
Males	**-0.32**	**0.13**	**-2.46**	**0.014**
Treatment (Control)				
Females	**-0.73**	**0.23**	**-3.12**	**0.0018**
Males	-0.24	0.21	-1.14	0.25
Treatment (None)				
Females	**-0.64**	**0.18**	**-3.49**	**0.0005**
Males	-0.14	0.13	-1.03	0.3
Foraging success x Mass	0.06	0.06	0.97	0.33
Age x Mass				
Females	-0.04	0.07	-0.62	0.54
Males	**-0.19**	**0.07**	**-2.68**	**0.007**
Age x Sex (M)	**0.66**	**0.19**	**3.49**	**0.00048**
Mass x Sex (M)	**-0.47**	**0.13**	**-3.63**	**0.0003**
Group size x Sex (M)	0.16	0.08	1.89	0.059
Treatment (Control) x Sex (M)	0.49	0.31	1.58	0.11
Treatment (None) x Sex (M)	**0.5**	**0.22**	**2.23**	**0.023**
Age x Treatment (Control)				
Females	-0.11	0.22	-0.5	0.61
Males	0.03	0.29	0.11	0.91
Age x Treatment (None)				
Females	0.17	0.16	1.03	0.3
Males	-0.02	0.14	-0.12	0.9
Age x Mass x Sex	-0.14	0.09	-1.51	0.13
Age x Treatment (Control) x Sex (M)	0.14	0.34	0.42	0.67
Age x Treatment (None) x Sex (M)	-0.19	0.21	-0.91	0.36

Data used in these analyses were 192 observations of relative pup feeding contributions to 26 litters produced in 7 groups across 3 years recorded from 101 subordinate meerkats.

Table S6. Effect of dominant female treatments on plasma cortisol concentrations. Data are from a linear mixed-effects model where the response variable is plasma cortisol concentrations (ln transformed) of the subordinate meerkat. The model contained random intercept terms for individual nested within their birth litter ($\sigma^2 = 0.042$), and capture group ($\sigma^2 = 0.000$).

Fixed Effect	b	SE	t	df	P-value
Intercept					
Females	**3.18**	**0.29**	**11.14**	**120**	**<0.0001**
Males	**3.32**	**0.26**	**12.68**	**111**	**<0.0001**
Sampling time	**1.15**	**0.10**	**11.74**	**260**	**<0.0001**
Sampling time^2	**-0.24**	**0.03**	**-7.27**	**267**	**<0.0001**
Time of day	-0.2	0.14	-1.37	275	0.17
Sample year (2015)	0.46	0.24	1.89	220	0.06
Sex (M)	0.19	0.2	0.91	37	0.37
Age					
Females	-0.1	0.24	-0.41	266	0.68
Males	0.12	0.21	0.57	250	0.57
Body mass	-0.19	0.21	0.89	230	0.37
Foraging success					
Females	-0.1	0.12	-0.84	226	0.4
Males	-0.11	0.11	-0.96	262	0.34
Group size	0.055	0.11	0.49	240	0.62
Group size^2	0.048	0.08	0.58	211	0.56
Pups in group	-0.28	0.18	-1.53	275	0.12
Group sex ratio	**0.19**	**0.08**	**2.35**	**183**	**0.02**
Relatedness	-0.1	0.18	-0.57	41	0.57
Weather (PC1)	-0.1	0.11	-0.94	275	0.35
Treatment (Cortisol)					
Females	-0.03	0.25	-0.11	39	0.91
Males	-0.31	0.19	-1.59	33	0.12
Sex (M) x Age	0.22	0.22	1	272	0.32
Sex (M) x Foraging success	-0.01	0.14	-0.07	257	0.94
Sex (M) x Treatment (Cortisol)	-0.29	0.3	-0.93	34	0.36
Age x Treatment (Cortisol)					
Females	-0.39	0.24	-1.61	275	0.11
Males	**-0.49**	**0.19**	**-2.59**	**274**	**0.01**
Group size x Pups Present (Yes)	-0.05	0.15	-0.37	215	0.71
Group size2 x Pups Present (Yes)	0.07	0.12	0.55	264	0.58
Group size x Weather (PC1)	-0.04	0.08	-0.49	273	0.62
Age x Sex (M) x Treatment (Cortisol)	-0.097	0.31	-0.31	274	0.75

Data used in these analyses were 299 measures of plasma cortisol concentrations from 49 subordinate meerkats produced in 14 litters from 10 different groups. Reference levels (intercept) for "Sex" was female, "Pups in group" was Yes, and for Relatedness was "No parent was dominant".

Table S7. Effect of dominant female treatments on faecal glucocorticoid metabolite (fGCM) concentrations. Data are from a linear mixed-effects model where the response variable fGCM concentrations (ln+1 transformed) of the subordinate meerkat. The model contained random intercept terms for individual nested within their birth litter ($\sigma^2 = 0.10$) and collection group ($\sigma^2 = 0.000$).

Fixed Effect	b	SE	t	df	P-value
Intercept					
Females	**5.49**	**0.27**	**20.1**	**32**	**<0.0001**
Males	**5.4**	**0.25**	**21.6**	**18**	**<0.0001**
Time of day	**-0.14**	**0.04**	**3.27**	**514**	**0.001**
Sample year (2015)	0.02	0.18	0.13	142	0.89
Sex (M)	-0.9	0.21	-0.44	21	0.67
Age					
Females	**0.81**	**0.27**	**3.03**	**167**	**0.0029**
Males	0.33	0.2	1.68	131	0.096
Body mass	-0.3	0.19	-1.5	317	0.13
Foraging success					
Females	0.21	0.13	1.57	394	0.12
Males	0.06	0.06	1.13	518	0.26
Group size	**0.37**	**0.12**	**3.04**	**113**	**0.003**
Group size2	-0.04	0.11	-0.37	462	0.71
Pups in group	-0.14	0.17	-0.84	325	0.4
Group sex ratio	**0.15**	**0.07**	**2.03**	**190**	**0.04**
Relatedness	0.18	0.29	0.6	14	0.55
Weather (PC1)	0.1	0.07	250	1.47	0.14
Treatment (Cortisol)					
Females	-0.15	0.33	-0.46	14	0.65
Males	-0.06	0.28	-0.22	7	0.83
Sex (M) x Age	**-0.48**	**0.18**	**-2.69**	**230**	**0.008**
Sex (M) x Foraging success	-0.14	0.14	-1.02	395	0.3
Sex (M) x Treatment (Cortisol)	0.09	0.34	0.26	18	0.79
Age x Treatment (Cortisol)					
Females	**-0.65**	**0.22**	**-2.95**	**481**	**0.0033**
Males	-0.07	0.15	-0.49	507	0.62
Group size x Pups Present (Yes)	-0.17	0.12	-1.39	272	0.17
Group size2 x Pups Present (Yes)	0.11	0.12	488	0.96	0.34
Group size x Weather (PC1)	0.06	0.06	1	202	0.32
Age x Sex (M) x Treatment (Cortisol)	**0.57**	**0.25**	**2.3**	**456**	**0.022**

Data used in these analyses were 542 faecal samples (n= 355 from controls, n = 187 from cortisol treated litters) from 34 subordinate meerkats (control: n = 12 females, n = 11 male2; cortisol-treated: n = 5 females, n = 6 males) produced in 10 litters from 7 different groups. Reference levels (intercept) for "Sex" was female, "Pups in group" was Yes, and for Relatedness was "No parent was dominant".