Orthogonal regulation of ferritin translation by human eIF3 and the iron response

proteins

Mia C. Pulos[1], Daniel N. Srole[1], David T. McSwiggen[1], Nicholas T. Ingolia[1,4], Jamie H. D.

Cate[1,2,3,4]

[1]Department of Molecular & Cell Biology, University of California, Berkeley, Berkeley, California

94720, USA

[2]Department of Chemistry, University of California, Berkeley, Berkeley, California 94720, USA.

[3]Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, California

94720, USA.

[4] California Institute for Quantitative Biosciences 3 (QB3), University of California, Berkeley,

Berkeley, California 94720, USA.

Corresponding author: Jamie H. D. Cate[1,2,3,4] jcate@lbl.gov

Abstract

Iron is essential to life, but excess iron is detrimental as it catalyzes reactive hydroxyl radical

production. To prevent iron toxicity, intracellular iron homeostasis is regulated by ferritin, a

protein complex composed of a mixture of two subunits: ferritin light chain (FTL) and ferritin

heavy chain (FTH). Ferritin expression is regulated post-transcriptionally by the iron response

proteins (IRPs), which bind an RNA hairpin — the iron responsive element (IRE) — located in

the *FTL* and *FTH* mRNA 5'-untranslated region (5'-UTR). Here, we show *FTL* translation is also

regulated in an IRP-independent manner by eukaryotic translation initiation factor 3 (eIF3). We

find that eIF3 represses *FTL* translation by interacting with a region of the 5'-UTR immediately

downstream of the IRE. Furthermore, we find that loss of eIF3-mediated regulation of *FTL*

translation likely contributes to a subset of clinically-observed hyperferritinemia cases.

Introduction

Iron is essential for a spectrum of metabolic pathways and cellular growth. However, if

not properly managed, excess iron catalyzes the production of harmful radicals that lead to lipid

peroxidation, protein and DNA damage, and ultimately cell death (Palmer, Roberts, and Bero

1994). To safeguard against these damaging effects, cells ubiquitously express ferritin, an iron-

sequestering protein complex. Ferritin is composed of a dynamic mixture of 24 individual

subunits of two types, the ferritin heavy chain (FTH) and the ferritin light chain (FTL), which

assemble into a hollow shell that encapsulates and stores up to 4,500 Fe^{3+} atoms (Harrison and

Arosio 1996) (Anderson and McLaren 2012). Though the two subunits that make up this

complex are structurally similar, they are not functionally interchangeable (Harrison and Arosio

1996). The FTH subunit catalyzes the oxidation of Fe^{2+} to Fe^{3+}, while the FTL subunit lacks

catalytic activity and instead serves as a nucleation site for mineralization, increases complex

stability, and aids in the long-term storage of iron (F. M. Torti and Torti 2002). While ferritin is

ubiquitously expressed, the FTH:FTL ratio changes substantially in different tissues to match

cell-type specific requirements for iron mineralization and metabolism. For example, cardiac

tissue is FTH rich, with ratios of approximately ten FTH to one FTL, whereas hepatic tissue is

generally FTL rich, with ratios of one FTH to seven FTL (Harrison and Arosio 1996).

Iron storage by ferritin not only maintains an iron reserve for the cell, but also protects

against the accumulation of insoluble ferric oxide and toxic iron byproducts. To maintain iron

level homeostasis, FTL and FTH are regulated post-transcriptionally via iron regulatory proteins

(Knovich et al. 2009). Both the *FTL* and *FTH* mRNAs have a highly conserved RNA hairpin in

their 5'-UTR known as the iron responsive element (Theil 1994). Under low iron conditions, the

IRE is bound by IRP1 or IRP2. This interaction generates steric hindrance that blocks 40S

ribosomal subunit recruitment and consequently inhibits translation. This inhibition prevents the

presence of excess ferritin and ensures sufficient levels of accessible iron. In response to an

increase in the labile iron pool (LIP) or under high iron conditions, IRP1 is loaded with an iron

sulfur cluster produced by mitochondria that prevents IRE binding, and IRP2 is targeted for

proteasomal degradation (Fig. 1A,B) (Wilkinson and Pantopoulos 2014). These processes

relieve the transcript from IRP/IRE generated steric hindrances, allowing for translation to

proceed and for the excess iron to be deposited into newly formed ferritin complexes. IRP-IRE

interactions are known to be disrupted in clinically relevant cases of hyperferritinemia, in which

single nucleotide mutations throughout the IRE and *FTL* transcript hinder IRP binding, leading to

a buildup of serum ferritin in the absence of iron overload (Cazzola et al. 1997).

Although the IRP-IRE interactions have been thought to be the sole post-transcriptional

form of regulation of ferritin expression, recent studies have provided strong evidence that

currently uncharacterized factors may impart another layer of regulation during *FTL* translation.

For example, the FTL subunit composition of ferritin is altered in response to environmental

factors such as hypoxia, but this change cannot be fully attributed to changes in the IRP-IRE

interaction or the addition of an iron analogue (Sammarco et al. 2008). Furthermore, preferential

translation of a single ferritin subunit type likely requires the presence of uncharacterized factors

because the IRP-IRE interaction is observed in the 5'-UTRs of both *FTL* and *FTH*. We recently

reported that, using photoactivatable ribonucleoside-enhanced cross-linking and

immunoprecipitation (PAR-CLIP), translation initiation factor eIF3 cross-links to the 5'-UTR of a

select set of cellular mRNA, affecting their translation in a context-dependent manner (Lee,

Kranzusch, and Cate 2015). In general translation, eIF3 acts as a scaffold that stimulates the

organization of other initiation factors and the 40S subunit of the ribosome on mRNAs,

promoting the subsequent scanning to the start codon (Hinnebusch 2006). However, we

showed that eIF3 also can act as either an activator or repressor of translation of specific

transcripts through its interactions with RNA structured elements, located primarily in the 5'-

UTRs of these mRNAs, as exemplified by *JUN* and *BTG1* (Lee, Kranzusch, and Cate 2015).

Notably, we were surprised to find that *FTL*, an mRNA with well-characterized post-translational

regulation mediated by the binding of IRP, also cross-links to eIF3. Whether eIF3 regulates *FTL*

translation through its PAR-CLIP identified interactions with the *FTL* 5'-UTR has not been

established.

Here we report that FTL mRNA utilizes an interaction with eIF3 to provide additional

regulation of its translation. We probed the eIF3-*FTL* mRNA interaction, both *in vitro* and in

cells. We used *Renilla* luciferase reporters to identify translation regulatory features in the 5'-

UTR of *FTL* mRNA that extend beyond the IRE-IRP interaction. Furthermore, we used CRISPR-

Cas9 genome editing to endogenously delete the eIF3 interaction site in *FTL*, allowing us to

monitor direct phenotypic responses in cells. Lastly, we used competitive IRP binding assays to

explore the potential role of eIF3 in hyperferritinemia. These experiments reveal that eIF3 exerts

a novel regulatory role on *FTL* translation, orthogonal to the repressive IRP-IRE interaction, that

likely contributes to a subset of hyperferritinemia cases.

Results

Identification of the eIF3-*FTL* mRNA interaction site

In order to understand the functional effect of the interaction between eIF3 and *FTL*

mRNA, we utilized *Renilla* luciferase (rLuc) reporter mRNAs where the 5' UTR from *FTL* is

placed upstream of the *Renilla* coding sequence (Fig 1D). To measure the importance of the

region identified by PAR-CLIP, various mutations were introduced to disrupt eIF3 binding. The

eIF3 binding site in the 5'-UTR of *FTL*, as determined by PAR-CLIP, spans a 24 nucleotide

sequence within a secondary structural element that overlaps with the last five nucleotides of

the annotated sequence for the *FTL* IRE (Sup. Fig 1A, 1C) (Lee, Kranzusch, and Cate 2015).

Notably, no eIF3 cross-linking site was observed in the 5'-UTR of the mRNA encoding *FTH*

(Sup. Fig 1B) (Lee, Kranzusch, and Cate 2015). Two deletions overlapping the eIF3 interaction

site were constructed, the first designed directly alter the PAR-CLIP data (Δ*PAR*) and the

second developed in order to fully maintain the IRE sequence (Δ*3RE*) (Fig. 1C) (Theil 2015).

The removal of the eIF3 cross-linking site in both cases dramatically increases translation of

rLuc when compared to the full length wild type 5'-UTR, 110 times in the case of the Δ*PAR*

mRNA and 20 times in the case of the Δ*3RE* mRNA (Fig. 1E). These results suggest that eIF3

binding represses *FTL* translation.

Decoupling the repressive role of eIF3 on *FTL* mRNA from that of IRP

Due to the close proximity between the eIF3 interaction site and the *FTL* IRE,

accompanied by the fact that the 5'-UTR of *FTL* is prone to drastic structural rearrangements

(Martin et al. 2011), we tested whether the de-repression observed in the Δ*PAR* and Δ*3RE*

mRNAs is a direct result of altering eIF3 binding and not the IRE-IRP interaction. To evaluate

the effect of the deletions (∆*PAR*, ∆*3RE*) on IRP binding, we carried out RNA-binding assays in

which *FTL* RNA containing a near-IR labeled nucleotide analogue was incubated with

recombinant IRP1 and the interaction was resolved on a nondenaturing gel. As expected, IRP1

bound to the wild type full-length 5'-UTR of *FTL* (Fig. 2A). IRP1 also bound the ∆*3RE* RNA but

failed to efficiently bind the ∆*PAR* RNA (Fig. 2A). The loss of IRP1 binding to ∆*PAR* RNA could

be attributed to disrupted RNA folding, or to a greater importance for the region of overlap

between the IRE and predicted eIF3 binding site than previously thought. In contrast, the

observation that IRP1 still binds the ∆*3RE* RNA suggests the alleviation of repression of

luciferase translation from the ∆*3RE* mRNA (Fig. 1E) is due to disruption of eIF3 binding. To

explore this model in more depth, all further experiments used the ∆*3RE* mutation.

To further ensure the alleviation of repression on the *FTL* 5'-UTR by the ∆*3RE* mutation

is due to disrupting eIF3 binding and not the IRE-IRP interactions, we modulated the location of

the IRE in the 5'-UTR of *FTL*. It had been shown previously that when the IRE in *FTH* was

moved further than 60 nucleotides (nts) from the 5' m^7G-cap, IRP only partially inhibited

translation (Goossen and Hentze 1992). Inhibition is lost because because bound IRP could no

longer provide the steric hindrance needed to prevent the recruitment of the 43S preinitiation

complex on the mRNA (Muckenthaler, Gray, and Hentze 1998) (Paraskeva et al. 1999). To

further investigate if this holds for *FTL* mRNA, we used the *FTL* and ∆*3RE* luciferase reporter

constructs and placed the characteristic C bulge of the IRE to be either 32 nts away (native) or

70 nts away (extended) from the 5' cap (Fig. 2B). Although overall translation was much higher

from the ∆*3RE* mRNAs (Fig 2C), moving the IRE further from the cap alleviated translation

repression of both the wild type *FTL* and the ∆*3RE* mRNAs (Fig. 2C). These results suggest that

IRP-mediated inhibition of translation is maintained in the ∆*3RE* mRNA, and also support the

hypothesis that the eIF3 cross-linking site confers an additional level of repression.

Phenotypic response to loss of eIF3-based repression

To investigate the phenotypic outcomes that result from the deletion of the eIF3

interaction site in the 5' UTR of endogenous *FTL* mRNA, we genetically engineered either

HEK293T or HepG2 cells to generate the Δ*3RE* 5' UTR in the *FTL* gene. This was achieved

using a tandem CRISPR-Cas9 enzyme programmed with single-guide RNAs (sgRNAs)

targeting the *FTL* gene, along with a single stranded DNA (ssDNA) oligonucleotide that was

homologous to the regions spanning the deleted Δ*3RE* sequence (Sup. Fig. 2A). This method

achieved an editing efficiency of nearly 100% in HEK293T cells and roughly 85% in HepG2 cells

(Sup. Fig 2B). We isolated separate clonal cell lines from the HepG2 editing for further analysis.

Through the use of EIF3B immunoprecipitations and qPCR on these edited cell lines, we found

that the Δ*3RE* mutation abolished the preferential interaction of eIF3 with *FTL* mRNA (Sup. Fig

2D,E). FTL protein production increases dramatically in the Δ*3RE* cell lines, as expected from

removing the predicted eIF3 repressive element (Fig 3A). Notably, the increase in FTL protein

levels is not due to increases in mRNA levels (Sup. Fig 3A), indicating that the de-repression

occurs post-transcriptionally.

Remarkably, the increase in FTL levels occurs with a concurrent reduction in FTH levels

(Fig. 3B). This decrease in FTH protein levels is also not due to changes in *FTH* mRNA levels

(Sup. Fig 3B). We therefore investigated whether the lack of eIF3-based repression of *FTL*

translation and concomitant decrease in FTH protein levels had any effects on the ferritin

complex. We purified the ferritin complex from either wild-type HepG2 cells or the Δ*3RE* HepG2

cell line using a fractional methanol precipitation protocol (Cham et al. 1985). We observed that

the Δ*3RE* cell line not only has substantially more ferritin complex, but this complex is also more

stable. Eight-fold more wild-type HepG2 cells were required to produce equivalent amounts of

the ferritin complex isolated from the Δ*3RE* cell line. Furthermore, without ferric ammonium

citrate (FAC) treatment, which has been suggested to stabilize the complex, the ferritin complex

purified from WT cells consistently degraded, unlike the stable complexes from the Δ*3RE* cell

line (Sup. Fig 5). This implicates eIF3 in regulating not only the abundance of the two ferritin

subunits, but also the dynamics and stability of the ferritin complex. Taken together, these

results support the hypothesis that removal of the eIF3 interaction site in the 5'-UTR of the *FTL*

mRNA depresses *FTL* translation and can have a dramatic effect on the ferritin subunit

composition in the cell.

To test whether IRP-mediated repression was maintained in the Δ*3RE* cell lines, we

subjected the cells to either FAC or Desferoxamine (DFO) treatment. FAC serves as an iron

donor, while DFO serves as an iron chelator (Sup Fig 4). Both of these treatments have been

used to modulate the IRP-IRE interaction in human cells (Schneider and Leibold 2003). The

Δ*3RE* cell lines responded to FAC and DFO treatment in a comparable manner to the unedited

(WT) cell lines, providing further evidence that the Δ*3RE* transcript retains IRP-dependent

translational regulation (Fig 3C,D, Sup Fig 4). This persistent iron-responsive regulation occurs

even though the basal levels of FTL protein are higher in the Δ*3RE* cells compared to the WT

cells, further supporting our hypothesis that the observed de-repression in the Δ*3RE* cells is due

to the modulation of eIF3-based regulation of *FTL* translation.

eIF3-mediated repression of FTL and its relation to hyperferritinemia

Mutations that disrupt normal ferritin regulation lead to human genetic diseases such as

hereditary hyperferritinemia cataract syndrome (HHCS), a clinical condition characterized by

constitutive synthesis of FTL and elevated serum ferritin in the absence of abundant iron stores

(Millonig, Muckenthaler, and Mueller 2010). This autosomal dominant condition primarily results

in early onset of cataracts due to ferritin amassing in the lens. HHCS arises from single

nucleotide polymorphisms (SNPs) or deletions in the 5'-UTR and occasionally in the coding

region of *FTL*, which are thought to disrupt the IRP-IRE interaction preventing proper regulation

of the *FTL* transcript. For example, mutations observed in the apical loop of the IRE directly

disrupt main contact sites essential for IRP binding to the IRE (Fig. 4A) (Cazzola et al. 1997).

Interestingly, two SNPs identified in certain patients with hyperferritinemia, G51C and G52C,

disrupt the nucleotides one and two bases upstream of the annotated eIF3 PAR-CLIP site (Fig.

4A) (Camaschella et al. 2000) (Luscieti et al. 2013). Although the PAR-CLIP methodology maps

the region of interaction between an RNA and protein of interest, it does not always capture the

full interaction site due to the requirement for 4-thiouridine cross-linking and subsequent RNase

digestion to generate fragments for deep sequencing (Ascano et al. 2011) (Hafner et al. 2010).

Thus, the proximity of the G51C and G52C mutations to the PAR-CLIP site led us to test

whether the G51C and G52C SNPs could potentially impact the interaction between eIF3 and

FTL mRNA. We generated luciferase reporter mRNAs with mutations in the IRE loop or with

either of the G51C and G52C SNPs and used mRNA transfections to test their effects on

luciferase translation levels. All of these mutations led to an increase in luciferase levels,

indicating that they alleviated translational repression (Fig 4B). We also observed that the G51C

and G52C mRNAs do not interact with eIF3, based on EIF3B immunoprecipitation from

transfected HEK293T cells (Fig. 4C).

Furthermore, the G51C and G52C SNPs in the context of the full length human *FTL* 5'-

UTR have little impact on IRP binding (Fig. 4D). We used RNA binding competition followed by

native gel electrophoretic mobility shift assays to determine the level of 5'-UTR binding to IRP1.

We first ensured the RNA competition experiments reached equilibrium – which required at

least 11 hours of incubation – by measuring the approximate dissociation rate constant (k_{off}) of

WT *FTL* 5'-UTR from IRP1 (see methods). In the competition assay, the IRE-loop mutation

completely abolished the interaction between IRP1 and the 5'-UTR element (Fig. 4D, Sup. Fig.

7). By contrast, the G51C and G52C SNPs maintained near-WT levels of IRP1 binding (Fig. 4D,

Sup. Fig. 7). The IC$_{50}$ values of the G51C and G52C SNP-containing RNAs are at most 2-fold

weaker than the wild-type *FTL* sequence (Fig. 4E). Altogether, these results implicate the loss of

the repressive interaction between eIF3 and the 5'-UTR of *FTL* as causative, at least in part, for

certain cases of hyperferritinemia.

Discussion

Here we have shown that eIF3 represses the translation of *FTL* mRNA, mediated by

binding a region of the *FTL* 5'-UTR immediately adjacent to the IRE. We had previously shown

that eIF3 can serve as an activator or repressor of translation, mediated by binding RNA

secondary structures in the 5'-UTR elements of select mRNAs, using *JUN* and *BTG1* as

prototypical mRNAs (Lee, Kranzusch, and Cate 2015). However, the role of eIF3 in regulating

other mRNAs through the specific interactions we identified by PAR-CLIP remains to be

determined. Using luciferase reporters and cell lines edited at the *FTL* locus, we found that the

FTL interaction with eIF3 can be disrupted without affecting the iron-responsiveness of *FTL*

translation (Fig. 3), or binding of IRP to the remainder of the 5'-UTR containing the IRE (Fig. 2,

Fig. 4). However, the basal level of FTL production increases dramatically in the absence of the

eIF3 binding site (i.e. Fig. 3). Taken together, these results expand the classical model of *FTL*

mRNA post-transcriptional regulation to include the iron-responsive IRP/IRE interaction, and an

orthogonal eIF3-dependent repressive mechanism (Fig. 5). Given the close proximity of the IRE

and eIF3 binding sites in the 5'-UTR of *FTL* mRNA, the interplay of eIF3 and IRP in carrying out

this dual repression of *FTL* translation remains to be determined. It will be interesting in the

future to determine whether eIF3 physically interacts with IRP, or if these two repressors act on

FTL mRNA simultaneously or compete to bind the transcript.

Moreover, we present compelling evidence implicating the disruption of eIF3-mediated

regulation of *FTL* translation as a cause for certain cases of hyperferritinemia. The specific

genetic mutations we analyzed (G51C, G52C) map to the less-conserved lower stem of the IRE,

and are predicted to have no direct interaction with IRP1 (PDB: 3SNP)(Walden et al. 2006).

These mutations minimally interfere with IRP's interaction with the IRE, whereas they greatly

impact the eIF3-based interaction and *FTL* translational repression (Fig 4C-E). While our results

connect eIF3 translational control to specific examples of hyperferritinemia, they also suggest a

broader role for eIF3 in other hyperferritinemias and ferritin-related diseases. For example, we

observed that derepression of *FTL* translation by disruption of the eIF3 interaction site leads to a

concomitant decrease in FTH levels, driving an overall increase in ferritin complex abundance

and stability (Fig. 3, Sup. Fig. 5). Ferritin is a known mediator of the inflammatory response,

raising the question of whether eIF3 may contribute to ferritin's role in inflammation (K.

Morikawa, Oseko, and Morikawa 2009) (Recalcati et al. 2008).

The inherent need for a dual repressive system involving IRPs and eIF3 in normal

human health remains to be determined. In the experiments presented here, eIF3 regulates *FTL*

translation in an iron-independent manner. It is known that different tissue types express

different ratios of FTL to FTH (Harrison and Arosio 1996), and our hypothesis is that eIF3

regulates FTL expression to adjust the composition of ferritin subunits in response to immediate

changes in the environment. For example, during reperfusion in the heart and brain, excessive

oxidative stress accounts for a majority of tissue damage. Maintaining proper iron storage could

be critical to mitigate excess radical production. In the Δ*3RE* cell lines, FTL translation increases

substantially, while FTH levels decrease (Fig. 7B, Fig. 8C,D). We propose that in stress

conditions like reperfusion, eIF3 could work to preserve high levels of FTH to FTL (F. M. Torti

and Torti 2002). This is supported by the fact that the subunit composition of ferritin shifts

towards greater FTH levels in the brain during hypoxic and reperfusion conditions (F. M. Torti

and Torti 2002; Chi et al. 2000). Additionally, it is not known whether or how this eIF3 based

regulation of FTL varies across various cell types. Determining how eIF3 responds to

environmental changes will enable a deeper understanding of how eIF3-dependent repression

of *FTL* translation is triggered, and how eIF3- and IRP-dependent repression mechanisms are

coordinated in different tissue contexts.

Methods

Plasmids

The *FTL* 5'-UTR was amplified from human cDNA, and cloned into the pcDNA4 vector

with a modified Kozak sequence (Kranzusch et al. 2014), by using In-Fusion® HD Cloning Kit

(Takara, Cat.# 638911) to generate the starting luciferase reporter plasmids. Additional

mutations were generated through around-the-horn cloning using either blunt primers for

deleting regions or primers with overhangs to introduce single or double nucleotide mutations.

The IRP1 protein expression plasmid was generated by amplifying the human IRP1 sequence

from human cDNA and inserting it into the 2B-T vector (Addgene, plasmid# 29666) following a

His$_6$ tag and TEV protease cleavage site.

***In vitro* transcription**

RNAs were transcribed using T7 RNA polymerase prepared in-house. For luciferase

reporter mRNAs, 5'-capping and 3'-polyadenylation were performed co-transcriptionally by

including 3′-O-Me-m^{7}G(5′)ppp(5′)G RNA Cap Structure Analogue (NEB, Cat.# S1411L) and

using linearized plasmid template that had a sequence encoding a poly-A tail. Non-labeled

RNAs for the IRP1 electrophoresis mobility shift assays were generated in the same manner,

except the templates were not polyadenylated. Additionally, RNAs with the 38-nt extension were

constructed using a random nucleotide sequence. The exact nucleotide composition prior to the

IRE was previously reported to not significantly impact IRP binding (Goossen and Hentze 1992).

RNAs were purified after DNA digestion by phenol-chloroform extraction followed by ethanol

precipitation.

For genome editing using CRISPR-Cas9-sgRNA RNA-protein complexes (RNPs) (Kim

et al. 2014), sgRNAs were designed using the CRISPR-MIT program. The DNA for transcription

was synthesized by appending the sgRNA sequence downstream of a T7 RNA polymerase

promoter. The DNA was then purified using phenol-chloroform extraction followed by

isopropanol precipitation. After transcription, the RNA products were treated with DNase I

(Promega, Cat.# M6101), run on a 10% denaturing polyacrylamide gel (6M urea), and extracted

from the gel using the crush and soak method, and then subjected to ethanol precipitation.

Luciferase reporter transfections

Human HepG2 cells were maintained in DMEM (Invitrogen 11995-073) with 10% FBS

(Seradigm) and 1% Pen/Strep (Gibco, Cat.# 15140122). Transfections of the luciferase reporter

mRNAs were done using the Mirus *Trans*IT®-mRNA Transfection Kit (Cat.# MIR 2250), with the

following protocol modifications. The day prior to transfection, HepG2 cells were seeded into

opaque 96-well plates so that they would reach 80% confluence at the time of transfection. At

this point, 200 ng of 5'-capped and 3'-polyadenylated mRNA was added to room temperature to

OptiMEM media (Invitrogen, Cat.# 31985-088) to a final volume of 10 µL. 0.6 µL of both Boost

reagent and TransIT mRNA reagent were added to the reaction and left to incubate at room

temperature for 3 min. The TransIT-mRNA Reagent:mRNA Boost:RNA complex was distributed

to the cells in a drop wise manner. Luciferase activity was assayed either 8 hr or 18 hr post-

transfection using the *Renilla* Luciferase assay kit (Promega, Cat.# E2820) and a Microplate

Luminometer (Veritas).

Cell line generation

To generate *FTL*-eIF3 interaction site null cell lines (Δ*3RE*), we used tandem CRISPR-Cas9

enzymes programmed with single-guide RNAs (sgRNAs) targeting the *FTL* gene, along with a

single-stranded DNA (ssDNA) donor with homology to the regions spanning the deleted Δ*3RE*

sequence. The sgRNAs were generated as described above, and targeted regions on both

sides of the eIF3 interaction site (Sup. Fig 2A). The RNP complex was generated by incubating

100 pmol Cas9 with the two sgRNAs at a 1:1.2 Cas9 to total sgRNA ratio. This mixture was

heated to 37 °C for ten minutes and then kept at room temperature until use. The ssDNA donor

was 90 nucleotides long, with 45-nucleotide homology on either side of the predicted double-

strand cut sites allowing it to have perfect homology to the predicted edited sequence.

The Cas9-sgRNA RNP complexes, along with 500 pmol of the ssDNA donor, were

transfected into either $5x10^5$ HEK293T or HepG2 cells using the Lonza 96 well shuttle system

SF kit (Cat. # V4SC-2096). The nucleofection programs used were as follows: CM-130 for

HEK293T and EH-100 for HepG2. The transfected cells were left to incubate for either 48 hrs

for HEK293T cells or 72 hrs for HepG2 cells before harvesting and extracting gDNA using

QuickExtract (Epicentre: QE09060). The editing efficiency of individual Cas9-sgRNA RNPs was

determined using a T7 endonuclease 1 assay (Reyon et al. 2012). The efficiency of the Dual-

sgRNA editing approach was determined by PCR-amplifying a 180-base pair region around the

eIF3 interaction site, and analyzing the resulting products on a 6% non-denaturing 29:1

polyacrylamide gel. Monoclonal populations of edited cells were sorted using FACS, screened,

and the final edited sequence was determined using TOPO TA cloning (Ramlee et al. 2015).

Transcript level abundance

Total transcript abundance was determined by lysing 1.25 x 10^6 cells with Qiazol lysis

buffer followed by using the Directzol RNA extraction kit (Zymo Research, Cat. # R2061),

according to the manufacturer's instructions. The cDNA was generated by reverse transcription

using 350 ng of RNA, random hexamers, and Superscript IV (Thermo Fisher scientific, Cat. #

18091050). Primers for the qPCR were as follows: FTL forward: 5'-

ACCTCTCTGGGCTTCTAT-3', FTL reverse: 5'-AGCTGGCTTCTTGATGTCCT-3'(Cozzi et al.

2013), ACTB forward: 5'-CTCTTCCAGCCTTCCTTCCT-3', ACTB reverse: 5'-

AGCACTGTGTTGGCGTACAG-3', PSMB6 forward: 5'-GGACTCCAGAACAACCACTG-3',

PSMB6 reverse: 5'-CAGCTGAGCCTGAGCGACA-3'(Mokany et al. 2013) , FTH forward: 5'-

CGCCAGAACTACCACCA-3', FTH reverse: 5'-TTCAAAGCCACATCATCG(Liu et al. 2012),18S

forward: 5'-GGCCCTGTAATTGGAATGAGTC-3', 18S reverse: 5'-

CCAAGATCCAACTACGAGCTT-3', (Lee et al. 2016) RLUC forward: 5'-

GGAATTATAATGCTTATCTACGTGC-3', RLUC reverse: 5'-

CTTGCGAAAAATGAAGACCTTTTAC-3'(Kong et al. 2008). Run conditions were: 95 ˚C for 15

sec, followed by 40 cycles of 95 ˚C for 15 sec, 60 ˚C for 60 sec, 95 ˚C for 1 sec.

RNA immunoprecipitation and qPCR

RNA immunoprecipitations were adapted from (Lee, Kranzusch, and Cate 2015) with the

following deviations. One 15 cm plate of either HEK293 or HepG2 cells was used to prepare cell

lysate using a NP40 lysis buffer (50 mM HEPES- KOH pH 7.5, 150 mM KCl, 2 mM EDTA, 0.5%

Nonidet P-40 alternative, 0.5 mM DTT, 1 Complete EDTA-free Proteinase Inhibitor Cocktail

tablet per 50 ml of buffer). The lysate was precleared with 15 µL of Dynabeads preloaded with

rabbit IgG (Cell Signaling, Cat.# 2729) for one hour at 4 °C. The lysate was collected and then

incubated with a fresh 15 µL aliquot of Dynabeads and 7.5 µL of anti-EIF3B antibody (Bethyl

A301-761A) for two hours at 4 °C. Preparation of cDNA and qPCR primers are described and

listed above.

For EIF3B immunoprecipitations of transfected mRNAs, 2.15 µg of mRNA was

transfected into 1 well of either HEK293T or HepG2 cells in a 12-well plate using the protocol

above. Cells were then left to incubate for 8 hours before harvesting. The cells were lysed using

the same buffer as described in (Lee, 2015), and precleared with 2 µL of rabbit IgG-coated

Dynabeads for one hour at 4 °C. The lysate was collected and then incubated with a fresh 2 µL

aliquot of Dynabeads and 4 µL of anti-EIF3B antibody (Bethyl A301-761A) for two hours at 4 °C.

RNA was collected, cDNA prepared and qPCR carried out with the following primers and run

conditions as described above.

Iron level modulation and ROS treatment

In order to modulate the iron levels in cells. HepG2 and HEK293T cells were treated with

a final concentration of either 50 µg/mL of ferric ammonium citrate (FAC) (an iron donor) or 200

µM of Desferoxamin (DFO) (an iron chelator) for 24 hours before harvesting.

IRP1 purification

IRP1 was purified based on the protocols in (Carvalho and Meneghini 2008) (Basilion et

al. 1994) with modifications. The IRP1-encoding 2B-T plasmid was transformed into chemically-

competent BL21 Rosetta pLysS *E. coli*, using heat shock at 42 °C, and grown on Ampicillin

plates. A single colony was used to inoculate a 5 mL LB culture containing Ampicillin, which was

then used to inoculate a 50 mL starter culture that was allowed to reach saturation overnight.

Approximately 4x10 mL of the overnight culture was used to inoculate 4x1L cultures using

ZY5052 media lacking the 1000x trace metal mix (30mM HEPES, 5% glycerol, 43 mM KCl,

0.5mM EDTA, 0.5 mM DTT) (Studier 2005)+ Carbomicillin. The 1L cultures were grown at 37 °C

to OD$_{600}$ = 0.36, at which point the temperature was lowered to 18 °C, and allowed to grow at 18

°C for 36 hours prior to harvest.

Pelleted *E. coli* cells were lysed using sonication in the lysis buffer (30 mM HEPES

pH=7.5, 400 mM KCl, 5% Glycerol and 1 mM DTT) along with Protease inhibitor (Roche, Cat. #

5056489001) tablets. Lysate was loaded on a 5 mL Ni-NTA pre-packed HiTrap column (GE,

Cat. # 17-5248-02), allowed to incubate at 4 °C for 1 hour, before eluting using 600 mM

imidazole in the same buffer as above. Pooled fractions from the elution were then dialyzed

overnight into ion-exchange (IEX) buffer (30mM HEPES pH7.5, 1 mM DTT, 5% Glycerol, and 1

mM EDTA), for subsequent purification using a 5 mL HiTrap Q-column (GE, Cat. # 17-1154-01).

Sample was loaded on the Q column using IEX buffer, and the column was washed with eight

column volumes of IEX buffer without KCl. IRP1 was eluded using 800 mM KCl in IEX buffer.

IRP1 electrophoresis mobility shift assays

To detect IRP1 binding by native gel shifts, RNA samples were transcribed using the

Atto-680 RNA Labeling Kit (Jena Bioscience, FP-220-680) and subsequently purified using RNA

Clean & Concentrator-25 columns (Zymo, R1018). This form of labeling has been shown not to

disrupt protein-RNA interactions (Kohn et al. 2010). Unlabeled RNA was transcribed and

purified as described above.

Binding experiments were carried out with a final concentration of 300 pM of labeled

RNA and 225 nM of recombinant human IRP1, which facilitated a 1:1 ratio of RNA binding to

IRP1. Heparin was included at the beginning of the reaction at a final concentration of 4.5

mg/mL. The initial binding reaction was carried out in a 1x RXN buffer (40 mM KCl, 20 mM Tris-

HCL, pH7.4, 2 mM MgCl$_2$, 2 mM DTT, and 5% Glycerol) for 30 minutes at room temperature

(Goforth et al. 2009; Fillebeen, Wilkinson, and Pantopoulos 2014). For competition experiments,

unlabeled RNA was then added in concentrations 1000x-100,000x that of the labeled RNA. The

reactions were then incubated for an additional 18 hours at room temperature, to allow the

reactions to reach equilibrium, based on the off-rate of WT *FTL* 5'-UTR determined in separate

experiments. In preliminary experiments, we found the k_{off} to be roughly 0.006 min^{-1} using an 8-

hour incubation time course with competitor. We then tested the fraction of IRP bound after 11

hr and 18 hr incubations with competitor *FTL* RNA and observed no changes in the residual

fraction of IRP bound to RNA (~15%). Thus, subsequent experiments with competitor RNAs

were carried out for 18 hr, after the first 30 minute pre-incubation in the absence of competitor.

The reactions were resolved on Tris-glycine gels (Thermo Fisher Scientific, Cat.#

XP04122BOX) and gels was visualized using a Odyssey Licor set to 700 nM wavelength and an

intensity of 6.5. Band intensity quantification was carried out using Image Studio (Licor). The

IC$_{50}$ values for each competitor RNA was determined using Graph Pad Prism 7 (Graph Pad

Software) from a set of triplicate experiments.

Ferritin complex purification

The ferritin complex purification procedure was adapted from (Ridgeway, 1985), with

slight modifications. Either one 15-cm dish of *Δ3RE* cells grown in normal media or eight 15-cm

dishes of wild-type HepG2 cells that had been treated with 50 ng/mL FAC for 24 hours were

harvested, weighed, and lysed in 4x weight/volume NP40 lysis buffer (50 mM HEPES-KOH

pH=7.5, 150 mM KCl, 2 mM EDTA, 0.5% Nonidet P-40 alternative, 0.5 mM DTT, 1 Complete

EDTA-free Protease Inhibitor Cocktail tablet per 10 mL of buffer). Samples were incubated on

ice for ten minutes and then centrifuged for 10 minutes at 15000 rpm at 4 °C. Samples were

diluted in 1:2 phosphate buffered saline (PBS), and methanol was added to the diluted lysate to

a final concentration of 40% (v/v). The sample was then heated to 75 °C for 10 min. After

cooling on ice for 5 min, the samples were centrifuged in a microfuge 20R at 3650 RPM for 15

min at 4 °C. The resulting supernatant was collected and concentrated using a 100k MW cutoff

Amicon filter (Cat.# UFC510024) by centrifugation for 10 min at 14000xg. The sample was

washed once with PBS and spun for an additional 4 min at 14000xg. The sample was then

collected by inverting the column and centrifuging the sample for 2 min at 1000xg at 4°C. All

samples collected were brought to a final volume of 80 µL with PBS. The purity of the sample

was determined by running the sample on a native gel (Thermo Fisher Scientific, Cat.#

XP04200BOX) followed by Coomassie staining.

Western blots

The following antibodies were used for Western blot analysis: anti-EIF3B (Bethyl A301-

761A) at 1:1000; anti-FTL (Abcam, ab69090) at 1:800; anti-FTH (Santa Cruz, sc-25617) at

1:400; anti-IRP1 (Abbexa, abx004618) at 1:400; anti-IRP2 (Abcam, ab129069) at 1:800; and

anti-b-Actin (Abcam, ab227) at 1:1000.

Authors Contributions

M. C. P. designed the main conceptual ideas and carried out the majority of the experiments.

M. C. P., D. T. M., J. H. D. C. conceived the project. D. N. S. assisted in carrying out

experiments related to screening and phenotypically characterizing clonal cell lines, as well as

construct generation. J. H. D. C. and N. T. I. supervised and aided in the planning of the overall

direction for the project. M. C. P. and J. H. D. C. drafted the manuscript with support from N. T.

I., D. N. S., and D. T. M.

Acknowledgments:

We are immensely grateful to Bruno Martinez for providing us with purified IRP1, Amy Lee for

helpful discussion and mentorship, Luisa Arake De Tacca for help optimizing the eIF3b

immunoprecipitations, Hector Nolla for help with flow cytometry and single cell sorting, and

Alison Killilea and the Berkeley Tissue culture facility for cells and advice.

Funding Sources

453   P50 GM102706/GM/NIGMS NIH HHS/United States

454   R01 GM065050/GM/NIGMS NIH HHS/United States

455   AHA predoctoral fellowship 16PRE30140013/United States

References

Anderson, Gregory J, and Gordon D McLaren, eds. 2012. *Iron Physiology and Pathophysiology*

in Humans. New York, NY: Humana Press. doi:10.1007/978-1-60327-485-2.

Ascano, Manuel, Markus Hafner, Pavol Cekan, Stefanie Gerstberger, and Thomas Tuschl.

2011. "Identification of RNA-Protein Interaction Networks Using PAR-CLIP." *Wiley*

Interdisciplinary Reviews: RNA 3 (2): 159–77. doi:10.1002/wrna.1103.

Basilion, J P, T a Rouault, C M Massinople, R D Klausner, and W H Burgess. 1994. "The Iron-

Responsive Element-Binding Protein: Localization of the RNA-Binding Site to the Aconitase

Active-Site Cleft" 91 (2): 574–78.

Camaschella, Zecchina, Lockitch, Roetto, Campanella, Arosio, and Levi. 2000. "A New Mutation

(G51C) in the Iron-Responsive Element (IRE) of L-Ferritin Associated with

Hyperferritinaemia-Cataract Syndrome Decreases the Binding Affinity of the Mutated IRE

for Iron-Regulatory Proteins." *British Journal of Haematology* 108 (3). Wiley/Blackwell

(10.1111): 480–82. doi:10.1046/j.1365-2141.2000.01920.x.

Carvalho, H, and R Meneghini. 2008. "Increased Expression and Purification of Soluble Iron-

Regulatory Protein 1 From Escherichia Coli Co-Expressing Chaperonins GroES and

GroEL." *Brazilian Journal of Medical and Biological Research* 41 (4). Brazilian Journal of

Medical and Biological Research: 270–76. doi:10.1590/S0100-879X2008005000009.

Cazzola, M, G Bergamaschi, L Tonon, E Arbustini, M Grasso, E Vercesi, G Barosi, P E Bianchi,

G Cairo, and P Arosio. 1997. "Hereditary Hyperferritinemia-Cataract Syndrome:

Relationship Between Phenotypes and Specific Mutations in the Iron-Responsive Element

of Ferritin Light-Chain mRNA." *Blood* 90 (2): 814–21.

Cham, B E, H P Roeser, A Nikles, and K Ridgway. 1985. "A Procedure for the Purification of

Ferritin From Human Liver by Heating a Methanol-Treated Homogenate." *Analytical*

Biochemistry 151 (2): 561–65.

Chi, S I, C K Wang, J J Chen, L Y Chau, and T N Lin. 2000. "Differential Regulation of H- and L-

Ferritin Messenger RNA Subunits, Ferritin Protein and Iron Following Focal Cerebral

Ischemia-Reperfusion." *Neuroscience* 100 (3): 475–84. doi:10.1016/S0306-4522(00)00317-

1.

Cozzi, Anna, Paolo Santambrogio, Daniela Privitera, Vania Broccoli, Luisa Ida Rotundo,

Barbara Garavaglia, Rudolf Benz, et al. 2013. "Human L-Ferritin Deficiency Is

Characterized by Idiopathic Generalized Seizures and Atypical Restless Leg Syndrome.."

The Journal of Experimental Medicine 210 (9). Rockefeller University Press: 1779–91.

doi:10.1084/jem.20130315.

Fillebeen, Carine, Nicole Wilkinson, and Kostas Pantopoulos. 2014. "Electrophoretic Mobility

Shift Assay (EMSA) for the Study of RNA-Protein Interactions: the IRE/IRP Example."

Journal of Visualized Experiments, no. 94: 1–9. doi:10.3791/52230.

Goforth, J B, S A Anderson, C P Nizzi, and R S Eisenstein. 2009. "Multiple Determinants Within

Iron-Responsive Elements Dictate Iron Regulatory Protein Binding and Regulatory

Hierarchy." *Rna* 16 (1): 154–69. doi:10.1261/rna.1857210.

Goossen, Brifta, and Matithias W Hentze. 1992. "Position Is the Critical Determinant for

Function of Iron-Responsive Elements as Translational Regulators." *Molecular and Cellular*

Biology 12 (5): 1959–66.

Hafner, Markus, Markus Landthaler, Lukas Burger, Mohsen Khorshid, Jean Hausser, Philipp

Berninger, Andrea Rothballer, et al. 2010. "PAR-CliP - a Method to Identify Transcriptome-

Wide the Binding Sites of RNA Binding Proteins." *Journal of Visualized Experiments*, no. 41:

1–5. doi:10.3791/2034.

Harrison, Pauline, and Paolo Arosio. 1996. *Harrison PM, Arosio PThe Ferritins: Molecular*

Properties, Iron Storage Function and Cellular Regulation. Biochim Biophys Acta 1275:161-

203. Vol. 1275.

Hinnebusch, Alan G. 2006. "eIF3: a Versatile Scaffold for Translation Initiation Complexes."

Trends in Biochemical Sciences 31 (10): 553–62. doi:10.1016/j.tibs.2006.08.005.

Kim, S, D Kim, S W Cho, J Kim, and J S Kim. 2014. "Highly Efficient RNA-Guided Genome

Editing in Human Cells via Delivery of Purified Cas9 Ribonucleoproteins." *Genome*

Research 24 (6): 1012–19. doi:10.1101/gr.171322.113.

Knovich, Mary Ann, Jonathan a Storey, Lan G Coffman, Suzy V Torti, and Frank M Torti. 2009.

"Ferritin for the Clinician." *Blood Reviews* 23 (3): 95–104. doi:10.1016/j.blre.2008.08.001.

Kohn, M, M Lederer, K Wachter, and S Huttelmaier. 2010. "Near-Infrared (NIR) Dye-Labeled

RNAs Identify Binding of ZBP1 to the Noncoding Y3-RNA." *Rna* 16 (7): 1420–28.

doi:10.1261/rna.2152710.

Kong, Yi Wen, Ian G Cannell, Cornelia H de Moor, Kirsti Hill, Paul G Garside, Tiffany L

Hamilton, Hedda A Meijer, et al. 2008. "The Mechanism of Micro-RNA-Mediated Translation

Repression Is Determined by the Promoter of the Target Gene." *Pnas* 105 (26). National

Academy of Sciences: 8866–71. doi:10.1073/pnas.0800650105.

Kranzusch, Philip J, Amy S Y Lee, Stephen C Wilson, Mikhail S Solovykh, Russell E Vance,

James M Berger, and Jennifer A Doudna. 2014. "Structure-Guided Reprogramming of

Human cGAS Dinucleotide Linkage Specificity." *Cell* 158 (5). Elsevier Inc.: 1011–21.

doi:10.1016/j.cell.2014.07.028.

Lee, Amy S Y, Philip J Kranzusch, and Jamie H D Cate. 2015. "eIF3 Targets Cell-Proliferation

Messenger RNAs for Translational Activation or Repression." *Nature* 522 (7554): 111–14.

doi:10.1038/nature14267.

Lee, Amy S Y, Philip J Kranzusch, Jennifer A Doudna, and Jamie H D Cate. 2016. "eIF3d Is an

mRNA Cap-Binding Protein That Is Required for Specialized Translation Initiation." *Nature*

Publishing Group 536 (7614). Nature Publishing Group: 96–99. doi:10.1038/nature18954.

Liu, Huan-Dong, Wei Li, Zhen-Rui Chen, Meng-Liang Zhou, Zong Zhuang, Ding-Ding Zhang,

Lin Zhu, and Chun-Hua Hang. 2012. "Increased Expression of Ferritin in Cerebral Cortex

After Human Traumatic Brain Injury." *Neurological Sciences* 34 (7): 1173–80.

doi:10.1007/s10072-012-1214-7.

Lizio, Marina. 2015. "Gateways to the FANTOM5 Promoter Level Mammalian Expression Atlas,"

January, 1–14. doi:10.1186/s13059-014-0560-6.

Luscieti, Sara, Gabriele Tolle, Jessica Aranda, Carmen Benet Campos, Frank Risse, Érica

Morán, Martina U Muckenthaler, and Mayka Sánchez. 2013. "Novel Mutations in the

Ferritin-L Iron-Responsive Element That Only Mildly Impair IRP Binding Cause Hereditary

Hyperferritinaemia Cataract Syndrome.." *Orphanet Journal of Rare Diseases* 8: 30.

doi:10.1186/1750-1172-8-30.

Martin, J S, M Halvorsen, L Davis-Neulander, J Ritz, C Gopinath, A Beauregard, and A

Laederach. 2011. "Structural Effects of Linkage Disequilibrium on the Transcriptome." *Rna*

18 (1): 77–87. doi:10.1261/rna.029900.111.

Millonig, Gunda, Martina U Muckenthaler, and Sebastian Mueller. 2010. "Hyperferritinaemia–

Cataract Syndrome: Worldwide Mutations and Phenotype of an Increasingly Diagnosed

Genetic Disorder." *Human Genomics* 4 (4): 250–62.

Mokany, E, Y L Tan, S M Bone, C J Fuery, and A V Todd. 2013. "MNAzyme qPCR with

Superior Multiplexing Capacity." *Clinical Chemistry* 59 (2): 419–26.

doi:10.1373/clinchem.2012.192930.

Morikawa, Keiko, Fumimaro Oseko, and Shigeru Morikawa. 2009. "A Role for Ferritin in

Hematopoiesis and the Immune System." *Leukemia & Lymphoma* 18 (5-6): 429–33.

doi:10.3109/10428199509059641.

Muckenthaler, Martina, Nicola K Gray, and Matthias W Hentze. 1998. "IRP-1 Binding to Ferritin

mRNA Prevents the Recruitment of the Small Ribosomal Subunit by the Cap-Binding

Complex eIF4F." *Molecular Cell* 2 (September): 383–88.

Palmer, C, R L Roberts, and C Bero. 1994. "Deferoxamine Posttreatment Reduces Ischemic

Brain Injury in Neonatal Rats.." *Stroke; a Journal of Cerebral Circulation* 25 (5): 1039–45.

Paraskeva, Efrosyni, Nicola K Gray, Britta Schl Ger, Kristina Wehr, and Matthias W Hentze.

1999. "Ribosomal Pausing and Scanning Arrest as Mechanisms of Translational Regulation

From Cap-Distal Iron-Responsive Elements" 19 (1): 807–16.

Ramlee, Muhammad Khairul, Tingdong Yan, Alice M S Cheung, Charles T H Chuah, and Shang

Li. 2015. "High-Throughput Genotyping of CRISPR/Cas9-Mediated Mutants Using

Fluorescent PCR-Capillary Gel Electrophoresis," October. Nature Publishing Group, 1–13.

doi:10.1038/srep15587.

Recalcati, Stefania, Pietro Invernizzi, Paolo Arosio, and Gaetano Cairo. 2008. "New Functions

for an Iron Storage Protein: the Role of Ferritin in Immunity and Autoimmunity." *Journal of*

Autoimmunity 30 (1-2): 84–89. doi:10.1016/j.jaut.2007.11.003.

Reyon, Deepak, Shengdar Q Tsai, Cyd Khayter, Jennifer A Foden, Jeffry D Sander, and J Keith

Joung. 2012. "FLASH Assembly of TALENs for High-Throughput Genome Editing." *Nature*

Biotechnology 30 (5): 460–65. doi:10.1038/nbt.2170.

Sammarco, Mimi C, Scott Ditch, Ayan Banerjee, and Ed Grabczyk. 2008. "Ferritin L and H

Subunits Are Differentially Regulated on a Post-Transcriptional Level." *Journal of Biological*

Chemistry 283 (8): 4578–87. doi:10.1074/jbc.M703456200.

Schneider, Brian D, and Elizabeth a Leibold. 2003. "Effects of Iron Regulatory Protein

Regulation on Iron Homeostasis During Hypoxia." *Regulation* 102 (9): 3404–11.

doi:10.1182/blood-2003-02-0433.Supported.

Studier, F William. 2005. "Protein Production by Auto-Induction in High-Density Shaking

Cultures." *Protein Expression and Purification* 41 (1): 207–34.

doi:10.1016/j.pep.2005.01.016.

Theil, E C. 1994. "Iron Regulatory Elements (IREs): a Family of mRNA Non-Coding

Sequences.." *Biochemical Journal* 304 (Pt 1): 1–11.

Theil, Elizabeth C. 2015. "IRE mRNA Riboregulators Use Metabolic Iron (Fe2+) to Control

mRNA Activity and Iron Chemistry in Animals." *Metallomics* 7 (January). Royal Society of

Chemistry: 15–24. doi:10.1039/C4MT00136B.

Torti, Frank M, and Suzy V Torti. 2002. "Regulation of Ferritin Genes and Protein." *Blood* 99

(10): 3505–16. doi:10.1182/blood.V99.10.3505.

Walden, William E, Anna I Selezneva, Jerome Dupuy, Anne Volbeda, Juan C Fontecilla-Camps,

and Elizabeth C Theil. 2006. "Structure of Dual Function Iron Regulatory Protein 1

Complexed with Ferritin IRE-RNA ." *Science* 314 (5807): 1903–8.

doi:10.1126/science.1133116.

Wilkinson, Nicole, and Kostas Pantopoulos. 2014. "The IRP/IRE System in Vivo: Insights From

Mouse Models." *Frontiers in Pharmacology* 5 JUL (July): 1–15.

doi:10.3389/fphar.2014.00176.

597
598
Figure 1: Post-transcriptional regulation of *FTL* mRNA. A, B) Iron-responsive regulation

mediated by binding of Iron Response Proteins (IRPs) to Iron Response Element (IRE) RNA

structures in the 5'-UTR in (A) low iron conditions and (B) high iron conditions. **C)** Mapping of

the IRP and eIF3 interaction sites on the experimentally-determined secondary structure of *FTL*

mRNA (Martin et al. 2011). The transcription start site is derived from the FANTOM5 database

(Lizio 2015). **D)** General schematic of the luciferase reporter mRNAs. The eIF3 PAR-CLIP site

in *FTL* mRNA spans nucleotides 53-76 and the 3RE region spans nucleotides 58-90. **E)**

Luciferase activity in HepG2 cells transfected with luciferase reporter mRNAs. The results are

from biological duplicates carried out in technical triplicate, normalized to luciferase

luminescence from luciferase reporter mRNA with wild-type *FTL* 5'-UTR.

609
Figure 2: Maintenance of 5'-cap and IRP-dependent regulation of 3RE deletions in the

FTL 5'-UTR. A) Representative native RNA gel shift showing IRP1 binding activity. Near IR

(NIR) labeled RNAs corresponding to the full-length WT *FTL* 5'-UTR and the 5'-UTR with either

of two deletions of the predicted eIF3 interaction site in *FTL* 5'-UTRs were incubated with

recombinant IRP1. The RNA complexes were resolved on a native tris- glycine gel and imaged

on an odyssey Licor. **B)** Schematics showing modulation of the distance of the IRE from the 5'-

cap of the RNA (Goossen and Hentze 1992; Muckenthaler, Gray, and Hentze 1998) and the

effects the change has on IRP regulation of translation initiation. **C)** The corresponding

luciferase activity of HepG2 cells transfected with mRNAs containing the native and extended

spacing between the 5'-cap and IRE, with or without the 3RE site, in HepG2 cells. The results

are from biological duplicates, each carried out in technical triplicate, and are normalized to the

luciferase luminescence of cells transfected with WT *FTL* mRNA with native spacing.

622
623

Figure 3: Physiological effects of the endogenous removal of the 3RE repressive

element. A,B) Representative Western blots of (A) FTL and (B) FTH levels in the edited (*Δ3RE*)

and WT HepG2 cells under normal iron conditions. Serial dilutions were used in order to better

visualize the significance of the changes in FTL and FTH protein abundance. **C, D)**

Representative Western blots of FTL levels in the edited (*Δ3RE*) and WT HepG2 cells under

high or low iron conditions. (C) Iron donor treatment with FAC at 50 µg/mL for 24 hr, and (D)

Iron chelation treatment with DFO at 200 µM for 24hr. The asterisk (*) indicates that lysate from

Δ3RE cells were diluted two-fold, due to the higher overall levels of FTL in these cells.

634
Figure 4: Role of eIF3 in select cases of Hyperferritinemia. A) Diagram of IRP binding to the

IRE in *FTL* mRNA (Anderson and McLaren 2012). Hyperferritinemia mutations are highlighted in

orange (Cazzola et al. 1997), green (Camaschella et al. 2000), and blue (Luscieti et al. 2013)

with their corresponding serum ferritin levels listed. The 3RE area is indicated in purple.

Nucleotides that directly interact with the IRP are also identified (i.e. A15, G16, U17). **B)**

Luciferase activity of HepG2 cells transfected with mRNAs encoding the WT *FTL* 5'-UTR or

various Hyperferritinemia mutations. The results are from biological duplicates; each carried out

in technical triplicate, and normalized to WT *FTL* reporter luciferase luminescence. **C)** Binding of

eIF3 to luciferase reporter mRNAs with WT or mutant forms of the *FTL* 5'-UTR, using EIF3B

Immunoprecipitation (IP), followed by RNA extraction RT-qPCR. Cells were harvested 8 hrs

post-transfection. Data shown as the percent in the IP compared input levels. **D)** Dose-response

curve of three RNA competition assays, based on gel shifts of NIR-labeled WT IRE-containing

RNA, with WT, G51C, G52C, or Loop mutant (A15G/G16C) RNAs serving as cold competitors.

Fold excess of competitors extended up to 100,000x. **E)** The calculated IC_{50} values of the

various competitor RNAs, based on the data in (**D**), with the error bars representing the

standard deviation. N.A., the IC_{50} value could not be determined for the Loop mutant due to lack

of any detectable competition.

Figure 5: Model of post-transcriptional regulation of *FTL* mRNA. IRPs repress FTL

translation in an iron-dependent manner, whereas eIF3 represses FTL translation in an iron-

independent manner. Coordination between IRP repression eIF3 repression is not known.

657
658 **Sup. Fig. 1: Sites of eIF3 interaction with *FTL* and *FTH* mRNAs. A, B)** eIF3 PAR-CLIP

659 cluster identified for (**A**) *FTL* and (**B**) *FTH* (Lee 2015). **C)** Mapping of the IRP and PAR-CLIP

660 derived eIF3 interaction sites on the secondary structure of the 5'-UTR in *FTL* mRNA,

661 determined by chemical probing (Martin et al. 2011). The transcription start site is derived from

662 the FANTOM5 database (Lizio 2015).

 Sup. Fig. 2: CRISPR-Cas9 editing to remove the proposed eIF3-*FTL* interaction site. A)

CRISPR-Cas9 RNP editing schematic in which two guide RNAs target sequences immediately

adjacent to the eIF3 repressive element (3RE). Reactions were supplemented with 90-nt ssDNA

that had full homology to the intended edited sequence. **B)** Editing efficiency in both HEK293T

and HepG2 cells, based on PCR of the region of interest and analysis on a native

polyacrylamide gel. **C)** Three successfully edited HepG2 clonal populations, as determined by

sequencing the *FTL* locus. **D, E)** Determination of the preferential binding of eIF3 to *FTL* mRNA

via EIF3B immunoprecipitation (IP) followed by RNA extraction and RT-qPCR in both (D)

HEK293T and (E) HepG2 cell lines. Control RNAs used to normalize IPs were *PSMB6* and

ACTB.

Sup. Fig. 3: mRNA levels in edited cell lines. A, B) mRNA levels of either (**A**) *FTL* or (**B**)

FTH as determined by RNA extraction from HepG2 (solid black bars) or HepG2 *Δ3RE* (black

outlined bars) cells and RT-qPCR. The mRNA levels for *FTL* and *FTH* were normalized to

ACTB mRNA. Cells were grown in normal media, 50 μg/mL FAC treatment for 24 hr, or 200 μM

DFO treatment for 24 hr.

A FAC treatment

B DFO treatment

680 **Sup. Fig. 4: Physiological effects of the endogenous removal of the 3RE on FTH levels. A,**

681 **B)** Representative Western blots of FTH levels in the edited (*Δ3RE*) and WT HepG2 cells under

682 various iron conditions. (**A**) Iron donor treatment with FAC at 50 µg/mL for 24 hr; (**B**) Iron

683 chelation treatment with DFO at 200 µM for 24hr.

Sup. Fig. 5: Analysis of the ferritin complex upon deletion of the 3RE in *FTL* mRNA.

Representative native 4-12% gradient Tris-Glycine gel resolving ferritin complexes purified

using methanol, from either 8x 15-cm dishes of WT HepG2 cells treated with 50 µg/mL FAC for

24 hrs or a single 15-cm dish of *Δ3RE* HepG2 cells under normal media conditions.

Sup. Fig. 6: Native gels resolving RNA-IRP1 complexes formed after competition

experiments. Near IR (NIR) labeled RNA corresponding to the full-length WT *FTL* 5'-UTR were

incubated with recombinant IRP1 and increasing molar excess concentrations of unlabeled

competitor RNAs (1,000x, 2,000x, 4,000x, 8,000x, 12,000x, 16,000x, 20,000x, 50,000x,

75,000x), as indicated. F, lane with only NIR-labeled *FTL* RNA; N, no addition of competitor

RNA to the NIR-RNA-IRP1 complex.