*Title of article:*

Flow modification associated with reduced genetic health of a river-breeding frog, *Rana*

*boylii*

*Authors' names:*

Ryan A. Peek[1†], Sean M. O'Rourke[2], Michael R. Miller[2]

[1] Center for Watershed Sciences, University of California, Davis, CA 95616, USA

[2] Department of Animal Science, University of California, Davis, CA 95616, USA

[†] E-mail: rapeek@ucdavis.edu

**Abstract**

River regulation or flow modification—the hydrological alteration of flow by dams and

diversions—has been implicated as a cause of fundamental change to downstream

aquatic ecosystems. Flow modification changes the patterns and functionality of the

natural flow regime, and has the potential to restrict population connectivity and gene

flow in river-dependent organisms. Since population connectivity and the maintenance

of genetic diversity are fundamental drivers of long-term persistence, understanding the

extent flow modification impacts these critical attributes of genetic health is an

important goal for long-term conservation. Foothill yellow-legged frogs (*Rana boylii*)

were historically abundant throughout many western rivers but have declined since the

onset of regulation. However, the extent to which *R. boylii* populations in rivers with

altered flow regimes are maintaining connectivity and genetic diversity is unknown.

Here we use genetic methods to investigate the impacts of flow alteration on *R. boylii* to

explore their potential for long-term persistence under continued flow modification. We

found *R. boylii* in rivers with flow modification showed striking patterns of isolation and

trajectories of genetic diversity loss relative to unregulated rivers. For example, flow

modification explained the greatest amount of variance in population genetic

differentiation compared with other covariates including geographic distance.

Importantly, patterns of connectivity and genetic diversity loss were observed

regardless of flow alteration level but were most prominent in locations with the

greatest flow modification intensity. Although our results do not bode well for long-term

persistence of *R. boylii* populations under current flow regulation regimes, they do

highlight the power of genetic monitoring for assessing population health in aquatic

organisms.

**Introduction**

River regulation, or the hydrological alteration of flow by dams and diversions, impacts the seasonal and interannual flow variability within a watershed. This flow modification changes the natural flow regime and dramatically alters geomorphic and hydrologic connectivity of watersheds (Poff et al. 2007), which may restrict population connectivity (Schick and Lindley 2007, Shaw et al. 2016) and gene flow (Deiner et al. 2007, Cazé et al. 2016, Thompson et al. 2019). River regulation can change flow frequency, magnitude, duration, timing, and rate of change, which can have significant impacts on aquatic organisms and ecological processes (Poff et al. 2007, Yarnell et al. 2010). Flow modification associated with hydropower generation has been implicated as a cause of fundamental changes to downstream aquatic ecosystems (Power et al. 1996, Bunn and Arthington 2002, Moyle et al. 2011). The hydrological regimes of over half of the world's largest rivers have been altered by large dams (Nilsson et al. 2005) and only recently has the extent of flow alteration and the associated ecosystem-level impacts been acknowledged (Pringle 2001, Dudgeon et al. 2006, Murchie et al. 2008).

Rivers simultaneously connect and carve the landscapes through which they flow. Rivers provide corridors of connectivity for riparian and aquatic organisms such as fish, amphibians, and macroinvertebrates (Wiens 2002, Pringle 2003), while also acting as physical barriers on the landscape for many terrestrial organisms (Voelker et al. 2013, Cazé et al. 2016). In Mediterranean climates, rivers have strong seasonal patterns associated with cold, wet winters and warm, dry summers. Native aquatic organisms have evolved life histories well adapted to these natural flow patterns, which are both

predictable and seasonal (Yarnell et al. 2010, Tonkin et al. 2017). Thus, changes to

temporal and spatial abiotic river processes caused by river regulation can have a

substantial impact on biotic communities. The negative effects of flow alteration on

migration and loss of spawning habitat (Lind et al. 1996, Fuller et al. 2011, Kupferberg et

al. 2012, Rolls and Bond 2017), reductions in population abundances and diversity (Lind

et al. 1996, Zhong and Power 1996, Vörösmarty et al. 2010, Fuller et al. 2011, Scribner et

al. 2016, Sabo et al. 2017, Guzy et al. 2018), and fragmentation (Zhong and Power 1996,

Vörösmarty et al. 2010, Werth et al. 2014, Scribner et al. 2016, Sabo et al. 2017, Guzy et

al. 2018) have been well documented. However, most rivers have not been regulated for

long periods (e.g., typically less than 100 years) compared to the time aquatic

organisms had to adapt to pre-anthropogenic flow conditions. In flow-altered rivers that

organisms still occupy, it remains unknown whether populations can persist long-term

with continued flow modification. In other words, while some species may have

persisted since flow regulation began in a system (e.g., typically four to five decades),

this does not necessarily mean these populations will persist into the future under

current altered flow regimes. Thus, exploring the potential for long-term persistence of

populations under different altered flow regimes is a crucial component for guiding

conservation efforts, yet it remains a significant gap.

One tool that can help address this gap is the integration of genetics and

hydrology to better assess the impact of flow alteration on aquatic organisms (Scribner

et al. 2016). Although aquatic organisms are often difficult to count and monitor by

conventional methods, genetic monitoring can be a powerful tool to assess population

health by revealing factors such as genetic fragmentation and population declines. It is

widely recognized that reductions in population connectivity can increase inbreeding,

leading to a potential "extinction vortex" (Gilpin and Soule 1986), yet there is limited

understanding of how flow alteration may impair the processes crucial for maintenance

of genetic variation and thus adaptive capacity. In addition, there is a pressing need for

more effective and flexible watershed management tools, particularly in relation to

monitoring aquatic populations, and assessment of environmental flows (Grantham et

al. 2010, Yarnell et al. 2020). Thus, population genetics could be a powerful tool to

understand the influence of different flow regimes on population health and this

information could facilitate improved flow management to better protect aquatic

populations.

The river-breeding foothill yellow-legged frog (*Rana boylii*) historically occurred in

lower and mid-elevation streams and rivers from Southern Oregon to northern Baja

California west of the Sierra-Cascade crest (Stebbins 2003). *Rana boylii* are intimately

linked with river hydrology because they have evolved to spawn in synchrony with

natural flow cues associated with seasonal spring snowmelt or rain recession periods

(Kupferberg 1996, Yarnell et al. 2010, 2016, Bondi et al. 2013). However, population

declines have been documented across the former range of this species, particularly in

southern California and the Sierra Nevada where it has been extirpated from

approximately 50 percent of its historical range (Jennings and Hayes 1994, Davidson et

al. 2002). In California, particularly in the Sierra Nevada, flow alteration may be a

significant environmental stressor (Lind et al. 1996, Kupferberg et al. 2012). Regulated

river reaches typically alter flows by augmenting or diverting winter and spring runoff,

thereby disrupting natural flow regimes and reducing or eliminating flow cues.

Aseasonal flow fluctuation from flow alteration can scour (i.e., detach from substrate)

or desiccate *R. boylii* egg masses, and the loss of clutches may have a significant

demographic impact because only one egg mass is laid per year. In many regulated

rivers in the Sierra Nevada, *R. boylii* populations are now restricted to small unregulated

tributaries flowing into the regulated mainstem (Peek 2018).

Here, we investigate the impacts of flow modification on genetic health of *R. boylii*

populations across three different flow regimes. Given that population connectivity and

genetic diversity are known to be play critical roles in long-term species persistence, our

primary questions were:

1. How has flow alteration impacted seasonality and predictability of flow patterns

in regulated river reaches?

2. Do populations of *R. boylii* in flow altered river reaches have less genetic diversity

and decreased measures of genetic connectivity than populations in unregulated

river reaches?

3. Are metrics of connectivity and diversity in *R. boylii* associated with different

levels of flow alteration?

Our goal was to assess the genetic health of *R. boylii* under different flow regimes to

better inform the potential for long-term persistence with flow alteration. Addressing

this issue will help to inform management and conservation efforts for *R. boylii*, through

evaluation and prioritization of conservation actions for populations with decreased

genetic health. In addition, our approach demonstrates the potential utility of genetics

for future conservation monitoring and assessment efforts in aquatic species in altered

systems.

**Material And Methods**

To begin investigating the impact of river regulation on *R. boylii*, we collected frog

tissue and buccal samples from 30 locations in six rivers representing three different

flow alteration levels associated with hydropower generation (Table 1). The three flow

regimes assessed were: 1) hydropeaking, where flows are pulsed on most days from

late spring through fall to provide electricity during peak-use hours and for recreational

whitewater rafting; 2) bypass, which diverts river flows from an upstream portion of the

basin to the downstream power generation facilities; and 3) unregulated, a largely

natural flow regime where no upstream controls exist to regulate flows (Figure 1A).

Flow data were obtained for each river reach using proximal USGS gaging stations

(Data S1). We analyzed *R. boylii* samples from sites in three major watersheds (Yuba,

Bear, and American) in the northern Sierra Nevada of California (Figure 1A; Table 1, Data

S2). The six study rivers share a similar Mediterranean climate, underlying geology,

watershed aspect (west-slope), stream morphology (riffle-pool), and vegetative

communities, but differ in the intensity of flow alteration (Steel et al. 2017). Although

flow alteration occurs in all three of the study watersheds, both the North Yuba and

North Fork (NF) American are unregulated whereas the Middle Fork (MF) American is

the only river that has a hydropeaking flow regime (Figure 1B, Figure 1C).

***Assessment of hydrologic impacts of river regulation on patterns of seasonality***

To evaluate how flow alteration has impacted seasonal flow patterns in

regulated river reaches, we used a wavelet analysis to derive metrics of predictability

and Colwell's information-theoretic metrics to define seasonality following methods

outlined in Tonkin et al. (2017). Seasonality—how much the environment varies during

the course of a single year—was measured using Colwell's M/P metric, which is the

contingency (M) divided by within season predictability (P) (Colwell 1974). Predictability

was measured using the standardized Morlet wavelet power at the 12-month time

interval across the entire data series (Torrence and Compo 1998). Daily average flow

data from all available gages within the study area were used to compare the inter-

annual (wavelet) and intra-annual (Colwell's indices) flow patterns in each of the study

reaches (Data S1, Data S3). In regulated reaches where data was available, metrics

were estimated from the full period of record prior to initiation of flow alteration (pre-

regulation) as well as from current altered flow regimes.

***Sample collection and DNA extraction***

To evaluate genetic diversity and connectivity in *R. boylii* populations, a total of

345 tadpole tail clips, buccal swabs, and tissue samples were compiled. The samples

were collected between 1992 and 2017, from watersheds across the west slope of the

Sierra Nevada (Data S2). Field sampling was conducted following Heyer et al. (1994),

under CDFW SCP Permit #0006881, with IACUC protocol #19327. Individual post-

metamorphic frogs were buccal-swabbed following established protocols (Goldberg et

al. 2003, Pidancier et al. 2003, Broquet et al. 2007). Each post-metamorphic individual

was comprehensively swabbed underneath tongue and inside of both cheeks for

approximately 30 sec to one minute. Swabs were air dried for approximately five

minutes and placed in 1.5 mL microcentrifuge tubes while in the field or placed in lysis

buffer (Goldberg et al., 2003). Dried samples were stored in the laboratory at -80°C until

DNA extraction. Where possible, tail clips from tadpole larvae were collected, and

tadpoles greater than 15 mm total length were targeted (Parris et al., 2010; Wilbur &

Semlitsch, 1990). One clip was taken per individual tadpole and dried on Whatman filter

paper (grade 1) or placed in 95% ethanol and stored at room temperature. DNA was

extracted from samples in ethanol and lysis buffer using Qiagen DNeasy Blood & Tissue

kits following manufacturer protocol and stored at -20°C. DNA was extracted from dried

buccal swabs and tail clips using an Ampure magnetic bead-based protocol (Ali et al.,

2016) and stored at -20°C.

***RAD Sequencing and RAD-Capture (Rapture) bait design***

To produce a genomic resource for frog species with large genome sizes, we

interrogated a significant fraction of the *R. boylii* genome using RAD sequencing with

SbfI (Miller et al. 2007, Baird et al. 2008, Ali et al. 2016). Paired-end Illumina sequence

data were generated using 24 *R. boylii* individuals (Data S4) and de novo locus

discovery and contig extension were carried out as previously described (Miller et al.

2012, Sağlam et al. 2016, Peek et al. 2019) using the alignment program Novoalign and

the PRICE assembler (Ruby et al. 2013). This resulted in a set of 77,544 RAD contigs

ranging from 300 to 800 bp which served as a de novo partial reference assembly for all

subsequent downstream analyses (Data S5). We next removed loci with five or more

SNPs to reduce potential paralogs and chimeras, and randomly selected 10,000 loci

from the remaining subset. Of these 10,000 loci, 8,533 were successfully designed into

120 bp RAD capture baits by Arbor Biosciences (Data S6).

***Rapture Sequencing and Genomic Analysis***

A total of 345 individual frog samples were prepared for sequencing following

the RAD Capture (Rapture) methods outlined in Ali et al. (2016) and detailed in Peek et

al. (2019). We generated RAD libraries from the samples, quantified the libraries using a

Fragment Analyzer (Agilent Technologies, Santa Clara, CA) and pooled them, performed

capture on the pooled library with the 120 bp baits described above, and sequenced the

resulting Rapture library with paired-end Illumina sequencing.

The mean number of filtered alignments across all 345 samples was 324,928,

and sequence coverage depth was measured at the mid-point (i.e., base-pair 60) of

each 120 bp Rapture bait locus using the depth function in SAMtools (Li et al. 2009, Li

2011). For downstream analysis, we selected individuals that had greater than 100,000

alignments (n=277), which provided sufficient data to investigate population genetic

attributes at broad and fine geographic scales (Table 2, Data S2). *Rana boylii* are cryptic,

and often occur in low densities within the study area. Thus, we retained a minimum of

three individuals per site, and the mean number of samples per site was approximately

nine (Table 2). With genomic data, population genetic parameters can be accurately

estimated from even low sample numbers (Hotaling et al. 2018), and genomic analyses

in non-model organism often use fewer loci than analyzed here (Narum et al. 2013).

Thus, the sequence data we obtained should be appropriate for population genetic

analyses across our study area.

The probabilistic framework implemented in Analysis of Next Generation

Sequencing Data (ANGSD) (Korneliussen et al. 2014) was used for all population

genetic analyses as it does not require calling genotypes and is suitable for low-

coverage sequencing data (Korneliussen et al. 2013, Fumagalli et al. 2013). ANGSD

analyses were conducted following details from Peek et al. (2019).

**Principal component analysis to assess population structure and connectivity**

To assess *R. boylii* population structure across the collection locations, we used

ANGSD (Korneliussen et al. 2014) to perform principal component analysis (PCA).

Settings used in ANGSD for PCA to identify polymorphic sites for PCA included a

SNP_pval of $1\times10^{-6}$, inferring major and minor alleles (doMajorMinor 1), estimating allele

frequencies (doMaf 2) (Kim et al. 2011), retaining SNPs with a minor allele frequency of

at least 0.05 (minMaf), genotype posterior probabilities were calculated with a uniform

prior (doPost 2), and the doIBS 1 and doCov 1 options were used to generate PCA data.

Principal components (PC) summarizing population structure were derived from classic

eigenvalue decomposition and were visualized using the {ggplot2} package (Wickham

2016) in R (R Core Team 2017).

**Genetic differentiation and diversity estimates**

To investigate patterns of genetic diversity and the overall trajectory (stable,

increasing, or decreasing) of diversity in a population, we summarized patterns of

genetic variation using two estimators of θ (4Nμ): Tajima's θ ($\theta_\pi$) is based on the

average number of pairwise differences (Tajima 1983), and Watterson's θ ($\theta_S$) is based

on the number of segregating sites (Watterson 1975). Estimates of θ were made using

the empirical Bayes method in ANGSD (Korneliussen et al. 2014) with the site frequency

spectrum (SFS) to calculate each statistic for each site, which were then averaged to

obtain a single value for each statistic (Korneliussen et al. 2013). These estimators are

influenced by the demographic history of a population and provide information on the

trajectory of changes in genetic diversity (Tajima 1989, Wakeley 2009). When genetic

diversity has been stable, these estimates should be equal; when genetic diversity has

been decreasing, $\theta_\pi > \theta_S$; and when genetic diversity has been increasing, $\theta_\pi < \theta_S$. Scaled

$\Delta\theta$ ($[\theta_\pi - \theta_S]$ / ($[\theta_\pi + \theta_S]$ / 2 )) was used to assess trajectories of genetic diversity in each

population.

To assess how patterns of genetic differentiation and connectivity are

associated with flow alteration in the study, we calculated pairwise $F_{ST}$ (Wright 1943)

between all collection locations within a river for all six rivers. Genome-wide $F_{ST}$

between location pairs was estimated by first calculating a site frequency spectrum

(SFS) for each population (doSaf) (Nielsen et al. 2012) with ANGSD. The two-

dimensional SFS and global $F_{ST}$ between each population pair were then estimated

using realSFS (Korneliussen et al. 2014). We then plotted the scaled pairwise $F_{ST}$ ($F_{ST}$ /

[1-$F_{ST}$]) (Rousset 1997) for each location against the river distance (the distance along

the river network from each collection location to every other location within that study

river). We calculated the river distances (distance along river network) between

locations within watersheds using the {riverdist} package in R (Tyers 2017). These

values were plotted and a generalized linear model was fitted (Scaled $F_{ST}$ ~ River

Distance) in R (R Core Team 2017). Furthermore, each location was assigned flow

alteration type based on either the regulation type at the mainstem sample location or

at the nearest mainstem; for pairwise comparisons the type was assigned based on

whichever was greatest (i.e., hydropeaking is of greater intensity than bypass).

***Boosted regression tree modeling of variance in $F_{ST}$***

We used boosted regression tree (BRT) models with the R packages {gbm}

(Ridgeway 2015) and {dismo} (Hijmans et al. 2017) to assess the relative influence of

flow alteration type as compared to other environmental covariates. Boosted regression

trees (BRT) are suitable frameworks for large and complex ecological datasets because

they do not assume normality, nor linear relationships between predictor and response

variables and they ignore non-informative predictor variables (Graham et al. 2008, Steel

et al. 2017). BRTs use iterative boosting algorithms to combine simple decision trees to

improve model performance (De'ath 2007) and provide a robust alternative to many

traditional statistical methods (Phillips et al. 2006, Guisan et al. 2007). BRTs assess the

relative impact of modeled variables by calculating the number of times a variable is

selected for splitting a tree across all folds of the cross validation. Following Steel *et al*.

2017, estimates of relative influence for each predictor variable were used to evaluate

the relative contribution a variable had in predicting the response. To evaluate the

relative influence of covariates on $F_{ST}$, models were trained using river distance (km),

elevation (m), upstream drainage area ($km^2$), Strahler stream order, and number of

samples per location. Stream segment data on elevation, length, slope, stream order,

and drainage area were derived from NHD Plus attributes (U.S. Geological Survey,

National Hydrography Dataset, Digital data, accessed, August 2017 at

http://nhd.usgs.gov/data.html). In addition, scaled Δθ was included to assess the effect

of genetic diversity trajectory on $F_{ST}$ across regulation types.

Model training and fitting were conducted following methods previously

described in Steel et al. (2017). To reduce overfitting, the learning rate (also known as

the shrinking rate) was set to 0.001. Stochastic gradient boosting was utilized to reduce

prediction error (De'ath 2007) and the fraction of training data sampled to build each

tree was 0.75, within the range as recommended by (Brown et al. 2012). Tree

complexity was set to three to allow for second and third order interaction effects. The

minimum number of observations required in the final nodes of each tree was three. A

ten-fold cross-validation technique allowed us to determine the number of trees at

which prediction error was minimized using the cross-validation deviance. Model

performance was evaluated using the minimum estimated cross-validation deviance

which maximized the estimated deviance explained.

## Results

### *Flow alteration reduces the strength and predictability of seasonality patterns*

Flow patterns between flow-altered and unregulated river reaches could be

clearly differentiated by flow regime (hydropeaking, bypass, unregulated). Wavelet

analysis and Colwell's M/P metric indicated flow alteration reduced seasonality and

predictability of flow patterns in regulated (hydropeaking and bypass) river reaches

(Figure 1B, 1C, 2). These patterns were distinctly different before and after flow

alteration began in regulated study reaches (Figure 2, Data S3). In all flow-altered

reaches, seasonality was reduced compared to pre-regulation flow patterns, and

predictability of these flow patterns was also reduced in flow-altered data (Figure 2).

### *Anomalous genetic pattern in highly regulated reach of Middle Fork American watershed*

To assess *R. boylii* population structure across the collection locations, we used

principal component analysis to provide a dimensionless comparison of all samples.

The first two principal components revealed four main groups corresponding to the

Yuba, Bear, North Fork (NF) American, and Middle Fork (MF) American samples (Figure

3A). Unlike the Yuba watershed where all rivers clustered as one group, the two rivers

within the American watershed (the NF American and MF American) were separated by

both PC1 and PC2. Although the NF American watershed clustered closely with the

adjacent Bear watershed, the MF American showed a surprisingly high degree of

genetic differentiation from other locations (Figure 3A). These data suggest that there

is less genetic differentiation between the NF American and the Bear watersheds, than

between the NF and MF American watersheds. We conclude that measurements of

overall genetic differentiation in *R. boylii* from our study area largely conform to

watershed and geographic expectations, with the exception of the American watershed,

which shows a high degree of genetic differentiation between the North (unregulated)

and Middle (hydropeaking) Forks.

To further investigate patterns of genetic variation within the American

Watershed, we performed two PCAs, one on samples from the NF American, and the

other on samples from the MF American. The PCA of the NF American showed minimal

differentiation among locations, with different study sites blending together and weak

patterns of population structure (Figure 3B). In contrast, PCA of the MF American

showed strong differentiation between sites (Figures 3C, 3D). The MF American PCA

completely resolved all sites, with the first component (PC1) strongly differentiating the

samples in the hydropeaking reach from all other sites in the MF American. This pattern

may be due to the differential river regulation between the two rivers; the NF American

is unregulated and has weak PCA differentiation, whereas the MF American has a

higher level of river regulation and all sites form distinct genetic clusters, indicative of

reduced gene flow among sites within the MF American.

***Flow regime is the strongest predictor of genetic isolation with R. boylii in the Northern***

***Sierra***

To explore the relationship between the flow alteration type and genetic

differentiation across the entire study area, we plotted scaled $F_{ST}$ against river distance

between sampling sites for each flow alteration type assessed. While there was a clear

relationship between scaled $F_{ST}$ and river distance (as shown by the slope of regression

lines in Figure 4A), there was a striking pattern of elevated $F_{ST}$ by flow alteration type

(Figure 4A, Data S7). Even the bypass flow alteration type showed a distinct pattern of

elevated $F_{ST}$. For instance, sites in hydropeaking flow regimes separated by less than

10km had scaled $F_{ST}$ values comparable to unregulated locations separated by river

distances over 50 km. Hydropeaking was the most extreme pattern of the three flow

alteration types and showed highly elevated $F_{ST}$ values compared with the unregulated

and bypass flow types. The centroid $F_{ST}$ of each flow type increased with increasing

flow alteration intensity from unregulated (centroid $F_{ST}$=0.144, se=0.007) to bypass

(centroid $F_{ST}$=0.218, se=0.012) to hydropeaking (centroid $F_{ST}$=0.284, se=0.017). The

centroid $F_{ST}$ for hydropeaking was nearly double compared with unregulated centroid

$F_{ST}$. Furthermore, the centroid of the river distance between site pairs was comparable

between flow alteration types (unregulated=20.9 km, bypass=21.4 km,

hydropeaking=21.6 km), further supporting patterns of genetic differentiation are

associated with flow alteration and not solely isolation by distance. This suggests a

greater degree of isolation within sites in reaches with altered flow regimes compared

with *R. boylii* populations in unregulated reaches, as larger $F_{ST}$ values represent

reductions in heterozygosity due to population subdivision (Slatkin 1987). We conclude

*R. boylii* in rivers with altered flow regimes show patterns of greater population isolation

and elevated $F_{ST}$, (i.e., reduced heterozygosity) compared to populations in unregulated

locations.

To investigate the relative influence of flow alteration compared to other

covariates such as river distance on genetic differentiation (i.e. $F_{ST}$), we used boosted

regression tree (BRT) modeling. Covariates included flow regime alteration type, river

distance, watershed variables derived from National hydrology data (NHD), topographic

data, and genetic diversity trajectory (i.e., scaled $\Delta\theta$). We found flow alteration

explained the greatest amount of variance in scaled $F_{ST}$ (Figure 4B). Thus, flow

alteration has a larger relative influence on genetic differentiation than river distance

between sampling locations in altered flow reaches. We conclude there is a pattern of

elevated $F_{ST}$ between populations in reaches with altered flow.

***Flow alteration is strongly correlated with decreasing genetic diversity in R. boylii***

To investigate the association between flow and genetic diversity trajectory(

whether genetic diversity is stable, increasing, or decreasing), we summarized patterns

of genetic variation using two estimators of $\theta$ (4N$\mu$): Tajima's $\theta$ ($\theta_\pi$) and Watterson's $\theta$

($\theta_S$). We found zero populations sampled within flow altered watersheds had evidence

of increasing genetic diversity (i.e., a $\theta_\pi - \theta_S$ and scaled $\Delta\theta$ ([$\theta_\pi - \theta_S$] / mean $\theta$)) that was

less than zero) (Figure 5A). The altered flow locations showed a clear trajectory of

genetic diversity loss (Figure 5A, 5B). Three of the four hydropeaking locations had the

highest values of scaled $\Delta\theta$, and the global mean was significantly different from other

flow alteration types. Although some tributary populations within unregulated

watersheds also showed signs of genetic diversity loss, the mean genetic diversity

trajectory at unregulated locations was largely stable (Figure 5B). This indicates many

populations in the northern Sierra Nevada which are already limited in number, are

losing genetic variation, and flow alteration is exacerbating these patterns. We conclude

there is evidence of recent genetic diversity loss across populations in the flow altered

river systems, regardless of alteration type.

**Discussion**

Although massive parallel sequencing (MPS) technologies have the potential to

facilitate collection of high-quality genetic data in virtually any species, a number of

challenges still remain for many species including low quality or non-existent reference

genomes, large/complex/repetitive genomes, and high cost of processing/sequencing

in studies with many samples. Amphibians are particularly challenging as many species

have very large genome sizes (Nunziata et al. 2017) for example, there are less than 10

frog reference genome assemblies available as of 2020 (Hellsten et al. 2010, Sun et al.

2015, Session et al. 2016, Hammond et al. 2017, Edwards et al. 2018, Rogers et al. 2018,

Seidl et al. 2019) despite over 7,000 recorded frog species (AmphibiaWeb 2020). Our

results demonstrate that Rapture (Ali et al. 2016) is a suitable method to rapidly and

inexpensively discover a large number of loci in a frog species with a complex genome.

In this study, we used new RAD sequencing and RAD capture (Rapture) methods (Ali et

al. 2016) to generate high-quality genomic data suitable for discovering and genotyping

many single nucleotide polymorphisms (SNPs) in *R. boylii*. Based on this dataset, we

were able to successfully characterize patterns of genetic variation within *R. boylii* as

well as design a set of RAD capture baits that can be used as a genetic monitoring

resource for *R. boylii* (and likely other ranid species). This highlights that the collection

of genetic information, even from large numbers of samples or in complex genomes, is

no longer a limitation with current genomic methods such as RAD and Rapture.

Demographic connectivity is widely recognized as a fundamental driver of long-

term population persistence (Fahrig and Merriam 1985, Taylor et al. 1993). Populations

must adapt over time and connectivity is a major way to transfer genetic information.

For example, previous studies have shown that adaptation can occur by spreading

specific alleles across large geographic distances (Miller et al. 2012, Prince et al. 2017).

In many regulated river reaches in the Sierra Nevada, *R. boylii* now occur in isolated

locations, breeding in tributaries rather than mainstem habitats. However, since these

frogs have the potential to move long distances (*R. boylii* have been observed moving

over 1 km per day (Bourque 2008)), the extent to which current population connectivity

has been lost due to river regulation was not understood. Examining pairwise $F_{ST}$,

revealed a major decrease in connectivity in populations in regulated systems, even with

low intensity flow alteration (i.e., bypass reaches). The primary factor influencing

genetic differentiation among these rivers is hydrologic alteration (Figure 3B). Thus,

despite being able to move long distances, *R. boylii* have not been able to maintain

population connectivity in rivers with altered flow regimes. This demonstrates that even

in species that can move relatively long distances and pass potential physical barriers

(e.g., infrastructure such as dams, canals, and reservoirs likely do not represent barriers

to movement of *R. boylii*) loss of connectivity may still occur and can be revealed with

genetic analysis.

Genetic diversity is also a critical component for long-term population

persistence because it is closely related to the evolutionary capacity for adaptation to

environmental changes (Lande and Shannon 1996, Frankham 2002, Hoffmann and Sgrò

2011, Ishiyama et al. 2015). In some cases, isolated populations can maintain genetic

diversity when they are sufficiently sized (Whiteley et al. 2010), however, species of

conservation concern typically have small and/or declining populations and thus may

be susceptible to genetic diversity loss (Frankham 2002, Krohn et al. 2018). Throughout

the Sierra Nevada, *R. boylii* have largely disappeared from regulated mainstem rivers,

but the extent to which existing populations have been able to maintain genetic diversity

is unclear. Strikingly, our analysis revealed that every single population within the

regulated watersheds exhibits a trajectory of genetic diversity loss. Thus, genomic

analysis of molecular variation provides a powerful lens to discover and assess

trajectories of genetic diversity.

Our analyses, using metrics that serve as a reasonable proxy for genetic health,

does not bode well for the long-term persistence of *R. boylii* populations in regulated

rivers in the Sierra Nevada. Many of these *R. boylii* populations are already losing

genetic diversity and the effects of inbreeding will likely exacerbate their problems given

their small size and reduced connectivity. *Rana boylii* have evolved in river systems with

consistent hydrologic seasonality and predictability, despite inter-annual variation. Flow

regulation has altered patterns of hydrologic seasonality and predictability in many

watersheds (Kupferberg et al. 2012). Long-term population persistence may still be

possible if conservation efforts utilize methods that promote or maintain genetic health

and increase population connectivity. For example, simulations by Botero et al. (2015)

demonstrated adaptation persisted in modeled populations through large

environmental changes—if phenotypic strategies were appropriate before and after the

change—but modeled populations declined rapidly when the current strategy was a

mismatch to the current environment. Thus, *R. boylii* conservation efforts should focus

on river reaches where flow management may provide opportunities to more closely

mimic local natural flow regimes and thus improve hydrologic connectivity.

Detecting evolutionary responses to within- and among-year changes in an

ecological or hydrological context has previously been difficult. However, utilizing

genetic data can fill these gaps and provide a highly informative process for identifying

the impacts of anthropogenic and environmental change on the process of adaptation

(Kahilainen et al. 2014, Botero et al. 2015). We demonstrate that an aquatic species that

has adapted to local hydrology patterns shows poor genetic health (i.e., clear patterns

of decreased connectivity and trajectories of genetic diversity loss). Our results

highlight the impact of river regulation on aquatic organisms and their potential for long

term persistence. In the future, similar genetic approaches could be used in many other

contexts to explore the impacts of river regulation on aquatic organisms. Taken

together, our results demonstrate that genetic monitoring can be a powerful tool for

assessment of population health and should be a critical component of conservation

management in aquatic organisms.

**Acknowledgements**

This research builds on work from Amy Lind, Sarah Kupferberg, Jennifer Dever,

and Sarah Yarnell. Many thanks to all of them for insight, and to many who helped

collect/prepare samples: Corey Luna, Kristen Hein Strohm, Rick Wachs, and Isaac

Chellman. This research could not have been conducted without access to specimens

from Brad Shaffer at the Krebb Museum, and field samples from Sarah Mussulman,

Caren Goldberg, and Mallory Bedwell. The authors have no conflicts of interest to

declare.

**Literature Cited**

Ali, O. A., S. M. O'Rourke, S. J. Amish, M. H. Meek, G. Luikart, C. Jeffres, and M. R. Miller.

2016. RAD Capture (Rapture): Flexible and Efficient Sequence-Based Genotyping.

Genetics 202:389–400.

AmphibiaWeb. 2020. Species numbers.

https://amphibiaweb.org/amphibian/speciesnums.html.

Baird, N. A., P. D. Etter, T. S. Atwood, M. C. Currey, A. L. Shiver, Z. A. Lewis, E. U. Selker,

W. A. Cresko, and E. A. Johnson. 2008. Rapid SNP discovery and genetic

mapping using sequenced RAD markers. PLoS One 3:e3376.

Bondi, C. A., S. M. Yarnell, and A. J. Lind. 2013. Transferability of habitat suitability

criteria for a stream breeding frog (*Rana boylii*) in the Sierra Nevada, California.

Conservation and Biology 8:88–103.

Botero, C. A., F. J. Weissing, J. Wright, and D. R. Rubenstein. 2015. Evolutionary tipping

points in the capacity to adapt to environmental change. Proceedings of the

National Academy of Sciences of the United States of America 112:184–189.

Bourque, R. M. 2008, December. Spatial ecology of an inland population of the Foothill

yellow-legged frog (*Rana boylii*) in Tehama County, California. MSc, Humboldt

State University, USA.

Broquet, T., L. Berset-Braendli, G. Emaresi, and L. Fumagalli. 2007. Buccal swabs allow

efficient and reliable microsatellite genotyping in amphibians. Conservation

Genetics 8:509–511.

Brown, L. R., J. T. May, A. C. Rehn, P. R. Ode, I. R. Waite, and J. G. Kennen. 2012.

Predicting biological condition in southern California streams. Landscape and

Urban Planning 108:17–27.

Bunn, S. E., and A. H. Arthington. 2002. Basic principles and ecological consequences of

altered flow regimes for aquatic biodiversity. Environmental Management

30:492–507.

Cazé, A. L. R., G. Mäder, T. S. Nunes, L. P. Queiroz, G. de Oliveira, J. A. F. Diniz-Filho, S. L.

Bonatto, and L. B. Freitas. 2016. Could refuge theory and rivers acting as barriers

explain the genetic variability distribution in the Atlantic Forest? Molecular

Phylogenetics and Evolution 101:242–251.

Colwell, R. K. 1974. Predictability, Constancy, and Contingency of Periodic Phenomena.

Ecology 55:1148–1153.

Davidson, C., H. B. Shaffer, and M. R. Jennings. 2002. Spatial Tests of the Pesticide

Drift, Habitat Destruction, UV-B, and Climate-Change Hypotheses for California

Amphibian Declines. Conservation Biology 16:1588–1601.

De'ath, G. 2007. Boosted trees for ecological modeling and prediction. Ecology 88:243–

251.

Deiner, K., J. C. Garza, R. Coey, and D. J. Girman. 2007. Population structure and genetic

diversity of trout (*Oncorhynchus mykiss*) above and below natural and man-made

barriers in the Russian River, California. Conservation Genetics 8:437–454.

Dudgeon, D., A. H. Arthington, M. O. Gessner, Z.-I. Kawabata, D. J. Knowler, C. Lévêque,

R. J. Naiman, A.-H. Prieur-Richard, D. Soto, M. L. J. Stiassny, and C. A. Sullivan.

2006. Freshwater biodiversity: importance, threats, status and conservation

challenges. Biological Reviews of the Cambridge Philosophical Society 81:163–

182.

Edwards, R. J., D. E. Tuipulotu, T. G. Amos, D. O'Meally, M. F. Richardson, T. L. Russell,

M. Vallinoto, M. Carneiro, N. Ferrand, M. R. Wilkins, F. Sequeira, L. A. Rollins, E. C.

Holmes, R. Shine, and P. A. White. 2018. Draft genome assembly of the invasive

cane toad, *Rhinella marina*. GigaScience 7.

Fahrig, L., and G. Merriam. 1985. Habitat Patch Connectivity and Population Survival.

Ecology 66:1762–1768.

Frankham, R. 2002. Introduction to Conservation Genetics. Cambridge University Press,

Cambridge, UK ; New York.

Fuller, T. E., K. L. Pope, D. T. Ashton, and H. H. Welsh. 2011. Linking the Distribution of

an Invasive Amphibian (*Rana catesbeiana*) to Habitat Conditions in a Managed

River System in Northern California. Restoration Ecology 19:204–213.

Fumagalli, M., F. G. Vieira, T. S. Korneliussen, T. Linderoth, E. Huerta-Sánchez, A.

Albrechtsen, and R. Nielsen. 2013. Quantifying population genetic differentiation

from next-generation sequencing data. Genetics 195:979–992.

Gilpin, M., and M. Soule. 1986. Minimum Viable Populations: Processes of Species

Extinction, Conservation Biology. Sunderland, Massachusetts: Sin. Assoc.:19–34.

Goldberg, C. S., M. E. Kaplan, and C. R. Schwalbe. 2003. From the frog's mouth: buccal

swabs for collection of DNA from amphibians. Herpetological Review 34:220–

221.

Graham, C. H., J. Elith, R. J. Hijmans, A. Guisan, A. Townsend Peterson, B. A. Loiselle,

and The Nceas Predicting Species Distributions Working Group. 2008. The

influence of spatial errors in species occurrence data used in distribution

models. The Journal of Applied Ecology 45:239–247.

Grantham, T. E., A. M. Merenlender, and V. H. Resh. 2010. Climatic influences and

anthropogenic stressors: an integrated framework for streamflow management

in Mediterranean-climate California, USA. Freshwater Biology 55:188–204.

Guisan, A., N. E. Zimmermann, J. Elith, C. H. Graham, S. Phillips, and A. T. Peterson.

2007. What Matters for Predicting the Occurrences of Trees: Techniques, Data, or

Species' Characteristics? Ecological Monographs 77:615–630.

Guzy, J. C., E. A. Eskew, B. J. Halstead, and S. J. Price. 2018. Influence of damming on

anuran species richness in riparian areas: A test of the serial discontinuity

concept. Ecology and Evolution 8:2268-2279.

Hammond, S. A., R. L. Warren, B. P. Vandervalk, E. Kucuk, H. Khan, E. A. Gibb, P. Pandoh,

H. Kirk, Y. Zhao, M. Jones, A. J. Mungall, R. Coope, S. Pleasance, R. A. Moore, R.

A. Holt, J. M. Round, S. Ohora, B. V. Walle, N. Veldhoen, C. C. Helbing, and I. Birol.

2017. The North American bullfrog draft genome provides insight into hormonal

regulation of long noncoding RNA. Nature Communications 8:1433.

Hellsten, U., R. M. Harland, M. J. Gilchrist, D. Hendrix, J. Jurka, V. Kapitonov, I.

Ovcharenko, N. H. Putnam, S. Shu, L. Taher, I. L. Blitz, B. Blumberg, D. S.

Dichmann, I. Dubchak, E. Amaya, J. C. Detter, R. Fletcher, D. S. Gerhard, D.

Goodstein, T. Graves, I. V. Grigoriev, J. Grimwood, T. Kawashima, E. Lindquist, S.

M. Lucas, P. E. Mead, T. Mitros, H. Ogino, Y. Ohta, A. V. Poliakov, N. Pollet, J.

Robert, A. Salamov, A. K. Sater, J. Schmutz, A. Terry, P. D. Vize, W. C. Warren, D.

Wells, A. Wills, R. K. Wilson, L. B. Zimmerman, A. M. Zorn, R. Grainger, T.

Grammer, M. K. Khokha, P. M. Richardson, and D. S. Rokhsar. 2010. The genome

of the Western clawed frog *Xenopus tropicalis*. Science 328:633–636.

Heyer, W. R., M. A. Donnelly, R. W. McDiarmid, L.-A. C. Hayek, and M. S. Foster. 1994.

Measuring and monitoring biological diversity. Standard methods for

amphibians. Smithsonian Institution Press, Washington DC.

Hijmans, R. J., S. Phillips, J. Leathwick, and J. Elith. 2017. dismo: Species Distribution

Modeling. R package.

Hoffmann, A. A., and C. M. Sgrò. 2011. Climate change and evolutionary adaptation.

Nature 470:479–485.

Hotaling, S., C. C. Muhlfeld, J. J. Giersch, O. A. Ali, S. Jordan, M. R. Miller, G. Luikart, and

D. W. Weisrock. 2018. Demographic modelling reveals a history of divergence

with gene flow for a glacially tied stonefly in a changing post-Pleistocene

landscape. Journal of Biogeography 45:304–317.

Ishiyama, N., I. Koizumi, T. Yuta, and F. Nakamura. 2015. Differential effects of spatial

network structure and scale on population size and genetic diversity of the

ninespine stickleback in a remnant wetland system. Freshwater Biology 60:733–

744.

Jennings, M. R., and M. P. Hayes. 1994. Amphibian and reptile species of special

concern in California. California Department of Fish and Game, Rancho Cordova.

Kahilainen, A., M. Puurtinen, and J. S. Kotiaho. 2014. Conservation implications of

species–genetic diversity correlations. Global Ecology and Conservation 2:315–

323.

Kim, S. Y., K. E. Lohmueller, A. Albrechtsen, Y. Li, T. Korneliussen, G. Tian, N. Grarup, T.

Jiang, G. Andersen, D. Witte, T. Jorgensen, T. Hansen, O. Pedersen, J. Wang, and

R. Nielsen. 2011. Estimation of allele frequency and association mapping using

next-generation sequencing data. BMC Bioinformatics 12:231.

Korneliussen, T. S., A. Albrechtsen, and R. Nielsen. 2014. ANGSD: Analysis of Next

Generation Sequencing Data. BMC Bioinformatics 15:356.

Korneliussen, T. S., I. Moltke, A. Albrechtsen, and R. Nielsen. 2013. Calculation of

Tajima's D and other neutrality test statistics from low depth next-generation

sequencing data. BMC Bioinformatics 14:289.

Krohn, A. R., C. J. Conroy, R. Pesapane, K. Bi, J. E. Foley, and E. B. Rosenblum. 2018.

Conservation genomics of desert dwelling California voles (*Microtus californicus*)

and implications for management of endangered Amargosa voles (*Microtus*

*californicus scirpensis*). Conservation Genetics 19:383–395.

Kupferberg, S. J. 1996. Hydrologic and geomorphic factors affecting conservation of a

river-breeding frog (*Rana boylii*). Ecological Applications 6:1332–1344.

Kupferberg, S. J., W. J. Palen, A. J. Lind, S. Bobzien, A. Catenazzi, J. Drennan, and M. E.

Power. 2012. Effects of flow regimes altered by dams on survival, population

declines, and range-wide losses of California river-breeding frogs. Conservation

Biology 26:513–524.

Lande, R., and S. Shannon. 1996. The role of genetic variation in adaptation and

population persistence in a changing environment. Evolution 50:434–437.

Li, H. 2011. A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data.

Bioinformatics 27:2987–2993.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis,

and R. Durbin. 2009. The Sequence Alignment/Map format and SAMtools.

Bioinformatics 25:2078–2079.

Lind, A. J., H. H. Welsh Jr, and R. A. Wilson. 1996. The effects of a dam on breeding

habitat and egg survival of the foothill yellow-legged frog (*Rana boylii*) in

Northwestern Calfifornia. Herpetological Review 27:62–66.

Miller, M. R., J. P. Brunelli, P. A. Wheeler, S. Liu, C. E. Rexroad 3rd, Y. Palti, C. Q. Doe, and

G. H. Thorgaard. 2012. A conserved haplotype controls parallel adaptation in

geographically distant salmonid populations. Molecular Ecology 21:237–249.

Miller, M. R., J. P. Dunham, A. Amores, W. A. Cresko, and E. A. Johnson. 2007. Rapid and

cost-effective polymorphism identification and genotyping using restriction site

associated DNA (RAD) markers. Genome Research 17:240–248.

Moyle, P. B., J. V. E. Katz, and R. M. Quiñones. 2011. Rapid decline of California's native

inland fishes: A status assessment. Biological Conservation 144:2414–2423.

Murchie, K. J., K. P. E. Hair, C. E. Pullen, T. D. Redpath, H. R. Stephens, and S. J. Cooke.

2008. Fish response to modified flow regimes in regulated rivers: research

methods, effects and opportunities. River Research and Applications 24:197–

217.

Narum, S. R., C. A. Buerkle, J. W. Davey, M. R. Miller, and P. A. Hohenlohe. 2013.

Genotyping-by-sequencing in ecological and conservation genomics. Molecular

Ecology 22:2841–2847.

Nielsen, R., T. Korneliussen, A. Albrechtsen, Y. Li, and J. Wang. 2012. SNP calling,

genotype calling, and sample allele frequency estimation from New-Generation

Sequencing data. PloS One 7:e37558.

Nilsson, C., C. A. Reidy, M. Dynesius, and C. Revenga. 2005. Fragmentation and flow

regulation of the world's large river systems. Science 308:405–408.

Nunziata, S. O., S. L. Lance, D. E. Scott, E. M. Lemmon, and D. W. Weisrock. 2017.

Genomic data detect corresponding signatures of population size change on an

ecological time scale in two salamander species. Molecular Ecology 26:1060–

1074.

Peek, R. A. 2018. Population Genetics of a Sentinel Stream-breeding Frog (*Rana boylii*).

Ecology, University of California, Davis.

Peek, R. A., M. Bedwell, S. M. O'Rourke, C. Goldberg, G. M. Wengert, and M. R. Miller.

2019. Hybridization between two parapatric ranid frog species in the northern

Sierra Nevada, California, USA. Molecular Ecology 28:4636–4647.

Phillips, S. J., R. P. Anderson, and R. E. Schapire. 2006. Maximum entropy modeling of

species geographic distributions. Ecological Modelling 190:231–259.

Pidancier, N., C. Miquel, and C. Miaud. 2003. Buccal swabs as a non-destructive tissue

sampling method for DNA analysis in amphibians. The Herpetological Journal

13:175–178.

Poff, N. L., J. D. Olden, D. M. Merritt, and D. M. Pepin. 2007. Homogenization of regional

river dynamics by dams and global biodiversity implications. Proceedings of the

National Academy of Sciences of the United States of America 104:5732–5737.

Power, M. E., W. E. Dietrich, and J. C. Finlay. 1996. Dams and Downstream Aquatic

Biodiversity: Potential Food Web Consequences of Hydrologic and Geomorphic

Change. Environmental Management 20:887–895.

Prince, D. J., S. M. O'Rourke, T. Q. Thompson, O. A. Ali, H. S. Lyman, I. K. Saglam, T. J.

Hotaling, A. P. Spidle, and M. R. Miller. 2017. The evolutionary basis of premature

migration in Pacific salmon highlights the utility of genomics for informing

conservation. Science Advances 3:e1603198.

Pringle, C. 2003. What is hydrologic connectivity and why is it ecologically important?

Hydrological Processes 17:2685–2689.

Pringle, C. M. 2001. Hydrologic Connectivity and the Management of Biological

Reserves: A Global Perspective. Ecological Applications 11:981–998.

R Core Team. 2017. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Ridgeway, G. 2015. gbm: Generalized Boosted Regression Models. http://CRAN.R-

project.org/package-gbm.

Rogers, R. L., L. Zhou, C. Chu, R. Márquez, A. Corl, T. Linderoth, L. Freeborn, M. D.

MacManes, Z. Xiong, J. Zheng, C. Guo, X. Xun, M. R. Kronforst, K. Summers, Y.

Wu, H. Yang, C. L. Richards-Zawacki, G. Zhang, and R. Nielsen. 2018. Genomic

Takeover by Transposable Elements in the Strawberry Poison Frog. Molecular

Biology and Evolution 35:2913–2927.

Rolls, R. J., and N. R. Bond. 2017. Environmental and Ecological Effects of Flow

Alteration in Surface Water Ecosystems. Pages 65–82 *in* Water for the

Environment. Elsevier.

Rousset, F. 1997. Genetic differentiation and estimation of gene flow from F-statistics

under isolation by distance. Genetics 145:1219–1228.

Ruby, J. G., P. Bellare, and J. L. Derisi. 2013. PRICE: software for the targeted assembly

of components of (Meta) genomic sequence data. G3 3:865–880.

Sabo, J. L., A. Ruhi, G. W. Holtgrieve, V. Elliott, M. E. Arias, P. B. Ngor, T. A. Räsänen, and

S. Nam. 2017. Designing river flows to improve food security futures in the Lower

Mekong Basin. Science 358.

Sağlam, İ. K., J. Baumsteiger, M. J. Smith, J. Linares-Casenave, A. L. Nichols, S. M.

O'Rourke, and M. R. Miller. 2016. Phylogenetics support an ancient common

origin of two scientific icons: Devils Hole and Devils Hole pupfish. Molecular

Ecology 25:3962–3973.

Schick, R. S., and S. T. Lindley. 2007. Directed connectivity among fish populations in a

riverine network. The Journal of Applied Ecology 44:1116–1126.

Scribner, K. T., W. H. Lowe, E. Landguth, G. Luikart, D. M. Infante, G. E. Whelan, and C. C.

Muhlfeld. 2016. Applications of Genetic Data to Improve Management and

Conservation of River Fishes and Their Habitats. Fisheries 41:174–188.

Seidl, F., N. A. Levis, R. Schell, D. W. Pfennig, K. S. Pfennig, and I. M. Ehrenreich. 2019.

Genome of Spea multiplicata, a Rapidly Developing, Phenotypically Plastic, and

Desert-Adapted Spadefoot Toad. G3 9:3909–3919.

Session, A. M., Y. Uno, T. Kwon, J. A. Chapman, A. Toyoda, S. Takahashi, A. Fukui, A.

Hikosaka, A. Suzuki, M. Kondo, S. J. van Heeringen, I. Quigley, S. Heinz, H. Ogino,

H. Ochi, U. Hellsten, J. B. Lyons, O. Simakov, N. Putnam, J. Stites, Y. Kuroki, T.

Tanaka, T. Michiue, M. Watanabe, O. Bogdanovic, R. Lister, G. Georgiou, S. S.

Paranjpe, I. van Kruijsbergen, S. Shu, J. Carlson, T. Kinoshita, Y. Ohta, S.

Mawaribuchi, J. Jenkins, J. Grimwood, J. Schmutz, T. Mitros, S. V. Mozaffari, Y.

Suzuki, Y. Haramoto, T. S. Yamamoto, C. Takagi, R. Heald, K. Miller, C.

Haudenschild, J. Kitzman, T. Nakayama, Y. Izutsu, J. Robert, J. Fortriede, K.

Burns, V. Lotay, K. Karimi, Y. Yasuoka, D. S. Dichmann, M. F. Flajnik, D. W.

Houston, J. Shendure, L. DuPasquier, P. D. Vize, A. M. Zorn, M. Ito, E. M. Marcotte,

J. B. Wallingford, Y. Ito, M. Asashima, N. Ueno, Y. Matsuda, G. J. Veenstra, A.

Fujiyama, R. M. Harland, M. Taira, and D. S. Rokhsar. 2016. Genome evolution in

the allotetraploid frog *Xenopus laevis*. Nature 538:336–343.

Shaw, E. A., E. Lange, J. D. Shucksmith, and D. N. Lerner. 2016. Importance of partial

barriers and temporal variation in flow when modelling connectivity in

fragmented river systems. Ecological Engineering 91:515–528.

Slatkin, M. 1987. Gene flow and the geographic structure of natural populations.

Science 236:787–792.

Stebbins, R. C. 2003. A Field Guide to Western Reptiles and Amphibians. 3rd edition.

Houghton Mifflin Harcourt, Boston.

Steel, A. E., R. A. Peek, R. A. Lusardi, and S. M. Yarnell. 2017. Associating metrics of

hydrologic variability with benthic macroinvertebrate communities in regulated

and unregulated snowmelt-dominated rivers. Freshwater Biology 63:844-858.

Sun, Y.-B., Z.-J. Xiong, X.-Y. Xiang, S.-P. Liu, W.-W. Zhou, X.-L. Tu, L. Zhong, L. Wang, D.-D.

Wu, B.-L. Zhang, C.-L. Zhu, M.-M. Yang, H.-M. Chen, F. Li, L. Zhou, S.-H. Feng, C.

Huang, G.-J. Zhang, D. Irwin, D. M. Hillis, R. W. Murphy, H.-M. Yang, J. Che, J.

Wang, and Y.-P. Zhang. 2015. Whole-genome sequence of the Tibetan frog

Nanorana parkeri and the comparative evolution of tetrapod genomes.

Proceedings of the National Academy of Sciences of the United States of

America 112:E1257-62.

Tajima, F. 1983. Evolutionary relationship of DNA sequences in finite populations.

Genetics 105:437–460.

Tajima, F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA

polymorphism. Genetics 123:585–595.

Taylor, P. D., L. Fahrig, K. Henein, and G. Merriam. 1993. Connectivity Is a Vital Element

of Landscape Structure. Oikos 68:571–573.

Thompson, T. Q., M. R. Bellinger, S. M. O'Rourke, D. J. Prince, A. E. Stevenson, A. T.

Rodrigues, M. R. Sloat, C. F. Speller, D. Y. Yang, V. L. Butler, M. A. Banks, and M.

R. Miller. 2019. Anthropogenic habitat alteration leads to rapid loss of adaptive

variation and restoration potential in wild salmon populations. Proceedings of

the National Academy of Sciences of the United States of America 116:177–186.

Tonkin, J. D., M. T. Bogan, N. Bonada, B. Rios-Touma, and D. A. Lytle. 2017. Seasonality

and predictability shape temporal species diversity. Ecology 98:1201–1216.

Torrence, C., and G. P. Compo. 1998. A Practical Guide to Wavelet Analysis. Bulletin of

the American Meteorological Society 79:61–78.

Tyers, M. 2017. riverdist: River Network Distance Computation and Applications.

Voelker, G., B. D. Marks, C. Kahindo, U. A'genonga, F. Bapeamoni, L. E. Duffie, J. W.

Huntley, E. Mulotwa, S. A. Rosenbaum, and J. E. Light. 2013. River barriers and

cryptic biodiversity in an evolutionary museum. Ecology and Evolution 3:536–

545.

Vörösmarty, C. J., P. B. McIntyre, M. O. Gessner, D. Dudgeon, A. Prusevich, P. Green, S.

Glidden, S. E. Bunn, C. A. Sullivan, C. R. Liermann, and P. M. Davies. 2010. Global

threats to human water security and river biodiversity. Nature 467:555–561.

Wakeley, J. 2009. Coalescent Theory: An Introduction. W. H. Freeman.

Watterson, G. A. 1975. On the number of segregating sites in genetical models without

recombination. Theoretical Population Biology 7:256–276.

Werth, S., M. Schödl, and C. Scheidegger. 2014. Dams and canyons disrupt gene flow

among populations of a threatened riparian plant. Freshwater Biology 59:2502–

2515.

Whiteley, A. R., K. Hastings, J. K. Wenburg, C. A. Frissell, J. C. Martin, and F. W.

Allendorf. 2010. Genetic variation and effective population size in isolated

populations of coastal cutthroat trout. Conservation Genetics 11:1929–1943.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York.

Wiens, J. A. 2002. Riverine landscapes: taking landscape ecology into the water.

Freshwater Biology 47:501–515.

Wright, S. 1943. Isolation by Distance. Genetics 28:114–138.

Yarnell, S. M., E. D. Stein, J. A. Webb, T. Grantham, R. A. Lusardi, J. Zimmerman, R. A.

Peek, B. A. Lane, J. Howard, and S. Sandoval-Solis. 2020. A functional flows

approach to selecting ecologically relevant flow metrics for environmental flow

applications. River Research and Applications 9:12.

Yarnell, S. M., J. H. Viers, and J. F. Mount. 2010. Ecology and Management of the Spring

Snowmelt Recession. Bioscience 60:114–127.

Yarnell, S., R. Peek, G. Epke, and A. Lind. 2016. Management of the Spring Snowmelt

Recession in Regulated Systems. Journal of the American Water Resources

Association 52:723–736.

Zhong, Y., and G. Power. 1996. Environmental impacts of hydroelectric projects on fish

resources in China. Regulated Rivers: Research & Management 12:81–98.

***Data Archiving Statement:*** Should the manuscript be accepted, the data supporting the

results will be archived in an appropriate public repository such as Dryad, and the data

DOI will be appended to the end of the article.

Table 1. Locality information for population genomic analysis of *R. boylii* in the Yuba,

Bear, and American Watersheds.

| Locality | Site ID | River | Watershed | Regulation Type | Lat. (WGS84) | Long. (WGS84) | Elev. (m) | NHD[†] Stream Order | NHD Total Drainage Area (sq. km) |
|---|---|---|---|---|---|---|---|---|---|
| American Canyon | 1 | MF American | American | Hydropeaking | 38.93396 | -120.94356 | 240 | 2 | 9 |
| Gas Canyon | 2 | MF American | American | Hydropeaking | 38.96651 | -120.93252 | 241 | 1 | 13 |
| Todd Creek | 3 | MF American | American | Hydropeaking | 38.96385 | -120.92158 | 367 | 2 | 10 |
| NFMFA-Skunk Canyon | 4 | MF American | American | Hydropeaking | 39.02237 | -120.73693 | 521 | 2 | 6 |
| Rubicon-US Powerhouse | 5 | MF American | American | Bypass | 38.99928 | -120.72333 | 360 | 5 | 816 |
| Rubicon-Long Canyon | 6 | MF American | American | Bypass | 38.98887 | -120.68996 | 415 | 5 | 806 |
| Ponderosa Bridge | 7 | NF American | American | Unregulated | 38.99994 | -120.9406 | 240 | 5 | 857 |
| Bunch Canyon | 8 | NF American | American | Unregulated | 39.03762 | -120.91028 | 286 | 3 | 27 |
| Shirttail Creek | 9 | NF American | American | Unregulated | 39.04446 | -120.89937 | 525 | 4 | 141 |
| Indian Creek | 10 | NF American | American | Unregulated | 39.05665 | -120.90854 | 296 | 2 | 24 |
| Slaughter Ravine | 11 | NF American | American | Unregulated | 39.09865 | -120.92548 | 356 | 2 | 6 |
| Robbers Ravine | 12 | NF American | American | Unregulated | 39.10451 | -120.92672 | 400 | 1 | 4 |
| Iowa Hill Mainstem | 13 | NF American | American | Unregulated | 39.11115 | -120.91684 | 386 | 4 | 605 |
| Euchre Bar | 14 | NF American | American | Unregulated | 39.18492 | -120.76204 | 579 | 4 | 508 |
| Sailor Canyon | 15 | NF American | American | Unregulated | 39.21694 | -120.49595 | 1005 | 3 | 166 |
| Greenhorn Creek | 16 | Bear | Bear | Bypass | 39.23206 | -120.90189 | 820 | 2 | 11 |
| Steep Hollow Creek DS | 17 | Bear | Bear | Bypass | 39.20231 | -120.87537 | 736 | 2 | 52 |
| Hawkins Ravine | 18 | Bear | Bear | Bypass | 39.18833 | -120.8981 | 706 | 2 | 4 |
| Steep Hollow Creek US | 19 | Bear | Bear | Bypass | 39.19444 | -120.88778 | 704 | 2 | 45 |
| Chicago Powerhouse | 20 | Bear | Bear | Bypass | 39.17484 | -120.89978 | 665 | 4 | 136 |
| Shady Creek | 21 | South Yuba | Yuba | Bypass | 39.35433 | -121.059 | 675 | 2 | 15 |
| Spring Creek | 22 | South Yuba | Yuba | Bypass | 39.33233 | -120.989 | 595 | 3 | 24 |
| Rock Creek | 23 | South Yuba | Yuba | Bypass | 39.32983 | -120.98634 | 593 | 4 | 710 |
| Missouri Canyon | 24 | South Yuba | Yuba | Bypass | 39.36096 | -120.88138 | 1094 | 2 | 5 |
| Logan Creek | 25 | South Yuba | Yuba | Bypass | 39.36914 | -120.85258 | 1201 | 1 | 5 |
| US Canyon Creek | 26 | South Yuba | Yuba | Bypass | 39.35387 | -120.73419 | 889 | 4 | 365 |
| Oregon Creek | 27 | Middle Yuba | Yuba | Bypass | 39.44188 | -121.05753 | 620 | 4 | 375 |

| Locality | Site ID | River | Watershed | Regulation Type | Lat. (WGS84) | Long. (WGS84) | Elev. (m) | NHD[†] Stream Order | NHD Total Drainage Area (sq. km) |
|---|---|---|---|---|---|---|---|---|---|
| US Our House Dam | 28 | Middle Yuba | Yuba | Bypass | 39.41305 | -120.99035 | 624 | 4 | 375 |
| Rocky Rest Mainstem | 29 | North Yuba | Yuba | Unregulated | 39.51191 | -120.97738 | 704 | 5 | 669 |
| Slate Creek | 30 | North Yuba | Yuba | Bypass | 39.689128 | -120.93889 | 1330 | 3 | 58.7 |

782    [†]NHD refers to the National Hydrography Dataset by USGS (U.S. Geological Survey, National Hydrography

783    Dataset, Digital data, accessed, August 2017).

785　Table 2. Sampling information for population genomic analysis of *R. boylii*, the number

786　of individuals (n) is given for the total number sequenced per location and the number

787　of individuals that were retained after filtering (see methods) across the 8,533 baits.

| Locality | Site ID | n initial | n retained |
|---|---|---|---|
| American Canyon | 1 | 16 | 6 |
| Gas Canyon | 2 | 6 | 6 |
| Todd Creek | 3 | 11 | 9 |
| NFMFA-Skunk Canyon | 4 | 18 | 18 |
| Rubicon-US Powerhouse | 5 | 11 | 11 |
| Rubicon-Long Canyon | 6 | 9 | 8 |
| Ponderosa Bridge | 7 | 5 | 5 |
| Bunch Canyon | 8 | 15 | 14 |
| Shirttail Creek | 9 | 16 | 15 |
| Indian Creek | 10 | 12 | 11 |
| Slaughter Ravine | 11 | 8 | 8 |
| Robbers Ravine | 12 | 30 | 11 |
| Iowa Hill Mainstem | 13 | 36 | 30 |
| Euchre Bar | 14 | 13 | 11 |
| Sailor Canyon | 15 | 8 | 5 |
| Greenhorn Creek | 16 | 15 | 6 |
| Steep Hollow Creek DS | 17 | 13 | 12 |
| Hawkins Ravine | 18 | 3 | 3 |
| Steep Hollow Creek US | 19 | 7 | 6 |
| Chicago Powerhouse | 20 | 6 | 6 |
| Shady Creek | 21 | 14 | 12 |
| Spring Creek | 22 | 4 | 4 |
| Rock Creek | 23 | 3 | 3 |
| Missouri Canyon | 24 | 8 | 6 |
| Logan Creek | 25 | 5 | 4 |
| US Canyon Creek | 26 | 6 | 6 |
| Oregon Creek | 27 | 15 | 13 |
| US Our House Dam | 28 | 13 | 12 |
| Rocky Rest Mainstem | 29 | 15 | 12 |
| Slate Creek | 30 | 4 | 4 |

Figure 1. Sampling locations and flow characteristics. A) Map of sampling locations

across three watersheds. B) Comparison of annual mean daily discharge from 1981–

2016 for three flow types. C) Comparison of hourly discharge in three different flow

regimes in April through July 2012, Bypass (South Yuba), Hydropeaking (Middle Fork

American), and Unregulated (North Fork American). See Table S1 for USGS gaging

station information.

Figure 2. A biplot of seasonality (Colwell's M/P [contingency/predictability] versus

predictability (average wavelet power at the 12-month time period following Tonkin et

al. (2017). Each point is labeled with the river acronym and USGS gage ID, followed by

the end year of the flow dataset used (See Table S3). Data from the same gage but from

two time periods (before flow alteration and with contemporary alteration) are

connected by arrows.

Figure 3. Principal component analysis of Rapture sequencing data. A) Northern Sierra

Nevada watersheds; B) Unregulated NF American; C) and D) Hydropeaking MF

American Reach.

Figure 4. Relationship between river distance and genetic differentiation in *R. boylii*. A)

pairwise scaled $F_{ST}$ vs. river distance (km) for each location pair, centroid of each flow

alteration type delineated as large point with black outline. Numbers on points

correspond to "sitepairID" in Data S7; B) Relative influence of covariates on scaled $F_{ST}$

from boosted regression tree models.

Figure 5. Relationship between river regulation and genetic diversity trajectory in *R*.

*boylii*. A) Assessment of genetic diversity trajectories using scaled Δθ ($\theta_\pi - \theta_S$ / [$\theta_\pi + \theta_S$]

/ 2) for each sampling location; B) Boxplots of scaled Δθ and pairwise significance

between regulation groups using a pairwise Wilcoxon rank sum test with Bonferroni

correction ($P < 0.05$). Negative values represent trends of increasing genetic diversity,

positive values represent trajectories of diversity loss, values near zero are stable.