Methylation patterns reveal cryptic structure and a pathway for adaptive divergence in a
panmictic carnivore

Justin B. Johnson*[1], Dennis L. Murray[1,2], Aaron B.A. Shafer[1,3]

1. Environmental and Life Sciences Graduate Program, Trent University, K9J 7B8
Peterborough, Canada
2. Biology Department, Trent University, K9J 7B8 Peterborough, Canada
3. Forensics Department, Trent University, K9J 7B8 Peterborough, Canada

*corresponding author

Address: 1600 W Bank Drive. Peterborough, ON, Canada, K9J 7B8.

e-mail: justinjohnson@trentu.ca

Summary

Determining molecular signatures of population divergence is a fundamental component of

evolutionary biology. Identifying divergence is particularly challenging between populations of

highly mobile species that undergo substantial gene flow, such as the Canada lynx (*Lynx*

canadensis), where populations are considered panmictic when only neutral genetic markers

are considered. Here, we used high-throughput bisulfite sequencing to examine the

environmental determinants of methylation structure across the distributional range of Canada

lynx. Despite a high degree of genetic similarity among mainland populations, epigenetic

structure did not mirror these neutral patterns, instead showing environmental associations and

high levels of divergence in the peripheral Newfoundland and Alaskan populations. Interestingly,

a disproportionate number of genes related to body-size were hypermethylated on the island of

Newfoundland, providing a possible mechanism for adaptive evolution and the observed island

effect on organism size. Our results indicate that epigenetic modifications, specifically DNA

methylation, are powerful markers to investigate adaptive divergence and rapid evolutionary

response.

Keywords: Ecological genetics, genomics / methylomics, adaptation, non-model organisms

Background

Investigations into divergent selection have often relied on quantifying phenotypic variation and the heritability of such traits, the latter assumed to be genetic polymorphisms transmitted via an organism's DNA[1]. However, detecting adaptive divergence in species that experience high rates of gene flow is challenging due to the homogenization of genomic regions that are neutral or under weak selection[2]. Environmental conditions can be a powerful driver of adaptive divergence, where relationships are traditionally ascertained by correlating allele frequencies to environmental variation[3]. An aspect of adaptive molecular evolution that is undetected by standard genetic sequencing involves direct modifications to the structure of DNA. Epigenetic modifications like DNA methylation are influenced by environmental conditions, directly affect gene expression, and may be indicative of early adaptive divergence due to local adaptation[4–6]. DNA methylation could play a role in local adaptation due to its regulatory role in transcription by modifying chromatin structure, repressing transcription factors, or recruiting protein complexes that block transcriptional machinery, especially around CpG islands[7–9]. CpG islands are dense clusters of cytosine-guanine dinucleotides and frequently occur near the transcription start site of genes and have functional role with gene expression[10–12]. Consequently, DNA methylation, especially around CpG islands, could explain the molecular basis of local adaptation due to its regulatory function on gene expression, and might be an overlooked molecular marker of adaptive divergence and rapid evolutionary adaptation.

Here, we assessed whether environmental variation, geographic distance, or insularity were determinants of DNA methylation structure in a free-ranging carnivore. Our study species, the Canada lynx, (*Lynx canadensis*) is a mid-sized felid that is highly mobile and whose neutral genetic variation (i.e., microsatellites) exhibits low levels of genetic differentiation across the mainland, with divergent island populations[13,14]. Despite this low degree of overall genetic structure, two potential mechanisms might drive epigenetic divergence in Canada lynx. First,

allele frequencies are correlated to climatic gradients in both population time-series and fine-

scale genetic analyses[13,15,16], suggesting climate might also influence patterns of methylation in

lynx. Second, Canada lynx show a subtle cline in body size with larger individuals in Alaska[17] to

smaller individuals in insular populations including Newfoundland and Cape Breton Island[18].

Body size changes in island populations appear to be consistent with the 'Island Rule'[18], where

insular mammals are smaller in size compared to their mainland counterparts[19–22]. If functional

genes related to body size are repressed in geographically isolated populations, then epigenetic

modifications might be an underlying mechanism driving population divergence[23]. Based on

these two mechanisms (climate and island isolation), we tested two predictions: i) climatic

conditions are the driving force behind spatial epigenetic structuring[24]; and ii) patterns of DNA

methylation over CpG islands and gene bodies are more correlated with biogeographical

substructuring than methylation patterns over DNA of unknown function, due to the regulation of

transcription arising from adaptation to local conditions[25].

Results

We identified differential methylation at base-pair resolution across the genome in 95 Canada

lynx epidermal tissue samples from four populations (n = 23-24 per sampling area) across North

America, including one insular population in Newfoundland (Figure 1; Supplemental Table S4).

The sampled populations have a wide geographical spread, with an average minimum distance

between populations (Québec and Newfoundland) of 1,158 km and a maximum distance

(Alaska and Newfoundland) of 5,520 km. Habitats around these populations present a dynamic

range of environmental conditions, ranging from 32 to 432 mm of winter precipitation and a

mean annual temperature range[26] of -6.3 to 4.7° C. To determine associations between these

environmental conditions and genome-wide patterns of DNA methylation, we created a reduced

representation bisulfite sequencing (RRBS) library (full protocol in Supplemental). Paired-end

sequencing on a HiSeq2500 generated a total of 210,773,612 filtered and demultiplexed reads

that were aligned to the domestic cat genome (85.0% average mapping success; Supplemental

Table S4; *Felis catus*; NCBI felcat9.0) and variants were called using specially designed

software for bisulfite converted reads[27]. We mapped our reads to the human (*Homo sapiens*)

and lambda phage genomes to rule out contamination (2.7% and <0.1% success, respectively),

and assessed bisulfite conversion efficiency by including non-methylated lambda phage DNA in

the sequencing lane. Furthermore, we quantified the temporal effects of methylation, the

ramifications of missing data, and model parameterization with sensitivity analyses, with no

implications on overall inferences (Supplemental Figures S5-S8).

Pronounced population structure revealed by methylation patterns

To examine regions with putative regulatory function, CpG islands were first bioinformatically

identified *de novo* (N = 28,127) using hidden Markov models and contained an average GC

content of 59.2% with a posterior probability of observed-to-expected GC content (CpG$_{o/e}$) of

1.13. Our DNA methylation analysis was broken into two subsets based on proximity to genomic

features, with identical filtering parameters for each subset to maximize comparative inferences.

Our subset over CpG islands and gene bodies contained 329 5,00-bp windows with 4,611 CpG

positions, while the unannotated dataset contained 376 5,000-bp windows with 5,031 CpG

positions (Supplemental Table S6). Qualitatively, population structure between geographically

peripheral populations was most pronounced in DNA methylation patterns over CpG islands and

gene bodies compared to methylation patterns over regions of unknown function. The first two

axes of a principal coordinates analysis (PCoA) on a Euclidean distance matrix summarizing

CpG island and gene body methylation (PCoA1 = 6.2%, PCoA2 = 5.0% variation explained)

distinctly cluster Alaska from the remaining mainland populations, while the mid-continental

populations show no structure (Figure 2). Similar, but less distinct patterns were seen in the

ordination of methylation patterns over unannotated regions of the genome, which showed more

distinct structure between the Québec and Manitoba populations (Figure 2). Analogous

population-level trends between the datasets were further confirmed with Pearson and

Spearman's rank coefficients of the first PCoA axis (r = 0.91; ρ = 0.88).

Environmental variation associated with epigenetic structure

For both methylation datasets, model selection identified three significant variables in our db-

RDA analysis: geographic distance, climate and a binary variable representing insularity for the

Newfoundland population (pseudo-F = 16.26 - 16.27; adjusted R^2 = 0.33; all p = 0.001). Tree

cover (p = 0.13 - 0.50) and a randomly-generated numeric variable to assess the effect of noise

(p = 0.64 – 0.78) added no explanatory power to either model (Supplemental Table S11).

Collinearity was low between all retained variables (VIF = 2.09 – 3.77; Supplemental Table

S12). We examined the explanatory power of each variable independently using partial db-

RDAs (Figure 4), which revealed similar trends between both methylation datasets for

geographic distance (pseudo-F = 15.72 – 17.47; adjusted R^2 = 0.11 – 0.12; p = 0.001) and

climate (pseudo-F = 6.75 – 6.90; adjusted R^2 = 0.04; p = 0.001). However, methylation patterns

over CpG islands and gene bodies were more strongly associated with the insular

Newfoundland population (pseudo-F = 13.66; adjusted R^2 = 0.09; p = 0.001) than with

methylation patterns over unannotated regions (pseudo-F = 10.03; adjusted R^2 = 0.07; p =

0.001), suggesting stronger epigenetic divergence in putatively regulatory regions of the

genome in insular Canada lynx (Figure 3).

Differential methylation over genes related to morphology

We identified differential methylation directly over CpG islands and gene bodies using beta

regressions (n = 329), with population as an explanatory variable (i.e. windows with > 10%

overall difference in methylation and p-value < 0.001). Of the 16 gene regions identified, 11

were significantly differentiated in the Newfoundland population (Table 1). All differentially

methylated regions with associated morphological function were hypermethylated, as well as

genes with direct epigenetic function with morphological interactions (i.e. transcription

regulation, DNA-binding transcription factors; Table 1). The 11 differentially methylated regions

in the insular population have a difference in methylation of 14%-40% compared to mainland

populations, which suggests that morphological divergence in the insular population might be

regulated by repressed gene expression due to DNA methylation. Additionally, we identified

three differentially methylated regions in the Alaskan population related to spectrin,

carbohydrate, and ATP binding, suggesting differential expression of genes with metabolic

function.

Genetic structure driven by insular and climatic divides

We examined the relationship between neutral SNPs, environmental variation, geographic

distance, and insularity by first performing a principal coordinates analysis on a Euclidean

distance dissimilarity matrix of 489 genomic SNPs identified using a Bayesian wildcard

algorithm with conservative calling and filtering parameters[28]. Consistent with previous research

on population structure in Canada lynx[13,29], our SNP data analysis indicated substantial genetic

separation of the insular Newfoundland population (F_{ST} = 0.10 – 0.13). We complemented our

F_{ST} analysis with a null model approach that calculated the relative differentiation between

populations, and mainland populations were relatively similar (F_{ST} = 0.005 – 0.024) with

Newfoundland as the most segregated (F_{ST} = 0.096). Consistent with previous research[14], gene

diversity was lowest in the Newfoundland population (H_e = 0.224). An analysis of molecular

variance (AMOVA) identified substantial genetic variation between populations, which explained

8.58% of the total variation in SNP data (σ = 5.84, p = 0.01). Outlier detection with Bayescan

v2.0[30] (false discovery rate > 0.05) identified no loci under selection.

We determined relationships between genetic data and environmental variables by

performing a db-RDA step-wise model selection on the retained PCoA axes, and identified only

climate (pseudo-F = 51.13; p-value = 0.001) and insularity (pseudo-F = 26.33; p-value = 0.001)

as important for explaining genetic variation (Supplemental Figure S11). Geographic distance

(p-value = 0.31), tree cover (p-value = 0.95), and the randomly generated numeric variable (p-

value = 0.33) were all dropped during the model step-selection. The first axis of the db-RDA

was strongly representative of both climate (db-RDA1 = -0.86) and insularity (db-RDA1 = 0.99),

while the second axis primarily summarized genetic variation associated with climate (db-RDA2

= 0.51). Partial db-RDAs identified a substantial amount of the net variation was explained by

the mainland – Newfoundland divide (pseudo-F = 26.3; adjusted R^2 = 0.15; p-value = 0.001),

while climatic variation explained much of the variation seen in mainland populations (pseudo-F

= 13.70; adj. R^2 = 0.08; p-value = 0.001). We identified a stronger correlation between SNP data

and non-coding region methylation (r = -0.78; ρ = -0.62) than with CpG island and gene body

methylation (r = -0.74; ρ = -0.61) from the first axes of PCoAs using Pearson's and Spearman's

ranking coefficients, respectively.

Discussion

A new chapter in Canada lynx population structure

We have shown that despite seemingly panmictic populations across the mainland at the

genetic level, epigenetic structure is substantial and might be an underlying mechanism driving

local adaptation. Row et al. showed a minimal barrier effect of the Rocky Mountains to Canada

lynx[13], but it remains possible that distinct environmental variation might be driving unique

patterns of functionally important DNA methylation at the range margins, including Alaska and

Newfoundland. Despite the lack of genetic structure between mainland populations[29], we see

distinct patterns of DNA methylation that differentiate Alaska from mid- and east-continental

populations. Furthermore, we identified variation that is associated by trends in winter

precipitation and temperature that is not accounted for by geographic distance.

The differentiated DNA methylation patterns in the Newfoundland population suggest a

molecular pathway for morphological differences between mainland and island populations, as

predicted by the Island Rule. Evidence for this adaptive divergence was observed in disparities

between methylation in CpG islands/gene bodies and DNA of unknown function and specific

genes. We identified specific genes (ZEB1 and HDAC9) that may underlie differences in body

size between mainland and Newfoundland lynx. The hypermethylated HDAC9 gene is of

particular interest because downregulation in HDAC9 is associated with diminished body

mass[31]. Commonly, insular mammalian carnivores are smaller in size[22], which is consistent with

existing data on the insular population of Canada lynx[17,18].

An overlooked marker for examining rapid evolution and adaptive divergence

Our results indicate that DNA methylation can detect population divergence despite substantial

gene flow, and can be a powerful marker to examine cryptic population structure in species that

seem genetically panmictic. Identifying the molecular basis of adaptive divergence between

populations has relied on comparing allele frequencies or isolating outlier loci under

selection[32,33]. These results suggest that epigenetic modifications increase resolution when

defining population structure and are a useful marker for examining subtle differentiation. With

time, divergent methylation patterns should lead linked SNPs to act as if under directional

selection, and thus could be factored into the working model of speciation[2]. Divergent

methylation patterns also have implications for conservation. For example, seemingly

genetically homogenous individuals could be epigenetically adapted to local conditions[34], and

thus, greater care would be needed to select individuals for translocations[35]. These results

indicate that epigenetic modifications play a role in defining population structure, with a putative

relationship to adaptation.

For decades, the adaptive role of epigenetic modifications in evolution has been

controversial[36–38]. Our data provide evidence that epigenetic modifications are a mechanism

underlying rapid evolutionary change. The evolutionary history of Canada lynx on Newfoundland

is unclear, yet evidence suggests post-glacial colonization (10,000 years)[13]. The

disproportionate levels of differential methylation in the smaller Newfoundland lynx, particularly

over genes with morphological significance, suggest an epigenetic pathway for rapid

evolutionary responses to geographic isolation[36]. Insular populations of mammals generally

have notable phenotypic differentiation[22,39], and DNA methylation serves as a potential

mechanism driving insular phenotypic divergence due to its plasticity to environmental

conditions[40]. Collectively, our results suggest that epigenetic modifications like DNA methylation

are an overlooked molecular signature of population divergence and local adaptation, and offer

a putative mechanism underlying rapid evolutionary response.

Methods

Sample Acquisition and Reduced-Representation Bisulfite Sequencing (RRBS) Library

Preparation

Georeferenced Canada lynx tissue samples were collected from fur auction houses throughout

eastern North America from dried pelts (North American Fur Auctions, Fur Harvester's Auctions,

Inc.). Individuals from four geographic locations were chosen for this study, spanning the

longitudinal and latitudinal range of the species and consistent with previous research on the

Canada lynx system (Rueness et al 2003). Tissue was consistently taken from the same

morphological location from each adult-sized pelt. Genomic DNA was isolated using a

MagneSil® (Promega Corporation) Blood Genomic Max Yield System a JANUS® workstation

(PerkinElmer, Inc.) and quantified and standardized to 20 ng/µl with a Quant-iT PicoGreen® ds-

DNA assay using manufacturer's instructions (Thermo Fisher Scientific). We adapted an

existing reduced representation bisulfite sequencing library preparation workflow designed for

multiplexed high-throughput sequencing[41]. Genomic DNA (400 ng) for 95 Canada lynx samples

was digested with NsiI and AseI restriction enzymes overnight and subsequently ligated with

methylated adapters (Supplemental Table S1). An individual sample of completely non-

methylated lambda phage genomic DNA (200 ng; Sigma-Aldrich – D3654) with a unique

barcode was included to assess bisulfite conversion efficiency (Supplemental Table S7).

Barcoded samples were then combined into eight pools to ensure consistent reaction

environments for the entire library using a QIAquick PCR Purification Kit (Qiagen, Valencia, CA,

USA) following manufacturer's instructions and 10 µL of 3M NaAc was added to neutralize pH.

We then performed a SPRI size selection on each pool (0.8x volume ratio) with Agencourt®

AMPure® XP beads (Beckman Coulter, Inc.). Nicks between the 3' fragment overhang and the

5' non-phosphorylated adapter nucleotide were repaired with DNA polymerase I and bisulfite

conversion was performed on each pool using an EZ DNA Methylation-Lightning Kit™ (Zymo

Research) with a 20-minute desulphonation time.

Pools were amplified in three separate PCRs to mitigate stochastic differences in

amplification and were subsequently concentrated using a QIAquick PCR Purification Kit

(Qiagen, Valencia, CA, USA). The eight pools were quantified with Qubit 3.0® (Thermo Fisher

Scientific) and appropriate amounts were added to a final super-pool for equal weighting. A final

magnetic bead clean-up (0.8x volume ratio) was performed to remove any adapter dimer. We

checked final library concentration and fragment distribution with a Qubit® 3.0 (ThermoFisher

Scientific) and Hi-Sense Bioanalyzer 2100 chip (Agilent), respectively. Paired-end 125-bp

sequencing was performed on a single lane on an Illumina HiSeq 2500 platform at the Centre

for Applied Genomics at the Hospital for Sick Children (Toronto, Ontario, Canada). Raw

sequence data FastQ files are available on the Sequence Read Archive (ID Pending).

Bioinformatics – quality checks and SNP calling

We assessed sequencing success as well as removed adapter and low-quality reads via

FastQC[42] and Cutadapt[43] implemented in TrimGalore! v0.4.4[44]. Individuals were demultiplexed

using python scripts[41]. Paired-ends reads for Canada lynx samples were initially aligned to

several genomes (*Felis catus, Homo sapiens,* Lambda Phage) using Bismark[27] to assess

mapping efficiency and contamination (Supplemental Table S4). Paired-end reads for

downstream analyses were aligned to the domestic cat genome with Bismark, using relaxed

mapping parameters (score_min L,0,-0.6)[27].

SNPs were called from indexed BAM files with CGmapTools[28] using a coupled Bayesian

wildcard algorithm with a conservative 0.01 error rate and a static 0.001 p-value for calling

variant sites, which generated variant call files (VCFs). VCF files were indexed, merged, and

filtered using VCFtools[45] for bi-allelic loci with a sequencing depth of at least five, and were

shared between at least 50% of the individuals (max-missing 0.5) and a minor allele frequency

(maf) of > 0.001. Sites were further filtered by removing any variants out of Hardy-Weinberg

equilibrium within populations (p-value < 0.05). A pair-wise Euclidean dissimilarity (distance)

matrix was computed on the SNP data using the function *daisy* within the package *cluster*[46]

using R v3.4.2[47]. This dissimilarity matrix was then summarized in a principal coordinates

analysis (PCoA) using the *dudi.pco* function in *adegenet*[48]. Missing SNP data was imputed by

mean allele at a population level. Pairwise F_{ST} was calculated using StAMPP (Supplemental

Figure S2) and relative F_{ST} was measured using a null model Bayesian approach implemented

in BayeScan v2.0[30] (Supplemental Figure S2). AMOVA between populations was computed

using the *poppr.amova* function within *poppr*[49], and heterozygosity was assessed with

adegenet.

Bioinformatics – DNA methylation

We identified methylated and non-methylated positions by first filtering BAM files for incomplete

bisulfite conversion based on reads containing more than three methylated positions in a CHH

or CHG context. In the remaining reads, methylated positions in a CpG context with a

sequencing depth of at least five were extracted with Bismark[27], truncating the last two bases of

the forward mate-paired reads (R1) and the first two bases of the reverse mate-paired (R2)

reads. Methylation polymorphisms in areas of overlap between read pairs were extracted only

once. After confirming that our non-directional library contained roughly equal reads for all

possible amplified DNA strands, we proceeded to analysis.

We first generated a custom CpG island annotation track using hidden Markov models

based on $CpG_{o/e}$ implemented in *makeCGI*[50]. Only islands with a calculated posterior probability

greater than 99.5% were retained for analysis, based on $CpG_{o/e}$. Mapped and extracted

methylated sites were then imported into Seqmonk[51] using the generic text importer, and raw

data was qualitatively visualized against the annotated domestic cat genome (felCat9.0). We

analyzed DNA methylation over CpG islands and gene bodies by creating 5,000-bp running

windows directly over and 25,000-bp upstream of gene bodies, combined with windows directly

over CpG islands. Each window was assigned a methylated percentage score based on the

overall ratio of methylated to non-methylated bases within the feature. We filtered this window-

set for regions that had at least one CpG and equal representation from each population. This

process was repeated for our second subset of analyses that investigated methylation patterns

of unannotated regions of the genome. For this analysis, we created 5,000-bp running windows

across the entire genome and removed any windows that overlapped with the initial windows

over CpG islands and gene bodies by more than 1%. Data was filtered, exported, and

summarized in the same way as the first analysis.

Sensitivity analyses

To determine the implications of missing data and PCoA axis retention thresholds, we

performed two sensitivity analyses. The first analysis examined the effects of missing data by

repeating all analyses using a subset of the top 10 individuals per population (N = 40), and

again with the top 6 individuals per population (N = 24), across all three datasets (Supplemental

Figure S5-S6, Table S8). Overall trends in explanatory effects (adjusted R^2) and qualitative

inferences (PCoA clustering) were investigated and no change in inferences were determined.

We examined the implications of arbitrary PCoA axis retention by repeating all analyses, but

instead using different cumulative variation explained thresholds as response variables. We

performed a number of db-RDAs using all axes explaining 30%, 50%, 75%, and 95%

cumulative variation as response variables, and results were qualitatively similar regardless of

axis retention (Supplemental Figure S8, Table S10).

Quantifying environmental associations

To determine if patterns of DNA methylation and genomic variability could be explained by

macro-scale climatic conditions, geographic distance, or insular divergence, we performed a

distanced-based redundancy analysis on the summarized SNP and DNA methylation data.

Meaningful axes explaining > 30% of the cumulative variation in the data were used as

response variables in a distance-based redundancy analysis (db-RDA) conducted in *vegan*[52],

using variables that putatively describe the environmental determinants of population structure

in Canada lynx. Our covariates included a binary variable of insularity, which identified the

Newfoundland population against mainland populations and was used to describe the largely

impassable barrier of the Strait of Belle Isle between Newfoundland and mainland Labrador[53]. A

variable of geographic distance was included which was simply the first axis of a principal

coordinates analysis (PCoA) on a Euclidean distance matrix of latitude and longitude (PCo1 =

99.7% of the variation). In addition to the geographic variables, we included a biotic variable of

percent tree cover[54], a randomly generated numerical variable to assess the effect of noise, and

a climate variable. For the climate variable, we performed a PCA on climate data to prevent

multi-collinearity, which reduced annual temperature ranges, winter precipitation, and minimum

coldest temperature to a single PCA axis[26] (PC1 = 85.6% of the variation). Linearity was

confirmed between response and explanatory variables, and multi-collinearity between

explanatory variables was assessed using the VIF and any variables > 4 were removed

(Supplemental Table S12). Step-wise model selection using the function *ordistep* within *vegan*[52]

was performed to isolate the best overall model using a QR decomposition technique based on

p-values (Supplemental Table S11). To isolate the individual explanatory power of each

variable, we performed partial distanced-based redundancy analyses (p-dbRDAs) on the

variables that were identified as significant in the full db-RDA.

Differentially methylated regions and gene ontology

We identified differentially methylated regions with functional biological correlates by performing

beta-regressions on windows over CpG islands and gene bodies for all 95 individuals with

percent methylation as the response variable and population as the explanatory variable. Beta

regressions are appropriate for proportion or percentage data[55], and we set an alpha threshold

at conservative levels seen in similar studies[34] (p-value < 0.001). Overall differences in percent

methylation were calculated, and genes were identified as either hypermethylated or

hypomethylated based on their relative degree of methylation compared to other populations.

Direct overlap between our differentially methylated regions and the felcat9.0 gene annotations

were extracted using Seqmonk[51]. We then identified functional associations by searching

UniProt (https://www.uniprot.org) by gene name against the database of genes in the domestic

cat genome using the search term "organism: *'Felis catus* (Cat) (*Felis silvestris catus*) (9685)'".

Acknowledgements

The authors thank Niels C.A.M. Wagemaker for his assistance with library preparation and his

generous donation of methylated adapters. We would also like to thank Felix Krueger and

Sibelle Torres Vilaça for their assistance with bioinformatics. We would like to thank Joe M.

Northrup (OMNR), Elizabeth M. Kirepka, Lynne E. Beaty, Angela R. Eads, and Melanie R.

Boudreau for their invaluable statistical advice and friendly manuscript reviews. We thank the

Natural Resources DNA Profiling and Forensic Centre (NRDPFC) for their assistance with

extractions and laboratory equipment, and the Centre for Applied Genomics with sequencing

troubleshooting. This work was funded by the CFI-JELF (358-47; A.B.A.S.), ComputeCanada

(GME-665-01; A.B.A.S.), NSERC Discovery Grants (D.L.M., A.B.A.S.) and CREATE-enviro

(J.B.J.).

Author Contributions

All authors contributed equally to this work.

Competing Interests

The authors declare no competing interests with this work.

Materials and Correspondence

Inquiries can be directed to the corresponding author, J.B.J.

References

1.   Watson, J. & Crick, F. Molecular structure of nucleic acids. *Nature* **171,** 737–8 (1953).

2.   Feder, J. L., Egan, S. P. & Nosil, P. The genomics of speciation-with-gene-flow. *Trends*

360   *Genet.* **28,** 342–350 (2012).

361   3.   Shafer, A. B. A. & Wolf, J. B. W. Widespread evidence for incipient ecological speciation:

362   A meta-analysis of isolation-by-ecology. *Ecol. Lett.* **16,** 940–950 (2013).

363   4.   Jones, P. A. & Takai, D. The role of DNA methylation in mammalian epigenetics. *Science*

364   *(80-.).* **293,** 1068–1070 (2001).

365   5.   Dubin, M. J. *et al.* DNA methylation in Arabidopsis has a genetic basis and shows

366   evidence of local adaptation. *Elife* **4,** 1–23 (2015).

367   6.   Artemov, A. V. *et al.* Genome-Wide DNA Methylation Profiling Reveals Epigenetic

368   Adaptation of Stickleback to Marine and Freshwater Conditions. *Mol. Biol. Evol.* **34,**

369   2203–2213 (2017).

7.   Fujita, N. *et al.* Methyl-CpG binding domain 1 (MBD1) interacts with the Suv39h1-HP1

heterochromatic complex for DNA methylation-based transcriptional repression. *J. Biol.*

Chem. **278,** 24132–24138 (2003).

8.   Lorincz, M. C., Dickerson, D. R., Schmitt, M. & Groudine, M. Intragenic DNA methylation

alters chromatin structure and elongation efficiency in mammalian cells. *Nat. Struct. Mol.*

Biol. **11,** 1068–1075 (2004).

9.   Maunakea, A. K. *et al.* Conserved role of intragenic DNA methylation in regulating

alternative promoters. *Nature* **466,** 253–257 (2010).

10.   Wutz, A. & P. Barlow, D. Imprinting of the mouse Igf2r gene depends on an intronic CpG

island. *Mol. Cell. Endocrinol.* **140,** 9–14 (1998).

11.   Han, L., Su, B., Li, W.-H. & Zhao, Z. CpG island density and its correlations with genomic

features in mammalian genomes. *Genome Biol.* **9,** R79 (2008).

12.   Jones, P. A. Functions of DNA methylation: Islands, start sites, gene bodies and beyond.

Nat. Rev. Genet. **13,** 484–492 (2012).

13.   Row, J. R. *et al.* Dispersal promotes high gene flow among Canada lynx populations

across mainland North America. *Conserv. Genet.* **13,** 1259–1268 (2012).

14.   Prentice, M. B. *et al.* Selection and drift influence genetic differentiation of insular Canada

lynx (Lynx canadensis) on Newfoundland and Cape Breton Island. *Ecol. Evol.* **7,** 3281–

3294 (2017).

15.   Stenseth, N. C. *et al.* Common Dynamic Structure of Canada Lynx Populations Within

Three Climatic Regions. (1999).

16.   Row, J. R. *et al.* The subtle role of climate change on population genetic structure in

Canada lynx. *Glob. Chang. Biol.* **20,** 2076–2086 (2014).

17.   Van Zyll De Jong, C. G. Differentiation of the Canada lynx, Felis (Lynx) canadensis

subsolana, in Newfoundland. *Can. J. Zool.* **53,** 699–705 (1975).

18.   Khidas, K., Duhaime, J. & Huynh, H. M. Morphological Divergence of Continental and

Island Populations of Canada Lynx. *Northeast. Nat.* **20,** 587–608 (2013).

19.   Foster, J. B. Evolution of mammals on Islands. *Nature* **202,** 234–235 (1964).

20.   Van Valen, L. Pattern and the balance of nature. *Evol. Theory* **1,** 31–49 (1973).

21.   Wayne, R. K. *et al.* A morphologic and genetic study of the Island fox, Urocyon littoralis.

Evolution (N. Y). **45,** 1849–1869 (1991).

22.   Lomolino, M. V. Body size evolution in insular vertebrates: Generality of the island rule. *J.*

Biogeogr. **32,** 1683–1699 (2005).

23.   Flatscher, R., Frajman, B., Schönswetter, P. & Paun, O. Environmental Heterogeneity

and Phenotypic Divergence: Can Heritable Epigenetic Variation Aid Speciation? *Genet.*

Res. Int. **2012,** 1–9 (2012).

24.   Stenseth, N. C. *et al.* Snow conditions may create an invisible barrier for lynx. *Proc. Natl.*

Acad. Sci. U. S. A. **101,** 10632–4 (2004).

25.   Keller, T. E., Lasky, J. R. & Yi, S. V. The multivariate association between genomewide

DNA methylation and climate across the range of Arabidopsis thaliana. *Mol. Ecol.* **25,**

1823–1837 (2016).

26.   Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. *Int. J. Climatol.* **37,** 4302–4315 (2017).

27.   Krueger, F. & Andrews, S. R. Bismark: A flexible aligner and methylation caller for

Bisulfite-Seq applications. *Bioinformatics* **27,** 1571–1572 (2011).

28.   Guo, W. *et al.* CGmapTools improves the precision of heterozygous SNV calls and

supports allele-specific methylation detection and visualization in bisulfite-sequencing

data. *Bioinformatics* **34,** 381–387 (2017).

29.   Rueness, E. K. *et al.* Ecological and genetic spatial structuring in the Canadian lynx.

Nature **425,** 69–72 (2003).

30.   Foll, M. & Gaggiotti, O. A genome-scan method to identify selected loci appropriate for

both dominant and codominant markers: A Bayesian perspective. *Genetics* **180,** 977–993

(2008).

31.   Chatterjee, T. K. *et al.* HDAC9 knockout mice are protected from adipose tissue

dysfunction and systemic metabolic disease during high-fat feeding. *Diabetes* **63,** 176–

187 (2014).

32.   Beaumont, M. A. & Balding, D. J. Identifying adaptive genetic divergence among

populations from genome scans. *Mol. Ecol.* **13,** 969–980 (2004).

33.   Jones, F. C. *et al.* The genomic basis of adaptive evolution in threespine sticklebacks.

Nature **484,** 55–61 (2012).

34.   Le Luyer, J. *et al.* Parallel epigenetic modifications induced by hatchery rearing in a

Pacific salmon. *Proc. Natl. Acad. Sci.* **114,** 12964–12969 (2017).

35.   Murray, D. L., Steury, T. D. & Roth, J. D. Assessment of Canada Lynx Research and

Conservation Needs in the Southern Range: Another Kick at the Cat. *J. Wildl. Manage.*

72, 1463–1472 (2008).

36.   Ho, M. W. & Saunders, P. T. Beyond neo-Darwinism-an epigenetic approach to evolution.

J. Theor. Biol. **78,** 573–591 (1979).

37.   Jablonka, E. & Raz, G. Transgenerational epigenetic inheritance: prevalence,

mechanisms, and implications for the study of heredity and evolution. *Q. Rev. Biol.* **84,**

131–76 (2009).

38.   Robertson, M. & Richards, C. Opportunities and challenges of next-generation

sequencing applications in ecological epigenetics. *Mol. Ecol.* **24,** 3799–3801 (2015).

39.   Meiri, S., Dayan, T. & Simberloff, D. Area, isolation and body size evolution in insular

carnivores. *Ecol. Lett.* **8,** 1211–1217 (2005).

40.   Lea, A. J., Altmann, J., Alberts, S. C. & Tung, J. Resource base influences genome-wide

DNA methylation levels in wild baboons (Papio cynocephalus). *Mol. Ecol.* **25,** 1681–1696

(2016).

41.   van Gurp, T. P. *et al.* epiGBS: reference-free reduced representation bisulfite

sequencing. *Nat. Methods* **13,** 322–4 (2016).

42.   Andrews, S. FastQC: A quality control tool for high throughput sequence data. Available

at: https://www.bioinformatics.babraham. (2010).

43.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17,** 10 (2011).

44.   Krueger, F. Trim Galore! : A wrapper tool around Cutadapt and FastQC to consistently

apply quality and adapter trimming to FastQ files. Available at:

https://www.bioinformatics.babraham. (2012).

45.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27,** 2156–2158

(2011).

46.   Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M. & Hornik, K. cluster: Cluster Analysis

Basics and Extensions. R package version 2.0.7-1 (2018).

47.   R Core Team. R: A language and environment for statistical computing. R Foundation for

461   Statistical Computing, Vienna, Au (2017).

462   48.   Jombart, T. Adegenet: A R package for the multivariate analysis of genetic markers.

463   *Bioinformatics* **24,** 1403–1405 (2008).

464   49.   Kamvar, Z. N., Tabima, J. F. & Grünwald, N. J. *Poppr*: an R package for genetic analysis

465   of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2,** e281

466   (2014).

467   50.   Wu, H., Caffo, B., Jaffee, H. A., Irizarry, R. A. & Feinberg, A. P. Redefining CpG islands

468   using hidden Markov models. *Biostatistics* **11,** 499–514 (2010).

469   51.   Andrews, S. Seqmonk: A tool to visualise and analyse high throughput mapped sequence

470   data. Available at https://www.bioinformatics.babraham.a (2007).

471   52.   Dixon, P. VEGAN, a package of R functions for community ecology. *J. Veg. Sci.* **14,** 927–

472   930 (2003).

473   53.   Koen, E. L., Bowman, J. & Wilson, P. J. Isolation of peripheral populations of Canada

474   lynx (*Lynx canadensis*). *Can. J. Zool.* **93,** 521–530 (2015).

475   54.   DeFries, R. S., Hansen, M. C., Townshend, J. R. G., Janetos, A. C. & Loveland, T. R. A

476   new global 1-km dataset of percentage tree cover derived from remote sensing. *Glob.*

477   *Chang. Biol.* **6,** 247–254 (2000).

478   55.   Ferrari, S. L. P. & Cribari-Neto, F. Beta regression for modelling rates and proportions. *J.*

479   *Appl. Stat.* **31,** 799–815 (2004).

480   56.   Weber, K. L., Fischer, R. S. & Fowler, V. M. Tmod3 regulates polarized epithelial cell

481   morphology. *J. Cell Sci.* **120,** 3625–3632 (2007).

482   57.   Saykally, J. N., Dogan, S., Cleary, M. P. & Sanders, M. M. The ZEB1 transcription factor

483   is a novel repressor of adiposity in female mice. *PLoS One* **4,** (2009).

Table 1. List of differentially methylated regions that have direct overlap with annotated
genes. Differentially methylated regions were identified using beta regressions with a p-
488   value < 0.001 and when methylation levels differed by more than 10% between the
identified population and the others. Hyper- and hypo-methylation is relative to the other
489   three populations. GO annotated functions were retrieved from UniProt (www.uniprot.org).

Population	Gene	UniProt ID	P-Value	% Difference	Hyper/Hypo-Methylated	GO Annotated Function
Alaska	ADGRL2	M3XFT6_FELCA	7.58E-07	20.17	Hyper	Carbohydrate binding, G-protein coupled receptor activity
	ANK2	M3WXH8_FELCA	2.41E-05	16.10	Hyper	Spectrin binding, maintenance of cytoskeletal structure
	ATP8A1	M3WM63_FELCA	0.000701	14.99	Hyper	ATP binding, magnesium ion binding
Manitoba	NRG3		0.000216	25.43	Hypo	No Uniprot Annotation
	RWDD1	M3XFU2_FELCA	1.86E-05	17.61	Hyper	Cytoplasmic translation
Newfoundland	CDH18	M3WJ78_FELCA	2.83E-05	21.23	Hypo	Calcium ion binding
	CENPU	M3W3F1_FELCA	0.000895	13.71	Hyper	Embryonic development
	CRISPLD1	M3W2H3_FELCA	0.000337	19.82	Hyper	Face morphogenesis
	DCC	M3WHY3_FELCA	0.000399	37.36	Hyper	Spinal cord ventral commissure morphogenesis
	FAM35A	M3WZ20_FELCA	0.000641	19.62	Hyper	No Uniprot Annotation
	HDAC9	A0A2I2U9G1_FELCA	4.40E-05	39.60	Hyper	Histone deacetylase; transcription regulation; downregulation associated with diminished body mass, adaptive thermogenesis[31]
	LOC101091724		0.000698	26.54	Hypo	lncRNA
	LOC109493917		0.000332	38.44	Hypo	lncRNA
	PBX3	M3WKS0_FELCA	3.21E-05	26.60	Hypo	DNA binding transcription factor activity
	TMOD2	M3X1Q9_FELCA	0.000847	29.43	Hyper	Myofibril assembly, muscle contraction, epithelial cell morphology[56]
	ZEB1	A0A2I2UAU6_FELCA	5.73E-05	37.60	Hyper	DNA-binding, transcription factor, regulation of adipose tissue mass[57]

494 **Figure 1.** Distribution of 95 Canada lynx (*Lynx canadensis*) samples across North America, used for high-throughput bisulfite sequencing. All four populations are
495 delineated by colour and include 24 individuals, except Alaska (n = 23).

Figure 2. Principal coordinates analysis (PCoA) plots of variation across three molecular marker datasets, with individuals as single circles and populations delineated by colour. All molecular data was summarized with a pair-wise Euclidean dissimilarity matrix. Methylation was summarized with 5,000-bp running windows over CpG islands and gene bodies (n = 329) and over unannotated regions (n = 376). SNP variants were called from bisulfite converted reads and reflect unstructured mainland populations (n = 496).

Figure 3. Distance-based redundancy analysis (db-RDA) on DNA methylation data over CpG islands and gene bodies, with population delineated by colour. The axes of a principal coordinates analysis (PCoA) were used a response variable to determine biogeographical relationships. The explanatory variables included a distance variable (the first axis of a PCoA on latitude and longitude); insularity (a binary variable distinguishing the Newfoundland island population); and climate (the first axis of a PCA summarizing winter precipitation, annual temperature ranges, and coldest minimum temperature).

Figure 4. Visual depiction of partial distance-based redundancy analyses (p-db-RDAs). The effect sizes indicate the independent explanatory effects of each variable on explaining methylation patterns, subtracted from the effect of any other variable. The effect size is adjusted R^2, (adj. R^2) and the test-statistic is a pseudo-F generated using QR decomposition within *vegan*.