**A legacy role for DNA binding of Lon protects against genotoxic stress**

Rilee D. Zeinert[1#], Justyne L. Ogdahl[1], Jing Liu[1], Qiyuan Yang[2], Yunguang Du[2], Benjamin B.

Barros[1], Peter L. Freddolino[3], Cole M. Haynes[2], Peter Chien[1*]

**Affiliations:**

[1]Department of Biochemistry and Molecular Biology, Molecular and Cellular Biology Program,

University of Massachusetts Amherst, Amherst MA

[2]Department of Molecular, Cell and Cancer Biology.

University of Massachusetts Medical School, Worcester MA

[3]Department of Biological Chemistry and Department of Computational Medicine &

Bioinformatics, University of Michigan, Ann Arbor, MI 48109, USA

#current address: National Institute of Child Health and Human Development, 49 Convent

Drive, Bethesda, MD 20892-4417

* pchien@umass.edu

**Abstract**

DNA binding proteins are essential for cellular life, but persistently bound complexes have

toxic consequences. Here we show that the proteotoxic responsive bacterial protease Lon

clears proteins from DNA to promote genotoxic stress resistance. Purified Lon binds DNA and

degrades neighboring bound proteins, while a fully active DNA-blind Lon variant does not.

This variant can degrade substrates as normal during unstressed growth, complements

pleotropic phenotypes of Δ*lon*, including proteotoxic resilience, but remains sensitive to

genotoxic stresses and fails to degrade proteins efficiently during DNA damage. Transposon

sequencing reveals that Δ*lon* is vulnerable to loss of protein-DNA eviction factors and we use

dynamic nucleoid occupancy profiling to show that chromosome-wide protein turnover relies

on Lon DNA binding. Finally, disrupting Lon binding to mitochondria genomes also results in

genotoxic stress sensitivity, consistent with the bacterial ancestry of this organelle. We

propose that clearance of persistent proteins from DNA by Lon originated in free-living $\alpha$-

proteobacteria and maintained during the evolution of mitochondria.

**Summary**

DNA binding by the Lon protease protects against genotoxic damage in a manner preserved

from bacteria to mitochondria.

**Introduction**

Maintenance, replication, and readout of genomes requires DNA-binding proteins, which perform their functions with specific dynamics. However, proteins that dwell on DNA beyond their normal lifetimes can result in toxic consequences such as collisions of DNA polymerase with these roadblocks. Given that many proteins are known to be associated with the DNA, cells must have dedicated protein eviction pathways to prevent DNA-Protein roadblocks. In eukaryotes, chromatin remodeling factors evict nucleosomes and recent discoveries of the Sprtn proteases that remove covalently bound protein adducts from DNA represent the most extreme version of this process (Reinking et al., 2020). In bacteria, chromatin is condensed into the nucleoid which is highly occupied with both specific binding proteins (e.g., transcription factors) and less specific factors (e.g. HU, H-NS, etc). Like in eukaryotes, clearance of these proteins is needed, however, what factors collaborate to control this process in bacteria is less understood.

In bacteria, the AAA+ protease Lon degrades misfolded proteins (Goldberg, 1972; Gur and Sauer, 2008) and seminal work in *E. coli* revealed that loss of Lon sensitized cells to genotoxic stresses (Mizusawa and Gottesman, 1983; Witkin, 1946). Lon also plays a role during normal growth. For example, in the alpha-proteobacteria *Caulobacter crescentus (Caulobacter)*, the Lon protease degrades the epigenetic regulator CcrM (Wright et al., 1996), the transcriptional regulator SciP (Gora et al., 2013), and the replication initiator DnaA (Jonas et al., 2013), in addition to its role in protein quality control (Lu et al., 2007). However, whether Lon plays a role in genotoxic tolerance in bacteria other than *E. coli* remains poorly understood. In mitochondria, which have an alphaproteobacteria origin (Gray, 2012), Lon has been clearly shown to manage the response to protein misfolding but it is unclear if it regulates mtDNA stress responses (Bota and Davies, 2002; Lu et al., 2007; Nargund et al., 2012).

Intriguingly, early biochemical work with Lon shows that it can bind DNA (Charette et al., 1984;

Chung and Goldberg, 1982; Liu et al., 2004; Lu et al., 2007) and more recent work identified

specific residues important for DNA binding (Karlowicz et al., 2017), but the biological

consequences of this binding remains unclear.

Here we report that Lon binding to DNA is important for genotoxic stress responses but

not proteotoxic stress responses. Using transposon sequencing, we find that factors important

for evicting persistent protein complexes from DNA show synthetic fitness defects with loss of

Lon DNA binding. In vitro, purified Lon degrades neighboring proteins when bound to the

same DNA region and genome-wide proteome occupancy shows that Lon binding to DNA

affects the lifetime of proteins on the chromosome. Finally, we show the importance of Lon

DNA binding in genotoxic stress tolerance is preserved in mitochondria, highlighting the link

between these organelles and their α-proteobacteria ancestor. Overall, our work supports a

model where Lon degrades lingering neighbor proteins when bound to chromosomal DNA, a

process that is especially important during the response to genotoxic stresses.

**Results**

*DNA binding of Lon is important for genotoxic stress tolerance*

We generated strains expressing only a DNA-blind variant of Lon (Lon4E) by mutating

residues implicated in DNA binding based on homology to the *E. coli* protein (Karlowicz et al.,

2017). Cells lacking Lon exhibit profound morphological defects including filamentation, loss of

motility in soft agar (Wright et al., 1996), and sensitivity to proteotoxic stresses such as

canavanine (Zeinert et al., 2020) (Figure 1A). Expression of Lon4E largely rescues these

phenotypes (Figure 1A,C; Figure S1). Unlike WT Lon, purified Lon4E fails to bind DNA (Figure

1B) but is completely active for degradation of model unstructured substrates such as casein

(Figure S1). *In vivo*, loss of Lon results in increased levels of its substrates DnaA and CcrM

(Jonas et al., 2013; Wright et al., 1996) and Lon4E restores steady-state levels to essentially

wildtype levels (Figure S1).

Interestingly, although they are resistant to proteotoxic stresses, Lon4E strains remain

sensitive to DNA damaging stresses, most particularly genotoxic agents that yield complex

damages such as mitomycin C (Figure 1A). Consistent with this, the prolific filamentation upon

DNA damage associated with Δlon strains is also seen with Lon4E but not WT strains (Figure

1C). Our general interpretation is that the Lon4E allele is fully capable of complementing

wildtype Lon activity during normal and proteotoxic stress but fails to fully protect against

genotoxic stress.

*Loss of protein eviction factors is synthetically sick with loss of Lon activity*

We had previously used transposon sequencing to identify genes that are differentially

critical in a Lon-dependent manner (Figure 2A; (Zeinert et al., 2020)). Given the extreme

sensitivity of Lon-deficient strains to genotoxic agents (Figure 1), we hypothesized that many

members of the DNA damage response (DDR) pathway would be essential in Δlon. However,

we found that most genes in the DDR were unaffected, including the sentinel protein RecA,

translesion polymerase (*imuB*) and mitomycin C specific responders (*mmcA*) (Figure 2 A,B),

which we confirmed by generating viable double mutants (Figure 2 C).

Only two DDR genes passed our filter as more essential in a Δlon background (Figure 2

A,B). The first of these is *uvrD*, a helicase originally known for its role in DNA damage repair

(Ogawa et al., 1968). UvrD can evict proteins bound to DNA, such as disassembling RecA

nucleofilaments (Petrova et al., 2015; Veaute et al., 2005) and stalled RNAP complexes

(Epshtein et al., 2014). While we could delete *uvrD* from wildtype cells, we could not delete *uvrD*

from Δ*lon* cells and depletion of Lon in a Δ*uvrD* strain was lethal (Figure 2C). Similarly, we found

*xseB*, a subunit of exonuclease VII that is important for reactivation of stalled replication forks

and mismatch repair (Lovett, 2011) also results in synthetic sickness when Lon is depleted

(Figure 2C; Figure S2).

One pathway where UvrD and ExoVII converge is in clearing protein roadblocks from DNA

(by UvrD) before they cause stalling of replication forks that must be restarted (initiated by

XseB). Interestingly, while Lon4E strains were equally fit as wildtype strains under normal

conditions, loss of UvrD in a Lon4E background compromised growth compared to the parental

Δ*uvrD* strain (Figure 2D). Taken together, we hypothesize that Lon binding to DNA aids in the

clearance of bound proteins, a function of particular importance during response to DNA

damage. We therefore looked at how protein dynamics *in vivo* could be affected by Lon DNA

binding.

*Lon binding to DNA affects condition dependent protein degradation*

Degradation of known Lon substrates, such as DnaA and CcrM, during nondamaging

conditions was the same in Lon4E strains as wildtype (Figure 3A). However, when MMC was

added to induce DNA damage, wildtype cells maintained robust DnaA turnover, while in

Lon4E, DnaA is degraded at ~50% of the wildtype rate. This stabilization did not extend to a

different Lon substrate, CcrM (Figure 3A), nor was this effect seen with the ClpXP substrate

CtrA (Jenal and Fuchs, 1998; Joshi et al., 2015)  (Figure S3).

*In vitro*, purified Lon degrades DnaA poorly on its own consistent with prior work showing

that unfolded substrates activate Lon degradation of DnaA (Jonas et al., 2013). Interestingly,

addition of a dsDNA with canonical DnaA binding sites results in robust degradation of DnaA

by Lon (Figure 3C). This effect required both an intact DnaA binding motif and a sufficiently

long adjacent region for Lon to bind as well (Figure 3C). Consistent with a requirement for

DNA binding by Lon for this activity, purified Lon4E does not degrade DnaA as well as

wildtype in the presence of this dsDNA scaffold (Figure S3). We conclude that Lon binding to

DNA is needed for robust degradation of neighboring substrates, such as DnaA, an effect

particularly important during genotoxic stress.

*Locus specific protein occupancy of the chromosome is regulated by DNA binding of Lon*

Our working model is that DNA binding AAA+ proteases like Lon can clear persistent

protein complexes. If this is true, then Lon mutants deficient in DNA binding should show

persistent protein occupancy on the chromosome. To explore this, we used a modification of

IPOD, a technique where fragmented crosslinked protein-DNA is purified and sequenced to

reveal protein occupancy of the bacteria chromosome (Freddolino et al., 2021). We monitored

chromosome occupancy in wildtype and Lon4E strains following translation inhibition, to reveal

loci where DNA-bound proteins are removed by AAA+ proteases.

Our dynamic IPOD approach revealed many regions of the chromosome that become

less occupied after translational inhibition in a wildtype strain - many of these regions

remained occupied in a Lon4E strain after similar treatment (Figure 3D). Correlation analyses

show that the occupancies of pre-inhibition wildtype, pre-inhibition Lon4E, and post-inhibition

Lon4E were more like each other compared to the occupancy seen in the wildtype post-

inhibition sample (Figure 3C). Closer examination of occupancy data suggests a role for Lon

clearing gene regulatory regions, such as those present in the promoter regions of *ccrM* and

CCNA_02622, a putative amidase located adjacent to *ftsZ* (Figure 3D). Taken together, these

data suggest that regions of the genome are bound with proteins that are cleared by Lon when

it binds to DNA.

*DNA binding properties of Lon are important for genotoxic stress responses in mitochondria*

Mitochondria have an endosymbiotic origin (Margulis, 1970) and the current model is that a free-living alpha-proteobacterium was engulfed by the ancestor of the modern eukaryotic cell (Gray, 2012). Because *Caulobacter* is an $\alpha$-proteobacteria, we explored if the need for DNA binding of Lon in tolerating DNA damage was preserved in mitochondria. In *C. elegans,* mitochondrial Lon (LONP-1) is required for development and growth (Nargund et al., 2012). Based on homology, we engineered a variant of *lonp-1* with mutations in the corresponding predicted DNA binding residues to generate LONP-1[4E] and used CRISPR/Cas9 to produce worms expressing only this allele at the endogenous locus. Importantly, we found that worms expressing only the LONP-1[4E] variant at endogenous levels were healthy, grew and developed normally (Figure 4A,D). Because LONP-1 has a known role in mitochondrial protein quality control by recognizing and degrading diverse proteins that incur oxidative damage (Bota and Davies, 2002), we infer that LONP-1[4E] is capable of maintaining proteostasis during animal development. Pull-down experiments indicated that LONP-1 robustly binds mitochondrial genomes *in vivo* as previously suggested (Fu et al., 1997; Lu et al., 2007), but LONP-1[4E] does not (Figure 4B). Therefore, DNA binding of mitochondria Lon is not essential for normal development or growth.

The mitochondrial unfolded protein response (UPR[mt]) couples the state of mitochondrial protein homeostasis to an upregulation of nuclear encoded chaperone genes. Induction of the UPR[mt] requires the transcription factor ATFS-1, which is targeted to the nucleus when mitochondria undergo stress. Complete depletion of LONP-1 activates the UPR[mt], as determined by upregulation of the transcriptional mitochondrial chaperone reporter *hsp-6pr::gfp* (Yoneda et al., 2004) (Figure 4C) consistent with a role for LONP-1 in the maintenance of mitochondrial function (Nargund et al., 2012). LONP-1[4E] worms do not activate the UPR[mt]

suggesting mitochondrial protein homeostasis is intact (Figure 4D). However, LONP-1[4E]

worms exposed to cisplatin showed a dramatic activation of UPR[mt] that is ATFS-1 dependent

while wildtype worms show no response (Figure 4D). Importantly, mitochondrial dysfunction

driven by depletion of the protein quality control protease SPG-7 caused UPR[mt] activation in

both wildtype and LONP-1[4E] animals to the same degree (Figure 4E), as does exposure to

paraquat which induces mitochondrial oxidative stress (Figure 4E). These results show that,

similar to what is seen in *Caulobacter*, loss of DNA binding of Lon in mitochondria results in a

specific intolerance to genotoxic stress.

**Discussion**

The role of DNA binding in Lon has been enigmatic since its initial observation (Chung

and Goldberg, 1982; Zehnbauer et al., 1981). Our working model is that Lon binding to DNA

allows it to recognize neighboring proteins and serves as an additional layer of quality control

to reduce the residence time of persistent DNA binding complexes (Figure 4F). This oversight

works in concert with other protein eviction factors such as UvrD and seems particularly

important during DNA damaging conditions (Figure 4F). Using a genome wide approach, we

find evidence for clearance of protein complexes during normal growth that relies on Lon

binding to DNA. Finally, we have shown that the same properties of DNA binding in Lon that

are important for genotoxic stress responses in *Caulobacter* are also important in

mitochondria, consistent with the endosymbiotic $\alpha$-proteobacteria ancestry of this organelle.

Overall, our working model is that Lon plays a canonical role in proteotoxic stress

response by degrading misfolded or damaged proteins. This function occurs regardless of

DNA binding. However, DNA binding of Lon also allows for local control of neighboring

proteins that is particularly important during genotoxic stress conditions. Our working model is

that Lon clearance of persistently bound proteins prevents the accumulation of protein

roadblocks that limit replication or transcription. This is particularly important in bacteria and

bacterially derived organelles, as replication, transcription and translation occur concurrently in

the same compartment. Because Lon recognizes substrates based on general chemical

properties as opposed to highly specific sequence motifs, Lon would be ideally suited to serve

in this capacity as the identity of the protein roadblock will likely depend on the specific

circumstance that generated it.

**Figure Legends**

**Figure 1**. **A**. Serial dilutions (10-fold dilutions left to right) of wildtype (WT), Lon deficient

($\Delta lon$), or DNA binding deficient Lon expressing (Lon4E) strains on media containing various

concentrations of either proteotoxic (L-canavanine (L-can)) or genotoxic (mitomycin C (MMC),

cisplatin (Cis), methyl methanosulfone (MMS), 4-nitroquinolone (4NQO)) compounds. **B.** Lon

binding to fluorescently labeled DNA oligo as measured by change in fluorescence

polarization. Data was fit to a hyperbolic binding model, $K_D$= 286 nM (95% CI: 90-1200 nM).

Concentration is reported as hexamers of Lon. inset: polarization measurement of labeled

DNA in the presence of wildtype Lon or Lon4E. **C.** Morphology of cells during normal growth

and after damaging with mitomycin C. Scalebar showing 5 μm in first panel, all other panels

are the same magnification.

# Figure 1

**Figure 2. A.** Transposon-sequencing reveals genes that differentially affect fitness in wildtype

or Δlon strains (Zeinert et al., 2020), highlighting genes annotated as DNA damage repair

genes (red).  **B.** Snapshots of transposon insertion sites for DNA damage repair genes in

either wildtype (WT) or Δlon strains.  **C.** above: Induction (+) or depletion (-) of Lon in either

wildtype, ΔuvrD, or ΔexoVII strains, showing synthetic sickness.  below: Deletions of DNA

damage repair genes *recA*, *imuB*, or *mmcA* can be generated in Δlon strains without loss of

viability and as expected are generally important for response to MMC.  D.  Lon4E strains

grow as well as wildtype based on competition, but deletion of *uvrD* reduces fitness in a Lon4E

background.

**Figure 2**

**Figure 3.  A.**  Degradation of DnaA and CcrM in wildtype, Lon4E or Δlon strains as monitored

by translational shutoff experiments either in nondamaging or DNA damaging (+Mitomycin C)

conditions.  **B.**  Cartoon illustrating dsDNA oligonucleotides used to test for scaffolding.  **C.**

Degradation of purified DnaA by Lon in the presence of different dsDNA scaffolds.  **D.**

Spearman correlation matrix of genome occupancy in wildtype (WT) or strains expressing

Lon4E (Lone4E) as monitored by IPOD (Freddolino et al., 2021) either before (Pre) or two

hours after (Post) translational shutoff.  Degree of correlation is shown by color and inset

values.  **E.**  Representative tracks showing changes in protein occupancy at *ftsZ* and *ccrM*

neighboring regions in either WT or Lon4E strains.

# Figure 3

**Figure 4.** DNA binding mutants of mitochondria Lon. **A**. Immunoblots of lysates from WT

worms, or strains expressing LONP-1-FLAG or LONP-1$^{4E}$-FLAG. Tubulin serves as a loading

control.  **B**. ChIP of mtDNA with FLAG antibody in WT, LONP-1-FLAG and LONP-1$^{4E}$-FLAG

strains. **C.**  Cartoon illustrating hsp-6 reporter of mitochondria stress. **D.** Photomicrographs of

*hsp-6$_{pr}$::gfp* WT or LONP-1$^{4E}$ worms raised on DMSO or cisplatin, scale bar, 0.5 mm.  **E.**

Photomicrographs of *hsp-6$_{pr}$::gfp* WT or LONP-1$^{4E}$ worms raised on paraquat (PQ) or *spg-*

*7*(RNAi). **F.** Model for how Lon binding to DNA clears persistent proteins (gray) together with

other evicting enzymes such as UvrD (red).

# Figure 4

A

FLAG

Tubulin

WT    lonp-1::FLAG    lonp-1$^{4E}$::FLAG

100kD

25kD

B    amount of DNA from α-FLAG IP
(normalized to WT)

n=3

lonp-1::FLAG

lonp-1$^{4E}$::FLAG

WT

C

normal mitochondria → stressed mitochondria

ATFS-1

hsp-6

UPRmt

D

| 0 µg/ml Cis | 1 µg/ml Cis | 10 µg/ml Cis | 10 µg/ml Cis |

atfs-1 (RNAi)

WT    WT    WT    WT

atfs-1 (RNAi)

LONP-1$^{4E}$    LONP-1$^{4E}$    LONP-1$^{4E}$    LONP-1$^{4E}$

hsp-6$_{pr}$::gfp

E

| control | 0.3 mM PQ | spg-7 (RNAi) |

WT    WT    WT

LONP-1$^{4E}$    LONP-1$^{4E}$    LONP-1$^{4E}$

hsp-6$_{pr}$::gfp

F

displacing
enzymes
(UvrD)

DNA
Dmg

Lon

removal    degradation

Damage Response

**Acknowledgements:**

We thank Lucy Shapiro for supplying the anti-CcrM antibody and members of the Chien lab

and the Protein Homeostasis theme of the Center for Models to Medicine for helpful feedback.

**Funding:** This work was supported by HHMI (C.M.H.), the Mallinckrodt Foundation (C.M.H)

and National Institutes of Health grants (R01AG040061) to C.M.H. and (R35GM130320) to

P.C.  R.Z. and J.L. were funded in part by the UMass Chemistry Biology Interface Training

Program (NIH T32GM139789).  J.O. was funded in part by the UMass Biotechnology training

program (NIH T32GM135096). **Author contributions:**  P.C., J.L., and R.Z. conceptualized the

study.  B.B., J.L., J.O., R.Z, Q.Y., and Y.D. performed experiments.  P.C., C.H., R.Z, and J.L

wrote the initial draft.  All authors reviewed the final draft.  **Data and materials availability**:

All data is available in the main text or the supplementary materials.

**Materials and Methods**

Bacteria strains and media

*C. crescentus* was grown on peptone yeast extract (PYE) at 30°C. 1.5% agar was used for

solid media. Carbon sources were supplemented at final concentrations of 0.2% glucose or

xylose. Antibiotics were used at the following concentrations for solid media growth:

Kanamycin 25 $\mu$g/ml, Spectinomycin 100 $\mu$g/ml, Tetracycline 1.5 $\mu$g/ml and Gentamycin 25

$\mu$g/ml. For growth of *E. coli* in solid or liquid media, we used lysogeny broth (LB) at 37°C.

Antibiotics were used at the following concentrations: Chloramphenicol 30 $\mu$g/ml, Kanamycin

50 $\mu$g/ml, Tetracycline 15 $\mu$g/ml and Gentamycin 50 $\mu$g/ml. After initial selection steps for

strain construction, antibiotic selection was excluded from all physiological related studies.

Strain construction

Deletion strains were constructed using the two-step recombination procedure with counter

selection by sucrose/*sacB*, using pNPTS138 plasmids containing deletion cassettes

comprised of 1000 bp upstream and downstream of the coding regions flanking a gentamycin

resistance cassette. Following transformation, initial primary selection was on PYE with

Kanamycin 25 $\mu$g/ml. Primary colonies were grown overnight in PYE without selection to allow

for secondary recombination. Overnight cultures were plated onto solid media supplemented

with 3% (w/v) sucrose and 25 $\mu$g/ml Gentamycin. To validate gene deletion, resulting strains

were tested for antibiotic sensitivity to Kanamycin and resistance to Gentamycin.  Phage

transduction of marked alleles was performed using PhiCr30.

Allelic replacement of Lon4E was performed by transforming pNPTS138-Lon4E into CPC546

(lon::specR) with primary selection on PYE with Kanamycin 25 $\mu$g/ml and secondary selection

on PYE + 3% (w/v) sucrose. The resulting strains were genotyped for antibiotic sensitivity to

Kanamycin and Spectinomycin. Clones harboring the correct genotype were then confirmed

by PCR amplification of the *lon* locus, Sanger sequencing, and whole genome resequencing of

the strain.

Protein Purification and *in vitro* degradation assays

Lon, Lon4E, and DnaA were purified with protocols as previously described (Jonas et al.,

2013) using hydroxyapatite resin (Sigma) instead of P11 cellulose for Lon/Lon4E.

FITC casein degradation was performed as previously described (Vieux et al., 2013). Briefly,

degradation assays (30°C) were as follows unless noted otherwise: Lon/Lon4E hexamer 0.15

$\mu$M, 10 $\mu$g/ml FITC casein type II (Sigma), ATP 4 mM, 75 $\mu$g/ml creatine kinase, 5mM creatine

phosphate.  Final concentrations of components in degradation assays (30°C) were as follows

unless noted otherwise: Lon/Lon4E 0.15 $\mu$M (hexamer concentration), DnaA 2.5 $\mu$M, DNA 2

$\mu$M, ATP 4 mM, 75 $\mu$g/ml creatine kinase, 5 mM creatine phosphate.

Fluorescent polarization studies

ssDNA (AACGGATGATCCACAGGAGAGTCTGGCGCAGGGCGAGAG) labeled with

fluorescein (FAM) was ordered from Integrated DNA Technologies. Wildtype Lon was titrated

in two-fold dilutions starting from 1.25 $\mu$M (all concentrations in hexamer).  For inset in Figure

1, 300 nM of Lon or Lon4E hexamer was incubated with 200nM FAM-DNA in 20mM HEPES,

10mM $MgCl_2$, 100mM KCl with 0.05% Tween. Polarization measurements were read from 40

uL of these mixtures using opaque non-binding black 384-well plates (Corning) and a

SpectraMax M5 plate reader (Molecular Devices), with excitation and emission wavelengths

set at 460 nm and 540 nm respectively.

<u>Bacteria Characterization: Motility, Morphology, Viability and Stress Assays</u>

For motility assays, single colonies were picked from PYE agar and inoculated into 0.3% PYE

agar. Plates were grown at 30°C for 2 days before imaging.

Microscopy images are phase contrast images of exponentially growing *Caulobacter* cells

$OD_{600} \approx 0.4$ (Zeiss AXIO ScopeA1) mounted on 1% PYE agar pads using a 100X objective.

Caulobacter strains were cultured in PYE medium overnight. Cultures were diluted to $OD_{600}$

0.05 and grown to $OD_{600} \approx 0.4$, after which 10-fold serial dilutions were spotted onto regular

PYE agar or agarose (5-Azacytidine) plates containing the indicated concentrations of

Mitomycin C, Cisplatin, L-canavanine, Methyl methanosulfonate, and 4-Nitroquinoline N-oxide

(Sigma) followed by incubation at 30°C for 72 hours prior to imaging. Mitomycin C, L-

canavanine, and Cisplatin was prepared at a stock concentration of 0.4 mg/ml, 100 mg/ml, and

20 mM respectively in water followed by filter sterilization. 4-NQO was prepared at a saturated

solution in acetonitrile.

For synthetic lethality experiments with Lon depletion, xylose was present in all cultures at a

final concentration of 0.2%. Overnight cultures were back diluted to $OD_{600}$ 0.05 and outgrown

to $OD_{600} \approx 0.4$. Cells were 10-fold serially diluted, plated onto media containing 0.2% glucose

or 0.2 xylose, grown for 72 hours at 30°C and imaged.

<u>Western Blot Analysis</u>

Strains were cultured in PYE medium overnight. Cultures were diluted to $OD_{600}$ 0.05 and

grown to $OD_{600} \approx 0.4$. For steady state or translation shutoffs (30 μg/ml chloramphenicol) cells

were normalized and loaded onto a 10% BIS Tris pH 6.8 gel. Samples were run at 150V for 60

mins. Gels were transferred to nitrocellulose membranes at 100V for 50 minutes. Following

transfer, blots were blocked with TBST with 5% milk for 45 minutes and probed with anti-rabbit

primary CcrM, DnaA, and ClpP at 1:5,000 dilution overnight at 4°C. Blots were washed three

times for 5 minutes in TBST. Followed by incubation with Goat anti-rabbit LiCOR 800CW

secondary for 1 hour. Blots were washed three times in TBST and imaged using a LiCOR

Odyssey Imager, quantification was performed using ImageJ (NIH).

<u>Worm strains and bacterial food sources</u>

Hermaphrodite worms were raised on the OP50 strain of *E. coli* unless treated with *atfs-*

*1*(RNAi), in which the HT115 *E. coli* strain harboring the *atfs-1*(RNAi) plasmid was used

(Haynes et al., 2010). The UPR$^{mt}$ reporter worm strain harboring the *hsp-6$_{pr}$::gfp* transgene

used in the *C. elegans* imaging experiments was described previously (Yoneda et al., 2004).

All new strains were confirmed by sequencing, and backcrossed three times prior to crossing

into the *hsp-6$_{pr}$::gfp* background.

*C. elegans* cisplatin experiments

Worm strains were synchronized by bleaching. The eggs were transferred to plates containing

the described concentration of cisplatin (Sigma), allowed to hatch and develop for 36 hours,

when they were harvested and imaged as described (Nargund et al., 2012).

<u>CRISPR/Cas9 genome editing</u>

Via CRISPR-Cas9, a nucleic acid sequence encoding the amino acids -

GS<u>DYKDDDDK</u>GS<u>DYKDDDDK</u>GS<u>DYKDDDDK</u>-, which yields 3xFLAG epitope sites

(underlined) interspersed by -GS- linkers, was inserted at the 3' end of the endogenous *lonp-1*

open reading frame at the native locus as described (Friedland et al., 2013; Katic et al., 2015;

Paix et al., 2017). crRNAs and donor templates were generated and purchased from

Integrated DNA Technologies.

**LONP-1::FLAG**

crRNA sequence:

/AlTR1/rArArGrArArArGrGrGrArCrArCrGrUrArUrUrArUrGrUrUrUrArGrArGrCrUrArUrGrCrU/

AlTR2/

Donor template:

ATATTCGATTCGTTTCGCATTATGATGAGCTCTACGAGCATCTCTTCCAAGGATCCGACTA

CAAGGACGATGACGACAAGGGATCTGACTACAAGGACGATGACGACAAGGGATCTGACT

ACAAGGACGATGACGACAAGTAATAATACGTGTCCCTTTCTTTCACCCCCAGTTAATACCT

TTCAGATAGATTTTG

**LONP-1$^{4E}$**

To generate LONP-1$^{4E}$

Amino acid residues 467 (Arginine), 468 (Arginine), 473 (Lysine) and 474 (Lysine) were all

altered to aspartic acid residues via CRISPR-Cas9 as described above.

crRNA sequence:

/AlTR1/rArArCrUrGrArArArGrArCrGrArCrGrArUrUrCrUrGrUrUrUrUrArGrArGrCrUrArUrGrCrU/

AlTR2/

Donor template:

CTACCTGGAATGGTTGACATCAGTACCCTGGGGACTTACATCACCCGAGAATGAGGAACT

TTCAGTTGCAGAAGAGGCTCTTGATGAAGGACATTATGGA

*C. elegans* western blots and mtDNA ChIP

Western blots and ChIP experiments were performed as described (*51*), with several

modifications. Synchronized worms were cultured in liquid and harvested at the L4 stage by

sucrose flotation. Worms were resuspended in cold PBS + protease inhibitors (Roche) and

homogenized. The homogenate was treated with 1.85% formaldehyde for 15 min to cross-link

DNA and protein. Glycine was then added to a final concentration of 125mM for 5 min at room

temperature to quench the formaldehyde. The pellets were washed twice in cold PBS +

protease inhibitors, centrifuged at 2500 x G. Samples were sonicated using a Bioruptor

(Diagenode) for 15 min at 4°C on high intensity (30s on, 30s off). Samples were transferred to

microfuge tubes and spun at 13,000 x G for 15 min at 4°C. The supernatants were precleared

with pre-blocked ChIP-grade Pierce™ magnetic protein A/G beads (Thermo Scientific) and

then incubated with Monoclonal ANTI-FLAG® M2 antibody (Sigma, F1804) or Mouse mAb

IgG1 Isotype Control (Cell Signaling Technology, G3A1) overnight at 4°C. The antibody-

chromatin complex was then concentrated with pre-blocked ChIP-grade Pierce™ magnetic

protein A/G beads. After washing, DNA-protein crosslinks were reversed by incubation 65°C

overnight followed by treatment with RNaseA at 37°C for 2 hours, and then proteinase K at

55°C for 2 hours. Finally, immunoprecipitated and input DNA were purified with ChIP DNA

Clean & Concentrator (Zymo Research, D5205) and used as templates for qPCR. Relative

enrichment folds of detected regions were determined after normalization to input and total

mtDNA primers as described (Nargund et al., 2015).

qPCR primers to quantify *C. elegans* mtDNA following ChIP:

5'-GCAGTCTTAGCGTGAGGACATTA -3'

5'- AAACATAAAACATAAATAGAACTAACCA-3';

Dynamic IPOD

*In vivo* protein occupancy (IPOD) was measured as described (Freddolino et al., 2021) in

wildtype or Lon4E *Caulobacter crescentus* at initial and two hours after addition of

chloramphenicol to shut off translation.  For Spearman correlation, occupancy scores were

averaged across overlapping 50 kb intervals across the genome and comparisons made

between timepoints and strains as shown.  Wig files for occupancy data are provided in

Supplemental information.

**REFERENCES**

Bota, D.A., and Davies, K.J.A. (2002). Lon protease preferentially degrades oxidized

mitochondrial aconitase by an ATP-stimulated mechanism. Nature Cell Biology *4*, 674–680.

Charette, M.F., Henderson, G.W., Doane, L.L., and Markovitz, A. (1984). DNA-stimulated

ATPase activity on the lon (CapR) protein. J Bacteriol *158*, 195–201.

Chung, C.H., and Goldberg, A.L. (1982). DNA stimulates ATP-dependent proteolysis and

protein-dependent ATPase activity of protease La from Escherichia coli. Proc Natl Acad Sci U

S A *79*, 795–799.

Epshtein, V., Kamarthapu, V., McGary, K., Svetlov, V., Ueberheide, B., Proshkin, S., Mironov,

A., and Nudler, E. (2014). UvrD facilitates DNA repair by pulling RNA polymerase backwards.

Nature *505*, 372–377.

Freddolino, P.L., Amemiya, H.M., Goss, T.J., and Tavazoie, S. (2021). Dynamic landscape of

protein occupancy across the Escherichia coli chromosome. PLoS Biol *19*, e3001306.

Friedland, A.E., Tzur, Y.B., Esvelt, K.M., Colaiácovo, M.P., Church, G.M., and Calarco, J.A.

(2013). Heritable genome editing in C. elegans via a CRISPR-Cas9 system. Nat Methods *10*,

741–743.

Fu, G.K., Smith, M.J., and Markovitz, D.M. (1997). Bacterial Protease Lon Is a Site-specific

DNA-binding Protein. Journal of Biological Chemistry *272*, 534–538.

Goldberg, A.L. (1972). Degradation of abnormal proteins in Escherichia coli (protein

breakdown-protein structure-mistranslation-amino acid analogs-puromycin). Proc Natl Acad

Sci U S A *69*, 422–426.

Gora, K.G., Cantin, A., Wohlever, M., Joshi, K.K., Perchuk, B.S., Chien, P., and Laub, M.T.

(2013). Regulated proteolysis of a transcription factor complex is critical to cell cycle

progression in Caulobacter crescentus. Mol Microbiol *87*, 1277–1289.

Gray, M.W. (2012). Mitochondrial evolution. Cold Spring Harb Perspect Biol *4*, a011403–

a011403.

Gur, E., and Sauer, R.T. (2008). Recognition of misfolded proteins by Lon, a AAA(+) protease.

Genes Dev *22*, 2267–2277.

Haynes, C.M., Yang, Y., Blais, S.P., Neubert, T.A., and Ron, D. (2010). The matrix peptide

exporter HAF-1 signals a mitochondrial UPR by activating the transcription factor ZC376.7 in

C. elegans. Mol Cell *37*, 529–540.

Jenal, U., and Fuchs, T. (1998). An essential protease involved in bacterial cell-cycle control.

EMBO J *17*, 5658–5669.

Jonas, K., Liu, J., Chien, P., and Laub, M.T. (2013). Proteotoxic stress induces a cell-cycle

arrest by stimulating Lon to degrade the replication initiator DnaA. Cell *154*, 623–636.

Joshi, K.K., Bergé, M., Radhakrishnan, S.K., Viollier, P.H., and Chien, P. (2015). An Adaptor

Hierarchy Regulates Proteolysis during a Bacterial Cell Cycle. Cell *163*, 419–431.

Karlowicz, A., Wegrzyn, K., Gross, M., Kaczynska, D., Ropelewska, M., Siemiątkowska, M.,

Bujnicki, J.M., and Konieczny, I. (2017). Defining the crucial domain and amino acid residues

in bacterial Lon protease for DNA binding and processing of DNA-interacting substrates. J Biol

Chem *292*, 7507–7518.

Katic, I., Xu, L., and Ciosk, R. (2015). CRISPR/Cas9 Genome Editing in Caenorhabditis

elegans: Evaluation of Templates for Homology-Mediated Repair and Knock-Ins by Homology-

Independent DNA Repair. G3 (Bethesda) *5*, 1649–1656.

Liu, T., Lu, B., Lee, I., Ondrovičová, G., Kutejová, E., and Suzuki, C.K. (2004). DNA and RNA

Binding by the Mitochondrial Lon Protease Is Regulated by Nucleotide and Protein Substrate.

Journal of Biological Chemistry *279*, 13902–13910.

Lovett, S.T. (2011). The DNA Exonucleases of Escherichia coli. EcoSal Plus *4*.

Lu, B., Yadav, S., Shah, P.G., Liu, T., Tian, B., Pukszta, S., Villaluna, N., Kutejová, E.,

Newlon, C.S., Santos, J.H., et al. (2007). Roles for the Human ATP-dependent Lon Protease

in Mitochondrial DNA Maintenance. Journal of Biological Chemistry *282*, 17363–17374.

Margulis, L. (1970). Origin of eukaryotic cells: evidence and research implications for a theory

of the origin and evolution of microbial, plant, and animal cells on the Precambrian earth (New

Haven: Yale University Press).

Mizusawa, S., and Gottesman, S. (1983). Protein degradation in Escherichia coli: the lon gene

controls the stability of sulA protein. Proc Natl Acad Sci U S A *80*, 358–362.

Nargund, A.M., Pellegrino, M.W., Fiorese, C.J., Baker, B.M., and Haynes, C.M. (2012).

Mitochondrial import efficiency of ATFS-1 regulates mitochondrial UPR activation. Science

*337*, 587–590.

Nargund, A.M., Fiorese, C.J., Pellegrino, M.W., Deng, P., and Haynes, C.M. (2015).

Mitochondrial and nuclear accumulation of the transcription factor ATFS-1 promotes OXPHOS

recovery during the UPR(mt). Mol Cell *58*, 123–133.

Ogawa, H., Shimada, K., and Tomizawa, J. (1968). Studies on radiation-sensitive mutants of

E. coli: I. Mutants defective in the repair synthesis. Molec. Gen. Genet. *101*, 227–244.

Paix, A., Folkmann, A., and Seydoux, G. (2017). Precision genome editing using CRISPR-

Cas9 and linear repair templates in C. elegans. Methods *121–122*, 86–93.

Petrova, V., Chen, S.H., Molzberger, E.T., Tomko, E., Chitteni-Pattu, S., Jia, H., Ordabayev,

Y., Lohman, T.M., and Cox, M.M. (2015). Active displacement of RecA filaments by UvrD

translocase activity. Nucleic Acids Res *43*, 4133–4149.

Reinking, H.K., Hofmann, K., and Stingele, J. (2020). Function and evolution of the DNA-

protein crosslink proteases Wss1 and SPRTN. DNA Repair (Amst.) *88*, 102822.

Veaute, X., Delmas, S., Selva, M., Jeusset, J., Le Cam, E., Matic, I., Fabre, F., and Petit, M.-A.

(2005). UvrD helicase, unlike Rep helicase, dismantles RecA nucleoprotein filaments in

Escherichia coli. EMBO J *24*, 180–189.

Vieux, E.F., Wohlever, M.L., Chen, J.Z., Sauer, R.T., and Baker, T.A. (2013). Distinct

quaternary structures of the AAA+ Lon protease control substrate degradation. Proc Natl Acad

Sci U S A *110*, E2002–E2008.

Witkin, E.M. (1946). Inherited Differences in Sensitivity to Radiation in Escherichia Coli. Proc

Natl Acad Sci U S A *32*, 59–68.

Wright, R., Stephens, C., Zweiger, G., Shapiro, L., and Alley, M.R. (1996). Caulobacter Lon

protease has a critical role in cell-cycle control of DNA methylation. Genes & Development *10*,

1532–1542.

Yoneda, T., Benedetti, C., Urano, F., Clark, S.G., Harding, H.P., and Ron, D. (2004).

Compartment-specific perturbation of protein handling activates genes encoding mitochondrial

chaperones. J. Cell Sci. *117*, 4055.

Zehnbauer, B.A., Foley, E.C., Henderson, G.W., and Markovitz, A. (1981). Identification and

purification of the Lon+ (capR+) gene product, a DNA-binding protein. Proc Natl Acad Sci U S

A *78*, 2043–2047.

Zeinert, R.D., Baniasadi, H., Tu, B.P., and Chien, P. (2020). The Lon Protease Links

Nucleotide Metabolism with Proteotoxic Stress. Mol. Cell.

SUPPLEMENTAL INFORMATION

**Supplemental Figure Legends.**

**Figure S1**. Supporting main Figure 1.  **A.** Three biological replicates of wildtype, Δlon and

Lon4E strains were individually grown to exponential phase. Cell length and stalk were

measured for each.  Cell lengths were determined automatically using MicrobeJ, Stalk lengths

were measured manually using ImageJ (NIH).  Individual measurements are shown as well as

a notched box plot for these values.  **B.** Growth in 0.3% Agar PYE media of WT, Δlon and

Lon4E strains on left.  Western blot of labeled proteins shown on right. **C.**  Degradation of

CcrM is restored in Lon4E.  Chloramphenicol was added to exponentially growing cells and

timepoints were taken at timepoints indicated.  Quantification was performed as described in

methods (n=3).  **D.** Purified Lon or Lon4E can degrade fluorescently labeled casein (FITC-

casein; Sigma) with similar efficiency.  Degradation of FITC-casein results in increased

fluorescence due to loss of self-quenching.  Lon binding to certain single-stranded DNA

oligonucleotides suppresses degradation of FITC-casein, but Lon4E is not responsive.  **E.**

Additional concentrations and replicates of growth under stress similar to main Figure 1.

# Figure S1

550 **Figure S2.** Supporting main Figure 2. Cells expressing a xylose inducible Lon as the sole copy

551 *(∆lon Pxyl:lon)* can tolerate loss of *uvrD* under inducing, but not repressing conditions.

552 Similarly, depletion of Lon in a *∆exoVII* strain also results in reduced fitness.  Loss of Lon

553 shows no synthetic effects with loss of *recA*, *imuB* or *mmcA*, although loss of these genes

554 dramatically increases sensitivity to DNA damage produced by mitomycin C.

**Figure S2**

557 **Figure S3.** Supporting main Figure 3. A. CtrA degradation in WT, Lon4E and Δlon strains with

558 and without mitomycin C damage. B. Degradation of DnaA by dsDNA is increased with WT

559 Lon, but not with Lon4E. n.b., DnaAR357A is used here as it is unable to hydrolyze ATP and

560 therefore is persistently locked in the ATP-bound / DNA-binding state.

# Figure S3

## A

| | no damage | + Mitomycin C |
|---|---|---|
| | 0  30  60  90 | 0  30  60  90 |

CtrA

WT
Lon4E
Δlon

## B

WT Lon     Lon4E

time        time

Lon/Lon4E
DnaA
R357A
Creatine
Kinase

- wt
- Lon4E

Time (mins)