Vascular brassinosteroid receptors confer drought resistance without penalizing plant growth

Norma Fàbregas[1,*], Fidel Lozano-Elena[1,*], David Blasco-Escámez[1], Takayuki Tohge[2,8], Cristina Martínez-Andújar[3], Alfonso Albacete[3], Sonia Osorio[4], Mariana Bustamante[1], José Luis Riechmann[1,9], Takahito Nomura[5], Takao Yokota[6], Ana Conesa[7] , Francisco Pérez Alfocea[3], Alisdair R. Fernie[2] and Ana I. Caño-Delgado[1, ¶]

[1]*Centre for Research in Agricultural Genomics (CRAG) CSIC-IRTA-UAB-UB, Barcelona 08193, Spain.*

[2]*Max Planck Institute of Molecular Plant Physiology, Potsdam-Golm D-14476, Germany.*

[3]*Department of Plant Nutrition, CEBAS-CSIC, Murcia 30100, Spain.*

[4]*Instituto de Hortofruticultura Subtropical y Mediterránea "La Mayora". University of Málaga-Consejo Superior de Investigaciones Científicas. Department of Molecular Biology and Biochemistry, Málaga 29071, Spain.*

[5]*Center for Bioscience Research and Education, Utsunomiya University, Minemachi, Utsunomiya 321-8505, Japan.*

[6]*Department of Biosciences, Teikyo University,Toyosatodai, Utsunomiya 320-8551, Japan.*

[7]*Microbiology and Cell Science Department, IFAS, Genetics Institute, University of Florida, Gainesville 32603, USA.*

[8]*Current Address: NAIST Graduate school of Biological Sciences, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan.*

25   [9]*Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, 08010,*

26   *Spain*

27   [*] Both authors contributed equally to this work.

28   [¶] Correspondence to: ana.cano@cragenomica.es

ABSTRACT

Drought represents a major threat to food security. Mechanistic data describing

plant responses to drought have been studied extensively and genes conferring

drought resistance have been introduced into crop plants. However, plants with

enhanced drought resistance usually display lower growth, highlighting the need

for strategies to uncouple drought resistance from growth. Here, we show that

overexpression of BRL3, a vascular-enriched member of the brassinosteroid

receptor family, can confer drought stress in Arabidopsis. Whereas loss-of-

function mutations in the ubiquitously expressed BRI1 receptor leads to drought

resistance at the expense of growth, overexpression of BRL3 receptor confers

drought tolerance without penalizing overall growth. Systematic analyses reveal

that upon drought stress, increased BRL3 triggers the accumulation of

osmoprotectant metabolites including proline and sugars. Transcriptomic

analysis suggests that this results from differential expression of genes in the

vascular tissues. Altogether, this data suggests that manipulating BRL3

expression could be used to engineer drought tolerant crops.

INTRODUCTION

Drought is responsible for at least 40% of crop losses worldwide and this proportion is dramatically increasing due to climate change [1]. Understanding cellular responses to drought stress represents the first step toward the development of better-adapted crops, something which is a great challenge for the field of plant biotechnology [2]. Classical approaches aimed at examining how plants cope with limited water led to the identification of regulators involved in the signal transduction cascades of the abscisic acid (ABA)-dependent and ABA-independent pathways [3]. Adaptation to drought stress has been associated with the presence of proteins that protect cells from dehydration, such as late-embryogenesis–abundant (LEA) proteins, osmoprotectants and detoxification enzymes [4,5]. These studies provided deep insights into the molecular mechanisms underlying abiotic stress [2], showing that drought resistance is a complex trait simultaneously controlled by many genes. While genetic approaches have succeeded in conferring stress resistance to plants, this generally comes at the cost of reduced growth [6,7]. Therefore, understanding how cellular growth is coupled to drought stress responses is essential for engineering plants with improved growth in rainfed environments.

Receptor-like kinases (RLKs) play an important role in optimizing plant responses to stress [8,9]. Brassinosteroid (BR) hormones directly bind to BRI1 (BR-INSENSITIVE 1) leucine-rich repeat (LRR)-RLK family members on the plasma membrane [10-14]. Ligand perception triggers BRI1 to interact with the co-receptor BAK1 (BRI1 ASSOCIATED RECEPTOR KINASE 1) [15-17], which is essential for early BR signaling events [18]. This BRI1-BAK1 heterodimerization initiates a signaling cascade of phosphorylation events that control the

expression of multiple BR-regulated genes mainly via the BRI1-EMS-

SUPPRESSOR1 (BES1) and BRASSINAZOLE RESISTANT1 (BZR1)

transcription factors [19-21].

Although BRs modulate multiple developmental and environmental stress

responses in plants, the exact role of BRs under stress conditions remains

controversial. Whereas the exogenous application of BRs and the

overexpression of the BR biosynthetic enzyme DWF4 both confer increased

plant adaptation to drought stress [22-24], suppression of the BRI1 receptor also

results in drought-resistant phenotypes [25,26]. Intriguingly, ABA signaling inhibits

the BR signaling pathway after BR perception, and crosstalk between the two

pathways upstream of the BIN2 (BRASSINOSTEROID-INSENSITIVE 2) kinase

has been reported [27,28]. Further crosstalk has been described downstream

mediated by the overlapping transcriptional control of multiple BR- and ABA-

regulated genes [29,30] such as *RESPONSE TO DESICCATION 26* (*RD26*) [26].

Recently, greater attention is being placed on the spatial regulation of hormonal

signaling pathways in attempt to further understand the coordination of plant

growth and stress responses [26,31-34]. For instance, while the BRI1 receptor is

widely localized in many tissues [35], the BRI1-LIKE receptor homologues BRL1

and BRL3 signal from the innermost tissues of the plant and thereby contribute

to vascular development [12,33,36]. BR receptor complexes are formed by different

combinations of BRI1-like LRR-RLKs with the BAK1 co-receptor in the plasma

membrane [33]. Despite BRI1 being a central player in plant growth and

adaptation to abiotic stress [26,37,38], the functional relevance of vascular BRL1 and

BRL3 is only just beginning to be explored [33,39]. For example, in previous

proteomic approaches we found abiotic stress-related proteins within BRL3

signalosome complexes [33], but the exact role of the BRL3 pathway in drought

remains elusive.

Here, we show that the different BR receptors modulate a plant´s adaptive

response to multiple drought stress traits in both the roots and shoots. While the

traits controlled by the BRI1 pathway are intimately linked to growth arrest, we

found that increased levels of vascular-enriched BRL3 receptors confer drought

resistance without penalizing overall plant growth. Moreover, metabolite

profiling revealed that the overexpression of the BRL3 receptor triggers the

production of an osmoprotectant signature (i.e., proline, trehalose, sucrose, and

raffinose family oligosaccharides) in the plant and the specific accumulation of

the osmoprotectant metabolites in the roots during periods of drought.

Subsequent transcriptomic profiling showed that this metabolite signature is

transcriptionally regulated by the BRL3 pathway in response to drought. An

enrichment of deregulated genes in root vascular tissues, especially in the

phloem, further supports a preferential accumulation of osmoprotectant

metabolites to the root. Overall, this study demonstrates that overexpression of

the BRL3 receptor boosts the accumulation of sugar and osmoprotectant

metabolites in the root to overcome drought-associated growth arrest, thereby

uncovering a strategy to protect crops against drought.

RESULTS

BR receptors control osmotic stress sensitivity in the root

To determine the contribution of the BR complexes in the response to drought,

we performed a comprehensive characterization of different combinations of

mutants of all the BR receptors and the BAK1 co-receptor. For each

combination, we first analyzed primary root growth (Fig. 1a). As previously

described [17,33,40], seven-day-old roots of *bak1*, *brl1brl3bak1*, *bri1*, and

bri1brl1brl3 displayed shorter roots than the Col-0 wild type (WT). We also

found that the primary roots of the quadruple mutant *bri1brl1brl3bak1* (hereafter

quad) were the shortest and the most insensitive to BRs (Fig. 1a,b and

Supplementary Fig. 1). Conversely, plants overexpressing BRL3 (*35S:BRL3-*

GFP, hereafter *BRL3ox*) not only exhibited longer roots than WT (Fig. 1a,b) but

also showed increased receptor levels in root vascular tissues [33]

(Supplementary Fig. 2). These results agree with the previously reported role of

BR receptors in promoting root growth [40,41]. We then subjected Arabidopsis

seedlings to osmotic stress by transferring them to sorbitol-containing media

and subsequently quantified the level of inhibition of root growth in sorbitol

relative to control conditions (see Methods). A significantly lower level of relative

root growth inhibition mediated by osmotic stress was observed in *bri1* (27%),

bri1brl1brl3 (28%) and *quad* (27%) mutants compared to the WT (39%; Fig.

1a,b). In contrast, no differences were found in *brlbrl3* and *brl1brl3bak1* root

growth inhibition when compared to WT (Fig. 1a,b). Similarly, the roots of

BRL3ox plants were like those of WT in terms of relative root growth inhibition

(Fig. 1a,b).

Previous experimental evidences unveiled that the water stress-induced cell

death in Arabidopsis roots is localized and undergoes programmed cell death

(PCD) [42]. As shown by the incorporation of propidium iodine (PI) into the nuclei

(Fig. 1c,d), a short period of osmotic stress (24h) caused cell death in the

elongation zone of WT roots. In comparison, a reduced amount of cell death

was observed in the roots of *bri1, bri1brl1brl3* and *quad* mutants (Fig. 1c,d),

thereby indicating less sensitivity towards osmotic stress. Conversely, plants with increased levels of BRL3 receptor showed a massive amount of cell death in the root tips compared to the WT, indicating an increased sensitivity to short osmotic stress (Fig 1c,d). These results point towards a role for BR receptors in triggering osmotic stress responses in the plant root.

Since root hydrotropism represents a key feature for adaptation to environments scarce in water [43], we investigated the capacity of roots to escape imposed osmotic stress by bending towards water-available media (Fig. 2a). We found that BR receptors loss-of-function mutants had reduced hydrotropic responses compared to WT plants. For instance, while no significant differences were found under control conditions (mock) (Supplementary Fig. 3), the roots of BR receptor mutants grew straighter than WT roots towards the sorbitol-containing media (Fig. 2a-c). Interestingly, *brl1brl3bak1* mutants were the most insensitive to osmotic stress in terms of hydrotropism, showing lower root curvature angles than the *quad* roots (Fig. 2b). Consistently, compared to WT roots, an enhanced hydrotropic response was observed in *BRL3ox* (Fig. 2a-c). Furthermore, exogenous application of the BR synthesis inhibitor brassinazole [44] reverted the hydrotropic response of WT roots (Supplementary Fig. 3). For better visualization, we generated a drought multi-trait matrix for all the BR receptor mutants analyzed in this study (Fig. 2d; Supplementary Table 1). From this matrix, it can be seen that the BRL3/BRL1/BAK1 receptors in the vascular tissues enhances hydrotropic sensitivity.

BRL3ox confers drought resistance without penalizing growth

To investigate if the impaired responses to abiotic stress observed in root

seedling were preserved in mature plants, we next analyzed the phenotypes of

plants exposed to severe drought. After 12 days of withholding water, dramatic

symptoms of drought stress were observed in WT, *brl1brl3* and *brl1brl3bak1*

mutants. In contrast, other BR mutants showed a remarkable degree of drought

resistance. In particular, *bak1*, *bri1*, *bri1brl1brl3*, and *quad* mutant plants were

the most resistant to the severe water-withholding regime (Fig. 3a). As these

mutants exhibited some degree of dwarfism (Fig. 3a), we confirmed their

resistance to drought by examining their survival rates after re-watering (Fig.

3b). To correct for the delayed growth seen in BR-deficient mutants, plants were

submitted to a time course of drought stress in which water use, photosynthesis

and transpiration parameters were monitored under similar relative soil water

content (Fig. 3c-e). The WT plants took just 9 days to use 70% of the available

water (field capacity) during the drought period (Fig. 3c). In comparison, BR

loss-of-function mutant plants *bri1*, *bri1brl1brl3* and *quad* took 15 days. All

subsequent measurements were done at the same soil water content for each

genotype. We found that the relative water content (RWC) in WT plants was

reduced during drought, while RWC in BR mutant leaves remained as in well-

watered conditions (Fig. 3d). In addition, compared to WT plants, BR mutants

sustained higher levels of photosynthesis and transpiration during the drought

period (Fig. 3e and Supplementary Fig. 4). Altogether our results indicate that

the dwarf BR receptor mutant plants are more resistant while consuming less

water, likely through avoiding the effects of drought (Fig. 3f).

Strikingly, we found that *BRL3ox* plants were more resistant than WT plants to

severe drought stress as shown by increased survival rates (Fig. 3a,b). Plants

with increased BRL3 receptors showed reduction of RWC during drought

similarly to WT plants (Fig. 3d). Interestingly the rate of photosynthesis was

lower in *BRL3ox* compared to WT at basal conditions, but together with

transpiration, was more stable than in WT plants during the drought period (Fig.

3,e and Supplementary Fig. 4). This indicates that *BRL3ox* plants are healthier

than WT under the same water consumption conditions. These results suggest

that the BRL3 pathway actively promotes drought tolerance without penalizing

plant growth (Fig. 3f).

BRL3 triggers the production of osmoprotectant metabolites

To further investigate the cause behind BRL3 receptor-conferred drought

tolerance, we performed metabolite profiling of *BRL3ox* plants and compared it

to the profile of WT and *quad* plants in a time course drought experiment. Roots

were separated from shoots to address possible changes in metabolite

accumulation from source to sink tissues. The complete metabolic fingerprints

are provided in Supplementary Fig. 5 and 6 and Supplementary Data 1 and 2.

Metabolite profiling of mature *BRL3ox* plants grown in control conditions (time

0) revealed an increment in the production of osmoprotectant metabolites. Both

shoots (Fig. 4a) and roots (Fig. 4b) of the *BRL3ox* plants exhibited metabolic

signatures enriched in proline and sugars, metabolites which have previously

been reported to confer resistance to drought [45-47]. This suggests that the BRL3

receptor promotes priming [48]. Importantly, the levels of these metabolites were

lower in *quad* mutant plants (Fig. 4a,c).

Compared to in WT, sugars including fructose, glucose, galactinol, galactose,

maltose, and raffinose were accumulated in the shoots of *BRL3ox* (Fig. 4a).

Conversely, whereas glucose levels were lower in the roots, sucrose, trehalose,

myo-inositol, and maltose appeared to accumulate here (Fig. 4b) suggesting

that the BRL3 pathway promotes sugar accumulation preferentially in the roots.

We then analyzed the dynamics of each metabolite in response to drought (see

Methods). In this time course, a rapid accumulation of osmoprotectant

metabolites was observed in *BRL3ox* plants (Fig. 4c,d). In the shoots, proline

showed the highest levels respect WT along the entire drought time course (Fig.

4c,f). In contrast, glucose, galactose and *myo*-inositol increased at similar or

slightly lower rates than in the shoots of WT plant (Fig. 4c,e,g). However, in

roots, an accumulation of trehalose, sucrose, proline, and raffinose was

observed in *BRL3ox* mutants subjected to drought stress (Fig. 4d), and this

accumulation showed steeper exponential dynamics than in WT plants (Fig.

4h). Additionally, glucose, galactose, fructose, and *myo*-inositol linearly

increased in WT roots but exponentially increased in *BRL3ox* roots (Fig. 4j).

Interestingly, throughout this time course, the levels of these metabolites were

lower in the *quad* mutant plants compared to in WT (Supplementary Fig. 7).

Altogether, these findings uncover a key role for BR receptors in promoting

sugar metabolism, and support the idea that BRL3 triggers the accumulation of

osmoprotectant metabolites in the root to promote growth during periods of

drought.

BRL3 transcriptionally controls the drought metabolic response

The accumulation of osmolytes in the roots of plants with increased levels of

BRL3 suggests faster sugar mobilization from shoots to roots. Thus, we next

investigated whether these metabolic pathways are transcriptionally regulated

by BRL3 in the root. RNAseq of *BRL3ox* roots revealed 759 deregulated genes

at basal conditions (214 upregulated and 545 downregulated; FC>1.5,

FDR<0.05; Supplementary Data 3) and 1,068 deregulated genes in drought

conditions (378 upregulated and 690 downregulated; FC>1.5, FDR<0.05;

Supplementary Data 4). In control conditions, a high proportion of the

deregulated genes belonged to the response to water stress, oxygen-containing

compounds (ROS) and response to ABA GO categories (Fig. 5a, 5c,

Supplementary Data 5 and 6). We next deployed the genes falling into the

response to stress category, which included classical drought stress markers

such as *RD22* and *RAB18* that were already upregulated in basal conditions

(Fig 5b). An enrichment of genes belonging to the response to hormone

category indicated altered hormonal responses in *BRL3ox* plants under drought

(Fig 5a,c and Supplementary Data 7 and 8). Further analyses of specific

hormonal responses revealed that the ABA and jasmonic acid (JA) were the

most altered responses (Supplementary Fig. 8). Repression of JA biosynthesis

genes resulted in decreased levels of JA in basal conditions (Supplementary

Fig. 8).

In order to uncover differential drought responses between WT and *BRL3ox*

roots, we constructed a linear model accounting for the interaction between

genotype and drought (Supplementary Data 9). Taking the 200 most

significantly affected genes, we grouped them in (i) genes more activated in

BRL3ox under drought compared to WT (Supplementary Data 10) and (ii)

genes more repressed in *BRL3ox* under drought compared to WT

(Supplementary Data 11). GO enrichment analysis of this genotype-drought

interaction revealed (i) secondary metabolism, response to stress and response

to water deprivation in the first group and (ii) brassinosteroid mediated signaling

pathway in the second group (Fig. 5d). Importantly, the expression levels of

dehydration response genes kept repressed in *quad* mutant plants during

drought (Supplementary Fig. 7 and Supplementary Data 12-15). The

expression levels of two key BR biosynthesis genes, *CPD* and *DWF4* were

analyzed by RT qPCR. Consistently, within the drought time course,

transcription levels of *CPD* and *DWF4* were increased in *quad* and reduced in

BRL3ox compared to WT plants. Quantification of the bioactive BR hormone

Castasterone (CS) showed similar trends and BL hormone was only detected in

quad, suggesting that BL is accumulated in *quad* more than in WT and *BRL3ox*

plants (Supplementary Fig. 9).

Analysis of transcription factors revealed 29 of them with differential responses

to drought between *BRL3ox* and WT roots. Interestingly, the drought-responsive

transcription factor *RD26* showed an enhanced response in *BRL3ox* roots

during stress, whereas several vascular-specific transcription factors remained

repressed under drought (Supplementary Fig. 10). Given that the BRL3

receptor is natively expressed at the phloem-pole pericycle and enriched in

vascular tissues when overexpressed [33], we analyzed the spatial distribution of

the derregulated genes within the root tissues in our RNAseq dataset [49]. The

deregulated genes were preferentially enriched in specific vascular tissues such

as the pericycle and phloem pole pericycle but also in lateral root primordia

(which initiates from pericycle) and root hair cells, (Fig. 6a, see Methods).

Interaction-affected genes concentrated in pericycle and phloem but also in

columella and cortex (Fig. 6b). Among the phloem-enriched genes, we found

two trehalose phosphatases (*TPPs*) and one galactinol synthases (*GolS2*) that

show increased expression in *BRL3ox* roots at basal conditions and in response

to drought (Fig. 6d). These enzymes drive the synthesis of the osmoprotectant

metabolites — trehalose, myo-inositol and raffinose — we found to be

accumulated in *BRL3ox* roots. Together, these results support the importance

of increasing BRL3 signaling in the phloem for sustaining drought resistance.

Furthermore, a statistical analysis revealed a significant link between the whole

transcriptomic and metabolomic signatures, both in basal conditions and under

drought (p=0.017 and p=0.001 respectively; see Methods). This indicates that

the metabolic signature of *BRL3ox* plants is transcriptionally controlled. We

used the metabolic and transcriptomic signatures to identify deregulated

metabolic pathways using Paintomics [50]. This analysis revealed a constitutive

control of sucrose metabolism by the vascular BRL3 pathway that was

enhanced during drought stress in *BRL3ox* plants. We also found that BRL3

controls galactose metabolism under periods of drought, including the raffinose

family of oligosaccharides (RFOs) synthesis pathway (Supplementary Fig. 11,

Supplementary Data 16 and 17). Collectively, these results support that the

BRL3 receptor promotes drought tolerance, mainly by controlling sugar

metabolism.

DISCUSSION

Our study addresses the importance of the BRL3 receptor in preventing growth

arrest during severe periods of drought. This is accomplished through the

transcriptional control of metabolic pathways that produce osmoprotectant

metabolites that are accumulated in the roots. While spatial BR signaling has

been shown to contribute to stem cell replenishment in response to genotoxic

stress [31,34], here we uncover a role for vascular-enriched BRL3 receptors in

promoting growth during drought. Altogether, our results highlight the

importance of the spatial regulation of BRL3 signaling in plant stress responses

and open a window for studying drought tolerance in plants at a cell-specific

resolution.

The exogenous application of BR compounds has been used widely in

agriculture to extend growth under different abiotic stresses [22,51], yet how these

molecules precisely activate growth in challenging conditions remains largely

unknown. The analysis of BR signaling and BR synthesis mutant plants

subjected to stress failed to provide a linear picture of the involvement of BR in

drought stress adaptation. For instance, although overexpression of the

canonical BRI1 pathway and the BR biosynthesis gene DWF4 can both confer

abiotic stress resistance [24,38], *BRI1* loss-of-function mutants also showed

drought stress resistance [2526]. However, increased levels of BR-regulated

transcription factors trigger antagonistic effects in drought stress responses [26,52],

thus depicting a complex scenario for the role of BRs in abiotic stress. Given the

spatiotemporal regulation of the BR signaling components [39] and the complexity

of drought traits [7], it is plausible to hypothesize that drought traits are under the

control of cell type-specific BR signaling.

Our study unveils that the BR family of receptors, in addition to promoting

growth, guides phenotypic adaptation responses to a myriad of drought stress

traits. The drought resistance phenotypes of BR loss-of-function mutants (Fig.

3a) are likely caused by a reduced exposure of these plants to the effect of

drought. This phenomenon, known as drought avoidance, is linked to growth

arrest and stress insensitivity that maintains transpiration, leaf water status and

photosynthesis along the drought (Fig. 3 and Supplementary Fig. 4). The

reduced levels of ABA and canonical stress-related metabolites, together with

the downregulation of stress genes, further support the insensitivity of *quad*

plants to stress (Supplementary Figs. 7 and 8).

In contrast, the phenotypes observed in *BRL3ox* plants indicate an active

drought tolerance mechanism driven by the BRL3 receptor. First, *BRL3ox* roots

showed induced water stress-induced PCD in the root tip compared to WT (Fig.

1c,d), which has been proposed to modify the root system architecture and

thereby enhance drought tolerance [49]. Second, the enhanced hydrotropic

response of *BRL3ox* roots (Fig. 2a-c) could function during water-limited

conditions by modifying root architecture for increased acquisition of water,

favoring plant growth and survival under drought conditions as previously

described [53]. Third, at same RWC in leaves, the rate of photosynthesis and

transpiration were more stable in *BRL3ox* than in WT plants during drought (Fig.

3d,e and Supplementary Fig. 4). Altogether, these findings indicate that BRL3

actively promotes drought tolerance without penalizing plant growth.

We found the expression of the drought-response transcription factor *RD26* to

be enhanced in *BRL3ox* roots when subjected to drought (Supplementary Fig.

10). *RD26* has been shown to antagonize with the BR canonical transcription

factor BES1 [26], thereby suggesting that BRL3 activates alternative pathways.

These alternative pathways may be derived from a spatial specialization of BR

functions within the root. Indeed, we found enriched transcriptional responses

within the vascular tissues of *BRL3ox* roots, especially within phloem-related

cell types (Fig. 6a,b). The localization of the native BRL3 protein in phloem cells

[33] and the metabolic signature found in *BRL3ox* supports its implication in

phloem loading during drought. Moreover, metabolic enzymes implicated in trehalose and RFO metabolism were enriched in vascular tissues and either upregulated in *BRL3ox* roots in basal conditions or strongly responding to drought (Fig. 6c,d). Thus, BRL3 function not only loads/unloads the phloem, but also directly controls metabolic pathways. This is the case for the trehalose phosphate phosphatase family (*TPPs*) [54,55] and galactinol synthase 2 (*GolS2*) [56], which are both described to impact drought responses and are involved in trehalose and RFO synthesis respectively. In addition to controlling enzymes at the vascular level, our analyses also uncover a role for BRL3 in mediating non-vascular enzymes important for metabolism and drought responses. These enzymes include hexokinases such as *HXK3* or *HKL1*, the sucrose synthases *SUS3* and *SPS2F*, and proline dehydrogenase genes such as the early response to dehydration 5 (*ERD5)* which is involved in stress tolerance [57]. In light of our findings and given that *Bes1-D* gain-of-function mutants exhibit drought hypersensitivity [26], we propose that vascular BRL3 receptors may act independently of the canonical growth-promoting BRI1 pathway.

Our data further suggest that *BRL3ox* plants accumulate sugars in the sink tissues to enable plant roots to grow and escape drought by searching for water within the soil. In support of these findings, we also observed reduced levels of photosynthesis in well-watered leaves of *BRL3ox* plants (Fig. 3e). These results, together with the higher levels of sucrose in roots compared with in shoots (Fig. 4a) and higher levels of glucose and fructose in the shoots suggest that the BRL3 pathway promotes sugar mobilization from the leaves (source) to the roots (sink). In fact, previous work reported that BRs promote the flow of

assimilates in crops from source to sink via the vasculature [58] and via sucrose

phloem unloading [59].

In control conditions, *BRL3ox* plants exhibited a metabolic signature enriched in

proline and sugars. Proline and sugar accumulation classically correlates with

drought stress tolerance, osmolytes, ROS scavengers, and chaperone functions

[5,45-47,60,61], suggesting that the BRL3 receptor promotes priming [48,62]. In addition,

BRL3ox plants also accumulated succinate, fumarate and malate. Importantly,

all these metabolites were decreased in *quad* mutant plants. Altogether, these

data uncover a role for BRL3 signaling in the promotion of the tricarboxylic acid

(TCA) cycle, sugar and amino acid metabolism.

In drought stress conditions, *BRL3ox* shoots displayed increased levels of the

amino acids proline, GABA and tyrosine. In contrast, trehalose, sucrose, *myo*-

inositol, raffinose, and proline were the most abundant metabolites in the

BRL3ox roots along the stress time course. Importantly, all these metabolites

have previously been linked to drought resistance [45,46,60]. In addition, the levels

of the RFO metabolites raffinose and *myo*-inositol, which are involved in

membrane protection and radical scavenging [63], were higher in the roots of

BRL3ox plants under drought, yet reduced in the roots of *quad* plants.

Our data indicate that the roots of *BRL3ox* plants are loaded with

osmoprotectant metabolites and are thus better prepared to alleviate drought

stress via a phenomenon previously referred to as priming [48,62]. Further

supporting our hypothesis, the BRL3 signalosome is enriched in abiotic stress

proteins [33]. Altogether these data suggest that drought stress responses are

correlated with BRL3 receptor levels in the root vasculature, especially within

the phloem, and that this is key for the greater survival rates of *BRL3ox* plants.

Future cell type-specific engineering of signaling cascades stands out as a

promising strategy to circumvent growth arrest caused by drought stress.

METHODS

Plant materials

Seeds were sterilized with 35% NaClO for 5 min and washed five times for 5min

with sterile dH$_2$O. Sterile seeds were vernalized 48 h at 4ºC and grown in half-

strength agar Murashige and Skoog (MS1/2) media with vitamins and without

sucrose. Plates were grown vertically in long day (LD) conditions (16 h of light /

8 h of dark; 22°C, 60% relative humidity). Genotypes used in this study:

Columbia-0 WT (Col-0 WT), *brl1-1brl3-1* (*brl1brl3*), *bak1-3* (*bak1*), *bri1-301*

(*bri1*), *bri1-301brl1-1brl3-1* (*bri1brl1brl3*), *bri1-301bak1-3brl1-1brl3-1 (quad)* and

35S:BRL3-GFP (BRL3ox) [33]*.* DNA rapid extraction protocol [64] was used for all

the plant genotyping experiments. Supplementary Table 2 describes the

primers used for genotyping of the BR mutant plants.

Brassinolide and sorbitol sensitivity assays in roots

For hormone treatments, seeds were continuously grown in concentration

series of brassinolide (BL,Wako, Japan). For sorbitol assays, three-day-old

seedlings were transferred to either control or 270 mM sorbitol media for four

additional days. The root length of seven-day-old seedlings was measured

using Image J (http://rsb.info.nih.gov/ij/) and compared with automatically

acquired data from the MyROOT [65] software (Supplementary Fig. 13). Four-day-

old roots grown in control conditions or in 24 h of sorbitol were stained with

propidium iodide (10 ug/ml, PI, Sigma). PI stains the cell wall (control) and DNA

in the nuclei upon cell death (sorbitol). Images were acquired with a confocal microscope (FV1000 Olympus). Cell death damage in primary roots was measured in a window of 500 µm from QC in the middle root longitudinal section (Image J). As an arbitrary setting to measure the stained area, a color threshold ranging from 160 to 255 in brightness was selected.

Root hydrotropism

Seedlings were germinated in MS1/2 without sucrose for six days. Then, the lower part of the agar was removed from the plates and MS1/2 with 270 mM sorbitol was added to simulate a situation of reduced water availability. The media was placed in 45-degree angle to scape gravitropism effect. When indicated, 1 µM of brassinazole [44] was added to sorbitol media. Root curvature angles were measured and analyzed using the Image J software (http://rsb.info.nih.gov/ij/).

Drought stress for scoring plant survival

One-week-old seedlings grown in MS1/2 agar plates were transferred individually to pots containing 30±1 g of substrate (plus 1:8 v/v vermiculite and 1:8 v/v perlite). For each biological replicate, 40 plants of each genotype were grown in LD conditions for three weeks. Three-week-old plants were subjected to severe drought stress by withholding water for 12 days followed by re-watering. After the seven-day recovery period, the surviving plants were photographed and manually counted (two-sided chi-squared test, p-value <0.01).

Metabolite profiling analyses

One-week-old seedlings were placed in individual pots with 30 g of autoclaved soil and grown under LD photoperiodic conditions. After three weeks growing, half of the plants were subjected to severe drought (withholding water) for six days and the other half were watered normally (well-watered control conditions). A total of five biological replicates were collected every 24 h during the time course (from day 0 to day 6) both in drought and watered conditions and for each genotype (WT, *quad* and *BRL3ox*). Four independent plants were bulked in each biological replicate. Roots were manually separated from shoots. Four entire shoots were grinded using the Frosty Cryogenic grinder system (Labman). Four entire root samples were grinded in the Tissue Lyser Mixer-Mill (Qiagen). Roots were aliquoted into 20 mg samples and shoot into 50 mg samples (the exact weight was annotated for data normalization). Primary metabolite extraction was carried as follows [66]. One Zirconia and 500 µl of 100% Methanol premixed with Ribitol (20:1) were added and samples were subsequently homogenized in the Tissue Lyser (Qiagen) 3 min at 25 Hz. Samples were centrifuged 10 min at 14000 rpm (10 ºC) and resulting supernatant was transferred into fresh tubes. Addition of 200 µl of CHCl3 and vortex ensuring one single phase followed by the addition of 600 µl of H20 and vortex 15 sec. Samples were centrifuged 10 min at 14000 rpm (10 ºC). 100 µl from the upper phase (polar phase) were transferred into fresh eppendorf tubes (1.5 ml) and dried in the speed vacuum for at least 3 h without heating. 40 µl of derivatization agent (methoxyaminhydrochloride in pyridine) were added to each sample (20 mg/ml). Samples were shaken during 3 h at 900 rpm at 37 ºC. Drops on the cover were shortly spun down. One sample vial with 1 mL MSTFA

+ 20 µl FAME mix was prepared. Addition of 7 0µl MSTFA + FAMEs in each

sample was done followed by shaking 30 min at 37 ºC. Drops on the cover were

shortly spun down.

Samples were transferred into glass vials specific for injection in GC-TOF-MS.

The GC-TOF-MS system comprised of a CTC CombiPAL autosampler, an

Agilent 6890N gas chromatograph, and a LECO Pegasus III TOF-MS running in

EI+ mode. Metabolites were identified by comparing to database entries of

authentic standards [67]. Chromatograms were evaluated using Chroma TOF 1.0

(Leco) Pegasus software was used for peak identification and correction of RT.

Mass spectra were evaluated using the TagFinder 4.0 software [68] for metabolite

annotation and quantification (peak area measurements). The resulting data

matrix was normalized using an internal standard, Ribitol, in 100% methanol

(20:1), followed by normalization with the fresh weight of each sample.

Metabolomics data from control (well-watered) conditions at day 0 were

analyzed with a two-tailed t-test, p-value<0.05 (no multiple testing correction).

Data from the time course was analyzed with R software using the maSigPro

package [69]. Briefly, the profile of each metabolite under each condition was

fitted to a polynomial model of maximum degree 3. The curves of each

genotype were statistically compared taking into account the fitting value and

correcting the p-value (Benjamini-Hochberg method). Significant metabolites (p-

value < 0.05) having a differential profile between genotypes were plotted to

visualize their behavior under the drought time course. Clustering analysis was

performed using the maSigPro package and the *hclust* R core function.

Transcriptomic profiling analysis

For microarray analysis, a drought stress time course was carried out in WT and *quad* mutant three-week-old plants. Entire plants grown under drought stress and control conditions were collected every 48 h during the time course (Day 0, Day 2 and Day 4). Two biological replicates composed of five independent rosettes were collected. RNA was extracted with the Plant Easy Mini Kit (Qiagen) and quality checked using the Bioanalyser. A Genome-Wide Microarray platform (Dual color, Agilent) was performed by swapping the color hybridization of each biological replicate (Cy3 and Cy5). Statistical analysis was performed with the package "limma" [70], and the "mle2/"normexp" background correction method was used. Different microarrays were quantile-normalized and a Bayes test used to identify differentially expressed probes. The results were filtered for adjusted p-value<0.05 (after Benjamini-Hochberg correction) and Log2 FC >|1.5|. For RNAseq analysis, three-week-old roots were detached from mature plants grown in soil under control conditions and five days of drought. RNA was extracted as described above. Stranded cDNA libraries were prepared with TruSeq Stranded mRNA kit (Illumina). Single-end sequencing, with 50-bp reads, was performed in an Illumina HiSeq500 sequencer, at a minimum sequencing depth of 21 M. Reads were trimmed 5 bp at their 3' end, quality filtered and then mapped against the TAIR10 genome with "HISAT2". Mapped reads were quantified at the gene level with "HtSeq". For differential expression, samples were TMM normalized and statistical values calculated with the "EdgeR" package in R. Results were filtered for adjusted p-value (FDR) <0.05 and FC >|2| in the pairwise comparisons. For the evaluation of differential drought response between WT and *BRL3ox* roots, a lineal model accounting for the interaction genotype and drought was constructed with "EdgeR" package.

The interaction term was evaluated. A gene was considered to be affected by

the interaction if p-value (uncorrected) < 0.0025. Heatmaps were performed in

R with the heatmap.2 function implemented in the "gplots" package.

For the Rt qPCR, cDNA was obtained from RNA samples by using the

Transcriptor First Strand cDNA Synthesis Kit (Roche) with oligo dT primers.

qPCR amplifications were performed from 10ng of cDNA using LightCycler 480

SYBR Green I master mix (Roche) in 96-well plates according the manufacturer

recommendations. The Real Time PCR was performed on a LightCycler 480

System (Roche). Ubiquitin (AT5G56150) was used as housekeeping gene for

relativizing expression. Primers used are described in Supplementary Table 3.

Statistical methods, and omics data integration.

For root tissue enrichment analysis, deregulated genes were queried against

available lists of tissue-enriched genes [49]. For each tissue, a 2x2 contingency

table was constructed, counting the number of deregulated genes in the tissue

that were enriched and non-enriched and also the number of non-deregulated

genes (for either FDR>0.05 or logFC >/< in the RNAseq gene universe) that

were enriched and non-enriched. Statistical values of the enrichment were

obtained using a one-sided Fisher's test. To statistically evaluate the influence

of transcriptomic changes on the metabolic signature, both deregulated

enzymes and metabolites were queried in an annotation file of the metabolic

reactions of *Arabidopsis thaliana*, which included merged data from the KEGG

(http://www.genome.jp/kegg/) and BRENDA (www.brenda-enzymes.org)

databases. Then, the same approach of constructing a 2x2 contingency table

was taken. Significant and non-significant metabolites annotated in the

database were matched with differentially and non-differentially expressed

genes annotated in the database. The statistical value of the association

between regulated metabolites and genes was obtained through a two-sided

Fisher's exact test. Genes and metabolites were mapped onto the KEGG

pathways using the PaintOmics3 (http://bioinfo.cipf.es/paintomics/) according

the developer´s instructions [50].

Physiological parameters and chlorophyll fluorescence

One-week-old seedlings were placed in individual pots and watered with the

same volume of a modified Hoagland solution (one-fifth strength). Pots were

weighed daily during the experiment. Well-watered control plants were grown in

100% field capacity (0% of water loss). The time course drought stress assay

was started by withholding the nutrient solution until reaching 25, 50, 60 and

70% water loss. Photosynthesis (A) and transpiration (E) were measured in

control and drought plants at those time points. Four plants of each genotype

were harvested at 0, 50 and 70% water loss for biomass, water content and

hormone analyses. Drought experiments were repeated three times and at least

four plants per genotype and treatment were used in each experiment. RWC

was calculated according to the formula: RWC (%) = [(FW-DW) / (TW-DW)] x

100.

Plant hormones quantification

Plant hormones cytokinins (*trans*-zeatin), gibberellins (GA1, GA4 and GA3),

indole-3-acetic acid (IAA), abscisic acid (ABA), salicylic acid (SA), jasmonic acid

(JA), and the ethylene precursor 1-aminocyclopropane-1-carboxylic acid (ACC)

were analyzed as follows. 10 μl of extracted sample were injected in a UHPLC–MS system consisting of an Accela Series U-HPLC (ThermoFisher Scientific, Waltham, MA, USA) coupled to an Exactive mass spectrometer (ThermoFisher Scientific, Waltham, MA, USA) using a heated electrospray ionization (HESI) interface. Mass spectra were obtained using the Xcalibur software version 2.2 (ThermoFisher Scientific, Waltham, MA, USA). For quantification, calibration curves were constructed for each analyzed hormone (1, 10, 50, and 100 μg l−1) and corrected for 10 μg l−1 deuterated internal standards. Recovery percentages ranged between 92 and 95%.

For endogenous BR analysis plant materials (4 g fresh weight) were lyophilized and grinded. BL and CS were extracted with methanol and purified by solvent partitions by using a silica gel column and ODS-HPLC as follows. The endogenous levels of BL and CS were quantified by LC-MS/MS using their deuterated internal standards (2ng).

LC-MS/MS analysis was performed with a triple quadrupole/linear ion trap instrument (QTRAP5500; AB Sciex, USA) with an electrospray source. Ion source was maintained at 300 ₌C. Ionspray voltage was set at 4500 V in positive ion mode. MRM analysis were performed at the transitions of m/z 487 to 433 (Collision Energy, CE 30 V) and 487 to 451 (CE 21 V) for 2 H 6 -BL, m/z 481 to 427 (CE 30 V) and 481 to 445 (CE 30 V) for BL, m/z 471 to 435 (CE 23 V) and 471 to 453 (CE 25 V) for 2 H 6 -CS and m/z 465 to 429 (CE 23 V) and 465 to 447 (CE 25 V) for CS. Enhanced product ion scan was carried out at CE 21 V. HPLC separation was performed using a UHPLC (Nexera X2; Shimadzu, Japan) equipped with an ODS column (Kinetex C18, f2.1 ' 150 mm, 1.7 μm; Phenomenex, USA). The column oven temperature was maintained at 30ºC.

The mobile phase consisted of acetonitrile (solvent A) and water (solvent B),

both of which contained 0.1% (v/v) acetic acid. HPLC separation was conducted

with the following gradient at flow rate of 0.2 mL·min-1: 0 to 12 min, 20% A to

80% A; 12 to 13 min, 80% A to 100% A; 13 to 16 min, 100% A.

DATA AVAILABILITY

RNAseq and microarray data that support the findings of this study have been

deposited in Gene Expression Omnibus (GEO) with the GSE119382 and

GSGE119383 accession codes.

[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE119382"]

[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE119383]

COMPETING INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

We thank Riccardo Aiese Cigliano (Sequentia Biotech) for help with microarray

analysis, Tony Ferrar for critical manuscript revision and language editing

(http://www.theeditorsite.com)and Olga Moreno Pradas for graphic design

support. We acknowledge financial support from the Spanish Ministry of

Economy and Competitiveness, through the "Severo Ochoa Programme for

Centres of Excellence in R&D" 2016-2019 (SEV-2015-0533)", the CERCA

Programme from the Generalitat de Catalunya, and the Agencia de Gestió

d'Ajuts Universitaris i de Recerca (2014 SGR 1406). N.F. was funded by

Fundación Renta Corporación, by EMBO short-term postdoctoral fellowship

(ASTF 422-2015) and by BIO2013-43873; F.L.E was funded by BIO2013-43873

and BIO2016-78150-P; and D.B. was funded by BIO2013-43873 and ERC-

2015-CoG-GA 683163. CMA, AA and FPA acknowledge financial support from

the Spanish MINECO-FEDER (project AGL2014-59728-R) and from the

European Union's Seventh Framework Programme for research technological

development and demonstration under grant agreement no. 289365 (project

ROOTOPOWER). S.O. was supported by Ministry of Science and Innovation

and University of Malaga (Spain) through the grant Ramón y Cajal program

(Sonia Osorio, RYC-09170). M.B was funded by BES-2012-053274 and J.L.R

laboratory is supported by grant BFU2014-58289-P from the Spanish Ministry of

Economy and Competitiveness. A.I.C.-D. – A.C. collaboration was funded by

the European Regional Development Funds and Marie Curie IRSES Project

DEANN (PIRSES-GA-2013-612583). A.I.C.-D. is a recipient of BIO2013-43873

and BIO2016-78150-P grants from the Spanish Ministry of Economy and

Competitiveness. A.I.C.-D. is recipient of an ERC Consolidator Grant ; this

project has received funding from the European Research Council (ERC) under

the European Union's Horizon 2020 research and innovation program (grant

agreement No 683163).

AUTHOR CONTRIBUTIONS

A.I.C.-D conceived, designed and supervised the study. N.F., F.L.E, D.B.,

C.M.A. and F.P.A. performed the genetics and stress phenomic experiments

and data analyses. N.F., F.L.E., M.B., J.L.R. and A.C.D. designed the genome-

wide experiments. N.F., F.L.E., A.C. and A.C.D. performed the genome-wide

experiments and data analysis. N.F., F.L.E., T.T., S.O., and A.R.F. carried out

the metabolite and data analysis. A.A. performed the hormone profiling

experiments. T.N. and T.Y. carried the CS and BL quantification assays. N.F.,

F.L.E and A.C.D. wrote the manuscript. All authors discussed the results and

the manuscript.

REFERENCES

Boyer, J. S. Plant productivity and environment. *Science* **218**, 443-448, doi:10.1126/science.218.4571.443 (1982).

Todaka, D., Shinozaki, K. & Yamaguchi-Shinozaki, K. Recent advances in the dissection of drought-stress regulatory networks and strategies for development of drought-tolerant transgenic rice plants. *Front Plant Sci* **6**, 84, doi:10.3389/fpls.2015.00084 (2015).

Yoshida, T., Mogami, J. & Yamaguchi-Shinozaki, K. ABA-dependent and ABA-independent signaling in response to osmotic stress in plants. *Curr Opin Plant Biol* **21**, 133-139, doi:10.1016/j.pbi.2014.07.009 (2014).

Graether, S. P. & Boddington, K. F. Disorder and function: a review of the dehydrin protein family. *Front Plant Sci* **5**, 576, doi:10.3389/fpls.2014.00576 (2014).

Seki, M., Umezawa, T., Urano, K. & Shinozaki, K. Regulatory metabolic networks in drought stress responses. *Curr Opin Plant Biol* **10**, 296-302, doi:10.1016/j.pbi.2007.04.014 (2007).

Shao, H. B., Chu, L. Y., Jaleel, C. A. & Zhao, C. X. Water-deficit stress-induced anatomical changes in higher plants. *C R Biol* **331**, 215-225, doi:10.1016/j.crvi.2008.01.002 (2008).

Tardieu, F. Any trait or trait-related allele can confer drought tolerance: just design the right drought scenario. *J Exp Bot* **63**, 25-31, doi:10.1093/jxb/err269 (2012).

Marshall, A. *et al.* Tackling drought stress: receptor-like kinases present new approaches. *Plant Cell* **24**, 2262-2278, doi:10.1105/tpc.112.096677 (2012).

9   Smakowska-Luzan, E. *et al.* An extracellular network of Arabidopsis

leucine-rich repeat receptor kinases. *Nature* **553**, 342-346,

doi:10.1038/nature25184 (2018).

10   Li, J. & Chory, J. A putative leucine-rich repeat receptor kinase involved

in brassinosteroid signal transduction. *Cell* **90**, 929-938 (1997).

11   Wang, Z. Y., Seto, H., Fujioka, S., Yoshida, S. & Chory, J. BRI1 is a

critical component of a plasma-membrane receptor for plant steroids.

Nature **410**, 380-383, doi:10.1038/35066597 (2001).

12   Kinoshita, T. *et al.* Binding of brassinosteroids to the extracellular domain

of plant receptor kinase BRI1. *Nature* **433**, 167-171,

doi:10.1038/nature03227 (2005).

13   Hothorn, M. *et al.* Structural basis of steroid hormone perception by the

receptor kinase BRI1. *Nature* **474**, 467-471, doi:10.1038/nature10153

(2011).

14   She, J. *et al.* Structural insight into brassinosteroid perception by BRI1.

Nature **474**, 472-476, doi:10.1038/nature10178 (2011).

15   Karlova, R. *et al.* The Arabidopsis SOMATIC EMBRYOGENESIS

RECEPTOR-LIKE KINASE1 protein complex includes

BRASSINOSTEROID-INSENSITIVE1. *Plant Cell* **18**, 626-638,

doi:10.1105/tpc.105.039412 (2006).

16   Li, J. *et al.* BAK1, an Arabidopsis LRR receptor-like protein kinase,

interacts with BRI1 and modulates brassinosteroid signaling. *Cell* **110**,

213-222 (2002).

17   Nam, K. H. & Li, J. BRI1/BAK1, a receptor kinase pair mediating

brassinosteroid signaling. *Cell* **110**, 203-212 (2002).

18   Gou, X. *et al.* Genetic evidence for an indispensable role of somatic

embryogenesis receptor kinases in brassinosteroid signaling. *PLoS*

Genet **8**, e1002452, doi:10.1371/journal.pgen.1002452 (2012).

19   Yin, Y. *et al.* BES1 accumulates in the nucleus in response to

brassinosteroids to regulate gene expression and promote stem

elongation. *Cell* **109**, 181-191 (2002).

20   Wang, Z. Y. *et al.* Nuclear-localized BZR1 mediates brassinosteroid-

induced growth and feedback suppression of brassinosteroid

biosynthesis. *Dev Cell* **2**, 505-513 (2002).

21   He, J. X., Gendron, J. M., Yang, Y., Li, J. & Wang, Z. Y. The GSK3-like

kinase BIN2 phosphorylates and destabilizes BZR1, a positive regulator

of the brassinosteroid signaling pathway in Arabidopsis. *Proc Natl Acad*

Sci U S A **99**, 10185-10190, doi:10.1073/pnas.152342599 (2002).

22   Kagale, S., Divi, U. K., Krochko, J. E., Keller, W. A. & Krishna, P.

Brassinosteroid confers tolerance in Arabidopsis thaliana and Brassica

napus to a range of abiotic stresses. *Planta* **225**, 353-364,

doi:10.1007/s00425-006-0361-6 (2007).

23   Krishna, P. Brassinosteroid-Mediated Stress Responses. *J Plant Growth*

Regul **22**, 289-297, doi:10.1007/s00344-003-0058-z (2003).

24   Sahni, S. *et al.* Overexpression of the brassinosteroid biosynthetic gene

DWF4 in Brassica napus simultaneously increases seed yield and stress

tolerance. *Sci Rep* **6**, 28298, doi:10.1038/srep28298 (2016).

25   Feng, Y., Yin, Y. & Fei, S. Down-regulation of BdBRI1, a putative

brassinosteroid receptor gene produces a dwarf phenotype with

enhanced drought tolerance in Brachypodium distachyon. *Plant Sci* **234**,

163-173, doi:10.1016/j.plantsci.2015.02.015 (2015).

26   Ye, H. *et al.* RD26 mediates crosstalk between drought and

brassinosteroid signalling pathways. *Nat Commun* **8**, 14573, doi:10.1038/

ncomms14573 (2017).

27   Zhang, S., Cai, Z. & Wang, X. The primary signaling outputs of

brassinosteroids are regulated by abscisic acid signaling. *Proc Natl Acad*

Sci U S A **106**, 4543-4548, doi:10.1073/pnas.0900349106 (2009).

28   Gui, J. *et al.* OsREM4.1 Interacts with OsSERK1 to Coordinate the

Interlinking between Abscisic Acid and Brassinosteroid Signaling in Rice.

Dev Cell **38**, 201-213, doi:10.1016/j.devcel.2016.06.011 (2016).

29   Huang, D., Wu, W., Abrams, S. R. & Cutler, A. J. The relationship of

drought-related gene expression in Arabidopsis thaliana to hormonal and

environmental factors. *J Exp Bot* **59**, 2991-3007, doi:10.1093/jxb/ern155

(2008).

30   Chung, Y., Kwon, S. I. & Choe, S. Antagonistic regulation of Arabidopsis

growth by brassinosteroids and abiotic stresses. *Mol Cells* **37**, 795-803,

doi:10.14348/molcells.2014.0127 (2014).

31   Vilarrasa-Blasi, J. *et al.* Regulation of plant stem cell quiescence by a

brassinosteroid signaling module. *Dev Cell* **30**, 36-47,

doi:10.1016/j.devcel.2014.05.020 (2014).

32   Fàbregas, N. & Caño-Delgado, A. I. Turning on the microscope turret: a

new view for the study of brassinosteroid signaling in plant development.

Physiol Plant **151**, 172-183, doi:10.1111/ppl.12130 (2014).

33   Fàbregas, N. *et al.* The brassinosteroid insensitive1-like3 signalosome

complex regulates Arabidopsis root development. *Plant Cell* **25**, 3377-

3388, doi:10.1105/tpc.113.114462 (2013).

34   Lozano-Elena, F., Planas-Riverola, A., Vilarrasa-Blasi, J., Schwab, R. &

Cano-Delgado, A. I. Paracrine brassinosteroid signaling at the stem cell

niche controls cellular regeneration. *J Cell Sci* **131**, jcs204065

doi:10.1242/jcs.204065 (2018).

35   Friedrichsen, D. M., Joazeiro, C. A., Li, J., Hunter, T. & Chory, J.

Brassinosteroid-insensitive-1 is a ubiquitously expressed leucine-rich

repeat receptor serine/threonine kinase. *Plant Physiol* **123**, 1247-1256

(2000).

36   Caño-Delgado, A. *et al.* BRL1 and BRL3 are novel brassinosteroid

receptors that function in vascular differentiation in Arabidopsis.

Development **131**, 5341-5351, doi:10.1242/dev.01403 (2004).

37   Belkhadir, Y. & Jaillais, Y. The molecular circuitry of brassinosteroid

signaling. *New Phytol* **206**, 522-540, doi:10.1111/nph.13269 (2015).

38   Eremina, M. *et al.* Brassinosteroids participate in the control of basal and

acquired freezing tolerance of plants. *Proc Natl Acad Sci U S A* **113**,

E5982-E5991, doi:10.1073/pnas.1611477113 (2016).

39   Vragovic, K. *et al.* Translatome analyses capture of opposing tissue-

specific brassinosteroid signals orchestrating root meristem

differentiation. *Proc Natl Acad Sci U S A* **112**, 923-928,

doi:10.1073/pnas.1417947112 (2015).

40 González-García, M. P. *et al.* Brassinosteroids control meristem size by
promoting cell cycle progression in Arabidopsis roots. *Development* **138**,
849-859, doi:10.1242/dev.057331 (2011).

41 Hacham, Y. *et al.* Brassinosteroid perception in the epidermis controls
root meristem size. *Development* **138**, 839-848, doi:10.1242/dev.061804
(2011).

42 Duan, Y. *et al.* An endoplasmic reticulum response pathway mediates
programmed cell death of root tip induced by water stress in Arabidopsis.
New Phytol **186**, 681-695, doi:10.1111/j.1469-8137.2010.03207.x (2010).

43 Takahashi, N., Goto, N., Okada, K. & Takahashi, H. Hydrotropism in
abscisic acid, wavy, and gravitropic mutants of Arabidopsis thaliana.
Planta **216**, 203-211, doi:10.1007/s00425-002-0840-3 (2002).

44 Asami, T. *et al.* Characterization of brassinazole, a triazole-type
brassinosteroid biosynthesis inhibitor. *Plant Physiol* **123**, 93-100 (2000).

45 Singh, T. N., Aspinall, D. & Paleg, L. G. Proline accumulation and varietal
adaptability to drought in barley: a potential metabolic measure of
drought resistance. *Nat New Biol* **236**, 188-190 (1972).

46 Szabados, L. & Savoure, A. Proline: a multifunctional amino acid. *Trends
Plant Sci* **15**, 89-97, doi:10.1016/j.tplants.2009.11.009 (2010).

47 Durand, M. *et al.* Water Deficit Enhances C Export to the Roots in
Arabidopsis thaliana Plants with Contribution of Sucrose Transporters in
Both Shoot and Roots. *Plant Physiol* **170**, 1460-1479,
doi:10.1104/pp.15.01926 (2016).

48 Conrath, U., Pieterse, C. M. & Mauch-Mani, B. Priming in plant-pathogen
interactions. *Trends Plant Sci* **7**, 210-216 (2002).

49   Brady, S. M. *et al.* A high-resolution root spatiotemporal map reveals
dominant   expression   patterns.   *Science*   **318**,   801-806,
doi:10.1126/science.1146265 (2007).

50   García-Alcalde, F., García-López, F., Dopazo, J. & Conesa, A.
Paintomics: a web based tool for the joint visualization of transcriptomics
and   metabolomics   data.   *Bioinformatics*   **27**,   137-139,
doi:10.1093/bioinformatics/btq594 (2011).

51   Shakirova, F. *et al.* Involvement of dehydrins in 24-epibrassinolide-
induced protection of wheat plants against drought stress. *Plant*
Physiology   and   Biochemistry   **108**,   539-548,
doi:10.1016/j.plaphy.2016.07.013 (2016).

52   Chen, J. *et al.* Arabidopsis WRKY46, WRKY54, and WRKY70
Transcription Factors Are Involved in Brassinosteroid-Regulated Plant
Growth and Drought Responses. *The Plant Cell* **29**, 1425 (2017).

53   Iwata, S., Miyazawa, Y., Fujii, N. & Takahashi, H. MIZ1-regulated
hydrotropism functions in the growth and survival of Arabidopsis thaliana
under natural conditions. *Ann Bot* **112**, 103-114, doi:10.1093/aob/mct098
(2013).

54   Ge, L.-F. *et al.* Overexpression of the trehalose-6-phosphate
phosphatase gene OsTPP1 confers stress tolerance in rice and results in
the activation of stress responsive genes. *Planta* **228**, 191-201,
doi:10.1007/s00425-008-0729-x (2008).

55   Nuccio, M. L. *et al.* Expression of trehalose-6-phosphate phosphatase in
maize ears improves yield in well-watered and drought conditions.
Nature Biotechnology **33**, 862, doi:10.1038/nbt.3277 (2015).

56　Himuro, Y. *et al.* Arabidopsis galactinol synthase AtGolS2 improves

drought tolerance in the monocot model Brachypodium distachyon.

Journal of Plant Physiology **171**, 1127-1131,

doi:10.1016/j.jplph.2014.04.007 (2014).

57　Nanjo, T. *et al.* Antisense suppression of proline degradation improves

tolerance to freezing and salinity in Arabidopsis thaliana. *FEBS Letters*

461, 205-210, doi:10.1016/S0014-5793(99)01451-9 (1999).

58　Wu, C. Y. *et al.* Brassinosteroids regulate grain filling in rice. *Plant Cell*

20, 2130-2145, doi:10.1105/tpc.107.055087 (2008).

59　Xu, F., Xi, Z.-m., Zhang, H., Zhang, C.-j. & Zhang, Z.-w. Brassinosteroids

are involved in controlling sugar unloading in Vitis vinifera 'Cabernet

Sauvignon' berries during véraison. *Plant Physiology and Biochemistry*

94, 197-208, doi:10.1016/j.plaphy.2015.06.005 (2015).

60　Krasensky, J. & Jonak, C. Drought, salt, and temperature stress-induced

metabolic rearrangements and regulatory networks. *J Exp Bot* **63**, 1593-

1608, doi:10.1093/jxb/err460 (2012).

61　Urano, K. *et al.* Characterization of the ABA-regulated global responses

to dehydration in Arabidopsis by metabolomics. *Plant J* **57**, 1065-1078,

doi:10.1111/j.1365-313X.2008.03748.x (2009).

62　Ding, Y., Fromm, M. & Avramova, Z. Multiple exposures to drought 'train'

transcriptional responses in Arabidopsis. *Nat Commun* **3**, 740,

doi:10.1038/ncomms1732 (2012).

63　Nishizawa, A., Yabuta, Y. & Shigeoka, S. Galactinol and raffinose

constitute a novel function to protect plants from oxidative damage. *Plant*

Physiol **147**, 1251-1263, doi:10.1104/pp.108.122465 (2008).

64   Aljanabi, S. M. & Martinez, I. Universal and rapid salt-extraction of high

quality genomic DNA for PCR-based techniques. *Nucleic Acids Res* **25**,

4692-4693 (1997).

65   Betegón-Putze, I., González, A., Sevillano, X., Blasco-Escámez, D. &

Caño-Delgado, A. I. . MyROOT: A novel method and software for the

semi-automatic measurement of plant root length. Preprint at *bioRxiv*

doi:10.1101/309773 (2018).

66   Lisec, J., Schauer, N., Kopka, J., Willmitzer, L. & Fernie, A. R. Gas

chromatography mass spectrometry-based metabolite profiling in plants.

Nat Protoc **1**, 387-396, doi:10.1038/nprot.2006.59 (2006).

67   Kopka, J. *et al.* GMD@CSB.DB: the Golm Metabolome Database.

Bioinformatics **21**, 1635-1638, doi:10.1093/bioinformatics/bti236 (2005).

68   Luedemann, A., Strassburg, K., Erban, A. & Kopka, J. TagFinder for the

quantitative analysis of gas chromatography--mass spectrometry (GC-

MS)-based metabolite profiling experiments. *Bioinformatics* **24**, 732-737,

doi:10.1093/bioinformatics/btn023 (2008).

69   Conesa, A., Nueda, M. J., Ferrer, A. & Talon, M. maSigPro: a method to

identify significantly differential expression profiles in time-course

microarray experiments. *Bioinformatics* **22**, 1096-1102,

doi:10.1093/bioinformatics/btl056 (2006).

70   Ritchie, M. E. *et al.* limma powers differential expression analyses for

RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47,

doi:10.1093/nar/gkv007 (2015).

 FIGURE LEGENDS

Figure 1. BR perception mutant roots are less sensitive to osmotic stress.

(**a**) Seven-day-old roots of WT, BR mutants *bak1*, *brl1brl3, brl1brl3bak1*, *bri1,*

bri1brl1brl3, and *bri1brl1brl3bak1* (*quad*), and BR overexpressor line

35S:BRL3-GFP (BRL3ox) grown in control (-) or 270 mM sorbitol (+) conditions.

Scale bar: 0.5 cm. (**b**) Boxplots depict the distribution of seven-day-old root

lengths in control (dark green) or sorbitol (light green) conditions. Red line

depicts relative root growth inhibition upon stress (ratio sorbitol/control +/-

s.e.m.). Data from five independent biological replicates (n>150). Different

letters represent significant differences (p-value<0.05) in an ANOVA plus

Tukey's HSD test. (**c**) Four-day-old roots stained with propidium iodide (PI, red)

after 24 h in control (top) or sorbitol (bottom) media. Green channel (GFP)

shows the BRL3 membrane protein receptor under the 35SCaMV constitutive

promoter localizing to the vascular tissues in primary roots. Scale bar: 100 µm.

(**d**) Quantification of cell death in sorbitol-treated root tips. Boxplots represent

the relative PI staining (sorbitol/control) for each genotype. Averages from five

independent biological replicates (n>31). Different letters represent significant

differences (p-value<0.05) in an ANOVA plus Tukey's HSD test. Boxplots

represent the median and interquartile range (IQR). Whiskers depict Q1-

1.5*IQR and Q3+1.5*IQR and points experimental observation.

Figure 2. The BRL3 receptor promotes root hydrotropism responses.

(**a**) Root curvature (hydrotropic response) in seven-day-old roots after 24 h of

sorbitol-induced osmotic stress (270 mM). Scale bar: 0.2 cm. (**b**) Discrete

distribution of root hydrotropic curvature angles in the different genotypes.

Lightest green depicts roots curved between 0º and 10º, light green between

10º and 20º, dark grey between 20º and 30º, and darkest green depicts roots

that have a curvature of more than 30º as indicated in the color legend. (**c**)

Continuous distribution of root curvature angles. Different letters indicate a

significant difference (p-value<0.05) in a one-way ANOVA test plus Tukey's

HSD test. Boxplot represent the median and interquartile range (IQR). Whiskers

depict Q1-1.5*IQR and Q3+1.5*IQR and points experimental observations. Data

from four independent biological replicates (n>50). (**d**) Stress traits matrix for all

physiological assays performed on the roots and shoots of WT, BR loss-of-

function mutants and *BRL3ox*. Root growth in control conditions is highlighted in

green. Color bar depicts values for scaled data.

Figure 3. BRL3 overexpression confers drought tolerance.

(**a**) From top to bottom, three-week-old plant rosette phenotypes of WT,

brl1brl3, bak1, *brl1brl3bak1, bri1, bri1brl1brl3, quad* and *BRL3ox* grown in well-

watered conditions (left column), after 12 days of drought stress (middle

column) and after 7 days of re-watering (right column). (**b**) Plant survival rates

after 7 days of re-watering. Averages of five independent biological replicates

+/- s.e.m. (n>140). Asterisk indicates a significant difference (p-value<0.05) in a

chi-squared test for survival ratios compared to WT. (**c**) Bar plot shows the days

needed to reach different percentages of the soil field capacity for each

genotype used in the study. (**d**) Relative water content (RWC) of mature

rosettes at 0% (field capacity), 50% and 70% soil water loss. (**e**) Photosynthesis

efficiency (µmol/m2*s) at different percentages of soil water loss. (**d, e**) Boxplot

represent the median and interquartile range (IQR). Whiskers depict Q1-

1.5*IQR and Q3+1.5*IQR and points experimental observations (n=6). Different

letters depict significant differences within each genotype in a one-way ANOVA

plus a Tukey's HSD test. (**f**) Schematic representation of BR signaling levels,

adult plant size and drought resistance. Loss-of-function mutants passively

avoid stress (drought avoidance), whereas plants with increased levels of BRL3

act actively to avoid drought stress (drought tolerance).

Figure 4. BRL3 overexpression plants show a primed metabolic signature.

(**a**) Metabolites differentially accumulated in *BRL3ox* (dark green) *or quad (*light

green) shoots relative to WT at basal conditions. (**b**) Metabolites differentially

accumulated in *BRL3ox* (dark green) *or quad (*light green) roots relative to WT

at basal conditions. (**a, b**) Boxplot represent the median and interquartile range

(IQR). Whiskers depict Q1-1.5*IQR and Q3+1.5*IQR and points experimental

observations (n=5). Asterisks denote statistical differences in a two-tailed t-test

(p-value < 0.05) for raw data comparisons *BRL3ox* vs WT (panel right side) or

quad (panel left side). (**c**) Metabolites following differential dynamics between

BRL3ox and WT shoots along the drought time course. (**d**) Metabolites

following differential dynamics between *BRL3ox* and WT roots along the

drought time course. (**c, d**) Heatmap represents the log2 ratio of *BRL3ox*/WT.

(**e-j**) Clustering of the dynamics of relative metabolite levels along the drought

time course in shoots and roots. Solid lines show the actual metabolic profile

(averages) of the representative metabolite for each cluster while dashed lines

represent the polynomial curve that best fit the profile. Statistical significance

was evaluated with the maSigPro package. (**e**) Metabolites following a linear

increase during drought in shoots include Glucose, Glucose-1P, *myo*-inositol,

and Sinapate. (**f**) Proline follows a steeper exponential increase in *BRL3ox*

shoots. (**g**) Metabolites following an exponential increase in *BRL3ox* shoots but

nearly a linear increase in WT include galactose, GABA, phenylalanine,

tyrosine, 2-methylmalate, lysine, isoleucine, leucine, nicotinate, uracil, and

tryptophan. (**h**) Metabolites following a steeper exponential increase in *BRL3ox*

roots include trehalose, sucrose, proline and raffinose. (**i**) Metabolites following

a reduced linear increase until a certain maximum in *BRL3ox* roots include

glycerate and malate. (**j**) Metabolites following an exponential increase in

BRL3ox roots but a linear increase in WT include glucose, fructose, *myo*-

inositol, galactose, and asparagine.

1004

Figure 5. Stress genes are constitutively activated in *BRL3ox* roots.

(**a**) Most representative GO categories enriched in *BRL3ox* roots from the

upregulated genes at time 0 and after 5 days of drought (**b**) Deployment of

genes within "Response to stress" (GO:0006950) term that are also annotated

as responsive to water, salt, heat, cold, and light stress. Colors in the heatmap

represent the log2 fold change of *BRL3ox* vs. WT roots in control conditions (C)

or the differential drought response (log2(FC drought/CTRL in *BRL3ox*) –

log2(FCdrought/CTRL in WT)) if the gene is affected by the interaction

genotype*drought (Int.). Red color in the squared heatmaps on the right shows

that the gene has been previously identified as a direct target of BES1 or BZR1

transcription factors. (**c**) Most representative GO categories enriched in *BRL3ox*

roots from the upregulated genes at time 0 and after 5 days of drought. (**d**) Most

representative GO categories enriched among genes affected by the interaction

genotype-drought. GO categories enriched in genes activated in *BRL3ox* under

drought compared to WT (left column) in genes repressed in *BRL3ox* under

drought compared to WT (right column). Color bars: –log of p-value (adjusted

by Benjamini-Hochberg or non-adjusted).

1022

Figure 6. Enrichment of deregulated genes in *BRL3ox* root vasculature.

(**a**) Tissue enrichment for upregulated (red) or downregulated (blue) genes in

control conditions. Bars trespassing the p-value threshold (0.05) were

considered enriched in the dataset. (**b**) Tissue enrichment for genes affected by

the interaction genotype*drought Bars trespassing the threshold p-value<0.05

were considered enriched in the dataset. (**a-b**) Deregulated genes tissue

enrichment.   AGL42: Quiescent center, APL: Phloem + Companion cells,

COBL9: Root hair cells, CORTEX: Cortex, GL2: Non-hair cells, J2661:

Pericycle, JO121: Xylem pole pericycle, LRC: Lateral root cap, PET111:

Columella, RM1000: Lateral root primordia, S17: Phloem pole pericycle, S18:

Maturing Xylem, S32: Protophloem, S4: Developing xylem, SCR5: Endodermis,

SUC2: Phloem. y-axes represent the negative logarithm of one-tailed Fisher's

test. (**c**) Deregulated genes enriched in the Pericycle (J2261 marker).   (**d**)

Deregulated genes enriched in the Phloem Pole Pericycle (S17 marker). (**c,d**)

Bars represent the log2 fold-change of *BRL3ox* vs. WT roots in control (black)

or the difference of drought responses between *BRL3ox* and WT (FC

drought/CTRL in *BRL3ox* – FC drought/CTRL in WT) in the lineal model (gray).

Blue boxes highlight enzymes directly involved in the metabolism of deregulated

metabolites.

1042

1043

Figure 1. BR perception mutant roots are less sensitive to osmotic stress.

Figure 2. The BRL3 receptor promotes root hydrotropism responses

Figure 3. BRL3 overexpression confers drought tolerance.

Figure 4. BRL3 overexpression plants show a primed metabolic signature

Figure 5. Stress genes are constitutively activated in *BRL3ox* roots.

Figure 6. Enrichment of deregulated genes in *BRL3ox* root vasculature.