Vascular brassinosteroid receptors confer drought resistance without

penalizing plant growth

Norma Fàbregas[1,*], Fidel Lozano-Elena[1,*], David Blasco-Escámez[1], Takayuki Tohge[2,8],

Cristina Martínez-Andújar[3], Alfonso Albacete[3], Sonia Osorio[4], Mariana Bustamante[1],

José Luis Riechmann[1,7], Ana Conesa[5,6], Francisco Pérez Alfocea[3], Alisdair R. Fernie[2]

and Ana I. Caño-Delgado[1,¶]

[1]*Centre for Research in Agricultural Genomics (CRAG) CSIC-IRTA-UAB-UB,*

Barcelona E-08193, Spain.

[2]*Max Planck Institute of Molecular Plant Physiology, Potsdam-Golm D-14476,*

Germany.

[3]*Department of Plant Nutrition, CEBAS-CSIC, Murcia E-30100, Spain.*

[4]*Instituto de Hortofruticultura Subtropical y Mediterránea "La Mayora". University of*

Malaga-Consejo Superior de Investigaciones Científicas. Department of Molecular

Biology and Biochemistry, Málaga 29071, Spain.

[5]*Microbiology and Cell Science Department, IFAS, Genetics Institute, University of*

Florida, Gainesville, USA.

[6]*Genomics of Gene Expression Lab, Centro de Investigaciones Príncipe Felipe,*

Valencia, Spain.

[7]*Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, 08010, Spain*

[8]*Current Address: NAIST Graduate school of Biological Sciences, 8916-5 Takayama,*

Ikoma, Nara 630-0192, Japan.

[*] Both authors contributed equally to this work.

[¶]Correspondence to: ana.cano@cragenomica.es

ABSTRACT

Drought is the primary cause of global agricultural losses and represents a major threat

to worldwide food security. Currently, plant biotechnology stands as the most

promising strategy to increase crop growth in rain-fed conditions. The main

mechanisms underlying drought resistance have been uncovered from the study of plant

physiology and by engineering drought resistance genes into crops. However, plants

with enhanced drought resistance usually display lower growth, necessitating the search

for novel strategies to uncouple drought resistance from growth. Here, we show that the

brassinosteroid family of receptors, in addition to promoting growth, guides the

phenotypic adaptation responses to a great variety of drought stress traits herein

analyzed. Whilst mutations in the ubiquitously localized BRI1 receptor pathway show

an enhanced drought resistance at the expense of growth, we found that vascular-

localized BRL3 receptors confer drought tolerance without penalizing overall plant

growth. Systematic analyses reveal that upon drought stress the vascular BRL3 receptor

pathway triggers the synthesis and mobilization of osmoprotectant metabolites, mainly

proline and sugars, preferentially in the roots favoring overall plant growth. Altogether,

our results uncover a new role for the spatial control of BR signaling in drought

tolerance, while offer a novel strategy to address food security issues in an increasingly

water limited climate.

Summary

- Brassinosteroid receptors control the plant responses to multiple drought traits in different developmental stages of roots and shoots organs.

- The vascular BRL3 receptor pathway triggers the production of osmoprotectant metabolites, preferentially sugars and proline, that once accumulated in the root help the plant to tolerate drought stress.

- BRL3 signaling promotes the transcription of canonical water stress-responses and osmoprotectant metabolism genes in normal conditions and under drought.

- Plants with increased levels of BRL3 receptors in the vascular tissues are primed and resistant to severe drought without penalizing overall plant growth.

INTRODUCTION

Drought causes at least 40% of crop losses worldwide and this proportion is dramatically increasing due to climate change [1]. Understanding cellular responses to drought stress represents the first step toward the development of better-adapted crops, which is a grand challenge for plant biotechnology [2]. Classical approaches to unravel how plants cope with limited water led to the identification of regulators in the signal transduction cascades of the abscisic acid (ABA)-dependent [3] and ABA-independent pathways [4]. Drought-stress adaptation has been associated with the presence of proteins that protect cells from dehydration, such as late-embryogenesis–abundant (LEA) proteins, osmoprotectants and detoxification enzymes [5-7]. These studies provided deep insights into the molecular mechanisms underlying abiotic stress [2], while showing that drought resistance is a complex trait simultaneously controlled by many genes. Genetic approaches have succeeded in conferring stress resistance to plants, but this generally comes at the cost of reduced growth [8,9]. Therefore, understanding how cellular growth is coupled to drought stress responses is essential for engineering plants with improved growth in rain-fed environments.

Receptor-Like Kinases (RLKs) play an important role in optimizing plant responses to drought stress [10,11]. Brassinosteroid (BR) hormones directly bind to the BRI1 (BR-INSENSITIVE 1) plasma membrane leucine-rich repeat (LRR)-RLK family members [12-17]. Ligand perception triggers BRI1 interaction with the co-receptor BAK1 (BRI1 ASSOCIATED RECEPTOR KINASE 1) [18-20], which is essential for early BR signaling events [21]. This BRI1-BAK1 heterodimerization initiates a signaling cascade of phosphorylation events that control the expression of multiple BR-regulated genes mainly via the BRI1-EMS-SUPPRESSOR1 (BES1) and BRASSINAZOLE RESISTANT1 (BZR1) transcription factors [22-26].

BRs modulate multiple developmental and environmental stress responses in the plant

yet the role of BRs under stress conditions remains controversial. The exogenous

application of BRs and the overexpression of the BR biosynthetic enzyme DWF4 both

confer increased plant adaptation to drought stress [27-29], whereas drought resistant

phenotypes result from suppressing the BRI1 receptor [30,31]. Intriguingly, ABA signaling

inhibits the BR signaling pathway after BR perception and upstream of the BIN2

(BRASSINOSTEROID-INSENSITIVE 2) kinase [32,33] and further downstream crosstalk

is mediated by the overlapping transcriptional control of multiple BR- and ABA-

regulated genes [34,35], such as *RESPONSE TO DESICCATION 26* (*RD26*) [31].

The spatial regulation of hormonal signaling pathways is receiving increasing attention

in attempt to enhance understanding of coordination of plant growth and stress

responses [31,36-38]. The BRI1 receptor is widely localized in many tissues [39] and the

BRI1-LIKE receptor homologues BRL1 and BRL3 signal from the innermost tissues of

the plant contributing to vascular development [15,40,41]. The BR receptor complexes are

formed by combinations of homologous LRR-RLK receptors of the BRI1-like family

with the BAK1 co-receptor in the plasma membrane [41]. Despite BRI1 being a central

player in plant growth and adaptation to abiotic stress [31,42,43], the functional relevance of

vascular BRL1 and BRL3 receptors has only started to be unraveled [41,44,45]. In previous

proteomic approaches we identified abiotic stress-related proteins in the BRL3

signalosome complexes [41], yet the involvement of the BRL3 pathway in drought

remains elusive.

Here, we show that the different BR receptors modulate plant adaptation to multiple

drought stress traits in roots and shoots. While the traits controlled by the BRI1 pathway

are intimately linked to growth arrest, we found that vascular-enriched BRL3 receptors

confer drought resistance without penalizing overall plant growth. Metabolite profiling

unveils that the BRL3 signaling triggers the production of an osmoprotectant metabolite

signature (i.e. proline, trehalose, sucrose and raffinose family oligosaccharides) in the

plant, and their specific accumulation in the roots during drought periods. Subsequent

transcriptomic profiling shows that this metabolite signature is transcriptionally

regulated by BRL3 pathway in response to drought. The enrichment of deregulated

genes in vascular tissues, especially in the phloem, further supports the mobilization of

osmoprotectant metabolites to root tissues driven by BRL3 signaling under drought.

Overall, this study uncovers that BRL3 signaling boost sugar and osmoprotectant

metabolites accumulation in the root hacking drought-stress arrested growth, while offer

a novel strategy to protect crops from drought.

RESULTS

BR receptors control the sensitivity to osmotic stress responses since early stages of

plant growth

To investigate the contribution of the BR perception complexes to drought, we

performed a comprehensive characterization of different mutant combinations for all the

BR receptors and the BAK1 co-receptor. First, the primary root growth was analyzed

(Fig. 1a). Seven-day-old roots of *bak1*, *brl1brl3bak1*, *bri1* and *bri1brl1brl3* displayed

shorter roots than Col-0 wild-type (WT) as previously described [20,41,46,47]. We found that

the primary roots of the quadruple mutant *bri1brl1brl3bak1* (hereafter *quad*) were the

shortest and the most insensitive to BRs (Fig. 1a,b and Supplementary Fig. 1).

Conversely, plants overexpressing BRL3 (*35S:BRL3-GFP,* hereafter *BRL3ox*) show

increased receptor levels in the vascular tissues [41] and exhibited longer roots than WT

(Fig. 1a,b). These results are in agreement with the previously known root growth-

promoting role of BR receptors [46,48]. Then osmotic stress was imposed transferring

Arabidopsis seedlings to sorbitol-containing media and root-growth inhibition was

quantified relative to control conditions (methods). A significantly lesser root-growth

inhibition was observed in *bri1* (27%), *bri1brl1brl3* (28%) and *quad* (27%) mutants

when compared to WT (39%), whereas no differences were found in *brlbrl3* and

bak1brl1brl3 mutant roots (Fig. 1a,b). Roots of *BRL3ox* plants were similar to WT in

terms of root growth inhibition under osmotic stress (Fig. 1a,b). A shorter period of

osmotic stress (24h) caused cell death in the elongation zone of WT roots, as marked by

the propidium iodine (PI) staining incorporation into nuclei [49] (Fig. 1c,d). Consistently,

bri1, bri1brl1brl3 and *quad* mutants displayed reduced sensitivity to osmotic stress

compared to WT, as revealed by a reduced cell death in the roots (Fig. 1c,d).

Conversely, plants with increased levels of BRL3 receptor showed a massive cell death

in the root tips compared to the WT, indicating an increased sensitivity to short osmotic

stress treatments (Fig 1c,d). These results point to a positive role for BR receptors in

triggering osmotic stress responses in the plant root.

Since root hydrotropism represents a key feature for adaptation to water scarcity

environments [50,51], the capacity of the roots to escape from an imposed osmotic stress

bending towards water-available media was investigated (Fig. 2a). Loss-of-function

mutants in the BR receptors showed reduced hydrotropic responses, as roots grew

straighter in the sorbitol containing media than in the WT roots (Fig. 2a-c), while no

significant differences were found in control conditions (mock) (Supplementary Fig. 2).

Interestingly, *brl1brl3bak1* exhibited the most insensitive hydrotropic response to the

osmotic stress induction, showing lower root curvature angles than the *quad* roots (10º

and 16º respectively; Fig. 2b). Consistently, a hypersensitive hydrotropic response was

observed in *BRL3ox* (30º) compared to WT roots (24º) (Fig. 2b), whereas exogenous

application of the BR-synthesis inhibitor brassinazole [52] reverted the hydrotropic

response (Supplementary Fig. 2). A drought multi-trait matrix was generated for all the

BR receptor mutants analyzed in our study (Fig. 2d; Supplementary Table 1), which

highlights that BRL3/BRL1/BAK1 signalosome in the vascular tissues enhances

hydrotropic sensitivity.

BRL3 receptors confer drought resistance without penalizing overall plant growth

To investigate if the impaired responses to abiotic stress observed in root seedling were

preserved in mature plants, we next analyzed the phenotypes of plants exposed to severe

drought. After 12 days of withholding water, dramatic symptoms of drought stress were

observed in WT, *brl1brl3* and *brl1brl3bak1*, whereas other BR mutants showed a

remarkable drought resistance. In particular, *bak1*, *bri1*, *bri1brl1brl3* and *quad* mutant

plants were the most resistant to the severe water-withholding regime (Fig. 3a). As these

mutants exhibit some degree of dwarfism (Fig. 3a), survival rates after re-watering

drought-treated plants confirmed the resistance phenotypes (Fig. 3b). To correct for the

delayed plant growth of the BR deficient mutants, plants were further submitted to a

time-course drought stress in which water-use, photosynthesis and transpiration

parameters were monitored under similar relative soil water content (Fig. 3c-e). The BR

loss-of-function mutant plants *bri1*, *bri1brl1brl3* and *quad* took 15 days to use 70% of

the water available (field capacity) during the drought stress period, compared to the 9

days taken by WT (Fig. 3c). At similar relative soil water content, BR mutants sustained

higher transpiration, photosynthetic activity (Fig. 3d,e) and leaf Relative Water Content

(RWC) during the drought period (Supplementary Fig. 3). Altogether our results

indicate that the dwarf BR receptor mutant plants are more resistant while consuming

less water, likely through avoiding drought (Fig. 3f).

Strikingly, we found that *BRL3ox* plants were more resistant to severe drought stress

than WT plants as shown by increased survival rates (Fig. 3a,b). Plants with increased

BRL3 receptors reached the 70% soil field capacity two days later than the WT plants

(Fig. 3c). Interestingly photosynthesis rate was decreased in the *BRL3ox* at basal

conditions, yet it was more stable than the WT during the drought period, together with

transpiration. (Fig. 3d,e and Supplementary Fig. 3). These results support that BRL3

pathway actively promotes drought tolerance without penalizing plant growth (Fig. 3f).

BRL3 receptors control the production of osmoprotectant metabolites

To further investigate the cause of the drought tolerance conferred by BRL3 receptors,

we performed metabolite profiling of *BRL3ox* plants and compared it to WT and *quad*

plants in a time-course drought experiment. Roots were separated from shoots to

address possible changes in metabolite mobilization from source to sink tissues. The

complete metabolic fingerprints are provided in Supplementary Fig. 4 and 5 and

Supplementary Table 2 and 3. Metabolite profiling of *BRL3ox* mature plants grown in

control conditions (time 0) revealed an increment in the production of osmoprotectant

metabolites in the plant. Both shoots (Fig. 4a) and roots (Fig. 4c) of *BRL3ox* plants

exhibited a metabolic signature enriched in proline and sugars, which have been

previously reported to confer resistance to drought [6,53-57] suggesting that BRL3 receptor

promotes priming [58,59]. Importantly, those metabolites were decreased in *quad* mutant

plants (Fig. 4a,c).

The sugars fructose, glucose, galactinol, galactose, maltose and raffinose were

accumulated in *BRL3ox* shoots (Fig. 4a). Conversely, sucrose, trehalose, *myo*-inositol

and maltose were accumulated in *BRL3ox* roots, while glucose levels were decreased

(Fig. 4c). The fact of increased glucose and fructose levels in the shoots while higher

levels of sucrose levels in the roots suggest that BRL3 pathway promotes sugar

mobilization to the roots (Fig. 4a). In response to drought, the dynamics of each

metabolite was analyzed along a time-course [60]. A rapid mobilization of osmoprotectant

metabolites was observed in *BRL3ox* plants (Fig. 4b,d). In shoots exposed to drought,

proline showed the highest levels respect WT along the entire drought time course (Fig.

4b,f). Instead, glucose, galactose and *myo*-inositol increased at similar or lower rates in

BRL3ox than in WT plant shoots (Fig. 4e,g). However in roots, an accumulation of

trehalose, sucrose, proline and raffinose was observed in *BRL3ox* under drought stress

time-course (Fig. 4d), following a steeper exponential dynamics than in WT plants (Fig.

4h). Additionally glucose, galactose, fructose and *myo*-inositol were linearly increased

in WT while exponentially accumulated in *BRL3ox* roots (Fig. 4j). Interestingly, along

the drought-time course these metabolite levels were decreased in the *quad* mutant

plants, compared to WT (Supplementary Fig. 6). Together, the drought time course

metabolite profiling uncovers a key role of BR receptors in promoting sugar metabolism

while supports that BRL3 triggers the accumulation of osmoprotectant metabolites in

the root, promoting growth under drought conditions.

BRL3 transcriptional control of metabolite production in response to drought

The osmolyte accumulation observed in roots of plants with increased BRL3 receptor

levels suggested a faster sugar mobilization from shoot to roots. We next investigated

whether those metabolic pathways were regulated transcriptionally by BRL3 in the root.

RNAseq of *BRL3ox* roots revealed 759 genes deregulated at basal conditions (214 up

and 545 down; FC>1.5, FDR<0.05; Fig. 5a; Supplementary Table 4) and 1068 genes

deregulated in drought conditions (378 up and 690 down; FC>1.5, FDR<0.05; Fig. 5a,

Supplementary Table 4). Both sets of deregulated genes overlapped only a 13% (Fig.

5c). In control conditions, a high proportion of deregulated genes were annotated in GO

category of response to water stress, oxygen-containing compounds (ROS) and

response to ABA (Fig. 5d, 5e, Supplementary Table 5). We next classified the genes in

the response to stress category, which included classical drought stress markers such as

RD22 or *RAB18* already upregulated in basal conditions (Fig 5b). The enrichment of

response to hormone genes indicated altered hormonal responses in *BRL3ox* plants (Fig

5d). Further analyses of specific hormonal pathways revealed ABA and jasmonic acid

(JA) as the most altered responses (Supplementary Fig. 7). Repression of the JA

signaling and synthesis genes resulted in decreased levels of JA in basal conditions

(Supplementary Fig. 7). An exclusive GO category being overrepresented in drought

was cell wall organization. Enzymes that down-regulated in *BRL3ox* under drought

(Fig. 5e) included a nourished group of expansins, pectin-lyases and cell wall

invertases, suggesting that *BRL3ox* plants are able to guarantee a sugar supply to the

root and keep cell wall integrity upon drought (Supplementary Fig. 8).

Analysis of transcription factors (TF) showed 29 TFs with differential response to

drought between *BRL3ox* and WT roots. Interestingly, the drought-responsive TF *RD26*

showed and enhanced response in *BRL3ox* roots during stress, whereas several vascular-

specific TFs upon drought remained repressed (Supplementary Fig. 9).

Given that the native expression domain of BRL3 receptor is located at phloem-pole

pericycle and the enriched localization in vascular tissues when overexpressed [41], the

spatial distribution of deregulated genes in the root tissues was analyzed in our RNAseq

dataset [61]. The deregulated genes were preferentially enriched in vascular tissues, such

as phloem and pericycle, but also in lateral root primordia and root hair cells, especially

during the drought stress (Fig. 6a,b, See methods). Among phloem-enriched genes we

found two galactinol synthases (*GolS*) and trehalose phosphatases (*TPPs*) with

increased expression in *BRL3ox* roots both, in basal conditions and drought (Fig. 6 c, d).

These enzymes drive the synthesis of the osmoprotectant metabolites trehalose, myo-

inositol and raffinose we found accumulated in *BRL3ox* roots. Together, these results

support the importance of BRL3 signaling in the phloem to sustain drought resistance.

Furthermore, by statistical analysis a significant link between the whole transcriptomic

and metabolomic signatures was found, both in basal conditions and under drought

(p=0.017 and p=0.001 respectively, methods), indicating that the metabolic signature of

BRL3ox plants is transcriptionally controlled. We used the metabolic and transcriptomic

signatures to identify deregulated metabolic pathways using Paintomics [62]. The analysis

revealed a constitutive control of sucrose metabolism by the vascular BRL3 pathway

that sustained during drought stress. We also found that BRL3 controls the galactose

metabolism under drought, which includes the Raffinose Family of Oligosaccharides

(RFOs) synthesis pathway (Supplementary Fig. 10, Supplementary Table 8).

Collectively, these results support that BRL3 receptor promotes drought tolerance,

mainly controlling sugar metabolism.

DISCUSSION

Our study address the importance of vascular BRL3 signaling in the transcriptional

control of metabolic pathways responsible for the production of osmoprotectant

metabolites in the leaves, that are mobilized to the roots preventing growth arrest during

severe drought periods (Supplementary Fig. 11). Spatial BR signaling has been shown

to contribute to stem cell replenishment in response to genotoxic stress[36,37], while the

present study uncovers a role for vascular-enriched BRL3 receptors in promoting

growth during drought periods. Together, these results show the importance of the

spatial regulation of BR signaling in plant stress responses and open a new window for

the study of drought tolerance with cell-specific resolution in plants.

Exogenous application of BR compounds has been widely used in agriculture to extend

growth under different abiotic stresses [27,30,63], yet how these molecules precisely are

able to activate growth in challenging conditions remains largely unknown. The

analysis of BR signaling and synthesis mutant plants in responses to stress failed to

provide a linear picture for their involvement in drought stress adaptation. The

overexpression of the canonical BRI1 receptor pathway and the BR-biosynthesis gene

DWF4 can both confer abiotic stress resistance [29,43] whereas *BRI1* loss-of-function

mutants also showed drought stress resistance [31]. However, increased levels of BR-

regulated transcription factors trigger antagonistic effects in drought stress responses

[31,64], thus depicting a complex scenario for BRs role on abiotic stress. Given the

spatiotemporal regulation of the BR-signaling components [45] and the complexity of

drought traits [9], it is plausible to hypothesize that drought traits are under control of cell-

type specific BR-signaling.

Our study unveils that the BR family of receptors, in addition to promoting growth,

guides the phenotypic adaptation responses to a myriad of drought stress traits. The

drought resistance phenotypes of BR loss-of-function mutants (Fig. 3a) are likely

caused by drought avoidance linked to growth arrest and a stress-insensitivity that

maintains transpiration, leaf water status and photosynthesis (Fig. 3e,f). The reduced

levels of ABA and canonical stress related metabolites together with the down-

regulation of drought stress genes further support the *quad* plants insensitivity to stress

(Supplementary Figs. 6 and 7). We found the drought-response RD26 transcription

factor expression enhanced in *BRL3ox* roots upon drought (Supplementary Fig. 9).

RD26 was shown to antagonize with the BRs canonical TF BES1 [31], thus suggesting

alternative pathways being activated by BRL3. The alternative pathways may be

derived of a spatial specialization of the BR functions within the root. Indeed we found

enriched transcriptional responses in the vascular tissues of the *BRL3ox* roots, especially

within phloem-related cell types (Fig. 6a,b). The native expression domain of BRL3 and

the metabolic signature profile found in *BRL3ox* supports the hypothesis of its

implication in phloem loading during drought. Moreover metabolic enzymes implicated

in trehalose and RFO metabolism were enriched in vascular tissues and upregulated in

BRL3ox roots (Fig. 6c,d). Thus, BRL3 function is not limited to the phloem

loading/unloading but also directly controlling metabolic pathways. Such is the case of

trehalose phosphate phosphatase family (*TPPs*) [65-67] and galactinol synthase 2 (*GolS2*)

[68-70], involved in trehalose and RFOs synthesis respectively, which have been described

to impact drought responses. In addition to the vascular findings, our analysis uncover

the control of non-vascular enzymes important in metabolism and drought responses, i.e

the control of hexokinases such as *HXK3* or *HKL1* and sucrose synthases, *SUS3* and

SPS2F or the constitutive downregulation of proline dehydrogenase genes, as the early

response to dehydration 5 (*ERD5*) [71,72]. In light of our findings, we propose that

vascular BRL3 receptors action may partially occur independently of the canonical

growth-promoting BRI1 pathway, given the drought hypersensitivity found in *Bes1-D*

gain-of-function mutants [31].

Our data further suggest that *BRL3ox* plants accumulate sugars in sink tissues to provide

plant root growth to escape drought by searching for water in the soil. Supporting these

findings, reduced photosynthesis was observed in the well-watered *BRL3ox* plant shoots

(Fig. 3e). These results, together with the higher levels of sucrose levels in roots over

shoots (Fig. 4a) suggest that the BRL3 pathway promotes sugar mobilization from the

leaves (source) to the roots (sink). Previous work reported that BRs promote the flow of

assimilates from source to sink via the vasculature [73] and sucrose phloem unloading [74]

in crops.

In control conditions *BRL3ox* plants exhibited a metabolic signature enriched in proline

and sugars. Proline and sugars accumulation classically correlates with drought stress

tolerance, osmolyte, ROS scavenger and chaperone functions [6,53-57] suggesting that

BRL3 receptor promotes priming [58,59]. In addition, *BRL3ox* plants also accumulated

succinate, fumarate and malate. Importantly, all these metabolites were decreased in

quad mutant plants. Altogether, these metabolic data uncover a role of BR signaling in

promoting the tricarboxylic acid (TCA) cycle, sugar and amino acid metabolism [75].

In drought stress conditions, *BRL3ox* shoots displayed increased levels of the amino

acids proline, GABA and tyrosine.   In the *BRL3ox* roots, trehalose, sucrose, *myo*-

inositol, raffinose and proline were the most accumulated metabolites along the stress

time course, and all have previously been linked to drought resistance [53-55]. In addition,

the RFO metabolites raffinose and *myo*-inositol, involved in membrane protection and

radical scavenging [76], were increased in *BRL3ox,* yet reduced in *quad* mutant roots

under drought.

Our data thus indicate that *BRL3ox* roots are loaded with osmoprotectant metabolites,

thus better prepared to alleviate drought stress, via a phenomenon previously referred to

as priming [58,59]. Further supporting our hypothesis, the BRL3 signalosome is enriched in

abiotic stress proteins [41]. Altogether these data suggest that sensing drought stress

responses is correlated with BRL3 receptor levels in the root vasculature, especially in

the phloem [77] is key in contributing to greater survival rates in *BRL3ox* shoots. Future

cell-type specific engineering of signaling cascades stands as promising strategy to

surpass the growth arrest caused by drought stress.

METHODS

Plant materials

Seeds were sterilized with 35% NaClO for 5 min and washed 5 times for 5min with sterile dH$_2$O. Sterile seeds were vernalized 48 h at 4ºC and grown in half-strength agar Murashige and Skoog (MS1/2) media with vitamins and without sucrose. Plates were grown vertically in long day (LD) conditions (16 h of light / 8 h of dark; 22°C, 60% relative humidity). Genotypes used in this study: Columbia-0 WT (Col-0 WT), *brl1-1brl3-1 (brl1brl3)*, *bak1-3 (bak1)*, *bri1-301 (bri1)*, *bri1-301brl1-1brl3-1 (bri1brl1brl3)*, *bri1-301bak1-3brl1-1brl3-1 (quad)* and *35S:BRL3-GFP (BRL3ox)* [41]. DNA rapid extraction protocol [78] was used for all the plant genotyping experiments. Primers used for genotyping the BR mutant plants are described in Table 1.

Brassinolide and sorbitol sensitivity assays in roots

For hormone treatments, seeds were continuously grown in concentrations series of brassinolide (BL,Wako, Japan). For sorbitol assays, three-day-old seedlings were transferred to either control or 270mM sorbitol media four additional days. Seven-day-old seedlings root length was measured with Image J (http://rsb.info.nih.gov/ij/) and compared to automatically acquired data with MyROOT [79] software (Supplementary Fig. 12). Four-day-old roots grown in control conditions or in 24h of sorbitol treatment were stained with Propidium iodide (10ug/ml, PI, Sigma) that stains the cell wall (control) and DNA in the nuclei upon cell death (sorbitol). Images were acquired with confocal microscope (FV1000 Olympus). Cell death damage in primary roots was measured in a window of 500μm from QC in the middle root longitudinal section (Image J). As arbitrary setting to measure the stained area, a color threshold ranging from 160 to 255 in brightness was selected.

Root hydrotropism

Seedlings were germinated in MS1/2 for six days. The lower part of the plates was removed and MS1/2 with 270mM sorbitol was added, to simulate a situation with reduced water availability. 1μM of BRZ_{220} [80] was added to sorbitol media when indicated. The media was placed in 45-degree angle to scape gravitropism effect as previously described [50,51]. Root curvature angles were measured and analyzed with Image J software (http://rsb.info.nih.gov/ij/).

Drought stress for scoring plant survival

One-week-old seedlings grown in MS1/2 agar plates were transferred individually to pots containing 30±1 g. of substrate (plus 1:8 v/v vermiculite and 1:8 v/v perlite). For each biological replicate, 40 plants of each genotype were grown in LD conditions for three weeks. Three-week-old plants were subjected to severe drought stress by withholding water during 12 days followed by re-watering. After the 7 days of recovery period, the surviving plants were pictured and manually counted (two-sided chi-squared test, $P < 0.01$).

Metabolite profiling analyses

One-week-old seedlings were placed in individual pots with 30g of autoclaved soil and grown in LD photoperiodic conditions. After 3 weeks growing, half of plants were subjected to a severe drought period of six days (withholding water) and other half were watered normally (well-watered control conditions). A total of five biological replicates were collected every 24h during the time course (from day 0 to day 6) both in drought and watered conditions for each genotype (WT, *quad* and *BRL3ox*). Four independent

plants were bulked in each biological replicate. Roots were manually separated from

shoots. Root samples were aliquoted in 20mg and shoot samples in 50mg (exact weight

was annotated for data normalization). Primary metabolites extraction was carried as

previously described [81]. Samples were transferred into glass vials specific for injection

in GC-TOF-MS. The GC-TOF-MS system comprised a CTC CombiPAL autosampler,

an Agilent 6890N gas chromatograph, and a LECO Pegasus III TOF-MS running in EI+

mode. Metabolites were identified in comparison to database entries of authentic

standards [82]. Chromatograms were evaluated using Chroma TOF 1.0 (Leco) Pegasus

software for peak identification and correction of RT. Mass spectra were evaluated

TagFinder 4.0 software [83] for metabolite annotation and quantification (peak area

measurements). The resulting data matrix was normalized using an internal standard

Ribitol in 100% Methanol (20:1) followed by normalization with the fresh weight of

each sample. Control well-watered conditions (day 0) metabolomic data was analyzed

with a two-tailed t-test, p-value<0.05 (no multiple testing correction). Time course

analysis data was analyzed with R software, using maSigPro package [60]. Briefly the

profile of each metabolite in each condition was fitted to a polynomial model of

maximum degree 3. Statistical comparison of curves between genotypes was done

taking into account the fitting value, then the p-value was corrected (Benjamini-

Hochberg method) and significant metabolites (p-value < 0.05) having a differential

profile between genotypes were plotted to visualize their behavior under the drought

time course. Clustering analysis was done by maSigPro package itself, using the *hclust*

R core function.

Transcriptomic profiling analysis

For microarray analysis, a drought stress time-course analysis was carried in WT and

quad mutant three-week-old mature plants. Entire plants were collected every 48 hours

during the time course (Day 0, Day 2 and Day 4) under drought stress and control

conditions. Two biological replicates composed of 5 independent rosettes were

collected. RNA was extracted with Plant Easy Mini Kit (Qiagen) and quality checked

by Bioanalyser. Genome-Wide Microarray platform (Dual color, Agilent) was done by

swapping color hybridization of each biological replicate (Cy3 and Cy5). Statistical

analysis was performed with the package "limma" [84], the "mle2/"normexp" background

correction method was used. Quantile normalization was used to normalize different

microarrays then a Bayes test was used to identify differentially expressed probes. The

results were filtered for adjusted p-value<0.05 (after Benjamini-Hochberg correction)

and Log2 FC >|1.5|. For RNAseq analysis, three-week-old roots were detached from

mature plants grown on soil at control conditions and exposed to 5 days of drought.

RNA was extracted as described above. Stranded cDNA libraries were prepared with

TruSeq Stranded mRNA kit (Illumina). Single-end sequencing, with 50-bp reads, was

performed in an Illumina HiSeq500 sequencer, at a minimum sequencing depth of 21M.

Reads were trimmed 5bp at their 3' end and quality filtered then mapped against the

TAIR10 genome with "HISAT2". Mapped reads were quantified at gene level with

"HtSeq". For differential expression samples were TMM normalized and statistical

values calculated with "EdgeR" package in R. Results were filtered for adjusted p-value

(FDR) <0.05 and FC >|2|. Heatmap representations were performed in R with the

heatmap.2 function implemented in the "gplots" package.

Statistical methods and metabolic and transcriptomic data integration.

For root tissue enrichment analysis, deregulated genes were queried against available

tissue-enriched genes lists [61]. For each tissue, a 2x2 contingency table was constructed

counting for the number of deregulated genes enriched and not enriched in the tissue

and also for the number of not deregulated genes (for either FDR>0.05 or logFC >/< in

the RNAseq gene universe) enriched and not enriched in the tissue. Statistical value of

the enrichment was obtained with one-sided Fisher's test. To statistically evaluate the

influence of transcriptomic changes in the metabolic signature, both deregulated

enzymes and metabolites were queried in an annotation file for Arabidopsis thaliana's

metabolic   reactions,   which   included   merged   data   from   KEGG

(http://www.genome.jp/kegg/) and BRENDA (www.brenda-enzymes.org) databases.

Then same approach of constructing a 2x2 contingency table was taken. Significant and

not significant metabolites annotated in the database were matched with differential or

not differential expressed genes annotated in the database. The statistical value of the

association between regulated metabolites and genes was obtained through a two-sided

Fisher's exact test. Genes and metabolites were mapped into the KEGG pathways using

the   PaintOmics3   (http://bioinfo.cipf.es/paintomics/)   according   the   developer

instructions [62].

Physiological analysis and Chlorophyll fluorescence parameters

One-week-old seedlings were placed in individual pots and watered with the same

volume of a modified one-fifth-strength Hoagland solution. Pots were weighed daily

during the experiment. Well-watered control plants were grown at 100% of field

capacity (0% of water loss). Time-course drought stress assay started by withholding

nutrient solution until reaching 25, 50, 60 and 70% of water loss. Photosynthesis (A)

and transpiration (E) were measured in control and drought plants at those time points.

Four plants of each genotype were harvested at 0, 50 and 70% of water loss for biomass,

water content and hormone analyses. Drought experiments were repeated three times

and at least 4 plants per genotype and treatment were used in each experiment. RWC

was calculated according to the formula: RWC (%) = [(FW-DW) / (TW-DW)] x 100.

Plant hormones cytokinins (*trans*-zeatin), gibberellins (GA1, GA4 and GA3), indole-3-

acetic acid (IAA), abscisic acid (ABA), salicylic acid (SA), jasmonic acid (JA) and the

ethylene precursor 1-aminocyclopropane-1-carboxylic acid (ACC) were analyzed in the

leaves as previously described [85].

ACKNOWLEDGEMENTS

We acknowledge financial support from the Spanish Ministry of Economy and

Competitiveness, through the "Severo Ochoa Programme for Centres of Excellence in

R&D" 2016-2019 (SEV-2015-0533)", the CERCA Programme from the Generalitat de

Catalunya, and the Agencia de Gestió d'Ajuts Universitaris i de Recerca (2014 SGR

1406). N.F. is funded by Fundación Renta Corporación and by EMBO short-term

postdoctoral fellowship (ASTF 422-2015). F.L.E and D.B are funded by BIO2013-

43873 grant from the Spanish Ministry of Economy and Competitiveness in A.I.C.-D.

laboratory. CMA, AA and FPA acknowledge financial support from the Spanish

MINECO-FEDER (project AGL2014-59728-R) and from the European Union's

Seventh Framework Programme for research technological development and

demonstration under grant agreement no. 289365 (project ROOTOPOWER). SO was

supported by Ministry of Science and Innovation and University of Malaga (Spain)

through the grant Ramón and Cajal program (Sonia Osorio, RYC-09170). M.B and

J.L.R are supported by grant BFU2014-58289-P from the Spanish Ministry of Economy

and Competitiveness. A.C.D. − A.C. collaboration was funded by the European

Regional Development Funds and Marie Curie IRSES Project DEANN (PIRSES-GA-

2013-612583). A.I.C.-D. is a recipient of a BIO2013-43873 grant from the Spanish

Ministry of Economy and Competitiveness and European Research Council, ERC

Consolidator Grant (ERC-2015-CoG – 683163).

521 AUTHOR CONTRIBUTIONS

A.I.C.-D conceived, designed and supervised the study. N.F., F.L.E, D.B., C.M.A. and

F.P.A. performed the genetics and stress phenomic experiments and data analyses. N.F.,

F.L.E., M.B., J.L.R. and A.C.D. designed the genome-wide experiments. N.F., F.L.E.,

A.C. and A.C.D. performed the genome-wide experiments and data analysis. N.F.,

F.L.E., T.T., S.O., and A.R.F. carried out the metabolite and data analysis. A.A.

performed the hormone profiling experiments. N.F. and A.C.D. wrote the manuscript.

All authors discussed the results and the manuscript.

Table 1. Primers used for genotyping

primer	Sequence (5' - 3')	Mutant
bri2.11	GTGAGAAACGAAGGTGGAACAGACTGCAG	*brl1-1*
bri2.12	CTTCTTGGCATGGATACGGGAGGTAATGG	*brl1-1*
JMLB	GGCAATCAGCTGTTGCCCGTCTCACTGGTG	*brl1-1*
bri4.2	TTTAGGGTGAGCATGAGATCTCGTGGGCCG	*brl3-1*
bri4.3	GAAATCCCTGTAGGAATCGGAAAGCTTGAG	*brl3-1*
JMRB	GCTCATGATCAGATTGTCGTTTCCCGCCTT	*brl3-1*
bak1-3 F	GCACTGAAAAACAGTTTAGC	*bak1-3*
bak1-3 R	GATGCAGGAAGGGGAGTCAACTTGGTG	*bak1-3*
T-DNA R	GCGTGGACCGCTTGCTGCAACT	*bak1-3*
bri1-301 F	GGAAACCATTGGGAAGATCA	*bri1-301* *
bri1-301 R	GCTGTTTCACCCATCCAA	*bri1-301* *

531 *PCR product was digested by *DpnII* enzyme (R0543, New England Biolabs)

REFERENCES

Boyer, J. S. Plant productivity and environment. *Science* **218**, 443-448, doi:10.1126/science.218.4571.443 (1982).

Todaka, D., Shinozaki, K. & Yamaguchi-Shinozaki, K. Recent advances in the dissection of drought-stress regulatory networks and strategies for development of drought-tolerant transgenic rice plants. *Front Plant Sci* **6**, 84, doi:10.3389/fpls.2015.00084 (2015).

Cutler, S. R., Rodriguez, P. L., Finkelstein, R. R. & Abrams, S. R. Abscisic acid: emergence of a core signaling network. *Annu Rev Plant Biol* **61**, 651-679, doi:10.1146/annurev-arplant-042809-112122 (2010).

Yoshida, T., Mogami, J. & Yamaguchi-Shinozaki, K. ABA-dependent and ABA-independent signaling in response to osmotic stress in plants. *Curr Opin Plant Biol* **21**, 133-139, doi:10.1016/j.pbi.2014.07.009 (2014).

Graether, S. P. & Boddington, K. F. Disorder and function: a review of the dehydrin protein family. *Front Plant Sci* **5**, 576, doi:10.3389/fpls.2014.00576 (2014).

Seki, M., Umezawa, T., Urano, K. & Shinozaki, K. Regulatory metabolic networks in drought stress responses. *Curr Opin Plant Biol* **10**, 296-302, doi:10.1016/j.pbi.2007.04.014 (2007).

Umezawa, T. *et al.* CYP707A3, a major ABA 8'-hydroxylase involved in dehydration and rehydration response in Arabidopsis thaliana. *Plant J* **46**, 171-182, doi:10.1111/j.1365-313X.2006.02683.x (2006).

Shao, H. B., Chu, L. Y., Jaleel, C. A. & Zhao, C. X. Water-deficit stress-induced anatomical changes in higher plants. *C R Biol* **331**, 215-225, doi:10.1016/j.crvi.2008.01.002 (2008).

Tardieu, F. Any trait or trait-related allele can confer drought tolerance: just design the right drought scenario. *J Exp Bot* **63**, 25-31, doi:10.1093/jxb/err269 (2012).

Marshall, A. *et al.* Tackling drought stress: receptor-like kinases present new approaches. *Plant Cell* **24**, 2262-2278, doi:10.1105/tpc.112.096677 (2012).

Smakowska-Luzan, E. *et al.* An extracellular network of Arabidopsis leucine-rich repeat receptor kinases. *Nature* **553**, 342-346, doi:10.1038/nature25184 (2018).

Clouse, S. D., Langford, M. & McMorris, T. C. A brassinosteroid-insensitive mutant in Arabidopsis thaliana exhibits multiple defects in growth and development. *Plant Physiol* **111**, 671-678 (1996).

Li, J. & Chory, J. A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell* **90**, 929-938 (1997).

Wang, Z. Y., Seto, H., Fujioka, S., Yoshida, S. & Chory, J. BRI1 is a critical component of a plasma-membrane receptor for plant steroids. *Nature* **410**, 380-383, doi:10.1038/35066597 (2001).

Kinoshita, T. *et al.* Binding of brassinosteroids to the extracellular domain of plant receptor kinase BRI1. *Nature* **433**, 167-171, doi:10.1038/nature03227 (2005).

Hothorn, M. *et al.* Structural basis of steroid hormone perception by the receptor kinase BRI1. *Nature* **474**, 467-471, doi:10.1038/nature10153 (2011).

She, J. *et al.* Structural insight into brassinosteroid perception by BRI1. *Nature* **474**, 472-476, doi:10.1038/nature10178 (2011).

18   Karlova, R. *et al.* The Arabidopsis SOMATIC EMBRYOGENESIS
RECEPTOR-LIKE KINASE1 protein complex includes BRASSINOSTEROID-
INSENSITIVE1. *Plant Cell* **18**, 626-638, doi:10.1105/tpc.105.039412 (2006).

19   Li, J. *et al.* BAK1, an Arabidopsis LRR receptor-like protein kinase, interacts
with BRI1 and modulates brassinosteroid signaling. *Cell* **110**, 213-222 (2002).

20   Nam, K. H. & Li, J. BRI1/BAK1, a receptor kinase pair mediating
brassinosteroid signaling. *Cell* **110**, 203-212 (2002).

21   Gou, X. *et al.* Genetic evidence for an indispensable role of somatic
embryogenesis receptor kinases in brassinosteroid signaling. *PLoS Genet* **8**,
e1002452, doi:10.1371/journal.pgen.1002452 (2012).

22   Yin, Y. *et al.* BES1 accumulates in the nucleus in response to brassinosteroids to
regulate gene expression and promote stem elongation. *Cell* **109**, 181-191
(2002).

23   Wang, Z. Y. *et al.* Nuclear-localized BZR1 mediates brassinosteroid-induced
growth and feedback suppression of brassinosteroid biosynthesis. *Dev Cell* **2**,
505-513 (2002).

24   He, J. X., Gendron, J. M., Yang, Y., Li, J. & Wang, Z. Y. The GSK3-like kinase
BIN2 phosphorylates and destabilizes BZR1, a positive regulator of the
brassinosteroid signaling pathway in Arabidopsis. *Proc Natl Acad Sci U S A* **99**,
10185-10190, doi:10.1073/pnas.152342599 (2002).

25   He, J. X. *et al.* BZR1 is a transcriptional repressor with dual roles in
brassinosteroid homeostasis and growth responses. *Science* **307**, 1634-1638,
doi:10.1126/science.1107580 (2005).

26   Yin, Y. *et al.* A new class of transcription factors mediates brassinosteroid-
regulated gene expression in Arabidopsis. *Cell* **120**, 249-259,
doi:10.1016/j.cell.2004.11.044 (2005).

27   Kagale, S., Divi, U. K., Krochko, J. E., Keller, W. A. & Krishna, P.
Brassinosteroid confers tolerance in Arabidopsis thaliana and Brassica napus to
a range of abiotic stresses. *Planta* **225**, 353-364, doi:10.1007/s00425-006-0361-
6 (2007).

28   Krishna, P. Brassinosteroid-Mediated Stress Responses. *J Plant Growth Regul*
22, 289-297, doi:10.1007/s00344-003-0058-z (2003).

29   Sahni, S. *et al.* Overexpression of the brassinosteroid biosynthetic gene DWF4
in Brassica napus simultaneously increases seed yield and stress tolerance. *Sci*
Rep **6**, 28298, doi:10.1038/srep28298 (2016).

30   Feng, Y., Yin, Y. & Fei, S. Down-regulation of BdBRI1, a putative
brassinosteroid receptor gene produces a dwarf phenotype with enhanced
drought tolerance in Brachypodium distachyon. *Plant Sci* **234**, 163-173,
doi:10.1016/j.plantsci.2015.02.015 (2015).

31   Ye, H. *et al.* RD26 mediates crosstalk between drought and brassinosteroid
signalling pathways. *Nat Commun* **8**, 14573, doi:10.1038/ncomms14573 (2017).

32   Zhang, S., Cai, Z. & Wang, X. The primary signaling outputs of brassinosteroids
are regulated by abscisic acid signaling. *Proc Natl Acad Sci U S A* **106**, 4543-
4548, doi:10.1073/pnas.0900349106 (2009).

33   Gui, J. *et al.* OsREM4.1 Interacts with OsSERK1 to Coordinate the Interlinking
between Abscisic Acid and Brassinosteroid Signaling in Rice. *Dev Cell* **38**, 201-
213, doi:10.1016/j.devcel.2016.06.011 (2016).

34   Huang, D., Wu, W., Abrams, S. R. & Cutler, A. J. The relationship of drought-
related gene expression in Arabidopsis thaliana to hormonal and environmental
factors. *J Exp Bot* **59**, 2991-3007, doi:10.1093/jxb/ern155 (2008).

35 Chung, Y., Kwon, S. I. & Choe, S. Antagonistic regulation of Arabidopsis
growth by brassinosteroids and abiotic stresses. *Mol Cells* **37**, 795-803,
doi:10.14348/molcells.2014.0127 (2014).

36 Vilarrasa-Blasi, J. *et al.* Regulation of plant stem cell quiescence by a
brassinosteroid signaling module. *Dev Cell* **30**, 36-47,
doi:10.1016/j.devcel.2014.05.020 (2014).

37 Lozano-Elena, F., Planas-Riverola, A., Vilarrasa-Blasi, J., Schwab, R. & Caño-
Delgado, A. I. Paracrine brassinosteroid signaling at the stem cell niche controls
cellular regeneration. *Journal of Cell Science* **131**, doi:10.1242/jcs.204065
(2018).

38 Fàbregas, N. & Caño-Delgado, A. I. Turning on the microscope turret: a new
view for the study of brassinosteroid signaling in plant development. *Physiol*
Plant **151**, 172-183, doi:10.1111/ppl.12130 (2014).

39 Friedrichsen, D. M., Joazeiro, C. A., Li, J., Hunter, T. & Chory, J.
Brassinosteroid-insensitive-1 is a ubiquitously expressed leucine-rich repeat
receptor serine/threonine kinase. *Plant Physiol* **123**, 1247-1256 (2000).

40 Caño-Delgado, A. *et al.* BRL1 and BRL3 are novel brassinosteroid receptors
that function in vascular differentiation in Arabidopsis. *Development* **131**, 5341-
5351, doi:10.1242/dev.01403 (2004).

41 Fàbregas, N. *et al.* The brassinosteroid insensitive1-like3 signalosome complex
regulates Arabidopsis root development. *Plant Cell* **25**, 3377-3388,
doi:10.1105/tpc.113.114462 (2013).

42 Belkhadir, Y. & Jaillais, Y. The molecular circuitry of brassinosteroid signaling.
New Phytol **206**, 522-540, doi:10.1111/nph.13269 (2015).

43 Eremina, M. *et al.* Brassinosteroids participate in the control of basal and
acquired freezing tolerance of plants. *Proc Natl Acad Sci U S A* **113**, E5982-
E5991, doi:10.1073/pnas.1611477113 (2016).

44 Nakamura, A. *et al.* The role of OsBRI1 and its homologous genes, OsBRL1
and OsBRL3, in rice. *Plant Physiol* **140**, 580-590, doi:10.1104/pp.105.072330
(2006).

45 Vragovic, K. *et al.* Translatome analyses capture of opposing tissue-specific
brassinosteroid signals orchestrating root meristem differentiation. *Proc Natl*
Acad Sci U S A **112**, 923-928, doi:10.1073/pnas.1417947112 (2015).

46 González-García, M. P. *et al.* Brassinosteroids control meristem size by
promoting cell cycle progression in Arabidopsis roots. *Development* **138**, 849-
859, doi:10.1242/dev.057331 (2011).

47 Albrecht, C., Russinova, E., Kemmerling, B., Kwaaitaal, M. & de Vries, S. C.
Arabidopsis SOMATIC EMBRYOGENESIS RECEPTOR KINASE proteins
serve brassinosteroid-dependent and -independent signaling pathways. *Plant*
Physiol **148**, 611-619, doi:10.1104/pp.108.123216 (2008).

48 Hacham, Y. *et al.* Brassinosteroid perception in the epidermis controls root
meristem size. *Development* **138**, 839-848, doi:10.1242/dev.061804 (2011).

49 Duan, Y. *et al.* An endoplasmic reticulum response pathway mediates
programmed cell death of root tip induced by water stress in Arabidopsis. *New*
Phytol **186**, 681-695, doi:10.1111/j.1469-8137.2010.03207.x (2010).

50 Takahashi, N., Goto, N., Okada, K. & Takahashi, H. Hydrotropism in abscisic
acid, wavy, and gravitropic mutants of Arabidopsis thaliana. *Planta* **216**, 203-
211, doi:10.1007/s00425-002-0840-3 (2002).

51 Galvan-Ampudia, C. S. *et al.* Halotropism is a response of plant roots to avoid a
saline environment. *Curr Biol* **23**, 2044-2050, doi:10.1016/j.cub.2013.08.042
(2013).
52 Nagata, N., Min, Y. K., Nakano, T., Asami, T. & Yoshida, S. Treatment of dark-
grown Arabidopsis thaliana with a brassinosteroid-biosynthesis inhibitor,
brassinazole, induces some characteristics of light-grown plants. *Planta* **211**,
781-790 (2000).
53 Singh, T. N., Aspinall, D. & Paleg, L. G. Proline accumulation and varietal
adaptability to drought in barley: a potential metabolic measure of drought
resistance. *Nat New Biol* **236**, 188-190 (1972).
54 Szabados, L. & Savoure, A. Proline: a multifunctional amino acid. *Trends Plant*
Sci **15**, 89-97, doi:10.1016/j.tplants.2009.11.009 (2010).
55 Krasensky, J. & Jonak, C. Drought, salt, and temperature stress-induced
metabolic rearrangements and regulatory networks. *J Exp Bot* **63**, 1593-1608,
doi:10.1093/jxb/err460 (2012).
56 Urano, K. *et al.* Characterization of the ABA-regulated global responses to
dehydration in Arabidopsis by metabolomics. *Plant J* **57**, 1065-1078,
doi:10.1111/j.1365-313X.2008.03748.x (2009).
57 Durand, M. *et al.* Water Deficit Enhances C Export to the Roots in Arabidopsis
thaliana Plants with Contribution of Sucrose Transporters in Both Shoot and
Roots. *Plant Physiology* **170**, 1460-1479, doi:10.1104/pp.15.01926 (2016).
58 Conrath, U., Pieterse, C. M. & Mauch-Mani, B. Priming in plant-pathogen
interactions. *Trends Plant Sci* **7**, 210-216 (2002).
59 Ding, Y., Fromm, M. & Avramova, Z. Multiple exposures to drought 'train'
transcriptional responses in Arabidopsis. *Nat Commun* **3**, 740,
doi:10.1038/ncomms1732 (2012).
60 Conesa, A., Nueda, M. J., Ferrer, A. & Talon, M. maSigPro: a method to
identify significantly differential expression profiles in time-course microarray
experiments. *Bioinformatics* **22**, 1096-1102, doi:10.1093/bioinformatics/btl056
(2006).
61 Brady, S. M. *et al.* A High-Resolution Root Spatiotemporal Map Reveals
Dominant Expression Patterns. *Science* **318**, 801 (2007).
62 García-Alcalde, F., García-López, F., Dopazo, J. & Conesa, A. Paintomics: a
web based tool for the joint visualization of transcriptomics and metabolomics
data. *Bioinformatics* **27**, 137-139, doi:10.1093/bioinformatics/btq594 (2011).
63 Shakirova, F. *et al.* Involvement of dehydrins in 24-epibrassinolide-induced
protection of wheat plants against drought stress. *Plant Physiology and*
Biochemistry **108**, 539-548, doi:https://doi.org/10.1016/j.plaphy.2016.07.013
(2016).
64 Chen, J. *et al.* Arabidopsis WRKY46, WRKY54, and WRKY70 Transcription
Factors Are Involved in Brassinosteroid-Regulated Plant Growth and Drought
Responses. *The Plant Cell* **29**, 1425 (2017).
65 Ge, L.-F. *et al.* Overexpression of the trehalose-6-phosphate phosphatase gene
OsTPP1 confers stress tolerance in rice and results in the activation of stress
responsive genes. *Planta* **228**, 191-201, doi:10.1007/s00425-008-0729-x (2008).
66 Jang, I.-C. *et al.* Expression of a Bifunctional Fusion of the
Escherichia coli Genes for Trehalose-6-Phosphate
Synthase and Trehalose-6-Phosphate Phosphatase in Transgenic Rice Plants
Increases Trehalose Accumulation and Abiotic Stress Tolerance without
Stunting Growth. *Plant Physiology* **131**, 516 (2003).

67   Nuccio, M. L. *et al.* Expression of trehalose-6-phosphate phosphatase in maize
ears improves yield in well-watered and drought conditions. *Nature*
Biotechnology **33**, 862, doi:10.1038/nbt.3277
https://www.nature.com/articles/nbt.3277 - supplementary-information (2015).
68   Gu, L. *et al.* ZmGOLS2, a target of transcription factor ZmDREB2A, offers
similar protection against abiotic stress as ZmDREB2A. *Plant Molecular*
Biology **90**, 157-170, doi:10.1007/s11103-015-0403-1 (2016).
69   Himuro, Y. *et al.* Arabidopsis galactinol synthase AtGolS2 improves drought
tolerance in the monocot model Brachypodium distachyon. *Journal of Plant*
Physiology **171**, 1127-1131, doi:https://doi.org/10.1016/j.jplph.2014.04.007
(2014).
70   Selvaraj, M. G. *et al.* Overexpression of an Arabidopsis thaliana galactinol
synthase gene improves drought tolerance in transgenic rice and increased grain
yield in the field. *Plant Biotechnology Journal* **15**, 1465-1477,
doi:10.1111/pbi.12731 (2017).
71   Kiyosue, T., Yoshiba, Y., Yamaguchi-Shinozaki, K. & Shinozaki, K. A nuclear
gene encoding mitochondrial proline dehydrogenase, an enzyme involved in
proline metabolism, is upregulated by proline but downregulated by dehydration
in Arabidopsis. *The Plant Cell* **8**, 1323 (1996).
72   Nanjo, T. *et al.* Antisense suppression of proline degradation improves tolerance
to freezing and salinity in Arabidopsis thaliana. *FEBS Letters* **461**, 205-210,
doi:10.1016/S0014-5793(99)01451-9 (1999).
73   Wu, C.-y. *et al.* Brassinosteroids Regulate Grain Filling in Rice. *The Plant Cell*
20, 2130-2145, doi:10.1105/tpc.107.055087 (2008).
74   Xu, F., Xi, Z.-m., Zhang, H., Zhang, C.-j. & Zhang, Z.-w. Brassinosteroids are
involved in controlling sugar unloading in Vitis vinifera 'Cabernet Sauvignon'
berries during véraison. *Plant Physiology and Biochemistry* **94**, 197-208,
doi:https://doi.org/10.1016/j.plaphy.2015.06.005 (2015).
75   Schroder, F. *et al.* Consequences of induced brassinosteroid deficiency in
Arabidopsis leaves. *BMC Plant Biol* **14**, 309, doi:10.1186/s12870-014-0309-0
(2014).
76   Nishizawa, A., Yabuta, Y. & Shigeoka, S. Galactinol and raffinose constitute a
novel function to protect plants from oxidative damage. *Plant Physiol* **147**,
1251-1263, doi:10.1104/pp.108.122465 (2008).
77   Salazar-Henao, J. E., Lehner, R., Betegon-Putze, I., Vilarrasa-Blasi, J. & Cano-
Delgado, A. I. BES1 regulates the localization of the brassinosteroid receptor
BRL3 within the provascular tissue of the Arabidopsis primary root. *J Exp Bot*
67, 4951-4961, doi:10.1093/jxb/erw258 (2016).
78   Aljanabi, S. M. & Martinez, I. Universal and rapid salt-extraction of high quality
genomic DNA for PCR-based techniques. *Nucleic Acids Res* **25**, 4692-4693
(1997).
79   Betegón-Putze, I., González, A., Sevillano, X., Blasco-Escámez, D. & Caño-
Delgado, A. I. MyROOT: A novel method and software for the semi-automatic
measurement of plant root length. *bioRxiv* (2018).
80   Asami, T. *et al.* Characterization of brassinazole, a triazole-type brassinosteroid
biosynthesis inhibitor. *Plant Physiol* **123**, 93-100 (2000).
81   Lisec, J., Schauer, N., Kopka, J., Willmitzer, L. & Fernie, A. R. Gas
chromatography mass spectrometry-based metabolite profiling in plants. *Nat*
Protoc **1**, 387-396, doi:10.1038/nprot.2006.59 (2006).

82 Kopka, J. *et al.* GMD@CSB.DB: the Golm Metabolome Database.
Bioinformatics **21**, 1635-1638, doi:10.1093/bioinformatics/bti236 (2005).
83 Luedemann, A., Strassburg, K., Erban, A. & Kopka, J. TagFinder for the
quantitative analysis of gas chromatography--mass spectrometry (GC-MS)-
based metabolite profiling experiments. *Bioinformatics* **24**, 732-737,
doi:10.1093/bioinformatics/btn023 (2008).
84 Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-
sequencing and microarray studies. *Nucleic Acids Res* **43**, e47,
doi:10.1093/nar/gkv007 (2015).
85 Albacete, A. *et al.* Hormonal changes in relation to biomass partitioning and
shoot growth impairment in salinized tomato (Solanum lycopersicum L.) plants.
Journal of Experimental Botany **59**, 4119-4131, doi:10.1093/jxb/ern251 (2008).
86 Yu, X. *et al.* A brassinosteroid transcriptional network revealed by genome-wide
identification of BESI target genes in Arabidopsis thaliana. *Plant J* **65**, 634-646,
doi:10.1111/j.1365-313X.2010.04449.x (2011).
87 Sun, Y. *et al.* Integration of brassinosteroid signal transduction with the
transcription network for plant growth regulation in Arabidopsis. *Dev Cell* **19**,
765-777, doi:10.1016/j.devcel.2010.10.010 (2010).
795

A

WT   bak1   brl1brl3   bak1 brl1brl3   bri1   bri1 brl1brl3   quad   BRL3ox

B

WT   bak1   brl1brl3   bak1 brl1brl3   bri1   bri1 brl1brl3   quad   BRL3ox

C

control

sorbitol

100μm

WT   bak1   brl1brl3   bak1 brl1brl3   bri1   bri1 brl1brl3   quad   BRL3ox

D

WT   bak1   brl1brl3   bak1 brl1brl3   bri1   bri1 brl1brl3   quad   BRL3ox

Figure 1. BR perception loss-of-function mutant roots are less sensitive to osmotic stress

Figure 1. BR perception loss-of-function mutant roots are less sensitive to osmotic stress.

(**A**) Seven-day-old roots of WT, BR mutant *bak1, brl1brl3, brl1brl3bak1, bri1, bri1brl1brl3* and *bri1brl1brl3bak1* (*quad*) and BR overexpressor line *35S:BRL3-GFP (BRL3ox)* grown in control (-) or sorbitol (+) conditions (270mM). (**B**) Seven-day-old root length in control (grey) or sorbitol (black) conditions. Green line depicts relative root growth inhibition upon stress (ratio sorbitol/control root length). Averages of 5 independent biological replicates +/- S.E. (n>150). Different letters mean significant differences in an ANOVA plus Tukey's HSD test. (**C**) Four-day-old roots stained with propidium iodide (PI, red) treated 24h in control (top) or sorbitol (bottom) media. Green channel (GFP) depicts the localization of BRL3 membrane protein receptor under the 35SCaMV constitutive promoter localized in the vascular tissues in primary roots. (**D**) Quantification of cell death in sorbitol-treated root tips. Red dots depict the mean relative PI staining (sorbitol/control) for each genotype. Averages of 5 independent biological replicates +/- S.E. (n>31). Different letters mean significant differences in an ANOVA plus Tukey's HSD test.

Figure 2. BRL3 receptor signalosome promotes root hydrotropism responses

Figure 2. BRL3 receptor signalosome promotes root hydrotropism responses.

(**A**) Root curvature hydrotropic response in seven-day-old roots after 24h of sorbitol stress induction (270mM). (**B**) Hydrotropic curvature angles distribution. (**C**) Mean values of root curvature angles in WT (24º), *brl1brl3* (23º), *bri1* (20º), *bri1brl1brl3* (18º), *quad* (16º), *bak1* (15º), *br1brl3bak1* (10º) and *BRL3ox* (30º). Averages of 4 independent biological replicates +/- S.E. (n>50). Different letters indicate a significant difference (p-value<0.05) in one-way ANOVA test plus Tukey's HSD test. (**D**) Stress traits matrix for all physiological assays performed on WT, BR loss-of-function mutant and *BRL3ox* roots and shoots. Root growth in control conditions is highlighted in green. Color bar depicts values for scaled data.

Figure 3. BR perception loss-of-function and overexpression mutants show higher survival rates upon drought stress

Figure 3. BR perception loss-of-function and overexpression mutants show higher survival rates upon drought stress.

(**A**) From top to bottom, three-week-old plant rosette phenotypes of WT, *brl1brl3, bak1*, *brl1brl3bak1*, *bri1*, *bri1brl1brl3, quad* and *BRL3ox* grown in well-watered conditions (left column), after 12 days of drought stress (middle column) and plant survival phenotypes after 7 days of re-watering (right column). (**B**) Plant survival rates after 7 days of re-watering. Averages of 5 independent biological replicates +/- S.E. (n>140). Asterisk indicates a significant difference (p-value<0.05) in chi-squared test of mean survival ratios. (**C**) Bar plot shows the days needed to reach different percentages of the soil field capacity for each genotype used in the study. (**D**) Photosynthesis efficiency (μmol/m2*s) at different percentages of field capacity. Bars represent means +/- S.E (n=6). Different letters depict significant differences within each genotype in a one-way ANOVA plus a Tukey's HSD test. (**E**) Transpiration rates (mmol/m2*s) at different percentages of field capacity. Bars represent means +/- S.E (n=6). Different letters depict significant differences within each genotype in a one-way ANOVA plus a Tukey's HSD test. (**F**) Schematic representation on BR signaling level and adult plant size and drought resistance. Loss-of-function mutants avoid the stress passively; whereas plants with increased BRL3 receptors act actively deploying drought stress.

Figure 4. BRL3 overexpression plants show a primed metabolic signature

Figure 4. BRL3 overexpression plants show a primed metabolic signature.

(**A**) Metabolite profiling of *BRL3ox* shoots in control conditions unveil a significant accumulation of 18 metabolites compared to WT. Metabolite profiling of *quad* shoots in control conditions show significant decreased levels of 7 metabolites compared to WT roots. (**B**) Metabolites following differential dynamics between *BRL3ox* and WT shoots along drought time course. Heatmap represent the log2 ratio of *BRL3ox*/WT. (**C**) Metabolite profiling of *BRL3ox* roots in control conditions unveil a significant accumulation of 20 metabolites compared to WT. Metabolite profiling of *quad* shoots in control conditions show significant decreased levels of 11 metabolites compared to WT roots. (**A-C**) Dotplots represent the means of each metabolite level (in *BRL3ox* and quad) relativized with the median of WT levels +/- S.D. (n=5). Asterisks denote statistical differences in a two-tailed t-test (p-value < 0.05) of the raw data (upper part comparison *BRL3ox* vs. WT, lower part comparison *quad* vs. WT). (**D**) Metabolites following differential dynamics between *BRL3ox* and WT roots along drought time course. Heatmap represent the log2 ratio of *BRL3ox*/WT. (**E-J**) Clustering of the dynamics of metabolite relative levels along drought time course in shoots and roots. Solid lines represent the median profile in the cluster while dashed lines represent the best polynomial curve that fits the data points in the cluster. Statistical significance was evaluated with maSigPro package[92]. (**E**) Metabolites following linear increase in drought in shoots include Glucose, Glucose-1P, myo-inositol and Sinapate. (**F**) Proline follows a stepper exponential increase in *BRL3ox* shoots. (**G**) Metabolites following exponential increase in BRL3ox shoots but linear in WT includes Galactose, GABA, Phenylalanine, Tyrosine, 2-methylmalate, Lysine, Isoleucine, Leucine, Nicotinate, Uracil and Tryptophan. (**H**) Metabolites following steeper exponential increase in *BRL3ox* roots include Trehalose, Sucrose, Proline and Raffinose. (**I**) Metabolites following a reduced linear increase until a maximum in *BRL3ox* roots include Glycerate and Malate. (**J**) Metabolites following exponential increase in *BRL3ox* roots but linear in WT include Glucose, Fructose, *myo*-inositol, Galactose and Asparagine.

Figure 5. Transcriptomics analysis of BRL3ox roots reveals activated drought stress responses already at basal conditions

Figure 5. Transcriptomics analysis of *BRL3ox* roots reveals activated drought stress responses already at basal conditions

(**A**) Number of genes upregulated or downregulated in control and after five days of water holding. Comparison of *BRL3ox* vs. WT roots. (**B**) Deployment of genes within "Response to stress" (GO:0006950) term that are also annotated as responsive to water, salt, heat, cold and light stress. Colors in the heatmap represent the log2 fold change of BRL3ox roots vs. WT in control (C) or drought (D) conditions. Red color in the squared right-sided heatmaps means that the gene has been previously identified as direct target of BES1 [82],BZR1 [83] transcription factors. (**C**) Overlap between deregulated genes in the control dataset and the drought data set. In both set only overlap a 13% of the total number of deregulated genes. (**D**) Most representative GO categories enriched in *BRL3ox* roots up-regulated genes at time 0 and after 5 days of drought. (**E**) Most representative GO categories enriched in *BRL3ox* roots down-regulated genes at time 0 and after 5 days of drought. Color bars: –log of adjusted p-value (Benjamini-Hochberg).

Figure 6. Enrichment of deregulated genes in BRL3ox in the root vasculature

Figure 6. Enrichment of deregulated genes in *BRL3ox* in the root vasculature.

(**A**) Tissue enrichment for upregulated (red) or downregulated (blue) genes in control conditions. Bars trespassing the p-value tresholed (0.05) were considered enriched in the data set. (**B**) Tissue enrichment for upregulated (red) or downregulated (blue) genes after five days of drought. Bars trespassing the p-value tresholed (0.05) were considered enriched in the data set. (**A-B**) Deregulated genes tissue enrichment [59]. AGL42: Quiescent center, APL: Phloem + Companion cells, COBL9: Root hair cells, CORTEX: Cortex, GL2: Non-hair cells, J2661: Pericycle, JO121: Xylem pole pericycle, LRC: Lateral root cap, PET111: Columella, RM1000: Lateral root, S17: Phloem pole pericycle, S18: Maturing Xylem, S32: Protophloem, S4: Developing xylem, SCR5: Endodermis, SUC2: Ploem companion cells. y-axis represent the negative logarithm of one-tailed Fisher's test. (**C**) Genes deregulated enriched in Pericycle (J2261 marker). Blue boxes highlight enzymes involved in the metabolism of deregulated metabolites. (**D**) Genes deregulated enriched in the Phloem Pole Pericycle (S17 marker). Blue boxes highlight enzymes involved in the metabolism of deregulated metabolites. (**E**) Genes deregulated enriched in the Phloem Companion Cells (SUC2 marker). (**F**) Genes deregulated enriched in all Phloem (APL marker). (**C-F**) Bars represent the log2 fold-change of *BRL3ox* vs. WT roots in control (black) or drought conditions (gray).

Supplementary Figure 1. BR loss-of-function quad mutant roots are insensitive to BRs

Supplementary Figure 1. BR loss-of-function *quad* mutant roots are insensitive to BRs.

(**A**) Five-week-old *quad* mutants mature plants phenotype. (**B**) Six-day-old roots of WT, *bri1*, *bri1brl1brl3* and *quad* seedlings in a dose-response curve assay at 0.0004, 0.004, 0.04, 0.4, 4, 40 and 400nM of brassinolide (BL) treatment. (**C**) Relative root length of six-day-old WT, *bri1*, *bri1brl1brl3* and *quad* seedlings within BL dose-response curve. Root length in BL was relativized to control root length for each concentration and genotype. BL resistance is observed in *bri1*, *bri1brl1brl3* and *quad* roots. Results are average of three independent biological replicates +/- S.E. (n=100).

Supplementary Figure 2. BRZ₂₂₀ chemical inhibitor reverts WT root hydrotropic response in sorbitol

Supplementary Figure 2. BRZ$_{220}$ chemical inhibitor reverts WT root hydrotropic response in sorbitol.

(**A**) Curvature angles of roots when the bottom part was replaced with the same media (mock). A small curvature in all genotypes was found (around 7º) but not different from WT. (**B**) Curvature angles of roots when the bottom part was replaced with sorbitol containing media. Note the difference in the scale units with respect A. (**A-B**) Bars represent angle means +/- S.E. (n>50). Different letters mean significant differences in one-way ANOVA plus Tukey's HSD test. (**C**) Distribution graphs of root curvature angles of WT plants in control conditions show 90% of roots curved less than 10º while WT plants treated with 270mM sorbitol show only 8% of roots less than 10º and 68% of roots in between 0 and 30º. Exogenous application of BRZ$_{220}$ reverted the sorbitol-induced response showing 33% of roots curved below 10º and 90% of the roots below 30º, similar to the *quad* mutants insensitive root phenotypes.

Supplementary Figure 3. Physiological parameters of mature plants under drought stress

Supplementary Figure 3. Physiological parameters of mature plants under drought stress

(**A**) Phenotypes of mature plants after drought. Rosettes were photographed when a 70% of soil water loss was reached (P70%). (**B**) Relative water content (RWC) of mature rosettes at 0% (field capacity), 50% and 70% soil water-loss. (**C**) Photosystem II (PSII) efficiency quantification of mature rosettes at 0% (field capacity), 50% and 70% soil water-loss. (**B-C**) Bars represent means +/- S.E. (n=6). Different letters depict significative differences within each genotype in a one-way ANOVA plus a Tukey's HSD test. (**D**) Plot of photosynthesis efficiency against soil water loss. Data was fitted to lineal model and the slops taken as indicators of drought-induced photosynthesis inhibition. Dots are the means for each genotype and dashed lines the fitted model.

A

Arabidopsis shoots primary metabolite sample distribution

B

	WT		BRL3ox		quad	
	watered	drought	watered	drought	watered	drought
days	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6

Raffinose
Proline
Tyrosine
Galactose
Glucose
Fructose
Ascorbate
Guanidine
Citrate
Phosphorate
Glycine
Succinate
4_hydroxyproline
Putrescine
Glutamine
Asparagine
Alanine
Citramalate
Threonine
Glycerate
Salicylate
Isocitrate
myo-inositol
Fumarate
Tryptophan
Octadecanoate
Palmitate
Arginine
Ornithine
Urea
Valine
Ribulose-5-phosphate
Aspartate
Benzylalcohol
Glycolate
Benzoate
1_6_hydroglucose
Pyroglutamate
Erythritol
Nicotinate
Methionine
Glycerol
Pyruvate
Glutamate
Ethanolamine
Serine
2_oxoglutarate
Erythronate
Malate
Histidine
Sucrose
Sinapate
Phenylalanine
Threonate
Galactinol
Maltose
Dehydroascorbate
Spermidine
2_methylmalate
Glucose-1-phosphate
GABA
beta-Alanine
Uracil
Lysine
Isoleucine
Leucine

-2 -1 0 1 2
log2 (relative abundance)

Supplementary Figure 4. WT, BRL3ox and quad mutant metabolite profiling under drought stress in the shoot

Fàbregas et al., MS.

Supplementary Figure 4. WT, *BRL3ox* and *quad* mutant metabolite profiling under drought stress in the shoot

(**A**) Distribution of complete sample set of WT, *BRL3ox* and *quad* shoots within a drought time course. Data show a uniform distribution without normalization. Samples which median was outside the average interquartile range were not considered for further analysis. (**B**) Heat map depicts the metabolic fingerprint of WT, *BRL3ox* and *quad* shoots within time course at control conditions and under drought stress. Homogeneous metabolite relative levels are shown in well-watered conditions, whereas deregulated metabolite patterns appeared upon drought exposure. Color bar depicts row-scaled values for the relative metabolite abundance.

Arabidopsis roots primary metabolite sample distribution

A

B

Supplementary Figure 5. WT, BRL3ox and quad mutant metabolite profiling under drought stress in the root

Supplementary Figure 5. WT, *BRL3ox* and *quad* mutant metabolite profiling under drought stress in the root

(**A**) Distribution of complete sample set of WT, *BRL3ox* and *quad* roots within a drought time course. Data show a uniform distribution without normalization. Samples which median was outside the average interquartile range were not considered for further analysis. (**B**) Heat map depicts the metabolic fingerprint of WT, *BRL3ox* and *quad* roots within time course at control conditions and under drought stress. Homogeneous metabolite relative levels are shown in well-watered conditions, whereas deregulated metabolite patterns appeared upon drought exposure. Color bar depicts row-scaled values for the relative metabolite abundance.

A *quad*/WT shoots

B *quad*/WT roots

D Upregulated in *quad vs.* WT

days of drought

cellular response to reactive oxygen species
response to iron ion
iron ion transport
brassinosteroid homeostasis
extracellular matrix organization
response to biotic stimulus
response to stress
cellular response to oxygen levels
programmed cell death
response to oxidative stress

-log(pvalue)

E Downregulated in *quad vs.* WT

days of drought

response to stress
response to jasmonic acid
response to water
response to abiotic stimulus
response to osmotic stress
growth
response to oxidative stress
cell wall organization
root development
response to iron ion
response to oxygen-containing compound
cell growth
carbohydrate metabolic process

-log(pvalue)

Supplementary Figure 6. Metabolomic and transcriptomics analysis of quad mutants under drought stress

Supplementary Figure 6. Metabolomic and transcriptomics analysis of *quad* mutants under drought stress

(**A**) Metabolites following differential dynamics between *quad* and WT shoots along drought time course. (**B**) Metabolites following differential dynamics between *quad* and WT roots along drought time course. (**A-B**) Clusters summarizing the drought dynamics of a group of metabolites are represented to the right. Heatmap represents the log2 ratio of *quad*/WT. (**C**) Microarray analysis of quad mature rosettes along drought time course. Heatmap represents row-normalized expression values. Black bars columns at the right depict if the gene has been identified as BES1 or BZR1 direct target. (**D**) Representative GO categories enriched among *quad* upregulated genes at 0,2,4 or 6 days of drought. (**E**) Representative GO categories enriched among *quad* upregulated genes at 0,2,4 or 6 days of drought. (**D-E**) Heatmaps represent the –log of corrected p-value for the enrichment (Benjamini-Hochberg).

Supplementary Figure 7. Deregulated genes responding to hormones in BRL3ox roots and hormone profiling

Supplementary Figure 7. Deregulated genes responding to hormones in *BRL3ox* roots and hormone profiling.

(**A**). Deployment of genes within "Response to hormone" (GO:0009725) term that are also annotated as responsive to brassinosteroids (BR), jasmonic acid (JA), gibberelines (GB), ethylene (ET), abscisic acid (ABA), auxin (AUX), citokinin (CK) or salicylic acid (SA). Colors in the heatmap represent the log2 fold change of *BRL3ox* roots vs. WT in control (C) or drought (D) conditions. Red color in the squared right-sided heatmaps means that the gene has been previously identified as direct target of BES1 [82],BZR1 [83] transcription factors. (**B**) Jasmonic acid (JA) levels. (**C**) *trans*-Zeatin levels, a type of cytokinin. (**D**) Indole-3-acetic acid (IAA) levels, the most common auxin. (**E**) Salicylic acid (SA) levels. (**F**) Total gibberellic acid (GAs) levels. (**G**) Abscisic acid (ABA) levels. (**H**) 1-aminocyclopropane-1-carboxylic acid (ACC) levels, a precursor of ethylene biosynthesis. (**B-H**) Data from 5 biological replicates. Error bars denote the average +/- S.E. Different letters indicate a significant difference (p-value<0.05), either in control (capital letters) or drought (lower-case letters) conditions. One-way ANOVA plus Tukey's HSD test.

A

Cell wall organization genes

Supplementary Figure 8. Genes of cell wall organization (GO category) deregulated upon drought in BRL3ox roots

Supplementary Figure 8. Genes of cell wall organization (GO category) deregulated upon drought in *BRL3ox* roots.

(**A**) Deployment of genes within "Cell wall organization" (GO:0071555) term that are deregulated in *BRL3ox* roots in control and drought conditions. Colors in the heatmap represent the log2 fold change of *BRL3ox* roots vs. WT. Red color in the squared right-sided heatmaps means that the gene has been previously identified as direct target of BES1 [82],BZR1 [83] transcription factors

A

Supplementary Figure 9. Transcription factor analysis of deregulated genes in BRL3ox roots upon drought

Supplementary Figure 9. Transcription factor analysis of deregulated genes in *BRL3ox* roots upon drought.

(**A**) Families of deregulated transcription factors in control (left) and drought (right) conditions. Color key represents the proportion of the family in the total number of deregulated transcription factors. (**B**) Transcription factors with a differential response between *BRL3ox* and WT roots in front of drought (lineal model accounting for interaction). Root tissue-specific [59] transcription factors are denoted in colors; APL: Phloem and phloem companion cells, COBL9: Root hair cells, J2261: Pericycle, S18: Maturing xylem, SUC2: Phloem companions cells.

A

STARCH AND SUCROSE METABOLISM

B

GALACTOSE METABOLISM

Supplementary Figure 10. Upon drought Sucrose and Galactose metabolic pathways are transcriptionally controlled by BRL3

Fàbregas et al., MS.

Supplementary Figure 10. Upon drought Sucrose and Galactose metabolic pathways are transcriptionally controlled by BRL3.

(**A**) Starch and Sucrose metabolism (KEGG ath00500). Deregulated genes in *BRL3ox* roots are depicted in colored boxes between metabolites. (**B**) Galactose metabolism (KEGG ath00052). Deregulated genes in *BRL3ox* roots are depicted in colored boxes between metabolites. Metabolic pathways with the deregulated enzymes were generated with PaintOmics.

Supplementary Figure 11. Role of vascular BRL3 receptors in drought stress

Supplementary Figure 11. Role of vascular BRL3 receptors in drought stress.

Drought resistance is generally associated to arrested plant growth. Here, we uncover a role for vascular BR receptors in conferring drought tolerance without penalizing overall plant growth. BRL3 receptors are enriched in the vascular (phloem) tissues where they control the division of stem cells. During drought, BRL3 signaling triggers the production of an osmoprotectant metabolic signature that are mobilized to the roots preventing growth arrest during severe drought periods.

A

Supplementary Figure 12. Comparison of root length measurements using ImageJ and MyROOT software

Supplementary Figure 12. Comparison of root length measurements using ImageJ and *MyROOT* software.

(A) Seven-day-old root length of WT, BR mutants and *BRL3ox* in control and sorbitol (sorb) conditions. Measurements were done manually using ImageJ and compared to data obtained by the semi-automatic software *MyROOT*. No significant differences (t-test, p-value > 0.05) in the mean root lengths were observed between both methods. Bars represent means +/- S.E.