Contrasting effects of acute and chronic stress on the transcriptome, epigenome, and immune response of Atlantic salmon

Tamsyn M. Uren Webster[1], Deiene Rodriguez-Barreto[1], Samuel A.M. Martin[2], Cock van Oosterhout[3], Pablo Orozco-terWengel[4], Joanne Cable[4], Alastair Hamilton[5], Carlos Garcia de Leaniz[1], Sofia Consuegra[1]

1. College of Science, Swansea University, Swansea, SA2 8PP, UK

2. College of Life Sciences & Medicine, University of Aberdeen, Aberdeen, AB24 2TZ, UK

3. School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

4. School of Biosciences, Cardiff University, Cardiff, CF10 3AX, UK

5. Landcatch Natural Selection Ltd, Stirling, FK9 4NF, UK

Abstract

Early-life stress can have long-lasting effects on immunity, but the underlying molecular

mechanisms are unclear. We examined the effects of acute stress (cold-shock during

embryogenesis) and chronic stress (absence of tank enrichment during larval-stage) on the

gill transcriptome and methylome of Atlantic salmon four months post-hatch. While only

chronic stress induced pronounced transcriptional effects, both acute and chronic stress

caused lasting, and contrasting, changes in the methylome. Crucially, we also found that

acute stress enhanced transcriptional response to an immunological challenge

(lipopolysaccharide), but chronic stress suppressed this response. Our results suggest that

epigenetic regulation may contribute to these effects on immuno-competence. Stress-

induced changes in promoter or gene-body methylation were associated with altered

expression of a small proportion of genes, and we also identified wider epigenetic regulation

of signalling pathways involved in immune response. These findings have important

implications for the management of stress and disease in wild and farmed fish.

Introduction

Stress response is a fundamental survival mechanism that provides a critical adaptive response to a range of environmental challenges (Chrousos 2009). Aspects of stress response may compromise the immune system, for example through immunosuppressive effects of glucocorticoid release, but in some circumstances mild stress may also have an enhancive effect on immunity (Calcagni & Elenkov 2006; Tort 2011). The precise impacts of environmental stress on the immune system are highly dependent on the timing, duration, magnitude and nature of the stressor and stress response (Tort 2011). Chronic stressors, lasting for days or weeks, dysregulate the immune response by inducing long-term changes in energetic metabolism, continuous low level inflammation, and by suppressing the release and function of immune cells and cytokines (Dhabhar 2014). In contrast, acute stressors, lasting for minutes to hours, are less likely to impair immune function and may even have enhancive effects through stimulating the maturation and redistribution of immune cells and cytokine secretion (Dhabhar 2009).

Developmental plasticity during critical periods of differentiation and maturation of the vertebrate nervous and immune systems increases sensitivity to stress during early life (Cao-Lei et al. 2017). In mammalian systems, it is well established that early-life stress can have long-lasting adverse effects on health and disease susceptibility. Maternal stress during pregnancy, for example, predisposes the offspring to developmental, immunological and behavioural abnormalities throughout their life, and post-natal trauma is also associated with an increased risk of depression, obesity, diabetes and cardiovascular disease (Silberman et al. 2016; Vaiserman 2015). However, exposure to mild stress during early life can have beneficial effects later in life. This effect includes enhanced immune function as part of a primed, more efficient stress response to future harsh environmental conditions (Monaghan & Haussmann 2015), and it has also been suggested that this effect could be harnessed in a clinical setting to boost protective immunity (Dhabhar 2014). However, the molecular mechanisms contributing to this effect have yet to be fully established.

To date, research into the long-lasting effects of early-life stress on latter-life health has

been conducted predominantly on the brains of humans and mice, due to the importance of

stress-related psychiatric disorders (Nasca et al. 2015a). These studies suggest that

epigenetic modifications mediate a neuronal memory of stress exposure, resulting in lasting

structural and transcriptional changes in the brain (Hunter et al. 2015; Nasca et al. 2015b).

Crucially, the epigenome is very sensitive to environmental factors, especially during early-

life, providing a way for stress to have long-term effects on transcriptional regulation and

phenotype throughout an individual's lifetime and, in some cases, on its progeny (Cao-Lei et

al. 2017; Silberman et al. 2016). In particular, promoter silencing activity, whereby DNA

methylation in promoter regions negatively regulates gene transcription, has been

convincingly demonstrated to mediate lasting stress-induced effects on physiology,

behaviour and disease in mammals (McGowan et al. 2009; Nasca et al. 2015b). However, at

a genome-wide level, the association between DNA methylation and gene expression is far

from straightforward, with complex interactions between different targets, cell types and

layers of epigenetic regulation (Hunter et al. 2015; Silberman et al. 2016).

Beyond these critical effects on the brain and behaviour, stress may also have far-

reaching effects on whole-organism physiology, including metabolism, nutrition, immunity

and reproduction (Papadopoulou et al. 2015). However, the potential for long-lasting effects

of stress on these wider processes is largely unknown, particularly for non-model species.

For fish, despite a wealth of research documenting the effects of both acute and chronic

stress, few studies have investigated how stress experienced during very early life can affect

later health outcomes beyond that of growth performance. Early developmental stages of

fish are clearly different from that of mammals, with embryos and alevins being readily

exposed to a range of environmental stressors, in both the natural environment and as part

of normal aquaculture practice. Most fish species display a type III survivorship curve

characterised by very high embryo mortality and a critical time for survival that coincides with

the transition from endogenous to exogenous feeding (Elliott 1989). It seems likely that these

early-life stages, coinciding with critical periods of immune development and epigenetic

plasticity, may be especially vulnerable to stress disruption, with the potential for long-lasting

effects on immune status and disease resistance (Castro et al. 2015). Although epigenetics

research in fish is in its infancy, Moghadam et al. (2017) recently reported that stress during

early life modulated the methylome and transcriptome of Atlantic salmon (*Salmo salar*) in a

developmental-stage specific manner, suggesting that epigenetic mechanisms may mediate

lasting effects of stress on fish health, as for mammals. Yet, whether the response of fish to

acute and chronic stress is similar to that shown by mammals, and how it influences the

immune system and disease resistance is unknown. Understanding the way in which fish

respond to environmental stress, and the mechanisms driving these effects, is essential in

order to improve the sustainability of both natural and farmed fish populations which are

facing increasing environmental pressures through overexploitation, climate change, habitat

loss and fragmentation, pollution and aquaculture intensification.

Here, we compared the effects of acute and chronic stress during early life on both the gill

transcriptome and methylome, and on transcriptional immune response to a model

pathogenic challenge (bacterial lipopolysaccharide, LPS) in Atlantic salmon. This species is

threatened by overexploitation and climate change, and is also one of the most widely

farmed fish worldwide. We identified contrasting effects on the baseline transcriptome and

on transcriptional response to LPS four months after the start of the stress experiment, with

acute and chronic stress enhancing and suppressing inflammatory immune response,

respectively. We also identified distinct DNA methylation patterns resulting from each type of

stress, which may contribute to the altered transcriptional response to pathogenic challenge

observed.

Materials and methods

Ethics statement

All experiments were performed with the approval of the Swansea Animal Welfare and

Ethical Review Body (AWERB; approval number IP-1415-6) and infection challenges were

approved by Cardiff University Animal Ethics Committee and conducted under a UK Home

Office License (PPL 302876).

Fish maintenance and stress experiments

Atlantic salmon eggs were obtained from Landcatch Natural Selection and assigned at

random to three experimental groups: control, acute environmental stress and chronic

environmental stress. To average out potential family effect, eggs were obtained from 10

different families (1:1 crosses) and equally assigned to each group. Each experimental

group consisted of two replicate vertical tray incubators (each containing 500 eggs), and

larvae from each replicate tray were transferred to a separate replicate fry trough (length 100

cm, width 40 cm, depth 8 cm) following hatching (at 460 degree days, DD).

We used an acute stressor which has previously been shown to induce repeatable effects

on the teleost transcriptome and methylome, and is relevant to temperature-shocking

procedures commonly used in aquaculture quality screen eggs (Moghadam et al. 2017).

During late embryogenesis (360 DD) embryos were immersed in iced water (0.2 °C) for five

minutes and then exposed to air (12 °C) for five minutes before being returned to the normal

water (9 °C). For the chronic stress group, hatched larvae were reared in fry troughs without

the artificial hatching substrate (Astroturf) used in the control and acute stress groups for the

duration of the experiment (four months). Artificial substrate is routinely used in aquaculture

to mimic natural substrate and provide support and shelter to fish larvae, and rearing in bare

tanks is known to increase stress (measured by cortisol levels), developmental abnormalities

and to impair growth (Bates et al. 2014; Hansen 1985).

Egg incubators and fry troughs were supplied with a constant flow (~ 3 L/min) of aerated

de-chlorinated tap water in a recirculation system. Upon commencement of feeding, fry were

fed Nutraplus salmonid feed (Skretting, UK). Water oxygen saturation was maintained above

90%, and ammonia (<0.02 mg/L), nitrate (<0.01 mg/L), nitrite (<15 mg/L) and pH (7.5 ± 0.2)

levels were maintained within the optimal range for the species. Water temperature was

increased from 9 °C to 11 °C and photoperiod was adjusted from 10:14h to 14:10h light:

dark over the 4 month experimental period, to reflect seasonal variation. Daily mortalities of

embryos, larvae and fry were recorded, and growth was monitored based on a subset of 20

euthanised individuals from each of the six replicate troughs at four time-points; 492, 748,

1019 and 1323 DD. Fish were euthanised via an overdose of anaesthetic (Phenoxyethanol;

0.5 mg/L), followed by destruction of the brain according to UK Home Office regulations. At

the final sampling point (1532 DD), mass and fork length of 20 fish from each tank were

determined and used to calculate Fulton's condition factor (Froese 2006). All gill arches

from both sides of each fish were dissected out and stored in RNAlater (Sigma Aldrich, UK)

at -20 °C for subsequent RNA/DNA extraction. Separately, at each of the five sampling

points, effects of stress treatment (control, acute, chronic) on fish mass, as well as condition

factor at the final sampling point, was assessed using linear mixed effect models (lme

function in nmle (Pinheiro et al. 2017)) in R version 3.3.3 (R_Core_Team 2014) using tank

identity as a random factor to account for variation between replicate tanks.

Immuno-stimulation experiment

To assess the effect of acute and chronic environmental stress on immune response, a

subset of salmon fry from each stress group and the control group were exposed to

lipopolysaccharide (LPS), a pathogen-associated molecular pattern, mimicking a bacterial

infection, at the end of the experimental period (1532 DD). Six fry from each replicate tank

(12 per group) were exposed to 20 µg/ml LPS obtained from *Pseudomonas aeruginosa*

(Sigma Aldrich, UK) for 24 h in 0.5 L tanks, each containing a static volume of aerated water

(Sundaram et al. 2012). Fish were visually monitored during the course of the experiment to

assess any effect on behaviour or physiology (for example, swimming ability, activity levels,

colouration, gill opercula movements and gasping). After exactly 24h exposure, fry were

euthanised as above, weighed and measured, and all gill arches were dissected out and

stored in RNAlater. Gills were selected as a target tissue for analysis because they have a

critical role in immune defence, and are a primary target of waterborne pathogen exposure

(Uribe et al. 2011).

Transcriptome and methylome sequencing

Matched transcriptome and methylome analysis of the gill was performed at the final sampling point (1532 DD) for a total of eight fish in each of the two stress groups and the control group (24 fish in total, including four from each replicate tank). Transcriptome analysis was also performed on the gills of eight LPS-exposed fish from each of the three experimental groups (24 fish; four per replicate tank). RNA and DNA were simultaneously extracted using the Qiagen AllPrep DNA/RNA Mini Kit, according to the manufacturer's instructions and then nucleic acid concentration was assessed using a Qubit 3.0 Fluorometer (Thermo Fisher Scientific, UK).

Transcriptomic analysis was conducted using RNA-seq. RNA purity and integrity was assessed using a NanoDrop NS-100 Spectrophotometer (NanoDrop Technologies, USA) and an Agilent 2100 Bioanalyzer (Agilent Technologies, USA). All RNA used for library construction was of high quality with 260/230 and 260/230 ratios > 1.8 and RIN scores >8. Library preparation was performed using the Illumina TruSeq RNA Library Prep Kit v2, with 1 µg total input RNA. Sequencing of the 48 libraries was performed on an Illumina NextSeq500 platform at University of Aberdeen, across a total of eight flow cells, using 76 bp paired-end sequencing.

Methylation analysis was performed using Reduced Representation Bisulfide Sequencing (RRBS). Bisulphite converted genomic DNA libraries were prepared using the Diagenode Premium RRBS Kit, including fully methylated and un-methylated spike-in controls to monitor bisulphite conversion efficiencies. Briefly, 100 ng of genomic DNA were digested with *MspI*, followed by end-repair, adaptor ligation and bead-based fragment size selection. 24 samples were multiplexed in pools of six following the manufacturers' recommendations. After multiplexing, the pooled samples were bisulphite-converted, amplified by enrichment PCR, and quality checked using the Agilent D1000 ScreenTape System. The libraries were then sequenced across 6 lanes on Illumina NextSeq 500 platform at Cardiff University using 76 bp single-end sequencing.

All raw sequence reads have been deposited to the European Nucleotide Archive under

the accession numbers PRJEB25636 and PRJEB25637. All analyses were conducted on a

local server running NEBC Bio-Linux 8.

Transcriptomic analysis

Quality screening of raw reads was performed using FastQC (Andrews 2010).

Trimmomatic (Bolger et al. 2014) was used to remove contaminating adaptor sequences and

the last three bp of all reads (containing nucleotide bias), and trim poor quality bases from

the 3' end using a sliding window of 4 bp, and specifying Q≥20. All reads < 30 bp in length

were discarded. High quality paired reads from each sample were then aligned to the

Atlantic salmon genome (v GCF_000233375.1_ICSAG_v2; (Davidson et al. 2010; Lien et al.

2016)) using HISAT2 (v 2.1.0; (Kim et al. 2015)), after first converting the GFF3 annotation

file to a GTF file using gffread (Pertea 2016), to enable splice-aware alignment and novel

transcript discovery. SAM alignment files were converted to sorted BAM files using samtools

v 1.5 (Li et al. 2009), and transcript reconstruction and assembly was then performed using

StringTie (v1.3.3) using recommended parameters (Pertea et al. 2015). Non-normalised

read counts for the set of merged assembled transcripts were then extracted using the

prepDE.py script supplied in the StringTie package.

Differential expression analysis was performed using DeSeq2 (Love et al. 2014) for

78,229 putative loci with at least 1 read count in more than one sample. Differentially

expressed genes in response to stress and LPS exposure were identified using a

multifactorial design, including the main effects of stress and LPS exposure, and their

interaction, and accounting for potential variation between replicate tanks. Within the model,

independent filtering of loci with low coverage depth was applied, optimising power for

identification of differentially expressed transcripts at a threshold of alpha=0.05. Default

settings were applied for outlier detection and moderation of gene-wise dispersion estimates.

Genes were considered significantly differentially expressed at FDR <0.05, and the contrast

function was used to extract differentially expressed gene identities.

For visualisation and cluster analysis, raw counts were normalised using variance

stabilising transformation within DeSeq2. Multidimensional scaling (MDS) analysis was

performed using pairwise distances between all samples, and visualised using ggplot2

(Wickham 2009). Heat maps were constructed to visualise transcriptional response to LPS,

and the effect of each type of stress on transcriptional response to LPS. Hierarchical

clustering of all genes significantly regulated by LPS, and all genes for which a significant

interaction between stress and LPS response was identified, was performed using an

Euclidean distance metric and visualised using the Pheatmap package in R (Kolde 2015).

Functional enrichment analysis of differentially regulated genes was performed using

DAVID (v 6.8; (Huang et al. 2008)), using all expressed genes in the gill transcriptome as

background. Due to the lack of gene ontology terms attached to Atlantic salmon gene ids,

the enrichment analysis was conducted on zebrafish orthologs which were identified using

Blastx against the Uniprot peptide database (specifying a minimum e value cutoff of $1e^{-15}$).

Methylation analysis

Following initial quality assessment of raw sequences using FastQC (Andrews 2010),

TrimGalore (Kreger 2016) was used to trim low-quality base calls and adapter contamination

using the '-- rrbs' option. High quality reads were aligned to the Atlantic salmon reference

genome and cytosine methylation calls extracted using Bismark v 0.17.0 (Krueger &

Andrews 2011) using the '--comprehensive' parameter. Mapped data were then processed

using SeqMonk (Andrews 2007) within R, considering only methylation within CpG context.

Only CpGs with a minimum coverage of 10 reads in each of the 24 samples were included in

the analysis, and extreme coverage outliers (>99.9th percentile) were removed. CpG island

locations were predicted using the newcpgreport tool within EMBOSS (Rice et al. 2000),

defining islands as regions of ≥200 bp with GC % ≥ 55% and an observed-to-expected CpG

ratio of ≥ 65%, island shores as regions located up to 2 kb of a CpG island, and shelves as 2

kb regions from a CpG shore.

Differentially methylated CpGs (DMCpGs) were identified using logistic regression tests

on the quantitated normalised data based on $p<0.01$ after multiple testing correction

(Benjamini-Hochberg) and >20% minimal CpG methylation difference ($|\Delta M|$). For each

DMCpG, we identified the genomic location (within gene body, promoter region (≤1500 bp

upstream of the transcription start site (TSS)), or intergenic region) and the context location

(CpG island, CpG shore, CpG shelf). For the DMCpGs that were within a gene, or within 2

kb (upstream or downstream) of the TSS or transcription termination site (TTS) respectively,

we also performed enrichment analysis based on gene function using DAVID v 6.8 (Huang

et al. 2008), after converting to zebrafish ids as described above. To generate a more

stringent list of DMCpGs for further cluster analysis between stress groups, we additionally

ran t-tests for each paired comparison using a threshold of $p<0.01$, to identify DMCpGs

shared by both statistical methods.

Transcriptome-methylome integration

Next we explored the relationship between the methylome and the transcriptome. We

performed targeted DNA methylation analysis for putative gene regulatory regions and for

gene bodies, and in each case investigated the relationship between total methylation level

and gene expression. As the regulatory element landscape for the Atlantic salmon reference

genome is unavailable (Macqueen et al. 2017), we defined the putative promoter regions

(p.promoters) as windows of 2500 bp surrounding the TSS (1500 bp upstream to 1000 bp

downstream), in accordance with previous studies (e.g. Rowley et al. 2017; Shahmuradov et

al. 2017). For the analysis of gene bodies, we only used gene bodies containing ≥ 5 CpGs,

each with ≥ 10 reads per CpG, in all 24 samples (10,017 genes covered out of 61,274

overall expressed genes; 16.3%). To increase the number of p.promoter regions included in

the analysis, we used a lower threshold (regions containing ≥ 3 CpGs, each with ≥ 5 reads

per CpG, in all 24 samples; 5,422 gene promoter regions covered out of 61,274 overall

expressed genes; 8.8%).

Firstly, we examined the transcriptome-wide relationship between gene expression and

DNA methylation within p.promoters and, separately, within gene bodies. We plotted mean

expression against mean DNA methylation for all covered genes in the control group, and

performed a Spearman correlation.

We then aimed to identify genes for which early life stress influenced both DNA

methylation and gene transcription. For all expressed genes at the baseline time-point (i.e.

fish not exposed to LPS), we plotted gene expression difference in each of the stress groups

relative to the control group (delta expression) against the respective difference in gene body

methylation and in p.promoter methylation (delta methylation). We identified genes with a

marked effect of stress on both expression and methylation (>2 fold delta expression and

>5% difference in methylation), based on previously described thresholds (e.g. Lussier et al.

2018; Muurinen et al. 2017) and performed functional enrichment analysis as before.

We also aimed to investigate the possible role of stress-induced changes in DNA

methylation in influencing transcriptional response to the immune challenge (LPS). We did

not perform methylation analysis for the fish exposed to LPS, but we hypothesised that

baseline promoter and/or gene body methylation status might influence the rapid

transcriptional response induced by LPS exposure. Therefore, for all genes for which a

significant interaction between stress and transcriptional response to LPS exposure was

identified, we performed a similar targeted methylation analysis. For each stress group, we

plotted delta expression following exposure to LPS relative to that in the control group

against delta baseline methylation relative to the control group.

<u>Results</u>

Survival and growth

Overall Atlantic salmon hatching success was 95.3 ± 1.1% and post-hatch larval survival

rate was 89.1 ± 1.1%, with no significant difference between the control, acute stress or

chronic stress groups (hatching; $F_{3,2}=1.38$, $P=0.377$, survival; $F_{3,2}=0.19$, $P=0.836$). There

was a significant effect of treatment on growth rate during pre- and early-feeding (748 and

1019 DD), whereby fish exposed to chronic stress had significantly reduced mass compared

to those in the control and acute stress groups, but this effect was no longer apparent at

later sampling points (492 DD: $F_{3,114}=0.37$, $P=0.718$; 748 DD: $F_{3,114}=15.82$, $P=0.025$; 1019

DD: $F_{3,114}=15.42$, $P=0.026$; 1323 DD: $F_{3,114}=1.63$, $P=0.330$, 1532 DD: $F_{3,114}=4.78$, $P=0.117$;

Table S1), and there was no significant difference in condition factor between groups at the

final sampling point $(F_{3,114}=2.183, P=0.260)$. Exposure to LPS for 24h caused no mortalities,

or any apparent physiological or behavioural changes.

Transcriptomic analysis

After quality filtering, an average of 27.8 million paired end reads (91.8%) were retained

per sample. Of these, a total of 94.5% were mapped to the Atlantic salmon genome,

including 84.1% unique alignments (Table S2). Following transcript reconstruction and novel

assembly, we obtained a total of 201,433 transcripts in 104,528 putative loci, and statistical

expression analysis was performed for 78,229 genes with nonzero read counts.

Expression analysis with DeSeq2, accounting for between-tank variation, identified a total

of 19 genes significantly differentially expressed (FDR <0.05) between the acute stress

group and the control fish. In the chronically-stressed group, there were 206 differentially

expressed genes compared to the control group, the vast majority (190, 92.2%) of which

were up-regulated, and a functional analysis of these genes revealed strong enrichment of

ribosome structural constituents and translation, as well as muscle development, energy

metabolism and bacterial defence response (Figure S1).

LPS exposure had a very marked effect on gene transcription in the gills. MDS analysis

based on the whole transcriptome clearly separated all LPS-exposed individuals from non-

exposed fish (Figure 1a). The main effect of LPS exposure in the (non-stressed) control

group consisted of 14,833 up-regulated and 10,636 down-regulated genes (FDR <0.05) in

fish exposed to 20 µg/ml LPS compared to non-exposed fish (Figure S2). These included a

large number of genes encoding proteins typically associated with inflammatory immune

response including a large number of mucins; interleukins, interferons, TNF, chemokines

and their regulatory factors and receptors; complement factors, immunoglobins and damage-

inducible molecular chaperones. Fold changes for a selected list of these significantly

differentially expressed genes with direct immune function is provided in the supporting

information (Table S3). Overall functional analysis for the up-regulated genes revealed

strong enrichment of GO terms related to cellular stress response (Figure S3). Enriched

terms included those related to cell adhesion, cellular signal transduction, regulation of

transcription, protein modification, response to LPS and response to cytokines (Biological

Process), extracellular matrix (Cellular Component), and transcription factor activity, protein

kinase/phosphatase activity, and binding of signalling molecules (Molecular Function).

Enriched KEGG pathways included extracellular matrix-receptor interaction, as well as

specific pathogen recognition pathways (NOD-like, RIG-like, TOLL-like receptor signalling).

For down-regulated genes enriched terms were related to general cell maintenance

processes. These included terms related to DNA-replication and repair, redox reactions, cell

cycle   and translation (Biological Process), ribosome and mitochondria (Cellular

Component), ribosome constituent, redox activity and metabolic enzyme activity (Molecular

Function), and DNA replication, ribosome and metabolic pathways (KEGG pathway).

We also identified a significant interaction between stress and transcriptional response to

pathogen challenge. Compared to the main effect of LPS exposure identified in control fish

(described above), we found that acute and chronic stress during early life altered

transcriptional response to LPS in contrasting ways (Figure 1b). In the acutely stressed fish,

a total of 194 genes were regulated by LPS in a significantly different way (FDR <0.05)

compared to in control fish. Of these genes, the vast majority (140, 72.2%) were up-

regulated relative to the control group in response to LPS and 41 (21.1%) were down-

regulated relative to the control group in response to LPS. Only 9 (4.6%) genes that were

regulated by LPS in the control group were not significantly responsive in the acutely-

stressed group, and 4 (2.1%) genes were regulated in the opposite direction. Functional

analysis of the genes with enhanced responsiveness to LPS in acutely-stressed fish

revealed further enrichment of processes identified for the main LPS response, together with

pathways related to lipid metabolism and mucin production (Figure S4a).

For chronically-stressed fish, a total of 347 genes were expressed in a significantly

different way following LPS exposure compared to the main effect of LPS exposure identified

in non-stressed control fish. The majority of these genes (218, 62.8%) were not significantly

regulated by LPS, or were significantly less responsive to LPS exposure relative to the

control group (Figure 1b). Functional analysis of these less responsive genes revealed

enrichment of a number of processes identified as part of the main LPS response (including

cell adhesion, signal transduction, response to bacterium/lipopolysaccharide and p53

signalling), and amongst the more down-regulated genes there was a strong enrichment of

ribosome and translation (Figure S4b). Only 28 (8.1%) and 57 (16.4%) genes, respectively,

were up-regulated and down-regulated to a greater extent in chronically-stressed fish than in

the control group, which is in stark contrast to the enhancive effect of acute stress on

transcriptional response to LPS.

Methylation analysis

After quality filtering, a total of 1534 million high quality single end reads, averaging 64

million/sample, were retained. A total of 90.6% of these were mapped to the reference

genome, with a unique alignment rate of 43.5% (Table S4). Analysis of spike-in methylation

controls revealed an overall bisulphite conversion efficiency of 99.7%, with 2.0%

inappropriate conversion. In total, we identified 21.3 million CpG sites in our libraries, but

only 1.1 million were covered in all samples, representing 2.75% of total CpGs in the Atlantic

salmon genome, which is a comparable percentage to that previously reported for RRBS

experiments in zebrafish (5.3%; Chatterjee et al. 2013) and rainbow trout (< 1%; Baerwald et

al. 2016). Methylation analysis was conducted only using CpGs covered by at least 10 reads

in all libraries (335,996 CpG sites).

The majority of the CpGs surveyed mapped to gene bodies (53%) or intergenic regions

(45%), with 2% located in putative promoter regions. In terms of CpG context, 12% of CpGs

were located in CpG islands, 17% in CpGshores, and 11% in CpGshelves. CpG methylation

level dropped progressively in the region upstream of the TSS, then increased sharply within

the gene body (Figure 2a). Genome-wide CpG methylation displayed a bimodal distribution,

whereby the majority of CpGs within gene bodies (60.08%) were highly methylated (>80%

methylation), while a large proportion (24.6%) were un-methylated or hypo-methylated

(<20% methylation) (Figure 2b). Across the putative promoter regions, there was a higher

proportion of hypo-methylated CpGs (64.1%) (Figure 2c). Genome-wide, the average

methylation of all CpG sites was 74.87% ± 0.38 in the control group, 74.04% ± 0.71 in the

acute stressed fish and 74.94% ±0.82 in the chronically stressed group, with no significant

difference among groups ($F_{2,21}=0.55, P=0.59$).

Compared to the control fish, a total of 1895 and 1952 differentially methylated CpG sites

(DMCpGs) were identified using logistic regression (FDR< 0.05, |ΔM|≥20%) in the acute

stress and chronic stress groups, respectively. The genomic distribution and context of the

DMCpGs largely mirrored the wider methylome landscape, with half of the DMCpGs

overlapping intragenic regions (52%). DMCpGs overlapped or neighboured (up to 2 Kb from

TSS) a total of 907 genes for the acute stress group, and 925 genes for the chronic stress

group, including 242 common genes shared in both groups. For both stress groups, the most

strongly enriched functional processes amongst these genes were related to cellular

adhesion and cellular signalling pathways (Figure S5). These included terms related to cell

adhesion, intracellular signal transduction, Rho protein signalling, calcium ion transport and

signalling and ion transport (Biological Process), plasma membrane, cell junction, myosin

complex (Cellular Component) and ion channel and transported activity, GTPase and

guanyl-nucleotide exchange activity (Molecular Function). A more stringent list of 1004 total

DMCpGs were identified across both types of stress using both statistical methods (logistic

regression and t-tests). Unsupervised hierarchical clustering of these DMCpGs revealed a

distinctive methylation profile for both the acutely and chronically stressed groups with

respect to the controls and to each other, although there was greater resemblance between

the control and acute stress groups (Figure 2d).

Firstly, we examined the transcriptome wide association between gene expression and

DNA methylation within p.promoter regions and within gene bodies. There was a significant

negative correlation between p.promoter mean methylation and gene expression (Spearman

rho= -0.37; P< $2.2e^{-16}$, based on 5422 total genes), but no overall linear correlation between

gene body methylation and gene expression (Spearman rho= -0.03). We fitted a smoothed

spline curve to illustrate the heterogeneous relationship between gene body methylation and

gene expression (Figure 3a,b).

We then identified genes for which there was a notable effect of early life stress on both

DNA methylation (p.promoters, gene bodies) and on gene expression (>2 FC delta

expression and >5% methylation difference) (Figure 3c-f). For p.promoters, there was

evidence of unequal distribution of genes between hyper-methylated/up-regulated, hypo-

methylated/up-regulated, hyper-methylated/down-regulated and hypo-methylated/down-

regulated groups (Acute stress: χ^2= 3.64, P=0.056, Chronic stress: χ^2= 9.32, P=0.002) with

greater numbers of genes with an inverse relationship between delta methylation and delta

expression (Table S5). Therefore, and given the overall negative relationship between

methylation in p.promoter regions and gene expression we focused only on these genes with

an inverse relationship. However, there was no evidence of a similar effect between delta

gene body methylation and delta expression (Acute stress: χ^2= 0.67, P=0.407, Chronic

stress: χ^2= 0.07, P=0.791), therefore we included all genes above the threshold (Table S6).

Combined functional analysis of these genes revealed enrichment of processes related to

ion/calcium ion transport and signal transduction (Figure S6).

Finally, we examined the potential for stress-induced changes in baseline DNA

methylation to contribute to the observed altered transcriptional response to LPS. Of the

genes that showed a significant interaction between stress treatment and response to LPS,

28 (acute stress) and 57 (chronic stress) met the coverage criteria for targeted analysis of

baseline p.promoter methylation (Figure S7). For acutely-stressed fish we identified three

genes with hypo-methylation ($|\Delta M|>5\%$) and increased-expression in response to LPS treatment relative to the control group (*lrrn4cl, usp54a, st3gal1l3*), and for chronically-stressed fish we identified three genes with hyper-methylation and reduced expression (*yaf2, casp3a, ddx56*). For gene body methylation, 42 (acute stress) and 63 (chronic stress) genes met the criteria for targeted analysis. Of these in acutely stressed fish we identified three hypo-methylated genes with respect to the control group (*cer, chpf, ahnakl*), and in chronically stressed fish we identified two hypo-methylated genes with respect to the control group (*nocta* and E3 ubiquitin-protein ligase KEG-like).

Discussion

Our study indicates that while acute stress experienced during early life had limited long-term effects on the gill transcriptome, chronic stress was associated with distinct lasting transcriptional changes in Atlantic salmon fry. However, both acute and chronic stress caused lasting, and contrasting, changes in the gill methylome. Crucially, early-life stress also altered transcriptional response to a model pathogen challenge in a stress-specific way; acute and chronic stress respectively enhanced and suppressed inflammatory response to LPS. Our data suggest that epigenetic changes may contribute to the modulatory effects of early life stress on immuno-competence observed. We identified a small proportion of genes for which there was a notable association between stress-induced changes in promoter or gene body methylation and changes in expression, suggestive of a direct regulatory relationship. Furthermore, gene enrichment analysis revealed broader epigenetic modification of critical cellular signalling pathways involved in immune response.

Lasting effects of stress on both the transcriptome and epigenome

Acute stress during late embryogenesis caused a lasting significant difference in the expression of fewer than 20 genes in the gill of salmon fry four months later. Previous studies have reported pronounced changes in transcription occurring immediately after

similar acute temperature challenges in fish embryos and in hatched fry (Donaldson et al.

2008; Moghadam et al. 2017). However, given that the acute stress was applied four months

earlier and we observed no effects on survival and growth, this suggests direct

transcriptional response to this acute, sub-lethal stress is short-lived. This is consistent with

apparent physiological and transcriptional recovery over time from acute temperature stress

and other stressors (Ankley & Villeneuve 2015; Donaldson et al. 2008). In contrast, over 200

genes were differentially expressed in response to chronic stress and, functionally, these

changes suggested an apparent up-regulation of active protein synthesis. Although

immediate stress responses have often been associated with reduced protein synthesis in

fish and mammals (Moghadam et al. 2017; Patel et al. 2002; Uren Webster & Santos 2015),

cellular stress response is extremely complex and it is possible that these transcriptional

changes represent a compensatory response to chronic stress. There was also evidence of

up-regulation of energy metabolism, muscle differentiation and insulin-like growth factor

signalling, which may be associated with a compensatory increase in growth rate in this

group after the initial weight loss observed.

With regards to the epigenome, both acute and chronic stress during early life induced

significant changes in gill DNA methylation profiles compared to controls. It was clear that

the two different stressors induced distinct and specific alterations in the methylation profile

of individual CpGs. Chronic stress induced a greater epigenetic change relative to the

controls, but acute stress also caused distinct and lasting effects on the methylome, even in

the absence of lasting transcriptional effects in this group. Stress has been shown to cause

long-lasting alterations in methylation profiles throughout an individual's lifetime that appear

to depend on the intensity and timing of the stressor (Cao-Lei et al. 2017; Moghadam et al.

2017; Vaiserman 2015). However, to our knowledge no study has examined the contrasting

effects of acute vs chronic stress on the fish transcriptome and epigenome. Notably, while

the acute and chronic stress resulted in quite distinct methylation patterns of individual

CpGs, functional analysis of the genes which contained or neighboured DMCpGs revealed

enrichment of very similar cellular processes for both types of stress. A large number of

terms related to cellular signalling pathways and their regulation were enriched for both

stress groups, particularly glutamate, calcium and Rho-GTPase signalling. Interestingly,

epigenetic modification has been reported to explain the dysregulation of the

neurotransmitter glutamate commonly associated with stress-induced disorders in the

mammalian brain (Nasca et al. 2015b), but glutamate also has an important signalling role in

peripheral tissues, including fish gill (Sundin et al. 2003), suggesting that this might be an

important, wider target of epigenetic regulation. Cellular adhesion was also one of the most

enriched terms in both stress groups, reflecting differential methylation of CpGs in a large

number of genes encoding cadherins and protocadherins, as well as integrins, laminins and

fibronectin. Cellular adhesion is critical for signal transduction as well as maintaining

structure in multicellular tissues, and altered epigenetic regulation of these components has

been reported for several autoimmune disorders and cancers (Zhang & Zhang 2015). This

differential methylation of similar signalling pathways by both acute and chronic early life

stress suggests that intra- and inter-cellular signal transduction may be a common target of

stress-induced epigenetic regulation, with the potential to influence an extremely diverse

array of cellular processes. However, it seems likely that the precise location and nature of

CpG methylation change, which was distinct between acute and chronic stress, accounts for

the fine tuning of epigenetic regulation and the resultant specific effects on transcription.

Stress-induced changes in immune response to a model pathogen exposure

There was a very pronounced transcriptional effect of LPS in the gill in all exposed fish

across treatment groups, characterising an extensive inflammatory response. Pro-

inflammatory cytokines (TNF, IFNγ, TGFβ, IL-1b), which are typical markers of LPS

regulation (Frost et al. 2002), together with many other cytokines, their regulatory factors and

receptors were differentially expressed. Several pathogen-associated molecular pattern

(PAMP) recognition signalling pathways (NOD-like receptor, RIG-I-like receptor and Toll-like

receptor signalling) were also functionally enriched, although there is no known specific

recognition receptor for LPS in fish (Sepulcre et al. 2009). Mucus provides a vital first line of

pathogen defence (Linden et al. 2008), and a number of mucins (muc-2,5,7,12,17 and 19)

were amongst the most up-regulated genes, together with other mucus components

essential for immune response including antimicrobial factors (hepcidin, cathepsin,

cathelicidin), lectins and complement factors. Transcriptional regulation and cellular

signalling pathways were strongly enriched amongst genes up-regulated by LPS exposure,

reflecting the enormous diversity and complexity of the immune response. In particular,

protein tyrosine kinases and phosphatases that are key regulators of signal transduction

cascades (Lemmon & Schlessinger 2010) were extensively up-regulated. Furthermore,

processes associated with the extracellular matrix (ECM), which provides cellular support

and facilitates cellular signalling and adhesion (Bonnans et al. 2014; Naba et al. 2016;

Sorokin 2010), were strongly enriched. Remodelling of the ECM is stimulated by

inflammatory cytokines and has a crucial role in inflammatory response, through immune cell

recruitment, activation and proliferation (Sorokin 2010). A large number of genes encoding

structural components of the ECM, including many collagens, elastins, integrins, cadherins,

laminins, thromobospondins and fibronectins, were up-regulated by LPS. We also found

enrichment of protein transport and exocytosis, which are involved in secretion of these ECM

components, and seriene/threonine proteases and matrix metallopeptidases which are

important for their cleavage and activation (Naba et al. 2016). In contrast, processes related

to cell division (cell cycle, DNA replication and repair) as well as protein, lipid and energy

metabolism, were most strongly enriched amongst down-regulated genes. It seems likely

that compensatory suppression of these processes, which are critical for the normal

maintenance of tissue function and order (Ragkousi & Gibson 2014), facilitate the

pronounced, acute immune response observed (Kan & Hodgkin 2014).

Transcriptional response to LPS exposure in the control group characterised a typical

inflammatory immune response. Crucially, we identified a significant interaction between

stress and LPS response, for both the acute and chronic stressors. For fish exposed to

acute cold shock during late embryogenesis, transcriptional response to LPS was of greater

magnitude, but functionally similar to that of the control group. The vast majority of LPS-

responsive genes identified in control fish were also differentially expressed in acutely

stressed fish, and a considerable number of these were regulated to a significantly greater

extent. Additional responsive genes included a number of other cytokines and their

receptors, mucins and ECM components. Processes related to lipid biosynthesis were

particularly over-represented amongst additional LPS-responsive genes in acutely stressed

fish, including genes involved in *srebp1* signalling and sphingolipid metabolism, both of

which are critical in mediating the membrane-dependent receptor based regulation of the

innate immune response (Köberlin et al. 2015). This suggests that acute stress during late

embryogenesis enhanced subsequent immune response, while in contrast chronic stress

appeared to depress the transcriptional response to LPS, as more than 200 genes were

significantly less responsive to the pathogen challenge than in the control group. These

included a number of the typical pro-inflammatory response markers and many genes

involved in processes such as signal transduction and ECM reorganisation, which were

identified as central to the main LPS response in control fish. These results are consistent

with previous reports of enhancing and suppressive effects of acute and chronic stress

respectively on immuno-competence in mammals and fish (Dhabhar 2014; Glaser & Kiecolt-

Glaser 2005; Smith et al. 2016; Tort 2011). While chronic stress is widely known to impair

immune function, by altering the balance and activity of immune cells and cytokines, acute

physiological stress can have an adaptive role, preparing organisms to deal with subsequent

challenges (Monaghan & Haussmann 2015). It is thought that mild, acute stress can

enhance both innate and adaptive immunity, by increasing the production and maturation of

immune cells and cytokines, especially when applied during key periods of immune

activation (Dhabhar 2014; Glaser & Kiecolt-Glaser 2005; Tort 2011).

An epigenetic basis for the lasting effect of stress?

On a gene specific basis, epigenetic mechanisms are known to mediate the lasting

effects of early life stress on physiology, behaviour and disease outcomes in mammalian

models (Cao-Lei et al. 2017; Vaiserman 2015). For example, reduced methylation in the

promoter of the glucocorticoid receptor, *Nr3c1,* due to early life stress is known to cause an

increase in its expression in the brain, with lasting physiological and behavioural effects

(Turecki & Meaney 2016). However, interpreting genome-wide associative patterns between

DNA methylation and gene-expression is exceptionally challenging due to the complexity of

the different layers of epigenetic regulation (Hunter et al. 2015). Existing evidence suggests

that the relationship between DNA methylation and gene expression varies widely across the

genome, and occurs on a gene-specific basis (Hu et al. 2016; Jones 2012). Here we found

evidence of a significant, transcriptome-wide negative correlation between DNA methylation

level in putative promoters and gene expression, which is consistent with previous reports in

mammals and fish (Huang et al. 2014; Wang et al. 2017; Zhong et al. 2016). In contrast,

there was no linear relationship between gene-body methylation and gene expression, which

again is consistent with previous reports in the literature for mammals and plants, where

non-monotonic relationships between gene methylation and expression have been reported

in some cases (Jjingo et al. 2012; Lim et al. 2017; Wang et al. 2015; Zemach et al. 2010).

However, transcriptome-wide, the relationship between gene expression and DNA

methylation was very variable among individual genes.

Given the marked effects of both acute and chronic early life stress on the gill methylome

we hypothesised that stress-induced changes in DNA methylation of putative promoter

regions and/or gene bodies could influence baseline transcription and also the rapid

transcriptional response to a pathogenic challenge. We identified a small proportion of genes

for which there was an association between stress-induced changes in baseline DNA

methylation and transcription. These included 20 different lncRNAs, perhaps reflecting the

complex and interactive nature of epigenetic modifications, since lncRNAs constitute another

layer of epigenetic regulation at the transcriptional and post-transcriptional level (Zhao et al.

2016). There were also 12 genes which were similarly influenced by both types of stress,

and overall functional analysis again revealed enrichment of ion transport and cellular

signalling pathways. Similarly, we found evidence of stress-induced methylation differences

(promoters or gene bodies) for a small proportion of the genes for which a significant

interaction between stress and LPS response was identified. Notably, these included a

number of genes involved in ubiquitination, which regulates a wide range of biological

processes including the immune system, and transcriptional regulation.

Our results suggest that direct associations between promoter or gene body methylation

and expression are likely to occur on a gene-specific basis. However, it is interesting that

this small proportion of genes consistently appear to include components of key signalling

and regulatory pathways, with the potential therefore to influence a diverse array of cellular

processes. We found limited direct evidence for stress-induced alterations in the methylome

corresponding to observed alterations in immune response to LPS. However, there was a

marked functional overlap between gene pathways with stress-induced changes in

methylation and those central to the inflammatory immune response to LPS, namely terms

related to cellular adhesion/the ECM and signal transduction. This suggests that less direct

mechanisms of epigenetic regulation involving DNA methylation may play a wider role in

mediating the long-lasting effects of both acute and chronic stress on the immune response

to pathogen challenge. These mechanisms may include DNA methylation in other features

such as lncRNAs and far-distant enhancer regions, which have variable and context-specific

regulatory effects on gene expression, as well as interactive effects between DNA

methylation and other epigenetic modifications.

Conclusions and potential implications

In summary, we found that acute and chronic stress during early life induced contrasting

effects on gill transcription and immune response in Atlantic salmon. Both stressors also

induced considerable changes in the baseline methylome and modulated similar cellular

signalling pathways, suggesting that these may be common targets of stress-induced

epigenetic regulation with the potential for far-reaching effects on cellular processes.

However, the specific patterns of methylation change at the individual CpG level were very

different between acute and chronic stress groups, suggesting distinction in fine level

epigenetic regulation. We found stress-induced changes in the methylome were only directly

associated with transcriptional differences, and transcriptional responses to LPS, for a small proportion of genes. However, at the gene-pathway level, we present evidence for stress-induced differential methylation in the key signalling and regulatory networks involved in transcriptional response to a pathogen challenge. This suggests that stress may influence the immune response through wider, less direct epigenetic mechanisms.

These results have important implications for health and disease management of natural and farmed fish populations, which are both commonly subject to multiple stressors and infection challenges. They highlight the importance of considering the long-lasting effects of early life stress, even when obvious effects on growth and condition are not apparent, and suggest that early-life stress has considerable effects on immuno-competence and disease susceptibility. This knowledge is important for risk assessment and stress management, and can perhaps even be used to harness the apparent stimulatory effects of acute stress on the immune system of Atlantic salmon and other commercially important fish. Our study provides the first evidence that direct and indirect epigenetic mechanisms may play a role in mediating the lasting effects of early-life stress in fish.

Acknowledgements

We are grateful to Sam Fieldwick for assistance with sampling and Dr Angela Marchbank for facilitating the RRBS sequencing. This work was funded by a BBSRC-NERC Aquaculture grant (BB/M026469/1) and by the Welsh Government and Higher Education Funding Council for Wales (HEFCW) through the Sêr Cymru National Research Network for Low Carbon Energy and Environment (NRN-LCEE).

References

Andrews, S. 2007 SeqMonk: A tool to visualise and analyse high throughput mapped
sequence data. *https://www.bioinformatics.babraham.ac.uk/projects/seqmonk/*.

Andrews, S. 2010 FastQC A Quality Control tool for High Throughput Sequence Data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Ankley, G. T. & Villeneuve, D. L. 2015 Temporal Changes in Biological Responses and
Uncertainty in Assessing Risks of Endocrine-Disrupting Chemicals: Insights from
Intensive Time-Course Studies with Fish. *Toxicological Sciences* **144**, 259-275.

Baerwald, M. R., Meek, M. H., Stephens, M. R., Nagarajan, R. P., Goodbla, A. M., Tomalty,
K. M. H., Thorgaard, G. H., May, B. & Nichols, K. M. 2016 Migration-related
phenotypic divergence is associated with epigenetic modifications in rainbow trout.
Molecular Ecology **25**, 1785-1800.

Bates, L., Boucher, M. & Shrimpton, J. 2014 Effect of temperature and substrate on whole
body cortisol and size of larval White Sturgeon (*Acipenser transmontanus*
Richardson, 1836). *Journal of Applied Ichthyology* **30**, 1259-1263.

Bolger, A. M., Lohse, M. & Usadel, B. 2014 Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-20.

Bonnans, C., Chou, J. & Werb, Z. 2014 Remodelling the extracellular matrix in development
and disease. *Nature Reviews Molecular Cell Biology* **15**, 786-801.

Calcagni, E. & Elenkov, I. 2006 Stress system activity, innate and T helper cytokines, and
susceptibility to immune-related diseases. *Annals of the New York Academy of*
Sciences **1069**, 62-76.

Cao-Lei, L., de Rooij, S. R., King, S., Matthews, S. G., Metz, G. A. S., Roseboom, T. J. &
Szyf, M. 2017 Prenatal stress and epigenetics. *Neuroscience & Biobehavioral*
Reviews Epub ahead of print.

Castro, R., Jouneau, L., Tacchi, L., Macqueen, D. J., Alzaid, A., Secombes, C. J., Martin, S.
A. M. & Boudinot, P. 2015 Disparate developmental patterns of immune responses to
bacterial and viral infections in fish. *Scientific Reports* **5**, 15458.

Chatterjee, A., Ozaki, Y., Stockwell, P. A., Horsfield, J. A., Morison, I. M. & Nakagawa, S.
2013 Mapping the zebrafish brain methylome using reduced representation bisulfite
sequencing. *Epigenetics* **8**, 979-89.

Chrousos, G. P. 2009 Stress and disorders of the stress system. *Nature Reviews*
Endocrinology **5**, 374-382.

Davidson, W. S., Koop, B. F., Jones, S. J., Iturra, P., Vidal, R., Maass, A., Jonassen, I., Lien,
S. & Omholt, S. W. 2010 Sequencing the genome of the Atlantic salmon *(Salmo*
salar). *Genome Biology* **11**, 403-410.

Dhabhar, F. S. 2009 Enhancing versus Suppressive Effects of Stress on Immune Function:
Implications for Immunoprotection and Immunopathology. *Neuroimmunomodulation*
16, 300-317.

Dhabhar, F. S. 2014 Effects of stress on immune function: the good, the bad, and the
beautiful. *Immunologic Research* **58**, 193-210.

Donaldson, M., Cooke, S., Patterson, D. & Macdonald, J. 2008 Cold shock and fish. *Journal*
of Fish Biology **73**, 1491-1530.

Elliott, J. 1989 The critical-period concept for juvenile survival and its relevance for
population regulation in young sea trout, *Salmo trutta*. *Journal of Fish Biology* **35**, 91-
98.

Froese, R. 2006 Cube law, condition factor and weight–length relationships: history, meta-
analysis and recommendations. *Journal of Applied Ichthyology* **22**, 241-253.

Frost, R. A., Nystrom, G. J. & Lang, C. H. 2002 Lipopolysaccharide regulates
proinflammatory cytokine expression in mouse myoblasts and skeletal muscle.
American Journal of Physiology-Regulatory, Integrative and Comparative Physiology
283, R698-R709.

Glaser, R. & Kiecolt-Glaser, J. K. 2005 Stress-induced immune dysfunction: implications for
health. *Nature Reviews Immunology* **5**, 243-251.

Hansen, T. 1985 Artificial hatching substrate: Effect on yolk absorption, mortality and growth
during first feeding of sea trout (*Salmo trutta*). *Aquaculture* **46**, 275-285.

Hu, Y., Huang, K., An, Q., Du, G., Hu, G., Xue, J., Zhu, X., Wang, C.-Y., Xue, Z. & Fan, G.
2016 Simultaneous profiling of transcriptome and DNA methylome from a single cell.
Genome Biology **17**, 88-94.

Huang, D. W., Sherman, B. T. & Lempicki, R. A. 2008 Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nature Protocols* **4**, 44-57.

Huang, Y.-Z., Sun, J.-J., Zhang, L.-Z., Li, C.-J., Womack, J. E., Li, Z.-J., Lan, X.-Y., Lei, C.-
Z., Zhang, C.-L., Zhao, X. & Chen, H. 2014 Genome-wide DNA Methylation Profiles
and Their Relationships with mRNA and the microRNA Transcriptome in Bovine
Muscle Tissue (*Bos taurine*). *Scientific Reports* **4**, 6546-6552.

Hunter, R. G., Gagnidze, K., McEwen, B. S. & Pfaff, D. W. 2015 Stress and the dynamic
genome: Steroids, epigenetics, and the transposome. *Proceedings of the National*
Academy of Sciences **112**, 6828-6833.

Jjingo, D., Conley, A. B., Yi, S. V., Lunyak, V. V. & Jordan, I. K. 2012 On the presence and
role of human gene-body DNA methylation. *Oncotarget* **3**, 462-474.

Jones, P. A. 2012 Functions of DNA methylation: islands, start sites, gene bodies and
beyond. *Nature Reviews Genetics* **13**, 484-92.

Kan, A. & Hodgkin, P. D. 2014 Mechanisms of cell division as regulators of acute immune
response. *Systems and Synthetic Biology* **8**, 215-221.

Kim, D., Langmead, B. & Salzberg, S. L. 2015 HISAT: a fast spliced aligner with low memory
requirements. *Nature Methods* **12**, 357-360.

Köberlin, Marielle S., Snijder, B., Heinz, Leonhard X., Baumann, Christoph L., Fauster, A.,
Vladimer, Gregory I., Gavin, A.-C. & Superti-Furga, G. 2015 A Conserved Circular
Network of Coregulated Lipids Modulates Innate Immune Responses. *Cell* **162**, 170-
183.

Kolde, R. 2015 Pretty Heatmaps. *R package version 3.1-131*.

Kreger, F. 2016 TrimGalore. A wrapper around Cutadapt and FastQC to consistently apply
adapter and quality trimming to FastQ files, with extra functionality for RRBS data.
https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/.

Krueger, F. & Andrews, S. R. 2011 Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics* **27**, 1571-1572.

Lemmon, M. A. & Schlessinger, J. 2010 Cell signaling by receptor tyrosine kinases. *Cell* **141**,
1117-1134.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J. & Homer, N. 2009 The Sequence
Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079.

Lien, S., Koop, B. F., Sandve, S. R., Miller, J. R., Kent, M. P., Nome, T., Hvidsten, T. R.,
Leong, J. S., Minkley, D. R., Zimin, A., Grammes, F., Grove, H., Gjuvsland, A.,
Walenz, B., Hermansen, R. A., von Schalburg, K., Rondeau, E. B., Di Genova, A.,
Samy, J. K. A., Olav Vik, J., Vigeland, M. D., Caler, L., Grimholt, U., Jentoft, S., Inge
Våge, D., de Jong, P., Moen, T., Baranski, M., Palti, Y., Smith, D. R., Yorke, J. A.,
Nederbragt, A. J., Tooming-Klunderud, A., Jakobsen, K. S., Jiang, X., Fan, D., Hu,
Y., Liberles, D. A., Vidal, R., Iturra, P., Jones, S. J. M., Jonassen, I., Maass, A.,
Omholt, S. W. & Davidson, W. S. 2016 The Atlantic salmon genome provides
insights into rediploidization. *Nature* **533**, 200-205.

Lim, Y. C., Li, J., Ni, Y., Liang, Q., Zhang, J., Yeo, G. S., Lyu, J., Jin, S. & Ding, C. 2017 A
complex association between DNA methylation and gene expression in human
placenta at first and third trimesters. *PloS One* **12**, e0181155.

Linden, S. K., Sutton, P., Karlsson, N. G., Korolik, V. & McGuckin, M. A. 2008 Mucins in the
mucosal barrier to infection. *Mucosal Immunology* **1**, 183-190.

Love, M. I., Huber, W. & Anders, S. 2014 Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550-555.

Lussier, A. A., Morin, A. M., MacIsaac, J. L., Salmon, J., Weinberg, J., Reynolds, J. N.,
Pavlidis, P., Chudley, A. E. & Kobor, M. S. 2018 DNA methylation as a predictor of
fetal alcohol spectrum disorder. *Clinical Epigenetics* **10**, 5-10.

Macqueen, D. J., Primmer, C. R., Houston, R. D., Nowak, B. F., Bernatchez, L., Bergseth,
S., Davidson, W. S., Gallardo-Escarate, C., Goldammer, T., Guiguen, Y., Iturra, P.,
Kijas, J. W., Koop, B. F., Lien, S., Maass, A., Martin, S. A. M., McGinnity, P.,
Montecino, M., Naish, K. A., Nichols, K. M., Olafsson, K., Omholt, S. W., Palti, Y.,
Plastow, G. S., Rexroad, C. E. R., Rise, M. L., Ritchie, R. J., Sandve, S. R., Schulte,
P. M., Tello, A., Vidal, R., Vik, J. O., Wargelius, A. & Yanez, J. M. 2017 Functional
Annotation of All Salmonid Genomes (FAASG): an international initiative supporting
future salmonid research, conservation and aquaculture. *BMC Genomics* **18**, 484-
490.
McGowan, P. O., Sasaki, A., D'alessio, A. C., Dymov, S., Labonté, B., Szyf, M., Turecki, G.
& Meaney, M. J. 2009 Epigenetic regulation of the glucocorticoid receptor in human
brain associates with childhood abuse. *Nature Neuroscience* **12**, 342-348.
Moghadam, H. K., Johnsen, H., Robinson, N., Andersen, Ø., H. Jørgensen, E., Johnsen, H.
K., Bæhr, V. J. & Tveiten, H. 2017 Impacts of Early Life Stress on the Methylome and
Transcriptome of Atlantic Salmon. *Scientific Reports* **7**, 5023.
Monaghan, P. & Haussmann, M. F. 2015 The positive and negative consequences of
stressors during early life. *Early Human Development* **91**, 643-647.
Muurinen, M., Hannula-Jouppi, K., Reinius, L. E., Söderhäll, C., Merid, S. K., Bergström, A.,
Melén, E., Pershagen, G., Lipsanen-Nyman, M., Greco, D. & Kere, J. 2017
Hypomethylation of HOXA4 promoter is common in Silver-Russell syndrome and
growth restriction and associates with stature in healthy children. *Scientific Reports* **7**,
15693-99.
Naba, A., Clauser, K. R., Ding, H., Whittaker, C. A., Carr, S. A. & Hynes, R. O. 2016 The
extracellular matrix: Tools and insights for the "omics" era. *Matrix Biology* **49**, 10-24.
Nasca, C., Bigio, B., Zelli, D., Nicoletti, F. & McEwen, B. S. 2015a Mind the gap:
glucocorticoids modulate hippocampal glutamate tone underlying individual
differences in stress susceptibility. *Molecular Psychiatry* **20**, 755-763.
Nasca, C., Zelli, D., Bigio, B., Piccinin, S., Scaccianoce, S., Nisticò, R. & McEwen, B. S.
2015b Stress dynamically regulates behavior and glutamatergic gene expression in
hippocampus by opening a window of epigenetic plasticity. *Proceedings of the*
National Academy of Sciences **112**, 14960-14965.
Papadopoulou, A., Siamatras, T., Delgado-Morales, R., Amin, N. D., Shukla, V., Zheng, Y.
L., Pant, H. C., Almeida, O. F. X. & Kino, T. 2015 Acute and chronic stress
differentially regulate cyclin-dependent kinase 5 in mouse brain: implications to
glucocorticoid actions and major depression. *Translational Psychiatry* **5**, e578.
Patel, J., McLeod, L. E., Vries, R. G., Flynn, A., Wang, X. & Proud, C. G. 2002 Cellular
stresses profoundly inhibit protein synthesis and modulate the states of

phosphorylation of multiple translation factors. *European Journal of Biochemistry*
269, 3076-85.

Pertea, G. 2016 gffread. *avaialble at: https://github.com/gpertea/gffread*.

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T. & Salzberg, S. L.
2015 StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nature Biotechnology* **33**, 290-295.

Pinheiro, J., Bates, D., DebRoy, S., Sarker, D. & Team, R. C. 2017 nlme: Linear and
Nonlinear Mixed Effects Models: R package version 3.1-131.

R_Core_Team. 2014 R: A language and environment for statistical computing. Vienna,
Austria.: R Foundation for Statistical Computing.

Ragkousi, K. & Gibson, M. C. 2014 Cell division and the maintenance of epithelial order. *The*
Journal of Cell Biology **207**, 181-188.

Rice, P., Longden, I. & Bleasby, A. 2000 EMBOSS: the European Molecular Biology Open
Software Suite. *Trends in Genetics* **16**, 276-277.

Rowley, M. J., Rothi, M. H., Böhmdorfer, G., Kuciński, J. & Wierzbicki, A. T. 2017 Long-
range control of gene expression via RNA-directed DNA methylation. *PLOS Genetics*
13, e1006749.

Sepulcre, M. P., Alcaraz-Pérez, F., López-Muñoz, A., Roca, F. J., Meseguer, J., Cayuela, M.
L. & Mulero, V. 2009 Evolution of lipopolysaccharide (LPS) recognition and signaling:
fish TLR4 does not recognize LPS and negatively regulates NF-κB activation. *The*
Journal of Immunology **182**, 1836-1845.

Shahmuradov, I. A., Umarov, R. K. & Solovyev, V. V. 2017 TSSPlant: a new tool for
prediction of plant Pol II promoters. *Nucleic Acids Research* **45**, e65-e65.

Silberman, D. M., Acosta, G. B. & Zorrilla Zubilete, M. A. 2016 Long-term effects of early life
stress exposure: Role of epigenetic mechanisms. *Pharmacological Research* **109**,
64-73.

Smith, B. L., Schmeltzer, S. N., Packard, B. A., Sah, R. & Herman, J. P. 2016 Divergent
effects of repeated restraint versus chronic variable stress on prefrontal cortical
immune status after LPS injection. *Brain, Behavior, and Immunity* **57**, 263-270.

Sorokin, L. 2010 The impact of the extracellular matrix on inflammation. *Nature Reviews*
Immunology **10**, 712-723.

Sundaram, A. Y., Consuegra, S., Kiron, V. & Fernandes, J. M. 2012 Positive selection
pressure within teleost Toll-like receptors tlr21 and tlr22 subfamilies and their
response to temperature stress and microbial components in zebrafish. *Molecular*
Biology Reports **39**, 8965-75.

Sundin, L., Turesson, J. & Taylor, E. W. 2003 Evidence for glutamatergic mechanisms in the
901   vagal sensory pathway initiating cardiorespiratory reflexes in the shorthorn sculpin
902   *Myoxocephalus scorpius. Journal of Experimental Biology* **206**, 867-876.

903   Tort, L. 2011 Stress and immune modulation in fish. *Developmental & Comparative*
904   *Immunology* **35**, 1366-1375.

905   Turecki, G. & Meaney, M. J. 2016 Effects of the Social Environment and Stress on
906   Glucocorticoid Receptor Gene Methylation: A Systematic Review. *Biological*
907   *Psychiatry* **79**, 87-96.

908   Uren Webster, T. M. & Santos, E. M. 2015 Global transcriptomic profiling demonstrates
909   induction of oxidative stress and of compensatory cellular stress responses in brown
910   trout exposed to glyphosate and Roundup. *BMC Genomics* **16**, 32-38.

911   Uribe, C., Folch, H., Enriquez, R. & Moran, G. 2011 Innate and adaptive immunity in teleost
912   fish: a review. *Veterinarni Medicina* **56**, 486-503.

913   Vaiserman, A. M. 2015 Epigenetic programming by early-life stress: Evidence from human
914   populations. *Developmental Dynamics* **244**, 254-265.

915   Wang, H., Beyene, G., Zhai, J., Feng, S., Fahlgren, N., Taylor, N. J., Bart, R., Carrington, J.
916   C., Jacobsen, S. E. & Ausin, I. 2015 CG gene body DNA methylation changes and
917   evolution of duplicated genes in cassava. *Proceedings of the National Academy of*
918   *Sciences* **112**, 13729-13734.

919   Wang, H., Wang, J., Ning, C., Zheng, X., Fu, J., Wang, A., Zhang, Q. & Liu, J.-F. 2017
920   Genome-wide DNA methylation and transcriptome analyses reveal genes involved in
921   immune responses of pig peripheral blood mononuclear cells to poly I:C. *Scientific*
922   *Reports* **7**, 9709-9715.

923   Wickham, H. 2009 *ggplot2: Elegant Graphics for Data Analysis*: Springer-Verlag New York.

924   Zemach, A., Kim, M. Y., Silva, P., Rodrigues, J. A., Dotson, B., Brooks, M. D. & Zilberman,
925   D. 2010 Local DNA hypomethylation activates genes in rice endosperm. *Proceedings*
926   *of the National Academy of Sciences* **107**, 18729-18734.

927   Zhang, Z. & Zhang, R. 2015 Epigenetics in autoimmune diseases: Pathogenesis and
928   prospects for therapy. *Autoimmunity Reviews* **14**, 854-63.

929   Zhao, Y., Sun, H. & Wang, H. 2016 Long noncoding RNAs in DNA methylation: new players
930   stepping into the old game. *Cell & Bioscience* **6**, 45.

931   Zhong, Z., Du, K., Yu, Q., Zhang, Y. E. & He, S. 2016 Divergent DNA Methylation Provides
932   Insights into the Evolution of Duplicate Genes in Zebrafish. *G3:*
933   *Genes|Genomes|Genetics* **6**, 3581-3591.

Figure 1. The impact of early life stress on transcriptional response to LPS.

A) Multidimensional scaling analysis illustrating the very significant effect of exposure to 20 µg/ml LPS on the entire gill transcriptome (78,229 putative loci) of fish from all stress treatment groups. **B)** Heat map illustrating the expression of all genes for which a significant interaction between stress and LPS response was identified (516 genes). Data presented are read counts for each individual normalised by library size, and by mean expression for each gene. Hierarchical clustering was performed using an Euclidian distance metric.

Figure 2. Visualisation of the Atlantic salmon gill methylome. A) Average CpG methylation percentage in gene bodies and within the 1.5 Kb upstream and downstream of the transcription start (TSS) and termination sites (TTS) for each stress group. **B-C)** Histograms of average methylation distribution within gene bodies and putative promoter regions. **D)** Heat map illustrating percentage methylation for all differentially methylated CpGs identified in each stress group (logistic regression q< 0.01 and |ΔM|>20%, and t.test p <0.01), using unsupervised hierarchical clustering.

Figure 3. Integration of transcriptome and methylome. Scatterplot and boxplot displaying mean gene expression and mean DNA methylation for **A)** putative promoters and **B)** gene bodies in control fish. **C-F)** Starburst plots displaying the effect of stress on the transcriptome and the methylome. For each type of stress relative to the control group, change in gene expression (log2fold change) is plotted against change in DNA methylation (ΔM) for **(C;E)** *putative promoters* and **(D;F)** *gene bodies*. Highlighted dots denote genes with ΔM > 5% and |FC |> 2; a full list of highlighted genes is provided in Table S5-S6.

Figure 1

Figure 2

Figure 3