Directed Bayesian Networks established functional differences between

breast cancer subtypes

Lucía Trilla-Fuertes[1¶], Andrea Zapater-Moros[2¶], Angelo Gámez-Pozo[1,2], Jorge M

Arevalillo[3], Guillermo Prado-Vázquez[1], Mariana Díaz-Almirón[4], María Ferrer-Gómez[2],

Rocío López-Vacas[2], Hilario Navarro[3], Enrique Espinosa[5,6], Paloma Maín[7], and Juan

Ángel Fresno Vara[1,2,7*]

[1]Biomedica Molecular Medicine SL, Madrid, Spain

[2]Molecular Oncology & Pathology Lab, Institute of Medical and Molecular Genetics-INGEMM,

La Paz University Hospital-IdiPAZ, Madrid, Spain

[3]Operational Research and Numerical Analysis, National Distance Education University (UNED),

Spain

[4]Biostatistics Unit, La Paz University Hospital-IdiPAZ, Madrid, Spain

[5]Medical Oncology Service, La Paz University Hospital-IdiPAZ, Madrid, Spain

[6]Biomedical Research Networking Center on Oncology-CIBERONC, ISCIII, Madrid, Spain

[7]Department of Statistics and Operations Research, Faculty of Mathematics, Complutense

University of Madrid, Madrid, Spain

¶ These authors contributed equally to this work.

*Corresponding Author:

E-mail: juanangel.fresno@salud.madrid.org (JAFV)

Short title: Directed networks established functional differences in breast cancer

Abstract

Breast cancer is a heterogeneous disease. In clinical practice, tumors are classified as hormonal receptor positive, Her2 positive and triple negative tumors. In previous works, our group defined a new hormonal receptor positive subgroup, the TN-like subtype, which has a prognosis and a molecular profile more similar to triple negative tumors. In this study, proteomics and directed Bayesian networks were used to characterize protein relationships in 106 breast tumor samples. Components obtained by these methods had a clear functional structure. The analysis of these components suggested differences in processes such as metastasis or proliferation between breast cancer subtypes, including our new subtype TN-like. In addition, one of the components, mainly related with metastasis, had prognostic value in this cohort. Functional approaches allow to build hypotheses about regulatory mechanisms and to establish new relationships among proteins in the breast cancer context.

Author Summary:

Breast cancer classification in the clinical practice is defined by three biomarkers (estrogen receptor, progesterone receptor and HER2) into hormone receptor positive, HER2+ and triple negative breast cancer (TNBC). Our group recently described a new ER+ subtype with molecular characteristics and prognosis similar to TNBC. In this study we propose a mathematical method, the directed Bayesian networks, as a useful tool to study protein interactions and differential biological processes in breast cancer subtypes, characterizing differences in relevant processes such as proliferation or metastasis and associated them with patient prognosis.

Introduction

Breast cancer is one of the most prevalent cancers in the world [1]. In clinical practice, breast cancer is classified according to the expression of hormonal receptors (estrogen or progesterone) and Her2, into positive hormonal receptor (ER+), HER2+ and triple negative (TNBC). In previous studies, our group defined a new ER+ molecular subgroup, named TN-like, with a molecular profile and a prognosis more similar to TNBC tumors [2]. The remaining ER+ tumors were considered as ER-true. We also found significant molecular differences among breast cancer subtypes. For instance, differences related with metabolism of glucose were described between ER-true, TN-like and TNBC tumors [2, 3].

Proteomics provides useful information about biological process effectors and may quantify thousands of proteins. Undirected probabilistic graphical models (PGM), based on a Bayesian approach, allow characterizing differences between tumor samples at functional level [2-5]. In this study we explored the utility of directed Bayesian networks in the molecular characterization of breast cancer. The main feature of targeted Bayesian networks is that they provide a hierarchical structure and targeted relationships between proteins.

In this work, we used proteomics and directed Bayesian networks to characterize protein relationships in a cohort of breast cancer tumor samples. These networks maintain a functional structure and it is possible to use this information to build prognostic signatures. This approach also reflected previously described interactions and it could be used to propose new hypotheses and mechanisms of regulation of these proteins.

Results

Patient characteristics

Clinical characteristics of this patient cohort have been previously described [2, 3, 6]. Briefly, one hundred and six patients were enrolled into the study. They all had node positive disease, Her2 negative and all had received adjuvant chemotherapy and hormonal therapy in the case of ER+ tumors. Among ER+ tumors, 50 patients were characterized as ER-true and 21 were defined as TN-like (Sup Table 1) [2].

MS analysis

Proteomics analyses from these samples has been previously described [3]. In summary, one hundred and two FFPE samples had enough protein to perform the MS analyses. After MS workflow, 96 samples provided useful protein expression data. After quality criteria, 1,095 proteins presented at least two unique peptides and detectable expression in at least 75% of the samples in at least one type of sample (either ER+ or TNBC).

Directed networks

Using proteomics data, directed acyclic graphs (DAG) were performed. Altogether, it was possible to establish 536 edges of which 414 are guided and 122 are undetermined. These edges formed 377 components formed by different number of nodes or proteins. An overview of the number of nodes (proteins) included in each component is provided in table 1.

Number of nodes	1	2	3	4	5	6	7	8	9	10	11	13	14	19	21	26	27	34	36
Number of components	229	70	24	12	15	2	6	3	4	2	1	2	1	1	1	1	1	1	1

Table 1: Characteristics of the components obtained from DAG. Number of nodes = number of proteins contained in each component, Number of components = directed components obtained.

We characterized components from DAG analysis. Components including less than 9 nodes

were dismissed because they were little informative. All components were named with the

number of nodes included by the DAG analysis, and the information was completed with

protein-protein interactions (PPI) based on experimental evidence obtained from Genes2FANS

(G2F) [7]. For example, component 14 includes 19 nodes, 14 nodes defined from our directed

Bayesian analysis and 5 extra nodes added by G2F.

Afterwards, components were interrogated for biological function. For components with less

than 27 nodes provided by DAG analysis, this functional analysis was performed by

bibliographical review. Thus, the main biological function of component 14 is metabolism.

Regarding components which had more than 27 nodes provided by the DAG analysis, gene

ontology analyses were done once they were completed by G2F information. Characteristics

about all components are supplied in table 2 and Supplementary File 1.

Component	Main function	Total number of nodes
Component 36	Extracellular exosome	65 nodes
Component 34	RNA processing and mTOR pathway	61 nodes
Component 19	Proliferation	26 nodes
Component 21	Proliferation & metastasis	34 nodes
Component 27	Metastasis	34 nodes
Component 14	Metabolism	19 nodes
Component 13	Proteasome	19 nodes
Component 11	None main function assigned	14 nodes
Component 10b	Proliferation	19 nodes
Component 9a	Growth	15 nodes
Component 9b	Proliferation & metastasis	22 nodes
Component 9c	Glycolysis	12 nodes
Component 9d	Transcription & ribosomes	20 nodes

96 **Table 2: Features of components obtained by DAG and G2F analysis.**

Component activity measurements

Component activities were calculated for each node. There were significant differences

between ER-true, TN-like and TNBC tumors in the component activity for component 34:

mRNA processing and mTOR, component 36: extracellular exosome, component 21:

proliferation and metastasis, component 27: metastasis, component 9a: growth, component

9b: proliferation and metastasis, component 9c: glycolysis, component 10b: metastasis,

component 13: proteasome and component 9d: transcription (Fig 1).

Fig 1: Component activity measurements for ER-true, TN-like and TNBC respectively.

Component 10b: metastasis

Component 10b activity showed prognostic value in our series, splitting our population into a

high and a low risk group and it can be used to make a distant metastasis-free survival (DMFS)

predictor (p=0.0068, HR= 0.2854, cut-off 40:60) (Fig 2). Additionally, dividing by molecular

subtype, this prognostic signature also split the population into low and high risk groups

although it is not statistically significant (Fig 3).

Fig 2: Component 10b activity prognostic value in the whole cohort.

Fig 3: Component 10b activity prognostic value by subtype.

Component 10b contains 10 proteins. Four out of nine edges (DYNLRB1-HSPH1; HSPH1-CFL1;

HBB-UCHL5; and UCHL5-CFL1) have been previously described in G2F database [7]. Then, G2F

added two more proteins (HSPH1 and UCHL5) to this component. Most of the proteins

included in this component are related with metastasis processes [8-12], whereas PBDC1

doesn´t have any associated function. On the other hand, HIST2H2BF is a histone and

HIST2H3PS2 is a histone pseudogene showing a directed relationship (Fig 4).

Fig 4: Component 10b merged with PPI information provided by Genes2FANS. Red arrows

indicate relationships from DAG; green lines indicate relationships from Genes2FANS database.

Orange nodes are common nodes between these two approaches, purple nodes are nodes

which come only from DAG and blue nodes are only from Genes2FANS.

Discussion

In this study, we used proteomics and DAG to characterize relationships between proteins in

breast cancer tumor samples. Unlike other approaches, such as G2F [7], our DAG method

supplies directed relationships between proteins and a hierarchical structure. Traditionally, PPI

networks are based in relationships described in the literature. However, we built a directed

network, i.e. a graph formed by edges with a direction, using protein expression data without

other *a priori* information, so it was possible to propose new hypotheses about protein

interactions. We used probabilistic graphical models (PGM) because they offer a way to relate

many random variables with a complex dependency structure.

Arrows in directed networks indicate causality between two proteins, i.e. if protein A and B are

connected and protein A changes its expression value, protein B changes its expression value

as well. This approach allows making hypotheses about causal relationships between proteins

and proposes a hierarchical structure. G2F relationships supplies additional information to our

directed networks, so it served as a validation of the network coherence, i.e., in some cases, an

experimental relationship between two proteins connected in the directed network had been

previously described. In component 10b, for instance, CFL1 and DYNRBL1 had a described

common nexus in HSPH1 [7]. It is remarkable that, in component 27, DAG analysis established

a relationship between PTRF and PRDXCDBP which is widely described [13].

We demonstrated in previous works that non-directed Bayesian graphs provided functional

information [2-5]. Interestingly, a functional structure also appeared in this type of networks.

Component activities suggested differences in functions such as metastasis, proliferation,

proteasome or glycolysis. We have previously described differences in proteasome and

glycolysis between ER-true and TN-like subtypes using non-directed Bayesian networks [2].

On the other hand, it is widely known the role that actin-cytoskeleton plays and its regulation

in directional migration and metastasis, Therefore, it is not surprising that the proteins that

make up component 10b had some relation to this biological function and had a prognostic

value. Using component 10b activity, based on these proteins related with metastasis, it is

possible to split our population into a low and a high risk of relapse groups and, interestingly,

this prognostic value is also showed in the analysis by subtype. In previous studies we have

used functional node activities from non-directed Bayesian network to develop prognostic

predictors [4, 5]. Now, this approach is also validated in directed networks.

Component 10b presents three proteins showing influence over others: HIST2H2BF, DYNLRB1

and RHOG. Dynein light chain roadblock-type 1 (DYNRBL1), also known as km23-1, is a

component of the cytoplasmic dynein 1 complex. In colon cells, its depletion can block events

known to be involved in cell invasion and tumor metastasis, and it is also an actin cytoskeletal

linker critical for the dynamic regulation of cell motility and invasion since it induces a highly

organized actin stress fiber network [14]. DYNRBL1 regulates RhoA and motility-associated

actin modulating proteins, such as cofilin1 (CFL1) and coronin [15], suggesting that DYNRBL1

may represent a novel target for anti-metastatic therapy [9].

On the other hand, Ras-homolog family member G (RhoG) encodes a member of the Rho

family of small GTPases. Constitutively active RhoG induces morphological and cytoskeletal

changes similar to those induced by the combination of active Rac and Cdc42 working

upstream of them. Rac is activated at the leading edge of motile cells and induces the

formation of actin-rich lamellipodia protrusions, which serves as a major driving force of

cancer cell. The major downstream proteins for Rac are the WAVE family proteins, the

activators of the Arp2/3 complex. In addition, RhoG induces translocation of Dock4-ELMO

complex to the plasma membrane and enhances Rac1 activation which promotes migration

[16]. Moreover, RNA interference-mediated knockdown of RhoG in HeLa cells reduced cell

migration in Transwell and scratch-wound migration assays [11].

HIST2H2BF has been proposed as a pancreatic ductal adenocarcinoma biomarker[17], but

there is no available information about its role in metastasis or breast cancer.

In component 10b, both DYNLRB1 and RHOG expression modulates cofilin1 (CFL1). CFL1 is a

widely distributed intracellular actin-modulating protein that binds and depolymerizes

filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent

manner. Increased phosphorylation of this protein by LIM kinase aids in Rho-induced

reorganization of the actin cytoskeleton [18]. CFL1 is an actin-severing protein that creates free

barbed ends in the actin-severing process. Arp2/3 binding to these barbed ends allows the

elongation of new actin filaments. This synergy between CFL1 severing activity and Arp2/3-

generated dendritic nucleation resulting in the formation of stable invadopods and a

directional migration, linking CFL1and ARP2/3, through RhoG, to tumor cell invasion [15, 19].

CFL1 pathway was previously related with metastases processes in breast cancer [18].

Furthermore, DYNRBL1 is required for TGFb1 secretion. It seems to be TGFb pathway plays an

important role in cell migration in this sense, because TGFbRII signaling regulates Rho GTPase

degradation and actin dynamics. Moreover, TGFb pathway induces expression of the GEF NET1

which accumulates and promotes actin polymerization and on the other hand, it induces

expression of tropomyosins that regulate assembly of a contractile apparatus that serves the

motility of the responding cell during the EMT process [15] Additionally, TGFß production by

macrophages or dendritic cells (DCs) that have engulfed apoptotic cells can promote the

generation of inducible regulatory T (Tregs) that play a known protumoral role [16].

Therefore, the relationships between CFL1, RHOG and DYNLRB1 are well-established.

Interestingly, DAG graph adds the decorin (DCN) to these edges. DCN is a small leucine-rich

proteoglycan that promotes matrix organization by decreasing collagen uptake/degradation,

which constitutes a physical barrier against migration/motility of cancerous cells. The

alteration of the matrix stiffness leads to differential integrin activation and changes in

cytoskeletal organization by Rac, affecting cell motility and invasiveness [20]. Moreover, DCN

acts as a matrikine whose interaction with CXC chemokine receptor 4 (CXCR4) impairs its bind

with stromal cell-derived factor-1a (SDF-1a), preventing directional migration [20]. In breast

cancer tumors, high DCN expression in stroma correlated with lower tumor grade, low Ki67

levels and ER positivity. On the other hand, high expression of DCN in the malignant epithelium

was correlated with LN positivity, higher number of positive lymph nodes and HER2-positive

status compared to patients with low DCN expression [21].

By last, lysyl-tRNA synthetase (KARS), also known by KRS, was recently shown to induce cancer

cell migration through its interaction with the 67-kDa laminin receptor (67LR) who binds

laminin on ECM. The interaction of KRS with 67LR enhances the membrane stability of 67LR,

which in turn results in an increase laminin-dependent cell migration in metastasis [18]. KARS,

causes incomplete epithelial-mesenchymal transition and ineffective cell‑extracellular matrix

adhesion for migration [22].

We used a mathematical method as DAG analysis and applied them to proteomics data of

breast cancer tumors in order to infer causal relationships between these proteins. This

method supplied some known relationships but also proposed new ones. Additionally, it

associated proteins with a similar function. Therefore, it seems that it is a good approach to

propose new hypotheses about mechanisms of action. Moreover, it was possible to associate

the results obtained by DAG analysis with prognosis and built a prognostic signature. As far we

know, this is the first time that this type of analysis is applied to clinical data and is associated

with clinical outcome.

Our study has some limitations. Proteomics provides complementary information to other

techniques such as genomics. However, an improvement in the number of detected proteins is

still necessary. Validation at the cellular level is also needed.

To sum up, in this study, we used proteomics and directed Bayesian networks to characterize

relationships between proteins in breast cancer tumors. This approach reflected some

previously described interactions and it could be used to propose new hypotheses and

mechanisms of action.

Material and methods

Samples

One hundred and six FFPE samples from patients with breast cancer were recovered from I+12

Biobank and from IdiPAZ Biobank, both integrated in the Spanish Hospital Biobank Network.

The histopathological characteristics were reviewed by a pathologist to confirm tumor

content. Samples had to include no less than half of tumor cells. The endorsement of the

study was obtained by Hospital Doce de Octubre and Hospital Universitario La Paz Ethical

Committees. These samples were utilized in previous studies [2, 3, 6].

Protein preparation

Proteins were extracted from FFPE samples as previously described [23]. Briefly, FFPE sections

were deparaffinized in xylene and washed twice with absolute ethanol. Protein extracts from

FFPE samples were set up in 2% SDS buffer using a protocol based on heat-induced antigen

retrieval. Protein concentration was quantified using the MicroBCA Protein Assay Kit (Pierce-

Thermo Scientific). Protein extracts (10 µg) were processed with trypsin (1:50) and SDS was

removed from digested lysates using Detergent Removal Spin Columns (Pierce). Peptide

samples were additionally desalted using ZipTips (Millipore), dried, and resolubilized in 15 µL

of a 0.1% formic acid and 3% acetonitrile solution before MS experiments.

Label-free proteomics

Samples were analyzed on a LTQ-Orbitrap Velos hybrid mass spectrometer (Thermo Fischer

Scientific, Bremen, Germany) coupled to NanoLC-Ultra system (Eksigent Technologies, Dublin,

CA, USA) as described previously [2, 3]. Briefly, after separation, peptides were eluted with a

gradient of 5 to 30% acetonitrile in 95 minutes. The mass spectrometer was operated in data-

dependent mode (DDA), followed by CID (collision-induced dissociation) fragmentation on the

twenty most intense signals per cycle. The acquired raw MS data were processed by MaxQuant

(version 1.2.7.4) [24], followed by protein identification using the integrated Andromeda

search engine [25]. Briefly, spectra were searched against a forward UniProtKB/Swiss-Prot

database for human, concatenated to a reversed decoyed fasta database (NCBI taxonomy ID

9606, release date 2011-12-13). The maximum false discovery rate (FDR) was set to 0.01 for

peptides and 0.05 for proteins. Label free quantification was calculated on the basis of the

normalized intensities (LFQ intensity). Quantifiable proteins were defined as those detected in

at least 75% of samples in at least one type of sample (either ER+ or TNBC samples) showing

two or more unique peptides. Only quantifiable proteins were considered for subsequent

analyses. Protein expression data were log2 transformed and missing values were replaced

using data imputation for label-free data, as explained in [26], using default values. Finally,

protein expression values were z-score transformed. Batch effects were estimated and

corrected using ComBat [27]. All the mass spectrometry raw data files acquired in this study

may be downloaded from Chorus (http://chorusproject.org) under the project name Breast

Cancer Proteomics.

Network construction

PGM are graph-based representations of joint probability distributions where nodes represent

random variables and edges (directed or undirected) represent stochastic dependencies

among the variables. In particular, we have used a type of PGM called Bayesian networks (BN)

[28]. With these models, the dependences between the variables in our data are specified by a

directed acyclic graph (DAG).

Firstly, we find the BN that best explains our data [29]. There are different algorithms to learn

a DAG from data but we have selected the well-known PC algorithm, a constraint-based

structure learning algorithm [30] based on conditional independence tests. The PC algorithm

was shown to be consistent in some high-dimensional settings [31] and have some other

statistical properties which make it very useful. In this way, our data are represented by a large

graph that can be partitioned into several connected components. Then, we focused on

finding suitable subgraphs that give us a much more clear understanding of the interrelations

therein. All these procedures are implemented in R [32] within packages *pcalg* [31] and *graph*

[33]. We used protein expression data without other *a priori* information.

With the aim of compare and complete the information provided by BN, the tool Genes2FANS

(G2F), developed by Ma'ayan 's group was used [7]. This software provides information about

protein-protein interactions (PPI) based on experimental evidence. A PPI network using G2F

was built for each component and finally, both networks, the DAG and the PPI network, were

merged.

Gene Ontology Analyses

Protein to Gene Symbol conversion was performed using Uniprot (www.uniprot.org) and

DAVID (www.david.ncifcrf.gov) [34]. Gene Ontology Analysis was also done in DAVID selecting

only "*Homo sapiens*" background and GOTERM-FAT, Biocarta and KEGG databases. As

concerns small connected components, a search in the literature was done so as to establish

component main functions.

Component activity measurements

Component activities were calculated as previously described [3, 4]. Briefly, activity

measurement was calculated by the mean expression of all the proteins of each component

related with the established major component function.

Statistical analyses

Network analyses were performed using Cytoscape software [35]. Statistical analyses were

done in GraphPad Prism v6. Prognostic signatures were developed using R v3.2.4 and BRB

Array Tools, developed by Richard Simon and BRB Array Tools Development Team [36]. P-

values under 0.05 were considered statistically significant.

Funding statement

This study was supported by Instituto de Salud Carlos III, Spanish Economy and Competitiveness Ministry, Spain and co-funded by the FEDER program, "Una forma de hacer Europa" (PI15/01310). LT-F is supported by the Spanish Economy and Competitiveness Ministry (DI-15-07614). GP-V is supported by Conserjería de Educación, Juventud y Deporte of Comunidad de Madrid (IND2017/BMD7783). The funders had no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions

All the authors have directly participated in the preparation of this manuscript and have approved the final version submitted. RL-V prepared the proteomics samples. JMA, MD-A, HN and PM contributed the directed graphical models. LT-F, AZ-M, AG-P, G-PV and MF-G performed the statistical analyses, the directed graphical model interpretation and the review of the literature. LT-F, AG-P, JAFV, and EE conceived of the study and participated in its design and interpretation. LT-F and AZ-M drafted the manuscript. AG-P, JAFV, and EE supported the manuscript drafting. AG-P and JAFV coordinated the study.

Competing interests

JAFV, EE and AG-P are shareholders in Biomedica Molecular Medicine SL. LT-F and GP-V are employees of Biomedica Molecular Medicine SL. The other authors declare no competing interests.

Data Availability Statement

All the mass spectrometry raw data files acquired in this study may be downloaded from Chorus (http://chorusproject.org) under the project name Breast Cancer Proteomics.

References

1.   Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. Int J Cancer. 2015;136(5):E359-86. Epub 2014/10/09. doi: 10.1002/ijc.29210. PubMed PMID: 25220842.

2.   Gámez-Pozo A, Trilla-Fuertes L, Berges-Soria J, Selevsek N, López-Vacas R, Díaz-Almirón M, et al. Functional proteomics outlines the complexity of breast cancer molecular subtypes. Scientific Reports. 2017;7(1):10100. doi: 10.1038/s41598-017-10493-w.

3.   Gámez-Pozo A, Berges-Soria J, Arevalillo JM, Nanni P, López-Vacas R, Navarro H, et al. Combined label-free quantitative proteomics and microRNA expression analysis of breast cancer unravel molecular differences with clinical implications. Cancer Res; 2015. p. 2243-53.

4.   de Velasco G, Trilla-Fuertes L, Gamez-Pozo A, Urbanowicz M, Ruiz-Ares G, Sepúlveda JM, et al. Urothelial cancer proteomics provides both prognostic and functional information. Sci Rep. 2017;7(1):15819. Epub 2017/11/17. doi: 10.1038/s41598-017-15920-6. PubMed PMID: 29150671; PubMed Central PMCID: PMCPMC5694001.

5.   Trilla-Fuertes L, Gamez-Pozo A, M Arevalillo J, Diaz-Almiron M, Prado-Vazquez G, Zapater-Moros A, et al. Molecular characterization of breast cancer cell response to metabolic drugs. Oncotarget2018.

6.   Gámez-Pozo A, Trilla-Fuertes L, Prado-Vázquez G, Chiva C, López-Vacas R, Nanni P, et al. Prediction of adjuvant chemotherapy response in triple negative breast cancer with discovery and targeted proteomics. PLoS One. 2017;12(6):e0178296. Epub 2017/06/08. doi: 10.1371/journal.pone.0178296. PubMed PMID: 28594844; PubMed Central PMCID: PMCPMC5464546.

7.   Dannenfelser R, Clark NR, Ma'ayan A. Genes2FANs: connecting genes through functional association networks. BMC Bioinformatics. 2012;13:156. doi: 10.1186/1471-2105-13-156. PubMed PMID: 22748121; PubMed Central PMCID: PMCPMC3472228.

8.   Kim DG, Choi JW, Lee JY, Kim H, Oh YS, Lee JW, et al. Interaction of two translational components, lysyl-tRNA synthetase and p40/37LRP, in plasma membrane promotes laminin-dependent cell migration. FASEB J. 2012;26(10):4142-59. Epub 2012/07/02. doi: 10.1096/fj.12-207639. PubMed PMID: 22751010.

9.   Jin Q, Pulipati NR, Zhou W, Staub CM, Liotta LA, Mulder KM. Role of km23-1 in RhoA/actin-based cell migration. Biochem Biophys Res Commun. 2012;428(3):333-8. Epub 2012/10/15. doi: 10.1016/j.bbrc.2012.10.047. PubMed PMID: 23079622; PubMed Central PMCID: PMCPMC3513371.

10.   Madak-Erdogan Z, Ventrella R, Petry L, Katzenellenbogen BS. Novel roles for ERK5 and cofilin as critical mediators linking ERα-driven transcription, actin reorganization, and invasiveness in breast cancer. Mol Cancer Res. 2014;12(5):714-27. Epub 2014/02/06. doi: 10.1158/1541-7786.MCR-13-0588. PubMed PMID: 24505128; PubMed Central PMCID: PMCPMC4020978.

11.   Katoh H, Hiramoto K, Negishi M. Activation of Rac1 by RhoG regulates cell migration. J Cell Sci. 2006;119(Pt 1):56-65. Epub 2005/12/08. doi: 10.1242/jcs.02720. PubMed PMID: 16339170.

12.   Järvinen TA, Prince S. Decorin: A Growth Factor Antagonist for Tumor Growth Inhibition. Biomed Res Int. 2015;2015:654765. Epub 2015/11/30. doi: 10.1155/2015/654765. PubMed PMID: 26697491; PubMed Central PMCID: PMCPMC4677162.

13.   Mohan J, Morén B, Larsson E, Holst MR, Lundmark R. Cavin3 interacts with cavin1 and caveolin1 to increase surface dynamics of caveolae. J Cell Sci. 2015;128(5):979-91. Epub 2015/01/14. doi: 10.1242/jcs.161463. PubMed PMID: 25588833.

14.   Jin Q, Ding W, Mulder KM. The TGFβ receptor-interacting protein km23-1/DYNLRB1
plays an adaptor role in TGFβ1 autoinduction via its association with Ras. Journal of Biological
Chemistry. 2012;287(31):26453-63.
15.   Moustakas A, Heldin C-H. Dynamic control of TGF-β signaling and its links to the
cytoskeleton. FEBS letters. 2008;582(14):2051-65.
16.   Green DR, Ferguson T, Zitvogel L, Kroemer G. Immunogenic and tolerogenic cell death.
Nature reviews Immunology. 2009;9(5):353.
17.   Castillo J, Bernard V, San Lucas FA, Allenson K, Capello M, Kim DU, et al. Surfaceome
profiling enables isolation of cancer-specific exosomal cargo in liquid biopsies from pancreatic
cancer patients. Ann Oncol. 2017. Epub 2017/09/25. doi: 10.1093/annonc/mdx542. PubMed
PMID: 29045505.
18.   Cho HY, Ul Mushtaq A, Lee JY, Kim DG, Seok MS, Jang M, et al. Characterization of the
interaction between lysyl-tRNA synthetase and laminin receptor by NMR. FEBS letters.
2014;588(17):2851-8.
19.   Hiramoto K, Negishi M, Katoh H. Dock4 is regulated by RhoG and promotes Rac-
dependent cell migration. Experimental cell research. 2006;312(20):4205-16.
20.   Feugaing DDS, Götte M, Viola M. More than matrix: the multifaceted role of decorin in
cancer. European journal of cell biology. 2013;92(1):1-11.
21.   Cawthorn TR, Moreno JC, Dharsee M, Tran-Thanh D, Ackloo S, Zhu PH, et al. Proteomic
analyses reveal high expression of decorin and endoplasmin (HSP90B1) are associated with
breast cancer metastasis and decreased survival. PloS one. 2012;7(2):e30992.
22.   Nam SH, Kang M, Ryu J, Kim HJ, Kim D, Kim DG, et al. Suppression of lysyl-tRNA
synthetase, KRS, causes incomplete epithelial-mesenchymal transition and ineffective
cell‑extracellular matrix adhesion for migration. Int J Oncol. 2016;48(4):1553-60. Epub
2016/02/08. doi: 10.3892/ijo.2016.3381. PubMed PMID: 26891990.
23.   Gámez-Pozo A, Ferrer NI, Ciruelos E, López-Vacas R, Martínez FG, Espinosa E, et al.
Shotgun proteomics of archival triple-negative breast cancer samples. Proteomics Clin Appl.
2013;7(3-4):283-91. doi: 10.1002/prca.201200048. PubMed PMID: 23436753.
24.   Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol.
2008;26(12):1367-72. Epub 2008/11/26. doi: 10.1038/nbt.1511. PubMed PMID: 19029910.
25.   Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M. Andromeda: a
peptide search engine integrated into the MaxQuant environment. J Proteome Res.
2011;10(4):1794-805. Epub 2011/01/25. doi: 10.1021/pr101065j. PubMed PMID: 21254760.
26.   Deeb SJ, D'Souza RC, Cox J, Schmidt-Supprian M, Mann M. Super-SILAC allows
classification of diffuse large B-cell lymphoma subtypes by their protein expression profiles.
Mol Cell Proteomics. 2012;11(5):77-89. Epub 2012/03/21. doi: 10.1074/mcp.M111.015362.
PubMed PMID: 22442255; PubMed Central PMCID: PMCPMC3418848.
27.   Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data
using empirical Bayes methods. Biostatistics. 2007;8(1):118-27. doi:
10.1093/biostatistics/kxj037. PubMed PMID: 16632515.
28.   Pearl J. Probabilistic reasoning in intelligent systems: networks of plausible inference.
Morgan Kaufmann2014.

29.   Neapolitan RE. Learning Bayesian Networks . Series in Artificial Intelligence. Prentice
Hall2004.

30.   Spirtes P, Glymour C, Scheines R. Causation, Prediction, and Search. Adaptive
Computation and Machine Learning. 2nd ed. The MIT Press2000.

31.   Kalisch M, Maechler M, Colombo D, Maathius MH, Bluehlmann P. Causal Inference
Using Graphical Models with the R Package pcalg. Journal of Statistical Software2012. p. 1-26.

32.   R Core Team. R: A language and environment for statistical computing. Vienna,Austria.
R Foundation for Stattistical Computing, 2013.
33.   Gentleman R, Whalen E, Huber W, Falcon S. graph: A package to handle graph data
structures.   R package version   1.54.0.

34.   Huang dW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. Nat Protoc. 2009;4(1):44-57. doi:
10.1038/nprot.2008.211. PubMed PMID: 19131956.
35.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
software environment for integrated models of biomolecular interaction networks. Genome
Res. 2003;13(11):2498-504. doi: 10.1101/gr.1239303. PubMed PMID: 14597658; PubMed
Central PMCID: PMCPMC403769.
36.   Simon R. Roadmap for developing and validating therapeutically relevant genomic
classifiers. J Clin Oncol. 2005;23(29):7332-41. Epub 2005/09/06. doi:
10.1200/JCO.2005.02.8712. PubMed PMID: 16145063.

433   Supporting information captions

434   **Sup Table 1: Clinical characteristics of patient cohort.**

435   **Sup File 1: Components defined by DAG analysis.**

ER-true

P value	0.0823	
Hazard Ratio (logrank)		A/B
Ratio (and its reciprocal)		0.3282
95% CI of ratio		0.1067 to 1.140

TN-like

P value	0.2409	
Hazard Ratio (logrank)		A/B
Ratio (and its reciprocal)		0.0
95% CI of ratio		0.04082 to 2.232

TNBC

P value	0.1873	
Hazard Ratio (logrank)		A/B
Ratio (and its reciprocal)		0.3650
95% CI of ratio		0.1100 to 1.530