1

Empirical single-cell tracking and cell-fate simulation reveal dual roles of p53 in tumor

suppression

Ann Rancourt[1,2], Sachiko Sato[1,3,4], and Masahiko S. Satoh[2,3,4]*

[1]Glycobiology and Bioimaging Laboratory of Research Center for Infectious Diseases and

Research Centre of CHU de Québec, [2]Laboratory of DNA Damage Responses and

Bioimaging, Research Centre of CHU de Québec, [3]The Bioimaging Platform of Research

Centre of CHU de Québec and [4]Faculty of Medicine, Laval University, 2705 Boulevard

Laurier, Québec, G1V 4G2, Canada

Corresponding author: Masahiko S. Satoh

TEL: 1-418-525-4444 ext. 47340

FAX: 1-418-654-2159

e-mail: masahiko.sato@crchudequebec.ulaval.ca

ABSTRACT

The tumor suppressor p53 regulates various stress responses via increasing its cellular levels. The

lowest p53 levels occur in unstressed cells; however, the functions of these low levels remain unclear.

To investigate the functions, we used empirical single-cell tracking of p53-expressing (Control) cells

and cells in which p53 expression was silenced by RNA interference (p53 RNAi). Here we show that

p53 RNAi cells underwent more frequent cell death and cell fusion, which further induced multipolar

cell division to generate aneuploid progeny. Those results suggest that the low levels of p53 in

unstressed cells indeed have a role in suppressing the induction of cell death and the formation of

aneuploid cells. We further investigated the impact of p53 silencing by developing an algorithm to

simulate the fates of individual cells. Simulation of the fate of aneuploid cells revealed that these cells

could propagate to create an aneuploid cell population. In addition, the simulation also revealed that

more frequent induction of cell death in p53 RNAi cells under unstressed conditions conferred a

disadvantage in terms of population expansion compared with Control cells, resulting in faster

expansion of Control cells compared with p53 RNAi cells, leading to Control cells predominating in

mixed cell populations. In contrast, the expansion of Control cells, but not p53 RNAi cells, was

suppressed when the damage response was induced, allowing p53 RNAi cells to expand their

population compared with the Control cells. These results suggest that, although p53 could suppress the

formation of aneuploid cells, which could have a role in tumorigenesis, it could also allow the

expansion of cells lacking p53 expression when the damage response is induced. p53 may thus play a

role in both the suppression and the promotion of malignant cell formation during tumorigenesis.

INTRODUCTION

The p53 gene is mutated in > 50% of cancers [1-3], and loss of p53 function is considered to be involved

in tumor progression, thereby defining *TP53* as a tumor suppressor gene [4, 5]. The cellular functions

mediated by p53 are mainly related to its cellular levels, which increase in response to stress, thus

activating various mechanisms e.g. cell cycle arrest, cell senescence, cell death, and DNA damage

responses [5-9]. Given that cell cycle arrest, cell senescence, and cell death could lead to the removal of

the damaged cells, these processes could act to suppress tumor formation [6, 10-15].

p53 levels are regulated via a mechanism mediated by Mdm2, which constantly degrades p53 by

ubiquitination to maintain a low level. p53 accumulation occurs when Mdm2 activity is suppressed by

stress-induced phosphorylation, which inhibits its ubiquitination activity, resulting in inhibition of p53

degradation [4, 6, 7, 16-18]. These observations suggest that increased stress levels are required to elevate

p53 levels sufficiently to allow it to exert its tumor suppressor function. Maintaining a background low

level of p53 may thus have evolved as a means of allowing cells to respond quickly to stress.

Alternatively, given that removing p53 by gene silencing increases the efficiency of induced

pluripotent stem cell formation [19], maintaining low levels of p53 in unstressed cells *per se* may have

functional implications. With this regard, it has been suggested that low levels of p53 activity may be

required to respond to a broad spectrum of stress [20] through suppression of apoptosis and regulation of

metabolisms [4, 6, 21, 22]. If low levels of p53 indeed play a role in regulating cellular functions, it is

plausible that loss of these functions could be involved in tumorigenesis. However, the effects of loss

of low levels of p53 on cell characteristics remain unclear.

In this study, we silenced the expression of low levels of p53 using small interfering RNA (siRNA)

and investigated the effects of this loss on the frequency of induction of cellular events using single-cell

tracking to detect rare cellular events and creating a cell-lineage database revealing the fates of

individual cells. Cell lineages can be generated by lineage-reconstruction approaches; however, these

assume that all cell divisions are bipolar [23-25], and thus cannot be used to analyze lineages including

cells that undergo cell death, multiple cell division, and cell fusion. To overcome this problem, we

developed a single-cell tracking method that can detect cell death, multipolar cell division, and cell

fusion events in individual cells, thus allowing the generation of accurate cell-lineage data. In addition

to silencing, we also used the alkylating agent, N-methyl -N'-nitroso-N-nitrosoguanidine (MNNG),

which acts as a DNA damaging agent by generating methylated bases[26-30]. Among MNNG-induced

methylated bases, O^6-methylguanine, which acts as a premutagenic DNA lesion that induces a G:C to

A:T transversion mutation, is repaired by O^6-methylguanine methyltransferase, and 7-methylguanine,

which causes the induction of cell death through the formation of DNA breaks, is repaired by base

excision repair [26-30]. It has been reported that MNNG can increase p53 levels in cells [31, 32] and activate

DNA repair processes that are not related to p53. We, therefore, studied how the fate trajectories of

wild-type and p53-silenced cells were affected by p53-mediated and non-mediated processes at single-

cell resolution. Furthermore, because single-cell tracking can reveal spatiotemporal information on

individual cells, we used this information to simulate the fates of individual cells beyond the empirical

limit of cell culture, by developing a cell-fate simulation algorithm.

Our results obtained by single-cell tracking and cell-fate simulation suggest that the low levels of

p53 in unstressed cells play a role in suppressing the induction of cell death and cell fusion, which may

lead to multipolar cell division and the production of aneuploid progeny. Cell-fate simulation analysis

revealed that some aneuploid progeny derived from p53-silenced (p53 RNAi) cells could propagate in

an environment dominated by p53-expressing (Control) cells, while p53 RNAi cells *per se* were unable

to gain an advantage over Control cells in terms of population expansion. In contrast, the balance of

expansion between Control and p53 RNAi cells was altered by induction of the damage response by

MNNG, resulting in the relative expansion of p53 RNAi cells. Thus, although low levels of p53 could

act as a tumor suppressor by inhibiting the formation of aneuploid cells, this role could largely depend

on the status of the cell population harboring the cells with impaired p53 function, and the stress-

damaged environment.

RESULTS

System to investigate the functional implications of maintaining low levels of p53 in unstressed

cells

We developed a system to determine if the low levels of p53 retained in unstressed cells had a

functional role in suppressing changes in cell characteristics. We, therefore, performed concurrent

video recordings of A549 p53 proficient lung carcinoma cells treated with scrambled siRNA or p53

siRNA (Control and p53 RNAi cells, respectively), and analyzed cells using single-cell tracking.

Distinct from other single-cell analyses; e.g., single-cell transcriptomics, which reveal the

characteristics of individual cells at a certain moment in time, data obtained by single-cell tracking

include information on spatiotemporal changes for individual cells, thus revealing the changes in cell-

population characteristics over time. Furthermore, because tracking of individual cells allows cellular

events to be detected regardless of the frequency of their occurrence, the response of a cell population

can be delineated at single-cell resolution with higher accuracy compared with analyses based on the

average responses of a cell population.

The first step in single-cell tracking analyses is the generation of a long-term live-cell imaging video

recording continuous cell movement. Images, typically $2,300 \times 2,300$ pixel area ($1,200 \times 1,200$ µm),

were acquired every 10 min (Figure 1). Phototoxicity, which can damage cells during long-term live-

cell imaging, was minimized by using differential interference contrast (DIC) imaging with near-

infrared light. The single-cell tracking videos were then started by selecting cells referred to as

progenitors in the video images at a tracking time of 0 min, and single-cell tracking of these progenitors

and their progeny was performed using our previous computerized single-cell tracking analysis

system[33]. This system created fully automated live-cell imaging videos (up to 16 videos concurrently)

and carried out image segmentation, automated single-cell tracking, database creation, data analysis,

and archiving. Because we excluded any data related to progenitors or progeny that moved out of the

field of view, the resulting single-cell tracking data contained complete information for progenies

derived from progenitors produced during the single-cell tracking period. The resulting single-cell

tracking data thus contained the position of each cell at every time point in an image, the type of events

that occurred in a cell (bipolar cell division, tripolar cell division, tetrapolar cell division [tripolar and

tetrapolar cell divisions were defined as multipolar cell division], cell death, and cell fusion (see Figure

1-Figure supplement 1 for still images of bipolar cell division, multipolar cell division, cell death, and

cell fusion; corresponding videos have been deposited in Dryad), and the relationships of a cell to its

parent and offspring cells. To organize single-cell tracking data, each progenitor and its progeny was

assigned a unique ID; i.e., a cell-lineage number. In this study, a group of cells (a progenitor and its

progeny cells) with the same ID was referred as to a cell lineage. A unique cell number was also

assigned to each cell constituting a cell lineage, and a database composed of multiple cell-lineage data

was referred as to a cell-lineage database. We then used the cell-lineage database to perform

quantitative bioinformatics analysis (Figure 1) at the single-cell and cell-lineage levels. Cell

population-level analysis could also be performed by collecting data for cells that belonged to different

cell lineages.

Some information in the cell-lineage database could be visually represented as a cell-lineage map

(Figure 1), containing multidimensional information, e.g. number of progeny produced from a

progenitor, type and time of events that occurred in a cell or in all cells comprising the lineage, and the

length of time between events. Quantitative bioinformatics analyses were performed to analyze specific

aspects of the information. Furthermore, given that the information stored in a cell-lineage database can

reveal a pattern of events that occurred in the cell, cell-lineage, and cell population, we used this

information to simulate the fates of cells by developing a cell-fate simulation algorithm.

Quantitation of cell numbers, areas, and densities

First, we prepared uniform cell cultures for siRNA treatment using plated cells occupying ~90% of the

culture surface, where > 99% of cells were attached to other cells. As shown in Figure 1-figure

supplement 2a, the number of cells/μm^2 increased linearly while the average cell surface area was

reduced, confirming that the number of cells could increase continuously in a linear manner during

video monitoring. To visualize this increase, density maps were also created at each imaging time point

(Figure 1-figure supplement 2b for still images and Figure 1-video 1) by assigning a value of 1 to

pixels within a 20-pixel diameter (10.43 µm) of the position of the cell (Figure 1-figure supplement 2c).

Cell positions were determined using data recorded in the cell-lineage database, and the pixel was

assigned the sum of the number of overlapped areas. When a cell was close to other cells, the number

of overlaps (i.e., the sum) increases, representing the cell density in an area. These sums were

converted into a heat map, which showed an increase in density during video monitoring (Figure 1-

figure supplement 2b and Figure 1-video 1).

Minimum number of cell lineages required to build a cell-lineage database

We then determined the minimum number of cell lineages (i.e., number of progenitors to be tracked)

needed per experiment to build a cell-lineage database. For this purpose, we first constructed 485 cell

lineages by performing single-cell tracking using 485 progenitor Control cells and then randomly

selected 50, 80, 120, 140, and 240 cell lineages from the 485 cell lineages to generate cell-lineage

databases containing the selected number of lineages. This selection was repeated three times for

triplicate analysis for each cell-lineage database. Variation in the rate of cell population expansion was

then evaluated by calculating the total number of cells at each time point using the cell-lineage

databases. The results for the 50, 80, 120, 140, and 240 cell lineages are shown in Figure 1-figure

supplement 3a-e. For example, the variation in the number of cells at the 4,000 min-determined cell-

lineage database containing 50 cell lineages (Figure 1-figure supplement 3a, black bar) was 51 cells,

and the variations became constant (7.5–6.0 cells for 150–240 cell lineages) when > 240 cell-lineages

were used (Figure 1-figure supplement 3f). This suggests that 240 cell lineages are sufficient for

reproducible single-cell tracking analysis, and we therefore used 335 cell lineages to create a cell-

lineage database.

Determination of length of time for single-cell tracking following siRNA and/or MNNG treatment

We determined the length of time required for single-cell tracking following siRNA transfection.

Because transfection can be monitored by observing the accumulation of particles (transfection

reagents) in cells (Figure 1-figure supplement 4a, DIC), we quantitated the number of particles in an

area of the cell culture by extracting particles with a grayscale value >150/255 from the video image,

and counted the number of particles with a size of 5–30 pixels (Figure 1-figure supplement 4a, Particles,

and Figure 1-video 2). The number of particles started to increase at about 1,200 min (20 h) after

transfection and peaked at 3,750 min (62.5 h) (Figure 1-figure supplement 4b). Western blotting

analysis performed 48 h after transfection confirmed silencing of p53 (Figure 1-figure supplement 4c;

Scrambled siRNA *vs.* p53 siRNA, ~70% reduction), the expression of low levels of p53 (Scrambled

siRNA, 0 µM MNNG) and the occurrence of p53 stabilization by exposure of cells to 7 µM MNNG

(Scrambled siRNA, 5.8-fold increase). Based on these parameters, we applied a 30-min pulse of

MNNG treatment after the start of particle accumulation (Figure 1-figure supplement 4b; 1,740 min (29

h) after transfection). Single-cell tracking was started at the end of MNNG treatment and continued for

4,000 min (66.7 h), by which time the number of particles was reduced to 70% of that at the peak.

Notably, the degree of p53-silencing may vary among cells, and it is possible that only certain groups

of cells may respond to silencing. This technique cannot reveal the level of silencing within the

individual cells, but the data for the population as a whole can reveal differences, providing a clue to

the function of the low levels of p53.

Analysis of rates of cell population expansion of Control and p53 RNAi cells, and cells treated

with MNNG using cell-counting and single-cell tracking approaches

The lowest levels of p53 could occur in unstressed cells [4, 6, 7, 16-18]. To gain insights into the functions of

such low levels of p53, we analyzed the expansion rate of a p53 RNAi cell population and their

response to MNNG exposure. The rate of cell population expansion, often referred as to the cell growth

curve, has generally been measured by counting cells at certain time intervals. We therefore initially

determined the rate of expansion using a classical, cell-counting approach (count cell numbers every

1,000 min) and compared the results with those obtained by single-cell tracking analysis (every 10 min)

to evaluate the accuracy of the analysis (Figure 2). For the cell-counting approach, cell population

expansion curves were determined manually by counting cells present in images of live-cell imaging

videos every 100 time points (1,000 min). To this end, images corresponding to every 100 time points

were divided into 25 (5 × 5) squares (~512 × 512 pixels per square, 266.24 × 266.24 µm per square),

and the number of cells within a square was counted (Figure 2-figure supplement 1). We used three

independently generated videos and randomly selected 15 squares. The results of counting of Control

and p53 RNAi cells are summarized in Figure 2a and b, respectively. The rates of expansion of the

Control and p53 RNAi cells and their responses to MNNG exposure were also compared (Figure 2c-f).

There was no significant difference in the expansion rates of Control and p53 RNAi cells (Figure 2c).

There was also no significant difference in the expansion rates of Control cells and Control cells

exposed to 1 µM MNNG (MNNG1) (Figure 2a, Control vs. MNNG1), while the expansion rate of p53

RNAi cells exposed to 1 µM MNNG (p53 RNAi-MNNG1) was reduced relative to Control cells

exposed to MNNG1 (MNNG1) and p53 RNAi cells (Figure 2b and d). When the dose of MNNG was

increased to 3 µM (MNNG3), the expansion rate of MNNG3 cells was reduced relative to Control cells

(Figure 2a) but was higher than that of p53 RNAi-MNNG3 cells (Figure 2e, MNNG3, and p53 RNAi-

MNNG3). At the highest MNNG dose used in this study (7 µM, MNNG7), at which significant

accumulation of p53 was observed) (Figure 1-figure supplement 4c), the rates of expansion of both the

MNNG7 and p53 RNAi-MNNG7 cell populations were significantly reduced compared with non-

exposed Control and p53 RNAi cells, respectively, but p53 RNAi cells were significantly less sensitive

to the MNNG dose than Control cells (Figure 2f). These results suggest that cells with different

sensitivities to MNNG compared with Control cells were generated by silencing the low levels of p53.

Furthermore, the relative sensitivities of Control and p53 RNAi cells to different doses of MNNG

differed, further suggesting that these p53 RNAi cells did not respond linearly to the different doses of

MNNG.

These results revealed how cell population size was influenced by p53 silencing and/or MNNG

treatment. However, this approach could not provide information on how each cell responded to the

treatment or how the response affected the size of the cell population. We then analyzed cell-lineage

databases obtained by single-cell tracking, given that the database contains information on events such

as cell death, fusion, and division, which influence the cell population expansion, at single-cell

resolution. Because the cell-lineage database generated by tracking contained information on individual

cells at each time point in the images (every 10 min), the total number of cells every 10 min was

plotted in Figure 2g-l (videos of the single-cell tracking processes, cell lineage maps, and cell lineage

data have been deposited in Dryad). Single-cell tracking analysis and cell counting yielded similar

results (Figure 2g and h); the relative differences in the rates of cell expansion between the Control and

p53 RNAi cells, and those treated with MNNG, were consistent between the counting and single-cell

tracking results (Figure 2j-l). Regarding the comparison between Control and p53 RNAi cells, although

counting every 1,000 min showed no significant differences, single-cell tracking, which can survey the

cell number every 10 min, revealed that the population expansion rate of p53 RNAi cells was

significantly lower than that of Control cells, confirming that the single-cell tracking analysis approach

produced equivalent results to the cell-counting approach (Figure 2c and i). However, in contrast to the

classical counting approach, which only represents the number of cells at a certain time point, cell-

lineage databases contain additional spatiotemporal information at single-cell resolution, which could

allow a more detailed analysis of how the cell population is expanded or reduced, what type of cellular

events were induced, and how they impacted the phenotype of a cell population with reduced p53

expression or DNA damage stress. We therefore conducted further analyses to gain insights into the

Control and p53 RNAi cell populations and those exposed to MNNG using computer-assisted single-

cell analysis.

Analysis of reproductive ability of cells using cell-lineage database

Cell populations are known to be composed of cells with different reproductive abilities [34], and the

overall rate of cell population expansion (Figure 2) is thus often influenced by the relative reproductive

abilities of the cells within the population. Cell population expansion curves *per se* do not provide any

insights regarding the reproductive abilities of individual cells to allow the impact of such variations in

the cell population to be assessed. We therefore further revealed the characteristics of Control, p53

RNAi, and MNNG-exposed cell populations by analyzing the reproductive abilities of the different

cells comprising each cell population.

To this end, we sorted each cell lineage into groups (Figure 3a). Given that the reproductive ability of

a cell could be represented by the number of cells (progenitor and its progeny) comprising a cell

lineage, we calculated the total number of cells comprising each cell lineage and sorted them according

to the number (the example cell lineages shown in Figure 3a are composed of 5 and 13 cells). We

grouped cell lineages that produced 1–2, 3–4, 5–6, 7–8, 9–10, 11–12, 13–14, and 15–16 cells as Groups

A, B, C, D, E, F, G, and H, respectively. The lineages in Figure 3a left and right were therefore sorted

as Groups C and G, respectively.

The total number of cells in each group was then calculated (Figures 3b and c). For example, if 20

cell lineages were composed of 11 cells (Group G), the total number shown in the Group G column

would be 220 cells. The columns in Figure 3b and c thus represent the composition of cells that could

have different reproductive abilities, categorized as A–H. In the case of Control cells, most cell

lineages belonged to Group D (7–8 cells), which accounted for 45% of the total number of Control

cells (Figure 3b). Most p53 RNAi cell lineages also belonged to Group D, but the number of cells with

higher reproductive capacity; i.e., Group F (11–12 cells), was increased relative to Control cells

(Control: 33 cells *vs* p53 RNAi: 143 cells), which might represent a change induced by silencing the

low levels of p53 (Figure 3c). The number of cell lineages in Group D was increased in the MNNG1

compared with the Control population (Control: 798 cells *vs* MNNG1: 1057 cells), while the number of

corresponding lineages in p53 RNAi-MNNG1 cells was reduced to 596 cells compared with p53 RNAi

823 cells (Figure 3b). This difference may reflect the difference in cell population-expansion rates

between MNNG1 and p53 RNAi-MNNG1 cells (Figure 2j). Because MNNG is a cytotoxic agent, it

could reduce the number of cells with a higher reproductive ability and increase cells with a lower

reproductive ability. Indeed, the numbers of Group B cell lineages were increased in MNNG3 and p53

RNAi-MNNG3 cells, but MNNG3 cells could maintain a higher ability to expand relative to p53

RNAi-MMNG3 cells, because of the larger content of Group D cell lineages (645 cells *vs* 435 cells)

(Figure 3b and c). Most lineages in MNNG7 and p53 RNAi-MNNG7 cells were Group A–B. However,

p53 RNAi-MNNG7 cells still contained Group C–D lineages, which could allow the cell population to

recover from the impact of 7 μM MNNG treatment. These results suggest that the Control cells did not

respond to different doses of MNNG in a simple dose-response manner and that silencing the low

levels of p53 also differentially affected the response patterns. Multipolar cell division, cell death, and

cell fusion, which could have a negative impact on the cell proliferation rate, could likely be involved

in determining the response patterns, and we therefore further analyzed the impact of p53 silencing and

MNNG exposure on these events.

Impact of p53-silencing and MNNG treatment on cell death, multipolar cell division, and cell

fusion

It has been reported that multipolar cell division occurs at a frequency of 1%–10% of all cell divisions

in established cultured cell lines[35, 36]. Cell death could be a less-frequent event in a growing cell

population, except in the presence of a cytotoxic agent. Although cell fusion is often induced during

differentiation; e.g., myogenesis, it is rarely induced in a growing cell population. These events would

thus be expected to occur less frequently than bipolar cell division. As noted above (Figure 1-figure

supplement 3f), more than 240 cell lineages were sufficient for reproducible single-cell tracking-based

analysis of a cell population expansion curve, which mainly represents the number of cell divisions, i.e.,

bipolar cell divisions. Although we started with 335 progenitors to perform single-cell tracking analysis,

it was not clear if tracking this number of progenitors and their progeny would produce enough

multipolar cell division, cell death, and cell fusion events for statistical analysis. To this end, we first

generated reference data by visually examining the videos and counting the numbers of multipolar cell

division and cell death events in three videos to gain an overview of the frequencies of these events in

the entire fields of view, and compared the results with those obtained using single-cell tracking

analysis (Figure 4). Notably, cell fusion was not included in the counting analysis because it was

difficult to detect by visual examination.

For counting, each time point of the video image was divided into four squares (Figure 4-figure

supplement 1a), and cell death and multipolar cell division events were marked. p53 silencing

significantly increased multipolar cell division and cell death events relative to Control cells (Figure 4a

and b, Figure 4-figure supplement 1b and c). This revealed that, even though Control and p53 RNAi

cells showed similar cell population expansion curves (Figure 2i), the cell population sizes (total cell

numbers) of the p53 RNAi and Control cell populations resulted from different processes. Increased

multipolar cell division and cell death events following p53 silencing were reduced by MNNG

treatment (Figure 4a, b, Figure 4-figure supplement 1b and c), with the degree of suppression of

multipolar cell division being more significant than that of cell death. Multipolar cell division and cell

death occurred less frequently in Control cells, even after exposure to MNNG, except for cells exposed

to 7 μM MNNG (Figure 4b). These results suggest that silencing of the low levels of p53 led to the

induction of multipolar cell division and cell death.

We then determined the numbers of multipolar cell division and cell death events using single-cell

tracking-based cell-lineage analysis. Given that the cell-lineage database contains information on not

only the events occurring in individual cells, but also their timing, it was possible to perform in-depth

analysis of other events that occurred before and after the events in question, as well as sibling

relationships. The areas selected for the analysis are indicated by black boxes in Figure 4-figure

supplement 1b and c. The results of analyses of multipolar cell division and cell death by single-cell

tracking analysis were consistent with the counting approach (Figure 4a-b (counting) vs. c-d (single-

cell tracking)), suggesting that the number of events determined by tracking the 335 progenitors and

their progeny was sufficient to represent the occurrence of these events in an entire field of view in the

videos. In one case; i.e., treatment of cells with 7 µM MNNG, the number of cell death events in

Control cells determined by counting was higher than that determined by single-cell tracking analysis.

Given that such events tend to occur in a non-uniform manner within the field of view (Figure 4-figure

supplement 1b and c), there may be some variation between the counting and single-cell tracking

approaches. Although such variations should be taken into account when interpreting the single-cell

tracking results, this approach still has an advantage over the counting approach, by detecting the

cellular events encountered by a cell and their trajectories. Cell fusion, which required the fused cells to

be followed for at least 30 time points (~5 h), can thus be detected by single-cell tracking analysis.

Similar to multipolar cell division and cell death, p53 silencing increased the frequency of cell fusion,

and this was counteracted by MNNG (Figure 4e). Taken together, these results suggest that silencing of

the low levels of p53 induced multipolar cell division, cell death, and cell fusion, and these were

reduced by exposure of the cells to MNNG.

Impact of cell death on cell population expansion of p53 RNAi cells using *in silico* generation of a

cell-lineage database

The rate of cell population expansion was similar in Control and p53 RNAi cells (Figures 2c and 2i).

However, detailed analysis revealed that silencing the low levels of p53 increased multipolar cell

division, cell death, and cell fusion (Figure 4). The probabilities of detecting bipolar cell division,

multipolar cell division, cell death, and cell fusion at a certain time point in an image were calculated as

0.54, 0.001, 0.004, and 0.004/100 lineages, respectively, for Control, and 0.54, 0.014, 0.037, and

0.023/100 lineages, respectively, for p53 RNAi cells. This implies that the more frequent occurrence of

these events in p53 RNAi cells relative to Control cells did not reflect the overall rate of cell population

expansion, even though the events, especially cell death, could counteract the expansion of the cell

population. In this regard, given that the p53 RNAi cell population contained an increased number of

cells with higher reproductive ability (e.g. group F cell lineages; Figure 3c) than Control cells, silencing

p53 may have increased cells with higher reproductive ability than Control cells, but this increase may

have been counteracted by the induction of cell death (Figure 4b and c), resulting in similar population

expansion curves for p53 RNAi cells and Control cells. In this case, if no cell death occurred in p53

RNAi cells, they would show a faster rate of cell population expansion than Control cells.

To test the effects of cell death on the rate of population expansion, we generated a cell lineage

database *in silico* (Figure 5a) by assuming that cells that undergo cell death continue to proliferate.

To this end, we searched the data for siblings of cells that underwent cell death in a cell-lineage

database. Given that their proliferation pattern was likely to be similar to their siblings, we replaced the

data for cells that underwent cell death with those for their siblings (Figure 5a). The cell population

expansion rate of the *in silico*-generated p53 RNAi cells without cell death (p53 RNAi-Silico(-)cell

death) (Figure 5b) showed that the cell population size was 7.4% larger than that of the p53 RNAi cells

at a tracking time of 4,000 min (Figure 5b; p53 RNAi *vs.* p53 RNAi-Silico(-)cell death). The number of

cell deaths that occurred in p53 RNAi cells are indicated in Figure 5b, and the p53 RNAi cell

population expansion rate was reduced around 2,000 min following cell death at 1,000–2,000 min of

tracking. This reduction was less evident in p53 RNAi-Silico(-)cell death cells, suggesting that cell

death in p53 RNAi cells did indeed counteract the expansion of this cell population. The cell

population size of the p53 RNAi-Silico(-)cell death cells was also larger than those of the Control and

in silico-generated Control cells (Control-Silico(-)cell death), where Control cells that underwent cell

death were similarly replaced with their surviving siblings. These results suggest that p53 silencing

promoted the reproductive ability of p53 RNAi cells, but this was counteracted by the induction of cell

death, resulting in the formation of p53 RNAi cell populations that were similar to or smaller than

Control cell populations (Figure 2i). This analysis also demonstrated that single-cell tracking provided

more nuanced and detailed information about the effects of silencing p53 and the processes involved in

determining the cell population size, by taking account of the occurrences of multipolar cell division,

cell death, and cell fusion, which are often difficult to put into spatiotemporal context in terms of the

individual fate of cells when using methods that only reveal the status of cells at a certain moment in

time.

Single-cell tracking revealed the most common event preceding multipolar cell division

While cell death counteracted the expansion of the cell population, multipolar cell division and cell

fusion could have greater impacts than cell death in terms of creating diversity in the cell population by

causing the formation of aneuploid cells[37-42], if the progenies of these processes survive. We further

examined how silencing the low levels of p53 altered the characteristics of cells by focusing on

multipolar cell division and cell fusion and the relationships between these events. To this end, we used

cell-lineage maps (Figure 1 and cell lineage maps deposited in Dryad) to identify cells that underwent

multipolar cell division, and traced back along the map to find an event that occurred prior to the

division (Figure 6a). We identified two patterns: multipolar cell division occurring after cell fusion

(Pattern 1a: Cell fusion ➤ Multipolar cell division), and multipolar cell division without cell fusion

(Pattern 1b: Multipolar cell division). Control and p53 RNAi cells demonstrated that 79.4% and 93.2%

multipolar cell divisions, respectively, occurred following cell fusion (Pattern 1a). It is likely that cells

with increased ploidy could have a higher chance of undergoing multipolar cell division and silencing

of p53 increased the incidence of Pattern 1a (Figure 6a, Pattern 1a, Control *vs.* p53 RNAi). We then

analyzed cells that fused by tracing further back along the cell-lineage maps (Figure 6b). Interestingly,

cell fusion was more frequent between sibling cells (Pattern 2a). In this regard, a previous study

reported that some persistent links remained after bipolar cell division leading to cell fusion [43].

However, given that cell fusion also occurred between non-siblings (Pattern 2b), the cell fusion

observed in this study was mediated by other processes. In summary, p53 silencing mainly promoted

the sequence of events: Bipolar cell division ➤ Cell fusion between siblings ➤ Multipolar cell

division (Figure 6b, Pattern 2a).

The progeny of multipolar cell division have been reported to be fragile, although some are capable

of entering a proliferation cycle [39, 40]. We, therefore, analyzed the survival of the progeny using cell-

lineage maps. In terms of percentage survival, 3.2%–3.6% of the total number of progeny produced by

multipolar cell division survived in both Control and p53 RNAi cells (Figure 6c). On the other hand,

given that silencing the low levels of p53 *per se* increased the number of multipolar cell divisions, the

number of surviving progenies produced by division in the p53 RNAi cell population was increased

10.1-fold (0.61 progeny of multipolar cell division/100 cell lineages of p53 RNAi cells *vs.* 0.06

progeny of multipolar cell division/100 cell lineages of Control cells). These results suggest that p53

silencing mainly promoted the number of events with the sequence: Bipolar cell division ➤ Cell

fusion between siblings ➤ Multipolar cell division ➤ Survival of aneuploid progeny, leading to the

creation of genetic diversity in the p53 RNAi cell population.

We plotted the number of multipolar cell divisions against the number of bipolar cell divisions in

Control and p53 RNAi cells and those treated with MNNG (Figure 6d). Although fewer multipolar cell

divisions occurred in Control cells relative to p53 RNAi cells, MNNG exposure resulted in a dose-

dependent linear reduction of bipolar vs. multipolar cell division, suggesting that the probability of

inducing multipolar cell division was proportional to the number of bipolar cell divisions. These results

also suggest that MNNG indirectly reduced the number of events in the sequence: Bipolar cell division

➤ Cell fusion between siblings ➤ Multipolar cell division ➤ Survival of aneuploid progeny.

Although MNNG is a mutagen that induces G-T to A-T transversion mutations through the formation

of O^6-methylguanine[31], it could act to suppress the formation of aneuploid cells in some contexts, with

possible relevance to the development of cancer[44].

Cell-fate simulation algorithm

The above data suggest that the fate of cell progeny following silencing of the low levels of p53 could

alter the characteristics of the cells and their sensitivity to MNNG. However, characterization of the

cells using a direct single-cell tracking approach was limited by various factors. For example, there is a

practical limit to the number of cells that can be analyzed by tracking and the duration that a cell

culture can be maintained without cell passage, which makes it difficult to follow the fate of a cell

beyond a certain duration. We, therefore, developed a cell-fate simulation algorithm to overcome this

limitation and further analyzed the effects of p53 silencing on the fate of the cell population. For the

simulation, we utilized the spatiotemporal information for each cell in the cell-lineage database records,

which revealed the cell's relationships with its parents, siblings, and offspring and events that occurred

in that cell. In addition to overcoming the limitation, cell-fate simulations can provide powerful and

flexible tools to model conditions that are not readily accessible by direct imaging, such as simulating

the fate of a cell in a mixed culture population, evaluating the response of cells to drug treatment

virtually, and testing existing models *in silico*.

The underlying concept of the cell-fate simulation algorithm is shown in Figure 7 (Figure 7-figure

supplement 1: overall scheme, and Figure 7-figure supplement 2: algorithms). We first decomposed a

cell lineage into units sandwiched between two events, such as bipolar cell division, multipolar cell

division, cell fusion, and cell death (Figure 7a). We referred to events that initiated and ended the unit

as Start and End events, respectively, and each unit was thus defined by the nature of its Start and End

events, and the length of time between the two events. An algorithm combining such units can thus be

used to generate a virtual cell lineage with a similar pattern to an empirically determined lineage.

Because each cell lineage in a cell population shows a variety of patterns, and the combination of these

variations defines the characteristics of the cell population, the simulation algorithm thus needs to

reflect this variation. To this end, we first produced a series of histograms of the distribution of the end

events for each Start event (Figure 7b). For example, if the Start event was bipolar cell division, we

created a histogram of the frequency of all possible End events; i.e., bipolar cell division, multipolar

cell division, cell death, and cell fusion. The histogram indicated the unique tendency of a cell

population. For example, the Figure 7b histogram indicates that the most frequent End event in the cell

population was bipolar division followed by multipolar division. The algorithm for the simulation thus

utilized the frequency to generate virtual cell lineages (see Figure 7-figure supplement 2b: 2 for the list

of Start events). We referred to the histogram data as Operation data-Events, which is composed of a

series of histograms corresponding to each Start event. The algorithm then reflected this distribution of

the histograms to choose the End event. The selection of the End event was performed by converting

the histogram data into an array by taking account of the frequency of the occurrence of each event to

generate a random number to select an event type from the array. In this manner, an event type that

occurred more frequently could have a higher chance of being selected, reflecting the unique

characteristics of a cell population. Once the Start and End events were chosen, the algorithm referred

to another set of histograms (Operation data-Time) that showed the distribution of the length of time

for a given unit (Figure 7c). For example, for a unit with bipolar cell division as the Start event and cell

fusion as the End event, the algorithm referred to the histogram data of time length distributions of the

corresponding event combination (see list of histograms for Figure 7-figure supplement 2b: 1). The

length of time was selected in a similar manner to the End event selection; i.e., converting histogram

data into an array and selecting an assigned length of time by generating a random number. These

processes were repeated until the cell lineage reached the desired time point. In addition to the process

of assigning event type and time length, the cell-doubling time of the parent cell was also taken into

account. If both the Start and End events of a parent cell were bipolar cell division, our experimental

results indicated that the length of time of its offspring was limited to ± 10% of that of the parent cell.

This restraint was determined by repeated stimulation with single-cell tracking data generated using

A549 cells. If a parent cell had a shorter length of time than others, its progeny could also have a

shorter doubling time, creating an actively growing cell lineage. In this manner, the algorithm could

generate lineages with different reproductive abilities, reflecting the characteristics of a cell population

obtained by empirical live-cell imaging. This restraint for the duration between divisions was only

applied to cells produced by bipolar cell division that underwent another bipolar division, given that

cells produced by other types of events; e.g., multipolar cell division and cell fusion, have less

contribution to forming a cell population than those that undergo bipolar cell division.

We summarized the overall scheme of analysis with single-cell tracking and cell-fate simulation

(Figure 7-figure supplement 1a). After the generation of live cell imaging videos, images were

segmented and the segmented data was used for automatic single-cell tracking. Then, the database was

used for cell-lineage map creation, data analysis, and generation of Operation data. Operation data was

then used for the cell-fate simulation (Figure 7-figure supplement 1b). After loading the cell-lineage

data, Operation data-Time and -Events were created. The length of time for progenitors and a primary

event type were assigned to each progenitor, followed by the generation of cell-lineage data by

assigning an event type and length of time to a cell. Example cell-lineage maps generated by empirical

single-cell tracking analysis and the algorithm are shown in Figure 7-figure supplement 3a and b,

respectively (cell-lineage maps generated by the algorithm have been deposited in Dryad). Furthermore,

because the simulation *per se* can be carried out using Operation data, various simulation options can

be created by modifying the Operation data content.

Cell-fate simulation options with Operation data

We created five different modes of simulations: Standard, Dose simulation, Mix culture, Switch, and

Mix culture-Switch. The Standard mode simulates the fate of one cell type using one Operation data

(Figure 8a). We applied this mode to simulate the expansion of a cell population for an extended period

of time, which cannot be achieved by *in vitro* cell culture (Figures 9 and 11). The Dose simulation

mode can be carried out by taking advantage of the fact that Operation data-Events and Operation data-

Time are histogram data, in which values can be modified by applying a certain rule, e.g. a linear

relationship, to create new Operation data. For example, if histogram arrays of Operation data-Events

A and B (in Figure 8b) obtained from the single-cell analysis of cells treated with 2 and 5 µM of a drug,

respectively, recorded 10 and 1 bipolar cell divisions following a Start event, the number of bipolar cell

divisions of a cell population exposed to 3 µM was calculated to be 7. In the example (Figure 8b), we

also calculated multipolar cell division, cell death, and cell fusion data stored in the histogram array.

Similar calculations were made for other histogram data stored in Operation data-Events and Operation

data-Time (Figure 7-figure supplement 2b), generating new Operation data-Events and Operation data-

Time. We applied this mode to simulate cellular responses to different doses of MNNG (Figure 10).

For example, the fate of cells exposed to 5 µM MNNG can be simulated by creating Operation data for

5 µM MNNG from Operation data for 3 µM and 7 µM MNNG. The Mixed culture mode (Figure 8c)

simulates a fate of a cell population composed of two or more types of cell populations using Operation

data corresponding to each type of cell population. We applied this mode to simulate the expansion of

cells under conditions in which 96% of cells were initially Control cells and 4% were p53 RNAi cells

(Figure 12). The Switch mode *per se* was not used in the current work, but this mode (Figure 8d) was

used to perform virtual drug treatments. For example, simulation using the Operation data for Control

cells was then switched to the second Operation data for cells treated with, for example, 3 µM MNNG,

allowing simulation of the effect of 3 µM MNNG on Control cells. The Mixed culture with Switch

mode (Figure 8e) allowed simulations to be started with a virtual cell population composed of two or

more different types of cell populations with virtual drug treatment. Thus, if two types of cell

population fates are to be simulated, two Operation data are used initially and then switched to

Operation data for drug-treated cells of each type (Figure 13). If the cell types have different treatment

sensitivities, the effects of this difference can be simulated by the Mixed culture with Switch mode. We

applied this mode to simulate the impact of repeated MNNG treatments on the expansion of a cell

population initially comprising 96% Control and 4% p53 RNAi cells at the start of the simulation

(Figure 13h). In summary, cell-fate simulation using Operation data allows the creation of various

simulation options and provides flexibility for designing virtual experiments that would otherwise be

difficult to perform empirically.

Verification of number of progenitors used for simulation

We characterized the algorithm by performing a series of test simulations. To distinguish data produced

by simulation from data produced by single-cell tracking, we referred to the cells as; e.g., Control-Sim

and MNNG7-Sim cells for the results produced by simulation (Figure 8-figure supplement 1). If the

corresponding Operation data needed to be indicated, the cell populations were referred to as; e.g.,

Operation data-Control and Operation data-MNNG7. We first performed simulations with 300, 500,

1,000, 1,500, and 2,000 progenitors with a simulation time of 15,000 min, repeated five times to detect

variations in the simulation. Cell population expansion curves of virtually created Control cells

(Control-Sim cells) are shown in Figure 8-figure supplement 1a. Under these conditions, the variation

was < 2% of the average number of cells when the simulation was carried out with > 1,000 progenitors.

The simulation performed using the Operation data-MNNG7 (Figure 8-figure supplement 1b) produced

a larger variation, but this converged to about 6% when the simulation was carried out with $> 1,000$

progenitors. We therefore performed the simulation with 1,000–2,000 progenitors and repeated the

simulation five to ten times (total of 5,000–10,000 progenitors).

Comparison between cell population expansion curves determined by simulation and single-cell

tracking

We then evaluated the accuracy of the simulation by referencing the cell population expansion curves

determined by single-cell tracking analysis (Figure 9). In the simulations, we used Standard mode with

Operation data generated from the cell-lineage database and performed simulation with 5,000

progenitors for 15,000 min. The number of progenitors was normalized to 100 to compare the results

obtained by simulation with those obtained by single-cell tracking analysis. During the simulation, a

Control-Sim cell population size increased 193.5-fold (5,000 to 967,281 cells), and about 2×10^6

virtual cells were created in the simulation period of 15,000 min. The Control-Sim, MNNG1-Sim,

MNNG3-Sim, and MNNG7-Sim cell population expansion curves (Figure 9a, 0–4,000 min) were

consistent with the curves determined using single-cell tracking analysis (Figure 2g), although the

Control-Sim cell curve intersected with the MNNG1-Sim cell curve at a simulation time of about

10,000 min (Figure 9b, 0–15,000 min). To evaluate the accuracy of the simulation, we calculated the

average percent difference between the cell numbers determined by the simulation and by single-cell

tracking analysis (Figure 9c-f). When the number of cells determined by single-cell tracking was

calculated as 100%, the average difference was 1%–6%. We then performed a similar analysis with

p53 RNAi cells and found that cell population expansion curves generated by the simulation showed

consistent patterns (Figure 9g, 0–4,000 min, and Figure 9h, 0–15,000 min) with those determined by

single-cell tracking analysis (Figure 2h). The percent differences were within 0–8% (Figure 9i-l). We

thus concluded that the simulation could be performed within a variation of maximal 8% compared

with the results of single-cell tracking analysis.

Doubling time of cells generated by the simulation

We also analyzed the doubling time of individual cells generated by single-cell tracking analysis and

compared them with ones generated by simulation. The average cell doubling time was prolonged

following the increased dose of MNNG (31.23 to 41.85 h, Figure 9-figure supplement 1a-d). The

doubling time of individual cells generated by the simulation (Figure 9-figure supplement 1e-h) was

similarly prolonged following the increased dose of MNNG (29.36 to 39.43 h), suggesting that the

algorithm could simulate the effect of MNNG on cell doubling. Similarly, the effect of MNNG

treatment on the cell doubling time of p53 RNAi cells (Figure 9-figure supplement 1i-l) could also be

simulated, showing the consistent results with the analysis of single-cell tracking analysis (Figure 9-

figure supplement 1m-p compared with i-l). On the other hand, the simulation tended to yield an

average cell doubling time about 2 h shorter than that determined by single-cell tracking analysis.

Given that the algorithm assigned a cell-doubling time to each cell by generating a random number

with Operation data-Time (Figure 7), a long cell-doubling time; e.g., 3,000 min, which occurred less

frequently, may have a lower chance of being assigned, resulting in the generation of a simulated cell

population with a cell-doubling time about 2 h shorter than that of cells analyzed by single-cell tracking.

Alternatively, the cell-doubling time distribution generated by the simulation would represent the

actual cell-doubling time of a cell population, given that the quantity of data generated by the

simulation was ~200 times larger than that by single-cell tracking. This shorter average cell-doubling

time could translate into an increased rate of cell population expansion, and the maximal 8% variation

in cell population expansion curves (Figure 9) could reflect the cell-doubling time of the simulated cell

population.

Simulation of occurrence of multipolar cell division, cell death, and cell fusion

We examined how the algorithm simulated the occurrence of multipolar cell division, cell death, and

cell fusion. Because these events were less frequent than bipolar cell division, simulations were

performed 10 times with 1,000 progenitors (total 10,000 progenitors). We compared the numbers of

these events generated within 4,000 min by the simulation with those obtained by single-cell tracking

analysis. These data were then shown as the percent of a total number of cells. The simulated results for

multipolar cell division (Figure 9-figure supplement 2a), cell death (Figure 9-figure supplement 2b),

and cell fusion (Figure 9-figure supplement 2c) showed similar tendencies to the single-cell tracking

results, and the differences between the tracking analysis and simulate results were within 1%. We,

therefore, concluded that the cell-fate simulation algorithm could perform virtual cell experiments with

similar accuracy to single-cell tracking analysis.

Virtual dose-response analysis and damage response

To begin the study with the fate-simulation, we reanalyzed the responses of cells transfected with

Scrambled siRNA or p53 siRNA to MNNG. To this end, Operation data were created for every 0.5 µM

of MNNG from 0–7 µM using Dose simulation mode to determine a more detailed dose-response curve.

The virtual dose-response curves determined by the simulation are shown in Figure 10a and b, and the

numbers of cells at simulation times of 4,000 and 15,000 min (indicated by blue and red arrow,

respectively, in Figure 10a and b) are shown as natural logarithm in Figure 10c and d, respectively. The

dose-response curves of cells containing low levels of p53 (Control cells) and p53 RNAi cells

determined at a simulation time of 4,000 min showed similar patterns to those determined by single-

cell tracking (Figure 10c), suggesting that the fate of cells exposed to 0.5 µM increments of MNNG can

be simulated by the Dose simulation mode. We then analyzed the effects of MNNG exposure on

Control and p53 RNAi cells at 15,000 min, to determine the long-term impact of MNNG exposure.

MNNG 7 µM reduced the cell population size of Control cells at 15,000 min from 193,928 to 5,810

cells (Figure 10d, difference: 188,118 cells, 97%), while the number of p53 RNAi cells was reduced

from 113,075 to 12,055 cells (difference: 101,020 cells, 89%), suggesting that Control cells showed a

stronger inhibitory response to MNNG than p53 RNAi cells in terms of cell population expansion,

possibly due to p53-induced responses (Figure 10d). These results suggest that the p53 RNAi cells

were less sensitive to MNNG due to the removal of the low levels of p53, resulting in a low or absent

p53-induced response triggered by MNNG.

MNNG induces its cytotoxic effects through the formation of DNA breaks and activation of the

response caused by the accumulation of p53 [26-32]. It is therefore conceivable that responses induced in

Control cells may be related to both DNA-break formation and the response induced by the DNA

damage-initiated accumulation of p53, while responses in p53 RNAi cells are only related to DNA-

break formation. To estimate the relative contributions of these two effects to the inhibition of cell-

population expansion, we plotted the population expansion curves of Control-Sim, MNNG7-Sim, p53

RNAi-Sim, and p53 RNAi-MNNG7-Sim cells and determined the population-doubling time (Figure

10e). We hypothesized that the difference in times between Control-Sim and MNNG7-Sim (7,150 min)

was due to both effects, while the difference between p53 RNAi-Sim and p53 RNAi-MNNG7-Sim

(4,510 min) represented the effect caused by the break formation. Based on these data, we estimated

that about 63% and 37% of inhibition of cell population expansion caused by 7μM MNNG exposure

were due to effects related to DNA-break formation and the accumulation of p53, respectively.

Although these percentages could be affected by other factors, dissecting the cellular response based on

the possible mechanism of cytotoxicity could provide a deeper understanding of how cells respond to

cytotoxic drugs.

Survival of the progeny of multipolar cell division

Next, we asked how frequently the progeny of multipolar cell division can survive and grow by

performing a simulation in Standard mode with 2,500 progenitors for a simulation time of 20,000 min

(13.8 days). Multipolar cell division-derived progeny, which lack reproductive ability (Figure 11a, left),

and progeny that underwent multipolar cell division followed by bipolar cell division, thereby

maintaining reproductive ability (Figure 11a, right), are shown in burgundy and green, respectively.

When the simulation was performed using the Operation data-Control (Figure 11b and c), no

reproductive progeny produced by multipolar cell division were found. To visually show such cells, we

created a computer animation (Figure 11-video 1 and Figure 11d for still image at 20,000 min), which

visualizes the population expansion of the virtual cells, showing only burgundy cells that eventually

underwent cell death or remained as non-growing cells. However, when the simulation was performed

using Operation data-p53 RNAi (Figure 11e and f), some progeny of multipolar cell division

underwent bipolar cell division, accounting for 0.72% of the cell population at a simulation time of

20,000 min. We showed the population expansion of the progeny of multipolar cell division visually by

creating a computer animation (Figure 11-video 2 and Figure 11g for still image at 20,000 min), which

showed the expansion of green cell populations (cells survived and expanded after multipolar cell

division) in accordance with the suggestion [39, 40] that an aneuploid cell population, which could have

distinct characteristics, could be generated by the survival of the progeny of multipolar cell division

promoted by p53 silencing.

Limiting expansion of p53 RNAi cells in the presence of Control cells

We then investigated how the presence of Control cell influenced the population expansion of p53

RNAi cells, given that cells that lack p53 likely arise within a Control cell population during

tumorigenesis. We performed a simulation based on the assumption that Control and p53 RNAi cells

comprised 96% and 4% of the cell population, respectively, using Mixed culture mode with Operation

data-Control and p53 RNAi. A total of 1.2×10^7 virtual cells were generated after 20,000 min of

simulation, and the percentage of p53 RNAi cell population was reduced from 4% to 2.4% (Figure 12a)

because of the frequent occurrence of multipolar cell division, cell death, and cell fusion (Figure 12b,

and Figure 12-video 1 for the animation; p53 RNAi cells shown as blue cells). These data suggested

that, although p53 RNAi cells could generate characteristic diversity in their cell population through

increased frequencies of multipolar cell division, cell death, and cell fusion compared with Control

cells, this conferred a disadvantage in terms of p53 RNAi cell population expansion when both cell

types co-existed in a population. We suggest that the occurrence of cell death in p53 RNAi cells may

be the main factor limiting the expansion of these cells (Figure 5b). Indeed, a simulation using

Operation data generated from p53 RNAi-Silico(-)cell death cells showed that the cell population

expanded at a similar rate to Control-Sim cells (Figure 12c), again suggesting that cell death acted as a

major factor limiting the population size of p53 RNAi cells.

Induction of damage response in Control cells and its impact on the expansion of p53 RNAi cell

population

The relative population-expansion rates of Control and p53 RNAi cells were altered by exposure to

MNNG (Figure 10: Dose simulation mode). We, therefore, simulated the responses of a cell population

comprising Control (96%) and p53 RNAi cells (4%) to virtual MNNG treatment using the Mixed

culture-Switch mode. We used Operation data-Control and p53 RNAi up to a simulation time of 9,900

min, and the Operation data was then switched to the one for relevant doses of MNNG and the

simulation was continued for another 10,000 min (total 20,000 min simulation) (Figure 13a). We

simulated the responses of cells to 1.5, 3, 5, 6, and 7 µM MNNG. p53 RNAi cells were still unable to

expand their population relative to Control cells following exposure to 1.5 and 3 µM MNNG (Figure

13b and c). However, p53 RNAi cells started to gain an advantage in terms of population expansion

over Control cells after exposure to 5 µM MNNG (Figure 13d), and started to expand their population

at doses of 6 and 7 µM MNNG (Figure 13e, f, and Figure 13-video 1). These results suggested that the

damage responses induced in Control cells reduced its expansion speed while the expansion of the p53

RNAi cells continued (Figure 13g). Indeed, when the exposure of cells to 7 µM MNNG was repeated

four times, 93% of the cell population was replaced with p53 RNAi cells (Figure 13h). These results

suggest that, although p53 could suppress the formation of aneuploid cells, which could have a role in

tumorigenesis, it could also allow the expansion of cells lacking p53 expression under damage-

response conditions. p53 may thus play a dual role in the suppression and promotion of malignant cell

formation during tumorigenesis.

DISCUSSION

Cell populations are known to be composed of cells with diverse phenotypic characteristics. Such

diversity could be generated by the formation of a cell with distinct characteristics and the subsequent

expansion of its progeny in the cell population. In this study, we revealed that silencing of the low

levels of p53 generated cells with distinct characteristics from Control cells, and also affected the

responses of cells to MNNG. These results suggest that a novel approach using single-cell tracking and

cell-fate simulation can provide unique insights into the function of p53, thus deepening our

understanding of its role in tumorigenesis.

Low levels of p53 and possible functions

p53 is required for cells to respond to stress, through regulation of its cellular content [4, 6, 7, 10-18]. The

equilibrium between p53 degradation (through ubiquitination mediated by Mdm2) and its synthesis can

be changed, depending on the degree of stress [4, 6, 7, 16-18]. Stronger stress tends to shift the equilibrium

towards increased p53 levels, leading to various responses; e.g., metabolic regulation, DNA damage

responses, autophagy, cell cycle regulation, and cell death [6, 10-15]. On the other hand, the functions of

the low levels of p53 present in unstressed cells remain unclear, but they may have a housekeeping

function. Indeed, several lines of evidence, including the promotion of induced pluripotent stem cells

by silencing p53 [19], and the spontaneous formation of tetraploid cells, as proposed cancer precursor

cells, in p53 knockout mice [10, 45], suggest that low levels of p53 are required for housekeeping and

homeostasis functions in cells. In addition, Vousden and Lane pointed out that the low levels of p53

may play a role in responding to "daily levels of stress" [20], and Valente et al. recently demonstrated that

p53 plays a role in responding to stress induced by physiological levels of oxygen tension [46].

Furthermore, the current study showed that low levels of p53 inhibited the induction of cell death and

cell fusion, which could lead to the induction of multipolar cell division, suggesting that the low levels

of p53 are involved in processes that ensure the accuracy of cell division.

Silencing of low levels of p53 and induction of cell death, cell fusion, and multipolar cell division

The different characteristics of Control and p53 RNAi cells are summarized in Figure 14a. Most

Control cells underwent bipolar cell division, resulting in expansion of the cell population. Although

cell death, cell fusion, and multipolar cell division occurred in Control cells, they had relatively little

effect on the expansion of the cell population. In contrast, although p53 RNAi cells could expand their

cell population through bipolar cell division, p53 silencing led to more frequent induction of cell death

and cell fusion.

The cell death events in p53 RNAi cells appeared to be induced by a p53-independent mechanism.

Although the mechanism underlying the induction of cell death remains to be investigated, it may

involve a natural defense mechanism to remove cells lacking p53 from a cell population. Indeed, we

demonstrated that cell death caused a 7.4% reduction in the rate of p53 RNAi cell population expansion.

Cell death thus conferred a disadvantage to p53 RNAi cells to expand their population relative to

Control cells. Simulation of the fates of cells in a population containing 4% p53 RNAi and 96%

Control cells showed that the proportion of p53 RNAi cells was reduced to 1.2% after 670 h (28 days)

of culture (Figure 14b). When such a simulation was initiated with 5,000 cells, the cell number reached

$\sim 1 \times 10^9$, with a mass equivalent to ~1 cm of cancer tissue, after 21 cell divisions. Within a cell tissue

mass, p53 RNAi cells may occur as dormant cells, given that the percentage of p53 RNAi cells in the

tissue would be reduced following the increase in tissue size. With regard to dormancy, many

metastatic cells are known to become dormant [47, 48], suggesting that their metastatic nature *per se* could

confer a survival disadvantage on these cells relative to normal or non-metastatic cells, as indicated by

the difference in expansion rates between p53 RNAi cells in a Control cell population.

Cell fusion was also increased following silencing of p53 (Figure 14a). Although the role of p53 in

the suppression of cell fusion is unknown, previous reports have suggested a link between the loss of

p53 and the induction of cell fusion. For example, fibroblasts derived from p53$^{-/-}$ mice rapidly became

tetraploid [10, 45]. Increased cell fusion of p53 RNAi cells, mainly between sibling cells produced by

bipolar cell division, could account for the formation of tetraploid cells (Figure 14a). Such fused cells

may be formed by failure of cytokinesis, mitotic slippage, or the formation of a link between siblings [43,

49-51]. However, low levels of p53 may block the process of fusion itself, given that we also observed

cell fusion following abscission and between non-sibling cells. During development, such as myotube

formation [52, 53], a cell fusion mechanism is activated, implying that cells can acquire a fusion-prone

status. In addition to the physiological process of cell fusion, if cells; e.g. siblings produced by bipolar

cell division, became fusion prone, it could lead to the formation of polyploid cells, which have been

suggested to be precursors of cancer cells [49, 54-57]. The low levels of p53 may thus function to prevent

cells from becoming fusion-prone, thereby reducing the risk of cells becoming cancer cells.

Loss of the low levels of p53 increased the induction of multipolar cell division following cell

fusion (Figure 14a). The division of a cell into three and more cells generates progeny with different

numbers of chromosomes. Although such aneuploid cells are generally fragile [39, 40], loss of p53

function has been suggested to contribute to the survival of aneuploid cells [58-60]. Indeed, we found that

some such aneuploid cells underwent bipolar cell division and the progeny could propagate, creating an

aneuploid cell population (Figure 14a). Even though gene mutation has been proposed as the main

cause of cancer, aneuploidy is a near-universal feature of human cancers [61] and is also involved in the

process of tumorigenesis [37, 50]. Furthermore, impaired p53 function is known to cause frequent

alterations in cell ploidy [49, 54-57]. The proposed process could thus be involved in the process of

tumorigenesis. In summary, the low levels of p53 found in unstressed cells could suppress cell fusion,

which leads to multipolar cell division, and loss of this function could thus create conditions favoring

the generation of cancer cells.

Response of p53-silenced cells to MNNG

The above-mentioned context reveals the potential roles of low levels of p53 under unstressed

conditions. One of the best-established roles of p53 is related to cellular responses induced following

its accumulation under stressed conditions, including its capacity to prevent the survival of damaged

cells. This capacity is closely associated with cell growth arrest and cell death. On the other hand,

induction of the damage response in Control cells may oppose the suppression of tumor formation.

Exposure of a cell population composed of 4% p53 RNAi cells and 96% Control cells to 7 μM MNNG,

which could lead to the accumulation of p53, was associated with growth suppression of the Control

cell population due to induction of the damage response, conferring an opportunity for expansion of the

p53 RNAi cell population, which lacks the p53-mediated damage response function. Indeed, our

simulation results suggested that 4% of p53 RNAi cells present at the beginning expanded to 93% of

the cell population (containing 0.6% aneuploid cells) after 40,000 min (27.78 days) if the population

was exposed to 7 μM of MNNG every 10,000 min (Figure 14 c). These results suggest that exposure of

tumor tissue composed of p53-proficient and -deficient cells to a reagent that induces a damage

response may lead to preferential expansion of the p53-deficient cell population that contains aneuploid

cells with reproductive activity. If the deficient cells have a malignant nature, p53 could indirectly

promote, rather than suppress the formation of a malignant tumor by suppressing the expansion of the

p53-proficient cell population. Understanding the role of p53 in tumorigenesis thus requires the

delineation of its role at both the cellular and population levels. Furthermore, because most cancer cells

carrying p53 gene mutations express mutated p53 in the cytoplasm instead of the nucleus, cytoplasmic

mutated p53 has been suggested to confer a cellular function, referred to as gain-of-function [62-67]. Such

cells may have a combined phenotype of both loss of wild type p53 and gain-of-function, or mutated

p53 may alter the phenotype caused by the loss of the wild type p53. Nevertheless, these need to be

considered in the context of p53-related tumorigenesis, and the current approach may be used to gain

clues to reveal the context.

Cell-fate simulation and its applications

Compared with other analytical methods, single-cell tracking analysis has the ability to accurately

detect various cellular events, regardless of the frequency of their occurrence, by creating

spatiotemporal data for individual cells. However, because single-cell tracking analysis is a

morphological observation-based technique, we were unable to determine the silencing levels of p53 in

individual cells. The availability of such information would allow deeper analysis; e.g., to examine the

relationship between p53-silencing levels and the chance of cell fusion. Although it is difficult to

monitor expression levels of p53, or other proteins or genes, in individual live cells using existing

methods, new technologies combining single-cell tracking analysis with spatial transcriptomics [68, 69],

for example, could be developed to expand the applicability of this type of analysis. Such expansion

could also provide additional flexibility for cell-fate simulations. We created a cell-fate simulation

algorithm to allow the flexible design of various types of virtual experiments by creating Operation

data, containing data generated by the categorization of event patterns. Such patterns could be further

enriched by the addition of information on the expression levels of proteins or genes in individual cells,

allowing the development of another new type of cell-fate simulation algorithm. Furthermore, given

that each type of cell population tends to show a specific cell population expansion profile and response

to treatment, the accumulation of public Operation data resources could be made accessible to various

research projects to simulate the fate of cells of interest, predict the responses of cells to treatments,

build models based on the simulations, and develop a theory-based model for cell behaviors. Because

single-cell tracking data contains information on the motility of each cell, it may also be possible to

simulate cell behaviors that could be influenced by cell-to-cell contact. This simulation-based biology

could thus help to overcome some of the limitations associated with empirical cell biological studies,

and provide greater flexibility in research with mammalian cells.

Conclusions

p53 may affect cell population dynamics, and loss of p53 could lead to the formation of an aneuploid cell population. In addition, the dynamics of cell populations containing both p53-null cells and wild-type cells may be affected by the status of the wild-type cells. Our results suggest that wild-type cells could gain a growth advantage over p53-null cells in low-stress environments, while the growth balance between the p53-null and wild-type cells may change under stress conditions, allowing p53-null cells to gain a growth advantage. The role of p53 in the process of tumorigenesis may thus depend on the environment surrounding the cell population, as well as the p53 status of individual cells within the population. Revealing the status of individual cells by single-cell tracking, cell-lineage analysis, and cell-fate simulation can thus further our understanding of the process of tumorigenesis.

   KEY RESOURCES TABLE

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
antibody (mouse)	Mouse anti-p53 (DO-1, monoclonal)	Invitrogen	AHO0152	1 µg/mL
antibody (goat)	Horseradish peroxidase conjugated goat anti-mouse antibody (polyclonal)	Abcam	Ab205719	1:25,000
chemical compound, drug	N-methyl -N'-nitroso-N-nitrosoguanidine	Sigma-Aldrich	129941	
cell line (human)	A549-luc-C8	Xenogen Corporation	N/A	
sequence-based reagent	p53 siRNA I	Cell Signaling Technology	6231	
sequence-based reagent	Control siRNA	Cell Signaling Technology	6568	
software, algorithm	MetaMorph (Quorum WaveFX, v7.8.12.0)	Quorum Technology	N/A	
software, algorithm	Image J.	Schneider et al., 2012	https://imagej.nih.ggov/ij/	
software, algorithm	Computerized single-cell lineage tracking analysis system software	Rancourt, Sato, and Satoh	BioRxiv. doi: https://doi.org/10.1101/508705	

software, algorithm	Cell fate simulation algorithm	This paper		Results: Cell-fate simulation algorithm; Materials and Methods: Overall flow of cell-fate simulation algorithm; Figure 7-Source code
other	Cell lineage data	This paper		Deposited in Dryad

MATERIALS AND METHODS

Cells and cell culture

A549-luc-C8 cells were purchased from Xenogen Corporation and cultured in RPMI containing 10%

fetal bovine serum (RPMI medium) in a humidified atmosphere with 5% CO_2. Cells were confirmed to

be mycoplasma negative. Cells were plated in the center of each well of a cover glass Lab-Tek II 8-

well chamber in 50 µL of cell suspension containing 3,500 cells and left to attach to the cover glass

surface. Culture was continued for 24 h to allow more than 99% of cells attached to other cells, and

culture medium (0.75 mL) was then added to each well and the chamber was viewed under a

microscope (Olympus IX81) after 24 h of plating.

Concurrent long-term live cell imaging, siRNA transfection, and MNNG treatment.

Concurrent long-term live cell imaging was performed as described previously [33, 36]. Briefly, images

were acquired under a microscope using near infrared DIC imaging with a ×20 dry objective

(UPlanSApo, 20×/0.75 NA, α/0.17/FN2G.5) and a ×1.5 coupler (Quorum Technologies) to generate a

×30-equivalent image. Images were acquired by area scanning (5×5 dimension, 512×512 pixels each,

$1.77mm^2$) using multi-dimensional acquisition mode in MetaMorph (Quorum Technology Inc.,

WaveFX, v7.8.12.0) with 34 ms exposure and XY piezo stage. The selected area was close to the

center of the cell population, given that the behavior of cells at the periphery may differ from those at

the center, mainly due to differences in cell density and the chance of intercellular attachments.

Typically, 30 z-planes were acquired every 1 µm, and 512×512 pixel multi-layered .tiff files were

created. Cells were maintained on the microscope stage in an environmental chamber (Live Cell

Instrument, Korea) in a humidified atmosphere with 7.5% CO_2 and images were acquired every 10 min.

Lipofection with scrambled siRNA and p53si RNA (Cell Signaling Technology) was performed

approximately 24 h after cell plating using Effectene Transfection Reagent (Qiagen) with 1 µg siRNA

in 742 µL transfection mixture, according to the supplier's instructions. Image acquisition was started

immediately after siRNA transfection. At 24 h after siRNA treatment, cells were exposed to various

doses of MNNG (Sigma-Aldrich) for 30 min in serum-free RPMI, and the medium was then replaced

with fresh RPMI medium. The evaporation rate of the medium was about 10 µl per 24 h, and each well

of a cover glass Lab-Tek II 8-well chamber thus contained 800 µl of RPMI medium to minimize the

impact of evaporation on cell growth. The RPMI medium contained phenol red to monitor the pH of

the medium, and 7.5% CO_2 was typically required to maintain an optimal pH

Image files corresponding to each field of view were saved using MetaMorph on the hard drive of a

Windows computer. The files were then transferred automatically to a Macintosh computer (OS15)

using in-house software [33]. Unique file names were assigned to each file for archiving, followed by the

creation of an all-in-focus image using in-house software [33]. The focused images were positioned and

the contrast was adjusted to create stitched images (25 images covering 5×5 dimension,

approximately 2500×2500 pixels). The images were then saved in chronological order for display.

File transferring, focused image creation, image positioning, and contrast adjustment were performed

automatically, and eight stitched images that corresponding to each Lab-Tek II 8-well chamber well

were typically created concurrently in an automatic manner [33]. The files were displayed using an in-

house movie player, to monitor the progress of live-cell imaging [33].

DIC image segmentation

Image segmentation was performed using our image segmentation software (for details see [33]). Image

segmentation can be performed by various methods, such as the region growing, by setting an optimal

parameter for an image. However, the optimal parameter for a time point of the image may be difficult

to use for other time points, given that the optimal parameter is affected by the cell density and size, the

relative location of a cell to other cells, the formation of cell debris, and image-quality variations

caused by the microscope system. We typically processed > 8,000 images, and our segmentation

software was therefore developed to perform DIC image segmentation automatically, regardless of

image quality [33]. Briefly, in a DIC image, cells appear as illuminated objects associated with a shadow,

and this rule can be applied to any cells in an image. For example, if a flat cell is located behind a bulky

cell, the flat cell appears as a darker cell, but the cell itself is brighter than its shadow. Our

segmentation software used these characteristics of the DIC images. The mean pixel value of a DIC

image (256 grayscale images) was first adjusted to a grayscale value of 100, and four images were then

created by applying four different pixel threshold values: e.g., > 200, > 180, > 160, and > 150. The

image with the highest threshold value was used first, and connected pixels, representing an area, were

identified by connectivity analysis. The connected pixels were then overlaid on an original DIC image

and the area of connected pixels was extended. The connected pixels were likely to represent the

illuminated cells in the image, and expansion was carried out to identify the possible borders of the

cells. The connected areas were then overlaid on the image created with the second highest grayscale

threshold value. Connectivity analysis was performed, except for the areas where a connected area had

already been set in the previous step. This process allowed us to identify darker cells in a DIC image

than those identified previously. Connected areas identified in the second step were overlaid on the

original DIC image and the area was expanded to find a border of darker cells. The same process was

repeated for the images created by the third and fourth thresholds to identify cells in an image. By this

approach, DIC images of various qualities could be segmented.

Single-cell tracking

Single-cell tracking analysis was performed using in-house software that tracked cells automatically,

detected cellular events, and allowed verification of the tracking results [33]. Single-cell tracking was

started by selecting a connected area (Area 1) that corresponded to a progenitor. Because the

segmentation pattern of the next image was often changed, automatic cell tracking was performed by

identifying the best-matched segmented area in the next image. Briefly, Area 1 was overlaid on the

connected areas in the next time-point image. If Area 1 overlapped with one area in the next image and

the size of the area was within the 2-fold of Area 1, the area in the next image was determined as the

area that corresponded to the cell that was being tracked. If Area 1 overlapped with one area in the next

image, but the size of the area was larger than twice Area 1, the area was likely divided into multiple

areas and one area that overlapped with Area 1 was determined as the area corresponding to the tracked

cell. If multiple areas in the next image overlapped with Area 1, the multiple areas were likely merged

and the merged area was determined as the area corresponding to the tracked cell. This process was

carried out for all selected progenitors and repeated until the predetermined time point. When the

tracking reached the target time point, manual data verification was performed to correct tracking

errors. Detected errors were corrected manually, and automatic single-cell tracking was restarted.

Typically, verification was performed every 100 time points to create error-free cell-lineage data. The

required time for verification was ~48 h for the tracking of 400 Control progenitors and their progeny

for 400 time points. Concerning event detection, different approaches were used for cell division, cell

fusion, and cell death. Bipolar and multipolar cell division were detected by first identifying a mitotic

round cell. Typically, such cells appeared as bright round objects in DIC images. The in-house cell-

tracking software thus built a data library of the shapes of mitotic cells, and determined if a tracked cell

underwent mitosis. During the manual verification step, if an error, such as the detection of a non-

mitotic cell as a mitotic one or failure to detect a mitotic cell, was found, the failed or missed pattern

was incorporated into the software to improve the subsequent detection accuracy. Following the

detection of mitotic cells, if a connected area overlapped with two, three, and four connected areas in

the next image, the software determined that bipolar, tripolar, and tetrapolar cell division, respectively,

had occurred. Concerning cell fusion, if two connected areas corresponding to two different cells

overlapped with the same connected area in the next image, and the area did not divide into multiple

areas in the next 10–20-time points, the software determined that two cells were fused. In the case of

cell death, dead cells typically formed remnants that appeared as bright, irregularly shaped objects. If a

connected area overlapped with an area corresponding to such an object, and the object was also found

in the next 10–20 time-point images, the software determined that cell death had occurred.

Automated cell tracking and manual verification can be used to generate an error-free cell-lineage

database. Furthermore, to ensure complete cell-lineage data, if any progeny of a cell lineage moved

outside of the field of view, all the data corresponding to that lineage were excluded from the analyses.

During single-cell tracking, information on cell-lineage number, cell number, the coordinates of the

cells, the types of an events occurring in the cell, the relationship of a cell to its parent cell, and

information related to fused cells were recorded in the cell-lineage database[33].

Western blotting

Cells were transfected with scrambled siRNA and p53 siRNA and treated with various doses of MNNG

after 24 h. At 48 h after treatment, the cells were harvested and proteins (15 µg) were separated by 10%

sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Western blotting was performed using anti-

p53 antibody (1 µg/mL, DO-1; Invitrogen) and horseradish peroxidase-conjugated goat anti-mouse

antibody (25,000-fold dilution; Abcam). Proteins were visualized using an ECL reagent

(Pierce). Quantitation of p53 was performed using Image J.

Generation of density maps

To create a cell density map, we assigned a value of one to a pixel within the 20-pixel diameter area for

cell density and 100-pixel diameter area for multipolar cell division and cell death density from the

position of the cell or position where multipolar cell division or cell death occurred. If an area

overlapped with other areas, a pixel was assigned the sum of the number of overlapped areas. These

values were used to generate heat maps.

Overall flow of the cell-fate simulation algorithm

The flow of the cell-fate simulation algorithm is shown in Figure 7-figure supplement 2a. The

simulation mode was selected from the available five modes; Standard, Dose simulation, Mix culture,

Switch, and Mix culture-Switch (Figure 7-figure supplement 2a, 1 and Figure 8). Standard mode

performed a simulation by setting a simulation time (Sim. Time end) and an initial number of cells (Init.

No. of cells; progenitors) followed by Operation data creation (Figure 7-figure supplement 2a, 2, and

Figure 7-figure supplement 2b) or uploading, and generation of simulation arrays (Figure 7-figure

supplement 2a, 3, and Figure 7-figure supplement 2b). Operation data held the categorized information

of events, and Simulation arrays (Figure 7-figure supplement 2a, 3) were used to assign the length of

time between two events (Ltime) and event type. Dose response mode was carried out by setting a

simulation time, the initial number of cells, and an intermediate dose. Two sets of Operation data were

then generated or uploaded and virtual Operation data corresponding to the intermediate dose were

created using those Operation data. For example, if the intermediate dose was 5 µM MNNG, virtual

Operation data were generated from Operation data for 3 µM and 7 µM MNNG. Mixed culture mode

performed a simulation for a virtual cell population composed of two cell types. A simulation time and

initial numbers of the first and second cell populations were set. Two sets of Operation data were used

to simulate the proliferation of the first and second cell populations individually. Two sets of

simulation arrays were also created. This mode could be used to simulate the proliferation of a cell

population surrounded by another cell population. Switch mode was used to perform virtual drug

treatment. The simulation times before and after treatment, and the initial number of cells were set.

Two sets of Operation data for the first and second doses were used, and simulation arrays

corresponding to each dose were created. Mixed culture-Switch mode was a combined version of

Mixed culture and Switch. This mode used four sets of Operation data and generated four sets of

simulation arrays. Initial processing was then performed by creating a cell data information array (CDI)

that included the Ltime and event type assigned to each cell (Figure 7-figure supplement 2a, 4, and

Figure 7-figure supplement 2d). Based on the information in the CDI, a virtual cell-lineage database

was created (Figure 7-figure supplement 2a, 5, and Figure 7-figure supplement 2e). After these initial

processes, a repeated cycle of assigning Ltime and event type (Figure 7-figure supplement 2f),

reassigning Ltime and event type to adjust for the effect of cell fusion on Ltime (Figure 7-figure

supplement 2g), and creating a virtual cell-lineage database (Figure 7-figure supplement 2h) were

carried out until the simulation time was reached (Figure 7-figure supplement 2a, 6). If Switch or

Mixed culture-Switch mode was used, the cell-lineage database end was trimmed (Figure 7-figure

supplement 2i), Operation data was switched to corresponding Operation data, and the process

proceeded to the second round of the repeated cycle (Figure 7-figure supplement 2a, 7). When

processing reached the simulation time, the simulation was terminated and the results were displayed.

Cell-fate simulation algorithm: Operation data

Processing related to Operation data, which comprised three types of data, is outlined in Figure 7-figure

supplement 2b. Operation data were either uploaded or generated from a cell-lineage database. When

the Operation data were generated, information held in the cell-lineage database was converted into an

Operation data-Time (Figure 7-figure supplement 2b, 1), Operation data-Event (Figure 7-figure

supplement 2b, 2), and Recovery data (Figure 7-figure supplement 2b, 3). The Operation data-Time

held categorized Ltime by event type; e.g., Ltime between bipolar cell division and cell death (BD-CD

data type) and Ltime from time 0 min to the time that an event occurred in a progenitor (First event

data type) (Figure 7-figure supplement 2b, 1). The Operation data-Event contained the frequency of

events occurred following, e.g. bipolar cell division (BD data type), or multipolar cell division (MD

data type) (Figure 7-figure supplement 2b, 2). Recovery data were used to simulate the rate of recovery

of a cell population following treatment with MNNG. Most cells adopted a static status following

treatment with MNNG; however, some cells regained their reproductive ability, as shown at the end of

the imaging procedure. Typically, such cells underwent bipolar cell division to produce two offspring.

However, live-cell imaging would be terminated before the offspring underwent cell division, as a

culture of reference, non-treated cells become confluent. In this case, it was not possible to determine

the cell-doubling time of the offspring. Recovery thus determined the percentage of cells that

underwent bipolar cell division over a certain period from the end of the tracking time. For example, if

the tracking time was 4,000 min, the percentage of cells was calculated by the number of bipolar cell

divisions that occurred within; e.g., 3,200–4,000 min, and the number was converted into a percentage

by dividing the total number of bipolar cell divisions that occurred during the tracking time and then

multiplying by 100 (Figure 7-figure supplement 2b, 3). Thus, if 10 bipolar cell divisions were found

between 3,200–4,000 min out of a total of 300 bipolar cell divisions (entire time), the result was 3.3%.

The simulation algorithm assigned bipolar cell division to 3.3% of non-dividing cells at around 4,000

min. Ltime, which was > 80% of tracking time, was then assigned to cells produced by bipolar cell

division. Using Dose response mode, virtual Operation data were created from two sets of Operation

data using the formula: virtual Operation data = Operation data 1 + [(Operation data 2 − Operation data

1)/(Dose 2 −Dose 1)]×(Intermediate dose − Dose 1) (Figure 7-figure supplement 2b, 4).

1009

Cell-fate simulation algorithm: Simulation arrays

The dataset stored in Operation data was then converted into Simulation arrays that were used to assign

Ltime and event types (Figure 7-figure supplement 2c). In the Operation data, each categorized

Operation data-Time was stored in the format; i.e., Ltime, and the number of Ltime that was found. For

example, 20 min of Ltime found four times was written as 20:4. In the Simulation arrays corresponding

to the Operation data-Time, this was converted to 20, 20, 20, 20, for example. If no corresponding type

of event occurred, the array for the event held no data. To generate a simulation array from the

Operation data-Event, the percentage of each event relative to the total number of cell divisions (total

of bipolar cell division and multipolar cell division) was calculated. The maximum number of entries

per array was 100. The total percentage of events could thus exceed 100, e.g. first event list (bipolar

cell division, cell death, multipolar cell division, and the number of cells with no cell division

(nonDiv)). In this case, the percentage of each event was adjusted to make a total of 100. If the

percentage of the event was < 100, 0 was filled to make the total entry for the array 100. If no

corresponding type of event occurred, an array for the event held no data. The Recovery percentage

was calculated as described above.

1025

Cell-fate simulation algorithm: Initial Ltime and event-type assignment.

In the first event assignment (Figure 7-figure supplement 2d), the CDI that held the status of each cell

throughout a simulation was created. In this step, the cell-lineage number was generated following the

initial number of cells set previously, and the cell number 0 was assigned to each progenitor. Using the

FirstEvent array, Ltime was then assigned to each cell. If the array was empty, the average time that

bipolar cell division occurred was calculated and −25% to +25% of the time selected by random

number (C++ rand ()) was assigned as Ltime, and bipolar cell division was then assigned to the cell. If

the FirstEvent array held data, Ltime was assigned by creating a random number, but if Ltime was

equal to the length of single-cell tracking, −30% to +30% of the length of the time of single-cell

tracking was assigned as Ltime and the event type was selected using the FirstEvent list array. If there

was no entry in the array, NonDiv was assigned, otherwise, bipolar cell division, multipolar cell

division, or cell death was assigned. If cell death was assigned, Ltime was reassigned using the

NonDivCD time data array.

1039

Cell-fate simulation algorithm: Initial cell-lineage database creation

The cell-lineage database was created using information held by the CDI (Figure 7-figure supplement

2e). This array contained information regarding Ltime and event type as a blueprint of the virtual cells

(Figure 7-figure supplement 2e). If Ltime exceeded the simulation time set previously, Ltime was

adjusted to a time equal to the simulation time, otherwise, a cell-lineage database was generated

composed of an X position, Y position, time point, event type, cell-lineage number, cell number, parent

cell information, cell number and the cell-lineage number of the cell that was fused in the event of cell

fusion. If the event type was bipolar cell division or multipolar cell division, information related to the

progeny created by bipolar cell division or multipolar cell division was entered into the CDI following

the generation of corresponding cell numbers for each progeny.

1050

Cell-fate simulation algorithm: Ltime and event-type assignment

In the repeated assignment cycle (Figure 7-figure supplement 2a, 6), the next Ltime and event type was

determined based on the current event type. If the current event in a cell was NonDiv (Figure 7-figure

supplement 2f, 1), a randomized number between 0 and 100 was generated, and if the number was

lower than the Recovery %, bipolar cell division was assigned to the cell. In the next cycle, this cell

was entered into a growing cycle. If the Ltime was 80%–100% of the length of the single-cell tracking

time, the Ltime was determined by generating random numbers between 80 and 100. Assignment of

bipolar cell division in this manner only occurred once in the first cycle of the assignment. If the

random value was higher than the Recovery %, either NonDiv or cell death was assigned, in which

case the % of cell death of total cell division was calculated and, if the random number between 0 and

100 was lower than the cell death %, a cell death was assigned. In this case, Ltime was assigned using

the NonDivCD time array. If the random number between 0 and 100 was higher than the cell death %,

NonDiv was assigned, and the length of single-cell tracking time was assigned as Ltime.

Next, if the current event was bipolar cell division, information related to its sibling was searched

for. If a sibling was found, either bipolar cell division, multipolar cell division, or cell death was

assigned using the Bipolar cell division list array, otherwise bipolar cell division, multipolar cell

division, cell fusion, or cell death was assigned. If the event assigned by the Bipolar cell division list

array was cell fusion, then its sibling was again searched. If no sibling was found, Ltime was assigned

by the BDCF time array. If the array was empty, the event type was changed to cell death and the

average time that cell death occurred was set as Ltime. If a sibling was found, Ltime was assigned

using the BDCF time array. If the array was empty, cell fusion was changed to cell death and the

average cell death time was set as Ltime. If Ltime was set, but it was longer than the Ltime of its

sibling, Ltime was made shorter than the sibling's Ltime. If the event assigned using the Bipolar cell

division list array was bipolar cell division, Ltime was assigned by the BDBD time array. If the array

was empty, the average bipolar cell division time was set as Ltime. If Ltime was set by the BDBD time

array but the Ltime was not within −10% to +10% of Ltime of its parent cell, Ltime was reselected

until the Ltime fell within this range. If Ltime assigned by the Bipolar cell division list array was

multipolar division, Ltime was assigned by the BDBD time array. If the array was empty, the average

bipolar cell division time was set as Ltime. If the event assigned by the Bipolar cell division list array

was cell death, Ltime was assigned the BDCD time array. If the array was empty, the average cell

death time was set as Ltime.

If the current event was multipolar cell division, information related to its siblings was searched for.

If siblings were found and cell fusion was assigned to two siblings, either bipolar cell division,

multipolar cell division, or cell death was assigned by the Multipolar cell division list array, otherwise

bipolar cell division, multipolar cell division, cell fusion, or cell death was assigned. If the event

assigned by the Multipolar cell division list array was cell fusion, then, event type of its sibling was

again searched. If no sibling was found, Ltime was assigned by the MDCF time array. If the array was

empty, the event type was changed to cell death, and the average time that cell death occurred was set

as Ltime. If only one sibling was found, Ltime was assigned by the MDCF time array. If the array was

empty, cell fusion was changed to cell death and the average cell death time was set as Ltime. If two

siblings were found and cell fusion was assigned to one of the siblings, Ltime was assigned by the

MDCF time array. If the array was empty, cell fusion was changed to cell death and the average cell

death time was set as Ltime. If Ltime was longer than the Ltime of its siblings, Ltime was made shorter

than its siblings. If cell fusion was not assigned to any of its siblings, Ltime was assigned by the MDCF

time array. If the array was empty, cell fusion was changed to cell death and the average cell death time

was set as Ltime. If the event assigned by the Multipolar cell division list array was bipolar cell

division, Ltime was assigned by the MDMD time array. If the array was empty, the average bipolar cell

division time was set as Ltime. If Ltime assigned by the Multipolar cell division list array was

multipolar cell division, Ltime was assigned by the MDMD time array. If the array was empty, the

average bipolar cell division time was set as Ltime. If the event assigned by the Multipolar cell division

list array was cell death, Ltime was assigned by the MDCD time array. If the array was empty, the

average cell death time was set as Ltime.

1103

Cell-fate simulation algorithm: Readjustment of Ltime in cells that underwent cell fusion

If cell fusion occurred in a cell, the Ltime of the cell being fused was often affected and the Ltime was

readjusted (Figure 7-figure supplement 2g). Fusion between non-siblings was not simulated by this

algorithm. If the current event was bipolar cell division, the next event was assigned by the BDCF lists.

If the next event was bipolar cell division or multipolar cell division, Ltime was assigned by the

DivCFDiv time array, and if the next event was cell death, Ltime was assigned by the BDCFCD array.

If the current event was multipolar cell division, the next event was assigned by the MDCF lists. If the

next event was bipolar cell division or multipolar cell division, Ltime was assigned by the DivCFDiv

time array, and if the next event was cell death, Ltime was assigned by the MDCFCD array.

1113

Cell-fate simulation algorithm: Cell-lineage database creation

The cell-lineage database was created using information held by the CDI (Figure 7-figure supplement

2h). If Ltime exceeded the simulation time set previously, Ltime was adjusted to a time equal to the

simulation time. If the event type was bipolar cell division or multipolar cell division, information

related to the progeny created by bipolar cell division or multipolar cell division was entered into the

CDI following the generation of corresponding cell numbers for each progeny. If the event type was

cell fusion, cells to be fused were searched and relevant information regarding cell-lineage number and

cell number was recorded for the cell.

1122

1123   **Cell-fate simulation algorithm: Trimming the end of the cell-lineage database**

1124   In Switch and Mixed culture-Switch modes, the end of the cell-lineage data was trimmed to the nearest

1125   cell division event (Figure 7-figure supplement 2i) before entering the second repeated cycle of

1126   assignment.

1127

1128   **Statistical analysis**

1129   Statistical analyses were performed by Prism 8.

1130

Naming rule for cells and Operation data

A549 cells treated with scrambled siRNA and p53 siRNA are referred to as Control and p53 RNAi

cells, respectively. Those cells treated with MNNG are named by adding; e.g., Control-MNNG1 (cells

treated with 1 μM of MNNG). If a cell population that is generated by cell fate simulation is referred, -

Sim was added; e.g., Control-Sim and Control-MNNG1-Sim. In the case that cells generated in silico

to remove cell death are referred, cells are named; e.g., Control-Silico(-)cell death. If an Operation

data-Time or Operation data-Events are required to be specified, the name of cells was added following

Operation data; e.g., Operation data-Control.

1139

1140   **ACKNOWLEDGEMENTS**

1141   We thank Amélie Fradet-Turcotte for critical reading of the manuscript. We like to acknowledge the

1142   Bioimaging Platform of Research Centre for Infectious Diseases, CHU de Québec Research Centre for

1143   the technical support of microscopes. This work was supported by the Canadian Foundation for

1144   Innovation and Canadian Institutes for Health Research.

1145

DECLARATION OF INTERESTS

1147 The authors declare no competing interests.

1148

DATA AVAILABILITY

All data generated or analyzed during this study are included in the paper and supporting file. Source

Data files have been provided for Figure 1-figure supplements 2–4, Figures 2, 3, and 4, Figure 4-figure

supplement 1, Figures 5 and 6, Figure 7-figure supplements 1–3, Figure 8-figure supplement 1, Figure

9, Figure 9-figure supplement 1 and 2, and Figures 10–13. Source code has been provided for Figure 7.

Figure 1-videos (cellular events), Figure 2-figure supplements (cell-lineage maps), Figure 2-videos

(single-cell tracking), Figure 2-source data (cell-lineage database), and Figure 7-figure supplement

(cell-lineage maps) have been deposited in Dryad.

REFERENCES

1.   Kandoth, C. et al. Mutational landscape and significance across 12 major cancer types. *Nature* **502**, 333-339 (2013).

2.   Hollstein, M., Sidransky, D., Vogelstein, B. & Harris, C.C. p53 mutations in human cancers. *Science* **253**, 49-53 (1991).

3.   Petitjean, A. et al. Impact of mutant p53 functional properties on TP53 mutation patterns and tumor phenotype: lessons from recent developments in the IARC TP53 database. *Hum Mutat* **28**, 622-629 (2007).

4.   Boutelle, A.M. & Attardi, L.D. p53 and Tumor Suppression: It Takes a Network. *Trends Cell Biol* **31**, 298-310 (2021).

5.   Levine, A.J. The many faces of p53: something for everyone. *J Mol Cell Biol* **11**, 524-530 (2019).

6.   Kastenhuber, E.R. & Lowe, S.W. Putting p53 in Context. *Cell* **170**, 1062-1078 (2017).

7.   Lavin, M.F. & Gueven, N. The complexity of p53 stabilization and activation. *Cell Death Differ* **13**, 941-950 (2006).

8.   Vogelstein, B., Lane, D. & Levine, A.J. Surfing the p53 network. *Nature* **408**, 307-310 (2000).

9.   Hafner, A., Bulyk, M.L., Jambhekar, A. & Lahav, G. The multiple mechanisms that regulate p53 activity and cell fate. *Nat Rev Mol Cell Biol* **20**, 199-210 (2019).

10.   Livingstone, L.R. et al. Altered cell cycle arrest and gene amplification potential accompany loss of wild-type p53. *Cell* **70**, 923-935 (1992).

11.   Kastan, M.B. et al. Levels of p53 protein increase with maturation in human hematopoietic cells. *Cancer Res* **51**, 4279-4286 (1991).

12.   Clarke, A.R. et al. Thymocyte apoptosis induced by p53-dependent and independent pathways. *Nature* **362**, 849-852 (1993).

13.   Lowe, S.W., Schmitt, E.M., Smith, S.W., Osborne, B.A. & Jacks, T. p53 is required for

radiation-induced apoptosis in mouse thymocytes. *Nature* **362**, 847-849 (1993).

14.   Campisi, J. Senescent cells, tumor suppression, and organismal aging: good citizens, bad

neighbors. *Cell* **120**, 513-522 (2005).

15.   Rufini, A., Tucci, P., Celardo, I. & Melino, G. Senescence and aging: the critical roles of p53.

Oncogene **32**, 5129-5143 (2013).

16.   Haupt, Y., Maya, R., Kazaz, A. & Oren, M. Mdm2 promotes the rapid degradation of p53.

Nature **387**, 296-299 (1997).

17.   Honda, R., Tanaka, H. & Yasuda, H. Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor

suppressor p53. *FEBS Lett* **420**, 25-27 (1997).

18.   Kubbutat, M.H., Jones, S.N. & Vousden, K.H. Regulation of p53 stability by Mdm2. *Nature*

387, 299-303 (1997).

19.   Guo, S. et al. Nonstochastic reprogramming from a privileged somatic cell state. *Cell* **156**, 649-

662 (2014).

20.   Vousden, K.H. & Lane, D.P. p53 in health and disease. *Nat Rev Mol Cell Biol* **8**, 275-283

(2007).

21.   Kruiswijk, F., Labuschagne, C.F. & Vousden, K.H. p53 in survival, death and metabolic health:

a lifeguard with a licence to kill. *Nat Rev Mol Cell Biol* **16**, 393-405 (2015).

22.   Lassus, P., Ferlin, M., Piette, J. & Hibner, U. Anti-apoptotic activity of low levels of wild-type

p53. *EMBO J* **15**, 4566-4573 (1996).

23.   Chow, K.K. et al. Imaging cell lineage with a synthetic digital recording system. *Science* **372**

(2021).

24.   Chapal-Ilani, N. et al. Comparing algorithms that reconstruct cell lineage trees utilizing

information on microsatellite mutations. *PLoS Comput Biol* **9**, e1003297 (2013).

25. Zafar, H., Lin, C. & Bar-Joseph, Z. Single-cell lineage tracing by integrating CRISPR-Cas9

mutations with transcriptomic data. *Nat Commun* **11**, 3055 (2020).

26. Wood, R.D. DNA repair in eukaryotes. *Annu Rev Biochem* **65**, 135-167 (1996).

27. Jacobs, A.L. & Schar, P. DNA glycosylases: in DNA repair and beyond. *Chromosoma* **121**, 1-

20 (2012).

28. Lindahl, T. Repair of intrinsic DNA lesions. *Mutat Res* **238**, 305-311 (1990).

29. Satoh, M.S. & Lindahl, T. Role of poly(ADP-ribose) formation in DNA repair. *Nature* **356**,

356-358 (1992).

30. Lindahl, T., Satoh, M.S., Poirier, G.G. & Klungland, A. Post-translational modification of

poly(ADP-ribose) polymerase induced by DNA strand breaks. *Trends Biochem Sci* **20**, 405-411

(1995).

31. Lindahl, T., Sedgwick, B., Sekiguchi, M. & Nakabeppu, Y. Regulation and expression of the

adaptive response to alkylating agents. *Annu Rev Biochem* **57**, 133-157 (1988).

32. Kim, W.J., Beardsley, D.I., Adamson, A.W. & Brown, K.D. The monofunctional alkylating

agent N-methyl-N'-nitro-N-nitrosoguanidine triggers apoptosis through p53-dependent and -

independent pathways. *Toxicol Appl Pharmacol* **202**, 84-98 (2005).

33. Sato, S., Rancourt, A. & Satoh, M.S. Identification of SNA-I-positive cells as stem-like cells in

an established cell line using computerized single-cell lineage tracking. *BioRxiv* **508705; doi:**

https://doi.org/10.1101/508705 (2018).

34. Puck, T.T., Marcus, P.I. & Cieciura, S.J. Clonal growth of mammalian cells in vitro; growth

characteristics of colonies from single HeLa cells with and without a feeder layer. *J Exp Med*

103, 273-283 (1956).

35. Ganem, N.J., Godinho, S.A. & Pellman, D. A mechanism linking extra centrosomes to

chromosomal instability. *Nature* **460**, 278-282 (2009).

36.   Sato, S., Rancourt, A., Sato, Y. & Satoh, M.S. Single-cell lineage tracking analysis reveals that

an established cell line comprises putative cancer stem cells and their heterogeneous progeny.

Sci Rep **6**, 23328 (2016).

37.   Boveri, T. Concerning the origin of malignant tumours by Theodor Boveri. Translated and

annotated by Henry Harris. *J Cell Sci* **121 Suppl 1**, 1-84 (2008).

38.   Keryer, G., Ris, H. & Borisy, G.G. Centriole distribution during tripolar mitosis in Chinese

hamster ovary cells. *J Cell Biol* **98**, 2222-2229 (1984).

39.   Holland, A.J. & Cleveland, D.W. Boveri revisited: chromosomal instability, aneuploidy and

tumorigenesis. *Nat Rev Mol Cell Biol* **10**, 478-487 (2009).

40.   Gisselsson, D. et al. When the genome plays dice: circumvention of the spindle assembly

checkpoint and near-random chromosome segregation in multipolar cancer cell mitoses. *PLoS*

One **3**, e1871 (2008).

41.   Saunders, W. Centrosomal amplification and spindle multipolarity in cancer cells. *Semin*

Cancer Biol **15**, 25-32 (2005).

42.   Emdad, L., Sarkar, D., Su, Z.Z. & Fisher, P.B. Emerging roles of centrosomal amplification and

genomic instability in cancer. *Front Biosci* **10**, 728-742 (2005).

43.   Shi, Q. & King, R.W. Chromosome nondisjunction yields tetraploid rather than aneuploid cells

in human cell lines. *Nature* **437**, 1038-1042 (2005).

44.   Rajagopalan, H. & Lengauer, C. Aneuploidy and cancer. *Nature* **432**, 338-341 (2004).

45.   Harvey, M. et al. In vitro growth characteristics of embryo fibroblasts isolated from p53-

deficient mice. *Oncogene* **8**, 2457-2467 (1993).

46.   Valente, L.J. et al. p53 deficiency triggers dysregulation of diverse cellular processes in

physiological oxygen. *J Cell Biol* **219** (2020).

47.   Husemann, Y. et al. Systemic spread is an early step in breast cancer. *Cancer Cell* **13**, 58-68

(2008).

48. Riethmuller, G. & Klein, C.A. Early cancer cell dissemination and late metastatic relapse:

clinical reflections and biological approaches to the dormancy problem in patients. *Semin*

Cancer Biol **11**, 307-311 (2001).

49. Eischen, C.M. Genome Stability Requires p53. *Cold Spring Harb Perspect Med* **6** (2016).

50. Ganem, N.J., Storchova, Z. & Pellman, D. Tetraploidy, aneuploidy and cancer. *Curr Opin*

Genet Dev **17**, 157-162 (2007).

51. Storchova, Z. & Kuffer, C. The consequences of tetraploidy and aneuploidy. *J Cell Sci* **121**,

3859-3866 (2008).

52. Hernandez, J.M. & Podbilewicz, B. The hallmarks of cell-cell fusion. *Development* **144**, 4481-

4495 (2017).

53. Shinn-Thomas, J.H. & Mohler, W.A. New insights into the mechanisms and roles of cell-cell

fusion. *Int Rev Cell Mol Biol* **289**, 149-209 (2011).

54. Deangelis, P. et al. P53 expression is associated with a high-degree of tumor DNA aneuploidy

and incidence of p53 gene mutation, and is localized to the aneuploid component in colorectal

carcinomas. *Int J Oncol* **3**, 305-312 (1993).

55. Dutertre, S., Hamard-Peron, E., Cremet, J.Y., Thomas, Y. & Prigent, C. The absence of p53

aggravates polyploidy and centrosome number abnormality induced by Aurora-C

overexpression. *Cell Cycle* **4**, 1783-1787 (2005).

56. Galipeau, P.C. et al. 17p (p53) allelic losses, 4N (G2/tetraploid) populations, and progression to

aneuploidy in Barrett's esophagus. *Proc Natl Acad Sci U S A* **93**, 7081-7084 (1996).

57. Ramel, S. et al. Inactivation of p53 and the development of tetraploidy in the elastase-SV40 T

antigen transgenic mouse pancreas. *Pancreas* **11**, 213-222 (1995).

58. Fujiwara, T. et al. Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-null

cells. *Nature* **437**, 1043-1047 (2005).

59. Thompson, S.L. & Compton, D.A. Examining the link between chromosomal instability and

aneuploidy in human cells. *J Cell Biol* **180**, 665-672 (2008).

60. Vitale, I. et al. Multipolar mitosis of tetraploid cells: inhibition by p53 and dependency on Mos.

EMBO J **29**, 1272-1284 (2010).

61. Li, R. & Zhu, J. Effects of aneuploidy on cell behaviour and function. *Nat Rev Mol Cell Biol*

(2022).

62. Acin, S. et al. Gain-of-function mutant p53 but not p53 deletion promotes head and neck cancer

progression in response to oncogenic K-ras. *J Pathol* **225**, 479-489 (2011).

63. Amelio, I. & Melino, G. Context is everything: extrinsic signalling and gain-of-function p53

mutants. *Cell Death Discov* **6**, 16 (2020).

64. Bargonetti, J. & Prives, C. Gain-of-function mutant p53: history and speculation. *J Mol Cell*

Biol **11**, 605-609 (2019).

65. Oren, M. & Rotter, V. Mutant p53 gain-of-function in cancer. *Cold Spring Harb Perspect Biol*

2, a001107 (2010).

66. van Oijen, M.G. & Slootweg, P.J. Gain-of-function mutations in the tumor suppressor gene p53.

Clin Cancer Res **6**, 2138-2145 (2000).

67. Yue, X. et al. Mutant p53 in Cancer: Accumulation, Gain-of-Function, and Therapy. *J Mol Biol*

429, 1595-1606 (2017).

68. Hou, X. et al. Integrating Spatial Transcriptomics and Single-Cell RNA-seq Reveals the Gene

Expression Profiling of the Human Embryonic Liver. *Front Cell Dev Biol* **9**, 652408 (2021).

69. Joglekar, A. et al. A spatially resolved brain region- and cell type-specific isoform atlas of the

postnatal mouse brain. *Nat Commun* **12**, 463 (2021).

1300

1301

 FIGURE LEGENDS

 Figure 1. System for investigating the functional role of maintaining low cell levels of p53. A549

 cells were cultured on multi-well chamber slides and the areas of interest were scanned. Cells were

 treated with scrambled siRNA or p53 siRNA in the presence or absence of various doses of MNNG,

 and the cell responses were monitored to generate live cell imaging videos. The videos were used for

 imaging and single-cell tracking, starting immediately after MNNG treatment. Progenitors were

 selected from the image designated as tracking time 0 min (T0), and the progenitors and their progeny

 were tracked empirically. Cell-lineage database comprised tracking results for one progenitor and their

 progeny, and any events that occurred in the cells. A cell-lineage map was generated from the database

 to represent the proliferation profile of a progenitor and its progeny through bipolar cell division (BD),

 multipolar cell division (MD), cell fusion (CF), and cell death (CD). If any progeny moved out from

 the areas of interest, data related to the lineage of that was not included in the cell-lineage database.

 Quantitative bioinformatics analysis was performed using the database, which was also used to develop

 a cell-fate simulation algorithm.

 Figure 2. Cell-population expansion rates determined by cell counting and single-cell tracking

 analysis. Control (**a**) and p53 RNAi (**b**) cells were exposed to 1, 3, and 7 µM MNNG (Control and p53

 RNAi cells exposed to those doses of MNNG are referred to as MNNG1, MNNG3, MNNG7, p53

 RNAi-MNNG1, p53 RNAi-MNNG3 and p53 RNAi-MNNG7, respectively). (**a-f**) Images were divided

 into a 5×5 squares and cell population expansion curves were determined by counting the number of

 cells in a square (~ 512×512 pixels) at 1,000 min intervals (see Figure 2-figure supplement 1). Fifteen

 squares were selected from three independently generated videos. Expansion curves were compared as

 follows: Control vs. p53 RNAi (**c**), MNNG1 vs. p53 RNAi-MNNG1 (**d**), MNNG3 vs. p53 RNAi-

 MNNG3 (**e**), and MNNG7 vs. p53 RNAi-MNNG7 (**f**) cells. Student's *t*-test (**c-f**) and ordinary one-way

 ANOVA (Tukey's) (**a** and **b**) were performed (n=15), and standard errors are shown. Cell population

expansion curves of Control (**g**) and p53 RNAi (**h**) cells were determined using a cell-lineage database.

Single-cell tracking was performed for 4,000 min with videos acquired at 10 min intervals (n = 399).

The number of cells every 10 min was extracted and plotted. Cell population expansion curves were

compared as follows: Control vs. p53 RNAi (**i**), MNNG1 vs. p53 RNAi-MNNG1 (**j**), MNNG3 vs. p53

RNAi-MNNG3 (**k**), and MNNG7 vs. p53 RNAi-MNNG7 (**l**) cells. (**i-l**) The data were analyzed by

Student's *t*-tests in the following approach. The difference in cell numbers between n and n + 10 min

was calculated (e.g., if the numbers of cells at 10 and 20 min were 100 and 103, the difference was 3).

Because the total number of imaging points was 400, the 399 differences in time points can be

calculated. Those differences for Control and p53 RNAi cells and those exposed to MNNG were used

for the statistical analyses. (**a-f** and **i-l**) NS: non-significant, $^{*}P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$, $^{****}P <$

0.0001.

1338

Figure 3. Number of progeny produced from a progenitor. (**a**) Example cell lineages are shown. If

a cell lineage was composed of one progenitor [1] and 4 progenies [2–5] (a: left), the cell lineage was

grouped as C (5–6). Similarly, if a cell lineage was composed of one progenitor [1] and 12 progenies

[2–13] (b: right), the lineage was grouped as G (13–14). Cells that underwent multipolar cell division,

cell death, and cell fusion were also included in this analysis. (**b** and **c**) The total number of cells

belonging to each group was calculated. For example, if 5 cell lineages were composed of 3 cells

(Group B), the number of cells in Group B was $5 \times 3 = 15$, which could represent the number of cells

with the reproductive ability to produce 3 cells within 4,000 min of culture in a cell population. **b**:

Control cells and cells exposed to MNNG. **c**: p53 RNAi cells and cells exposed to MNNG.

1348

Figure 4. Analysis of multipolar cell division, cell death, and cell fusion using counting and single-

cell tracking analysis. Control and p53 RNAi cells were exposed to 1, 3, and 7 μM MNNG. The

numbers of multipolar cell division (**a**) and cell death (**b**) events that occurred during the tracking time

(4,000 min) were determined by counting those events recorded in a video. Each time point of image

was divided into four areas (Figure 4-figure supplement 1), and three independently generated videos

were used. Thus, counting was performed in 12 squares (n = 12). The numbers of multipolar cell

division (**c**), cell death (**d**), and cell fusion (**e**) events that occurred during the tracking time (4,000 min)

were determined using single-cell tracking and cell-lineage database (n = 335). (**a-e**) Statistical analysis

was performed by ordinary one-way ANOVA (Tukey's). NS: non-significant, [*]$P < 0.05$, [**]$P < 0.01$,

[***]$P < 0.001$, [****]$P < 0.0001$. Standard errors are shown.

1359

Figure 5. *In silico* generation of cell-lineage database to evaluate the impact of cell death on the

cell population expansion. (**a**) To evaluate the impact of cell death on population expansion, we

created an *in silico* cell-lineage database, in which cells do not undergo cell death, to address the

impact of cell death on the cell population expansion. In the example cell lineages, the number of

survived cells was reduced by half as a result of cell death. If the siblings of the dead cell grew, the

dead cell was assumed to continue to grow similar to its sibling, and a cell-lineage database was created

in silico based on this assumption. (**b**) A cell population expansion curve was generated using the *in*

silico-created cell-lineage database for Control and p53 RNAi cells (Control-Silico(-)cell death) and

p53 RNAi-Silico(-)cell death, respectively). At a tracking time of 4,000 min, cell death resulted in a

7.4% reduction in the population size of p53 RNAi cells. The frequency of cell death (red lines)

occurring during tracking is also shown. Statistical analysis was performed by ordinary one-way

ANOVA (Tukey's). [*]$P < 0.05$.

1372

Figure 6. Events leading to the induction of multipolar cell division. (**a**) Cellular events that

occurred prior to multipolar cell division were analyzed by tracing the cell lineage backward. Pattern 1a

(cell fusion (CF) ➤ multipolar cell division (MD)): cell fusion occurred prior to multipolar cell

division. Pattern 1b (➤ MD): no cell fusion occurred. The numbers of each pattern occurring in 50 cell

lineages are shown. Percentages of Patterns 1a+1b in Control and p53 RNAi cells are also shown. (**b**)

Cells that underwent multipolar cell division were identified. If the cells were generated by cell fusion,

the origin of the fused cells was searched by tracing the cell lineage backward. Pattern 2a (BD ➤

Sibling CF ➤ MD): bipolar cell division occurred, cells produced by division were fused, followed by

multipolar cell division. Pattern 2b (BD ➤ Non-Sibling CF ➤ MD): one sibling produced by bipolar

cell division was fused with another non-sibling cell, followed by multipolar cell division. The numbers

of each pattern in 50 cell lineages are shown. Percentages of Patterns 2a+2b in Control and p53 RNAi

cells are also shown. (**c**) Progeny produced by multipolar cell division was tracked and the number of

reproductive progeny that underwent bipolar cell division was determined. The numbers of

reproductive progeny found in 100 cell lineages and the percentages of such progeny among the total

number of progeny produced by multipolar cell divisions are shown. (**d**) The number of multipolar cell

division events was plotted against the number of bipolar cell division events. Numbers in (**d**) indicate

doses of MNNG. (**a-d**) BD: bipolar cell division, MD: multipolar cell division, CD: cell death, and CF:

cell fusion. Because the number of multipolar cell divisions in Control cells was low, we searched three

videos to find multipolar cell divisions and then traced the cells back to their progenitors to determine

the events preceding multipolar cell division. To normalize the values, we assumed that multipolar cell

division occurred at a similar frequency to that recorded in the cell-lineage database.

1394

Figure 7. Schematic illustration of cell-fate simulation process. (**a**) A cell lineage is composed of a

progenitor and its progeny. Each of those cells is generated by a Start event (blue arrowhead), followed

after a certain length of time by an End event (red arrowhead), such as bipolar cell division (BD),

multipolar cell division (MD), cell death (CD), and cell fusion (CF). Cell-fate simulation was

performed by combining these components, types of event, and length of time between the two events.

To perform the simulation, cells were classified based on the combination of the Start and the End

events; e.g., BD-BD, and BD-MD (see Figure 7-figure supplement 2b for full list of classifications),

and two sets of histogram data, referred to as Operation data-Events and Operation data-Time were

created. (**b**) The Operation data-Events holds the frequency of events that occurred following a Start

event, such as BD, MD, or CF. In the case of the example shown in (**b**), if the Start event was BD, the

histogram data indicated that the chances of the occurrence of each End event type; i.e., BD, MD, CD,

and CF, were ten, five, two, and three, respectively. In the example, MD and BD were selected from

the histogram to assign an End event type to daughter cells produced by BD. Selection was performed

by generating a random value. (**c**) Operation data-Time was then used to assign the length of time

between the Start and End events using histogram data (Operation data-Time) for BD-BD and BD-MD,

which included the choices of length of time. The processes were repeated to generate cell-lineage data

for virtual cells.

1412

Figure 8. Operation modes. (**a**) Standard mode: cell-fate simulation with one type of cell population

performed with Operation data. (**b**) Dose simulation mode: Operation data created from other

Operation data. The example in (**b**) represents an Operation data-Events, which holds information on

the chance of the occurrence of subsequent events following an event, as outlined in Figure 7c. In the

case of Operation data A, generated from a cell-lineage database created by tracking cells exposed to a

dose of a drug (e.g., 2 µM MNNG), 10, 5, 2, and 4 bipolar cell division (BD), multipolar cell division

(MD), cell death (CD), and cell fusion (CF) events, respectively, occurred following the event. On the

other hand, Operation data B, generated from a cell-lineage database created by tracking cells in

another cell population exposed to a different dose (e.g., 5 µM) of the drug, assumed that 1, 5, 4, and

10 BD, MD, CD, and CF events would occur following the same type of events referred to in

Operation A. If a cell population was assumed to be exposed to the drug at 3 µM, Operation data for

the exposed cells could be created by calculating; e.g., BD (7) = BD for Operation data A (10) +

Rancourt et al.   62

[(Operation data B (1) – Operation data A (10)/(Dose 2 (×5) – Dose 1 (×2))) × (Intermediate dose (×3)

– Dose 1 (×2)). A similar calculation can be performed for all data stored in Operation data-Time and -

Events to create new Operation data (Operation data-virtual). Using the Operation data, the cell fate

could be simulated without the need for empirical analysis. (**c**) Mixed culture mode: assuming that

multiple cell types coexist in a culture, simulation was performed using corresponding Operation data

for each cell type. (**d**) Switch mode: cell-fate simulation is performed with Operation data and then

switched to another Operation data. If the second Operation data is for cells treated with a drug, this

simulation allow the effect of the drug treatment to be evaluated virtually. (**e**) Mixed culture-Switch

mode: this mode is a combined Mixed culture (**c**) and Switch (**d**) mode.

1434

Figure 9. Simulations of effects of p53 silencing and MNNG exposure on cell population

expansion. Simulations were performed using Operation data generated from cell-lineage data with a

total of 5,000 virtual progenitors for a simulation time of 15,000 min. (**a** and **b**) Virtual cell population

expansion curves of Control-Sim cells and cells exposed to various doses of MNNG (**a**: 0–4,000 min,

and **b**: whole scale, 0–15,000 min). (**c**, **d**, **e**, **f**) Data obtained from the simulation were compared with

cell population expansion curves determined by single-cell tracking analysis; (**c**) Control, (**d**) MNNG1,

(**e**) MNNG3, and (**f**) MNNG7. Difference (%) was determined by obtaining the sum of the number of

cells (single-cell tracking)/the number of Simulated cells determined at each simulation time point

divided by 1,500 simulation time × 100. For example, a difference of 2.7% implied that the simulation

generated 2.7% more cells compared with the single-cell tracking data. (**g** and **h**) Virtual cell

population expansion curves of p53 RNAi-Sim cells and cells exposed to various doses of MNNG (**g**:

0–4,000 min, and **h**: whole scale, 0–15,000 min). (**i**, **j**, **k**, **l**) These simulation data were compared with

the cell population expansion curves determined by single-cell tracking analysis; (**i**) p53 RNAi, (**j**) p53

RNAi-MNNG1, (**k**) p53 RNAi-MNNG3, and (**l**) p53 RNAi-MNNG7. Results of simulation obtained

using Operation data-p53 RNAi-Silico(-)cell death were also included in **g** and **h**.

1450

Figure 10. Virtual dose-response analysis. Virtual Operation data corresponding to each 0.5 µM dose of MNNG was generated in Dose simulation mode. (**a, b**) Virtual MNNG-dose-response cell population expansion curves of Control and p53 RNAi cells generated by the simulation using Operation data-Control to MNNG 1.5 (purple), 2–3 (pink), 3.5–6.5 (dark green), and 7 (marron) are shown. (**c**) The numbers of cells at simulation times of 4,000 min (indicated by a blue arrow in (**a**) and (**b**)) were plotted. Results of single-cell tracking analysis are also included. (**d**) The numbers of cells at simulation times of 15,000 min (indicated by a red arrow in (**a**) and (**b**)) were plotted. The difference of cell number between Control cells and the cells exposed to 7 µM MNNG, and p53 RNAi cells and the cells to the dose of MNNG are shown (Italic). The number of cells at 0 and 15,000 min were also shown. (**c** and **d**) Standard deviations are shown. A natural log scale was used. (**e**) Time necessary to reach a cell population to be doubled (100 to 200 cells) was determined by using Operation data-Control, MNNG7, p53 RNAi, and p53 RNAi-MNNG7. The difference in time between MNNG7-Sim (9,651 min) and Control-Sim (2,501 min) was 7,150 min. Similarly, the difference in doubling time between p53 RNAi-MNNG7-Sim and p53 RNAi-Sim was 4,510 min. Assuming that the difference in doubling time between MNNG7-Sim and Control-Sim represents the effect of MNNG induced by e.g. DNA breaks and p53-mediated responses, while that between p53 RNAi-MNNG7-Sim and p53 RNAi-Sim only represents the effect of MNNG induced by e.g. DNA breaks, 63% and 37% of suppression of cell population expansion in MNNG7-Sim were likely to be due to the effect of MNNG caused by e.g. DNA breaks and responses induced by the stabilization of p53, respectively.

1470

Figure 11. Simulation of proliferation of progeny of multipolar cell division. The simulation was performed using Operation data-Control and p53 RNAi in Standard mode. (**a**) The numbers of progeny of bipolar cell division (grey), multipolar cell division (burgundy), and progeny cells that first underwent multipolar cell division followed by bipolar cell division (green) were plotted. The

simulation was carried out for 2,500 progeny, which generated 6.6×10^6 cells during the 20,000 min of

simulation. (**b** and **c**) Whole-scale and magnified images generated by the simulation using the

Operation data-Control are shown in (**b**) and (**c**), respectively. (**d**) An animation was created to

visualize the proliferation of Control-sim cells (Figure 11-video 1) and an image at simulation time

20,000 min is shown. Grey, burgundy, and green dots represent cells that underwent bipolar division,

multipolar division, and multipolar division followed by bipolar division, respectively. (**e** and **f**) Whole-

scale and magnified images generated by the simulation using Operation data-p53 RNAi are shown in

(**e**) and (**f**), respectively. (**g**) An animation was created to visualize the proliferation of p53 RNAi-sim

cells (Figure 11-video 2) and an image at simulation time 20,000 min is shown. Bipolar cell division

(BD), multipolar cell division (MD), and multipolar cell division followed by bipolar cell division

(MD-BD).

1486

Figure 12. Limited expansion of p53 RNAi cells in a Control cell population. We simulated the

expansion of the p53 RNAi cell population (4% at the initial time point) in a Control cell population

(96% at the initial time point) using Mixed culture mode with Operation data-Control and p53 RNAi

for a simulation time of 20,000 min with 500 progenitors and repeating 10 simulations (total 5,000

progenitors). (**a**) Simulation of 480 Control progenitors was carried out using Operation data-Control

and 20 p53 RNAi progenitors with Operation data-p53 RNAi. The initial percentage of p53 RNAi cells

was 4%, which was reduced to 2.4% at 20,000 min. (**b**) The sums of multipolar cell division, cell death,

and cell fusion events (MD+CD+CF) are shown. (**c**) Simulation was performed using Operation data-

p53 RNAi-Silico(-)cell death.

1496

1497 **Figure 13. Expansion of p53 RNAi cells in a Control cell-dominant cell population subjected to**

1498 **virtual MNNG exposure.** We simulated the response of p53 RNAi cells in a Control cell-dominant

1499 cell population to MNNG exposure using Mixed culture- Switch mode. (**a**) The simulation was

1500 performed with 96% Control and 4% p53 RNAi cells, using Operation data-Control and p53 RNAi for

1501 9,900 min. The Operation data was then switched to; e.g., MNNG7 and p53 RNAi-MNNG7 for virtual

1502 MNNG treatment, and the simulation was continued up to 20,000 min. In the animation (Figure 13-

1503 videos 1), the pixel value of the image was reduced to indicate the virtual MNNG treatment. (**b-f**) In

1504 the simulation, Operation data corresponding to (**b**) 1.5, (**c**) 3, (**d**) 5, (**e**) 6, and (**f**) 7 µM MNNG were

1505 used to perform a virtual MNNG treatment. The numbers in the images indicate the percent of p53

1506 RNAi cells at simulation times 0 and 20,000 min. (**g**) The percentages of p53 RNAi cells in the

1507 population were plotted. Statistical analysis (n=10) was performed using data at simulation time 20,000

1508 min by ordinary one-way ANOVA. NS: nonsignificant, $^{*}P < 0.05$, $^{****}P < 0.0001$. (**h**) Based on the

1509 simulation data for 7 µM MNNG, percentage of p53 RNAi cells following repeated 7 µM MNNG

1510 exposure every 10,000 min (~7days) was calculated. Arrows indicate the time that the virtual exposure

1511 was performed.

1512

1513 **Figure 14. Role of the low levels of p53 in suppression of multipolar cell division, cell death, cell**

1514 **fusion, and damage response.** (**a**) Characteristics of Control and p53 RNAi cells. (**b**) The simulation

1515 of the fate of p53 RNAi cells in the Control cell population. (**c**) The simulation of the fate of p53 RNAi

1516 cells in the Control cell population, which were virtually exposed to 7 µM MNNG. The percentage of

1517 aneuploid cells is also shown.

1518

 SUPPLEMENTARY FIGURES

1520

Figure 1-figure supplement 1. Events occurring in the cells. Cellular events that occurred in A459

cells were categorized as follows: bipolar cell division, tripolar cell division, tetrapolar cell division,

cell death, and cell fusion. In the current study, tripolar and tetrapolar cell divisions were defined as

multipolar cell division. Green arrowheads indicate cells that underwent an event. Red arrowheads

indicate daughter cells, a dead cell, and a fused cell. Timestamps (min) relative to the first frame and a

scale bar (20 μm) are shown. Corresponding videos are deposited in Dryad.

1527

Figure 1-figure supplement 2. Quantitation of cell numbers, areas, and densities. (**a**) Cell density

and average cell size were plotted every 10 min. (**b**) Cell density maps were created using a heat map

scale. Still images were taken from Figure 1-video 1. Timestamps (min) and a scale bar (80 μm) are

shown. (**c**) The cell positions that were determined using single-cell tracking and recorded in the cell-

lineage database were used. Cell density map created by assigning value 1 to a pixel within the 20-pixel

diameter area from the position of a cell (blue); if an area overlapped with other areas, the pixel was

assigned the sum of the number of overlapped areas (light green). At the high cell density area, values

were increased according to the number of overlaps, allowing to generate a heat map. These values

were used to generate heat maps.

1537

Figure 1-figure supplement 3. Determination of minimum number of cell-lineages required to

build a cell-lineage database. A cell-lineage database containing 50–240 cell-lineages was used to

determine cell population expansion curves. Cell-lineage database was created by tracking of 485

progenitors of Control cells. (**a-e**) Cell population expansion curves determined by randomly selecting

50 (**a**), 80 (**b**), 120 (**c**), 140 (**d**), and 240 (**e**) cell-lineages from the cell-lineage database. The number of

cells every 10 min was calculated and plotted. The random selection was repeated three times (green,

orange, and red lines) to show variations at a tracking time of 4,000 min (black bars). Each repeated

selection was performed in a way that the same lineage was not selected, except for the selection of 240

lineages. The initial number of cell-lineages was normalized to 100. (**f**) Variations of 4,000 min were

plotted.

1548

Figure 1-figure supplement 4. Accumulation of particles following siRNA transfection and

analysis of p53 expression following transfection and/or MNNG exposure. (**a**) Particles formed

following siRNA transfection are shown. DIC and objects with pixel value 150–254 and size 5–30

pixels were extracted from the DIC image (Particles). (**b**) The numbers of particles at five different

locations in the image were plotted. Blue lines represent the average number of particles at each time

point per location and the red line shows the overall average. MNNG treatment was performed after

approximately 1,740 min of siRNA transfection, and continued for 30 min. The number of particles

peaked at 3,750 min after transfection. Single-cell tracking was started immediately after MNNG

treatment and continued for 4,000 min. (**c**) Western blotting with anti-p53 antibody was performed 48 h

after siRNA treatment. Amounts of p53 relative to Control are shown.

1559

1560   **Figure 2-figure supplement 1. Determination of cell-population expansion rate using video**

1561   **images.** A video image (typically 2,560 pixels × 2,560 pixels) was divided into 5 × 5 squares and the

1562   numbers of cells in selected squares were counted. Counting was performed every 1,000 min in video

1563   images from 0–4,000 min.

1564

Figure 4-figure supplement 1. Counting multipolar cell division and cell death using live-cell

imaging videos. (**a**) Each time point of the video image was divided into four squares, and cell death

and multipolar cell division in each square were marked to determine the numbers of those events that

occurred during imaging. Three independently generated videos were used, and total 12 squares were

thus counted. The positions of multipolar cell division and cell death were also stacked to generate

multipolar cell division and cell death density maps. Multipolar cell division (**b**) and cell death (**c**)

density maps were created using a similar approach to cell density determination (Figure 1-figure

supplement 2b). Density was shown using a heat map scale. Black squares correspond to the area

used for single-cell tracking.

1574

1575 **Figure 7-figure supplement 1. Single-cell tracking and cell-fate simulation processes.** (**a**) Overall

1576 scheme of single-cell tracking. (**b**) Processes of cell-fate simulation.

1577

Figure 7-figure supplement 2. Cell-fate simulation algorithm. Outlines of the cell fate simulation

algorithm are shown. (**a**) Overall summary of the algorithm, (**b**) creation of Operation data and its types,

(**c**) simulation arrays and its format, (**d**) initial event assignment, (**e**) initial cell-lineage database

generation, (**f**) main event assignment, (**g**) re-assignment of events in cases with cell fusion, (**h**) cell-

lineage database generation, and (**i**) trimming of cell-lineage data ends to switch from the first to

second Operation data. Bipolar cell division (BD), multipolar cell division (MD), cell death (CD), cell

fusion (CF), no cell division (NonDiv), and Ltime (LT).

1585

Figure 7-figure supplement 3. Cell-lineage maps generated by single-cell tracking and a cell-fate

simulation algorithm. (**a**) Cell-lineage maps generated by cell-lineage database Control cells. The

single-cell tracking analysis created 335 cell-lineage datasets, and 16 cell-lineage maps are shown as

examples. (**b**) Cell-lineage maps generated by simulation using Operation data-p53 RNAi. The

simulation created 2,500–5,000 virtual cell-lineage datasets, and 16 cell-lineage maps generated from

the virtually created cell-lineage datasets are shown as examples. Light blue circle, mitosis; pink square,

cell death; orange circle and orange vertical line, cell fusion; black vertical line, bipolar cell division;

red vertical line, multipolar cell division.

1594

Figure 8-figure supplement 1. Determination of number of progenitors required to perform a

reproducible simulation. The simulation was performed with 300–2,000 progenitors using either the

Operation data-Control (**a**) or Operation data-MNNG7 (**b**). Each simulation was repeated five times

and virtual cell-lineage databases for Control-Sim and MNNG7-Sim cells were created to determine

the population expansion curves of the virtual cells. The percentage of variation relative to the average

at simulation time 15,000 min is also shown in parentheses.

1601

Figure 9-figure supplement 1. Cell-doubling time distribution. Cell-doubling time distribution was

determined using the cell lineage database generated by single-cell tracking and simulation data. (**a**, **b**,

c, and **d**) Cell doubling times of (**a**) Control, (**b**) MNNG1, (**c**) MNNG3, and (**d**) MNNG7 cells

determined using the cell-linage database. (**e, f, g** and **h**) Simulations were performed using the

Operation data (**e**) Control, (**f**) MNNG1, (**g**) MNNG3, and (**h**) MNNG7. (**i**, **j**, **k**, and **l**) Cell doubling

time of (**i**) p53 RNAi, (**j**) p53 RNAi-MNNG1, (**k**) p53 RNAi-MNNG3, and (**l**) p53 RNAi-MNNG7. (**m**,

n, **o**, and **p**) Results of corresponding simulations are shown; the Operation data (**m**) p53 RNAi, (**n**)

p53 RNAi-MNNG1, (**o**) p53 RNAi-MNNG3, and (**p**) p53 RNAi-MNNG7. Arrow indicates the average

cell-doubling time, and the average cell-doubling times are shown.

1611

Figure 9-figure supplement 2. Simulation results for numbers of multipolar cell division, cell

death, and cell fusion events. The numbers of multipolar cell division (**a**), cell death (**b**), and cell

fusion (**c**) events in a virtual cell population generated by the simulation were compared with those

determined by single-cell tracking analysis. The simulation was performed with 1,000 progenitors for a

simulation time of 15,000 min and was repeated 10 times. The range of results obtained by the

simulation is also indicated.

1618

SUPPLEMENTARY VIDEOS

Figure 1-video 1. Visualization of cell density. Density of cultured cells visualized using a heat map color scale. Timestamps (min) and a scale bar (80 μm) are shown.

Figure 1-video 2. Visualization of progress of transfection. DIC and particles with 150–255 grayscale and size of 5–30 pixels are shown. Timestamps (min) and a scale bar (20 μm) are shown.

Figure 11-video 1. Cell-fate simulation of progeny of multipolar cell division in Control cells. Progeny of multipolar cell division and the progeny that underwent bipolar cell division are indicated in burgundy and green, respectively. Other cells are indicated in gray. The animation video was created using data from the virtual cell-lineage database and Operation data-Control. After assigning the initial positions of the cells, the cell movement was created programmatically. When cell death occurred, virtual cells were split into small particles.

Figure 11-video 2. Cell-fate simulation of progeny of multipolar cell division in p53 RNAi cells. Progeny of multipolar cell division and the progeny that underwent bipolar cell division are indicated in burgundy and green, respectively. Other cells are indicated in gray. The animation video was created using data from the virtual cell-lineage database and Operation data-p53 RNAi. After assigning the initial positions of the cells, the cell movement was created programmatically. When cell death occurred, virtual cells were split into small particles.

Figure 12-video 1. Cell-fate simulation of p53 RNAi cells in a Control cell population. The p53 RNAi cells are indicated in blue. Other cells are indicated in gray. The animation video was created

1643 using data from the virtual cell-lineage database and Operation data-Control and p53 RNAi. After

1644 assigning the initial position of the cells, cell movement was created programmatically. When cell

1645 death occurred, virtual cells were split into small particles.

1646

Figure 13-video 1. Cell-fate simulation of p53 RNAi cells in a Control cell population subjected to

virtual MNNG treatment. The p53 RNAi cells are indicated in blue. Other cells indicated in gray.

The animation video was created using data from the virtual cell-lineage database and Operation data-

Control and p53si, and MNNG7 and p53 RNAi-MNNG7. The pixel value of the image was changed

after virtual MNNG treatment at a simulation time of 10,000 min. After assigning the initial position of

the cells, cell movement was created programmatically. When cell death occurred, virtual cells were

split into small particles.

1654

SOURCE DATA

Figure 1-source data 1. Quantitation of cell numbers, areas, and densities

Data used in Figure 1-figure supplement 2.

Figure 1-source data 2. Determination of minimum number of cell lineages required to build a cell-lineage database.

Data used in Figure 1-figure supplement 3.

Figure 1-source data 3. Accumulation of particles following siRNA transfection and analysis of p53 expression following transfection and/or MNNG exposure.

Data used in Figure 1-figure supplement 4.

Figure 2-source data 1. Cell-population expansion rates determined by cell counting and single-cell tracking analysis.

Data used in Figure 2.

Figure 3-source data 1. Number of progeny produced from a progenitor.

Data used in Figure 3.

Figure 4-source data 1. Analysis of multipolar cell division, cell death, and cell fusion using counting and single-cell tracking analysis.

Data used in Figure 4.

1679   **Figure 5-source data 1.** *In silico* **generation of cell-lineage database to evaluate the impact of cell**

1680   **death on cell population expansion.**

1681   Data used in Figure 5.

1682

1683   **Figure 6-source data 1. Events leading to the induction of multipolar cell division.**

1684   Data used in Figure 6.

1685

1686   **Figure 8-source data 1. Determination of number of progenitors required to perform a**

1687   **reproducible simulation.**

1688   Data used in Figure 8-figure supplement 1.

1689

1690   **Figure 9-source data 1. Simulations of effects of p53 silencing and MNNG exposure on cell**

1691   **population expansion.**

1692   Data used in Figure 9.

1693

1694   **Figure 9-source data 2. Cell-doubling time distribution.**

1695   Data used in Figure 9-figure supplement 1.

1696

1697   **Figure 9-source data 3. Simulations of multipolar cell division, cell death, and cell fusion**.

1698   Data used in Figure 9-figure supplement 2.

1699

1700   **Figure 10-source data 1. Virtual dose-response analysis.**

1701   Data used in Figure 10.

1702

1703   **Figure 11-source data 1. Simulation of growth of the progeny of multipolar cell division.**

1704   Data used in Figure 11.

1705

1706   **Figure 12-source data 1. Limiting expansion of p53 RNAi cells in a Control cell population.**

1707   Data used in Figure 12.

1708

1709   **Figure 13-source data 1. Expansion of p53 RNAi cells in a Control cell population subjected to**

1710   **virtual MNNG exposure.**

1711   Data used in Figure 13.

1712